Mouse and Human Antibodies that Bind HLA-E-Leader Peptide Complexes and Enhance

NK Cell Cytotoxicity

Dapeng Li[1,5,8,9,*], Simon Brackenridge[2,8], Lucy C. Walters[2,8], Olivia Swanson[1], Karl Harlos[3],

Daniel Rozbesky[3,4], Derek W. Cain[1,5], Kevin Wiehe[1,5], Richard M. Scearce[1], Maggie Barr[1],

Zekun Mu[1], Robert Parks[1], Max Quastel[2], Robert J. Edwards[1,5], Yunfei Wang[1,5], Wes

Rountree[1,5], Kevin O. Saunders[1,6,7], Guido Ferrari[7], Persephone Borrow[2], E. Yvonne Jones[3], S.

Munir Alam[1,5], Mihai L. Azoitei[1,5,*], Geraldine M. Gillespie[2,*], Andrew J. McMichael[2,*], Barton F.

Haynes[1,6,*]

[1]Duke Human Vaccine Institute, Duke University School of Medicine, Durham, NC 27710, USA

[2]Nuffield Department of Clinical Medicine, University of Oxford, Oxford, OX3 7FZ, UK

[3]Division of Structural Biology, Wellcome Centre for Human Genetics, University of Oxford,

Oxford, OX3 7BN, UK

[4]Department of Cell Biology, Charles University, Prague, 12800, Czech Republic

[5]Department of Medicine, Duke University School of Medicine, Durham, NC 27710, USA

[6]Department of Immunology, Duke University School of Medicine, Durham, NC 27710, USA

[7]Department of Surgery, Duke University School of Medicine, Durham, NC 27710, USA

[8]Authors contributed equally

[9]Lead contact

*Address correspondence to dapeng.li@duke.edu, mihai.azoitei@duke.edu,

geraldine.gillespie@ndm.ox.ac.uk, andrew.mcmichael@ndm.ox.ac.uk and

barton.haynes@duke.edu

ABSTRACT (155 words)

HLA-E is a non-classical class Ib molecule that has limited polymorphism and binds HLA class Ia leader peptides (VL9). HLA-E-VL9 complexes interact with the natural killer (NK) cell receptors NKG2A-C/CD94 and regulate NK cell-mediated cytotoxicity. Here we isolated a murine IgM antibody 3H4, that specifically recognized HLA-E-VL9 bound complexes and enhanced killing of HLA-E-VL9-expressing cells by an NKG2A$^+$ NK cell line. Structural analysis revealed how 3H4 prevents CD94/NKG2A docking on HLA-E-VL9 by binding with an overlapping footprint. Upon *in vitro* maturation, an affinity-optimized 3H4 IgG showed enhanced NK killing of HLA-E-VL9-expressing cells. Remarkably, HLA-E-VL9-specific IgM autoantibodies with similar specificity and functions to 3H4 were subsequently isolated from naïve B cells of cytomegalovirus (CMV)-negative, healthy male human donors. Thus, a repertoire of germline low affinity HLA-E-VL9-reactive antibodies are present in both naïve human and murine B cell repertoires. These antibodies can enhance NK cell cytotoxicity and therefore have potential for therapeutic modulation of NK cell function.

INTRODUCTION

Natural killer (NK) cells play critical roles in immune surveillance by discriminating normal

from altered cells, and function by killing non-self malignant or pathogen-infected cells and by

producing inflammatory cytokines (Chiossone et al., 2018; Raulet, 2006; Yokoyama and Kim,

2006). Specific recognition of abnormal cells by NK cells relies on a series of activating and

inhibitory receptors, including the killer immunoglobulin-like receptor (KIR) family in humans and

NKG2/CD94 heterodimeric receptors (Andre et al., 2018; Chiossone et al., 2018). NK cell

inhibitory receptors ligate human lymphocyte antigen (HLA) or major histocompatibility complex

(MHC) class I molecules expressed on healthy cells as self. Conversely, cells lacking MHC

class I are recognized by NK cells as "missing-self" and are sensitive to NK cell-mediated killing

(Ljunggren and Karre, 1985, 1990). In humans, KIRs recognize specific classical HLA class Ia

molecules (Colonna and Samaridis, 1995; Karlhofer et al., 1992; Pende et al., 2019), whereas

the inhibitory NKG2A/CD94 heterodimeric receptor interacts with the non-classical HLA class Ib

molecule HLA-E and is balanced by an activating receptor NKG2C/CD94 (Braud et al., 1997;

Braud et al., 1998; Brooks et al., 1997). While KIR expression is heterogeneous, NKG2A/CD94

is expressed on ~40% of human NK cells (Andre et al., 1999; Mahapatra et al., 2017; Pende et

al., 2019). Unlike classical HLA class I molecules, HLA-E has limited polymorphism with only

two expressed variants, HLA-E*01:01 and HLA-E*01:03, that differ only in residue 107, which is

outside the peptide-binding groove (Kraemer et al., 2014). The NKG2A/CD94/HLA-E pathway is

an important immune checkpoint and has recently become a focus for NK cell-based

immunotherapeutic strategies (Andre et al., 2018; Hu et al., 2019; Kim et al., 2019; Souza-

Fonseca-Guimaraes et al., 2019; van Hall et al., 2019). A subset of CD8+ T cells also express

NKG2A/CD94, and inhibition of the NKG2A/CD94 - HLA-E interaction has similar application in

CD8+ T cell-based immunotherapy (Andre et al., 2018; van Montfoort et al., 2018).

HLA-E engages with NKG2A/CD94 via a restricted subset of peptides VMAPRT(L/V)

(V/L/I/F)L (designated VL9) that derive from the leader sequence of HLA-A, -C, -G and a third of

HLA-B molecules (Braud et al., 1997; Braud et al., 1998; Lee et al., 1998a; Lee et al., 1998b).

HLA-E binds VL9 peptides stabilizing its surface expression (Braud et al., 1997; Braud et al.,

1998). This indicates that HLA-Ia expression is not perturbed by a pathogenic process and

initiates recognition by NKG2A/CD94 or NKG2C/CD94 on NK cells. The binding affinity of HLA-

E-VL9 peptide complexes for NKG2A/CD94 is greater than that for NKG2C/CD94, so that the

inhibitory signal dominates to suppress aberrant NK cell-mediated cytotoxicity and cytokine

production (Aldrich et al., 1994; Braud et al., 1998; Kaiser et al., 2008; Llano et al., 1998; Rolle

et al., 2018). As a secondary function, HLA-E and its murine and rhesus macaque homologs are

capable of binding to a range of other host- and pathogen-derived peptides, including heat-

shock protein 60 (Hsp60)-derived peptides (Michaelsson et al., 2002), Mycobacterium

tuberculosis (Mtb) peptides (Joosten et al., 2010; van Meijgaarden et al., 2015), and simian

immunodeficiency virus (SIV) Gag peptides (Hansen et al., 2016; Walters et al., 2018).

However, the VL9 peptide binds HLA-E with higher affinity and dominates the peptidome

(McMurtrey et al., 2017). Only HLA-E-VL9 can engage CD94/NKG2A to protect cells from NK

cell cytotoxicity (Kraemer et al., 2015; Michaelsson et al., 2002; Sensi et al., 2009). Hence,

leader sequence VL9 peptides are essential not only for stabilizing HLA-E surface expression

but also for mediating HLA-E/NKG2A/CD94 regulation of NK cell self-recognition.

Natural autoantibodies are a specific class of antibodies that recognize self molecules and

populate the B cell germline repertoire in healthy individuals. These antibodies mostly comprise

non-mutated IgM isotypes and can participate in diverse immune functions ranging from

immune defense and regulation to immune pathology. Autoantibodies that recognize both

classical HLA-Ia (Alberu et al., 2007; Morales-Buenrostro et al., 2008) and HLA-E heavy chains

(Ravindranath et al., 2010a; Ravindranath et al., 2010b) have been reported previously. In

certain instances these antibodies have been implicated as contributors to allograft damage in

non-alloimmunised individuals (Hickey et al., 2016; McKenna et al., 2000). However, very little is

unknown beyond this, especially in relation to their frequencies, specificities and functions.

Here, we focused on HLA-E, and initially isolated a murine IgM monoclonal antibody (mAb)

3H4 that bound specifically to HLA-E-VL9 on target cells and enhanced NK cytotoxicity

mediated by an NKG2A+ NK cell line. Crystallographic analysis showed that 3H4 and

CD94/NKG2A cannot simultaneously bind to HLA-E-VL9 and that the heavy chain CDR3

residues at the 3H4-HLA-E-VL9 binding interface were germline-encoded. While 3H4 mAb

enhanced NK cytotoxicity as an IgM, the IgG form of the antibody did not. However,

mutagenized 3H4 IgG variants with enhanced HLA-E-VL9 binding affinity blocked NKG2A

mediated inhibition of NK cells. We subsequently screened healthy humans blood donors and

identified HLA-E-VL9-reactive, near-germline IgMs autoantibodies from human naïve B cell

repertoires. Some of these antibodies also enhanced NK cell killing as IgG subtypes. Thus, we

identified a group of near germline HLA-E-VL9-targeting antibodies in mice and male CMV

seronegative humans that have the potential to regulate NK cell function.

RESULTS

Isolation of murine HLA-E-VL9-specific mAb 3H4

With the original intention of raising monoclonal antibodies to the HIV-1 Gag peptide

RMYSPTSIL (RL9HIV) an HIV-1 Gag epitope previously described (Hansen et al., 2016), we

immunized human HLA-B27/β2-microglobulin (β2M) transgenic mice (Taurog et al., 1990)

(*Figure. S1a-b*) with 293T cells transfected with surface-expressed single-chain HLA-E-RL9HIV

complexes (Yu et al., 2002) (*Figure. S1c-d*). We produced hybridomas, and screened culture

supernatants for binding on a panel of 293T cells transfected with either single-chain HLA-E-

RL9HIV peptide complexes, or with single-chain HLA-E-VL9 peptide complexes as a control.

Unexpectedly, we isolated a subset of antibodies that specifically reacted with HLA-E-VL9

peptide, the most potent of which was the IgM mAb 3H4. Unlike the well-characterized pan-

HLA-E mAb 3D12 (Marin et al., 2003), 3H4 reacted specifically with HLA-E-VL9 (VMAPRTLVL)

and not with control, non-VL9 HLA-E-peptide complexes (*Figure. 1a*). Mab 3H4 also bound to

VL9 peptide-pulsed HLA-class I negative K562 cells transfected with HLA-E (Lampen et al.,

2013) (*Figure. 1b*) and also to soluble HLA-E refolded with synthetic VL9 peptide in both ELISA

(*Figure. 1c*) and surface plasmon resonance (SPR) assays (*Figure. 1d*). SPR measurents

showed that the 1:1 dissociation constants (K_Ds) of IgM 3H4 and human IgG1 backbone 3H4

for soluble HLA-E-VL9 were 8.1 and 49.8 µM, respectively (*Figure. S1e*).

Sequence analysis of 3H4 mAb revealed 1.04% heavy chain variable region (V_H) and

2.51% light chain viable region (V_L) mutations *(Table S1)*. We isolated 3 additional HLA-E-VL9

mouse mAbs from two additional immunization studies in mice (see Methods), and each of the

four HLA-E-VL9 antibodies were minimally mutated IgMs (mean VH and VL mutations, 1.21%

and 2.87%, respectively (*Table S1*). Negative stain electron microscopy showed that the 3H4

IgM hybridoma antibody was predominantly pentameric with a small proportion of hexamers

(*Figure. S1f-g*). In addition, 3H4 was not autoreactive in anti-nuclear antibody or clinical

autoantibody assays (*Figure. S1h-i*).

3H4 IgM recognized the α1/α2 domain of HLA-E and N-terminus of the VL9 peptide

To map the epitope on the HLA-E-VL9 complex recognized by 3H4, we tested 3H4 binding

to VL9 peptide presented by HLA-E, the rhesus ortholog Mamu-E, as well as two HLA-E/Mamu-

E hybrids – one with HLA-E α1/Mamu-E α2 (Hα1/Mα2), the other with Mamu-E α1/HLA-E α2

(Mα1/Hα2). 3H4 did not bind to Mamu-E/VL9 or Hα1/Mα2-VL9, and its staining of cells

expressing Mα1/Hα2-VL9 was weak (*Figure. 1e*), suggesting that 3H4 recognition involved

interaction with both α1 and α2 domains of HLA-E, and the epitope on α2 might be partially

conserved between human and rhesus. 3H4 also did not cross-react with mouse ortholog Qa-1b

(*Figure. S1j*). Moreover, VL9 mutations indicated that position 1 (P1) of the peptide was

important for 3H4 binding (*Figure. 1f*), with strong antibody recognition of VL9 peptide P1

variants with alanine, cysteine, isoleucine, serine, threonine, weak binding to histidine and

proline substitutions, but no interaction with arginine, glutamate, glycine, lysine, methionine,

asparagine, tryptophan, tyrosine or phenylalanine (*Figure. 1g and S1k*). These data suggested

that mAb 3H4 made contacts with both the HLA-E α1/α2 domain and the amino-terminal end of

the VL9 peptide.

Co-complex crystal structure of a 3H4 Fab bound to HLA-E-VL9

A co-complex structure of 3H4 Fab bound to HLA-E-VL9, which packed in the C2 space

group and diffracted to 1.8 Å (*Table S2a*), was obtained. One of the two copies present in the

asymmetric unit is discussed here. 3H4 docked onto the N-terminal region of the HLA-E-

peptide-binding groove making contacts with both the heavy chain α-helices in addition to VL9

peptide residues 1-4 (*Figure. 2a-b*). The 3H4-HLA-E interface mainly comprised electrostatic

interactions and was dominated by the 3H4 VH chain which created a total buried surface area

(BSA) of 1109.4 $Å^2$ and formed ten hydrogen bonds (H-bonds) and three salt bridges with HLA-

E α1-helix residues and one H-bond with T163 of the HLA-E α2-helix. The smaller 3H4 VL

chain-HLA-E interface buried 522.8 $Å^2$ and involved only three inter-molecular H-bonds and

three salt bridges (*Tables S2d-h*). Superposition of the 3H4-HLA-E-VL9 co-complex with a

published HLA-E-bound CD94/NKG2A structure (Kaiser et al., 2008; Petrie et al., 2008)

revealed steric clashes between the VH and VL domains of 3H4 and the CD94 and NKG2A

subdomains, respectively (*Figure. 2c-d*). Moreover, seven HLA-E heavy chain residues (α1

helix positions 58, 59, 62, 63 and α2 helix positions 162, 163 and 167) are shared 3H4-HLA-E

and CD94/NKG2A-HLA-E footprints (*Figure. 2E-F*). Consequently, simultaneous docking of

these two HLA-E binding partners, 3H4 and NKG2A/CD94, would likely be disallowed.

All four 3H4-derived residues that interfaced with the VL9 peptide (Y97, S100, S100A and

Y100B) resided within the VH CDR3 D-junction and were germline-encoded. This 3H4-VL9

interface was characterized by weak Van der Waals and hydrophobic contacts, for example,

Y100B (3H4) and V1 or P4 (VL9) (*Figure. 2g*). Further, positioning of the Y100B (3H4) side

chain directly above V1 (VL9) in part explained preference for small side chains at this peptide

position and the dramatic reductions in 3H4 binding to VL9 variants with larger H or F residues

at position 1 (*Figure. 1g*). Unique shape complementarity also featured at this interface with the

side chains of S100 and S100A (3H4) wrapping around the cyclic side chain of P4 (VL9).

The germline-encoded VH CDR3 D-junction residues that formed the 3H4-VL9 interface

(Y97, S100, S100A and Y100B), also mediated key HLA-E heavy chain contacts. The surface

loop residues A93-V102 swept across the HLA-E-peptide-binding groove forming H-bonds with

both the α1 and α2 helices; HLA-E α2 helix T163 formed an H-bond with S100 (3H4), and HLA-

E α1-helix R62 formed two H-bonds with the Y100B (3H4) mainchain and an additional H-bond

with the main chain of S100A (3H4) (*Figure. 2h*). Y100B (3H4) was involved in multiple polar pi

stacking interactions. Not only was the Y100B side chain sandwiched between R62 and W167

of the HLA-E α1 and α2 helices, respectively, R62 (HLA-E α1) was also positioned between the

aromatic rings of the VH CDR3 domain Y100B and W100D residues.

Key contacts outside the germline-encoded CDR3 D-junction region were also formed at

the 3H4 VH- and VL-HLA-E interfaces. For 3H4 HC, the VH CDR2 region (residues I51-T57)

was positioned above the HLA-E α1-helix where numerous inter-molecular H-bonds were

formed involving VH CDR2 residues G56 and N54 in addition to D50, Q61 and K64 of the VH

framework region (*Figure. 2h*). Critically, HLA-E α1-helix R65 residue formed four H-bonds with

the 3H4 VH and also mediated polar pi stacking interactions with W100D of the VH CDR3 loop.

For 3H4 LC, D92 and E93 of the VL CDR3 loop H-bond with K170 of the HLA-E α2-helix and

N30 of the VL CDR1 loop formed an H-bond with the α2-helix residue, E166, of HLA-E (*Figure.

2i*). Notably, the four key interfacing residues of the 3H4 VH CDR3 D-junction (Y97, S100,

S100A and Y100B) were germline-encoded (*Figure. 2j*).

3H4 IgM enhanced NK cell cytotoxicity against HLA-E-VL9-expressing target cells

Given the suppressive role of the HLA-E-VL9/NKG2A/CD94 pathway in NK cell function, we

tested whether the binding of mAb 3H4 to HLA-E-VL9 could enhance NK cell killing of target

cells (*Figure. 3a*). An NKG2A/CD94-positive, CD16/CD32/CD64-negative human NK cell line,

NK92 (*Figure. S2a-b*), exhibited significantly increased cytotoxicity against HLA-E-VL9-

transfected 293T cells (*Figure. 3b; P<0.0001,* mixed effects models) but not against non-HLA-

E-expressing 293T cells (*Figure. 3c*) in the presence of 3H4 IgM compared to an isotype

control IgM. In addition, we tested a combination of 3H4 with the NKG2A specific murine

antibody, Z199. While Z199 alone enhanced NK killing against HLA-E-VL9 expressing cells, no

additional elevation of killing was observed with the combination of mAbs 3H4 and Z199

(*Figure. 3d-e*), suggesting that killing enhancement was maximal with either 3H4 or Z199 alone.

These data demonstrated that HLA-E-VL9-specific IgM mAb 3H4 could enhance the killing

capacity of NKG2A+ NK cells *in vitro* by binding to HLA-E-VL9 on target cells.

The majority of multimeric IgM is restricted to serum and lymph and does not penetrate well

into tissues (Sathe and Cusick, 2021). Therefore, we constructed a recombinant 3H4 IgG in a

human IgG1 backbone and tested it for ability to enhance NK92 cell killing of HLA-E-VL9 target

cells. In contrast to 3H4 IgM, 3H4 IgG could not mediate enhancement of NK cell killing (*Figure.*

S3a). Thus, either the affinity of the 3H4 Fab on an IgG was too low (K_D = 49.8 µM; *Figure.*

S1e), or a multimeric antibody is needed for for efficient blocking of HLA-E-VL9 binding to

NKG2A/CD94 to enhance of NK killing.

To distinguish between the need for higher affinity versus multimerization of the IgM

antibody for enhanced NK killing activity, we developed and analyzed 3H4 antibody libraries

using high-throughput screening on the surface of yeast (*Figure. 4a*). A library was built that

contained ~1.1 million 3H4 scFv variants with amino acid diversity at sites that were determined

by structural analysis to interact with HLA-E-VL9 (*Figure. 4b*). Seventeen total residues located

in the CDR loops of 3H4 were randomized in groups of four based on their proximity, and all

possible combinations of amino acids were sampled at these sites (*Figure. S3b*). The resulting

3H4 scFv library was transformed into yeast and screened for three rounds by fluorescence-

activated cell sorting (FACS) for binding to fluorescently labeled HLA-E-VL9 tetramer (*Figure.*

4c). Eleven 3H4 variants were selected for experimental characterization as recombinant

human IgGs from the highly represented clones remaining in the library upon the final selection

round. These novel Abs (3H4 Gv1 to 3H4 Gv12) were mutated at positions 97-100 of the CDR

H3 loop. Compared to the original 3H4 mAb, the optimized antibodies predominantly contained

small amino acids at positions 97 and 98, a polar amino acid at position 99, and a large

aromatic at position 100 that is closest to the HLA-E-VL9 (*Figure. 4d*).

We next expressed all eleven 3H4 Gv antibodies recombinantnly as human IgGs, and

confirmed that they had higher binding than wild-type 3H4 IgG on cell surface expressed HLA-

E-VL9 (*Figure. 4e and S3c*). Two 3H4 variants, 3H4 G3v and 3H4 G6v, were selected for

affinity and functional analysis. SPR measurents showed that the 1:1 dissociation constants

(K_Ds) of the selected 3H4 variants for soluble HLA-E-VL9 were markedly improved compared to

that of wild-type 3H4 that had a K_D of 49.8 µM. 3H4 G3v showed the tightest HLA-E-VL9

binding, with a K_D of 220 nM, representing a ~226-fold improvement in affinity over the WT mAb

(*Figure. 4f*). In the NK cytotoxicity assay, the optimized 3H4 mAbs enhanced NK-92 cell killing

of HLA-E-VL9-transfected 293T cells at concentrations of 10 µg/ml and 1 µg/ml to levels

comparable to those observed for 3H4 IgM (*Figure. 4g and S3d*). Therefore, the higher affinity

of affinity-optimized 3H4 IgG for HLA-E-VL9 could compensate for the need for avidity effect of

3H4 IgM multimers to mediate *in vitro* NK enhancement.

Isolation of near-germline HLA-E-VL9-specific antibodies from CMV-negative, healthy

humans

We next asked if similar HLA-E-VL9 antibodies were present in the naïve B cell receptor

(BCR) repertoire in humans and whether they could enhance NK killing of target cells. Using

HLA-E-VL9 tetramers as probes, we identified B cells expressing HLA-E-VL9-specific B cell

receptors (BCRs) in four male, cytomegalovirus (CMV) seronegative human donors (*Figure. 5a*

and S4a, Table S3). We isolated 56 HLA-E-VL9-specific antibodies that reacted with HLA-E-

VL9 complexes but not control HLA-E-peptide complexes (**Figure. 5b and S4b-c, Table S4**); all

were IgM (**Table S4**). By performing more in-depth analysis of the binding profiles of four

representative HLA-E-VL9 antibodies - CA123, CA133, CA143 and CA147, we found that these

antibodies exhibited differential cross-reactivities with rhesus Mamu-E-VL9 or mouse Qa-1-VL9

complexes (**Figure. S4d**) in addition to distinct binding specificities to VL9 peptide variants

(**Figure. S4e**) . The apparent affinities (K_D) of CA123 and CA147 on a human IgG1 backbone to

soluble HLA-E-VL9 were 3.8 and 25.0 µM, respectively (**Figure. S4f**). Human HLA-E-VL9

antibodies CA147 and CA123 tested in functional NK killing assays as a recombinant human

IgG1. CA147 enhanced NK-92 cell cytotoxicity to HLA-E-VL9-expressing target cells (**Figure.**

5c), whereas CA123 had no enhancing effect (**Figure. S4g-h**), suggesting that NK killing-

enhancement function of the HLA-E-VL9 antibodies was determined by factors beyond binding

affinity.

In the four humans, the percentages of HLA-E-VL9-specific B cells in pan-B cells (CD3$^-$

CD235$^-$CD14$^-$CD16$^-$CD19$^+$) were 0.0009%-0.0023% (mean of 0.0014%) (**Figure. 5d**). HLA-E-

VL9-specific B cells were IgD$^+$IgM$^{+/-}$ B cells, in which four cell subsets were observed (**Figure.**

5e) – CD10$^-$CD27$^-$CD38$^{+/-}$ naïve B cells (71.4%), CD10$^+$CD27$^-$CD38^{++} immature or newly formed

B cells (Giltiay et al., 2019) (10.7%), and CD10$^-$CD27$^+$CD38$^-$ non-class-switched memory cells,

demonstrating that BCRs targeting HLA-E-VL9 peptide existed in the naïve B cell repertoire of

healthy humans.

V$_H$/V$_L$ gene usage of HLA-E-VL9-specific antibodies

To characterize the human antibody gene usage of HLA-E-VL9 antibodies, we analyzed the

paired heavy chain and light chain gene sequences of 56 human HLA-E-VL9 antibodies, and

found 1 multiple-member clone containing 6 antibodies in donor LP021 (Kepler et al., 2014)

(**Table S4**). Next, we compared the 51 HLA-E-VL9-specific B cell clones with a reference

human antibody repertoire (DeKosky et al., 2015). Over 45% of the heavy chain variable region

(V_H) genes were VH3-21 or VH3-11 in HLA-E-VL9 antibodies, whereas less than 7% of the

control B cells used these two genes (***Figure. 5f, Table S4***). HLA-E-VL9 antibody light chain

variable regions (V_κ/V_λ) also were skewed and preferentially utilized IGKV3-15, IGKV1-39 and

IGKV3-11 genes compared to controls (***Figure. 5g, Table S4***). No J chain gene usage

preference was observed (***Figure. S5a-d***). Moreover, HLA-E-VL9 antibodies showed a trend to

have shorter heavy chain complementarity determining region 3 (CDR3) lengths than reference

antibodies (***Figure. 5h***), while no difference was observed for light chain CDR3 (***Figure. 5i***).

Given that HLA-E-VL9 antibodies were IgMs derived primarily from naïve or immature B cells,

we compared the mutation frequencies of the 51 clones with a reference human antibody

repertoire containing both naïve and antigen-experienced antibodies (DeKosky et al., 2016).

Both HLA-E-VL9 antibody heavy and light chain variable region genes exhibited low somatic

mutation rates that were similar to naïve B cell controls (***Figure. 5j-k***). Thus, human HLA-E-

VL9-specific antibodies were IgM, minimally mutated and displayed skewed usage of V_H and

V_κ/V_λ genes.

DISCUSSION

In this study, we have isolated and characterized antibodies reactive with HLA-E-VL9

peptide complexes, and found these antibodies were derived from the naïve IgM B cell BCR

repertoire in mice as well as in human blood donors. We selected HCMV negative donors to

avoid possible cross reacting antibody responses to the VL9 sequence present in the signal

peptide of the UL40 protein, and male donors to exclude anti-HLA antibodies made in

pregnancy. Somatic mutations of these antibodies were minimal, and the affinities of these

antibodies for HLA-E-VL9 were low. The lack of class-switching in HLA-E-VL9-specific

antibodies may reflect self-tolerance of CD4 T cells and a lack of T cell help for affinity

maturation of these antibodies. While the mouse antibodies were selected in the setting of HLA-

E-unrelated peptide immunizations, they were minimally mutated IgM antibodies, as were the

antibodies isolated from human CMV-negative, healthy males. Structural analysis of the HLA-E-

VL9-3H4 Fab co-complex revealed that the murine 3H4 heavy chain made key contacts with

HLA-E and the VL9 peptide using germline-encoded residues in the CDR-H3 (D) region.

However, 3H4 is a mouse antibody that reacted with human HLA-E-VL9. The HLA-E equivalent

in C57BL/6xSJL mice is Qa1b which presents a similar class Ia signal peptide AMAPRTLLL but

3H4 did not bind to this HLA-E-peptide complex. Therefore, it remains unclear how 3H4 and

possibly the other HLA-E-VL9-specific IgM antibodies were genetically selected as germlines or

induced with minimal mutation in the setting of HLA-E-unrelated peptide immunizations. In

contrast, the human antibodies such as CA147 that bound to human HLA-E-VL9 were auto-

antibodies with near germline sequences that exist in some, possibly many, individuals.

Autoantibodies to HLA-Ia (Alberu et al., 2007; Morales-Buenrostro et al., 2008) and HLA-E

heavy chains (Ravindranath et al., 2010a; Ravindranath et al., 2010b) have been detected in

non-alloimmunized males, and can contribute to allograft damage (Hickey et al., 2016;

McKenna et al., 2000). It has been suggested that the HLA-E antibodies in non-alloimmunized

humans could be elicited by autoantigens derived from soluble HLA-E heavy chains that

become immunogenic without the β2M subunit, or viral/bacterial agents cross-reactive with

HLAs (Alberu et al., 2007; Hickey et al., 2016; Ravindranath et al., 2010a; Ravindranath et al.,

2010b). It is of interest that human cytomegalovirus (CMV), which encodes the VL9 sequence

VMAPRTLIL in the leader sequence of its UL40 gene. This peptide is processed in a TAP

independent manner and presented bound to HLA-E at the cell surface to inhibit NK cell killing

and evade innate immune responses (Tomasec et al., 2000). This has not been reported to

elicit antibody responses, but HLA-E-UL40 peptide-specific T cells have been described when

the limited polymorphism in the HLA A, B and C sequences mismatches that of the virally-

encoded VL9 peptide sufficiently to overcome self-tolerance (Sullivan et al., 2015). However,

the subjects in our study were all HCMV seronegative and male. The possibility that germline

encoded anti-HLA-E-VL9 antibodies play a physiological role in regulating NK cell function in

several species needs further exploration.

Harnessing NK cells to attack tumor cells has emerged as an attractive strategy for cancer

immunotherapies (Guillerey et al., 2016; Lowry and Zehring, 2017). A promising target for

therapeutic immune-modulation of NK cell functions is the inhibitory NKG2A/CD94-HLA-E-VL9

interaction. Monalizumab, the first-in-class monoclonal antibody checkpoint inhibitor targeting

NKG2A, enhances anti-tumor immunity by activating cytotoxic activities of effector CD8+ T cells

and NK cells (Andre et al., 2018; Creelan and Antonia, 2019; van Hall et al., 2019). In our study,

co-complex structural analysis revealed steric clashes between the 3H4 Fab and the NK

inhibitory receptor NKG2A/CD94 when docked onto HLA-E-VL9, showing how 3H4 IgM can

enhance NKG2A+ NK cell killing. Both mouse 3H4 IgM, the affinity-optimized 3H4 IgG, and the

recombinant IgG1 form of human CA147 enhanced the cytotoxicity of an NKG2A+ human NK

cell line NK92, which is a safe and established cell line that has been used for adoptive

immunotherapy in phase I clinical trials (Klingemann et al., 2016). Thus, HLA-E-VL9-targeting

antibodies 3H4 and CA147 could have therapeutic potential as NK checkpoint inhibitors.

METHODS

Detailed methods are provided in the supplemental online material.

ACKNOWLEDGMENTS

We thank Duke Human Vaccine Institute (DHVI) programs and finance staff for project

oversight and the contributions of technical staff at the DHVI, including Jordan Cocchiaro, Kelly

Soderberg, Ahmed Yousef Abuahmad, Yunfei Wang, Giovanna Hernandez, Esther Lee, Paige

Power, Aja Sanzone, Brenna Harrington, Andrew Foulger, Amanda Newman, Cindy Bowman,

Grace Stevens, Laura Sutherland, Margaret Deyton, Victoria Gee-Lai, Tarra Von Holle, Thad

Gurley, Madison Berry, Kara Anasti, Katayoun Mansouri, Erika Dunford, and Dawn Jones

Marshall. We thank the DHVI Flow Cytometry Core and Duke Cancer Institute Flow Cytometry

Shared Resource (FCSR) for technical assistance. This study was funded by the Collaboratory

of AIDS Researchers for Eradication (CARE; UM1AI126619) (B.F.H.) and the Bill and Melinda

Gates Foundation OPP1108533 (A.J.M.). P.B and A.J.M. are Jenner Institute Investigators.

ACCESSION NUMBERS

The 3H4-HLA-E-VL9 co-complex structure determined in this study has the PDB accession

code, 7BH8.

AUTHOR CONTRIBUTIONS

D.L. isolated and characterized antibodies, performed ^{51}Cr release assay, and analyzed

data. S.B. made single-chain trimer constructs, performed epitope mapping experiments and

analyzed data. G.M.G, L.W. and M.Q. prepared the antigens and HLA-E tetramers. L.W.,

G.M.G., D.R. and K.H performed structural experiments with oversight from E.Y.J.. O.S. and

M.L.A. performed antibody affinity optimization work. D.W.C. helped with flow cytometry sorting

set-up. R.S., R.P. and M.B. performed hybridoma experiments, ELISA assays and help with

^{51}Cr release assays. R.J.E. oversaw negative stain EM. M.A. supervised and interpreted the

SPR experiments. K.W. oversaw and performed antibody gene sequence analysis. Z.M. and

M.B. helped with antibody isolation. Y.W. and W.R. provided statistical analysis. K.O.S. oversaw

antigen production. G.F. and P.B. provided advice on NK cell assays and contributed to study

design and data interpretation. B.F.H. and A.J.M. conceived, designed, coordinated the study.

D.L. and B.F.H. wrote the manuscript, which was edited by L.W., S.B., O.S., M.L.A., P.B.,

G.M.G., and A.J.M., and reviewed by all authors.

DECLARATION OF INTERESTS

D.L., S.B., M.L.A., G.M.G., A.J.M. and B.F.H. have patents submitted on antibodies and

select methods in this paper.

REFERENCES

Alberu, J., Morales-Buenrostro, L.E., de Leo, C., Vargas-Rojas, M.I., Marino-Vazquez, L.A., and

Crispin, J.C. (2007). A non-allogeneic stimulus triggers the production of de novo HLA

antibodies in healthy adults. Transpl Immunol *18*, 166-171.

Aldrich, C.J., DeCloux, A., Woods, A.S., Cotter, R.J., Soloski, M.J., and Forman, J. (1994).

Identification of a Tap-dependent leader peptide recognized by alloreactive T cells specific for a

class Ib antigen. Cell *79*, 649-658.

Andre, P., Brunet, C., Guia, S., Gallais, H., Sampol, J., Vivier, E., and Dignat-George, F. (1999).

Differential regulation of killer cell Ig-like receptors and CD94 lectin-like dimers on NK and T

lymphocytes from HIV-1-infected individuals. Eur J Immunol *29*, 1076-1085.

Andre, P., Denis, C., Soulas, C., Bourbon-Caillet, C., Lopez, J., Arnoux, T., Blery, M.,

Bonnafous, C., Gauthier, L., Morel, A.*, et al.* (2018). Anti-NKG2A mAb Is a Checkpoint Inhibitor

that Promotes Anti-tumor Immunity by Unleashing Both T and NK Cells. Cell *175*, 1731-1743

e1713.

Braud, V., Jones, E.Y., and McMichael, A. (1997). The human major histocompatibility complex

class Ib molecule HLA-E binds signal sequence-derived peptides with primary anchor residues

at positions 2 and 9. Eur J Immunol *27*, 1164-1169.

Braud, V.M., Allan, D.S., O'Callaghan, C.A., Soderstrom, K., D'Andrea, A., Ogg, G.S., Lazetic,

S., Young, N.T., Bell, J.I., Phillips, J.H.*, et al.* (1998). HLA-E binds to natural killer cell receptors

CD94/NKG2A, B and C. Nature *391*, 795-799.

Brooks, A.G., Posch, P.E., Scorzelli, C.J., Borrego, F., and Coligan, J.E. (1997). NKG2A

complexed with CD94 defines a novel inhibitory natural killer cell receptor. J Exp Med *185*, 795-

800.

Chiossone, L., Dumas, P.Y., Vienne, M., and Vivier, E. (2018). Natural killer cells and other

innate lymphoid cells in cancer. Nat Rev Immunol *18*, 671-688.

Colonna, M., and Samaridis, J. (1995). Cloning of immunoglobulin-superfamily members

associated with HLA-C and HLA-B recognition by human natural killer cells. Science *268*, 405-

408.

Creelan, B.C., and Antonia, S.J. (2019). The NKG2A immune checkpoint - a new direction in

cancer immunotherapy. Nat Rev Clin Oncol *16*, 277-278.

DeKosky, B.J., Kojima, T., Rodin, A., Charab, W., Ippolito, G.C., Ellington, A.D., and Georgiou,

G. (2015). In-depth determination and analysis of the human paired heavy- and light-chain

antibody repertoire. Nat Med *21*, 86-91.

DeKosky, B.J., Lungu, O.I., Park, D., Johnson, E.L., Charab, W., Chrysostomou, C., Kuroda, D.,

Ellington, A.D., Ippolito, G.C., Gray, J.J.*, et al.* (2016). Large-scale sequence and structural

comparisons of human naive and antigen-experienced antibody repertoires. Proc Natl Acad Sci

U S A *113*, E2636-2645.

Giltiay, N.V., Giordano, D., and Clark, E.A. (2019). The Plasticity of Newly Formed B Cells. J

Immunol *203*, 3095-3104.

Guillerey, C., Huntington, N.D., and Smyth, M.J. (2016). Targeting natural killer cells in cancer

immunotherapy. Nat Immunol *17*, 1025-1036.

Hansen, S.G., Wu, H.L., Burwitz, B.J., Hughes, C.M., Hammond, K.B., Ventura, A.B., Reed,

J.S., Gilbride, R.M., Ainslie, E., Morrow, D.W.*, et al.* (2016). Broadly targeted CD8(+) T cell

responses restricted by major histocompatibility complex E. Science *351*, 714-720.

Hickey, M.J., Valenzuela, N.M., and Reed, E.F. (2016). Alloantibody Generation and Effector

Function Following Sensitization to Human Leukocyte Antigen. Front Immunol *7*, 30.

Hu, W., Wang, G., Huang, D., Sui, M., and Xu, Y. (2019). Cancer Immunotherapy Based on

Natural Killer Cells: Current Progress and New Opportunities. Front Immunol *10*, 1205.

Joosten, S.A., van Meijgaarden, K.E., van Weeren, P.C., Kazi, F., Geluk, A., Savage, N.D.,

Drijfhout, J.W., Flower, D.R., Hanekom, W.A., Klein, M.R.*, et al.* (2010). Mycobacterium

tuberculosis peptides presented by HLA-E molecules are targets for human CD8 T-cells with

cytotoxic as well as regulatory activity. PLoS Pathog *6*, e1000782.

Kaiser, B.K., Pizarro, J.C., Kerns, J., and Strong, R.K. (2008). Structural basis for NKG2A/CD94

recognition of HLA-E. Proc Natl Acad Sci U S A *105*, 6696-6701.

Karlhofer, F.M., Ribaudo, R.K., and Yokoyama, W.M. (1992). MHC class I alloantigen specificity

of Ly-49+ IL-2-activated natural killer cells. Nature *358*, 66-70.

Kepler, T.B., Munshaw, S., Wiehe, K., Zhang, R., Yu, J.S., Woods, C.W., Denny, T.N.,

Tomaras, G.D., Alam, S.M., Moody, M.A.*, et al.* (2014). Reconstructing a B-Cell Clonal Lineage.

II. Mutation, Selection, and Affinity Maturation. Front Immunol *5*, 170.

Kim, N., Lee, H.H., Lee, H.J., Choi, W.S., Lee, J., and Kim, H.S. (2019). Natural killer cells as a

promising therapeutic target for cancer immunotherapy. Arch Pharm Res *42*, 591-606.

Klingemann, H., Boissel, L., and Toneguzzo, F. (2016). Natural Killer Cells for Immunotherapy -

Advantages of the NK-92 Cell Line over Blood NK Cells. Front Immunol *7*, 91.

Kraemer, T., Blaszczyk, R., and Bade-Doeding, C. (2014). HLA-E: a novel player for

histocompatibility. J Immunol Res *2014*, 352160.

Kraemer, T., Celik, A.A., Huyton, T., Kunze-Schumacher, H., Blaszczyk, R., and Bade-Doding, C.

(2015). HLA-E: Presentation of a Broader Peptide Repertoire Impacts the Cellular Immune

Response-Implications on HSCT Outcome. Stem Cells Int *2015*, 346714.

Lampen, M.H., Hassan, C., Sluijter, M., Geluk, A., Dijkman, K., Tjon, J.M., de Ru, A.H., van der

Burg, S.H., van Veelen, P.A., and van Hall, T. (2013). Alternative peptide repertoire of HLA-E

reveals a binding motif that is strikingly similar to HLA-A2. Mol Immunol *53*, 126-131.

Lee, N., Goodlett, D.R., Ishitani, A., Marquardt, H., and Geraghty, D.E. (1998a). HLA-E surface

expression depends on binding of TAP-dependent peptides derived from certain HLA class I

signal sequences. J Immunol *160*, 4951-4960.

Lee, N., Llano, M., Carretero, M., Ishitani, A., Navarro, F., Lopez-Botet, M., and Geraghty, D.E.

(1998b). HLA-E is a major ligand for the natural killer inhibitory receptor CD94/NKG2A. Proc

Natl Acad Sci U S A *95*, 5199-5204.

Ljunggren, H.G., and Karre, K. (1985). Host resistance directed selectively against H-2-deficient

lymphoma variants. Analysis of the mechanism. J Exp Med *162*, 1745-1759.

Ljunggren, H.G., and Karre, K. (1990). In search of the 'missing self': MHC molecules and NK

cell recognition. Immunol Today *11*, 237-244.

Llano, M., Lee, N., Navarro, F., Garcia, P., Albar, J.P., Geraghty, D.E., and Lopez-Botet, M.

(1998). HLA-E-bound peptides influence recognition by inhibitory and triggering CD94/NKG2

receptors: preferential response to an HLA-G-derived nonamer. Eur J Immunol *28*, 2854-2863.

Lowry, L.E., and Zehring, W.A. (2017). Potentiation of Natural Killer Cells for Cancer

Immunotherapy: A Review of Literature. Front Immunol *8*, 1061.

Mahapatra, S., Mace, E.M., Minard, C.G., Forbes, L.R., Vargas-Hernandez, A., Duryea, T.K.,

Makedonas, G., Banerjee, P.P., Shearer, W.T., and Orange, J.S. (2017). High-resolution

phenotyping identifies NK cell subsets that distinguish healthy children from adults. PLoS One

12, e0181134.

Marin, R., Ruiz-Cabello, F., Pedrinaci, S., Mendez, R., Jimenez, P., Geraghty, D.E., and

Garrido, F. (2003). Analysis of HLA-E expression in human tumors. Immunogenetics *54*, 767-

775.

McKenna, R.M., Takemoto, S.K., and Terasaki, P.I. (2000). Anti-HLA antibodies after solid

organ transplantation. Transplantation *69*, 319-326.

McMurtrey, C., Harriff, M.J., Swarbrick, G.M., Duncan, A., Cansler, M., Null, M., Bardet, W.,

Jackson, K.W., Lewinsohn, D.A., Hildebrand, W.*, et al.* (2017). T cell recognition of

Mycobacterium tuberculosis peptides presented by HLA-E derived from infected human cells.

PLoS One *12*, e0188288.

Michaelsson, J., Teixeira de Matos, C., Achour, A., Lanier, L.L., Karre, K., and Soderstrom, K.

(2002). A signal peptide derived from hsp60 binds HLA-E and interferes with CD94/NKG2A

recognition. J Exp Med *196*, 1403-1414.

Morales-Buenrostro, L.E., Terasaki, P.I., Marino-Vazquez, L.A., Lee, J.H., El-Awar, N., and

Alberu, J. (2008). "Natural" human leukocyte antigen antibodies found in nonalloimmunized

healthy males. Transplantation *86*, 1111-1115.

Pende, D., Falco, M., Vitale, M., Cantoni, C., Vitale, C., Munari, E., Bertaina, A., Moretta, F., Del

Zotto, G., Pietra, G*., et al.* (2019). Killer Ig-Like Receptors (KIRs): Their Role in NK Cell

Modulation and Developments Leading to Their Clinical Exploitation. Front Immunol *10*, 1179.

Petrie, E.J., Clements, C.S., Lin, J., Sullivan, L.C., Johnson, D., Huyton, T., Heroux, A., Hoare,

H.L., Beddoe, T., Reid, H.H*., et al.* (2008). CD94-NKG2A recognition of human leukocyte

antigen (HLA)-E bound to an HLA class I leader sequence. J Exp Med *205*, 725-735.

Raulet, D.H. (2006). Missing self recognition and self tolerance of natural killer (NK) cells. Semin

Immunol *18*, 145-150.

Ravindranath, M.H., Kaneku, H., El-Awar, N., Morales-Buenrostro, L.E., and Terasaki, P.I.

(2010a). Antibodies to HLA-E in nonalloimmunized males: pattern of HLA-Ia reactivity of anti-

HLA-E-positive sera. J Immunol *185*, 1935-1948.

Ravindranath, M.H., Taniguchi, M., Chen, C.W., Ozawa, M., Kaneku, H., El-Awar, N., Cai, J.,

and Terasaki, P.I. (2010b). HLA-E monoclonal antibodies recognize shared peptide sequences

on classical HLA class Ia: relevance to human natural HLA antibodies. Mol Immunol *47*, 1121-

1131.

Rolle, A., Meyer, M., Calderazzo, S., Jager, D., and Momburg, F. (2018). Distinct HLA-E

Peptide Complexes Modify Antibody-Driven Effector Functions of Adaptive NK Cells. Cell Rep

24, 1967-1976 e1964.

Sathe, A., and Cusick, J.K. (2021). Biochemistry, Immunoglobulin M. In StatPearls (Treasure

Island (FL)).

Sensi, M., Pietra, G., Molla, A., Nicolini, G., Vegetti, C., Bersani, I., Millo, E., Weiss, E., Moretta,

L., Mingari, M.C.*, et al.* (2009). Peptides with dual binding specificity for HLA-A2 and HLA-E are

encoded by alternatively spliced isoforms of the antioxidant enzyme peroxiredoxin 5. Int

Immunol *21*, 257-268.

Souza-Fonseca-Guimaraes, F., Cursons, J., and Huntington, N.D. (2019). The Emergence of

Natural Killer Cells as a Major Target in Cancer Immunotherapy. Trends Immunol *40*, 142-158.

Sullivan, L.C., Westall, G.P., Widjaja, J.M., Mifsud, N.A., Nguyen, T.H., Meehan, A.C.,

Kotsimbos, T.C., and Brooks, A.G. (2015). The Presence of HLA-E-Restricted, CMV-Specific

CD8+ T Cells in the Blood of Lung Transplant Recipients Correlates with Chronic Allograft

Rejection. PLoS One *10*, e0135972.

Taurog, J., Hammer, R., Maika, S., Sams, K., El-Zaatari, F., Stimpson, S., and Schwab, J.

(1990). HLA-B27 transgenic mice as potential models of human disease. In Transgenic mice

and mutants in MHC research (Springer), pp. 268-275.

Tomasec, P., Braud, V.M., Rickards, C., Powell, M.B., McSharry, B.P., Gadola, S., Cerundolo,

V., Borysiewicz, L.K., McMichael, A.J., and Wilkinson, G.W. (2000). Surface expression of HLA-

E, an inhibitor of natural killer cells, enhanced by human cytomegalovirus gpUL40. Science *287*,

1031.

van Hall, T., Andre, P., Horowitz, A., Ruan, D.F., Borst, L., Zerbib, R., Narni-Mancinelli, E., van

der Burg, S.H., and Vivier, E. (2019). Monalizumab: inhibiting the novel immune checkpoint

NKG2A. J Immunother Cancer *7*, 263.

van Meijgaarden, K.E., Haks, M.C., Caccamo, N., Dieli, F., Ottenhoff, T.H., and Joosten, S.A.

(2015). Human CD8+ T-cells recognizing peptides from Mycobacterium tuberculosis (Mtb)

presented by HLA-E have an unorthodox Th2-like, multifunctional, Mtb inhibitory phenotype and

represent a novel human T-cell subset. PLoS Pathog *11*, e1004671.

van Montfoort, N., Borst, L., Korrer, M.J., Sluijter, M., Marijt, K.A., Santegoets, S.J., van Ham,

V.J., Ehsan, I., Charoentong, P., Andre, P.*, et al.* (2018). NKG2A Blockade Potentiates CD8 T

Cell Immunity Induced by Cancer Vaccines. Cell *175*, 1744-1755 e1715.

Walters, L.C., Harlos, K., Brackenridge, S., Rozbesky, D., Barrett, J.R., Jain, V., Walter, T.S.,

O'Callaghan, C.A., Borrow, P., Toebes, M.*, et al.* (2018). Pathogen-derived HLA-E bound

epitopes reveal broad primary anchor pocket tolerability and conformationally malleable peptide

binding. Nat Commun *9*, 3137.

Yokoyama, W.M., and Kim, S. (2006). How do natural killer cells find self to achieve tolerance?

Immunity *24*, 249-257.

Yu, Y.Y., Netuschil, N., Lybarger, L., Connolly, J.M., and Hansen, T.H. (2002). Cutting edge:

single-chain trimers of MHC class I molecules form stable structures that potently stimulate

antigen-specific T cells and B cells. J Immunol *168*, 3145-3149.

FIGURES

Figure. 1 Isolation of monoclonal antibody 3H4 specifically targeting HLA-E-VL9

complex.

a, 3H4 bound HLA-E-VL9 single chain trimer (SCT)-transfected 293T cells. All SCT

constructs express EGFP to indicate transfection efficiency. Transfected cells were stained with

test antibody and then an Alexa fluor 555 (AF555)-anti-mouse Ig(H+L) secondary antibody. A

control mouse IgM TE4 was used as a negative control. Anti-pan-HLA-E antibody 3D12 was

used as a positive control. Representative data from one of five independent experiments are

shown.

b, 3H4 bound VL9 peptide pulsed K562-HLA-E cells. RL9HIV, RL9SIV, Mtb44 peptides

served as peptide controls. TE4 and 3D12 were used as antibody controls. Peptide-pulsed cells

were stained with test antibody and then an Alexa fluor 647 (AF647)-anti-mouse Ig(H+L)

secondary antibody. Mean fluorescence intensity (MFI) of each sample is shown.

Representative data from one of three independent experiments are shown.

c-d, 3H4 specifically bound to soluble HLA-E-VL9 complexes as measured by ELISA and

SPR. **c,** ELISA plates were coated with 3H4 or control IgM TE4 in serial dilution, blocked, and

incubated with C-trap-stabilized HLA-E-VL9, HLA-E-RL9HIV, HLA-E-RL9SIV antigens. After

washing, antigen binding was detected by adding HRP-conjugated anti-human β2M antibody. **d,**

For SPR, biotinylated HLA-E-peptide complexes (HLA-E-VL9, HLA-E-RL9SIV, HLA-E-RL9HIV

and mock control) were bound to the immobilized streptavidin. Antibody 3H4 and control TE4

were flowed over sensor chips and antibody binding was monitored in real-time. Representative

data from one of two independent experiments are shown.

e, 3H4 recognized the α2 domain of HLA-E. Flow cytometry analysis of 3H4 and 2M2 (a

control β2M mAb) binding to 293T cells transfected with VL9 presented by HLA-E, Mamu-E, and

two HLA-E/Mamu-E hybrids - one with HLA-E α1/Mamu-E α2 (Hα1/Mα2), the other with Mamu-

E α1/HLA-E α2 (Mα1/Hα2) (green). Transfected cells were stained with test antibody and then

an AF647-anti-mouse Ig(H+L) secondary antibody. Isotype control stained cells were used as

negative controls (grey filled histograms). Representative data from one of three independent

experiments are shown.

f, 3H4 recognized position 1 (P1) of the VL9 peptide. 3H4 and 2M2 (a control β2M mAb)

staining of 293T cells transfected with HLA-E-VL9 (VMAPRTLLL) or HLA-E-VL9 with a mutation

at P1 (valine to arginine; RMAPRTLLL) (blue), and with HLA-E-RL9HIV (RMYSPTSIL) or HLA-

E-RL9HIV with a mutation at P1 (arginine to valine; VMYSPTSIL) (red). Transfected cells were

stained with test antibody and then an AF647-anti-mouse Ig(H+L) secondary antibody. Isotype

control stained cells were used as negative controls (grey filled histograms). Representative

data from one of three independent experiments are shown.

g, 3H4 recognized peptides with variants in P1. 293T cells were transfected with HLA-E

SCTs with VL9 peptides with single amino acid mutations at P1, then stained with 3H4 antibody

followed by AF647 conjugated anti-mouse IgG(H+L) secondary antibody. Cells were gated for

EGFP positive subsets. MFI of 3H4 staining on wildtype VL9 peptide was set as 100%, and the

percentages for binding to mutants calculated as (MFI of 3H4 binding on each P1 variant) / (MFI

of 3H4 binding on wildtype VL9) x 100%.

Figure. 2 3H4 Fab-HLA-E-VL9 co-complex structural visualization.

a-b, 3H4 Fab-HLA-E docking angles. HLA-E heavy chain and β2M light chain: grey; VL9

peptide: lime green; 3H4 heavy chain: light purple; 3H4 light chain: teal.

c-d, Superposition of 3H4 Fab and CD94/NKG2A docking sites on HLA-E. The HLA-E

complex and 3H4 Fab are color-coded according to A and B. CD94: orange; NKG2A: marine

blue.

e, Aerial view of the HLA-E-VL9 peptide binding groove surface. Non-interfacing HLA-E

residues: light grey; non-interfacing peptide residues: lime green; VL9 peptide residues involved

in the 3H4 interface: marine blue. Interfacing HLA-E residues that contact 3H4 heavy and light

chain: orange and teal, respectively; interfacing HLA-E residues that contact both 3H4 heavy

and light chains: violet. Residue positions are numbered on the HLA-E surface view.

f, Aerial view of the overlapping 3H4 and CD94/NKG2A footprints on the HLA-E peptide

binding groove. VL9 peptide residues involved in both the 3H4 and CD94/NKG2A interfaces:

marine blue; HLA-E heavy chain residues involved in both interfaces: violet. Peptide and HLA-E

heavy chain residues involved exclusively in the CD94/NKG2A interface: teal and orange,

respectively.

g, Binding interface of 3H4 HC/VL9 peptide. Interfacing residues (Y97, S100, S100A and

Y100B of the VH CDR3 loop and V1, M2, P4 and R5 of the VL9 peptide) are shown in ball and

stick-form with non-interfacing residues in cartoon form. VL9 peptide: lime green; HLA-E heavy

chain: grey; 3H4 heavy chain: light purple.

h-i, Binding interfaces of 3H4 HC/HLA-E heavy chain (h) and 3H4 LC/HLA-E HC (i).

Interfacing residues are displayed in ball-and-stick form, non-interfacing residues are displayed

in cartoon form and hydrogen bonds as dashed lines.

j, Key interfacing residues within the germline-encoded D-junction. 3H4 heavy chain VH

sequence were in purple and the CDR1/2/3 regions shaded grey. Germline-encoded residues

within the VH CDR3 D-junction are denoted. The 4 key interfacing residues (Y97, S100, S100A

and Y100B) within this germline-encoded D-junction that make contacts both the HLA-E heavy

chain and VL9 peptide are highlighted magenta in the sequence and illustrated as magenta

sticks in the PyMol visualization. HLA-E heavy chain: grey; VL9 peptide: green; hydrogen

bonds: magenta dashed lines; residues of the 3H4 heavy chain that are not germline-encoded

key interfacing residues: light purple.

Figure. 3 MAb 3H4 enhanced the cytotoxicity of the NKG2A+ NK cell line NK-92 against HLA-E-VL9 expressing 293T cells.

a, Schematic illustrating the hypothesis. Blockade of the inhibitory NKG2A/CD94/HLA-E pathway with anti-HLA-E-VL9 antibody (3H4) and/or anti-NKG2A antibody (Z199) could enhance target cell lysis by NK cells.

b-c, NK cell cytotoxicity against 3H4 IgM-treated target cells as assessed by ^{51}Cr release assay. Antibody was incubated with HLA-E-VL9 transfected 293T cells (b) and untransfected 293T cells (c) at final concentration of 10 µg/ml or 3 µg/ml, and NK92 cells were added into the

mixture as effector cells at effector: target (E:T) ratios of 20:1 and 6:1. Mouse IgM MM-30 was

used as an isotype controls. Dots represent the mean values of triplicate wells in eight

independent experiments. Statistical analysis was performed using mixed effects models.

Asterisks show the statistical significance between indicated groups: ns, not significant,

*P<0.05, **P<0.01, ***P<0.001, ****P<0.0001.

d-e, NK cell cytotoxicity in the presence of anti-NKG2A mouse IgG Z199 in combination

with TE4 control- or 3H4- treated target cells as assessed by ^{51}Cr release assay. Antibody

combinations of Z199 + IgM control (d) or Z199 + 3H4 (e) were incubated with HLA-E-VL9

transfected 293T cells and untransfected 293T cells at a final concentration of 10 μg/ml, and

NK92 cells were added into the mixture as effector cells. Dots represent the mean values of

triplicate wells in three independent experiments.

d

3H4 GWT	Y97	Y98	G99	S100
3H4 Gv2	T	A	R	Y
3H4 Gv3	S	A	Q	Y
3H4 Gv4	T	G	R	F
3H4 Gv5	G	A	Q	T
3H4 Gv6	G	A	Q	W
3H4 Gv7	G	A	R	W
3H4 Gv8	A	A	Q	H
3H4 Gv9	A	A	R	F
3H4 Gv10	G	A	R	H
3H4 Gv11	A	A	T	Y
3H4 Gv12	R	G	P	P

Figure. 4 Affinity maturation of HLA-E-VL9-specific antibody 3H4 on human IgG1

backbone.

a, Schematic illustration of the affinity maturation strategy. Libraries of 3H4 mAb variants

were transformed into *S. cerevisiae* and displayed on the surface of yeast cells as single-chain

fragment variable (scFv). APC-conjugated HLA-E-VL9 tetramers were used for FACS sorting.

b, Sites at the 3H4/HLA-E-VL9 interface where sequence optimization by library

screening provideed the most significant affinity gains. 3H4: *purple*; HLA-E: *green*; VL9

peptide: *orange*.

c, Enrichement of HLA-E-VL9+ library clones after three rounds of selection by

fluorescence-activated cell sorting (FACS). The yeast cells containing the scFv libraries were

sorted sequentially for binding to decreasing concentrations of fluorescently labeled HLA-E-VL-9

(50 µg/ml, *top*; 10 µg/ml, *middle*; or 0.6 µg/ml *bottom*).

d, Mutations at positions 97-100 in the eleven 3H4 variants chosen for additional

characterization upon library screening.

e, Binding of 3H4 Gwt and optimized variants to HLA-E-VL9 or HLA-E-Mtb44 transfected

293T cells. Representative flow cytometry data from one of three independent experiments are

shown.

f, SPR sensorgrams showing binding kinetics of 3H4 Gwt and optimized variants. Rate

constants (k_a, k_d) and dissociation constant K_D were determined by curve fitting analysis of SPR

data with a 1:1 binding model. Binding data are shown as colored lines, and the best fits of a 1:1

binding model are shown as black lines. Representative data from one of two independent

experiments are shown.

g, Enhanced NK-92 cell cytotoxicity by optimized IgG 3H4 Gv3 and 3H4 Gv6 on HLA-E-

VL9 transfected 293T cells and untransfected 293T cells, in compare with IgG 3H4 Gwt.

Dots represent the mean values of triplicate wells in four or five independent [51]Cr release

assays.

Statistical analysis was performed using mixed effects models. Asterisks show the statistical

significance between indicated groups: ns, not significant, *P<0.05, **P<0.01, ***P<0.001,

****P<0.0001.

Figure. 5 HLA-E-VL9-specific antibodies isolated from the B cell pool of healthy humans.

a, Scheme of isolating HLA-E-VL9-specific antibodies from healthy humans. Pan-B cells

were first isolated by negative selection from human leukapheresis PBMCs. A three-color

sorting strategy was used to sort single B cells that were positive for HLA-E-VL9 and negative

for HLA-E-RL9HIV or HLA-E-RL9SIV. Flow cytometry data showing the sorting of HLA-E-VL9

double positive, HLA-E-RL9HIV negative, HLA-E-RL9SIV negative B cells in PBMCs from four

donors (LP021, LP030, LP059 and LP060) are shown. Variable regions of antibody heavy and

light chain genes were isolated from the sorted B cells by PCR, and cloned into an expression

backbone with a human IgG1 constant region. Antibodies were produced by transient

transfection in 293i cells, and antibody binding specificities were analyzed by surface staining of

transfected 293T cells and high throughput screening (HTS) flow cytometry.

b, Binding specificities of the HLA-E-VL9-specific antibodies (n=56) from four donors

shown as a heatmap. The compensated MFIs of HLA-E-VL9-specific antibody staining on

HLA-E-VL9, HLA-E-RL9HIV, or HLA-E-RL9SIV transfected 293T cells at a concentration of 1

µg/ml were shown. Representative data from one of two independent experiments are shown.

c, NK cell cytotoxicity against CA147 IgG-treated target cells as assessed by [51]Cr release

assay. Human antibody CA147 was incubated with HLA-E-VL9 transfected 293T cells and

untransfected 293T cells at final concentration of 10 µg/ml or 3 µg/ml, and NK92 cells were

added into the mixture as effector cells at effector: target (E:T) ratio of 20:1 and 6:1. Human

antibody A32 was used as the isotype control. Dots represent the mean values of triplicate wells

in five independent experiments. Statistical analysis was performed using mixed effects models.

Asterisks show the statistical significance between indicated groups: ns, not significant,

*P<0.05, **P<0.01, ***P<0.001, ****P<0.0001.

d, Percentage of HLA-E-VL9-specific B cells in CD19[+] pan-B cells in four donors.

e, Phenotypes of HLA-E-VL9-specific B cells (n=56) shown as heatmap. Expression of

markers in each single B cell were determined from index sorting data and are shown as MFIs

after compensation. Compensated MFIs below zero were set as zero. Each row indicates one

single cell. The rows were clustered by K-means Clustering in R. Four subsets were observed:

CD10[-]CD27[-]CD38[+/-] naïve B cells, CD10[+]CD27[-]CD38[++] transitional B cells, CD10[-]CD27[+]CD38[-]

non-class-switched memory B cells, and CD10⁻CD27⁺CD38⁺ plasmablast cells. Detailed

information for each single cell and antibody is shown in *Table S4*.

f-g, Antibody gene usages. f, Heavy chain viable (V_H) region gene usage shown as a bar chart

(left) and pie chart (right). The top five V_H genes found in HLA-E-VL9-specific antibodies are

colored in the pie charts. **g,** Kappa chain variable (V_κ) and lambda chain variable (V_λ) region

gene usage shown as a bar chart (left) and pie chart (right). The top five V_κ/V_λ genes found in

HLA-E-VL9-specific antibodies are colored in the pie charts. Reference VH-VL repertoires

(n=198,148) from three healthy human donors from a previous study (*DeKosky, Nat Med 2015*)

were used as a control. The chi-square test of independence was performed to test for an

association between indicated gene usage and repertoire/antibody type in panels A-B. ****,

$p<0.0001$; *, $0.01<p<0.05$.

h-i, Comparison of heavy chain (h) and light chain (i) CDR3 (CDR-H3) length. HLA-E-VL9

antibody CDR-H3 length was compared with the reference (DeKosky et al., 2015) human

antibody CDR-H3 length.

j-k, Violin plots showing the mutation rates of heavy chains (j) and light chains (k). HLA-

E-VL9 antibody sequences (E-VL9) were compared with reference sequences from naïve and

antigen-experienced (Ag-Exp) antibody repertoires (n=13,780 and 34,692, respectively).

1   **Supplementary Materials for**

Murine and Human Antibodies that Bind HLA-E-Leader Peptide Complexes and

Enhance NK Cell Cytotoxicity

Dapeng Li[1,5,8,9,*], Simon Brackenridge[2,8], Lucy C. Walters[2,8], Olivia Swanson[1], Karl Harlos[3],

Daniel Rozbesky[3,4], Derek W. Cain[1,5], Kevin Wiehe[1,5], Richard M. Scearce[1], Maggie Barr[1],

Zekun Mu[1], Robert Parks[1], Max Quastel[2], Robert J. Edwards[1,5], Yunfei Wang[1,5], Wes

Rountree[1,5], Kevin O. Saunders[1,6,7], Guido Ferrari[7], Persephone Borrow[2], E. Yvonne Jones[3],

S. Munir Alam[1,5], Mihai L. Azoitei[1,5,*], Geraldine M. Gillespie[2,*], Andrew J. McMichael[2,*],

Barton F. Haynes[1,6,*]

[1]Duke Human Vaccine Institute, Duke University School of Medicine, Durham, NC 27710,

USA

[2]Nuffield Department of Clinical Medicine, University of Oxford, Oxford, OX3 7FZ, UK

[3]Division of Structural Biology, Wellcome Centre for Human Genetics, University of Oxford,

Oxford, OX3 7BN, UK

[4]Department of Cell Biology, Charles University, Prague, 12800, Czech Republic

[5]Department of Medicine, Duke University School of Medicine, Durham, NC 27710, USA

[6]Department of Immunology, Duke University School of Medicine, Durham, NC 27710, USA

[7]Department of Surgery, Duke University School of Medicine, Durham, NC 27710, USA

[8]Authors contributed equally

[9]Lead contact

*Address correspondence to dapeng.li@duke.edu, mihai.azoitei@duke.edu,

geraldine.gillespie@ndm.ox.ac.uk, andrew.mcmichael@ndm.ox.ac.uk and

barton.haynes@duke.edu

This PDF file includes:

Materials and Methods

References

Supplementary Figure. S1 to S4

Other Supplementary Material for this manuscript includes the following:

Supplementary Table S1-S4

MATERIALS AND METHODS

Cell Lines

K562-E cells (K562 cells stably expressing HLA-E) and K562-E/UL49.5 cells (with a

TAP-inhibitor UL49.5) are kindly provided by Dr. Thorbald van Hall from Leiden University

(Lampen et al., 2013). All the other cells used in this study are from ATCC. 293T cells

(ATCC CRL-3216) were maintained in Dulbecco's Modified Eagle's Medium (DMEM; Gibco,

Catalog# 10564) supplemented with 10% fetal bovine serum (FBS; Gibco, Catalog#

10099141) and 1% penicillin/streptomycin (Gibco, Catalog# 10378016). K562 cells (ATCC

CCL-243), K562-E cells and K562-E/UL49.5 cells were cultured in Iscove's Modified

Dulbecco's Medium (IMDM; Hyclone, Catalog# SH30228.01) supplemented with 10% FBS.

Jurkat, DU-4475 and U-937 cells were cultured in RPMI-1640 medium (Gibco, Catalog#

72400) supplemented with 10% FBS. SiHa cells were cultured in Minimum Essential

Medium (MEM; Gibco, Catalog# 11095080) supplemented with 10% FBS. The NK-92

human cell line (ATCC CRL-2407) was cultured in Alpha Minimum Essential medium (α-

MEM; Gibco, Catalog# 12561072) supplemented with 2 mM L-glutamine, 0.2 mM inositol,

0.1 mM 2-mercaptoethanol, 0.02 mM folic acid, 100 U/ml recombinant IL-2 (Biolegend,

Catalog# 589108), 12.5% horse serum (Gibco, Catalog# 16050122) and 12.5% FBS. All the

cells were maintained at 37°C, 5% CO_2 in humidified incubators.

Animals

Transgenic mice carrying human β2-microglobulin (β2m) and HLA-B*27:05 genes were

obtained from Jackson lab (B6.Cg-Tg(B2M,HLA-B*27:05)56-3Trg/DcrJ; stock# 003428).

Hemizygous mice were used in this experiment, as this strain is homozygous lethal. For

hemizygous mice genotyping, peripheral blood lymphocytes (PBLs) were isolated and

stained using mouse CD45 antibody (Biolegend, Catalog# 103122), human HLA class I

antibody (Biolegend, Catalog# 311406) and human β2m antibody (Biolegend, Catalog#

316312). All animal experiments were conducted with approved protocols from the Duke

University Institutional Animal Care and Use Committee.

Human Subjects

Human leukapheresis frozen vials were collected by the External Quality Assurance

Program Oversight Laboratory (EQAPOL) (Sanchez et al., 2014a; Sanchez et al., 2014b).

Samples from four male donors were used in this study. *Table S5* shows the clinical

characteristics of the individuals studied. All experiments that related to human subjects was

carried out with the informed consent of trial participants and in compliance with Institutional

Review Board protocols approved by Duke University Medical Center.

Peptide synthesis

The VL9 peptide (VMAPRTVLL) was synthesized to >85% purity via Fmoc (9-

fluorenylmethoxy carbonyl) chemistry by Genscript USA and reconstituted to 200mM in

DMSO.

HLA-E-peptide protein refolding and purification

β2-microglobulin, previously purified from inclusion bodies in a Urea-MES buffer, was

added to a refolding buffer to achieve a final concentration of 2 μM. The refold buffer

comprised 100 mM Tris pH8.0, 400 mM L-arginine monohydrochloride, 2 mM EDTA, 5 mM

reduced glutathione and 0.5 mM oxidized Glutathione and was prepared in MiliQ water. A 20

μM concentration of VL9 peptide (VMAPRTVLL), previously reconstituted to 200 mM in

DMSO, was added to the refolding buffer followed by HLA-E*0103 heavy chain, which was

pulsed into the refold to a final concentration of 1 µM. Once the refold had incubated for 72

hours at 4 °C it was filtered through a 1.0 µm cellular nitrate membrane and concentrated in

the VivaFlow 50R and VivaSpin Turbo Ultrafiltration centrifugal systems with 10 kDa

molecular weight cut-offs. The concentrated samples were injected onto a Superdex S75

16/60 column and refolded protein eluted according to size into phosphate buffered saline

(PBS). Eluted protein complexes were validated by non-reducing SDS-PAGE

electrophoresis on NuPAGE 12% Bis-Tris protein gels and further concentrated via VivaSpin

Turbo Ultrafiltration centrifugal device to 1.1 mg/mL.

HLA-E-peptide biotinylation and tetramer generation

HLA-E-peptide samples requiring biotinylation were subsequently buffered exchanged

on Sephadex G-25 PD10 columns (GE Healthcare, UK) into 10mM Tris buffer using

commercially available BirA enzyme (Avidity, USA) following the manufacturer's

instructions. Following overnight biotinylation, protein samples were subsequently

purified into 20mM Tris pH8,100mM NaCl buffer or PBS on a HiLoad 16/600 Superdex 75pg

column using an AKTA size exclusion fast protein liquid chromatography (FPLC) system.

Correctly folded β2m-HLA-E*01:03-peptide complexes were subsequently concentrated to 2

mg/mL and snap frozen.

HLA-E*01:03 tetramers were generated via conjugation to various fluorescent labels

including Extravidin-PE (Sigma), Streptavidin-bound APC (Biolegend, San Diego) or BV421

(Biolegend, San Diego) at a Molar ratio of 4:1 as previously described (Braud et al., 1998).

Immunization in HLA-B27/β2m Transgenic Mice

HLA-B27/β2m transgenic mice (*n*=23) were intramuscularly (i.m.) immunized with

pooled HLA-E-RL9HIV complex (12.5 µg/animal) and HLA-E-RL9SIV complex (12.5

µg/animal) adjuvanted with STR8S-C (Moody et al., 2014) at weeks 0, 2, 4, 6, 12 and 16.

MAb 3H4 was isolated from this study. In another experiment, HLA-B27/β2m transgenic

mice (n = 10) were i.p. immunized with either HLA-E-RL9HIV single chain trimer (SCT)

transfected 293T cells ($2x10^6$ cells/animal) or HLA-E-RL9SIV SCT transfected 293T cells

($2x10^6$ cells/animal) at weeks 0, 2, 4, 6, 17 and 19. MAb 13F11 was isolated from this study.

In the third experiment, HLA-B27/β2m transgenic mice (n=10) were i.m. immunized with

HLA-E-VL9 complex (25 μg/animal) adjuvanted with STR8S-C at Week 0, 2 and 4, following

by intraperitoneally (i.p.) immunization with HLA-E-VL9 SCT transfected 293T cells ($2x10^6$

cells/animal) at Week 14, 16 and 18. MAb 10C10 and 2D6 were isolated from this study.

Serum titers were monitored by ELISA Mice with high binding antibody titers were selected

for the subsequent spleen cell fusion and B cell sorting experiments.

Hybridoma Cell Line Generation and Monoclonal Antibody Production

Mice were boosted with the indicated priming antigen 3 days prior to fusion. Spleen cells

were harvested and fused with NS0 murine myeloma cells using PEG1500 to generate

hybridomas. After 2 weeks, supernatant of hybridoma clones were collected and screened

by flow cytometry-based high throughput screening (HTS). Specifically, we tested for

antibodies differentially binding 293T cells transiently transfected with plasmid DNA

expressing single chain peptide-HLA-E-ß2m trimers so that they expressed HLA-E-RL9HIV,

HLA-E-RL9SIV or HLA-E-VL9 at the cell surface. Hybridomas cells that secreted HLA-E-VL9

antibodies were cloned by limiting dilution for at least 5 rounds until the phenotypes of all

limiting dilution wells are identical. IgG mAbs were purified by protein G affinity

chromatography, while IgM mAbs were purified by ammonium sulfate precipitation and by

Superose 6 column size-exclusion chromatography in AKTA Fast Protein Liquid

Chromatography (FPLC) system. The VH and VL sequences of mAbs were amplified from

hybridoma cell RNA using primers reported previously (Tian et al., 2016; von Boehmer et al.,

2016).

Cell Surface Staining and High-Throughput Screening (HTS)

HLA-E SCT constructs encoding HLA-E-VL9, HLA-E-RL9HIV, or HLA-E-RL9SIV were

transfected into 293T cells using GeneJuice transfection reagent (Novagen, Catalog#

70967). For epitope mapping experiment, a panel of HLA-E-VL9 SCT constructs with single

amino acid mutations were transfected into 293T cells using the same method. Cells were

dissociated with 0.1% EDTA at 48 hours post-transfection and stained with a Fixable Near-

IR Dead Cell Stain Kit (Thermo Fisher, Catalog# L34976). After washing, primary antibodies

(supernatant from hybridoma cells, supernatant from transfected cells, or purified antibodies)

were added and incubated with cells for 1 hour at 4°C, following by staining with 1:1000

diluted secondary antibodies for 30 mins at 4°C. For mouse primary antibodies, we used

Alexa Fluor 555 (AF555) conjugated goat anti-mouse IgG (H+L) (Thermo Fisher, Catalog#

A32727) or Alexa Fluor 647 (AF647) conjugated goat anti-mouse IgG (H+L) (Thermo Fisher,

Catalog# A32728) as secondary antibodies; for human primary antibodies, we used AF555

conjugated goat anti-human IgG (H+L) (Thermo Fisher, Catalog# A-21433) or AF647

conjugated goat anti-human IgG (H+L) (Thermo Fisher, Catalog# A-21445) as secondary

antibodies. Cells were then washed 3 times and resuspended in fixation buffer (1%

formaldehyde in PBS, pH7.4). Data were acquired on a BD LSR II flow cytometer and

analyzed using FlowJo version 10.

3H4 Fab production

A humanized version of the 3H4 antibody (3H4-huIgG1) was digested to produce Fab

fragments using the Pierce Fab Preparation kit (ThermoFisher SCIENTIFIC). 3H4 Fab-

retrieved sample was further purified by size exclusion on a Superdex S75 16/60 column

and eluted into PBS buffer. Following concentration to 1.1mg/mL and SDS-PAGE gel-based

validation, 3H4 Fab purified material was incubated for 1 hours on ice with freshly purified

HLA-E-VL9. The combined 3H4:Fab-HLA-E-VL9 sample was concentrated to 7.5mg/mL

prior to crystallographic set-up.

Crystallization screening

Crystals were grown via sitting drop vapour-diffusion at 20 °C in a 200nL drop with a 1:1

protein to reservoir ratio (Walter et al., 2005). The 3H4 Fab-HLA-E(VL9) co-complex

crystallized in 20% PEG 8000, 0.1 M Na HEPES at pH 7, in the ProPlex sparse matrix

screen. Crystals were cryo-preserved in 25% glycerol and diffraction data were collected at

the I03 beamline of Diamond Light Source.

Crystallographic analysis

Two copies of the co-complex structure of 3H4 Fab bound to HLA-E-VL9 were present

in the asymmetric unit, a single copy constituted the focus of further discussion since root-

mean-square deviation (RMSD) calculations from Cα-atom pairwise alignment of the two

copies indicated minimal repositioning of interfacing residues at the HLA-E-3H4 binding site

(*Table S2B-F*). Additionally, pairwise alignment with the previously published non-receptor-

bound HLA-E coordinates (PDB ID: 1MHE) (O'Callaghan et al., 1998) revealed minimal

structural changes in HLA-E upon 3H4 engagement (*Table S2C*).

Diffraction data were merged and indexed in xia2 dials (Winter et al., 2018). Outer shell

reflections were excluded from further analysis to ensure the $CC_{1/2}$ value exceeded the

minimum threshold (>0.5) in each shell (Karplus and Diederichs, 2012). Sequential

molecular replacement was carried out in MolRep of the CCP4i suite using molecule one of

the previously published Mtb44-bound HLA-E structure with the peptide coordinates deleted

(PDB ID: 6GH4) and one molecule of the previously published anti-APP-tag Fab structure

(PDB ID: 6HGU) as phasing models (Vagin and Teplyakov, 2010; Winn et al., 2011). Rigid

body and retrained refinement were subsequently carried out by Phenix.refine (Afonine et

al., 2012) in between manual model building in Coot (Emsley et al., 2010). Model geometry

was validated by MolProbity (Chen et al., 2010) and structural interpretation was conducted

using the PyMOL Molecular Graphics System, version 2.0 (Schrödinger, LLC) in addition to

the PDBePISA (Krissinel and Henrick, 2007) and PDBeFOLD (Krissinel and Henrick, 2004)

servers.

Antigen-Specific Single B Cell Sorting

HLA-E-VL9-specific human B cells were sorted in flow cytometry using a three-color

sorting technique. Briefly, the stabilized HLA-E-β2M-peptide complexes were made as

tetramers and conjugated with different fluorophores. Human pan-B cells, including naïve

and memory B cells, were isolated from PBMCs of healthy donors using human pan-B cell

enrichment kit (STEMCELL, Catalog# 19554). The isolated pan-B cells were then stained

with IgM PerCp-Cy5.5 (Clone# G20-127, BD Biosciences, Catalog# 561285), IgD FITC

(Clone# IA6-2, BD Biosciences, Catalog# 555778), CD3 PE-Cy5 (Clone# HIT3a, BD

Biosciences, Catalog# 555341), CD235a PE-Cy5 (Clone# GA-R2, BD Biosciences,

Catalog# 559944), CD10 PE-CF594 (Clone# HI10A, BD Biosciences, Catalog# 562396),

CD27 PE-Cy7 (Clone# O323, eBioscience, Catalog# 25-0279), CD16 BV570 (Clone# 3G8,

Biolegend, Catalog# 302035), CD14 BV605 (Clone# M5E2, Biolegend, Catalog# 301834),

CD38 APC-AF700 (Clone# LS198-4-2, Beckman Coulter, Catalog# B23489), CD19 APC-

Cy7 (Clone# LJ25C1, BD Biosciences, Catalog# 561743) and tetramers at 2 µg/million cells

(including BV421-conjugated HLA-E-VL9 tetramer, PE-conjugated HLA-E-VL9 tetramer,

APC-conjugated HLA-E-RL9SIV tetramer and APC-conjugated HLA-E-RL9HIV tetramer).

The cells were then stained with a Fixable Aqua Dead Cell Stain Kit (Invitrogen, Catalog#

L34957). HLA-E-VL9-specific B cells were sorted in BD FACSAria II flow cytometer (BD

Biosciences) for viable CD3^{neg}/ CD14^{neg} /CD16^{neg} /CD235a^{neg}/CD19^{pos} / HLA-E-VL9$^{double-pos}$/

HLA-E-RL9HIV^{neg}/HLA-E-RL9SIV^{neg} subset as single cells in 96-well plates.

PCR Amplification of Human Antibody Genes

The $V_HD_HJ_H$ and V_LJ_L genes were amplified by RT-PCR from the flow cytometry-sorted

single B cells using the methods as described previously (Liao et al., 2009; Wrammert et al.,

2008) with modification. Primer details were listed in Tables S2. The PCR-amplified genes

were then purified and sequenced with 10 µM forward and reverse primers. Sequences were

analyzed by using the human library in Clonalyst for the VDJ arrangements of the

immunoglobulin *IGHV*, *IGKV*, and *IGLV* sequences and mutation frequencies (Kepler et al.,

2014). Clonal relatedness of $V_H D_H J_H$ and $V_L J_L$ sequences was determined as previously

described (Liao et al., 2013).

Expression of $V_H D_H J_H$ and $V_L J_L$ as Full-Length IgG Recombinant mAbs

Transient transfection of recombinant mAbs was performed as previously described

(Liao et al., 2009). Briefly, purified PCR products were used for overlapping PCR to generate

linear human antibody expression cassettes. The expression cassettes were transfected into

293i cells using ExpiFectamine (Thermo Fisher Scientific, Catalog# A14525). The

supernatant samples containing recombinant antibodies were used for cell surface staining

and HTS assay to measure the binding reactivities.

The selected human antibody genes were then synthesized and cloned (GenScript) in a

human IgG1 backbone with 4A mutations (Saunders, 2019). Recombinant IgG mAbs were

then produced in HEK293i suspension cells by transfection with ExpiFectamine and purified

using Protein A resin. The purified mAbs were run in SDS-PAGE for Coomassie blue

staining and western blot. Antibodies with aggregation were further purified in AKTA FPLC

system using a Superdex 200 size-exclusion column.

Surface Plasmon Resonance (SPR)

Surface plasmon resonance assays were performed on a BIAcore 3000 instrument, and

data analysis was performed with BIAevaluation 3.0 software as previously described (Liao

et al., 2006). Purified mAbs flowed over CM5 sensor chips at concentrations of 100 µg/ml,

and antibody binding was monitored in real-time at 25°C with a continuous flow of PBS at 30

µl/min. For SPR affinity measurements, antibody binding to HLA-E-VL9 complex protein was

performed using a BIAcore S200 instrument (Cytiva, formerly GE Healthcare, DHVI BIA

Core Facility, Durham, NC) in HBS-EP+ 1x running buffer. The antibodies were first

captured onto CM5 sensor chip to a level of ~9000 RU. The HLA-E-VL9 soluble proteins

were injected over the captured antibodies at a flow rate of 30uL/min. After dissociation, the

antibodies were regenerated using a 30 second pulse of Glycine pH2.0. Results were

analyzed using the Biacore S200 Evaluation software (Cytiva). Subsequent curve fitting

analyses were performed using a 1:1 Langmuir model with a local Rmax. The reported

binding curves are representative of two data sets.

ELISA

Direct binding ELISAs were conducted in 384-well ELISA plates coated with 2 µg/ml of

C-trap-stabilized HLA-E-VL9, C-trap-stabilized HLA-E-RL9HIV or C-trap-stabilized HLA-E-

RL9SIV in 0.1 M sodium bicarbonate overnight at 4°C. Plates were washed with PBS +

0.05% Tween 20 and blocked with 3% BSA in PBS at room temperature for 1 h. MAb

samples were incubated for 1 h in 3-fold serial dilutions starting at 100 µg/ml, followed by

washing with PBS-0.05% Tween 20. HRP-conjugated goat anti-human IgG secondary Ab

(SouthernBiotech, catalog# 2040-05) was diluted to 1: 10,000 in 1% BSA in PBS-0.05%

Tween 20 and incubated at room temperature for 1 h. For sandwich ELISA, 384-well ELISA

plates were coated with HLA-E-VL9 antibodies in a 3-fold dilution starting from 100 µg/mL in

0.1 M sodium bicarbonate overnight at 4°C. Plates were washed with PBS + 0.05% Tween

20 and blocked with 3% BSA in PBS at room temperature for 1 h. C-trap-stabilized HLA-E-

VL9, C-trap-stabilized HLA-E-RL9HIV, C-trap-stabilized HLA-E-RL9SIV, or diluent control

were then added at 2 µg/mL and incubated at room temperature for 1 h. After washing,

HRP-conjugated anti-human β2M antibody (Biolegend, catalog# 280303) were added at 0.2

µg/mL and incubated at room temperature for 1 h. These plates were washed for 4 times

and developed with tetramethylbenzidine substrate (SureBlue Reserve). The reaction was

stopped with 1 M HCl, and optical density at 450 nm (OD_{450}) was determined.

Antibody Poly-Reactivity Assays

All mAbs isolated from mice and human were tested for ELISA binding to nine

autoantigens - Sjogren's syndrome antigen A (SSA), Sjogren's syndrome antigen (SSB),

Smith antigen (Sm), ribonucleoprotein (RNP), scleroderma 70 (Scl-70), Jo-1 antigen,

double-stranded DNA (dsDNA), centromere B (Cent B), and histone as previously described

(Han et al., 2017; Liao et al., 2011). Indirect immunofluorescence assay of mAbs binding to

HEp-2 cells (Inverness Medical Professional Diagnostics, Princeton, NJ) was performed as

previously described (Haynes et al., 2005; Liao et al., 2011). MAbs 2F5 (Yang et al., 2013)

and 17B (Moore and Sodroski, 1996) were used as positive and negative controls,

respectively. All antibodies were screened in two independent experiments.

Negative Stain Electron Microscopy of IgM antibodies

FPLC purified IgM antibodies were diluted to 0.08 mg/ml in HEPES-buffered saline (pH

7.4) + 5% glycerol, and stained with 2% uranyl formate. Images were obtained with a Philips

EM420 electron microscope at 82,000 magnification and processed in Relion 3.0.

Peptide-Pulsing in K562-E Cells

K562-E cells and K562-E/UL49.5 cells were resuspended with fresh IMDM media with

10% FBS at $2x10^6$ cells/ml. Peptides were added into cell suspension at a final

concentration of 100 μM. The cell/peptide mixtures were incubated at 26°C with 5% CO_2 for

20-22 hours and were transferred to 37°C for 2 hours with 5% CO_2 before use. In the

following mAb staining experiment, medium with 100 μM peptides was used to maintain

peptide concentration.

NK Cell Cytotoxicity Assay

NK Cell Cytotoxicity was measured by ^{51}Cr release assay. A NKG2A-positive,

CD16/CD32/CD64-negative NK-92 cells were used as effector cells in our study.

Transfected or untransfected 293T cells were used as target cells. Target cells were

counted, washed, resuspended in R10 at $1×10^7$ cell/ml, and labeled with $Na_2{}^{51}CrO_4$ at 250

μCi/ml for 2 hours at 37°C. After washing three times using R10, cells were mixed with the

testing antibody and effector cells in a final effector to target (E:T) ratio of 20:1 and 6:1 in

triplicate wells in a flexible 96 well round bottom plates (PerkinElmer, Catalog# 1450-401).

The plates were inserted in flexible 96-well plate cassettes (PerkinElmer, Catalog# 1450-

101), sealed and incubated at 37°C for 4 hours. After the incubation, cells were pelleted by

centrifugation, and from the top of the well, add 25 ul of supernatant to a rigid 96 well

isoplates (PerkinElmer, Catalog#1450-514) containing 150 ul of Ultima Gold LSC Cocktail

(Sigma, Catalog# L8286). The plates were inserted in rigid 96-well plate cassettes

(PerkinElmer, Catalog# 1450-105), sealed and counted on Perkin Elmer Microbeta Triux

1450 counter. ^{51}Cr labeled target cells without effector cells were set as a spontaneous

release control, and ^{51}Cr labeled target cells mixed with detergent (2% Triton X-100) were

used as a maximum release control. The percentages of specific lysis were calculated with

the formulation: The Percentages of Specific Lysis (^{51}Cr Release %) = [(Experimental

Release – Spontaneous Release)/ (Maximum Release – Spontaneous Release)] x 100.

Development and screening of scFv libraries on the surface of yeast

A library was built that contained ~1.1 million 3H4 scFv variants with amino acid

diversity at sites that were determined by structural analysis to interact with HLA-E-VL9.

Seventeen residues (*Figure. S3*) located in the CDR loops of 3H4 were randomized in

groups of four based on their proximity and all the possible combinations of amino acids

were sampled at these sites. Library DNA was synthesized on a BioXP 3250 (Codex)

system and amplified with High Fidelity Phusion polymerase (New England Biolabs). PCR

products were gel extracted (Qiagen Gel Extraction kit) to select full length genes as per the

manufacturer's protocol. 3H4 scFv variants were displayed in library format on the surface of

yeast as previously described (Benatuil et al., 2010; Chao et al., 2006). Briefly, *S. cerevisiae*

EBY100 cells were transformed by electroporation with a 3:1 ratio of 12 μg scFv library DNA

and 4 μg pCTcon2 plasmid digested with BamHI, SalI, NheI (New England Biolabs). The

size of the transformed library, determined by serial dilution on selective plates, was 5×10^{7}

individual colonies. Yeast Libraries were grown in SDCAA media (Teknova) supplemented

with pen-strep at 30°C and 225 rpm. 80% of the sequences recovered from the transformed

libraries were confirmed to contain full length, in-frame genes by Sanger sequencing

(Genewiz). scFv expression on the surface of yeast was induced by culturing the libraries in

SGCAA (Teknova) media at a density of 1×10^7 cells/mL for 24-36 hours. Cells were washed

twice in ice cold PBSA (0.01M sodium phosphate, pH 7.4, 0.137M sodium chloride, 1g/L

bovine serum albumin) and labeled with APC conjugated HLA-E-VL9 tetramer and 1:100

anti-c-myc:FITC (ICL) and incubated for 1 hour at 4C. Initial selection was conducted with

50mg/mL labeling concentration of HLA-E-VL9 tetramer; the second round of selection was

done at 0.6mg/mL tetramer. Cells were washed twice with PBSA after incubation with the

fluorescently labeled probes and sorted on a BD FACS-DiVa. Double positive cells for APC

and FITC were collected and expanded in SDCAA media supplemented with pen-strep

before successive rounds of enrichment. FACS data was analyzed with Flowjo_v10.7

software (Becton, Dickinson & Company). All clones selected by FACS were expanded, and

their DNA was extracted (Zymo Research) for analysis by Sanger sequencing (Genewiz).

scFv encoding plasmids were recovered from yeast cultures by yeast miniprep with the

Zymoprep yeast plasmid miniprep II kit (Zymo Research). Isolated DNA was transformed

into NEB5α strain of *E. coli* (New England Biolabs) and the DNA of individual bacterial

colonies was isolated (Wizard Plus SV Minipreps, Promega) and analyzed by Sanger

sequencing (Genewiz).

Statistics Analysis

Data were plotted using Prism GraphPad 8.0 or visualized using the ComplexHeatmap

R package. SAS 9.4 (SAS Institute, Cary, NC) was used to perform the statistical analysis

with a p-value < 0.05 considered significant. For ^{51}Cr release assays, mixed effects models

were used to make comparisons of antibody to control using a random intercept for the

triplicates run within each experiment and fixed effects of E:T ratio, type (antibody or

control), and the interaction of E:T ratio by type. For human antibody gene usage

analysis, chi-square test of independence was used to compare differences between

groups.

REFERENCES

Afonine, P.V., Grosse-Kunstleve, R.W., Echols, N., Headd, J.J., Moriarty, N.W., Mustyakimov,

M., Terwilliger, T.C., Urzhumtsev, A., Zwart, P.H., and Adams, P.D. (2012). Towards

automated crystallographic structure refinement with phenix.refine. Acta Crystallogr D Biol

Crystallogr *68*, 352-367.

Benatuil, L., Perez, J.M., Belk, J., and Hsieh, C.M. (2010). An improved yeast transformation

method for the generation of very large human antibody libraries. Protein Eng Des Sel *23*,

155-159.

Braud, V.M., Allan, D.S., O'Callaghan, C.A., Soderstrom, K., D'Andrea, A., Ogg, G.S., Lazetic,

S., Young, N.T., Bell, J.I., Phillips, J.H.*, et al.* (1998). HLA-E binds to natural killer cell receptors

CD94/NKG2A, B and C. Nature *391*, 795-799.

Chao, G., Lau, W.L., Hackel, B.J., Sazinsky, S.L., Lippow, S.M., and Wittrup, K.D. (2006).

Isolating and engineering human antibodies using yeast surface display. Nature Protocols *1*,

755-768.

Chen, V.B., Arendall, W.B., 3rd, Headd, J.J., Keedy, D.A., Immormino, R.M., Kapral, G.J.,

Murray, L.W., Richardson, J.S., and Richardson, D.C. (2010). MolProbity: all-atom structure

validation for macromolecular crystallography. Acta Crystallogr D Biol Crystallogr *66*, 12-21.

Emsley, P., Lohkamp, B., Scott, W.G., and Cowtan, K. (2010). Features and development of

Coot. Acta Crystallogr D Biol Crystallogr *66*, 486-501.

Han, Q., Williams, W.B., Saunders, K.O., Seaton, K.E., Wiehe, K.J., Vandergrift, N., Von Holle,

T.A., Trama, A.M., Parks, R.J., Luo, K.*, et al.* (2017). HIV DNA-Adenovirus Multiclade

Envelope Vaccine Induces gp41 Antibody Immunodominance in Rhesus Macaques. J Virol

91.

Haynes, B.F., Fleming, J., St Clair, E.W., Katinger, H., Stiegler, G., Kunert, R., Robinson, J.,

Scearce, R.M., Plonk, K., Staats, H.F.*, et al.* (2005). Cardiolipin polyspecific autoreactivity in

two broadly neutralizing HIV-1 antibodies. Science *308*, 1906-1908.

Karplus, P.A., and Diederichs, K. (2012). Linking crystallographic model and data quality.

Science *336*, 1030-1033.

Kepler, T.B., Munshaw, S., Wiehe, K., Zhang, R., Yu, J.S., Woods, C.W., Denny, T.N.,

Tomaras, G.D., Alam, S.M., Moody, M.A.*, et al.* (2014). Reconstructing a B-Cell Clonal

Lineage. II. Mutation, Selection, and Affinity Maturation. Front Immunol *5*, 170.

Krissinel, E., and Henrick, K. (2004). Secondary-structure matching (SSM), a new tool for fast

protein structure alignment in three dimensions. Acta Crystallogr D Biol Crystallogr *60*, 2256-

2268.

Krissinel, E., and Henrick, K. (2007). Inference of macromolecular assemblies from crystalline

state. J Mol Biol *372*, 774-797.

Lampen, M.H., Hassan, C., Sluijter, M., Geluk, A., Dijkman, K., Tjon, J.M., de Ru, A.H., van

der Burg, S.H., van Veelen, P.A., and van Hall, T. (2013). Alternative peptide repertoire of

HLA-E reveals a binding motif that is strikingly similar to HLA-A2. Mol Immunol *53*, 126-131.

Liao, H.X., Chen, X., Munshaw, S., Zhang, R., Marshall, D.J., Vandergrift, N., Whitesides, J.F.,

Lu, X., Yu, J.S., Hwang, K.K.*, et al.* (2011). Initial antibodies binding to HIV-1 gp41 in acutely

infected subjects are polyreactive and highly mutated. J Exp Med *208*, 2237-2249.

Liao, H.X., Levesque, M.C., Nagel, A., Dixon, A., Zhang, R., Walter, E., Parks, R., Whitesides,

J., Marshall, D.J., Hwang, K.K.*, et al.* (2009). High-throughput isolation of immunoglobulin

genes from single human B cells and expression as monoclonal antibodies. J Virol Methods

158, 171-179.

Liao, H.X., Lynch, R., Zhou, T., Gao, F., Alam, S.M., Boyd, S.D., Fire, A.Z., Roskin, K.M.,

Schramm, C.A., Zhang, Z.*, et al.* (2013). Co-evolution of a broadly neutralizing HIV-1 antibody

and founder virus. Nature *496*, 469-476.

Liao, H.X., Sutherland, L.L., Xia, S.M., Brock, M.E., Scearce, R.M., Vanleeuwen, S., Alam,

S.M., McAdams, M., Weaver, E.A., Camacho, Z.*, et al.* (2006). A group M consensus envelope

glycoprotein induces antibodies that neutralize subsets of subtype B and C HIV-1 primary

viruses. Virology *353*, 268-282.

Moody, M.A., Santra, S., Vandergrift, N.A., Sutherland, L.L., Gurley, T.C., Drinker, M.S., Allen,

A.A., Xia, S.M., Meyerhoff, R.R., Parks, R.*, et al.* (2014). Toll-like receptor 7/8 (TLR7/8) and

TLR9 agonists cooperate to enhance HIV-1 envelope antibody responses in rhesus macaques.

J Virol *88*, 3329-3339.

Moore, J.P., and Sodroski, J. (1996). Antibody cross-competition analysis of the human

immunodeficiency virus type 1 gp120 exterior envelope glycoprotein. J Virol *70*, 1863-1872.

O'Callaghan, C.A., Tormo, J., Willcox, B.E., Braud, V.M., Jakobsen, B.K., Stuart, D.I.,

McMichael, A.J., Bell, J.I., and Jones, E.Y. (1998). Structural features impose tight peptide

binding specificity in the nonclassical MHC molecule HLA-E. Mol Cell *1*, 531-541.

Sanchez, A.M., DeMarco, C.T., Hora, B., Keinonen, S., Chen, Y., Brinkley, C., Stone, M.,

Tobler, L., Keating, S., Schito, M.*, et al.* (2014a). Development of a contemporary globally

diverse HIV viral panel by the EQAPOL program. J Immunol Methods *409*, 117-130.

Sanchez, A.M., Rountree, W., Berrong, M., Garcia, A., Schuetz, A., Cox, J., Frahm, N., Manak,

M., Sarzotti-Kelsoe, M., D'Souza, M.P.*, et al.* (2014b). The External Quality Assurance

Oversight Laboratory (EQAPOL) proficiency program for IFN-gamma enzyme-linked

immunospot (IFN-gamma ELISpot) assay. J Immunol Methods *409*, 31-43.

Saunders, K.O. (2019). Conceptual Approaches to Modulating Antibody Effector Functions

and Circulation Half-Life. Front Immunol *10*, 1296.

Tian, M., Cheng, C., Chen, X., Duan, H., Cheng, H.L., Dao, M., Sheng, Z., Kimble, M., Wang,

L., Lin, S.*, et al.* (2016). Induction of HIV Neutralizing Antibody Lineages in Mice with Diverse

Precursor Repertoires. Cell *166*, 1471-1484 e1418.

Vagin, A., and Teplyakov, A. (2010). Molecular replacement with MOLREP. Acta Crystallogr

D Biol Crystallogr *66*, 22-25.

von Boehmer, L., Liu, C., Ackerman, S., Gitlin, A.D., Wang, Q., Gazumyan, A., and

Nussenzweig, M.C. (2016). Sequencing and cloning of antigen-specific antibodies from

mouse memory B cells. Nat Protoc *11*, 1908-1923.

Walter, T.S., Diprose, J.M., Mayo, C.J., Siebold, C., Pickford, M.G., Carter, L., Sutton, G.C.,

Berrow, N.S., Brown, J., Berry, I.M.*, et al.* (2005). A procedure for setting up high-throughput

nanolitre crystallization experiments. Crystallization workflow for initial screening, automated

storage, imaging and optimization. Acta Crystallogr D Biol Crystallogr *61*, 651-657.

Winn, M.D., Ballard, C.C., Cowtan, K.D., Dodson, E.J., Emsley, P., Evans, P.R., Keegan, R.M.,

Krissinel, E.B., Leslie, A.G., McCoy, A.*, et al.* (2011). Overview of the CCP4 suite and current

developments. Acta Crystallogr D Biol Crystallogr *67*, 235-242.

Winter, G., Waterman, D.G., Parkhurst, J.M., Brewster, A.S., Gildea, R.J., Gerstel, M.,

Fuentes-Montero, L., Vollmar, M., Michels-Clark, T., Young, I.D.*, et al.* (2018). DIALS:

implementation and evaluation of a new integration package. Acta Crystallogr D Struct Biol

74, 85-97.

Wrammert, J., Smith, K., Miller, J., Langley, W.A., Kokko, K., Larsen, C., Zheng, N.Y., Mays,

I., Garman, L., Helms, C.*, et al.* (2008). Rapid cloning of high-affinity human monoclonal

antibodies against influenza virus. Nature *453*, 667-671.

Yang, G., Holl, T.M., Liu, Y., Li, Y., Lu, X., Nicely, N.I., Kepler, T.B., Alam, S.M., Liao, H.X.,

Cain, D.W.*, et al.* (2013). Identification of autoantigens recognized by the 2F5 and 4E10

broadly neutralizing HIV-1 antibodies. J Exp Med *210*, 241-256.

SUPPLEMENTAL FIGURES

Figure. S1 Isolation and characterization of monoclonal antibody 3H4.

a-b, Expression of human HLA-B27 and β2M in peripheral blood lymphocytes (PBLs)

of the transgenic (TG) mice. HLA-B27/β2M TG mice were used to minimize the induction

of antibodies to HLA class I and β2M. Mouse PBLs from TG mice and littermate control were

isolated and stained with anti-mouse CD45, anti-human HLA class I (A/B/C) and anti-human

β2m antibodies. Representative data (a) and the percentages of human HLA-B27$^+$β2M$^+$ cells

in CD45$^+$ PBLs from TG mice (n=6) and control mice (n=6) (b) were shown.

c, Schematic diagram of the immunization, splenocyte fusion and hybridoma

screening strategy. HLA-B27/β2M TG mice (n=10) were immunized with cell surface-

expressing HLA-E-RL9 peptide (a peptide derived from HIV-1; denoted RL9HIV hereafter)

single-chain trimer (SCT)-transfected 293T cells (indicated by red arrows). After

immunizations, spleen cells were harvested from the selected mouse and the fusion was

performed using NS0 cells to generate hybridoma cells. Supernatants from the hybridoma

cell candidates were screened for differential binding by surface staining on HLA-E-VL9,

HLA-E-RL9HIV or HLA-E-RL9SIV transfected 293T cells. Hybridomas producing antibodies

specific for HLA-E-VL9 but not others were selected for cloning and downstream analysis.

Monoclonal cells were cloned for at least five rounds.

d, Serum antibody binding ELISA. Serum antibodies to HLA-E-VL9, HLA-E-RL9SIV, HLA-

E-RL9HIV complexes were quantified by ELISA and shown as log AUC (area under curve).

Antigens used for immunizations and ELISA assays are all cysteine (C)-trap stabilized. Each

curve represents one animal, and the curve for animal that we used for splenocyte fusion

are shown in red.

e, Affinity of 3H4 binding to soluble HLA-E-VL9 complex. 3H4 as a mouse IgM or as a

recombinant human IgG1 were immobilized on CM5 sensor chips and soluble HLA-E-VL9

complex protein at the indicated concentrations was flowed over antibody immobilized

sensor chips. Binding data are shown as black lines, and the best fits of a 1:1 binding model

are shown as colored lines. Rate constants (k_a, k_d) and dissociation constant K_D were

measured following curve fitting analysis.

f, Purification of 3H4 by FPLC using Superose 6 size exclusion column. The arrowed

peak was collected and analyzed by negative staining.

g, Representative class average images of 3H4 negative stain electron microscopy

(NSEM).

h, Binding of 3H4 expressed in a human backbone G1.4A to a panel of autoantigens

by AtheNA assays. HIV-1 gp41 antibody 4E10 was set as a positive control, and a Flu

antibody Ab82 was used as a negative control. The dotted lines indicate the cutoff values

≥100 luminance units used to denote positivity.

i, Binding of 3H4 in human backbone G1.4A to HEp-2 epithelial cells in indirect

immunofluorescence staining assays. HIV-1 gp41 antibody 2F5 was set as a positive

control, and HIV-1 gp120 antibody 17B was used as a negative control. Antibody staining

concentration was 50 µg/ml, and data were collected at 40x objective for 8 seconds. Data

are representative from one of two independent experiments.

j, 3H4 does not cross-react with mouse Qa-1b-peptide complex. 293T cells were

transfected with HLA-E-VL9 (VMAPRTLLL), HLA-E-AL9 (AMAPRTLLL), mouse Qa-1b-VL9,

or mouse Qa-1b-AL9. Transfected cells were stained with 3H4 antibody or an anti-β2M

control antibody 2M2 followed by AF647 conjugated anti-mouse IgG(H+L) secondary

antibody. Data are representative from one of three independent experiments.

k, 3H4 recognizes peptides with variants in P1. 293T cells were transfected with HLA-E

SCTs with VL9 peptides with single amino acid mutations at P1, then stained with 3H4

antibody or an anti-β2M control antibody 2M2 followed by AF647 conjugated anti-mouse

IgG(H+L) secondary antibody (dark blue). Cells were gated for EGFP positive subsets.

Isotype control stained cells were used as a negative control (grey filled histograms), and the

wildtype VL9 peptide was a positive control (pale blue filled histograms). Data are

representative from one of three independent experiments.

Figure. S2 Phenotypic analysis of NK-92 cells.

a, NKG2A and CD94 expression in NK-92 cell line detected by flow cytometry. NK-92
cells were stained with PE-CD94 antibody or FITC-NKG2A antibody and analyzed in flow
cytometer. A PE-isotype and FITC-isotype antibodies were used as negative controls.

**b, Fc receptors CD16, CD32 and CD64 expression in NK-92 cell line detected by flow
cytometry.** NK-92 cells were stained with BV650-CD16 antibody, APC-CD32 antibody, or
BV421-CD64 antibody and analyzed in flow cytometer. Peripheral blood mononuclear cells
(PBMCs) were used as positive controls. Dot plots overlay of antibody stained cells (red)
and unstained control cells (grey) were shown. NK-92 cells were negative for CD16, CD32
or CD64, while a subset of PBMC cells were positive for each antibody. Data from a single
antibody phenotype experiment.

Figure. S3 Affinity optimization of 3H4 IgG.

a, NK cell cytotoxicity against wild-type 3H4 IgG-treated target cells measured by ^{51}Cr

release assay. Unoptimized, wild-type 3H4 on a human IgG1 backbone (3H4 Gwt) was

incubated with HLA-E-VL9 transfected 293T cells and untransfected 293T cells at final

concentration of 10 µg/ml, 1 µg/ml, or 0.1 µg/ml, and NK92 cells were added into the mixture

as effector cells at effector: target (E:T) ratios of 20:1 and 6:1. Human IgG1 CH65 was used

as an isotype controls. Dots represent the mean values of triplicate wells in four or five

independent experiments. Asterisks show the statistical significance between indicated

groups: ns, not significant.

b, 3H4 residues optimized for affinity improvements by library screening. Seven

different libraries were designed that simultaneously sampled group of 4 amino acids in the

CDR loops of 3H4 that interact with HLA-E-VL-9 by structural analysis. *Top*: Structural

mapping of the amino acids (*spheres*) sampled together in the different libraries. Residues

shown in the same color were randomized together. HLA-E: *green*; VL-9: *orange*; *Bottom*:

Amino acid sequence of the seven 3H4 libraries, with randomized residues marked with '*X'*

and colored as in the structural panels above.

c, Binding of wild-type 3H4 and variants on transfected 293T cells. Wild-type 3H4 and

variants were titrated on HLA-E-VL9-transfected or untranfected 293T cells. Mean

fluorescent intensity (MFI) from one of three independent experiments were shown.

d, Enhanced NK-92 cell cytotoxicity by optimized IgG 3H4 Gv5 and 3H4 Gv7 on HLA-

E-VL9 transfected 293T cells and untransfected 293T cells, in compare with IgG 3H4

Gwt. Dots represent the mean values of triplicate wells in four or five independent ^{51}Cr

release assays. Statistical analysis was performed using mixed effects models. Asterisks

show the statistical significance between indicated groups: ns, not significant, *P<0.05,

**P<0.01, ***P<0.001, ****P<0.0001.

Figure. S4 Isolation and characterization of human HLA-E-VL9-specific antibodies.

a-b, HLA-E-VL9-specific B cell phenotyping from mice. Splenic cells isolated from HLA-

B27/β2M TG mice (a) or C57BL/6 mice (b) were stained and analysed flow cytometry for B

cells (B220+CD19+) that are HLA-E-VL9 double positive, HLA-E-RL9HIV negative and HLA-

E-RL9SIV negative.

c, Gating strategy of the single cell sorting for HLA-E-VL9-specific B cells from a

Cytomegalovirus (CMV)-negative, male human. Human B cells were first enriched from

PBMCs by pan-B cell negative selection magnetic beads. The enriched cells were stained

and gated on viable/$CD14^{neg}$/$CD16^{neg}$/$CD3^{neg}$/$CD235a^{neg}$/$CD19^{pos}$/HLA-E-$VL9^{pos}$/HLA-E-

$RL9HIV^{neg}$/ HLA-E-$RL9SIV^{neg}$ subset as shown. Cells were single-cell sorted into 96-well

plates for the downstream PCR cloning. Representative data from one of the four donors

were shown.

d-e, Flow cytometry titration of purified HLA-E-VL9-specific mAbs isolated from a

CMV-negative, male human. Antibodies recovered from sorted B cells were constructed in

human IgG1 backbones and used for staining titration on both C-trap-stabilized and

unstabilized HLA-E-VL9, HLA-E-RL9SIV, HLA-E-RL9HIV transfected 293T cells. EGFP

expression indicates transfection efficiency. Transfected cells were stained with testing

antibodies at the concentration of 2 μg/ml, followed by secondary antibody AF555-anti-

human IgG staining. (d) Staining data of a representative antibody CA147 and a negative

control antibody CA136. (e) Summary of the MFI of antibody binding data shown as bar

chart. Data are representative from one of two independent experiments.

f, Cross-reactivities of human HLA-E-VL9 antibodies with rhesus Mamu-E-VL9 and

mouse Qa-1b-VL9 complex. 293T cells were transfected with HLA-E-VL9, Mamu-E-VL9,

two HLA-E/Mamu-E hybrids [HLA-E α1/Mamu-E α2 (Hα1/Mα2) and Mamu-E α1/HLA-E α2

(Mα1/Hα2)], and Qa-1b-VL9. Transfected cells were stained with human antibodies CA123,

CA133, CA143, and CA147, followed by AF647 conjugated anti-mouse IgG(H+L) secondary

antibody. Data are representative from one of three independent experiments.

g, Mapping of representative HLA-E-VL9-specific mAbs CA123, CA133, CA143 and

CA147 on 293T cells transfected with HLA-E-VL9 peptide variants. 293T cells were

transfected with HLA-E SCTs with VL9 peptides with single amino acid mutations at P1, then

stained with human antibodies CA123, CA133, CA143, and CA147, followed by AF647

conjugated anti-mouse IgG(H+L) secondary antibody (dark blue). Cells were gated for EGFP

positive subsets. MFI of the indicated antibody staining on wildtype VL9 peptide was set as

100%, and the percentages equals to (MFI of binding on each P1 variant) / (MFI of binding

on wildtype VL9) x 100%.

h, Affinity measurements of human HLA-E-VL9 antibodies binding to soluble HLA-E-

VL9 complex. Human antibodies CA123 or CA147 on human IgG1 backbone was

immobilized on CM5 sensor chips and soluble HLA-E-VL9 complex protein at the indicated

concentrations was flowed over the antibody immobilized sensor chips. Rate constants (k_a,

k_d) and dissociation constant K_D were measured following curve fitting analysis.

i-j, NK cell cytotoxicity against CA123 IgG-treated target cells as assessed by [51]Cr

release assay. Human antibody CA123 was incubated with HLA-E-VL9 transfected 293T

cells (i) and untransfected 293T cells (j) at final concentration of 30 µg/ml, 10 µg/ml or 3

µg/ml, and NK92 cells were added into the mixture as effector cells at effector: target (E:T)

ratio of 20:1 and 6:1. Human antibody A32 were used as the isotype control. Dots represent

the mean values of triplicate wells in five independent experiments. Statistical analysis was

performed using mixed effects models.

Figure. S5 J chain sequence analysis of HLA-E-VL9-specific antibodies (n=51).

Reference VH-VL repertoires (n=198,148) from three healthy humans from a previous study

(DeKosky et al., 2015) was used as a control.

a-b, Heavy chain (J_H) gene usage shown as bar chart (a) and pie chart (b).

c-d, Kappa chain (J_k) and lambda chain (J_L) gene usage shown as bar chart (c) and pie

chart (d).

 SUPPLEMENTAL TABLES

Table S1. Gene usage and mutation rate of four HLA-E-VL9-specific mAbs isolated

from immunized transgenic mice.

Table S2. Crystallographic data for the 3H4 Fab and VL9-bound HLA-E co-complex

structure.

a, Crystallographic data collection and refinement statistics. AS: Ammonium sulphate.

§ r.m.s.d.: root mean square deviation from ideal geometry. Statistics for outer shell

indicated in parentheses. AU: asymmetric unit. Rfree equals the R-factor against 5% of the

data removed prior to refinement.

b-f, Inter-chain RMSD and inter-molecular interfaces in the 3H4-HLA-E-VL9 structure.

b, Table detailing the total buried surface area of the interface in $Å^2$ between the 3H4 VH, VL

and the VL9-bound HLA-E complex. **c,** Table of RMSD (root mean square deviation) in Å

between chains of the 3H4-HLA-E-VL9 co-complex structure. Two copies of the 3H4 Fab-

HLA-E-VL9 co-complex were present in the asymmetric unit and thus RMSD between

chains related by non-crystallographic symmetry was calculated via Cα atom pairwise

alignment on the PDBePISA server. Average Cα atom RMSD following pairwise alignment is

also reported for the HLA-E heavy chain (HC) of 1MHE (Chain A), a previously published

non-receptor-bound VL9-loaded HLA-E complex, and the HLA-E HC from the 3H4-HLA-E-

VL9 structure reported here (Chain A). **d,** Table listing residues involved in the interface

between the 3H4 VH and the VL9 peptide. **e,** Table of interacting residues of the 3H4 VH

and HLA-E HC interface. **f,** Table of interacting residues of the 3H4 VL and HLA-E HC

interface.

g-h, Hydrogen bonding and salt bridges in the 3H4-HLA-E-VL9 structure. Table of

hydrogen bonds and salt bridges formed between the 3H4 heavy chain (HC) and the HLA-E

HC (**g**) and the 3H4 light chain and the HLA-E HC (**h**).

Hydrogen bonding cut-offs according to the PDBePISA default criteria. 3H4 chain numbering

is according to the Kabat scheme whereby alternate insertion codes (letters after the residue

number) are added to variable length regions of the antibody sequence. 3H4 residues within

the CDRs are shaded green and labelled 'CDR1/2/3'. The position of HLA-E HC residues

either on the α1 or α2 helix is also noted. Amino acid atom abbreviations: C - mainchain

Carbon atom, O - mainchain Oxygen atom, N - mainchain Nitrogen atom, CA - α-Carbon

atom, CB - β-Carbon atom, CD - δ-Carbon atom, CE - ε-Carbon atom, CG - γ-Carbon atom,

CH - η-Carbon atom,, CZ - ζ-Carbon atom, OD - δ-Oxygen atom, OE - ε-Oxygen atom, OG -

γ-Oxygen atom, OH - η-Oxygen atom, ND - δ-Nitrogen atom, NE - ε-Nitrogen atom, NH - η-

Nitrogen atom, NZ - ζ-Nitrogen atom.

Table S3. Information of human subjects used in this study.

Table S4. HLA-E-VL9-specific antibodies isolated from human. Index sorting MFI,

immunogenetics information, and transfected 293T cell staining MFI of the 60 HLA-E-VL9-

specific antibodies isolated from single B cell sorting were shown. The original isotypes of all

the antibodies were IgMs. Antibodies from the same clonal family were highlighted in yellow.