New insight into bacterial iron competition: A-to-I editing in ferric entero

bactin uptake

Wenhan Nie [1*], Sai Wang [1*], Qin Xu [2], Peihong Wang [1], Yan Wu [1], Rui He [3],

Ayizekeranmu Yiming [1], Jingling Liang [1], Iftikhar Ahmad [1,5], Luoyi Fu [1], Longbiao

Guo [4], Junhua Yuan [3], Bo Zhu [1$], Gongyou Chen [1$]

[1] Key Laboratory of Urban Agriculture by Ministry of Agriculture of China, School of

Agriculture and Biology, Shanghai Jiao Tong University, Shanghai, China

[2] State Key Laboratory of Microbial Metabolism, and SJTU-Yale Joint Center for

Biostatistics and Data Science, School of Life Sciences and Biotechnology, Shanghai

Jiao Tong University, Shanghai 200240, China

[3] Hefei National Laboratory for Physical Sciences at the Microscale and Department

of Physics, University of Science and Technology of China, Hefei, Anhui, China

[4] State Key Lab for Rice Biology, China National Rice Research Institute, Hangzhou

310006, PR China

[5] Department of Environmental Sciences, COMSATS University Islamabad,

Vehari-Campus, Vehari-61100, Pakistan

[*] Contribute equally

[$] To whom correspondence should be addressed

Contact: gyouchen@sjtu.edu.cn or bzhu1981@sjtu.edu.cn

Running Title: A-to-I editing in iron metabolism

ABSTRACT

Iron is an essential element for growth and survival of pathogenic bacteria. However, how these bacteria sense and respond to iron starvation or excess is still not fully understood. Here, we show that *xfeA* (a homolog of *fepA*) in *Xanthomonas oryzae*, can sense the extra-cytoplasmic iron concentration and change the 3D structure of ligand channel domain by A-to-I RNA editing. The percentage of A-to-I RNA editing was increased to 76.87% under iron-starvation, facilitating iron to pass through the XfeA channel without any interference to the strongest Iron(III)–siderophore $[Fe-(Ent)]^{3-}$. However, under high iron concentration, the percentage of A-to-I editing to *xfeA* was reduced, thereby restricting the Fe-Ent to pass iron through the channel of XfeA. The variations in concentration of Fe-Ent was sensed by methyl-accepting chemotaxis proteins (MCPs) at cytoplasmic membrane, which influences the expression of a number of chemotaxis related genes. This regulation drives the bacteria to move further towards the iron-rich direction. Overall, our results revealed a new signaling mechanism that bacterial iron sense and homeostasis are modulated through A-to-I RNA editing.

IMPORTANCE

A-to-I RNA editing, which is catalyzed by the adenosine deaminase, acts on RNA family of enzymes, is one of the most prevalent type of post-transcriptional modification in metazoans. However, the research of A-to-I editing in bacteria is limited and the importance of this editing is still underestimated. In this study, we found that bacteria may use A-to-I editing (*xfeA*, a homolog of *fepA*) as an alternative

1 strategy in iron metabolic uptake, as the editing event level can precisely regulate the

2 RNA and the protein sequence within relatively short physiological time scales

3 comparing with the traditional switch "on" and "off" regulation. To best of our

4 knowledge, it is a new identified inching-switch-like mechanism by which pathogens

5 may absorb iron better in order to compete with the hosts.

INTRODUCTION

Iron is one of the most abundant element in the Earth's crust and virtually

essential for all forms of life, including bacteria during all aspect of cellular processes

such as energy generation, respiration, photosynthesis, DNA replication to oxygen

transport and protection against various stresses (1, 2). Bacterial pathogens must

acquire iron within their hosts in order to survive and replicate. The competition of

iron for pathogenic bacteria is extremely important because they invade the host

plants or animals where iron is limited, the termed known as nutritional immunity (3).

However, the hosts retain iron and other nutrients by sequestration with various

proteins or small-molecular chemicals (4, 5). To compete for iron acquisition, bacteria

developed very efficient iron (usually ferric Fe^{3+}) uptake mechanisms based on

high-affinity iron-carriers, named siderophores (6), of which, enterobactin (Fe-Ent) is

one of the strongest iron chelator, which is produced and secreted by many pathogenic

bacteria (7).

Many Gram-negative bacteria including model plant pathogen *Xanthomonas*

oryzae, can produce and take up this siderophore to scavenge iron (8). Generally, this

molecule is first synthesized within the bacterial cell and secreted through the cell membranes to capture Fe(III) from the environment. The resulting Fe(III)-enterobactin complex (Fe-Ent) is recognized at the bacterial cell surface by highly selective outer-membrane TonB-dependent receptors (TBDR) and actively transported into the periplasm (9). FepA is the most well characterized receptor in previous study. In periplasm, Fe-Ent is sequestered by the periplasmic binding protein (PBP), such as FepB, which transfers Fe-Ent to inner membrane transporter for further transport into the cytoplasm (6). Once there, Fe-Ent is hydrolyzed to release Fe(III) and convert to Fe^{2+} for further usage in the cell (10). Interestingly, we recently found that A-to-I editing event on the mRNA of $fliC$ can enhance the Fe^{3+} uptake ability and thus, increase the biomass in $vivo$ and enhance the virulence (11). Post-transcriptional A-to-I editing has been proved to be important in eukaryote however, it was recently reported in bacteria (12) and barely known of the biological function. Since the A-to-I RNA editing in K^+ channel's pore of octopus has been reported that could alter the channel open state to respond to the extreme cold environment (13), this raises the possibility that bacteria may also use the similar mechanism to regulate the iron uptake.

In this study, our objectives were to identify the biological function of A-to-I RNA editing event (T408A, threonine to alanine in the 408[th] amino acid) occurred in the xoc_3486 ($Xanthomonas$ $fepA$ homolog, named $xfeA$) that may enhance the bacterial iron uptake capacity and the tolerance to iron depleted conditions, and to explore the occurrence conditions of the T408A editing event and demonstrate its

significance in bacterial chemotaxis and virulence.

RESULTS

The T408A editing level in *xoc_3486* (*xfeA*) depend on the environmental iron

concentration

Previously, we have reported that the A-to-I RNA editing event occurred in *xfeA*,

causing change in an amino acid from threonine to alanine at residue 408 (11). Here,

we have analyzed potential editing events in cDNA samples from BLS256 WT in the

presence of 2, 2'-dipyridyl (DP) and $FeCl_3$ as iron chelator and supplements,

respectively (Fig. 1a). The results suggest that T408A editing level changed

depending on the environmental iron concentration, generally the rate of A-to-I RNA

editing rise with the depletion of environmental iron concentration. Specifically, in the

presence of NB medium only, 21.27% of WT transcriptional samples showed editing

events; however, hardly any editing event was detected in the samples of NB plus

100μM $FeCl_3$. In the samples of iron depletion, 37.39%, 76.87% and 78.03% of the

cDNA samples exhibited the editing events in the presence of 50μM, 100μM and

150μM DP, respectively (Fig. 1a and 1b). To block the RNA editing, the synonymous

mutation (ACG to ACA, red arrow) containing strain named T408[silent] was constructed,

and hardly A-to-I RNA editing event was detected at this site (Fig. 1b and Fig. S5).

We analyzed the A-to-I RNA editing sites from RNA-seq data of T408A and

T408[silent] strain growth in NB medium by AIMAP. Interestingly, 9 of 12 A-to-I editing

sites were detected in both T408A and T408[silent] strain (Table S4, orange labeled) and

the RNA editing patterns were generally coherent (Fig. S5). Besides, we checked the

A/G rate on *xfeA* T408A site from the reads of RNA-seq data, and the result suggested

that the editing level of *xfeA* is 100% and 0% in T408A and T408[silent] strain (Table S4,

blue labeled and Fig. S5 red frame). Thus verifying the construction of T408[silent] strain

could block the A-to-I RNA editing of *xfeA* and highlighting the importance of the

motif.

Interestingly, the editing event is independent on the *tadA* (tRNA-specific

adenosine deaminase), in spite of both the TACG motif and the RNA secondary

structure (in a loop) coincide with the enzyme recognized (12) (Fig. 1b and Fig. S2).

In addition, no homologues of adenosine deaminase *rdrB* were found in *Xanthomonas*

oryzae genome, which was recently identified by Gao (14) and named phage

restriction by an adenosine deaminase acting on RNA (RADAR). Since the A-to-I

RNA editing was blocked in T408[silent] strain, we hypothesized that the *xfeA* T408A

mutation was induced by iron deficiency and catalyzed by an (or several) unknown

enzyme(s), as well as the recognition base on the motif.

T408A editing is predicted to impact the structure of XfeA

The XfeA protein is a homolog of FepA which is the ferric enterobactin (Fe-Ent)

outer membrane receptor protein (15). Furthermore, FepA plays an important role of

Fe-Ent entry which is the most important way of iron uptake in many gram negative

bacteria (16, 17). Thus, it was hypothesized that the T408A editing event could

change the efficiency of Fe-Ent entry, and thereby cause variations in the ability of

iron uptake. To further investigate the impact of the T408A exchange, a

three-dimensional homology model of XfeA was constructed based on the crystal

structure of a similar TBDR, PiuA from *A. baumannii* (PDB ID: 5FP1) (18). In this

model, XfeA is composed with a 22-stranded transmembrane β-barrel surrounding a

plug domain as other TBDRs (Fig. 2a). The mutated residue Thr408 is on the edge of

the entrance of the β-barrel, with its side chain hydrogen-bonded with those of

Glu416 and Gln453 (Fig. 2b in red circle). This local hydrogen bonding network may

help to fix the flexible extracellular segment of 408-420, with the long side chains of

the residues Arg418 and Arg419 pointing toward the entrance that may contribute to

the binding of the ligands.

T408A mutants show increased tolerance to iron depleted stress

Since the T408A editing event was induced by iron depletion, we hypothesized

that editing event may contribute in environmental iron deficiency. To further

investigate, the growth curve of BLS256 WT, T408^{silent} and the A-G point mutation

containing strain named T408A were compared without and with the presence of

50μM, 100μM and 150μM DP in NB medium (Fig. 3 a-d). The strains grown in NB

medium without or with 50μM DP showed similar growth patterns (Fig. 3a and 3b);

however, lower growth rate of log phase and longer time of lag phase were observed

in the WT and T408^{silent} strains grown in NB supplemented with 100μM and 150μM

DP (Fig. 3c and 3d). The T408A strains (fixed 100% editing level) in growth

interfering level of DP showed enhanced tolerance, while WT (free to edit) showed a

slightly higher tolerance to iron depletion when compared with T408^{silent} (block

editing). Notably, these results suggest that the level of T408A editing could help *Xoc*

adapt to environmental iron deficiency.

T408A editing enhance the iron uptake ability

The intracellular concentration of total iron in the T408A, T408^{silent} and WT strains were quantified with inductively coupled plasma optical emission spectrometry (ICP-OES). In the NB medium plus 50μM DP, the iron accumulation in *Xoc* cells of T408A and WT strains significantly increased by 76.0% and 46.9%, respectively of the level in the T408^{silent} strain (Fig. 3e). In the NB medium, the iron concentration in the T408A and WT strains also increased by 175.3% and 110.1%, respectively to the levels in the T408^{silent} strain. Under iron-replete condition (NB+100μM FeCl$_3$), the iron content of T408A mutation also 145.6% higher than T408^{silent} while the level of WT still 23.6% higher than T408^{silent} (Fig. 3e). This may be related to the decline of the WT editing level in high iron concentration condition. The sufficient free iron leads the T408A strain showed a better iron uptake performance in iron supplement, and iron uptake of other pathways may lead the smaller difference between T408A and T408^{silent} strains in iron limitation. These results exhibit the importance of T408A RNA editing on *xfeA* for iron uptake.

Streptonigrin (SNG) is a drug that the iron is indispensable for its bacteriostatic activity and its toxic effect is proportional to the intracellular iron concentration (2, 19). To further characterize the relationship between T408A editing and iron uptake, resistance of *Xoc* strains to SNG was measured. In the presence of 1 μg/mL SNG, the survival rate of T408A was strongly lower (by 69.8%), whereas the WT was lower (by 23.2%) than T408^{silent}. Under the iron-depleted condition (NB plus 50μM DP), the survival rates of three stains were varying degrees of upward (Fig. 3f). Whereas the

ordering of strains survival rate was unaltered (T408A < WT < T408$^{\text{silent}}$). Taken

together, these results supported the view that *xfeA* T408A editing event could help

Xoc to take-up iron, especially it could increase the tolerance to iron depleted stress.

T408A editing upregulates genes related to chemotaxis pathway

The RNA-seq was used to compare the T408A and T408$^{\text{silent}}$ strains to further

investigate the change of *xfeA* T408A editing for *Xoc*. However, before comparing

RNA-seq profiles of two strains, reproducibility was evaluated for two replicate

experiments using pairwise linear correlation analysis. The correlation coefficients (*r*)

between the two replicate experiments were 0.991 and 0.998, suggesting satisfactory

reproducibility of RNA-seq data under the experimental conditions. Based on the

cutoff as described in the M & M section, 138 differentially expressed genes were

identified (Fig 4a, Table S3). The KEGG pathway enrichment analysis suggested only

one pathway (Bacterial chemotaxis) was enriched in these DEGs (Table. S3).

Interestingly, 21 of 24 chemotaxis pathway genes were up-regulated at least 1.5-fold

(FDR < 0.01). Based on the annotations, 5 genes (*xoc_2278*, two-component system

response regulatory protein, CheB; *xoc_2280*, chemotaxis protein methyltransferase 2,

CheR; *xoc_2289*, *xoc_2291*, *xoc_2297*, methyl-accepting chemotaxis protein, MCP)

were selected to verify with qRT-PCR. The result suggested that the expression level

of chemotaxis pathway genes generally positive correlation with the environmental

iron concentration, and the level of T408A are overall higher than T408$^{\text{silent}}$ (Fig. 4b).

Besides, the expression of *xfeA* was independent of environmental iron content (Fig.

4b and Fig. S1). The chemotaxis pathway diagram of *Xoc* (based on KEGG) was

constructed to visually exhibit the influence of T408A editing on upregulation of

genes, while the color represents the gene expression level (the value is averaged if

have homologous gene). Interestingly, almost all genes in chemotaxis pathway were

up-regulated (Fig. 4c), which revealed that T408A editing event may induce more

sensitivity in the bacteria to chemoattractant.

To demonstrate the effect of chemotaxis pathway genes up-regulated, the

capillary assay was performed to determine the chemotactic potential of T408A,

T408^{silent} and WT strains towards glucose (2.0 mg/mL), serine (10 mg/mL), Fe-Ent

(10 μM and 100 μM) and PBS buffer (represent random diffusion). The relative (to

glucose) chemotaxis response for each strain was calculated, and the data showed that

T408A strain comparing to T408^{silent} strain had increased the sensitivity of Fe-Ent

(both 10 μM and 100 μM level), whereas the chemotactic ability of serine did not

show significant difference (Fig. 4d).

Based on the results of chemotaxis response, two of MCPs (methyl-accepting

chemotaxis proteins, XOC_2282 and XOC_2291) were expressed and purified to

determine whether the MCPs directly interact with $FeCl_3$ or Fe-Ent using Fortebio

Octet RED system (Fig. 5). In association and dissociation phases, the association

curves of Fe-Ent (green and blue) exceeded the 0 nm line, which shows that MCPs

could bind Fe-Ent directly (Fig 5c and 5d). Whereas, the curves of $FeCl_3$ (bronzing)

were lower than 0 nm line (Fig 5a and 5b), which conferred the lack of potential

binding affinity between the MCPs and Fe^{3+}. In addition, the binding kinetic analysis

demonstrated that the MCPs indeed directly bind Fe-Ent rather than Fe^{3+}, with a K_D

value of 1-10μM level (Table 1). In summary, T408A editing event could enhance the

sensitivity of bacterial chemotaxis towards Fe-Ent, thus it directly binds with MCPs to

stimulate the bacterial sense of iron.

T408A editing contributes to virulence of *Xoc*

Since the importance of iron uptake and chemotaxis for virulence (20-23), leaves

of six-week-old susceptible rice cv. Yuanfengzao were inoculated with *Xoc* strains and

mutants (Fig. 6a). At 14 days after inoculation, lesions induced by the T408A mutant

were 3.17□cm; this was significantly longer than 1.88 cm lesions induced by

T408[silent] and 2.33 cm of WT strain (Fig.□6b). Compare to WT, there was a clear

decrease in the virulence of the T408[silent] mutant, suggesting the level of T408A RNA

editing contributes to virulence. Furthermore, *in planta* growth assays indicated that

the colonial population of the T408A mutant was significantly higher than the other

strains, besides the colonizing capacity of T408[silent] was obviously lower than WT

strains (Fig. 6c). Overall, the T408A editing event could improve the niche adaption

of iron depletion; moreover, the increase of iron uptake efficiency mediated by T408A

editing could significantly enhance the virulence of the *Xoc* strain.

DISCUSSION

The detection of iron concentration is necessary in bacteria including pathogenic

bacteria to determine their successful survival and infection in the environment (2).

Currently, the most well studied example is Fur, which is an important cytoplasmic

regulator of iron homeostasis in bacteria (24). It can regulate expression of beneficial

or detrimental genes based on the limitation of ferrous iron (24). Interestingly, our

recent study (11) and this study suggest that bacteria may use our newly identified

strategy (post-transcriptional A-to-I modification) to regulate the uptake of the ferrous

iron efficiently without the regulation by Fur.

A-to-I RNA editing, which is catalyzed by the adenosine deaminase, acts on RNA

(ADAR) family of enzymes, is one of the most prevalent type of post-transcriptional

modification in metazoans (25, 26). In eukaryotes, this modification has been proved

to play an important role in all aspects of regulatory processes such as splicing,

microRNA targeting and processing and mRNA stability (27). The editing of nuclear

protein-coding transcripts was long thought to be absent in bacterial species until

recently that several cases were discovered (11, 12). From our recent and this study,

we found that bacteria may use A-to-I editing as an alternative strategy in iron

metabolic uptake, as the editing event level can precisely regulate the RNA and the

protein sequence within relatively short physiological time scales (12) comparing with

the traditional switch "on" and "off" regulation by Fur regulators.

Iron is an essential for bacteria, however, iron excess is also harmful to organisms.

To sense the iron is limited or excessed, many bacteria employ the ferric uptake

regulator (Fur) to regulate cellular Fe^{2+} levels (28). When at high levels, Fe^{2+} acts as a

co-repressor forming a Fur-Fe^{2+} homodimer that binds to *cis*-regulatory element of

iron uptake-associated genes to repress their transcription (29). In contrast, at low

Fe^{2+} levels, the Fur-Fe^{2+} homodimer is relieved to clear the promoter region and

allowing gene expression (28). Previous report has revealed that Fur-mediated

regulation controls the expression of dozens of genes including some virulence factors

(30). This result suggests that the change of the regulation of Fur will cause huge impact on all aspect of bacteria.

Because of the double-edge sword effect of iron for bacteria (2), an efficient and precise mechanism to regulate iron absorption is necessary. Our study revealed that in *X. oryzae* pv. *oryzacola*, XfeA is a TBDR that mediates Fe-Ent transport across the outer membrane (Fig. 7). Under low iron concentrations, the percentage of A-to-I editing of the *xfeA* was increased, causing changes in the 3D structure of XfeA, which is facilitating iron to pass through the XfeA channel without any interference to the Fe-Ent ligand (Fig. 7). However, under high iron concentration, the percentage of A-to-I modification to *xfeA* was reduced, thereby restricting the Fe-Ent to pass iron through the channel of XfeA (Fig. 7). The signal of changed concentration of Fe-Ent was received by MCP at cytoplasmic membrane, which modulated the expression of a number of chemotaxis related genes as identified by RNA-seq and direct binding kinetic analysis. Besides the editing is flexible and immediate (31), thus making the modification event like an inching-switch instead of switch on/off regulation. Based on these findings, it was hypothesized that this mechanism evolved due to adaptation of bacteria in the changing environment.

MCPs, which are involved in the regulation of diverse aspects of cellular activities, are the predominant chemoreceptors in bacteria (32). A typical MCP contains a ligand-binding domain (LBD), which can recognize their ligands directly (33). Many types of ligands such as serine, maltose, oxygen and irons have been proven to bind the MCPs, thereafter, transduce the signal to the downstream genes

(34). In this study, we found that rather inorganic Fe^{3+}, Fe-Ent can bind to MCP directly. To the best our knowledge, this is the first report that MCP can bind to Fe-Ent directly. Moreover, we also found that the signal of MCP binding to Fe-Ent can be transduced to regulate the activity of CheA and other chemotaxis related genes, confirmed by RNA-seq and qPCR (Fig. 4, Table S3). Based on the model of *E.coli*, the MCPs-CheW-CheA may form a signaling kinase complex and transduce the signal to response regulator CheY by phosphorylation. Phosphatase CheZ, methyltransferase CheR and methylesterase CheB constitute the regulator and MCPs modification pathway, therefore the seven different types of proteins constituting the chemotaxis signal transduction system in prokaryotes (35). From the above, this regulation and its consequences on the chemotaxis system can be served as precise sensory adaptation in bacterial chemotaxis through A-to-I RNA editing. The evidence of directly binding between MCP and Fe-Ent (instead of Fe^{3+}) can help to research of the iron chemotaxis and the key interacting residues could be found with further studies.

Overall, the results of this study revealed the significance of the A-to-I RNA editing for bacterial iron metabolic uptake. Using the RNA-seq and Sanger sequence, we demonstrated the rate of A-to-I modification rise with the depletion of environmental iron concentration, and the T408A editing event can improve the bacterial iron uptake capacity and the tolerance to iron depletion. We provide the evidence that MCPs could bind to Fe-Ent directly, and that T408A editing event plays an important role in the chemotaxis. Virulence and colonization assay of susceptible rice provide further evidences about the T408A editing event could improve the niche

adaption of iron depletion. Interestingly, we found that T408A editing in our study is

*tad*A independent. Recently, Gao and his colleagues found that there are other types

of A-to-I editing enzymes existing in bacterial genomes (14). Strikingly, we can't find

any homologs of these genes in the genome of *Xoc*. This result suggests that there

should be another unknown A-to-I editing gene existing in the genome of *Xoc*.

In conclusion, we have identified a new inching-switch-like mechanism by which

pathogens may absorb iron better in order to compete with the hosts, furthermore the

level of RNA editing is regulated by environmental iron concentration.

MATERIALS AND METHODS

Bacterial strains, plasmids, and plant materials

The bacterial strains and plasmids used in this study are described in Table S1.

Xoc BLS256 is the wild-type (WT) strain of *X. oryzae* pv. *oryzicola.* T408$^{\text{silent}}$

contains a synonymous mutation (TAC<u>G</u> to TAC<u>A</u>) in *xfeA* that blocks

post-transcriptional A-to-I modification at amino acid 408. Strain T408A is a

derivative of BLS256 containing a mutation (T<u>A</u>CG to T<u>G</u>CG) that changed the

threonine residue at amino acid 408 to alanine. Strain BL21-2282 and BL21-2291 is a

derivative of BL21 that harbor the pET-30a::*xoc_2282* and pET-30a::*xoc_2291* to

express protein of MCPs (XOC_2282 and XOC_2291). Further details of strains and

plasmids are provided in Table S1.

Escherichia coli strains were cultured in Luria-Bertani (LB) medium at 37°C (36).

Xoc BLS256 strains were cultured in nutrient broth (NB, 1% peptone, 0.5% yeast

extract, 1% sucrose) or in NB containing 1.5% agar at 28°C. The final concentrations

of the antibiotics were as follows: kanamycin, 25 μg/mL; cephalexin, 40 μg/mL and

streptonigrin (SNG) 1 μg/mL. Filter sterilized 2, 2'-dipyridyl (DP) and $FeCl_3$ as iron

chelator and supplements were prepared as 10mM stocks, added in 100- or 200-fold

volume NB medium when used.

Rice seeds were obtained from the International Rice Research Institute. All rice

plants were grown at 28°C with a 12-h photoperiod in a greenhouse located at

Shanghai Jiao Tong University.

Bacterial mutant and strain construction

Construction of bacterial mutant strains was performed as described by Baba (37)

with the following modifications. For site-directed mutagenesis, plasmids were

modified with the Fast Mutagenesis System (Transgen Biotech, Beijing, China) to

obtain clones containing the point mutations (T408A and T408^{silent}). Fragments

encoding *xfeA* in *Xoc* BLS256 were amplified with the corresponding *xoc3486* F/R

primers (Table S2), digested with *Sal*I/*Xho*I, and subcloned into the *Sal*I/*Xho*I site of

pKMS1(38). Mutations were introduced by PCR amplification with overlapping

primers containing the target A-G point mutations as follows: T408A F/R and

T408^{silent} F/R (Table S2). The PCR product was digested with the DMT enzyme

supplied in the Fast Mutagenesis System, and plasmids containing A-G or G-A point

mutations were transformed into DMT competent cells and verified by Sanger

sequencing. Sequence substitutions were performed by double homologous

recombination similar to the gene knockouts. In brief, for the T408A and T408^{silent}

strains, the point mutation carried fragments in pKMS1 were separately introduced

into *Xoc* BLS256 by electroporation (Bio-Rad Laboratories Inc., Hercules, CA, USA)

and subjected to kanamycin selection. A single transformant colony with kanamycin

resistance was picked and cultured for 8 h in NB medium without kanamycin and

sucrose; 10-fold dilutions were then inoculated to nutrient agar (NA) with 15%

sucrose to obtain sucrose-insensitive clones.

To construct the protein-expressing strain (BL21-xoc2282 and BL21-xoc2291),

the whole-length fragments of *xoc_2282* and *xoc_2291* were amplified with the *Pfu*

polymerase (TransGen Biotech, Beijing, China) using the xoc2282 and xoc2291 F/R

primers (Table S2). These fragments were cloned, digested with *BamHI/Hind*III, and

the two fragments were ligated into the *BamHI/Hind*III site of expression vector

pET-30a (+). The protein expression fragments in pET-30a (+) were separately

introduced into BL21 (DE3) by chemical transformation. In brief, 50μl BL21 (DE3)

chemically competent cell added 2.5μl of DNA, and then keep in the ice bath for 30

min. After heat-shock at 42□ for 45 sec, the tube was placed in the ice bath for 2 min,

and then add a 500μl LB medium for reviving and shaking for 1 h. 200μl of

transforming cell were inoculated to LB agar and cultured overnight to obtain protein

expressing clones.

RNA secondary structure prediction

A sequence consisting of residues from -25 to +25 bp from the edited site was

extracted for prediction of RNA secondary structure with the RNAfold web server

(http://rna.tbi.univie.ac.at/cgi-bin/RNAWebSuite/RNAfold.cgi) based on minimum

free energy.

3D structure prediction

For 3D modeling, the sequence of the XfeA was input into the RCSB Protein

Data Bank (PDB) to search for templates that could be used for homology modeling.

The X-ray structure of *A. baumannii PiuA (PDB ID: 5FP1)* was used as the template

for homology modeling of XfeA by an in-house program SAMM (39) with energy

minimization using AMBER10 force field (40). Default values were used for the

iteration limit and RMS gradient test.

Bacterial growth assays in response to iron deplete stress

Optical density (OD) was measured with a Bioscreen C instrument (Labsystem,

Helsinki, Finland). For each well, 99 μL of NB medium with or without DP was

inoculated with 1 μL of an overnight culture (1×10^9 CFU/mL). The OD values at

420-580 nm were measured every 15 min for 24 h with continuous shaking at 28℃.

Four independent replicates were carried out in this study. Pairwise comparisons of

growth curve (Logistic curve fitted by OD values) for each strain

and growth condition were analyzed with the *F*-test against the curve obtained for

T408^{silent}. Data progressing, fitting and drawing with OriginPro 2018C version

9.5.1.195.

Cellular iron concentration

Xoc cellular iron concentrations were detected by inductively coupled

plasma-optical emission spectroscopy (ICP-OES, Optima 8000, PerkinElmer, MA

USA), as reported previously (2, 41). The bacterial cells were grown to an OD_{600} of

1.0 in NB medium. Bacterial cells were collected by centrifugation and washed with

sterilized PBS buffer three times, and the stains were then inoculated into 2-fold

volume of fresh NB medium added 50 μM DP or 100 μM $FeCl_3$. After 3-5 h of

growth (until OD_{600}=0.6), cells were harvested by centrifugation and washed three

times with sterilized PBS buffer. The pelleted cells were oven-dried at 65□ for 48 h,

and digested with the acid solution (HNO_3–$HClO_4$, 4:1, v/v). The digest was then

transferred to a 25 mL volumetric tube and diluted to 25 mL with deionized water.

The iron atom content was then measured by ICP-OES. Samples did not treat with the

acid solution were used as parallel controls. The iron concentration of strain was

calculated by dividing the iron atom value for every 10^9 *Xoc* cells.

Streptonigrin survival assays

Streptonigrin survival assays were performed as described previously with minor

modifications (42). Log-phase (OD_{600}=0.6) *Xoc* cells were harvested by centrifugation,

washed with 0.01M PBS buffer (NaCl 8.5g/L, Na_2HPO_4 2.2g/L, NaH_2PO_4 0.4g/L,

pH=7.0) three times, and exposed to 1μg/mL streptonigrin for 16h at 28°C. Cultures

were subjected to serial 10-fold dilutions, inoculated to NA, and colonies were

counted after a 48 h-incubation at 28 °C. The percentage of survival rate was

calculated by comparing the treated group without streptonigrin. This assay was

performed in triplicate.

Exposure of *Xoc* strains to iron chelator/supplement and RNA extraction

Xoc strains were incubated as described previously with minor modifications (43).

BLS256 WT strain were incubated at 28□ to OD_{600}=1.0 in NB medium with DP or

$FeCl_3.$ Aliquots were withdrawn and pelleted by centrifugation at 4°C. Each cell pellet

was washed two times in cold PBS buffer, and total RNA was then extracted with the

RNeasy Protect Bacteria Mini Kit (Qiagen). Two biological replicates were examined

for this experiment.

qRT-PCR and PCR for cDNA

RNA was purified with EasyPure RNA Kit (Transgen Biotech) as recommended

by the manufacturer. RNA (1 μg) was used to synthesis cDNA with Magic 1st cDNA

Synthesis Kit (With dsDNase) (Magic Biotech, Hangzhou, China). Then, 20 μl of

synthesized cDNA was diluted to 100 μl and used for qRT☐PCR with Magic SYBR

Green qPCR Mix (Magic Biotech) and the ABI 7500 quantitative PCR system

(Applied Biosystems, Foster City, CA). Normalized expression levels of target genes

were calculated relative to rpoD using the ΔΔCT method, where CT is the threshold

cycle. Three independent biological replicates were carried out in this study.

PCR (c T408 F/R primers, Table S2) was performed to sequence the *xfeA*

transcript from cDNA samples of *Xoc* BLS256. The chromatograms were visualized

by Chromas Lite (Technelysium, Brisbane, Australia), and the frequency of editing

was estimated by ratiometric A/G measurement.

mRNA purification and cDNA synthesis for RNA sequence

Total RNA (10 μg/sample) was isolated and treated with the MICROBExpress

Bacterial mRNA Enrichment kit (Ambion) and RiboMinus™ Transcriptome Isolation

Kit (Bacteria) (Invitrogen) as recommended by the manufacturers. Bacterial mRNAs

were suspended in RNase-free water (15 μL) and chemically fragmented to 200-250

bp using 1X fragmentation solution (Ambion) for 2.5 min at 94°C. cDNA was

generated according to instructions provided in the SuperScript Double-Stranded

cDNA Synthesis Kit (Invitrogen). Briefly, mRNA samples were mixed with 100 pmol

of random hexamers, incubated at 65°C for 5 min, chilled in ice, and mixed with 4 μL

of First-Strand Reaction Buffer (Invitrogen), 2 μL 0.1 M DTT, 1 μL 10 mM

RNase-free NTP mix, and 1 μL SuperScript III reverse transcriptase (Invitrogen);

mixtures were then incubated at 50°C for 1 h. To generate the second strand, the

following Invitrogen reagents were added and samples were incubated at 16 °C for

2.5 h: 51.5 μL RNase-free water, 20 μL second-strand reaction buffer, 2.5 μL 10 mM

RNase-free dNTP mix, 50 U DNA polymerase, and 5 U RNase H.

RNA sequencing and RNA-seq data analysis

The Illumina Paired End Sample Prep kit was used to create RNA-Seq libraries as

described previously (44). After removing low quality reads and adaptors, RNA-Seq

reads were aligned to the corresponding *Xoc* BLS256 genome using Tophat 2.0.7 (45),

allowing for a maximum of two mismatched nucleotides. If reads mapped to more

than one location, only the site showing the highest score was retained. Reads

mapping to rRNA and tRNA regions were removed from further analysis. After

obtaining read numbers from every sample, edgeR with TMM normalization method

was used to determine differentially expressed genes (DEGs). Significant DEGs (FDR

value < 0.01) were selected for further analysis, and Cluster 3.0 and Treeview 1.1.6

were used to generate heatmap clusters based on the RPKM values (46, 47).

Capillary chemotaxis assay

For the chemotaxis assay, the capillary method was performed as described

previously with several modifications (20). Briefly, *Xoc* strains were grown in NB

medium to $OD_{600} = 1.0$. The cells were centrifuged at $800 \times g$ for 6 min, washed with

PBS buffer three times and resuspended. Sterilized capillary tubes containing 0.22-µm

filter-sterilized attractant 2.0 mg/mL glucose, 10 mg/mL serine, 10µM Fe-Ent, 100µM

Fe-Ent and 0.01M PBS buffer (as negative control) were incubated with cell

suspension of *Xoc* strains at 28 ℃ for 4 h. To determine the CFU of strains in the

capillaries, the content of the capillary was serially 10-fold dilution plated on PBS

buffer (The diagram see Fig. S3). Relative chemotaxis response (relative to glucose

for each strain respectively) was calculated by the number of migrated bacterial cells

in the capillary, exclude the cell number of PBS buffer containing because of bacterial

random motility or diffusion.

Direct binding kinetic analysis of MCPs to $FeCl_3$ and Fe-Ent

For the protein (XOC_2282 and XOC_2291) expression and purification,

6×His-tag recombinant protein was expressed by constructing the expression vector

(pET-30a) and purified from BL21 strain (Table S1) following the protocol of the

BeyoGoldTM His-tag purification Resin (Beyotime, Shanghai, China), and then the

proteins were concentrated with molecular sieve (Millipore China, Shanghai, China).

The binding kinetic analysis was performed as described previously with several

modifications (48). In brief, $FeCl_3$ or Fe-Ent was dissolved with PBS buffer at the

concentration of 50µM. Similarly, the products of protein concentration were diluted

with PBS buffer at the concentration of 10µM. After baseline step, the His-tag labeled

protein (XOC_2282 and XOC_2291) were bound to the Octet Red biosensors

(Ni-NTA sensor), then washed and blocked by PSA buffer (0.2% PSA and 0.02%

Tween-20 dissolve in PBS buffer), the biosensors were incubated for 300 sec with

$50\mu M$ $FeCl_3$ or Fe-Ent buffer to association, whereas the biosensors were incubated

with the PBS buffer for 600 sec to determine the dissociation rate. The Octet Data

Analysis HT software version 12 was used to fit the curve and calculate the value of

the dissociation constant (K_D).

Plant inoculation assays

Bacterial suspensions (OD_{600}=0.6) were used to inoculate 6-week-old rice with

needleless syringes, and lesion lengths were measured 14 days after inoculation.

Thirteen leaves were inoculated and evaluated for each *Xoc* strain.

For colony counting, three leaves, each infected with one *Xoc* strain, were excised

around the inoculation site and crushed with sterile beads. Each suspension was

subjected to a 10-fold serial dilution, and 5 μL aliquots from 10^{-1} to 10^{-6} dilutions

were inoculated to NA with cephalexin. Plates were incubated at 28°C until individual

colonies could be counted; statistical analysis was used to calculate viable bacteria per

leaf.

Data Availability

All sequence data generated in this study are deposited in NCBI under BioProject

number: PRJNA673071.

ACKNOWLEDGEMENTS

We thank Prof. Fu-Zhou Xu for providing Fe-Ent. We thank Prof. Wei Qian for

his valuable suggestion. This work was supported by National Key R&D Program of

REFERENCES

1. Skaar EP. 2010. The battle for iron between bacterial pathogens and their vertebrate hosts. PLoS Pathog 6**:**e1000949.

2. Wang L, Pan Y, Yuan Z-H, Zhang H, Peng B-Y, Wang F-F, Qian W. 2016. Two-component signaling system VgrRS directly senses extracytoplasmic and intracellular iron to control bacterial adaptation under iron depleted stress. PLoS Pathog 12**:**e1006133.

3. Hood MI, Skaar EP. 2012. Nutritional immunity: transition metals at the pathogen–host interface. Nat Rev Microbiol 10**:**525-537.

4. Kehl-Fie TE, Skaar EP. 2010. Nutritional immunity beyond iron: a role for manganese and zinc. Curr Opin Chem Biol 14**:**218-224.

5. Fischbach MA, Lin H, Liu DR, Walsh CT. 2006. How pathogenic bacteria evade mammalian sabotage in the battle for iron. Nat Chem Biol 2**:**132-138.

6. Raines DJ, Moroz OV, Blagova EV, Turkenburg JP, Wilson KS, Duhme-Klair A-K. 2016. Bacteria in an intense competition for iron: Key component of the Campylobacter jejuni iron uptake system scavenges enterobactin hydrolysis product. Proc Natl Acad Sci U S A 113**:**5850-5855.

7. Hider RC, Kong X. 2010. Chemistry and biology of siderophores. Nat Prod Rep 27**:**637-657.

8. Ryan RP, Vorhölter F-J, Potnis N, Jones JB, Van Sluys M-A, Bogdanove AJ, Dow JM. 2011. Pathogenomics of *Xanthomonas*: understanding bacterium–plant interactions. Nat Rev Microbiol 9**:**344-355.

9. **Schalk IJ, Mislin GL, Brillet K.** 2012. Structure, function and binding selectivity and stereoselectivity of siderophore–iron outer membrane transporters, p. 37-66, Current topics in membranes, vol. 69. Elsevier.

10. Raymond KN, Dertz EA, Kim SS. 2003. Enterobactin: an archetype for microbial iron transport. Proc Natl Acad Sci U S A 100**:**3584-3588.

11. Nie W, Wang S, He R, Xu Q, Wang P, Wu Y, Tian F, Yuan J, Zhu B, Chen G. 2020. A-to-I RNA editing in bacteria increases pathogenicity and tolerance to oxidative stress. PLoS Pathog 16**:**e1008740.

12. Bar-Yaacov D, Mordret E, Towers R, Biniashvili T, Soyris C, Schwartz S, Dahan O,

Pilpel Y. 2017. RNA editing in bacteria recodes multiple proteins and regulates an evolutionarily conserved toxin-antitoxin system. Genome Res 27**:**1696-1703.

13. Garrett S, Rosenthal JJ. 2012. RNA editing underlies temperature adaptation in K^+ channels from polar octopuses. Science 335**:**848-851.

14. Gao L, Altae-Tran H, Böhning F, Makarova KS, Segel M, Schmid-Burgk JL, Koob J, Wolf YI, Koonin EV, Zhang F. 2020. Diverse enzymatic activities mediate antiviral immunity in prokaryotes. Science 369**:**1077-1084.

15. Buchanan SK, Smith BS, Venkatramani L, Xia D, Esser L, Palnitkar M, Chakraborty R, Van Der Helm D, Deisenhofer J. 1999. Crystal structure of the outer membrane active transporter FepA from *Escherichia coli*. Nat Struct Biol 6**:**56-63.

16. Ma L, Kaserer W, Annamalai R, Scott DC, Jin B, Jiang X, Xiao Q, Maymani H, Massis LM, Ferreira LC. 2007. Evidence of ball-and-chain transport of ferric enterobactin through FepA. J Biol Chem 282**:**397-406.

17. Newton SM, Igo JD, Scott DC, Klebba PE. 1999. Effect of loop deletions on the binding and transport of ferric enterobactin by FepA. Mol Microbiol 32**:**1153-1165.

18. Moynié L, Luscher A, Rolo D, Pletzer D, Tortajada A, Weingart H, Braun Y, Page MG, Naismith JH, Köhler T. 2017. Structure and function of the PiuA and PirA siderophore-drug receptors from *Pseudomonas aeruginosa* and *Acinetobacter baumannii*. Antimicrob Agents Chemother 61.

19. Yeowell HN, White JR. 1982. Iron requirement in the bactericidal mechanism of *streptonigrin*. Antimicrob Agents Chemother 22**:**961-968.

20. Kumar Verma R, Samal B, Chatterjee S. 2018. *Xanthomonas oryzae* pv. *oryzae* chemotaxis components and chemoreceptor Mcp2 are involved in the sensing of constituents of xylem sap and contribute to the regulation of virulence◻associated functions and entry into rice. Mol Plant Pathol 19**:**2397-2415.

21. Yao J, Allen C. 2006. Chemotaxis is required for virulence and competitive fitness of the bacterial wilt pathogen *Ralstonia solanacearum*. J Bacteriol 188**:**3697-3708.

22. Garvis S, Munder A, Ball G, De Bentzmann S, Wiehlmann L, Ewbank JJ, Tümmler B, Filloux A. 2009. Caenorhabditis elegans semi-automated liquid screen reveals a specialized role for the chemotaxis gene *cheB2* in *Pseudomonas aeruginosa* virulence. PLoS Pathog 5**:**e1000540.

23. Brown JS, Gilliland SM, Holden DW. 2001. A *Streptococcus pneumoniae* pathogenicity island encoding an ABC transporter involved in iron uptake and virulence. Mol Microbiol 40**:**572-585.

24. Hassan HM, Troxell B. 2013. Transcriptional regulation by Ferric Uptake Regulator (Fur) in pathogenic bacteria. Front Cell Infect Microbiol 3**:**59.

25. Bazak L, Haviv A, Barak M, Jacob-Hirsch J, Deng P, Zhang R, Isaacs FJ, Rechavi G, Li JB, Eisenberg E. 2014. A-to-I RNA editing occurs at over a hundred million genomic sites, located in a majority of human genes. Genome Res 24**:**365-376.

26. Palladino MJ, Keegan LP, O'Connell MA, Reenan RA. 2000. A-to-I pre-mRNA editing in *Drosophila* is primarily involved in adult nervous system function and integrity. Cell 102**:**437-449.

27. Schaffer AA, Kopel E, Hendel A, Picardi E, Levanon EY, Eisenberg E. 2020. The cell line A-to-I RNA editing catalogue. Nucleic Acids Res 48**:**5849-5858.

28. Hantke K. 2001. Iron and metal regulation in bacteria. Curr Opin Microbiol 4**:**172-177.

29. Chen Z, Lewis KA, Shultzaberger RK, Lyakhov IG, Zheng M, Doan B, Storz G, Schneider TD. 2007. Discovery of Fur binding site clusters in Escherichia coli by information theory models. Nucleic Acids Res 35**:**6762-6777.

30. McHugh JP, Rodríguez-Quiñones F, Abdul-Tehrani H, Svistunenko DA, Poole RK, Cooper CE, Andrews SC. 2003. Global iron-dependent gene regulation in *Escherichia coli* A new mechanism for iron homeostasis. J Biol Chem 278**:**29478-29486.

31. Hanswillemenke A, Kuzdere T, Vogel P, Jékely Gsr, Stafforst T. 2015. Site-directed RNA editing *in vivo* can be triggered by the light-driven assembly of an artificial riboprotein. J Am Chem Soc 137**:**15875-15881.

32. Ud-Din AIMS, Roujeinikova A. 2017. Methyl-accepting chemotaxis proteins: a core sensing element in prokaryotes and archaea. Cell Mol Life Sci 74**:**3293-3303.

33. Milburn MV, Prive GG, Milligan DL, Scott WG, Yeh J, Jancarik J, Koshland DE, Kim S-H. 1991. Three-dimensional structures of the ligand-binding domain of the bacterial aspartate receptor with and without a ligand. Science 254**:**1342-1347.

34. Muok AR, Deng Y, Gumerov VM, Chong JE, DeRosa JR, Kurniyati K, Coleman RE, Lancaster KM, Li C, Zhulin IB. 2019. A di-iron protein recruited as an Fe [II] and oxygen sensor for bacterial chemotaxis functions by stabilizing an iron-peroxy species. Proc Natl Acad Sci U S A 116**:**14955-14960.

35. Wuichet K, Zhulin IB. 2010. Origins and diversification of a complex signal transduction system in prokaryotes. Science signaling 3**:**ra50.

36. **Maniatis T, Fritsch EF, Sambrook J.** 1982. Molecular cloning: a laboratory manual, vol. 545. Cold spring harbor laboratory Cold Spring Harbor, NY.

37. Baba T, Ara T, Hasegawa M, Takai Y, Okumura Y, Baba M, Datsenko KA, Tomita M, Wanner BL, Mori H. 2006. Construction of *Escherichia coli* K□12 in□frame, single□gene knockout mutants: the Keio collection. Mol Syst Biol 2**:**2006.0008.

38. Xie G-L, Zhang G-Q, Liu H, Lou M-M, Tian W-X, Li B, Zhou X-P, Zhu B, Jin G-L. 2011. Genome sequence of the rice-pathogenic bacterium *Acidovorax avenae* subsp. *avenae* RS-1. J Bacteriol 193**:**5013-5014.

39. Wang J-F, Wei D-Q, Lin Y, Wang Y-H, Du H-L, Li Y-X, Chou K-C. 2007. Insights from modeling the 3D structure of NAD (P) H-dependent D-xylose reductase of *Pichia stipitis* and its binding interactions with NAD and NADP. Biochem Biophys Res Commun 359**:**323-329.

40. Vanommeslaeghe K, Hatcher E, Acharya C, Kundu S, Zhong S, Shim J, Darian E, Guvench O, Lopes P, Vorobyov I. 2010. CHARMM general force field: A force field for drug□like molecules compatible with the CHARMM all□atom additive biological force fields. J Comput Chem 31**:**671-690.

41. Desrosiers DC, Sun YC, Zaidi AA, Eggers CH, Cox DL, Radolf JD. 2007. The general transition metal (Tro) and Zn^{2+} (Znu) transporters in Treponema pallidum: analysis of metal specificities and expression profiles. Mol Microbiol 65**:**137-152.

42. Si M, Zhao C, Burkinshaw B, Zhang B, Wei D, Wang Y, Dong TG, Shen X. 2017. Manganese scavenging and oxidative stress response mediated by type VI secretion system in *Burkholderia thailandensis*. Proc Natl Acad Sci U S A 114**:**E2233-E2242.

43. Upadhya R, Campbell LT, Donlin MJ, Aurora R, Lodge JK. 2013. Global transcriptome

profile of *Cryptococcus neoformans* during exposure to hydrogen peroxide induced
oxidative stress. PLoS ONE 8**:**e55110.
44. Fang Y, Wang H, Liu X, Xin D, Rao Y, Zhu B. 2019. Transcriptome analysis of
Xanthomonas oryzae pv. *oryzicola* exposed to H2O2 reveals horizontal gene transfer
contributes to its oxidative stress response. PLoS ONE 14**:**e0218844.
45. Trapnell C, Pachter L, Salzberg SL. 2009. TopHat: discovering splice junctions with
RNA-Seq. Bioinformatics 25**:**1105-1111.
46. Saldanha AJ. 2004. Java Treeview—extensible visualization of microarray data.
Bioinformatics 20**:**3246-3248.
47. de Hoon MJL, Imoto S, Nolan J, Miyano S. 2004. Open source clustering software.
Bioinformatics 20**:**1453-1454.
48. Li H, Tao Y, Zhao P, Ban X, Zhi D, Li G, Wang F, Yang X, Huai L. 2015. Recognization
of receptors on bone marrow-derived dendritic cells bound with *Pholiota nameko*
polysaccharides. Int J Biol Macromol 72**:**649-657.

Figure Legends

Figure 1. Analysis of the *xfeA* (T408A) A-to-I RNA editing level by Sanger

sequencing. (a) Sequence analysis of cDNA in edited regions of wild-type (WT) in

NB medium, NB+50μM 2,2'-dipyridyl (DP, as an iron chelator), NB+100μM DP,

NB+150μM DP, NB+100μM $FeCl_3$, and *ΔtadA* (tRNA-specific adenosine deaminase)

strain in NB medium. PCR fragments were obtained from cDNA (mRNA) using

primers described in Methods. Samples were incubated over-night at log-phase and

then subjected to Sanger sequencing. Three independent biological replicates were

carried out in this study (N.S. represent no significance). (b) The conserved editing

TACG motif is framed by the blue frame in the DNA sequence. Chromatograms show

editing level in different culture conditions, and the A/G percentage of the edited

sequence as determined by Sanger sequencing is indicated (A-to-I editing event site in

the red frame). For T408[silent] strain, the red arrow indicates the DNA mutation site of

TACG to TACA, and to demonstrate the change of motif could block the A-to-I

editing on the editing site (red frame).

Figure 2. Homology modeling of XfeA. (a) Homology model of XfeA using PiuA

from *A. baumannii* (PDB ID: 5FP1) as the template. The backbone is colored green

and magenta at the 22-stranded transmembrane β-barrel and plug domains of the

model, respectively. The threonine residue (Thr408) is depicted using spheres. (b)

Comparison of hydrogen bonding around the mutation site. The loss of several critical

H bonds (in red circle) in the Thr408Ala mutant (right) may disrupt the H-bonding

network surrounding Thr408 (left), which may diminish this local tether of this

segment. The carbons of the β-barrel are colored green and the plug domains residues

are colored magenta in both models.

Figure 3. T408A RNA editing modulates bacterial responses to iron-replete and

depleted conditions. (a) Bacterial growth curve of *Xoc* T408A, T408^silent and WT in

NB medium. Strains were grown in quadruplicate to mid-exponential phase, diluted to

OD_{600}=0.1, transferred to fresh NB and placed in a Bioscreen C apparatus at 28°C; (b)

Growth in NB supplemented with 50μM 2,2'-dipyridyl. (c) Growth in NB

supplemented with 100μM 2,2'-dipyridyl. (d) Growth in NB supplemented with

150μM 2,2'-dipyridyl. Error intervals (light shaded regions) indicate mean ± SE of *n*

= 4 replicates. In (a, b, c and d), the F-test against the fitting logistic function obtained

for T408A and WT compared with T408^silent growth curve, * and ** indicate the

significant difference is lower than 0.05 and 0.01 respectively. (e) Iron content in *Xoc*

T408A, T408^{silent} and WT. Bacterial cells were cultured in NB medium added 50μM

2,2'-dipyridyl or 100μM $FeCl_3$. The concentrations of total iron in *Xoc* were measured

with inductively coupled plasma spectroscopy (ICP-OES). (f) Survival rates of *Xoc*

subjected to 1μg/ml streptonigrin (SNG) treatment. Middle log-phase bacterial

cultures (OD_{600} = 0.5) in NB medium (with or without 50μM 2,2'-dipyridyl) were

treated with 1μg/ml streptonigrin (SNG) for 24 h and the survival rates were

calculated by counting the 10-fold serial dilution colony-forming unit (cfu). In (e and

f), error bars represent standard deviations (*n* = 3), * and ** indicate the significant

difference between the samples and the control group T408^{silent} is lower than 0.05 and

0.01 respectively (Student's *t*-test).

Figure 4. The upregulation of a chemotaxis gene pathway, which was detected by

RNA-seq, contributes to enhance the sensitivity of Fe-Ent. (a) Volcano plot showing

the FDR P-value and fold change of T408A versus T408^{silent}. Red dots indicate the

upregulated genes with FDR P < 0.01, while blue dots indicate downregulated genes

with FDR P < 0.01. (b) Normalized expression levels of selected chemotaxis pathway

genes were calculated relative to rpoD using the ΔΔCT method between T408A and

T408^{silent}, where CT is the threshold cycle (all samples normalized to T408^{silent} strain

grown on NB). Three independent biological replicates were carried out in this study.

The significant difference between T408A and T408^{silent} is lower than 0.01 (**) and

0.001 (***, Student's t-test). (c) The gene expression fold change (indicated by color)

of bacterial chemotaxis pathway and flagellar assembly based on the RNA-seq data.

(d) The relative chemotaxis response of *Xoc* strains with different chemoattractants.

The relative chemotaxis response value was calculated by the number of migrated

bacterial cells. The significant difference was calculated by Student's t-test.

Figure 5. Binding kinetic analysis of MCPs (*xoc_2282* and *xoc_2291*) to $FeCl_3$ (Fig

5a and 5b) or Fe-Ent (Fig 5c and 5d) by Fortebio Octet Red System (ForteBio

Analytics Shanghai Co., Ltd.). Real-time association and dissociation analysis of

MCPs to $FeCl_3$ or Fe-Ent. MCPs were bound to the Octet Red biosensors (Ni-NTA

sensor), and then washed and blocked by PSA buffer, the biosensors were incubated

with 50μM $FeCl_3$ or Fe-Ent buffer to association (bronzing: $FeCl_3$; blue: Fe-Ent

(XOC_2282); green: Fe-Ent (XOC_2291); and red: fitted curve), and then the

biosensors were incubated with the PBS buffer to determine the dissociation rate.

Figure 6. Virulence and *in planta* growth in rice cv.Yuanfengzao. Virulence was

assessed by inoculating six-week-old susceptible Yuanfengzao rice plants with

Xoc strains. (a) Symptoms on rice leaves inoculated with the T408A, T408^{silent} and

WT. (b) Leaves (*n*=13) were inoculated with needleless syringes and evaluated for

lesion length 14 days after inoculation. Values represent the mean lesion

length $\pm$ SD. The cross (×) indicates the abnormal value. The asterisks (*) indicate

significant differences between the data obtained for T408A and WT as compared to

the T408^{silent} strain (*, $P < 0.05$; ***, $P < 0.001$; ANOVA with Dunnett's

multiple test post-hoc correction compared with T408$^{\text{silent}}$). (c) Population dynamics

of T408A T408$^{\text{silent}}$ and WT cells *in planta* (means ± SD). Leaves (*n*=3) each infected

with one strain of *Xoc,* were excised around the inoculation site and then crushed with

sterile beads. Each suspension was subjected to a 10-fold serial dilution, and 10 μl

drops from the 10^{-1}–10^{-6} dilutions were placed on NB agar with cephalexin. *, *P* <

0.05; **, *P* < 0.01.

Figure 7. Proposed new inching switch mechanism that bacteria use to dynamically

regulate the iron homeostasis by A-to-I modification in *xfeA* adapt to shifty iron

concentration. Under iron-depleted or host plant environments, T408A editing level

within *xfeA* is enhanced, thus increasing the Fe-Ent uptake, and then rouse the

chemotaxis pathway genes to improve the sensitivity, thereby contributing to the

fitness of iron deficiency and virulence of the host plant. When *Xoc* under the

iron-depleted condition, the A-to-I modification level is reduced.

Appendixes

Figure S1. Normalized expression levels of selected chemotaxis pathway genes and

xfeA were calculated relative to rpoD using the ΔΔCT method in BLS256 wild-type

strain, where CT is the threshold cycle. Three independent biological replicates were

carried out in this study.

Figure S2. RNA secondary structure prediction of the T408A mutation in *Xoc* XfeA.

RNA secondary structure analysis (http://rna.tbi.univie.ac.at/) showed that the edited

site is embedded within a loop (see arrow).

Figure S3. The diagram of capillary chemotaxis test device and process.

Figure S4. Verifications of the *xfeA* T408A A-to-I RNA editing site by Sanger

sequencing (gDNA as the template). Chromatograms show that no A/G mixture exists

at the DNA level, and could exclude the possibility of DNA mutation in a

subpopulation that is further selected at experimental conditions.

Figure S5. Comparing of A-to-I RNA editing level between the RNA-seq data of

T408A and T408^{silent} strain in NB medium with two biological replicates. Because of

the DNA site-mutation, the editing level of *xfeA* is 100% and 0% in T408A and

T408^{silent} strain respectively (red frame), and verifying the construction of T408^{silent}

strain could block the A-to-I RNA editing of *xfeA*.

Table 1. Direct binding kinetics for MCPs binding $FeCl_3$ or Fe-Ent immobilized on Ni-NTA biosensors

Figure	Protein	Compound	Conc. (μM)	K_D (M)	K_{on} ($M^{-1} \cdot s^{-1}$)	K_{dis} (s^{-1})	R^2
a	Xoc_2282	$FeCl_3$	50	N.A.	N.A.	N.A.	0
b	Xoc_2291	$FeCl_3$	50	N.A.	N.A.	N.A.	0
c	Xoc_2282	Fe-Ent	50	$(4.88\pm1.02)\times10^{-6}$	$(5.21\pm0.08)\times10^{2}$	$(2.54\pm0.03)\times10^{-3}$	0.9962
d	Xoc_2291	Fe-Ent	50	$(8.74\pm2.35)\times10^{-6}$	$(2.78\pm0.04)\times10^{2}$	$(2.43\pm0.02)\times10^{-3}$	0.9335

Note: N.A. means the parameter is not available for the interaction combination.

a

T408A RNA editing level (%)

*** P < 0.001

b

DNA sequence ⋯ G T T G G T G C T A C G C A ⋯

50µM DP

20µM DP

NB

100µM FeCl₃

ΔtadA(NB)

* P < 0.05 ** P < 0.01

* P < 0.05 ** P < 0.01 *** P < 0.001