De novo **transcriptome assembly and functional annotation of cobia (*Rachycentron***

***canadum*): molecular approach as a base to nutritional studies**

David Aciole Barbosa [1] [#], Bruno C. Araújo [1] [#], Giovana S. Branco [2]; Alexandre W. S. Hilsdorf [1],

Luiz R. Nunes [3], Renata G. Moreira [2], Fabiano B. Menegidio [1]*

[1]Núcleo Integrado de Biotecnologia, Universidade de Mogi das Cruzes, Mogi das Cruzes, 08701-970,

SP, Brazil

[2]Departamento de Fisiologia, Instituto de Biociências, Universidade de São Paulo, São Paulo, 05508-

090, SP, Brazil

[3]Centro de Ciências Naturais e Humanas, Universidade Federal do ABC, São Bernardo do Campo,

09606-045, SP, Brazil

∗ To whom correspondence should be addressed: Fabiano B. Menegidio, PhD, Núcleo Integrado de

Biotecnologia, Universidade de Mogi das Cruzes (UMC), Av. Dr. Cândido X. de Almeida e Souza, 200

- Centro Cívico, Mogi das Cruzes - SP, 08780-911, Brazil;

E-mail: fabianomenegidio@umc.br

#. These authors contributed equally to this article

ABSTRACT: Cobia (*Rachycentron canadum*) is a marine teleost species with great productive potential worldwide. However, the genomic information currently available for this species in public databases is limited. This lack of information hinders gene expression assessments, which could bring forward novel insights into the physiology, ecology, evolution, and genetics of this aquaculture species. In this study, we report the first *de novo* transcriptome assembly of cobia liver to improve the availability of gene sequences from this important commercial fish. Thus, Illumina sequencing of liver transcripts generated 1,761,965,794 raw reads, which were filtered into 1,652,319,304 high-quality reads. *De novo* assembly resulted in 101,789 unigenes and 163,096 isoforms, with an average length of 950.61 and 1617.34 nt, respectively. Comparisons against six different databases resulted in functional annotation for 125,993 of these elements (77.3%), providing relevant information regarding the genomic content of *R. canadum*. We trust that the availability of these functionally annotated cobia elements will likely assist future nutrigenomics and breeding programs, involving this important fish farming species.

Keywords: *Rachycentron canadum*; Cobia; Transcriptome; Functional Annotation; Aquaculture; Liver.

1. Introduction

Currently, aquaculture is the most prominent food production industry, with significant growth worldwide. Global aquaculture production increased by 5.3% per year between 2001-2018, with a historical record of 114.5 million tons of farmed species, from which almost 17.7 million tons was due to finfish production. (FAO, 2020). Marine aquaculture plays an essential role in the effort of supplying the increased demand for animal-based protein. Cobia (*Rachycentron canadum*) is a carnivorous

marine fish species of worldwide distribution, and it is the sole representative of the Rachycentridae family. Currently, this species is regarded as the most promising marine fish species in Brazil, mostly due to its fast growth rate (reaching about 4 to 6 kg per year), excellent meat quality (excellent color, texture, and flavor), and high market value (Arnold *et al*., 2002; Benetti *et al*., 2008; Nunes, 2014). However, cobia production is still hindered by lacking of nutrition information, which prevents this species' productivity in industrial aquaculture operation (Fraser and Davies 2009).

Next-generation sequencing (NGS) studies have become an essential molecular tool in aquaculture, assisting in the production of several commercial fish species such as *Sparus aurata* (Calduch-Giner *et al.*, 2013), *Dicentrarchus labrax* (Magnanou *et al.*, 2014), and *Salmo salar* (Glencross *et al.*, 2015; Andrew *et al.*, 2020). *De novo* transcriptome assembly can be used in several different contexts, like genomics/gene expression analyses, which may be applied in many key areas of study, such as conservation genetics, selective breeding, reproductive biology, and nutrition (Leaver *et al.*, 2008; Calduch-Giner *et al.*, 2013; Fox *et al.*, 2014). For example, identifying genes associated with proteins and lipid metabolism in the liver can help the development of specific diets to improve the productive chain of commercial aquaculture species, including cobia. Unfortunately, genomic information in cobia is still scarce, limiting the development of such studies. Thus, to help overcome such limitations, the present manuscript describes an assembled/annotated reference transcriptome of hepatic cells in cobia juveniles. The information contained in this dataset contributes to improve our knowledge regarding the biological and physiological aspects of this fish species. Moreover, it establishes a solid foundation for future studies involving population genomics, breeding programs and nutrigenomics.

2. Data Description

2.1. Sample collection and RNA preparation

Ninety (90) cobia juveniles (128.85 ± 18.43 g) were obtained from a commercial hatchery (Redemar Alevinos, SP, Brazil) and randomly allocated in three 2.000 L tanks. The animals were kept under a mean temperature of 23±1.5 ºC and natural photoperiod throughout the trial, at the Marine Biology Center of the University of São Paulo (CEBIMar). Animals were equally hand-fed twice a day until apparent satiety with a commercial marine fish diet (Guabipirá, Guabi Nutrição e Saúde Animal S.A., SP, Brazil). After six weeks (42 days), fish were anesthetized with benzocaine concentration of 0.4 g $\square$ mL^{-1} and then euthanized by spinal cord section. Hepatic tissue samples from all experimental animals were collected, immediately frozen in liquid nitrogen, and subsequently stored at -80 °C for further analyses. Total RNA from hepatic tissue samples was extracted using PureLink® RNA Mini Kit (Invitrogen 12183018A), following the manufacturer's instructions. RNA samples had their concentration determined with a Nanodrop^{TM} Spectrophotometer (ThermoFisher) and the RNA integrity was determined by the 2100 Bioanalyzer (Agilent Technologies, CA, USA), only samples with RNA integrity number (RIN) above 6.5 were used. This study's experimental procedures were conducted according to the guidelines and approval of the Institutional Animal Care and Use Ethics Committee (#008/2017).

2.2. Library construction and sequencing

The construction of the library and the sequencing of the tissues were carried out at the GENIAL in Center for Research Support Facilities (CEFAP-USP), of the Institute of Biomedical Sciences of the University of São Paulo. RNA extracted from hepatic tissue samples from all animals were equally diluted to 1.000 ng µL^{-1} concentration and pooled for library construction using the TruSeq RNA

Sample Preparation kit, according to the manufacturer's specifications (Illumina Inc., USA). The library fragments were purified with the AM Pure XP system (Beckman Coulter, Beverly, MA, USA) and library quality was quantified by Qubit 2.0 before being subjected to sequencing Finally, paired-end sequencing (2×75 bp) of these cDNA libraries was performed on the NextSeq® equipment (Illumina, Inc., USA) to obtain the sequencing according to the manufacturer's recommendations. MIxS data (Minimum Information about any (x) Sequence) for this study is available in **Table 1.**

Information	Value
BioProject accession	PRJNA675281
BioProjects name	De novo transcriptome assembly and functional annotation of cobia (*Rachycentron canadum*) liver
BioSamples accession	SAMN16708758, SAMN16708759, SAMN16708760, SAMN16708761, SAMN16708762, SAMN16708763, SAMN16708764, SAMN16708765, SAMN16708766, SAMN16708767, SAMN16708768, SAMN16708769
SRAs accession	SRR13009897, SRR13009896, SRR13009895, SRR13009894, SRR13009893, SRR13009892, SRR13009891, SRR13009890, SRR13009889, SRR13009888, SRR13009887, SRR13009886
TSA accession	GIWT00000000
Sample Name	*Rachycentron canadum* Liver
Organism	*Rachycentron canadum*
Ecotype	"São Sebastião"
Collection date	01/jun/18
Broad-scale environmental context	ENVO_00000447 (marine biome)
Local-scale environmental context	ENVO_00001999 (marine water body)
Environmental medium	ENVO_00002010 (saline water)
Geographic location	Brazil:Sao Sebastiao,Sao Paulo
Latitude and longitude	23.858778 S 45.428833 W
Developmental stage	juvenile
Sequence method	Illumina Nextseq
Assembly method	Trinity
Annotation method	EnTAP, Diamond (RefSeq proteins, Nr proteins, Swiss-Prot, EggNOG)

Table 1. MIxS information for transcriptome assembly of *R. canadum*.

2.3. *De novo* assembly, functional annotation

FASTQ format raw sequencing data was processed in a Public Galaxy Server, available at

https://usegalaxy.eu. Initially, the quality of raw sequences was assessed using FastQC (Andrews,

2010) and MultiQC (Ewels *et al*., 2016). Fastp (Chen *et al*., 2018) was then used to remove low-quality

reads (Q<30), adapters, and other contaminant sequences. Trinity platform (Haas *et al.*, 2013) was then

used for *de novo* transcriptome assembly of filtered reads, and assembly metrics were obtained using

the TrinityStats script. Transcriptome completeness was finally assessed using the Benchmarking

Universal Single-Copy Orthologs (BUSCO) software (Seppey *et al*., 2019) based on OrthologDB

version 9 (Zdobnov *et al*., 2017).

Elements of the assembled transcriptome were functionally annotated by similarity searches

(blastn) performed against RefSeq RNA and the *"Lates/Seriola* database", a custom RefSeq transcripts

database containing sequences from cobia-related species, such as *Lates calcarifer, Seriola dumerili*,

and *Seriola lalandi dorsalis.* These sequences are also in the Kegg's Organisms Complete Genomes list

(available at https://www.genome.jp/kegg/catalog/org_list.html). Additional annotations were obtained

with the aid of the Eukaryotic Non-Model Transcriptome Annotation Pipeline (EnTAP) (Hart *et al.*,

2019).

Contigs were also queried (blastx) for homology against the National Center for Biotechnology

Information non-redundant protein database (NCBI nr), NCBI proteins reference database (RefSeq),

the curated Swiss-Prot database from UniProt Knowledgebase (UniProtKB) (UniProt Consortium,

2019) and the EggNOG proteins database (Huerta-Cepas *et al.*, 2016). The EggNOG hits also helped to

assign biological function to the putative cobia genes, identifying GO (Gene Ontology) (Ashburner *et*

al., 2000; Gene Ontology Consortium, 2019) and KEGG (Kyoto Encyclopedia of Genes and Genomes)

(Kanehisa & Goto, 2000; Kanehisa, 2019) terms. The EnTAP functional annotation process was carried

out using a Dugong container environment (Menegidio *et al.*, 2018). Additional transcriptome filtering

before NCBI submission was carried out with the aid of seqtk subseq from seqtk toolkit

(https://github.com/lh3/seqtk), based on sequence length (>200 nucleotides) and univec

(https://ftp.ncbi.nlm.nih.gov/pub/UniVec/).

2.4. Functional and structural characterization of transcriptome

Sequencing of the cDNA libraries derived from *R. canadum* liver material resulted in

1,761,965,794 raw reads. After high-quality reads selection and trimming, we were left with a total of

1,652,319,304 reads (93.77% of raw reads), which were used for *de novo* transcriptome assembly,

using Trinity software (Haas *et al.*, 2013). General features of the *R. canadum* liver transcriptome are

summarized in **Table 2**, consisting of 101,789 unigenes and 163,096 isoforms, displaying an average

length of 1,617.34 and 950.61 nt, respectively. The median length (N50) of these elements was 7,843 nt

for unigenes and 2,312 nt for isoforms. A total of 95,075 transcripts (58.29%) presented 500 nt or

more. Identification of 83.8% of the complete universal genes (3839 out of the total 4584 genes from

Actinopterygii odb9 lineage) supported the high quality and completeness of this transcriptome

assembly (**Fig. 1A**). Among the 3839 conserved BUSCO genes, 38.1% were single copy, while 45.7%

were duplicated (**Supplementary Table ST1-S11**).

Illumina sequencing	raw	high-quality
Number of reads	1,761,965,794	1,652,319,304

Assembly	unigenes	isoforms
Total sequences	101,789	163,096
N50	7,843	2,312
Average contig length (bp)	1,617.34	950.61
Median contig length (bp)	704	391
Total assembled bases	263,781,171	96,761,286
CG%	44.5	

Functional annotation	annotated sequences
NCBI nr	29,155
NCBI RefSeq RNA	122,741
Swiss-Prot	15,507
Lates/*Seriola*	118,570
GO-Biological Process	48,550
GO-Molecular Function	47,766
GO-Cellular Component	34,492
Kegg	24,100
Total	125,993

146   **Table 2.** Summary of the *de novo* transcriptome assembled for *R. canadum*.

Fig. 1. Completeness and homology search of *R. canadum* liver transcripts. (A) Percentage of completeness on the core
set of genes of *R. canadum* based on Actinopterygii database (orthodb9) from BUSCO. (B) Species annotation distribution
for the best hits from NCBI nr, NCBI RefSeq and Swiss Prot databases. (C) Gene ontology distribution for the 10 most
assigned biological process, cellular component and molecular function of assembled isoforms from *R. canadum* liver
transcriptome (Available in 10.6084/m9.figshare.13335698).

Transcriptome annotation against a series of databases (NCBI nr, NCBI RefSeq RNA,

SwissProt, GO, KEGG, *Lates*/*Seriola*) resulted in functional assignment for 125,993 transcripts

(77.3%). Most sequence homologies were found against NCBI RefSeq RNA (122,741 transcripts,

97.4%), followed by the *Lates*/*Seriola* database (118,570 transcripts, 94.1%), NCBI nr (29,155

transcripts, 23.14%), NCBI RefSeq (25,728, 20.42%) and Swiss-Prot (15,507 transcripts, 12.30%)

(**Table1**; **Supplementary Table ST1-S10**). Sequence homologies identified by EnTAP were

distributed throughout many bony fish species, of which *Seriola dumerili* was the most frequent (nr =

39.83%, refseq = 40.59%), followed by *Seriola lalandi dorsalis* (nr = 23,27%, refseq = 25.86%),

Echeneis naucrates (nr 17.9%, refseq = 18.37%) and *Larimichthys crocea* (nr = 4.99%, refseq =

3.13%) (**Fig. 1B**). Most transcripts (75,060) were functionally annotated with GO terms by eggNOG

(Huerta-Cepas *et al.*, 2016), which assigned 48,550 transcripts (65%) to biological processes, 34,492 to

cellular components (46%) and 47,766 to molecular functions (64%). The ten most representative

functional groups within each category are shown in **Fig. 1C**. A total of 24,100 isoforms were

annotated into at least one KEGG pathway term (**Table2**; **Supplementary Table ST1-S10**).

In summary, our findings provide an important tool for applied studies related to cobia

production. These outcomes will help to improve the knowledge regarding cobia physiology

consequently improving aquaculture operations leading to higher productivity.

2.5 Data availability

Sequencing raw data were deposited in the Sequence Read Archive (SRA) repository of the

National Center for Biotechnology Information (NCBI), under accession number SRR13009897,

SRR13009896, SRR13009895, SRR13009894, SRR13009893, SRR13009892, SRR13009891,

SRR13009890, SRR13009889, SRR13009888, SRR13009887 and SRR13009886, associated to the

BioProject numbers PRJNA675281 and BioSamples numbers SAMN16708758, SAMN16708759,

SAMN16708760, SAMN16708761, SAMN16708762, SAMN16708763, SAMN16708764,

SAMN16708765, SAMN16708766, SAMN16708767, SAMN16708768 and SAMN16708769. The

Transcriptome Shotgun Assembly (TSA) project has been deposited at DDBJ/EMBL/GenBank under

accession number GIWT00000000. The version described in this paper is the first version,

GIWT00000000.1. Supplementary Table S1 is available from the Figshare repository

(10.6084/m9.figshare.13242176). Additional data derived from this study (including all intermediate

data) are also available from the Open Science Framework (OSF) repository (DOI:

10.17605/OSF.IO/BV3WA).

Authors contributions

B.C.A. sampled the specimens. B.C.A., G.S.B., AWSH and R.G.M. performed molecular

analyses and sequencing. D.A.B., L.R.N. and F.B.M. assembled and evaluated the transcriptome. All

authors wrote the paper. All authors read and approved the final version of the manuscript.

Declaration of Competing Interest

The authors report no conflicts of interest. The authors alone are responsible for the content and

the writing of the paper.

Acknowledgements

This study was financed in part by the Fundação de Amparo à Pesquisa do Estado de São Paulo

(FAPESP) (grant # 2019/26018-0) and Conselho Nacional de Desenvolvimento Científico e

Tecnológico (CNPq) (grant # 305493/2019-1). D.A.B. and B.C.A. are recipients of scholarship grants

from Coordenação de Aperfeiçoamento de Pessoal de Nível Superior–Brasil-CAPES

(http://www.capes.gov.br/) (grant # 88882-365823/2019-01) and FAPESP (grant # 2016/12435-0),

respectively. A.W.S. Hilsdorf and R.G. Moreira are recipients of CNPq productivity scholarships

(grants # 304662/2017-8 and 305493/2019-1, respectively). We are grateful to the Facility GENIAL

(Genome Investigation and Analysis Laboratory) - Center for Research Support Facilities (CEFAP-

USP), of the Institute of Biomedical Sciences of the University of São Paulo.

206 Appendix A. Supplementary data

207   Supplementary data to this article can be found online at Open Science Framework:
208 https://osf.io/bv3wa/.

210 References

Ashburner, M., Ball, C.A., Blake, J.A., Botstein, D., Butler, H., Cherry, J.M., Davis, A.P., Dolinski, K.,

Dwight, S.S., Eppig, J.T. and Harris, M.A., 2000. Gene ontology: tool for the unification of

biology. Nat. Genet 25(1), 25-29. https://doi.org/10.1038/75556.

Andrew, S.C., Primmer, C.R., Debes, P.V., Erkinaro, J. and Verta, J.P., 2020. The Atlantic salmon

whole blood transcriptome and how it relates to major locus maturation genotypes and other

tissues. Mar. Genom., p.100809. https://doi.org/10.1016/j.margen.2020.100809.

Andrews S., 2010. FastQC: a quality control tool for high throughput sequence data. Available online

at: http://www.bioinformatics.babraham.ac.uk/projects/fastqc.

Arnold, C.R., Kaiser, J.B., Holt, G.J., 2002. Spawning of cobia *Rachycentron canadum* in captivity. J.

World Aquacult. Soc., *33*(2), 205-208. https://doi.org/10.1111/j.1749-7345.2002.tb00496.x.

Benetti, D.D., Orhun, M.R., Sardenberg, B., O'Hanlon, B., Welch, A., Hoenig, R., Zink, I., Rivera,

J.A., Denlinger, B., Bacoat, D., Palmer, K., 2008. Advances in hatchery and grow□out

technology of cobia *Rachycentron canadum* (Linnaeus). Aquac. Res. 39(7), 701-711.

https://doi.org/10.1111/j.1365-2109.2008.01922.x.

Calduch-Giner, J.A., Bermejo-Nogales, A., Benedito-Palos, L., Estensoro, I., Ballester-Lozano, G.,

Sitjà-Bobadilla, A., Pérez-Sánchez, J., 2013. Deep sequencing for de novo construction of a

marine fish (*Sparus aurata*) transcriptome database with a large coverage of protein-coding

transcripts. BMC genomics, 14(1),c178. https://doi.org/10.1186/1471-2164-14-178.

Chen, S., Zhou, Y., Chen, Y. and Gu, J., 2018. fastp: an ultra-fast all-in-one FASTQ preprocessor.

Bioinformatics, 34(17), i884-i890. https://doi.org/10.1093/bioinformatics/bty560

Ewels, P., Magnusson, M., Lundin, S., Käller, M., 2016. MultiQC: summarize analysis results for

multiple tools and samples in a single report. Bioinformatics, 32(19), 3047-3048.

https://doi.org/10.1093/bioinformatics/btw354.

Fox, S.E., Christie, M.R., Marine, M., Priest, H.D., Mockler, T.C. and Blouin, M.S., 2014. Sequencing

and characterization of the anadromous steelhead (Oncorhynchus mykiss) transcriptome. Mar.

Genom. 15, 13-15. doi: 10.1016/j.margen.2013.12.001.

FAO. 2020. The State of World Fisheries and Aquaculture 2020. Sustainability in action. Rome. 244 p.

Fraser, T.W. and Davies, S.J., 2009. Nutritional requirements of cobia, *Rachycentron canadum*

(Linnaeus): a review. Aquac. Res. 40(11), 1219-1234. https://doi.org/10.1111/j.1365-

2109.2009.02215

Gene Ontology Consortium, 2019. The gene ontology resource: 20 years and still GOing strong.

Nucleic Acids Res., 47(D1), d330-d338. https://doi.org/10.1093/nar/gky1055.

Glencross, B.D., De Santis, C., Bicskei, B., Taggart, J.B., Bron, J.E., Betancor, M.B. and Tocher, D.R.,

2015. A comparative analysis of the response of the hepatic transcriptome to dietary

docosahexaenoic acid in Atlantic salmon (*Salmo salar*) post-smolts. BMC genomics, 16(1), 684.

https://doi.org/10.1186/s12864-015-1810-z.

Haas, B.J., Papanicolaou, A., Yassour, M., Grabherr, M., Blood, P.D., Bowden, J., Couger, M.B.,

Eccles, D., Li, B., Lieber, M., MacManes, M.D., 2013. De novo transcript sequence

reconstruction from RNA-seq using the Trinity platform for reference generation and analysis.

Nat. Protoc 8(8), 1494-1512. https://doi.org/10.1038/nprot.2013.084.

Hart, A.J., Ginzburg, S., Xu, M., Fisher, C.R., Rahmatpour, N., Mitton, J.B., Paul, R., Wegrzyn, J.L.,

2020. EnTAP: bringing faster and smarter functional annotation to non☐model eukaryotic

transcriptomes. Mol. Ecol. Resour. 20(2), 591-604. https://doi.org/10.1111/1755-0998.13106.

Huerta-Cepas, J., Szklarczyk, D., Forslund, K., Cook, H., Heller, D., Walter, M.C., Rattei, T., Mende,

D.R., Sunagawa, S., Kuhn, M., Jensen, L.J., 2016. eggNOG 4.5: a hierarchical orthology

framework with improved functional annotations for eukaryotic, prokaryotic and viral sequences.

Nucleic Acids Res. 44(D1), D286-D293. https://doi.org/10.1093/nar/gkv1248.

Kanehisa, M. and Goto, S., 2000. KEGG: kyoto encyclopedia of genes and genomes. Nucleic Acids

Res, 28(1), 27-30. https://doi.org/10.1093/nar/28.1.27.

Kanehisa, M., 2019. Toward understanding the origin and evolution of cellular organisms. Protein Sci.

28(11), 1947-1951. https://doi.org/10.1002/pro.3715.

Leaver, M.J., Bautista, J.M., Björnsson, B.T., Jönsson, E., Krey, G., Tocher, D.R., Torstensen, B.E.,

2008. Towards fish lipid nutrigenomics: current state and prospects for fin-fish aquaculture. Rev

Fish Sci. 16, 73-94. https://doi.org/10.1080/10641260802325278.

Magnanou, E., Klopp, C., Noirot, C., Besseau, L., Falcón, J., 2014. Generation and characterization of
the sea bass Dicentrarchus labrax brain and liver transcriptomes. Gene, 544(1), 56-66.
https://doi.org/10.1016/j.gene.2014.04.032.

Menegidio, F.B., Jabes, D.L., Costa de Oliveira, R. and Nunes, L.R., 2018. Dugong: a Docker image,
based on Ubuntu Linux, focused on reproducibility and replicability for bioinformatics analyses.
Bioinformatics, 34(3), 514-515. https://doi.org/10.1093/bioinformatics/btx554.

Nunes, A.J.P. 2014. Ensaios com o beijupirá, *Rachycentron canadum*. Fortaleza: Ministério da Pesca e
Aquicultura/CNPQ/UFC, 2014. 352 p.

Seppey, M., Manni, M. and Zdobnov, E.M., 2019. BUSCO: assessing genome assembly and annotation
completeness. In Gene Prediction (pp. 227-245). Humana, New York, NY.
https://doi.org/10.1007/978-1-4939-9173-0_14.

UniProt Consortium, 2019. UniProt: a worldwide hub of protein knowledge. Nucleic Acids Res.
47(D1), D506-D515. https://doi.org/10.1093/nar/gky1049.

Zdobnov, E.M., Tegenfeldt, F., Kuznetsov, D., Waterhouse, R.M., Simao, F.A., Ioannidis, P., Seppey,
M., Loetscher, A., Kriventseva, E.V., 2017. OrthoDB v9. 1: cataloging evolutionary and
functional annotations for animal, fungal, plant, archaeal, bacterial and viral orthologs. Nucleic
Acids Res, 45(D1), D744-D749. https://doi.org/10.1093/nar/gkw1119.

A - Actinopterygii genes completeness

C:83.8%[S:38.1%,D:45.7%],F:4.4%,M:11.8%,n:4584

Complete BUSCOs (C)	3839
Fragmented BUSCOs (F)	200
Missing BUSCOs (M)	545
Complete and single-copy BUSCOs (S)	1746
Complete and duplicated BUSCOs (D)	2093
Total BUSCO groups searched	4584

Missing 11.8%
Complete 83.8%
Fragmented 4.4%
Duplicated 45.7%
Single-copy 38.1%

B - Species annotation distribution

NCBI nr
39.83%, 23.27%, 17.09%, 4.99%, 3.42%, 2.67%, 2.38%, 2.27%, 1.69%, 0.84%, 0.78%, 0.39%, 0.37%

Seriola dumerili
Seriola lalandi dorsalis
Echeneis naucrates
Larimichthys crocea
Scophthalmus maximus
Lates calcarifer
Epinephelus lanceolatus
Morone saxatilis
Perca flavescens
Acanthochromis polyacanthus
Amphiprion ocellaris
Anabas testudineus
Mastacembelus armatus

NCBI Refseq
40.59%, 25.86%, 18.37%, 3.13%, 2.57%, 2.03%, 1.88%, 1.66%, 1.41%, 1.14%, 0.34%, 1.01%

Seriola dumerili
Seriola lalandi dorsalis
Echeneis naucrates
Lates calcarifer
Epinephelus lanceolatus
Mastacembelus armatus
Larimichthys crocea
Anabas testudineus
Acanthochromis polyacanthus
Amphiprion ocellaris
Acinetobacter baumannii
Sparus aurata

Swiss-Prot
37.31%, 21.35%, 17.23%, 5.78%, 5.73%, 4.56%, 3.58%, 1.77%, 1.68%, 0.6%, 0.25%, 0.17%

Homo sapiens
Mus musculus
Danio rerio
Rattus norvegicus
Bos taurus
Gallus gallus
Xenopus laevis
Xenopus tropicalis
Pongo abelii
Sus scrofa
Macaca fascicularis
Takifugu rubripes

C - Top 10 GO annotation distribution

Biological Process

GO term	Transcripts
GO:0008150−biological process(L=0)	48550
GO:0009987−cellular process(L=1)	34189
GO:0008152−metabolic process(L=1)	30402
GO:0044699−single−organism process(L=1)	24047
GO:0071704−organic substance metabolic process(L=2)	23008
GO:0044238−primary metabolic process(L=2)	22316
GO:0044763−single−organism cellular process(L=2)	21075
GO:0044237−cellular metabolic process(L=2)	20914
GO:0065007−biological regulation(L=1)	18430
GO:0050789−regulation of biological process(L=2)	14125
GO:0043170−macromolecule metabolic process(L=3)	3366

Cellular Component

GO term	Transcripts
GO:0005575−cellular component(L=0)	34492
GO:0044464−cell part(L=2)	24115
GO:0005623−cell(L=1)	24115
GO:0005622−intracellular(L=3)	22270
GO:0044424−intracellular part(L=3)	19598
GO:0043226−organelle(L=1)	15984
GO:0043229−intracellular organelle(L=3)	15941
GO:0016020−membrane(L=1)	14091
GO:0043227−membrane−bounded organelle(L=2)	13250
GO:0043231−intracellular membrane−bounded organelle(L=4)	13213

Molecular Function

GO term	Transcripts
GO:0003674−molecular function(L=0)	47766
GO:0005488−binding(L=1)	31915
GO:0003824−catalytic activity(L=1)	24493
GO:0043167−ion binding(L=2)	23300
GO:0097159−organic cyclic compound binding(L=2)	19825
GO:1901363−heterocyclic compound binding(L=2)	19734
GO:0043169−cation binding(L=3)	13872
GO:0046872−metal ion binding(L=4)	13787
GO:0043168−anion binding(L=3)	11811
GO:0036094−small molecule binding(L=2)	9094
GO:0003676−nucleic acid binding(L=3)	2150

Transcripts