1

A secreted endoribonuclease ENDU-2 from the soma protects germline immortality in *C.*

** *elegans***

Wenjing Qi[1,#], Erika D v. Gromoff[1], Fan Xu[1], Qian Zhao[1], Wei Yang[1], Dietmar Pfeifer[4], Wolfgang

Maier[1], Lijiang Long[1] and Ralf Baumeister[1,2,3]

[1] Bioinformatics and Molecular Genetics (Faculty of Biology), University of Freiburg, Germany

[2] Center for Biochemistry and Molecular Cell Research (Faculty of Medicine), University of

Freiburg, Germany

[3] Signalling Research Centers BIOSS and CIBSS

[4] Department of Hematology, Oncology and Stem Cell Transplantation, Medical Center University

of Freiburg, Freiburg, Germany

[#] Correspondence: wenjing.qi@biologie.uni-freiburg.de

Bio3/Bioinformatics and Molecular Genetics, University of Freiburg, Schänzlestrasse 1, D-79104

Freiburg, Germany

Phone: +49-761-203 2728; Fax: +49-761-203 8352

Abstract

Multicellular organisms coordinate tissue specific response to environmental information via both

cell-autonomous and non-autonomous mechanisms. In addition to secreted ligands, secreted small

RNAs have recently been reported to regulate gene expression across tissue boundaries. Here we

show that the conserved poly-U specific endoribonuclease ENDU-2 is secreted from the soma and

taken-up by the germline to ensure germline immortality at elevated temperature in *C. elegans*.

ENDU-2 binds to mature mRNAs and negatively regulates mRNA abundance both in the soma

and the germline. While ENDU-2 promotes RNA decay in the soma directly via its

endoribonuclease activity, ENDU-2 prevents misexpression of soma-specific genes in the

germline and preserves germline immortality independent of its RNA-cleavage activity. In

summary, our results suggest that the secreted RNase ENDU-2 transmits environmental

information across tissue boundaries and contributes to maintenance of stem cell immortality

probably via retaining a stem cell specific program of gene expression.

Introduction

Germ cells are the only type of cells within an organism that deliver genetic and epigenetic material to the offspring. Germ cells are therefore distinct from the somatic cells in their ability to maintain totipotency to produce an entire organism upon fertilization and their immortality to allow reproduction for unlimited future generations. In *C. elegans*, loss of germline immortality typically results in sterility of initially fertile animals after a number of generations. To date, telomerase-mediated telomere maintenance, balanced heterochromatic H3K9 methylation and euchromatic H3K4/H3K36 methylation are the prevalent known molecular mechanisms involved in maintenance of germline immortality (Ahmed and Hodgkin, 2000; Katz et al., 2009; Lev et al., 2017; Weiser et al., 2017). Piwi RNA mutants and mutations in multiple components of a nuclear RNA interference (RNAi) pathway that promotes the inheritance of germline RNAi also result in a temperature-dependent mortal germline (Mrt) phenotype (Buckley et al., 2012; Simon et al., 2014). Failure to maintain germline immortality in Piwi RNA mutants has recently been explained by transgenerational silencing of histone genes (Barucci et al., 2020). These studies, however, focused almost exclusively on autonomous mechanisms in the germline. It is unknown whether somatic signaling may contribute to maintenance of germline immortality.

C. elegans ENDU-2 belongs to a conserved but less studied family of proteins containing proposed poly-U specific endoribonuclease (XendoU) domains. While human EndoU (PP11 placental protein 11) is used as a cancer marker gene, Viral EndoU Nsp15 is highly conserved in all known coronavirus and has been suggested to promote viral RNA replication and limit innate immunity response of the host cells (Hackbart et al., 2020; Ivanov et al., 2004). *C. elegans* ENDU-2 has been shown to regulate cold stress response (Ujisawa et al., 2018). Very recently, ENDU-2 is reported

to regulate nucleotides metabolism and germline proliferation in response to changes in nucleotide

levels and genotoxic stresses (Jia et al., 2020).

Here we report that secretion of the poly-U specific endoribonuclease ENDU-2 from the soma to

the germline preserves germline immortality at elevated temperature. We find that ENDU-2 binds

to mature mRNAs and down-regulates mRNA levels both in the soma and the germline. In

addition, RNA-binding and -cleavage are two separable activities utilized by ENDU-2 to control

gene expression via distinct mechanisms. In the soma, ENDU-2 relies on its endoribonuclease

activity to negatively regulate part of its mRNA targets. In the germline, ENDU-2 prevents

misexpression of soma-specific genes and ensures stem cell immortality primarily only via its

RNA-binding activity, suggesting an essential role of ENDU-2 in retaining the stem cell specific

program of gene expression. In summary, our data implicates that the soma sends ENDU-2 as a

messenger to the germline to control gene expression across tissue boundaries in response to

temperature alteration.

Loss of *endu-2* causes a temperature dependent Mrt phenotype

Freshly outcrossed *endu-2(lf)* mutants were similar to wild type animals in the first few generations, except slightly reduced germline proliferation, an egg-laying defect (Egl) due to abnormal development of the vulva, and a moderate reduction in adult lifespan at 20°C (Supplemental Fig. S1). Long-term strain maintenance at 20°C was difficult, since the *endu-2(lf)* mutants showed gradually increased sterility that, however, could be reset by new outcrosses with wild type animals. We examined the fertility of two *endu-2* alleles *tm4977* and *by188* over generations after four additional out-crosses. *tm4977* allele has a 620 bp deletion starting from the 5'UTR to the end of the third intron, while the 20 bp deletion in *by188* causes an early stop codon in the first exon. Therefore, these two alleles are probably null mutants. Both *tm4977* and *by188* alleles became sterile at 20°C after about 15-20 generations (Fig. 1A, Supplemental Fig. S2A). Since we did not observe a significant reduction of the number of germ nuclei in the mitotic zone across generations, sterility was not the consequence of declining germline proliferation (Supplemental Fig. S2B). Instead, *endu-2* day 1 adult animals in the generations with highly penetrant sterile phenotype showed pleiotropic defects in the germline. The most prominent defects were abnormal cell death (38%, n=105) and increased number of apoptotic corpses (31% with ≥ 2 corpses per gonad arm, n=105) (Fig. 1B). We also observed ongoing spermatogenesis 24 hours after mid-L4 stage (11%, n=105) whereas wild type animals had completely switched spermatogenesis to oogenesis, suggesting prolonged spermatogenesis. Furthermore, endomitosis occurred in the mitotic region (14%, n=25) (Fig. 1B). In the generations displaying strong sterile phenotype, *endu-2(lf)* additionally showed high incidence of male (Him, 14%, n=218) and increased occurrence of other phenotype aspects (Rol, Dpy, Sma phenotypes, in total 10%, n=218). The latter was probably caused by elevated somatic mutation rates, since these phenotypes were

not heritable. The sterile phenotype was 100% penetrant at 25°C already at generation 6-10, but

was not detected at 15°C (Supplemental Fig. S2C and S2D). Notably, the highly penetrate sterility

could be fully reversed within several generations by transferring the animals to 15°C (Fig. 1A).

Taken together, these results suggest an essential role of ENDU-2 in preserving germline

immortality at elevated temperature and existence of additional mechanisms to compensate for the

loss of *endu-2* at 15°C.

ENDU-2 is a secreted protein

A previous report has suggested a wide-spread expression of *endu-2* in somatic tissues (Ujisawa

et al., 2018), but did not report ENDU-2 localization in the germline. To retest the expression

pattern of ENDU-2, we generated several independent transcriptional, translational EGFP fusion

endu-2 reporters and a CRISPR/Cas9 EGFP knock-in strain at the endogenous *endu-2* locus

(expression constructs are shown in the Supplemental Fig. S3A). The transcriptional fusion

reporter gene, harboring 4 kb of *endu-2* upstream sequences, was expressed only in the intestine

(Supplemental Fig. S3B). A translational reporter harboring the same upstream sequences, and

also the entire genomic region of *endu-2*, fused to EGFP, rescued Egl, reduced germline

proliferation, and short lifespan phenotypes (Supplemental Fig. S1A, S1C and S1D). The

CRISPR/Cas9 EGFP knock-in strain displayed neither Mrt nor Egl phenotypes (Supplemental Fig.

S1E). These results suggest that ENDU-2::EGFP fusion proteins in the transgenic and the EGFP

knock-in strains are functional. We observed ENDU-2::EGFP protein in the cytoplasm of intestine,

somatic gonad and coelomocytes in both *endu-2::EGFP* transgenic and the CRISPR/Cas9 EGFP

knock-in strains (Fig. 2A and Supplemental Fig. S3C). Different from the study of Ujisawa et al.

(2018), we did not detect ENDU-2::EGFP in the musculature or in the neurons. Instead, we noticed

extracellularly localized ENDU-2::EGFP, such as in the interspace between uterine wall and

embryos (Fig. 2A and Supplemental Fig. S3C), suggesting that ENDU-2 may be a secreted protein.

ENDU-2 indeed contains a predicted N-terminal (1-19 amino acid) secretion signal peptide for

endoplasmic reticulum (ER) targeting, indicating that the protein is destined towards the secretory

pathway. Fusion of this secretion signal peptide to the N-terminus of a neuron specific expressed

EGFP reporter led to strongly decreased expression level of EGFP in neurons accompanied by

EGFP signal in extracellular space and coelomocytes, which endocytose secreted proteins existing

in the pseudocoelic fluid (Fig. 2B). In addition, removing the N-terminal secretion signal peptide

of ENDU-2 (Δ_{ss}ENDU-2::EGFP) resulted in strong expression of Δ_{ss}ENDU-2::EGFP only in the

intestine but not in the coelomocytes (Supplemental Fig. S3E and S4A). These results together

suggest that the secretion signal peptide composed of the first 19 amino acids of ENDU-2 is

necessary and sufficient to trigger secretion of a protein. Strikingly, we noticed that another

transgenic reporter expressing 3xFlag::ENDU-2::EGFP was expressed strongly in the intestinal

and also weakly in some head neurons, muscle cells in the head region and anal depressor muscle

cells, corroborating the report from Ujisawa et.al (Supplemental Fig. S3D). We speculate that the

N-terminal 3xFlag fusion may prevent secretion by impairing binding of the secretion signal

peptide by signal recognition particle (SRP), thus allowing detection of the weak expression in the

muscle and neuron. In addition, expression in the head neurons and muscle cells might possibly

be controlled by promoter signals localized in the first intron since this was the only sequence

absent in the transgene expressing $\Delta_{ss}endu-2::EGFP$ (Supplemental Fig. S3A), in which we failed

to observe muscle or neuronal localization. Moreover, when we expressed *endu-2::EGFP*

selectively either in the neurons (*unc-119* promoter), muscles (*myo-3* promoter) or intestine (*vha-*

6 promoter), ENDU-2::EGFP was always detected in the coelomocytes (Supplemental Fig. S4A),

indicative for its secretion from these tissues. Notably, expressing *endu-2::EGFP* specifically in

either neurons or muscles of heat-stressed animals resulted in ENDU-2::EGFP localization in the

pharynx (Supplemental Fig. S4B), suggesting that either secretion or uptake of ENDU-2 in these

tissues could be modulated by temperature.

Secretion of ENDU-2 from the soma to the gonad protects germline immortality

It is generally accepted that multi-copy extrachromosomal arrays are silenced and not expressed

in the *C. elegans* germline (Kelly et al., 1997). However, we found that extrachromosomal *endu-*

2::EGFP transgenes rescued the Mrt germline phenotype of *endu-2(lf)* animals (Fig. 3A),

suggesting that somatic ENDU-2 might preserve germline immortality across tissue boundaries.

Our hypothesis was that secreted ENDU-2 could be endocytosed by the gonad. To test this

assumption, we first tested whether ENDU-2 protein could be detected in the germline. All of our

endu-2::EGFP transgenic reporters displayed weak expression level unless secretion was blocked.

Therefore, we could only occasionally observe faint ENDU-2::EGFP in few oocytes

(Supplemental Fig. S4C). Via GFP antibody staining we detected ENDU-2::EGFP in punctate

structures in the germline at 15°C, 20°C and 25°C (Fig. 2C and Supplemental Fig. S5). In contrast,

Δ_{ss}ENDU-2::EGFP was not detectable in the germline despite of its high expression level

(Supplemental Fig. S3F). And fusion the secretion signal peptide of another secreted protein SEL-

1 (Grant and Greenwald, 1996) to Δ_{ss}ENDU-2::EGFP reactivated secretion and its localization in

the germline (Fig. 2C and Supplemental Fig. S3E). These observations indicate that secretion of

ENDU-2 is indispensable for its gonadal localization. Moreover, we performed single molecular

FISH (smFISH) staining to visualize *endu-2* mRNA. *endu-2* mRNA signals were visible

predominately in the intestine but clearly absent in the germline (Fig. 2D). Furthermore,

sequencing of RNA extracted from isolated gonads also revealed absence of *endu-2* mRNA in the

wild type germline (Supplemental Table S3). Taken together, these results implicate that

somatically expressed ENDU-2 protein can enter the gonad via its secretion and uptake.

Next, we asked from which tissue ENDU-2 ensures germline immortality. Expressing ENDU-2

specifically in the intestine or the neurons, but not the muscle or somatic gonad, were sufficient to

rescue the Mrt phenotype, indicating a non-cell-autonomous ENDU-2 signal from the neurons and

intestine to the germline (Fig. 3B). In addition, expressing the secretion deficient Δ_{ss}ENDU-

2::EGFP failed to rescue the Mrt phenotype (Fig. 3C). Furthermore, SS_{sel-1}:: Δ_{ss}ENDU-2::EGFP

rescued the Mrt phenotype (Fig. 3C), suggesting that guiding ENDU-2 into ER-Golgi secretory

pathway rather than the first 19 amino acids of ENDU-2 is essential for its function in the germline.

To directly test whether loss of somatic *endu-2* expression is sufficient to induce a mortal germline,

we performed *endu-2* RNAi knock-down in a *ppw-1* mutant background in which germline RNAi

does not function (Tijsterman et al., 2002). Although neither wild type nor *ppw-1* mutants upon

endu-2 RNAi displayed a fully penetrant Mrt phenotype within 15 generation, *endu-2* RNAi,

resulted in gradually reduced brood sizes over generations in both *ppw-1* and wild type

background, indicating that somatically expressed *endu-2* mRNA is required for normal

reproduction (Supplemental Fig. S6A). The failure to obtain a fully penetrant Mrt phenotype in

this experiment may be due to the relative inefficiency of RNAi in the nervous system that,

expresses *endu-2* to ensure germline immortality.

ENDU-2 affects the oocytes to preserve germline immortality independent of the nuclear

RNAi pathway

To know whether ENDU-2 affects oocytes or sperm to ensure germline immortality, we tested the

parental contribution of ENDU-2 during sexual reproduction. For this purpose, we crossed *endu-*

2(+/+) wild type parents with either males or hermaphrodites of *endu-2(-/-)* that had been grown

at 25°C for five to seven generations and displayed strongly reduced brood size and high

percentage of sterile phenotype (Fig. 3D). Whereas the heterozygous progeny of *endu-2(+/+)*

mothers had brood sizes with typically more than 100 F2 animals, the brood size of F1 cross-

progeny derived from *endu-2(-/-)* mothers was similarly low as that of their mothers. We conclude

that ENDU-2 function is required predominantly in the oocytes to preserve germline immortality.

The temperature dependent Mrt phenotype in *endu-2* mutants resembles that of *hrde-1* mutants in

the nuclear RNAi pathway (Buckley et al., 2012; Ni et al., 2016). In addition to loss of germline

immortality, *hrde-1* is also defective in multigenerational inheritance of germline RNAi. To know

whether ENDU-2 acts in the nuclear RNAi pathway, we examined *oma-1* RNAi inheritance, which

leads to suppression of the embryonic lethality of the *oma-1* gain-of-function mutant for several

generations (Alcazar et al., 2008). Unlike the *hrde-1* mutants that lost inheritance of *oma-1* RNAi

within two generations, effect of *oma-1* RNAi knock-down persisted for 5-6 generations both in

endu-2(lf) and wild type animals (Supplemental Fig. S6B), suggesting that ENDU-2 does not act

in the nuclear RNAi pathway to control transgenerational inheritance of germline RNAi.

Therefore, maintenance of germline immortality by ENDU-2 probably functions via a distinct

mechanism from that of the nuclear RNAi pathway.

mRNA-binding, but not mRNA-cleavage, by ENDU-2 is essential for maintaining germline

immortality

ENDU-2 harbors two XendoU domains, of which the C-terminally localized one is more similar

to both human EndoU and *Xenopus* XendoU (Supplemental Fig. S7). It had been suggested

recently that ENDU-2 might also be an RNA-binding protein (Ujisawa et al., 2018). To identify

the candidate RNAs associated with ENDU-2, we precipitated ENDU-2::EGFP and analyzed co-

immunoprecipitated RNAs by deep-sequencing (RIP-Seq). We also reasoned that, since wild type

ENDU-2 would potentially cleave its RNA targets, this might prevent enrichment of intact RNA

targets for identification. Although we had no direct proof of an RNA cleaving activity of ENDU-

2 yet, we thought that generation of an ENDU-2 variant that maintains RNA-binding, but loses

cleavage activity, should facilitate RNA target detection. *Xenopus* XendoU mutants with E to Q

exchanges at positions 161 or 167 of the EndoU domain lose RNA-cleavage without reducing

RNA-binding activity (Gioia et al., 2005). The second glutamic acid (167E) is conserved in both

XendoU domains of *C. elegans* ENDU-2 (175E and 460E), whereas the first (161E) is only found

in the second XendoU domain (454E) (Supplemental Fig. S7B). We expressed both ENDU-

2(E454Q)::EGFP and ENDU-2(E460Q)::EGFP variants in *endu-2(lf)* background and performed

RIP-Seq with these two strains in additional to wild type ENDU-2::EGFP. By plotting normalized

reads (RPM) of each transcript we were able to investigate RNA binding affinity of ENDU-2 under

different conditions. In general, all three ENDU-2 variants tested showed stronger RNA-binding

affinity at 15°C than at 25°C (Fig. 4A). E460Q showed weaker RNA binding already at 15°C than

wild type ENDU-2 and almost completely lost RNA-binding capacity at 25°C. In addition, ENDU-

2(E454Q) displayed stronger RNA binding activity than ENDU-2(wt) at 15°C. Therefore, we used

RIP-Seq data of ENDU-2(E454Q) at 15°C for detecting RNAs bound by ENDU-2. A total of 5,920

transcripts were co-immunoprecipitated with ENDU-2(E454Q) (Supplemental Table S1). Most of

them (>99%) were protein coding transcripts except few non-coding RNAs (5 snoRNAs, 27

pseudogenes, 5 ncRNAs and 1 lincRNA). In addition, reads distribution analysis showed > 99%

of sequenced reads were mapped to exonic regions, suggesting that ENDU-2(E454Q) primarily

bound to processed mRNAs (Fig. 4B). We performed *in vitro* mRNA binding assays with

recombinant ENDU-2 proteins and two selected mRNA targets from the RIP-Seq data and could

confirm that both ENDU-2 and ENDU-2(E454Q) directly interacts with these mRNAs (Fig. 4C).

Studies of *Xenopus* XendoU have shown that that the RNA hydrolysis activity of XendoU requires

Mn^{2+} or Ca^{2+} (Schwarz and Blower, 2014). ENDU-2 seemed to have a similar requirement, since

the addition of 5 mM Mn^{2+} or Ca^{2+} in the buffer medium, but not 5 mM Mg^{2+} led to degradation

of bulk RNA in the wild type worm lysates (Fig. 4D and Supplemental Fig. S8). Bulk RNA

degradation was strongly reduced in *endu-2(lf)* animals. In addition, expression of the wild type

endu-2::EGFP transgene, but not *endu-2(E454Q)::EGFP*, restored RNA decay in *endu-2(lf)*

mutants, supporting our hypothesis that ENDU-2(E454Q) lost RNA-cleavage capability despite

of the increased RNA-binding. Surprisingly, extrachromosomal expression of ENDU-2(E454Q),

but not of ENDU-2(E460Q), fully rescued the Mrt phenotype of *endu-2(lf)* at 25°C (Fig. 4E). We

conclude that mRNA-binding, rather than -cleavage activity of ENDU-2, is essential in the

germline to maintain stem cell immortality at elevated temperature.

ENDU-2 negatively regulates somatic mRNA abundance via its endoribonuclease activity

To test if ENDU-2 affects mRNA level, we performed microarray experiments at 25°C to

determine transcriptomic alterations mediated by ENDU-2 (procedure of sample preparation

illustrated in the Supplemental Fig. S9A). A comparison of differential gene expression in *endu-*

2(-) and *endu-2(+)* backgrounds suggested a similar numbers of transcripts as up- and down-

regulated by *endu-2* (fold-change >2) (Fig. 5A and Supplemental Tab. S2). Among the 258

transcripts down-regulated in *endu-2(lf)* animals, germline expressed genes (n=206) were over-

represented (Supplemental Table S2). qPCR quantifications of two selected genes *cav-1* and *trsc-*

1 confirmed our microarray result and revealed that their expression levels strongly decreased in

endu-2 mutants compared with wild type animals at 25°C (Supplemental Fig. S10C and S10D).

However, neither a germline expressed *cav-1::GFP* reporter nor smFISH staining of *trcs-1*

confirmed significantly altered germline expression *in situ* (Supplemental Fig. S10A, S10B and

S10D). As the gonads of *endu-2(lf)* mutants at 25°C became significantly smaller (Supplemental

Fig. S9B), we speculate that apparent down-regulation of germline mRNA within the

transcriptome data set may be caused by reduced sample size of the gonad in *endu-2* mutants.

Therefore, we put our focus on genes that were up-regulated in *endu-2(lf)* back ground. 32% (76

out of 237) of these transcripts were candidates for direct ENDU-2 targets, since they had been co-

immunoprecipitated with ENDU-2 (Supplemental Tab. S1). In addition, the vast majority of them

(62 out of 76) did not have increased expression in the gonad of *endu-2(lf)* animals (see the next

chapter, Supplemental Table S1). This suggests that, at minimum, these 62 transcripts are down-

regulated by ENDU-2 in the somatic tissues. GO term analyses implicated these somatic ENDU-

2 targets in regulation of various metabolic processes (Fig. 5A and Fig. S9C). Consistently, we

found loss of *endu-2* resulted in increased lipid content at 25°C but not at 15°C (Fig. 5B). Taken

together, these results indicate a putative regulatory role of ENDU-2 in limiting abundance of the

mRNAs that are primarily involved in metabolic functions in the soma.

As some aspects of the *endu-2(lf)* phenotype were temperature dependent, we asked whether

ENDU-2 regulates mRNA abundance in response to temperature alterations. To test this, we

performed qPCR to quantitate ten selected mRNAs which were bound and down-regulated by

ENDU-2. We found that mRNA levels of these targets were up-regulated in *endu-2* mutant

background, compared to wild type, already at standard growth temperature (20°C) and were even

more abundant at 25°C (Fig. 5C). Increased temperature, on the other hand, did not strongly affect

the abundance of most of these mRNAs in wild type background. These results suggest a

temperature dependent negative influence of ENDU-2 on the levels of these mRNA targets. We

also performed smFISH to inspect the influence of ENDU-2 on the mRNA level of yet another

target *fat-7*. *fat-7* mRNA was only detected in the intestine and *endu-2(-)* animals had higher *fat-*

7 transcript levels than wild type at both 20°C and 25°C (Fig. 5D). Moreover, we used a *fat-7::GFP*

translational fusion reporter to monitor FAT-7::GFP protein level. Wild type animals showed

reduced FAT-7::GFP expression at 25°C *vs.* 15°C (Fig. 5E). *endu-2(lf)* displayed stronger FAT-

7::GFP expression at both temperatures. Furthermore, transgenic expression of *endu-2(wt)* but not

the *endu-2(E454Q)* transgene (that has lost RNA cleavage activity) restored decreased FAT-

7::GFP expression (Fig. 5F). We conclude that ENDU-2 mediated mRNA-cleavage is required for

decreasing expression of at least some of its somatic target genes, such as *fat-7*.

ENDU-2 prevents misexpression of soma specific genes in the germline

To investigate germline transcriptomes regulated by ENDU-2, we isolated gonads from wild type

and *endu-2(lf)* animals which had been grown at 25°C from L1 stage, and sequenced two biological

replicates of polyadenylated RNA from each strain. A gene with normalized reads of RPKM > 1

was scored as expressed. We identified 8,356 expressed genes in the gonad of wild type animals

at 25°C, among which 88% (7,365) have been reported as expressed in the wild type gonad at

20°C (Ortiz et al., 2014). The following DESeq2 analysis showed that up-regulation of mRNA

levels was the prevalent change in the gonad of *endu-2* mutants (422 up-regulated, 35 down-

regulated, FC>2, P value<0.05; Fig. 6A, Supplemental Tab. S3). And only genes up-regulated in

endu-2(lf) background were enriched in ENDU-2 binding, as 218 out of 422 up-regulated genes

(P-value < 0.0001, Chi-Square test) and 11 out of 35 down-regulated genes (P-value = 0.7675,

Chi-Square test) were co-immunoprecipitated with ENDU-2. We considered them as direct targets

of ENDU-2 and grouped them in two classes, depending on whether they are up- (class I) or down-

regulated (class II) in *endu-2(lf)* background (Fig. 5B). Expression of a class I target is repressed

while class II genes are activated by wild type ENDU-2 activity. Consistently, we noticed that

most of the class I, but not the class II target genes were not, or very lowly, expressed in the gonad

of wild type animals (Fig. 5C) and tissue enrichment analysis suggested that the class I targets are

enriched expressed in the somatic tissues (intestine, neurons, muscles, somatic gonad and

pharynx). GO term enrichment analysis revealed that the major function of class I targets are

involved in immune and defense responses (Fig. 5B). We wondered whether the class I target

genes were preferentially soma specific genes that are misexpressed in the germline in the absence

of ENDU-2. To validate this assumption, we combined our wild type gonad transcriptome with a

previous published whole animal's transcriptome at 25°C (Gomez-Orte et al., 2018) and calculated

a soma enrichment factor (SEF) for each transcript to estimate the expression ratio between the

soma and the gonad. Class I but not class II targets had on average significantly higher SEF value

than the total mRNAs (Fig. 6D), suggesting that the class I genes are predominantly expressed in

the soma but only expressed at low level in the germline in wild type animals. As maintenance of

germline immortality by ENDU-2 does not require its RNA cleavage activity, we additionally

sequenced mRNA from isolated gonads of *endu-2(lf)* mutant carrying *endu-2(E454Q)::EGFP*

rescue transgene. Expressing *endu-2(E454Q)::EGFP* was sufficient to repress germline

expression of most all the class I targets (Fig. 6E and 6F). In summary, our data indicate the

inhibitory role of ENDU-2 to prevent germline expression of soma-specific genes is primarily

independent of its RNA cleavage activity.

Discussion

The task of germline is to maintain pluripotency and immortality for an accurate transmission of

genetic and epigenetic information between generations. Despite of emerging studies about

molecular mechanisms protecting the germline from the impacts of stress, one obvious open

question is whether the germline responds directly to environmental signals, such as nutrients or

stress, or rather reacts to alterations in somatic signaling. A direct stress response seems to be less

advantageous from an economical point of view, since this requires, in a pluripotent stem cell, the

expression of an entire hierarchy of mechanisms, involving the sensation of stress, signaling, as

well as responsive pathways to enable a dynamic and rapid response to a wide range of

environmental cues. In contrast, the second alternative would more beneficial as differentiated

somatic cells are specialized to sense environmental changes. In such a case, molecular

messengers, such as secreted hormones and ligands, are required to shuttle between somatic and

germline tissues to adjust a proper response in the germline. Previous studies have revealed that

microRNA could act as such messengers mediating communication between soma and germline

(Maciel and Mansuy, 2019; Sharma et al., 2018). Here we show secretion of a conserved

endoribonuclease ENDU-2 from the soma prevents misexpression of soma-specific gene in the

germline and preserve germline immortality at elevated temperature. This finding, together with

another recent study reporting regulation of germline proliferation by intestinal ENDU-2 in

response to thymidine imbalance (Jia et al., 2020), strongly suggests ENDU-2 as a crucial molecule

mediating non-cell-autonomous stress responses.

Our data implicate ENDU-2 in enabling long range communication between cells via its secretion

signal peptide (Fig. 2B). Whereas an earlier study had claimed neurons and muscles as the tissues

primarily expressing *endu-2* (Ujisawa et al., 2018), our data, consistent with the report from Jia et

al. (2020), suggest intestine as the most important organ to produce ENDU-2, whereas expression

in neuronal, muscular and somatic gonad tissues is rather weak (Fig. 2A and Supplemental Fig.

S3). We compared, for this analysis, both RNA and protein levels of ENDU-2 as well as different

ENDU-2 variants with truncations in the N-terminal secretion signal peptide (Δ_{ss}ENDU-2 and

SS$_{sel-1}$:: Δ$_{ss}$ENDU-2). We could demonstrate that the tissues containing ENDU-2 protein differ

from those that harbor *endu-2* mRNA and presence of a secretion signal peptide is necessary and

sufficient to target ENDU-2 to the secretory pathway to reach distant cells.

Our data strongly suggests that ENDU-2 production and function may occur in different tissues,

and that secretion of ENDU-2 allows the control of mRNA abundance in the distance. A simple

assumption would be that ENDU-2 has similar functions in any of its target tissues, no matter from

which cells it is expressed. This seems to be the case for some activities mediated by ENDU-2.

For example, the egg-laying defect observed as a consequence of abnormal vulva development in

an *endu-2(lf)* mutant is rescued by transgenic *endu-2* expression in the muscle, intestine or neuron

(data not shown). In contrast, only neuronally or intestinally expressed ENDU-2 was able to rescue

germline immortality (Fig. 3C), indicating existence of functional difference of ENDU-2 from

distinct origins. Tissue-specific interactors or modifiers of ENDU-2 are probably crucial to specify

its individual activities. It is currently not known whether there are isoform-specific functions of

ENDU-2, or whether ENDU-2 protein is modulated by protein modifications.

We distinguish two activities of ENDU-2: mRNA-binding and mRNA-cleavage. RIP-Seq and

transcriptomic results demonstrate that the levels of only about 10% of the mRNAs to which

ENDU-2 binds change in *endu-2(lf)* mutants at elevated temperature (Fig. 4-6), suggesting that

ENDU-2 is able to discriminate its regulated targets from the bound transcripts. Such scenarios

have also been proposed for SMG-2/UPF1, the core factor of the non-sense mediated decay

(NMD) complex that degrades only a small fraction of the RNAs it binds to (Muir et al., 2018).

ENDU-2 represses expression of its somatic target genes probably via direct RNA degradation

while it utilizes primarily RNA-binding but not cleavage activity to prevent misexpression of

soma-specific genes in the germline, suggesting rather an indirect mechanism of ENDU-2 in the

germline to restrict its target mRNA abundance. Why such different layers of regulatory

mechanisms exist is currently unknown. We speculate that one reason might be association of

ENDU-2 with distinct protein complexes in the soma and germline. Future study focusing on

dissecting tissue specific mechanisms will help to elucidate how ENDU-2 coordinates gene

expression in the soma and germline upon environmental stimuli.

Xenopus EndoU was suggested to both control snoRNA biogenesis in the nucleolus and cleave

mRNAs in the cytosol (Gioia et al., 2005; Schwarz and Blower, 2014). Mouse EndoU has been

reported to regulate peripheral B cell survival via reducing c-Myc mRNA level (Poe et al., 2014).

Inactivation of neuronal Drosophila DendoU causes neurotoxicity partially by down-regulation of

dTDP-43 (Laneve et al., 2017). SARS and SARS-Covid-2 endoribonucleases Nps15/NendoU

have been implicated in virus replication as well as limitation of host innate immune response,

although the detailed functions are still enigmatic (Kim et al., 2020). Our work here suggests that

C. elegans ENDU-2, in response to environmental stimuli such as temperature elevation, can exert

its function in tissues different from the ones in which its gene is expressed. Therefore, ENDU-2

is a novel candidate for transmitting environmental signals across tissue boundaries, most notably

involving signaling to the germline. A secretion signal peptide is also present in the human EndoU

homolog that is expressed in the placenta and detected in the serum (Inaba et al., 1980). Therefore,

the potential conservation of its non-cell-autonomous functions could open a new research area to

study intercellular communication. EndoU, thus, could be involved in a mechanism by which

environmental influences and experiences of the somatic tissues are transmitted into the

reproductive system.

Methods

Strains

The *C. elegans* N2 (Bristol) strain was used as wild type in all experiments in this study. Mutant strains: KHR83 *endu-2(tm4977),* outcrossed 8x with our N2. BR7130 *endu-2(by188)*, NL3511 *ppw-1(pk1425),* TX20 *oma-1(zu405)*, BR7510 *oma-1(zu405);endu-2(tm4977)*, BR8649 *oma-1(zu405);hrde-1(tm1200),* BR7205 *endu-2(by190[endu-2::EGFP])*, BR7089 *byEx1315[endu-2P::EGFP;rol-6(su1006)]*, BR8657 *endu-2(tm4977);byEx1814[endu-2P::endu-2::EGFP::$_{endu-2}$3'UTR;rol-6(su1006);]*, BR7295 *endu-2(tm4977);byEx1375[endu-2P::endu-2::EGFP;myo-2P::mCherry]*, BR5402 *byEx749[unc-119P::EGFP;rol-6(su1006)]*, BR8672 *byEx1821[unc-119P::SS$_{endu-2}$::EGFP;rol-6(su1006)]*, BR7827 *endu-2(tm4977);byEx1551[vha-6P::endu-2::EGFP::3xFlag;myo-2P::mCherry]*, BR7332 *endu-2(tm4977);byEx1379[fos-1P::endu-2::EGFP;myo-2P::mCherry]*, BR8551 *endu-2(tm4977);byEx1795[unc-119P::endu-2::EGFP::3xFlag;rol-6(su1006)]*, BR8662 *endu-2(tm4977);byEx1816[myo-3P::endu-2::EGFP::3xFlag;myo-2P::mCherry]*, BR7512 *endu-2(tm4977);byEx1449[endu-2P::Δ$_{ss}$endu-2::EGFP;rol-6(su1006);myo-2P::mCherry]*, BR8747 *endu-2(tm4977);byEx1847[endu-2P::Δ$_{ss}$endu-2::EGFP::$_{endu-2}$3'UTR;myo-2P::mCherry]*, BR8821 *endu-2(tm4977);byEx1875[endu-2P::SS$_{sel-1}$::Δ$_{ss}$endu-2::EGFP; myo-2P::mCherry]*, BR7680 *endu-2(tm4977);byEx1492[endu-2P::endu-2(E454Q)::EGFP::3xFlag;myo-2P::mCherry]*, BR7683 *endu-2(tm4977);byEx1495[endu-2P::endu-2(E460Q)::EGFP::3xFLAG;myo-2P::mCherry]*, DMS303 *nIs590[fat-7p::fat-7::GFP+lin15(+)]*, BR8317 *endu-2(tm4977);nIs590[fat-7p::fat-7::GFP+lin-15(+)]*, BR8754 *endu-2(tm4977);nIs590[fat-7p::fat-7::GFP+lin-15(+)];byEx1853[endu-2P::endu-2(E454Q);myo-2P::mCherry]*, BR8775 *endu-2(tm4977);nIs590[fat-7p::fat-7::GFP+lin-15(+)];byEx1860[endu-2P::endu-2;myo-2P::mCherry]*, RT688 *unc-119(ed3);pwIs28[pie-1p::cav-1::GFP(7)+unc-119(+)]*, BR8358

endu-2(tm4977);pwIs28[pie-1p::cav-1::GFP(7)+unc-119(+)]. For RIP-Seq the following strains

were used: BR7802 *endu-2(tm4977);byIs240[endu-2P::endu-2(E460Q)::EGFP::3xFLAG;myo-*

2p::mCherry],   BR7803   *endu-2(tm4977);byIs241[endu-2P::endu-*

2(E454Q)::EGFP::3xFLAG;myo-2P::mCherry], BR7205 *endu-2(by190[endu-2::EGFP])* and

Ex[ife-2P::GFP] (as IP control, gift from Tavernarakis Lab). For the RNA cleavage assay,

BR8311 *endu-2(tm4977);byIs267[endu-2P::endu-2::EGFP;myo-2::mCherry]* and BR7803 *endu-*

2(tm4977);byIs241[endu-2P::endu-2(E454Q)::EGFP::3xFlag;myo-2P::mCherry] were used in

additional to N2 and *endu-2(tm4977)* strains. *endu-2(tm4977)* allele was used for all experiments,

if not noted otherwise. Except for the Fig. 1A and Supplemental Fig. S2, all the *endu-2(lf)* mutants

were the granddaughter generation (G2) from a *endu-2(tm4977)* carrying transgenic *endu-2* rescue

strains whose parents (G1) had lost the rescue transgene. Information for extrachromosomal

transgenic strains generated in this study is summarized in the Supplemental Tab. S5.

Plasmids

To construct an EGFP transcriptional fusion reporter of *endu-2* (pBY3798), a 4691bp genomic

fragment upstream of the *endu-2* ATG was PCR amplified and inserted into pEGFP-N1 with

Eco47III/BglII sites. For translational fusion reporters, 5487 bp *endu-2* genomic region was cloned

into pBY3798 with *BglII/SmaI* sites to receive *endu-2P::endu-2::EGFP (*pBY3800), the *endu-2*

3'UTR was inserted into pBY3800 at *NotI* site to receive *endu-2P::endu-2::EGFP:: endu-2 3'UTR*

(pBY4137). *Δss endu-2::EGFP* (pBY3843) and *Δss endu-2::EGFP::endu-2 3'UTR* construct

(pBY4172) was generated by removing sequences encoding the N-terminal 2-19 amino acids from

pBY3800 and pBY4137, respectively. The *endu-2P::SS_{sel-1}::Δss endu-2::EGFP* construct

(pBY4194) was generated by insert the sequence encoding the 1-20 amino acids of *sel-1* to the

pBY3843 with *BglII* site. A *endu-2P::3xFlag::endu-2::EGFP:: endu-2 3'UTR* expressing construct

(pBY4138) was generated by inserting 3xFlag encoding sequence and *endu-2* genomic region into

pBY3798 with *BglII* and *BglII/SmaI* sites, respectively. The 3'UTR of *endu-2* was inserted via

Gibson ligation. The *unc-119P::EGFP* expressing construct (pBY2941) was made by fusing the

2.1 kb *unc-119* promoter region into pEGFP-N1 with *Eco47III/BglII* sites. The *unc-119P::SS$_{endu-}$*

$_2$::EGFP construct pBY4148 was generated by inserting the first 57 nucleotides of *endu-2* cDNA

into pBY2941 with *BglII/AgeI* sites. For expressing recombinant ENDU-2::EGFP in HEK 293T

cells, wild type *endu-2* cDNA was cloned into a 3xFlag containing pEGFP-N1 with *XhoI/SmaI*

(pBY3878). Site-directed mutagenesis was performed to receive *endu-2(E454Q)::EGFP::3xFlag*

(pBY3894) and *endu-2(E460Q)::EGFP::3xFlag* mutants (pBY3895). The fragments *endu-*

2::EGFP::3xFlag from pBY3878, endu-2(E454Q)::EGFP::3xFlag from pBY3894 and *endu-*

2(E460Q)::EGFP::3xFlag from pBY3895 were inserted into pBY3798 to receive constructs for

generating   *endu-2P::endu-2::EGFP::3xFlag*   (pBY3892),

endu-2P::endu-2(E454Q)::EGFP::3xFlag   (pBY3897)   and

endu-2P::*endu-2(E460Q)::EGFP::3xFlag* (pBY3898) reporter strains. *EGFP::3xFlag* fragments

in pBY3892 and pBY3897 were removed to receive pBY4188 *endu-2P*::*endu-2* and pBY4066

endu-2P::*endu-2(E454Q)*. Somatic gonad specific expression was achieved via PCR amplifying

and insertion of *endu-2* cDNA into a *fos-1P::GFP* plasmid (a gift from David Sherwood lab) with

SalI and *SmaI* sites (pBY3833). For intestinal specific expression, a 1.2 kb *vha-6* promoter was

inserted at *Eco47III/XhoI* sites into pBY3878 to receive *vha-6P::endu-2::EGFP::3xFlag*

(pBY3937). A 2.1 kb *unc-119* promoter was inserted into pBY3878 at *Eco47III/BglII* sites to

receive *unc-119P::endu-2(cDNA)::EGFP::3xFlag* (pBY4127) for neuronal expression. A 2.4 kb

myo-3 promoter region was cloned into pBY3878 to receive *myo-3P::endu-2::EGFP::3xFlag*

(pBY4135) for muscular specific *endu-2* expression.

Scoring of Mrt phenotype

Mrt phenotype was examined by quantification of brood size by separating 15 L4 animals on single plates and counting number of progeny per animal, or transferring six L4 larvae onto new agar plates and scoring percentage of the fertile animals 36 hours after the L4 stage in the next generation. Both *endu-2(tm4977)* and *endu-2(by188)* were further outcrossed 4x with N2 before scoring Mrt phenotype for Fig. 1A and Supplemental Fig. S2A. For brood size recovery at 15°C, wild type and outcrossed *endu-2(tm1977)* animals were maintained at 25°C for 6 generations until *endu-2(tm4977)* showed strong sterile phenotype. 15 L1-L2 animals from the 6th generation were shifted to 15°C and designated as G1. For Mrt rescue experiment, *endu-2(tm4977)* progenies of *endu-2(tm4977);byEx1375[endu-2::EGFP;myo-2P::mCherry]*, *endu-2(tm4977);byEx1449[endu-2P::Δ$_{ss}$endu-2::EGFP;myo-2P::mCherry]*, *endu-2(tm4977);byEx1875[endu-2P::SS$_{sel-1}$::Δ$_{ss}$endu-2::EGFP;myo-2P::mCherry]*, *endu-2(tm4977);byEx1488[endu-2P::endu-2(E454Q)::EGFP::3xFLAG;myo-2P::mCherry]* and *endu-2(tm4977);byEx1488[endu-2P::endu-2(E460Q)::EGFP::3xFLAG;myo-2P::mCherry]* animals (P0) were isolated and designated as G1.

Germline proliferation

Number of germ nuclei in the proliferation zone was quantified as described in a previous study (Qi et al., 2017).

Lifespan

Lifespan at 20°C was initiated at L4 stage. As *endu-2(tm4977)* animals display strong egg-laying defect due to abnormal vulval development, agar plates containing 200µM FUDR were used to during the first seven days of adulthood to avoid internal hatching.

Antibody staining

Antibody staining was performed with day one adult animals as described (Crittenden and Kimble,

2009). A monoclonal mouse anti-GFP antibody (Roche, Nr. 11814460001) was used to detect

ENDU-2::EGFP in the gonad.

oma-1 **RNAi inheritance assay**

oma-1 RNAi was initiated at L1 stage at 20°C (restrictive temperature) and these animals were

designated as P0. Six P0 day one adults animals were transferred onto OP50 seeded agar plates

and incubated at 20°C for scoring the embryonic lethality over generations. Embryonic lethality

was examined by transferring of about 100 embryos onto an OP50 seeded agar plate and

quantifying the number of hatched animals after 48 hours.

Oil Red O staining to quantify content of body fat

Worms were fixed in 500µl 60% isopropanol. After removal of supernatant, 500µl freshly filtered

ORO working solution was added to stain worms at 25°C in a wet chamber overnight. Animals

were washed once with 1ml 0.01% Triton-x100 containing M9 buffer and responded in 250µl M9-

Triton-x100 buffer. Stained worms can be stored at 4°C for at least one month. For quantification,

images were recorded with color camera. The original images were spitted into red and blue

channels and the level of Oil-Red-O was quantified by determining the excess intensity in the red

channel in comparison to the blue channel. The mean fatness per worm was calculated as the total

intensity within stained regions normalized by the area of the worm regions. Oil Red O working

solution was prepared as follows: 0.5 g of Oil Red O powder was dissolved in 100 ml isopropanol

solution and equilibrated for several days. The solution was then freshly diluted with 40% water

for a 60% stock and allowed to sit overnight at room temperature and filtered using 0.2 mm filters.

RIP-Seq

To isolate ENDU-2 associated RNAs, animals of mixed stage were lysed in Lysis-NP-40 buffer

(150 mM NaCl,50 mM Tris pH8.0, 1% NP40 + Protease Inhibitor+100 µl/ml RNase Inhibitor)

with SilentCrusher (x30 sec. 75000 rpm). Lysates were clarified by centrifuging at 14k rpm for 15

min. Supernatants were pre-cleared with Dynal Protein A magnetic beads (Invitrogen) and

incubated with anti-GFP antibody (Abcam) for 1 hour at 4ºC, followed by incubation with Dynal

Protein A Magnetic Beads for 2 h at 4ºC. Beads were washed 3 times with Lysis-NP-40 Puffer

before DNase I digestion (100 µl 1x DNase buffer, 1 µl DNase I, 1 µl RNase inhibitor) at 37 ºC

for 30 minutes. 10 µl 50 mM EDTA was added to stop DNase digestion. RNA was extracted with

5 volumes Trizol reagent (Invitrogen), followed by isopropanol precipitation. Precipitates were

resolved in RNase-free water for visualization on 12.5% denaturing acrylamide gel with SYBR

Gold (Thermo Scientific). Random hexamer primers were used for preparing TruSeq3 reverse-

stranded cDNA libraries for Illumina single-end 50bp sequencing. RIP-Seq to compare RNA

binding efficiency of ENDU-2(wt), ENDU-2(E454Q) and ENDU-2(E460Q) at 15 ºC and 25 ºC

were performed without cross-linking. RIP-Seq to determine RNA targets of ENDU-2 was carried

out with ENDU-2(E454Q)::EGFP expressing animals raised at 15 ºC with UV cross-linking.

Analysis of RIP-Seq data

The sequencing data were uploaded to the European Galaxy Server at https://usegalaxy.eu, and all

read quality controls, trimming, mapping and counting were performed through Galaxy (Afgan et

al., 2018). Specifically, after trimming with Trimmomatic 0.36 (Bolger et al., 2014), we mapped

the reads to the *C. elegans* genome using the Wormbase WS260 genomic sequence

(ftp://ftp.wormbase.org/pub/wormbase/releases/WS260/species/c_elegans/PRJNA13758/c_elega

ns.PRJNA13758.WS260.genomic.fa.gz)   and   the   WS260   canonical   gene   set

(ftp://ftp.wormbase.org/pub/wormbase/releases/WS260/species/c_elegans/PRJNA13758/c_elega

ns.PRJNA13758.WS260.canonical_geneset.gtf.gz) with the RNA aligner STAR 2.6.0b (Dobin et

al., 2013) with default settings, then counted reads mapping to individual genes using feature

Counts 1.6.2 (Liao et al., 2014). For a read to be counted as mapping to a particular gene, we

required a minimum read mapping quality of 12 (-Q 12 option of feature Counts) and an overlap

of at least 1 base between the read and any of the exons of the gene (--minOverlap 1).

The IP enrichment of a transcript was expressed as: FC=(IP RPM + 1)/(Mock IP RPM + 1), where

RPM= number of reads mapped to a gene/(number of all mapped reads) x 10ˆ6. The addition of 1

in the FC formula prevents infinite enrichment. ENDU-2 associated RNAs were defined as

transcripts with both FC>=4 and RPKM >=1 in ENDU-2(E454Q) IP sample. RPKM is calculated

as follows: RPKM = RPM/length of transcripts x 10ˆ3.

RNA-seq and data processing of isolated gonads

Animals were raised at 25°C from the L1 larval stage and the complete gonads were isolated 24 h

after mid-L4 stage. Total RNA of the gonads was extracted with RNeasy® Mini Kit (Qiagen,

Venlo, The Netherlands). Purification of poly-A containing RNA molecules, RNA fragmentation,

strand-specific random primed cDNA library preparation and Single-read sequencing (50 bp) on

an Illumina HiSeq 4000 were carried out by Eurofins Genomics. The RNA-seq results from two

independent biological replicate of dissected gonads were uploaded to the European Galaxy Server

at https://usegalaxy.eu. All read quality controls, trimming, mapping and counting were performed

through Galaxy (Afgan et al., 2018) using protocols like described for RIP-Seq (above). The

DESeq2 (Galaxy Version 2.11.40.6+galaxy1) was used to determine differentially expressed

features from count tables of differential transcript abundances. The RPKMs throughout this study

were normalized separately. Only the longest isoform of each gene was used as length of a

transcript. To estimate genes with enriched expression in the soma, RNA-seq data from wild type

animals raised at 25°C from a previous study (Gomez-Orte et al., 2018) were compared with our

transcriptome from isolated gonads of wild type animals. The soma enrichment factor (SEF) was

calculated with the formal SEF=(RPKM$_{\text{whole animal}}$ +2)/(RPKM$_{\text{gonad}}$ +2)]. Addition of factor 2 to the

formal minimize large SEF value caused by low RPKM values.

In *vitro*. RNA binding assay of recombinant ENDU-2 protein

The RNA binding assay was performed as described (Panda et al., 2016). *fat-7* and *trcs-1* mRNA

were transcribed from complementary DNA (cDNA) with biotinylated uracil. The presence and

purity of RNA were examined in an RNA gel. This biotin-labeled RNA was incubated with

recombinant   ENDU-2(wt)::EGFP::3xFlag,   ENDU-2(E454Q)::EGFP::3xFlag   and   ENDU-

2(E460Q)::EGFP::3xFlag proteins expressed in HEK 293T cells. The labeled RNA was captured

and isolated using streptavidin beads. Proteins binding the RNA were then visualized via Western

Blot.

RNA cleavage assay

Pellets of animals raised at 15°C in mixed stages were washed twice with 1 ml 10 mM EGTA

containing M9 buffer before resuspension in 0.7 ml NP-40 buffer. 3x Protease inhibitor was added

to the buffer and worms were lysed with SilentCrusher (2x30", 75,000 rpm). 5µl RNAsin Plus

RNAse Inhibitor (40U/µl) was added to the worm lysate and centrifuged by 12,000 rpm for 15

min at 4°C. For the RNA cleavage assay 95 µl supernatant of the worm lysate with 200 µg total

protein was incubated with 5 mM Ca^{2+}, Mg^{2+}, Mn^{2+} or 10 mM EGTA for 30 minutes at 7°C before

total RNA was extracted with Trizol. 0.2 µg RNA was loaded on 10% 8M Urea acrylamide gel

with SYBR Gold for visualization of RNA.

Microarray

Both *endu-2(tm4977)* and *endu-2(tm4977);byEx1375[endu-2P::endu-2::EGFP;myo-*

2::mCherry] animals used for microarray were granddaughter decedents of one single *endu-*

2(tm4977);byEx1375[endu-2P::endu-2::EGFP;myo-2::mCherry] animals (stain preparation see

Supplemental Fig. S8). Animals were raised at 25ºC for 48 h after hatching and total RNA was

purified by using RNeasy® Mini Kit (Qiagen, Venlo, The Netherlands). Microarray and the

following data analysis were performed as described before (Qi et al., 2017; Ritchie et al., 2015).

GO term enrichment analysis

GO term and tissue enrichment analysis was carried out with the online enrichment analysis tool

(https://wormbase.org/tools/enrichment/tea/tea.cgi)(Angeles-Albores et al., 2016).

Quantification of mRNA level via quantitative RT-PCR (q-PCR)

Total RNA was prepared from day one adult animals raised at 25ºC or 20ºC by using RNeasy®

Mini Kit (Qiagen, Venlo, The Netherlands). *endu-2(tm4977)* animals were granddaughter

decedents of *endu-2(tm4977);byEx1375[endu-2P::endu-2::EGFP;myo-2::mCherry]* animals.

act-4 was used as internal control. q-PCR primer sequences: *cav-1* forward aagtgctggtggagtagatgc,

cav-1 reverse tccgatagcgatgttctcttc, *clec-169* forward tggacacttgtaactgtgcaaga, *clec-169* reverse

ttttcatttgaccactttagatcg, *F38B6.4* forward cccttgtcctcggaaattaga, *F38B6.4* reverse

caggcccgaagatggtaat, *pfas-1* forward caagattggaagttccgaaga, *pfas-1* reverse ctcaatatccggacagtcgtc,

B0286.3 forward gacccgaaaacttggagctt, *B0286.3* reverse Tggaacgaatgagcacataatc, *atic-1* forward

tgctacaaaaatgccagcag, *atic-1* reverse cgtttaaaggaagaccaacagc, *acox-1.2* forward

agcggtgatctatggaagtga, *acox-1.2* reverse agctcagggatcttggacac, *abdh-3.2* forward

atgacaccaccccaattgtc, *abdh-3.2* reverse tgctgtcatgagtacttcctgtg, *col-176* forward

ccacaacctttgctccaatc, *col-176* reverse gagcacacttgatgcagtcg, *clec-15* forward tgcgccagaagggtattact,

clec-15 reverse cgcagaacgatctaaccaga, *asp-14* forward gctgcagttaccaacattacca, *asp-14* reverse

agcaacgaagaaggttgagg.

Single Molecule Fluorescent in Situ Hybridization (smFISH) and data analysis

38-48 probes of 20nt in length targeting each mature mRNA were designed with Stellaris RNA

FISH probe designer on gene target specificity. Probes were produced, conjugated to CAL Fluor

Red 610 and purified by Biocat (sequences of the probe sets are available in the supplemental

Table S4). Wild type and *endu-2(tm4977)* L4 animals were selected and dissected 24 hours later

for smFISH staining using a published protocol procedure (Lee et al., 2016). Images were acquired

with an Image Z1 fluorescence microscope. Exposure times and acquisition settings were identical

between replicates. mRNA puncta were quantified by using ImageJ 1.51s cell counter plugin.

Regions of interest for acquisition were defined by nuclei DAPI staining.

Statistical analysis

604 Experiments shown in this study were performed independently twice to four times. Details of the

605 particular statistical analyses used, precise *P* values, statistical significance, number of biological

606 replicas and sample sizes for all of the graphs are indicated in the figures or figure legends. *n*

607 represents the number of animals tested, unless mentioned otherwise. N means the number of

608 biological replicates. *** represents P-value < 0.0001.

Data availability.

Microarray data were submitted to the ArrayExpresss under accession number E-MTAB-7993.

RIP-seq data were submitted to BioProject under BioProject ID PRJNA544611. RNA-seq data

of isolated gonad were submitted to NCBI Sequence Read Archive with ID SUB7811674. The

remaining data for RIP-Seq and RNA-seq of isolated gonad are located in Supplemental Table

S1-S3. Correspondence and requests for materials should be addressed to B.R

(baumeister@celegans) or Q. W (wenjing.qi@biologie.uni-freiburg.de).

References

Afgan, E., Baker, D., Batut, B., van den Beek, M., Bouvier, D., Cech, M., Chilton, J., Clements,

D., Coraor, N., Gruning, B.A.*, et al.* (2018). The Galaxy platform for accessible, reproducible and

collaborative biomedical analyses: 2018 update. Nucleic Acids Res *46*, W537-W544.

Ahmed, S., and Hodgkin, J. (2000). MRT-2 checkpoint protein is required for germline

immortality and telomere replication in C. elegans. Nature *403*, 159-164.

Alcazar, R.M., Lin, R., and Fire, A.Z. (2008). Transmission dynamics of heritable silencing

induced by double-stranded RNA in Caenorhabditis elegans. Genetics *180*, 1275-1288.

Angeles-Albores, D., RY, N.L., Chan, J., and Sternberg, P.W. (2016). Tissue enrichment analysis

for C. elegans genomics. BMC Bioinformatics *17*, 366.

Barucci, G., Cornes, E., Singh, M., Li, B., Ugolini, M., Samolygo, A., Didier, C., Dingli, F., Loew,

D., Quarato, P.*, et al.* (2020). Small-RNA-mediated transgenerational silencing of histone genes

impairs fertility in piRNA mutants. Nat Cell Biol *22*, 235-245.

Bolger, A.M., Lohse, M., and Usadel, B. (2014). Trimmomatic: a flexible trimmer for Illumina

sequence data. Bioinformatics *30*, 2114-2120.

Buckley, B.A., Burkhart, K.B., Gu, S.G., Spracklin, G., Kershner, A., Fritz, H., Kimble, J., Fire,

A., and Kennedy, S. (2012). A nuclear Argonaute promotes multigenerational epigenetic

inheritance and germline immortality. Nature *489*, 447-451.

Crittenden, S., and Kimble, J. (2009). Preparation and immunolabeling of Caenorhabditis elegans.

Cold Spring Harb Protoc *2009*, pdb prot5216.

Dobin, A., Davis, C.A., Schlesinger, F., Drenkow, J., Zaleski, C., Jha, S., Batut, P., Chaisson, M.,

and Gingeras, T.R. (2013). STAR: ultrafast universal RNA-seq aligner. Bioinformatics *29*, 15-21.

Gioia, U., Laneve, P., Dlakic, M., Arceci, M., Bozzoni, I., and Caffarelli, E. (2005). Functional

characterization of XendoU, the endoribonuclease involved in small nucleolar RNA biosynthesis.

J Biol Chem *280*, 18996-19002.

Gomez-Orte, E., Cornes, E., Zheleva, A., Saenz-Narciso, B., de Toro, M., Iniguez, M., Lopez, R.,

San-Juan, J.F., Ezcurra, B., Sacristan, B.*, et al.* (2018). Effect of the diet type and temperature on

the C. elegans transcriptome. Oncotarget *9*, 9556-9571.

Grant, B., and Greenwald, I. (1996). The Caenorhabditis elegans sel-1 gene, a negative regulator

of lin-12 and glp-1, encodes a predicted extracellular protein. Genetics *143*, 237-247.

Hackbart, M., Deng, X., and Baker, S.C. (2020). Coronavirus endoribonuclease targets viral

polyuridine sequences to evade activating host sensors. Proc Natl Acad Sci U S A *117*, 8094-8103.

Inaba, N., Renk, T., and Bohn, H. (1980). Immunohistochemical location of placental proteins

(PP8, 9, 10, 11, 12) in human term placentae. Arch Gynecol *230*, 109-121.

Ivanov, K.A., Hertzig, T., Rozanov, M., Bayer, S., Thiel, V., Gorbalenya, A.E., and Ziebuhr, J.

(2004). Major genetic marker of nidoviruses encodes a replicative endoribonuclease. Proc Natl

Acad Sci U S A *101*, 12694-12699.

Jia, F., Chi, C., and Han, M. (2020). Regulation of Nucleotide Metabolism and Germline

Proliferation in Response to Nucleotide Imbalance and Genotoxic Stresses by EndoU Nuclease.

Cell Rep *30*, 1848-1861 e1845.

Katz, D.J., Edwards, T.M., Reinke, V., and Kelly, W.G. (2009). A C. elegans LSD1 demethylase

contributes to germline immortality by reprogramming epigenetic memory. Cell *137*, 308-320.

Kelly, W.G., Xu, S., Montgomery, M.K., and Fire, A. (1997). Distinct requirements for somatic

and germline expression of a generally expressed Caernorhabditis elegans gene. Genetics *146*,

227-238.

Kim, Y., Jedrzejczak, R., Maltseva, N.I., Wilamowski, M., Endres, M., Godzik, A., Michalska,

K., and Joachimiak, A. (2020). Crystal structure of Nsp15 endoribonuclease NendoU from SARS-

CoV-2. Protein Sci.

Laneve, P., Piacentini, L., Casale, A.M., Capauto, D., Gioia, U., Cappucci, U., Di Carlo, V.,

Bozzoni, I., Di Micco, P., Morea, V.*, et al.* (2017). Drosophila CG3303 is an essential

endoribonuclease linked to TDP-43-mediated neurodegeneration. Sci Rep *7*, 41559.

Lee, C., Sorensen, E.B., Lynch, T.R., and Kimble, J. (2016). C. elegans GLP-1/Notch activates

transcription in a probability gradient across the germline stem cell pool. Elife *5*.

Lev, I., Seroussi, U., Gingold, H., Bril, R., Anava, S., and Rechavi, O. (2017). MET-2-Dependent

H3K9 Methylation Suppresses Transgenerational Small RNA Inheritance. Curr Biol *27*, 1138-

1147.

Liao, Y., Smyth, G.K., and Shi, W. (2014). featureCounts: an efficient general purpose program

for assigning sequence reads to genomic features. Bioinformatics *30*, 923-930.

Maciel, E., and Mansuy, I.M. (2019). Extracellular Vesicles and their miRNA Cargo: A Means of

Communication between Soma and Germline in the Mammalian Reproductive System. Chimia

(Aarau) *73*, 356-361.

Muir, V.S., Gasch, A.P., and Anderson, P. (2018). The Substrates of Nonsense-Mediated mRNA

Decay in Caenorhabditis elegans. G3 (Bethesda) *8*, 195-205.

Ni, J.Z., Kalinava, N., Chen, E., Huang, A., Trinh, T., and Gu, S.G. (2016). A transgenerational

role of the germline nuclear RNAi pathway in repressing heat stress-induced transcriptional

activation in C. elegans. Epigenetics Chromatin *9*, 3.

Ortiz, M.A., Noble, D., Sorokin, E.P., and Kimble, J. (2014). A new dataset of spermatogenic vs.

oogenic transcriptomes in the nematode Caenorhabditis elegans. G3 (Bethesda) *4*, 1765-1772.

Panda, A.C., Martindale, J.L., and Gorospe, M. (2016). Affinity Pulldown of Biotinylated RNA

for Detection of Protein-RNA Complexes. Bio Protoc *6*.

Poe, J.C., Kountikov, E.I., Lykken, J.M., Natarajan, A., Marchuk, D.A., and Tedder, T.F. (2014).

EndoU is a novel regulator of AICD during peripheral B cell selection. J Exp Med *211*, 57-69.

Qi, W., Yan, Y., Pfeifer, D., Donner, V.G.E., Wang, Y., Maier, W., and Baumeister, R. (2017). C.

elegans DAF-16/FOXO interacts with TGF-ss/BMP signaling to induce germline tumor formation

via mTORC1 activation. PLoS Genet *13*, e1006801.

Ritchie, M.E., Phipson, B., Wu, D., Hu, Y., Law, C.W., Shi, W., and Smyth, G.K. (2015). limma

powers differential expression analyses for RNA-sequencing and microarray studies. Nucleic

Acids Res *43*, e47.

Schwarz, D.S., and Blower, M.D. (2014). The calcium-dependent ribonuclease XendoU promotes

ER network formation through local RNA degradation. J Cell Biol *207*, 41-57.

Sharma, U., Sun, F., Conine, C.C., Reichholf, B., Kukreja, S., Herzog, V.A., Ameres, S.L., and

Rando, O.J. (2018). Small RNAs Are Trafficked from the Epididymis to Developing Mammalian

Sperm. Dev Cell *46*, 481-494 e486.

Simon, M., Sarkies, P., Ikegami, K., Doebley, A.L., Goldstein, L.D., Mitchell, J., Sakaguchi, A.,

Miska, E.A., and Ahmed, S. (2014). Reduced insulin/IGF-1 signaling restores germ cell

immortality to Caenorhabditis elegans Piwi mutants. Cell Rep *7*, 762-773.

Tijsterman, M., Okihara, K.L., Thijssen, K., and Plasterk, R.H. (2002). PPW-1, a PAZ/PIWI

protein required for efficient germline RNAi, is defective in a natural isolate of C. elegans. Curr

Biol *12*, 1535-1540.

Ujisawa, T., Ohta, A., Ii, T., Minakuchi, Y., Toyoda, A., Ii, M., and Kuhara, A. (2018).

Endoribonuclease ENDU-2 regulates multiple traits including cold tolerance via cell autonomous

and nonautonomous controls in Caenorhabditis elegans. Proc Natl Acad Sci U S A *115*, 8823-

8828.

Weiser, N.E., Yang, D.X., Feng, S., Kalinava, N., Brown, K.C., Khanikar, J., Freeberg, M.A.,

Snyder, M.J., Csankovszki, G., Chan, R.C.*, et al.* (2017). MORC-1 Integrates Nuclear RNAi and

Transgenerational Chromatin Architecture to Promote Germline Immortality. Dev Cell *41*, 408-

423 e407.

Acknowledgements

We thank N. Tavernarakis and the *Caenorhabditis elegans* Genetics Center which is funded by

the NIH Office of Research Infrastructure Programs (P40 OD010440) for providing some strains;

Dr. S Mitani for providing *endu-2(tm4977)* strain, Cornelia Habacher for sharing the RIP-Seq

protocol, D. Sherwood for providing the *fos-1* promoter. We thank the staff of the Life Imaging

Center (LIC) of the Albert-Ludwigs-University Freiburg for their microscopy resources. This work

was funded by grants from the German Research Foundation (DFG) (SFB850, SFB1381),

Germany's Excellence strategy (CIBSS-EXC-2189 - Project ID 8 390939984) and the German

Excellence Initiative (BIOSS - EXC294) to R.B.

Author Contributions

W.Q. and R.B. designed the experiments and analyzed data. W.Q., E.D.v.G., F. X., Q. Z., L.L. and

W.Y. performed the experiments. D.P. performed and analyzed microarray. W.Q. and W.M.

analyzed RNA-Seq data. W.Q. and R.B. wrote the manuscript.

Competing interest

The authors declare no competing financial interests.

Correspondence and requests for materials should be addressed to W.Q.

(wenjing.qi@biologie.uni-freiburg.de)

Fig. 1 *endu-2* mutant shows a temperature dependent Mrt phenotype.

A. Upper graph: Brood size of *endu-2(tm4977)* animals decreases gradually after 4x initial

backcrosses with wild type (G0) over generations at 20°C. Lower graph: The defects of *endu-*

2(tm4977) animals in reproduction is reversible at 15°C. To obtain this graph, adult animals from

G6 that had been maintained at 25°C were shifted back to 15°C and counted as G0. As numbers

of generation to reach 100% sterility at 20°C among the 4 biological replicates varied from 10 to

20, only one represented replicate with n=15 animals for each generation is shown. The data are

mean ± SEM. **B**. The sterile *endu-2(tm4977)* animals display multiple defects in germline mitosis

and meiosis. I-IV) DIC images of wild type and sterile *endu-2(tm4977)* animals 24 hours after

mid-L4 stage. The white arrows point to (II) increased number of germline apoptotic corpses; (III)

empty gonad due to germ cell death; IV) prolonged spermatogenesis; (V) DAPI staining of

germline proliferating zone of day one wild type and (VI) sterile *endu-2(tm4977)* adults,

respectively. White arrows in VI point to abnormally large chromosomes due to endomitosis. Scale

bar 10 µm.

Fig. 2. ENDU-2 is a secreted protein.

A. Fluorescence micrographs of transgenic *endu-2(tm4977);byEx1814[endu-2P::endu-*

2::EGFP::$_{endu-2}$3'UTR] animals. ENDU-2::EGFP is detected in the intestine, the somatic gonad,

coelomocytes and extracellular space between the uterus and embryos. **B.** Fusion of the 1-19 amino

acids of ENDU-2 (SS$_{endu-2}$) to neuronal specific expressed EGFP is sufficient for secretion of

EGFP protein. Upper panels shows expression pattern of *unc-119P::EGFP* which is detected only

in neuronal cells (strong) of animals and embryos. Lower panels are images showing expression

pattern of *unc-119P::SS$_{endu-2}$::EGFP* which is detected in the neurons (weak), coelomocytes and

interspace between egg shell and embryos, indicative of efficient secretion of EGFP enforced by

the secretion signal peptide of ENDU-2. Images are representative for more than 20 animals,

analyzed by using a 40× objective. **C.** Fluorescence micrographs of GFP-antibody staining of

endu-2(tm4977);byEx1375[endu-2P::endu-2::EGFP],   *endu-2(tm4977);byEx1449[endu-*

2P::Δ$_{ss}$endu-2::EGFP]   and   *endu-2(tm4977);byEx1875[endu-2P::SS$_{sel-1}$::Δ$_{ss}$endu-2::EGFP]*

transgenic animals at 20°C. ENDU-2::EGFP (n=62, 100%) and SS$_{sel-1}$::Δ$_{ss}$ENDU-2::EGFP (n=21,

100%) but not Δ$_{ss}$ENDU-2::EGFP (n=26, 0%) is detected in the germline. N=2. **D.** smFISH

staining reveals presence of *endu-2* mRNA in the intestine (white arrow) but not in the germline.

Blue: DAPI stained nuclear DNA. Red: smFISH probes stained mRNA. N=4. Scale bar 10 µm for

all images in this Figure.

Fig. 3. Maternally provided ENDU-2 from the soma contributes to maintenance of germline

immortality.

A. *endu-2(wt)::EGFP* rescues the Mrt phenotype at 25°C. Left panel shows the experimental

strategy for all rescue experiments with *endu-2* transgenes in this study. *endu-2(-/-)* daughter

generation (G1) that has lost the extrachromosomal rescue constructs expressing *endu-2::EGFP*

was isolated. This and the following *endu-2(-/-)* generations (G2 to Gn) were compared to *endu-*

2;Ex[endu-2:.EGFP] derived from the same P0 animal. Data are mean ± SD, N=3. **B**. Expression

of *endu-2(wt)::EGFP* from neuron or intestine is sufficient to rescue the Mrt phenotype at 25°C.

Data are mean ± SD, N=3. **C**. *SS$_{sel-1}$::Δ$_{ss}$endu-2::EGFP* but not *Δ$_{ss}$endu-2::EGFP* transgene

rescues the Mrt phenotype at 25°C. Data are mean ± SD, N=3. **D**. ENDU-2 primarily affects oocyte

to maintain germline immortality. Data are pooled data from three biological replicates with

similar tendencies in results.

Fig. 4. mRNAs-binding but not –cleavage activity of ENDU-2 protects germline immortality.

A. Comparison of RNA binding activities of ENDU-2(wt)::EGFP, ENDU-2(E454Q)::EGFP, and

ENDU-2(E460Q)::EGFP at 15°C and 25°C. Shown are plotted normalized reads (RPM) from RIP-

Seq of each identified transcript under different conditions. **B.** Mapping of the RIP-Seq reads of

two representative binding targets of ENDU-2. Only fragments of mRNA exons (red boxes) but

not introns (black lines) were co-immunoprecipitated with ENDU-2. **C.** ENDU-2::EGFP variants

bind to selected mRNAs in *vitro*. Shown are Western Blots to detect proteins binding to *fat-7* and

trcs-1 mRNA, respectively. EGFP is a negative control. N= 3 biological replicates. **D.** Wild type

ENDU-2, but not ENDU-2(E454Q), leads to RNA decay (smear) in a Ca^{2+} and Mn^{2+} dependent

manner, N=3. Shown is fused images of two gels from one experiment. The results of additional

two biological replicates are shown in the Supplemental Fig. S8. **E.** *endu-2(E454Q)::EGFP* but

not *endu-2(E460Q)::EGFP* transgene rescues the Mrt phenotype. Data are mean ± SD, N=3. Both

e*ndu-2(tm4977)* and *endu-2(tm4977)* carrying *endu-2(E454Q)::EGFP* or *endu-2(E460Q)::EGFP*

transgenes were decedents of one single P0 animal carrying the respective transgenes.

Fig. 5. ENDU-2 negatively regulates mRNA abundance in the soma.

A. Volcano blot of microarray results comparing *endu-2(tm4977)* mutant and *endu-2(tm4977)*

animals carrying the rescuing *endu-2::EGFP* transgene at 25°C. N = 3 biological replicates. The

blue dots mark transcripts bound by ENDU-2. Transcripts with blue and black label were ENDU-

2 associated according to the RIP-Seq result and those with black labels were verified with q-PCR.

B. ENDU-2 negatively regulates lipid content at 25°C. Shown are pooled data of quantification of

relative Oil Red O stained signal in wild type, *endu-2(tm4977)* as well as *endu-*

2(tm4977);byEx1375[endu-2::EGFP] day one adult animals at different temperatures. N=3

replicates. Statistical test with One-way ANOVA. **C.** qPCR to quantitate selected ENDU-2 target

mRNAs in the soma at 20°C and 25°C. Data are mean ± SD, N= three biological replicates. **D.**

Fluorescent images of smFISH staining of the ENDU-2 target *fat-7* mRNA and quantification of

the smFISH stained mRNA foci per examined area in wild type and *endu-2(tm4977)* animals.

Scale bar 10 µm. Data are mean ± SD of pooled data from three biological replicates. Statistical

test with One-way ANOVA. **E.** Fluorescent images and quantification of fluorescence intensity of

FAT-7::GFP in wild type and *endu-2(tm4977)* day one adult animals at different temperatures.

Shown are pooled data from three biological replicates. Statistical test with One-way ANOVA. **F**.

Fluorescent images and quantification of fluorescence intensity of FAT-7::GFP in wild type, *endu-*

2(tm4977), *endu-2(tm4977);Ex[endu-2(wt)]* and *endu-2(tm4977);Ex[endu-2(E454Q)]* day one

adult animals at 25°C. Shown are pooled data from three biological replicates. Statistical test with

One-way ANOVA. *** in this Figure means P value < 0.0001.

Fig. 6. ENDU-2 prevents misexpression of soma specific genes in the germline.

A. Volcano plot of gonad expressed genes in *endu-2(tm4977)* relative to wild type day one adult

animals grown up at 25°C. 2 biological replicates for each strain. **B.** Comparison of the total

transcripts co-immunoprecipitated with ENDU-2 and ENDU-2 regulated genes in the gonad. **C.**

Comparison of expression levels (RPKM) of total gonad-expressed genes, ENDU-2 class I and

class II targets in wild type animals. Shown are Box-Plot of Min/Max whiskers. Statistical test

with One-way ANOVA. **D.** Comparison of relative expression between the soma and the gonad of

the total transcripts, ENDU-2 class I and class II targets in wild type animals. Shown are Box-Plot

of Min/Max whiskers. Statistical test with One-way ANOVA. **E.** The majority of the alerted

expression of the germline class I targets is rescued by *endu-2(E454Q)::EGFP* transgene. **F.** Heat

map of ENDU-2 repressed germline genes in wild type, *endu-2(tm4977)* and *endu-2;Ex[endu-*

2(E454Q)::EGFP] animals. *** in this Figure means P value < 0.0001

Fig.1

Fig.2

Fig.3

Fig.4

Fig.5

A

ENDU-2 soma targets

lipid metabolic process
organic acid metabolic process
nucleoside phosphate binding
molting cycle
extracellular space
hydrolyse activity
purine nucleotide metabolism

B

WT
endu-2(lf)
endu-2;Ex[endu-2::EGFP]

C

clec-15
atic-1
asp-14
abhd-3.2
clec-169

acox-1.2
col-176
B0286.3
F38B6.4
pfas-1

D

25°C WT   25°C endu-2(lf)

DAPI

fat-7 smFISH

fat-7 mRNA

E

FAT-7::GFP

WT 15°C   endu-2(lf) 15°C
WT 25°C   endu-2(lf) 25°C

F

FAT-7::GFP 25 °C

WT   endu-2(lf)
endu-2;Ex[endu-2]   endu-2;Ex[E454Q)]

WT
endu-2(lf)
endu-2;Ex[endu-2]
endu-2;Ex[endu-2(E454Q)]

Fig.6