Precision methylome and *in vivo* methylation kinetics characterization of *Klebsiella pneumoniae*

Jing Fu[1,2,#], Ju Zhang[2,#], Li Yang[2,3,#], Nan Ding[2,#], Liya Yue[2,#], Xiangli Zhang[2,3], Dandan Lu[2,3], Xinmiao Jia[2,4], Cuidan Li[2], Chongye Guo[2], Zhe Yin[5], Xiaoyuan Jiang[5], Yongliang Zhao[6,3], Fei Chen[2,3,*], Dongsheng Zhou[5,*]

[1] *Department of Oncology, Henan Provincial People's Hospital, People's Hospital of Zhengzhou University, People' s Hospital of Henan University, Zhengzhou, China*
[2] *CAS Key Laboratory of Genome Sciences and Information, Beijing Institute of Genomics, Chinese Academy of Sciences, and China National Center for Bioinformation, Beijing, China*
[3] *University of Chinese Academy of Sciences, Beijing, China*
[4] *Department of Medical Research Center, Peking Union Medical College Hospital, Peking Union Medical College & Chinese Academy of Medical Sciences, Beijing, China*
[5] *State Key Laboratory of Pathogen and Biosecurity, Beijing Institute of Microbiology and Epidemiology, Beijing, China*
[6] *CAS Key Laboratory of Genomic and Precision Medicine, Beijing Institute of Genomics, Chinese Academy of Sciences, and China National Center for Bioinformation, Beijing, China*

* Corresponding author.
  E-mail: chenfei@big.ac.cn (Chen F), dongshengzhou1977@gmail.com (Zhou D).
[#] Equal contribution.

Running title: *Fu J et al / Dynamic methylome of Klebsiella pneumoniae*

Abstract

Klebsiella pneumoniae (*K. pneumonia*) is an important pathogen that can cause severe hospital-/community-acquired infections. To panoramically analyze *K. pneumoniae*'s methylation features, we completed the whole genome sequences of 14 *K. pneumoniae* strains covering various serotypes, multilocus-sequence typings (MLSTs), clonal groups (CG), viscosity/virulence and drug-resistances, and further characterized their methylomes using Pacbio-SMRT and bisulfite technologies. We identified 15 methylation motifs (13 6mA and two 5mC motifs), among which eight were novel ones. Their corresponding MTases were further validated. Additionally, we analyzed the genomic distribution of GA<u>T</u>C and <u>C</u>CW<u>G</u>G methylation motifs shared by all strains, and identified differential distributive patterns of some hemi/un-methylated GA<u>T</u>C motifs tending to locate in the intergenic regions (IGRs). Specifically, we characterized the *in vivo* methylation kinetics at single-base resolution on a genome-wide scale by simulating the dynamic processes of replication-mediated passive demethylation and MTase-catalyzed re-methylation. The slower methylation-rates of the GA<u>T</u>C motifs in the replication origins (*oriC*) and IGRs suggest an epigenetic mechanism implicated in the regulation of replication-initiation and transcription. Our findings illustrate the first comprehensive dynamic methylome map of *K. pneumonia* at single base resolution, and provide an efficient means and important reference for a better understanding of epigenetic regulation in bacteria.

KEYWORDS: *Klebsiella pneumonia*; DNA methylation; epigenomics; 6mA; 5mC; Dam

Introduction

Klebsiella pneumonia (K. pneumoniae), the important member of the *Enterobacteriaceae*, can cause severe hospital- and community-acquired infections (*e. g.,* pneumonia, genitourinary tract infection, and septicemia). There are some typing methods for *K. pneumoniae* strains, such as serotype, multilocus-sequence typings (MLSTs), and clonal group (CG) [1,2]. The related studies indicated that the hypervirulence phenotype usually corresponded to K1/K2/K57 serotypes and CG23-ST23 [1-3], while multidrug-resistance (MDR) phenotype often corresponded to CG258- ST11/ST258 [4].

Several studies on DNA methylation of *K. pneumoniae* strains using molecular biological techniques have identified three DNA methylases (MTases) and corresponding motifs, including two restriction-modification (R-M) systems (M.KpnI: GG$\underline{T}$ACC; M.KpnBI: CAAAN$_6$R$\underline{T}$CA) and one orphan MTase (Dam: GA$\underline{T}$C) [5−7]. Further research on Dam indicated the epigenetic mechanism in regulating mismatch repair, virulence and pathogenicity of *K. pneumoniae* strains [8].

Recent rapid progress on high-throughput sequencing techniques, such as Pacific single-molecule real-time (SMRT) sequencing for accurate detection of modified bases (mainly 6mA) on a genome-wide scale and bisulfite sequencing for efficient analysis of genome-wide 5mC [9,10], has greatly facilitated investigations of DNA methylome in bacteria. It is well known that 6mA and 5mC are the two most important types of DNA methylation in prokaryote [9]. To date, many bacterial methylomes have been precisely determined using the above two techniques, including *E. coli*, *Mycoplasma genitalium*, *Bifidobacterium breve*, *Clostridium difficile*, *Campylobacter jejuni*, *Helicobacter pylori* [9−13] and *Mycobacterium tuberculosis* complexes (MTBC) reported by our group [14].

Through precisely and comprehensively analyzing the bacterial methylome, a lot of valuable information has been revealed, including methylation motifs and their corresponding methyltransferases, motif distributions on genomes, and the related epigenetic regulation mechanisms in bacteria [9,15,16]. Most identified MTases and the corresponding motifs belong to R-M system, which primary function is to prevent (cleave) the invading DNA and protect genomic DNAs through methylation-mediated mechanism [9−14]. Distinctively, some orphan MTases (no cognate restriction enzymes) and the corresponding motifs exert multiple epigenetic regulation functions

in bacteria [15–18]. Among them, Dam/GA**T**C motif is the most well-known because

of its prevalent existence in almost all the *Enterobacteriaceae* bacteria and

involvement in the epigenetic regulation of replication, transcription and mismatch

repair [15–17,19–21]. In particular, its regulatory role in replication initiation has

been studied in *E. coli*. Its *oriC* region contains five DnaA boxes and 11 GA**T**C motif

sites. The replication-mediated passive demethylation causes the hemi-methylated

GA**T**C motifs adjacent to the DnaA boxes that are specifically recognized and bound

by SeqA, further leading to the competitive occupation of the motif sites between

Dam and SeqA [17]. As a result, the re-methylation of the motifs is delayed, which in

turn prevents the initiation cascade for chromosome replication induced by *DnaA*

protein [17,19]. The re-methylation of the upstream GA**T**C motifs of the third and

fifth *DnaA* boxes are the rate-limiting steps for DNA replication initiation in *E.coli*

strains [17]. In addition, Dam also participates in the transcriptional regulation of the

downstream genes (*e. g*., *opvAB* in *Salmonella enterica*) [15].

Although several MTases and corresponding motifs have been revealed in *K.*

pneumoniae strains, the whole methylome has not been reported so far. Here, we

obtained whole-genome sequences of *K. pneumoniae* strains of 14 various types, and

then characterized their methylomes using SMRT combined bisulfite sequencing

techniques. Fifteen methylation motifs were identified (13 6mA and two 5mC

methylation motifs), including eight novel ones corresponding to eight novel MTases

termed as *kamA ~ G* (*K. pneumoniae* adenine methyltransferase A ~ G) and *kcmA* (*K.*

pneumoniae cytosine methyltransferase A). We further analyzed the distribution

pattern of the GA**T**C and C**C**W**G**G methylation motifs shared by all *K. pneumoniae*

strains. Importantly, by establishing a mathematical model to simulate the dynamic

process of passive demethylation and re-methylation of each motif in the exponential

phase, we characterized the genome-wide *in vivo* methylation kinetics at single-base

resolution. The motifs at different genomic locations showed different re-methylation

rates, and the GA**T**C motifs in *oriC* region and intergenic regions (IGRs) had slower

re-methylation rates. Our findings indicate potential roles of epigenetic regulation in

the replication initiation and transcription of *K. pneumonia* genome*,* and provide

important reference and insights into our better understanding of *K. pneumonia*

epigenomics.

Results

General bioinformatic analysis of 14 *K. pneumoniae* strains

We first obtained the whole genome sequences of 14 *K. pneumoniae* strains with various serotypes, MLSTs, CGs, viscosity/virulence and drug-resistances (Table S1, S5) by SMRT sequencing followed by correction using Illumina sequencing (**Figure 1**). The genome data have been deposited in National Genomics Data Center with the GSA accession number CRA003482 for strains of NTUH-K2044, 11492, 11420, 11454, 12208, 11311, 23, 11305, N201205880, 309074, 13190, 283747, 721005, 11021 and 11305. We then constructed a phylogenetic tree using 76 complete genomes of *K. pneumoniae* strains (14 from ours and 62 from online) (Figure S1, Table S5). Our 14 *K. pneumoniae* strains covered many common CGs and MLSTs of *K. pneumoniae* strains in China (*e. g*., CG23-ST23 and CG258-ST11) [22], indicating the well representativeness yet diversity of *K. pneumoniae* strains selected in our study.

The bioinformatic analyses provided the general genome information (**Table 1**, Figure 1), including genome size (5.20-5.54 Mb), GC content (57.4-57.9%), predicted protein-coding gene number (4990-5485), gene length (886-922 bp), and the percentage of coding region (88.26-90.23%) [23]. In addition, each *K. pneumoniae* strain contained 1-8 plasmids, with lower GC content (47.27-54.85%), lower percentage of coding region (75.98-88.63%), and shorter average gene length (596-831 bp) (Table 1).

Additionally, the relatively conserved genomic sequences and structures among the 14 *K. pneumoniae* strains were also indicated: ANI analysis revealed more than 99% identity among the 14 *K. pneumoniae* strains; no extensive translocations, duplications or inversions were found in the *K. pneumoniae* genomes except for strains of 11492 and 11454 each containing a large inverted fragment on the genomes (Figure S2).

Known and novel methylation motifs and corresponding methyltransferases in *K. pneumoniae* strains

A total of 13 6mA and two 5mC methylation motifs were identified in all *K. pneumoniae* strains (**Table 2-4**) by SMRT and bisulfite sequencing techniques (Table S2, S6), including seven known and eight novel methylation motifs (**Table 5**, Table

S10). It is worth noting that GA**T**C and CCW**G**G motifs were shared by all strains.

The other motifs were shared by at most two *K. pneumoniae* strains (Table 2-4).

Further analyses indicated the relationship between the motif and stain types (serotype,

MLST and CG). The MT**C**GAK motif existed in NTUH-K2044 and 11492 belonging

to K1 serotype and ST23-CG23, the common types of hypervirulent *K. pneumoniae*

strains. The CCA**A**YN$_7$**T**TYG motif was shared by two strains (11311 and 23) of

hypervirulent serotype K57 and ST412. The CCAGN$_7$R**T**TC motif was present in

strains 11305 and 13190 belonging to MDR CG147.

Modification analysis indicated that not all motif sites were fully-methylated

(methylated on both strands, Table 2-4). A minority of motif sites (< 30%) were

detected as being hemi-methylated (methylated on one strand only) or un-methylated

within the *K. pneumoniae* genomes. The only exception was the MT**C**GAK motif in

the NTUH-K2044 and 11492 genomes showing over half of hemi/un-methylated sites

(56.03-56.77%). Further analysis indicated that the un-methylated MT**C**GAK motif

prefered to have a guanine (G) in front of the motif (Figure S3, Table S11).

To search for the responsible MTases, we first predicted 22 MTases that might be

responsible for the 15 methylated motifs from *K. pneumoniae* genomes in the

REBASE database (http://rebase.neb.com/rebase/rebase.html) [24]. Among them,

seven MTases and their corresponding motifs had been verified in previous studies

(Table 5) [24]. To determine the MTases responsible for the eight newly detected

methylated motifs, we performed the restriction digestion and SMRT/bisulfite

sequencing using the plasmids containing the predicted MTase genes in

methyltransferase-free *E. coli* ER2796. Crossover validations identified the

corresponding eight MTases that could specifically recognize and methylate the

respective eight novel motifs (Table 5, Table S10, Figure S4-S5).

We further analyzed the distribution of 15 MTase genes on the genomes of the *K.*

pneumoniae strains. Thirteen genes were located on the chromosomes and two others

were located on the plasmids (**Figure 2**). In addition, within the 15 identified MTases,

there were 10 Type I MTases, three Type II MTases and two classical orphan MTases

(Dam and Dcm). Here *dam* and *dcm* genes were carried by all *K. pneumoniae* strains,

and responsible for the methylation of GA**T**C and CCW**G**G motif, respectively.

Among the three newly identified Type II MTases, KamC and KamD were predicted

to be Type IIG enzymes, which function of endonuclease and methyltransferase were

executed by a single gene (Table S10).

Differential distributions of GATC and CCWGG motifs on *K. pneumoniae*
genomes

Among the 15 methylation motifs, the GATC and CCWGG had the most extensive

distributions in all of the 14 *K. pneumoniae* strains, each containing ~30,000 GATC

and ~20,000 CCWGG modified sites (Table 2, 4). The distribution features of the

above two motifs on the genomes of *K. pneumoniae* strains were further analyzed,

and differential/uneven distribution on the genomes were observed (Figure 1). Both

motifs exhibited some high-density and low-density regions on the genomes, where

the genes were clustered into different COG functional categories (Figure S7).

Importantly, the GATC motif showed the highest distribution density (~34 sites/kb) in

the *oriC* region (average density: 5-6 sites/kb) of the 14 *K. pneumoniae* genomes

(Figure S8). In contrast, the CCWGG motifs didn't display such enrichment in the

oriC region.

We then compared the density distributions of GATC and CCWGG motifs on the

14 *K. pneumoniae* genomes and the simulated genome with the same base

composition (**Figure 3**, Figure S9-10). The result revealed that their distribution

densities on the *K. pneumoniae* genomes (1kb consecutive window) were higher than

those on the simulated genome, indicating the high-density/nonrandom distribution of

these two motifs on the *K. pneumoniae* genome. To explore the underlying causes for

their high-density distributions, we investigated the impact of selection pressure on

these two motifs by calculating the Ka/Ks ratios of the corresponding fragments in

GRs (more than 90% of the two motifs were located in GRs) (Figure S11). We

observed that the amino acid (AA) codons with two motifs (two AA codons for

GATC motif, two or three AA codons for CCWGG motif) were under strong

negative/purifying selection with Ka/Ks ≈ 0.09/0.09 in relative to the ratio of

0.39/0.54 for the scramble sequences in GRs.

Distribution patterns of the methylated, hemi-methylated and un-methylated
GATC and CCWGG motifs on *K. pneumoniae* genomes

215   We identified three methylation patterns (methylated, hemi-methylated and un-

216   methylated) for the GATC and CCWGG motifs. Most GATC (84.26-98.13%) and

CCWGG (77.58-98.84%) sites were detected as methylated state (Table 2, 4), whereas a small percentage of motif sites were hemi-methylated (< 15%) or un-methylated (< 10%) on the *K. pneumoniae* genomes. Further analysis demonstrated that the ratio of hemi-methylated GATC motifs were (~6.48%) much higher than that of un-methylated GATC motifs (~0.38%), while the hemi-methylated CCWGG motifs (~3.60%) showed a similar proportion as the un-methylated ones (~2.94%).

We then investigated the distribution ratio of methylated, hemi-methylated and un-methylated GATC and CCWGG motifs in gene regions (GRs) and IGRs. The hemi/un-methylated GATC motifs preferred to locate in IGRs, since its ratio (7.14%/34.11%) in IGRs was significantly higher than that of methylated GATC motifs (5.23%) ($P < 0.01$, **Figure 4**A, Table S12). The analysis of the "fraction of methylated reads" (FRAC value) for the motifs in GRs/IGRs also supported the above findings that the hemi/un-methylated GATC motifs were inclined to locate in the IGRs (Figure 4B). In contrast, the hemi/un-methylated CCWGG motifs (~7.49%) showed a little lower distribution ratio in IGRs compared with the methylated CCWGG motifs (~7.68%, Table S13).

The analysis of the GATC motifs' density in 5' upstream region (USR), coding sequence (CDS) and 3' downstream region (DSR) also supported the above conclusion: the methylated motifs displayed a higher density in GRs, while the hemi-/un-methylated motifs presented a higher density in IGRs (both 5' USR and 3' DSR) (Figure 4C). We further explored the hemi/un-methylated GATC sites shared in the 5' USR regions of 14 *K. pneumoniae* strains: 13 hemi/un-methylated GATC sites corresponding to 11 genes were observed (Table S14). We also detected 12 high-density hemi/un-methylated clusters (no less than three consecutive hemi/un-methylated motifs in at least two strains) in the 5' USR regions (Table S15).

We then analyzed the sequence conservation of the DNA fragments (20 nt on both sides) containing hemi/un-methylated or methylated motifs to determine the causes leading to their differential distributions in IGRs. The results showed that the fragments with hemi/un-methylated GATC motifs (44 nt) exhibit higher conservation than those with methylated GATC motifs in IGRs (Figure S12).

Methylation kinetic characteristics of GATC and CCWGG motifs during growth cycle of *K. pneumoniae* strains

Since there is no demethylase in bacteria, the *in vivo* methylation kinetics

characterization is based on the dynamic equilibrium between the replication-

mediated passive demethylation and MTases-catalyzed re-methylation [18]. To

explore the features of methylation kinetics of GA**T**C and CC**W**GG motifs during the

growth cycle, we first characterized the methylomes of the NTUH-K2044 and 11492

strains at the exponential phase (1 h), transition-to-stationary phase (4 h) and

stationary phase (24 h) (**Figure 5**A). By comprehensively analyzing genome-wide

sequencing coverages and the fractions of methylated reads (methylated-read

ratio/FRAC value), GA**T**C and CC**W**GG motifs were found to exhibit distinct kinetic

features during the growth cycle (Figure 5B, Figure S13). As for GA**T**C motifs, the

methylated-read ratios were more than 90% throughout the genomes in the three

phases (Figure 5B), although the sequencing coverages varied on the genomes during

the growth cycles (Figure S13), indicating that the GA**T**C motifs could be re-

methylated in a very short time after the passive demethylation caused by replication:

i.e., the re-methylated rate was almost identical to the passive demethylation rate.

In contrast, the CC**W**GG motif showed a much slower re-methylation rate. First, in

the exponential phase (1 h), the methylated-read ratio of CC**W**GG motifs in the *oriC*

regions ($55.66 \pm 18.54\%$) was much lower than that in the replication termination (*Ter*)

regions ($82.63 \pm 12.71\%$) (Figure S13B). Secondly, the average methylated-read ratio

in the transition-to-stationary phase ($54.50 \pm 27.14\%$, 4 h) was close to that of the

exponential phase ($60.50 \pm 17.71\%$, 1 h), but not the stationary phase (80.28 ± 19.05

%, 24 h) (Figure S13B). This finding suggested a much slower re-methylation rate

compared to the replication-mediated passive demethylation rate for CC**W**GG motifs,

which resulted in the above differences in the methylated-read ratios.

To quantify the re-methylation time per motif in the exponential phase, we first

investigated the replication kinetics of the aforementioned two *K. pneumoniae* strains.

Figure S13 showed that the genome sequencing coverage in the *oriC* regions was

more than twice as much as that of the *Ter* regions (2.93 and 3.02 folds for the 11492

and NTUH-K2044 strains, respectively), indicating that the next round of replication

was re-initiated before the replication termination in the exponential phase. We then

constructed a replication model (Figure 5C) by fitting genome coverage data, and

obtained the ratio (t_D/t_R) of doubling time (t_D) to replication termination time (t_R) (see

"Materials and methods" for detail). The doubling time (t_D) in the exponential phase

also reflects the re-initiation time. Interestingly, we obtained the same t_D/t_R values

(0.59) for the two *K. pneumoniae* strains (Figure 5D), suggesting a similar regulatory

mechanism for replication cycle, which should be intrinsically carried by *K.*

pneumoniae strains. Two t_D values were then calculated through fitting the growth

curves (~34.64 min for strain 11492; ~27.95 min for NTUH-K2044, Figure 5A),

therefore we could infer the t_R from the t_D/t_R values (~58.71 and ~47.36 min for strain

11492 and NTUH-K2044, respectively).

We further obtained the re-methylation time per motif (**Table 6**) through simulating

the dynamic processes of passive demethylation and re-methylation in the exponential

phase based on the five parameters, such as methylation-read fraction of each motif

($M_{(x)}$), initial methylation-read fraction of each motif ($M_{(x)}^0$), t_D, t_R and distribution

density of the first replication forks ($P_{(x_1)}$, see "Materials and methods" for detail).

In general, the mean re-methylation time of 6mA was shorter (3.52 and 3.46 min

for GA<u>T</u>C and GRA<u>C</u>RAC motifs, respectively) than that of 5mC (9.23 and 4.55 min

for C<u>C</u>W<u>G</u>G and MT<u>C</u>GAK motifs, respectively) in the exponential phase (Table 6).

As for a certain motif, the re-methylation time in the NTUH-K2044 strain was shorter

than that in the 11492 strain (Figure S14). In addition, the flanking bases could

influence the re-methylation rates of the GA<u>T</u>C and C<u>C</u>W<u>G</u>G motifs. When the

flanking bases were C/G rather than T/A, GA<u>T</u>C motif showed a faster re-methylation

rate, which was totally reversed for the C<u>C</u>W<u>G</u>G motif (Table 6, Figure S15).

Slower re-methylation rate of the GA<u>T</u>C motifs in intergenic and *oriC* regions at

the exponential phase

To investigate the role of GA<u>T</u>C motifs in transcriptional regulation of *K. pneumoniae*

strains, we analyzed the re-methylation rates of the motifs in IGRs (including 5' USRs

and 3' DSR) at the exponential phase (**Figure 6**A). The results indicated the slower

re-methylation motifs in IGR regions (3.94 ± 5.82 min) than that in GR regions (3.50

± 5.10 min). Since most 5' USRs in bacteria are overlapped with the promoter regions

and involved in the transcriptional regulation [25], we further explored the COG

functional category of genes with top 5% slow re-methylation sites (> 7.04 min/motif)

in 5' USRs (Table S16). Four enriched functional categories ("Cell cycle control, cell

division, chromosome partitioning", "Carbohydrate transport and metabolism",

"Intracellular trafficking, secretion, and vesicular transport" and "Translation,

ribosomal structure and biogenesis") were observed (Table S16), of which at least 5%

of genes contained a slow re-methylation sites (> 7.04 min/motif) in 5' USRs. Among

them, the "carbohydrate transport and metabolism" functional category owned the

most genes (22 genes) with slow re-methylation rates in 5' USRs, including three

hemi/un-methylated motifs shared in 14 *K. pneumoniae* strains and even one hemi/un-

methylated motif-clusters (Table S16).

To further explore the role of GA**T**C motifs in replication regulation of *K.*

pneumoniae strains, we analyzed the re-methylation rates of the motifs in *oriC*

regions (Figure 6B). The GA**T**C motifs in *oriC* region had the highest distribution

density (~34 sites/kb) (Figure S8) and slower re-methylation rates (10.35 ± 8.69 min).

Among the 14 GA**T**C motifs in *oriC* region, eight and seven motifs with slow re-

methylation rates (> 7.04 min/motif) were enriched in the upstream sequence of the

fourth DnaA binding site (DnaA box) and the adjacent AT-rich region in strain 11492

and NTUH-K2044, respectively (Figure S16).

Discussion

In this study, we characterized the precision methylome of 14 *K. pneumoniae* strains

with different serotypes, MLSTs, CGs, viscosity/virulence and drug-resistances, using

SMRT and the bisulfite sequencing techniques, and identified 15 DNA methylation

motifs (eight novel ones) including 13 6mA motifs and two 5mC motifs

corresponding to 13 R-M system MTases and two orphan MTases, respectively. Two

motifs (GA**T**C and C**C**W**G**G) and their respective orphan MTases (Dam and Dcm)

appeared to be the most important since they were present in all *K. pneumoniae*

strains with the most extensive distributions on genomes in relative to other modified

motifs (Table 2, 4). This feature was also reported in almost all members of the

Enterobacteriaceae family [21,24]. Functional analysis in previous reports indicated

that these two motifs, especially GA**T**C, could exert multiple functions including

transcriptional regulation, cell-cycle controlling and mismatch repair in *E.coli* and *S.*

enterica [15−17,19−21]. We also demonstrated their high-density distributions on *K.*

pneumoniae genome comparing with that on simulated genome (Figure 3, Figure S9-

10), which was probably caused by purifying selection (Figure S11) leading to their

evolutionary conservation (Figure S12) as reported in other bacterial genome [26].

This well-conserved feature also implies an essential role of these two motifs in *K.*

pneumoniae.

The hemi/un-methylated GA<u>T</u>C motifs were found to possess the tendency of IGRs

(including 5' USR and 3' DSR) localization (Figure 4A, C). Since promoters in *K.*

pneumoniae were also predicted to be located across the 5' USR regions (Figure S17),

thus, these hemi-/un-methylated GA<u>T</u>C motifs in 5' USR might be protected from the

methylation by competitively binding of some regulators to the promoter regions [17,

19]. In *E.coli* strains, this feature facilitates the epigenetic regulation of the

downstream gene expression [27]. The status of these motifs suggested similar

epigenetic mechanisms in *K. pneumoniae* strains, and might be the consequence of

selection during long-term evolution (Figure S11), which was supported by our

findings that the genomic fragments with hemi/un-methylated GA<u>T</u>C motifs in IGRs

had higher sequence conservation (Figure S12).

Importantly, by establishing a mathematical model to simulate the dynamic process

of passive demethylation and re-methylation of each motif in the exponential phase,

we obtained the re-methylation time of each motif throughout the whole genome

(Table 6, Figure S14). Our studies revealed that the motifs at different genomic

locations showed different re-methylation rates. We could reasonably infer that the

slower re-methylation rates in some sites/regions might also be due to the competitive

binding of certain proteins in order to prevent methylation [17, 19]. Thus, the slow re-

methylation rates could precisely reflect the methylation-mediated epigenetic

regulations at these sites/regions *in vivo*.

There are two methylation-mediated epigenetic regulations: transcription and

replication, which were examined by methylation kinetic analysis in our study (Figure

6, Figure S16). Firstly, transcriptional regulation analysis indicated that the GA<u>T</u>C

motifs in IGRs presented the slower re-methylation rates than those in GRs (Figure 6).

This is in agreement with the distribution characteristics of the GA<u>T</u>C motifs in IGRs:

lower methylated-read ratios and more hemi/un-methylated motifs (Figure 4A, 4B).

As described above, most IGRs in bacteria overlap with the promoter regions and

participate in the transcriptional regulation [25]. Thus, the promoter regions with slow

re-methylation motifs should be the locations where the GA<u>T</u>C motifs with hemi/un-

methylated status function as transcription regulators in *K. pneumoniae* strains.

Similarly, this should also be the consequence of the competitive binding between

Dam and some transcription regulatory proteins (such as OxyR) to these sites/regions

[15,27,28], which is analogous to the epigenetic transcriptional regulation mediated

by the competition between DNA methyltransferases and CTCF in CpG islands of

eukaryotic cells [29]. Secondly, replication regulation analysis identified 7 - 8 slow

GA<u>T</u>C motifs enriched in the upstream sequence of the fourth DnaA binding site

(DnaA box) and the adjacent AT-rich region (Figure S16). SeqA has been shown to

prefer to bind to these motifs for the purpose of lowering re-methylation rates [30]

and preventing the initiation cascade for chromosome replication [17,30]. Therefore,

re-methylation of these motifs in the fourth DnaA box and AT-rich regions should

represent the main rate-limiting steps for triggering initiation of DNA replication in *K.*

pneumoniae strains. Our findings also uncovered many promoter regions with slower

re-methylation motifs in *K. pneumoniae* strains (Figure 6), therefore, it is reasonably

speculated that epigenetic regulation in bacteria is very complex rather than simple as

we believed.

Previous studies on MTase kinetics mainly focused on *in vitro* analysis [31−33].

Several reported analyses about the *in vivo* methylation kinetics, including one

publication about bacteria, only studied the overall, but not single-motif methylation

kinetics [34−36]. Our study is the first to characterize the *in vivo* methylation kinetics

at single motif resolution in whole genome, thus, offer an efficient means and

valuable resources for a better understanding of epigenetic regulation in bacteria.

Materials and methods

Strain information, growth curve, and phenotypic characterization

The information of 14 *K. pneumoniae* strains used in this study was shown in Table

S1. The strains were cultured overnight in Luria-Bertani (LB) medium at 37 °C. 1 ml

of overnight culture suspensions was then transferred to a flask containing 200 ml of

LB medium and cultured in a shaker at 200 rpm. The growth curve of bacteria was

determined by recording OD_{600} values at different time-points. This experiment was

performed in triplicate.

Drug susceptibility test was performed by VITEK 2, and the drug-resistance

phenotypes were determined by Clinical and Laboratory Standards Institute (CLSI)

standards (https://clsi.org/standards/products/microbiology/documents/m100/). The

string test of *K. pneumoniae* strains was conducted, and hypermucoviscosity was

defined by viscous strings with a length of more than 5 mm [37].

Genomic DNA Extraction, sequencing, assembly, correction and annotation

The genomic DNA was extracted using TIANamp Bacteria Genomic DNA Kit (Tiangen Biotech Co. Ltd., Beijing, China). The whole-genome sequencing was implemented by PacBio RSII platform using P6/C4 chemistry. Each strain was sequenced with 1-2 SMART cells with genome coverage of more than 50× (Table S2−S3).

De novo assembly of the genome was performed using Hierarchical Genome Assembly Process 3 (HGAP3) in SMRT Portal (version 2.2.0). Gap closing was completed by PBJelly [38]. On the basis of Blast results, genome circularization was finished by manually removing the contigs overlapped regions.

To correct the polymer-errors, we re-sequenced the strains using Illumina sequencing (Table S4). Paired-end libraries were prepared as previous description, and clean reads were obtained after eliminating redundant and low-quality raw reads. Paired reads were extracted and then mapped onto the assembled genome sequences to obtain unique mapped reads using BWA. Pilon v.1.13 was subsequently used to polish genome sequences using unique map reads.

Genome sequences were annotated in the Subsystem Technology (RAST) pipeline with Rapid Annotation [39]. Unannotated genes were then predicted by alignment in the NCBI non-redundant (NR) database using BLAST. The protein functions were annotated by Clusters of Orthologous Groups (COG). tRNAs and rRNAs were predicted by tRNAscan-SE and "search_for_rnas" tools, respectively.

Genome structure and phylogenetic analysis

The average nucleotide identity (ANI) and coverage were calculated by ANI on EzGenome (http://www.ezbiocloud.net/tools/ani) and online BLAST (https://blast.ncbi.nlm.nih.gov/Blast.cgi). Multiple alignments of genomic sequences were performed using the Mauve multiple alignment software [23].

Single nucleotide polymorphisms (SNPs) were detected by MUMmer based on the 14 genomes in this study and 62 published genomes (Table S5) using HS11286 as reference. Prank [40] was used to annotate the protein coding genes of the 76 *K. pneumonia* genomes, and then used Roary to predict 3173 core genes[41]. The SNPs (117,142 SNPs) in core genes were used to construct a phylogenetic tree on the basis

of maximum likelihood using FastTree [42], followed by decoration using software

evolview v2 (http://www.evolgenius.info/evolview/) [43].

Genome-wide detection of 6mA and the related motifs using the SMRT

sequencing data

The SMRT Portal (version 2.2.0) was applied to detect genome-wide 6mA

modification and related motifs using the standard settings in the

RS_Modification_and_Motif_Analysis.1 protocol as previously described

(https://smrt-analysis.readthedocs.io/en/latest/SMRT-Pipe-Reference-Guide-v2.2.0/).

We then identified the motifs through selecting the top 1000 kinetic hits and subjected

a window of ± 20 base around the detected modified base to MEME-ChIP [44],

followed by comparing them with the predicted MTase targeting motif sequences in

REBASE [24].

Three methylation patterns (methylated, hemi-methylated and un-methylated) were

identified for DNA motifs within the *K. pneumoniae* genome. A methylated motif is

the sites were fully-methylated on both strands, whereas the hemi-methylated motif is

the sites were methylated on one strand only. The un-methylated motif is the sites

were not modified on both strands. A methylated site was defined as methylated-

read ratio > 0.8. The "methylated-read ratio" is the percentage of methylated reads in

the total number of reads mapped to the sites, which was obtained from "frac =

decimal" in "modification.gff" file.

Genome-wide detection of 5mC and the related motifs using the bisulfite

sequencing data

The 5mC methylation was detected by bisulfite sequencing as previously reported

(Table S6, S7). Trimmomatic (v0.32) [45] was used to trim adapters and low-quality

bases using default parameters. Clean reads were mapped on the reference genomes

by Bismark (version 0.12.2) [46]. We then identified the motifs through subjecting a

window of ± 20 base around the detected modified base to MEME-ChIP [44],

followed by comparing with the predicted MTase targeting motif sequences in

REBASE [24].

MTases cloning and verification

The predicted MTase genes were amplified from bacterial genomic DNAs using the

gene-specific primers (Table S9), and cloned into plasmid pRRS as previously

described [14]. The corresponding methylation sensitive restriction-site sequences

(used to detect the activity of MTases) were included in the 3'-end oligonucleotides.

The recombinant plasmids were then transformed into bacterial host ER2796 (not

containing known MTase genes), followed by bacterial culture overnight. The plasmid

DNAs were prepared using Qiagen Miniprep kit. The appropriate restriction enzymes

were then used to determine the presence or absence of methylated motifs in

recombinant plasmid. The digestion reaction were conducted for 4 h at 37 ºC and run

on 1% agarose gels. The methylation motifs were also double detected by sequencing

the recombinant plasmid by SMRT/bisulfite sequencing.

Density distributions of GATC and CCWGG motifs on the _K. pneumoniae_ and

simulated genomes

The simulated genome was generated based on the same length and GC content as

those in the _K. pneumoniae_ genomes. We then calculated the number of motifs in 1kb

consecutive segments, so as to determine the density of GATC and CCWGG motifs

across the _K. pneumoniae_ and simulated genomes.

We counted the number of motifs in each 2kb-size non-overlap sliding window

along the genome sequence of _K. pneumoniae_ strains. By fitting the in normal

distribution in R using the "pnorm" function, top 5% regions with higher motif counts

were selected as high-density regions, and the tail 5% regions with fewer motif counts

were selected as low-density regions.

Ka/Ks analysis of the GATC and CCWGG "motif sequences" and corresponding

"scramble sequences"

We firstly extracted the minimum DNA sequences (2~3 codons) containing the motifs

from the open reading frames of genes, in order to obtain "motif sequences". The

corresponding "scramble sequences" were obtained by random shuffling (eliminating

the "motif sequences"). The top-10 frequent "scramble sequences" were used in

subsequent analysis. We then identified the reference sequences of "motif sequences"/

"scramble sequences" on reference genome (HS11286) by multiple sequence

alignment (mafft) [47]. The "motif sequences", "scramble sequences" and

corresponding reference sequences of each strain were respectively concatenated, and

their Ka/ks ratios were calculated by paraAT [48].

Sequence conservation analysis of the GA<u>T</u>C and CCW<u>G</u>G motifs

We extracted the methylated motifs and their flanking sequences (20 nt) in IGRs from

one genome, and then obtained the corresponding sequences of 13 other genomes by

multiple sequence alignment. Using the above 14 sequences, the conservation scores

were calculated by phylop [49]. The conservation scores of the hemi-/un-methylated

motif with 20-nt flanking sequences in IGRs were calculated by the same method.

Calculation of the genomic replication time (t_R) through simulating genomic

coverage plots

The genomic coverage plots reflect the accumulated copy numbers across genomes.

Since the replication forks always proceed from origin of replication (*oriC*) to

doubling point (s_{DP}), only a few ones can get close to the doubling point in the

exponential phase. As a result, the sequencing coverage in *oriC* region is much higher

than that in doubling point. In each cell, the copy number of each site (*s*) depends on

its relative position to the first replication fork (s_1) and the minimal time of successive

initiations (re-initiation). The re-initiation time is consistent with the doubling time (t_D)

in the exponential phase, reflecting the growth rate of *K. pneumoniae* strains.

After initiation, the replication forks should go forward bi-directly at similar speeds

in the exponential phase. Therefore, the genome coverage plots are symmetrical near

the doubling point (s_{DP}). We divided the coverage plots into the left ($s \leq s_{DP}$) and

right ($s \geq s_{DP}$) parts $s \leq s_{DP} s \geq s_{DP}$ by doubling point, and simulated the curves

independently. We then obtained the relative position ($x \in (0,1]$) of each genomic site

shown as equation (1). S_G represents the half length of genome.

540
$$\begin{cases} x = s/S_G & (s \leq s_{DP}) \\ x = 2 - s/S_G & (s \geq s_{DP}) \end{cases} \tag{1}$$

541   We subsequently calculated the relative distance of the first replication fork when

542   another one at *oriC* is re-initiated (Δx_R) as equation (2). t_R represents the genomic

543   replication time.

544
$$\Delta x_R = t_D/t_R \qquad (0 \leq t_D \leq t_R) \tag{2}$$

By constructing the replication frequency matrix of genomic sites (*x*) with different

distances from the first replication fork (x_I), the copy number of the genomic site

($f_{(x,x_1)}$) can be determined as equation (3). When the genomic site is in front of the

first replication fork ($x \geq x_1$), its copy number is one; when the site is between the

first and second replication forks ($x_1 - \Delta x_R \leq x \leq x_1$), its copy number is two. The

rest was deduced by analogy.

$$f_{(x,x_1)} = 2^{(x_1-x)/\Delta x_R} \tag{3}$$

Next, we used the B distribution to assess cell population density with different

genomic positions of the first replication forks ($x_1 \in [0, 1]$). Since x is not continuous

data (with the step size of $1/S_G$), the cell population density of the first replication

forks ($P_{(x_1)}$) at each genomic site was determined by the difference of accumulated

densities ($I_x(\alpha, \beta)$) of the adjacent sites.

$$P_{(x_1)} = I_{(x_1)}(\alpha, \beta) - I_{\left(x_1 - \frac{1}{S_G}\right)}(\alpha, \beta) \tag{4}$$

The genome coverage plots could be determined as the integral of the copy

numbers of each genomic site in the cellular populations ($H(x)$) as shown in the

equation (5). The integral was achieved based on $P_{(x_1)}$ and the relative distance

between x and x_1 ($x_1 - x$). By substituting equation (2)-(4) into equation (5), we

established the mathematical model to fit the genome coverage plots as equation (6).

$$H(x) = \int_0^1 f_{(x,x_1)} * P_{(x_1)} d_{x_1} \tag{5}$$

$$H(x) = \int_0^1 2^{(x_1-x)*t_R/t_D} * \left(I_{(x_1)}(\alpha, \beta) - I_{\left(x_1 - \frac{1}{S_G}\right)}(\alpha, \beta) \right) d_{x_1} \tag{6}$$

We then substituted equation (1) into equation (6), and repeatedly fit the genome

coverage plots by selecting continuous parameters with step sizes of 0.1. We finally

obtained the optimal solutions of Δx_R and B distribution parameters (α,β) through

Goodness of Fit Test. Since t_D could be calculated from growth curve of each *K.*

Pneumoniae strain, we could obtain the replication time (t_R) by the equation (2).

Calculation of the re-modification time of motifs

In the exponential phase, methylation-read fraction ($M_{(x)}$) of each motif was

determined by its initial methylation-read fraction ($M_{(x)}^0$), replication-induced passive

de-methylation and methyltransferase-catalyzed re-methylation. Based on $M_{(x)}$ in the

exponential phase and B distribution parameters (α,β), we calculated the proceeding

distance of the corresponding replication fork when the hemi-methylated motif was

re-methylated (Δx_M). As shown in the equation (7), Δx_M corresponds to the ratio of

mean re-methylation time (t_M) to the genomic replication time (t_R).

$$\Delta x_M = t_M / t_R \tag{7}$$

When the motif x is located at downstream of replication fork ($x_j \leq x$), its

methylation-read fraction ($M_{(x)}$) remain unchanged because of no replication-induced

passive de-methylation. When the motif x is located at upstream of replication fork

with a longer distance ($x_j \geq x + \Delta x_M$), the methylation-read fraction is also

unchanged because of the completion of re-methylation of motif x. Thus, we only

considered the de-methylation effect of the replication forks in a certain range

($x \leq x_j \leq (x + \Delta x_M)$) on $M_{(x)}$ in the mathematical derivation. j represents the serial

number of the replication fork affecting $M_{(x)}$.

If the hemi-methylated motifs can be fast re-modified before next replication fork

($t_M \leq t_D$), we only need to evaluate the de-methylation effect of one replication for

all replication forks in the range of $x \leq x_j \leq x + \Delta x_M$. Here we include the first

replication forks (x_1) and other effective replication forks (x_j) in the range. We then

substituted the accumulated B distribution density of first replication forks ($x +$

$(j-1)\Delta x_R \leq x_1 \leq x + \Delta x_M + (j-1)\Delta x_R$) for that of the corresponding effective

replication forks ($x \leq x_j \leq x + \Delta x_M$). Due to semi-conservative replication, we

further obtain Δx_M from equation (8). N represents the upper limit of j, which is

determined by Δx_R and the relative genomic position (x) of methylated motifs.

$$\begin{cases} M(x) = M_{(x)}^0 * \left(1 - 0.5 * \sum_{j=1}^{N} \begin{pmatrix} I_{(x+\Delta x_M+(j-1)*\Delta x_R)}(\alpha, \beta) - \\ I_{(x+(j-1)*\Delta x_R)}(\alpha, \beta) \end{pmatrix}\right) \\ N = \lceil (1-x)/\Delta x_R \rceil \end{cases} \tag{8}$$

If the methylation rate is slow, the hemi-methylated motifs may not be re-modified

when the next replication fork crosses them ($t_M > t_D$). In this case, the effect of

multiple passive de-methylation on $M_{(x)}$ should be considered. n represents upper

limit of multiple passive de-methylation times (i) as equation (9), which is determined

by the ratio of mean re-methylation time (t_M) to re-initiation time (t_D).

$$n = \lceil t_M / t_D \rceil = \lceil \Delta x_M / \Delta x_R \rceil \tag{9}$$

We then converted genomic position of effective replication forks (x_j) into that

of the corresponding first replication forks (x_1), and substituted accumulated B

distribution density of x_1 ($x + (j-1) * \Delta x_R \leq x_1 \leq x + j * \Delta x_R$) for the

x_j population density ($x \leq x_j \leq x + \Delta x_M$). Considering the differential influences

on $M_{(x)}$ with the case of $j < n$ and $n \leq j \leq N$, we calculated Δx_M as the equation

(10) by comprehensive calculations.

$$
\begin{cases}
M(x) = M_{(x)}^0 * \begin{pmatrix} 1 - \sum_{j=1}^{n-1} \sum_{i=1}^{j} 0.5^i * \begin{pmatrix} I_{(x+j*\Delta x_R)}(\alpha,\beta) - \\ I_{(x+(j-1)*\Delta x_R)}(\alpha,\beta) \end{pmatrix} \\ - \sum_{j=n}^{N} \sum_{i=1}^{n-1} 0.5^i * \begin{pmatrix} I_{(x+j*\Delta x_R)}(\alpha,\beta) - \\ I_{(x+(j-1)*\Delta x_R)}(\alpha,\beta) \end{pmatrix} \\ - 0.5^n * \sum_{j=n}^{N} \begin{pmatrix} I_{(x+\Delta x_M+(j-n)*\Delta x_R)}(\alpha,\beta) - \\ I_{(x+(j-1)*\Delta x_R)}(\alpha,\beta) \end{pmatrix} \end{pmatrix} \\[6pt]
n = \lceil t_M/t_D \rceil = \lceil \Delta x_M/\Delta x_R \rceil \\[4pt]
N = \lceil (1-x)/\Delta x_R \rceil
\end{cases}
\tag{10}
$$

Based on the growth curve and genome coverage plots of the sequencing data, we

deduce some parameters in different *K. pneumoniae* strains, such as t_D, t_R and

$I_x(\alpha, \beta)$. The mean re-methylation time (t_M) of each motif could be further calculated

by substituting the above parameters, the equation (2) and (7), as well as the

methylation-read fraction in exponential phase ($M_{(x)}$) and stationary phase ($M_{(x)}^0$) into

the equation (10).

Data availability. The genome data have been submitted to GSA at the National

Genomics Data Center (CRA003482).

CRediT author statement

Jing Fu: Validation, Writing - Original Draft, Resources. **Ju Zhang**: Methodology,

Writing - Original Draft, Resources. **Li Yang**: Software, Formal analysis, Writing -

Original Draft. **Nan Ding**: Formal analysis, Writing - Original Draft, Visualization.

Liya Yue: Validation, Writing - Original Draft, Visualization. **Xiangli Zhang**: Formal

analysis, Visualization. **Dandan Lu**: Formal analysis, Visualization. **Xinmiao Jia**:

Visualization. **Cuidan Li**: Visualization, Data Curation. **Chongye Guo**: Visualization.

Zhe Yin: Resources. **Xiaoyuan Jiang**: Resources. **Yongliang Zhao**: Writing - Review

& Editing. **Fei Chen**: Conceptualization, Writing - Review & Editing, Funding

acquisition, Project administration, Supervision. **Dongsheng Zhou**: Conceptualization,

Writing - Review & Editing, Project administration, Supervision.

Competing interests

The authors have declared no competing interests.

Acknowledgements

This work was supported by the National Key Program for Infectious Disease of China [2018ZX10302301-004-003] and the National Natural Science Foundation of China [31770870, 31801093].

ORCID

ORCID: 0000-0001-7961-0888 (Fu, J)

ORCID: 0000-0003-3274-9844 (Zhang, J)

ORCID: 0000-0001-7353-4415 (Yang, L)

ORCID: 0000-0002-1045-1695 (Ding, N)

ORCID: 0000-0002-5752-6914 (Yue, L)

ORCID: 0000-0003-4816-6598 (Zhang, X)

ORCID: 0000-0001-6977-4301 (Lu, D)

ORCID: 0000-0002-6084-0146 (Jia, X)

ORCID: 0000-0002-8430-1155 (Li, C)

ORCID: 0000-0002-1918-8643 (Guo, C)

ORCID: 0000-0003-0530-7100 (Yin, Z)

ORCID: 0000-0002-8962-7170 (Jiang, X)

ORCID: 0000-0003-0121-1312 (Zhao, Y)

ORCID: 0000-0003-3895-6318 (Chen, F)

ORCID: 0000-0003-3827-5969 (Zhou, D)

References

[1] Guo Y, Wang S, Zhan L, Jin Y, Duan J, Hao Z, et al. Microbiological and clinical characteristics of hypermucoviscous *Klebsiella pneumoniae* isolates associated with invasive infections in China. Front Cell Infect Microbiol 2017;7:24.

[2] Lam MMC, Wyres KL, Duchene S, Wick RR, Judd LM, Gan YH, et al. Population genomics of hypervirulent *Klebsiella pneumoniae* clonal-group 23 reveals early emergence and rapid global dissemination. Nat Commun 2018;9:2703.

[3] Yeh KM, Kurup A, Siu LK, Koh YL, Fung CP, Lin JC, et al. Capsular serotype K1 or K2, rather than magA and rmpA, is a major virulence determinant for *Klebsiella pneumoniae* liver abscess in Singapore and Taiwan. J Clin Microbiol 2007;45:466–71.

[4] Bialek-Davenet S, Criscuolo A, Ailloud F, Passet V, Jones L, Delannoy-Vieillard AS, et al. Genomic definition of hypervirulent and multidrug-resistant *Klebsiella pneumoniae* clonal groups. Emerg Infect Dis 2014;20:1812–20.

[5] Hammond AW, Gerard GF, Chatterjee DK. Cloning the KpnI restriction-modification system in *Escherichia coli*. Gene 1991;97:97–102.

[6] Valinluck B, Lee NS, Ryu J. A new restriction-modification system, KpnBI, recognized in *Klebsiella pneumoniae*. Gene 1995;167:59–62.

[7] Mehling JS, Lavender H, Clegg S. A Dam methylation mutant of *Klebsiella pneumoniae* is partially attenuated. FEMS Microbiol Lett 2007;268:187–93.

[8] Fang CT, Yi WC, Shun CT, Tsai SF. DNA adenine methylation modulates pathogenicity of *Klebsiella pneumoniae* genotype K1. J Microbiol Immunol Infect 2017;50:471-7.

[9] Fang G, Munera D, Friedman DI, Mandlik A, Chao MC, Banerjee O, et al. Genome-wide mapping of methylated adenine residues in pathogenic *Escherichia coli* using single-molecule real-time sequencing. Nat Biotechnol 2012;30:1232–9.

[10] Bottacini F, Morrissey R, Roberts RJ, James K, van Breen J, Egan M, et al. Comparative genome and methylome analysis reveals restriction/modification system diversity in the gut commensal *Bifidobacterium breve*. Nucleic Acids Res 2018;46:1860–77.

[11] Lluch-Senar M, Luong K, Llorens-Rico V, Delgado J, Fang G, Spittle K, et al. Comprehensive methylome characterization of *Mycoplasma genitalium* and

Mycoplasma pneumoniae at single-base resolution. PLoS Genet 2013;9:e1003191.

[12] Hargreaves KR, Thanki AM, Jose BR, Oggioni MR, Clokie MR. Use of single molecule sequencing for comparative genomics of an environmental and a clinical isolate of *Clostridium difficile* ribotype 078. BMC Genomics 2016;17:1020.

[13] Murray IA, Clark TA, Morgan RD, Boitano M, Anton BP, Luong K, et al. The methylomes of six bacteria. Nucleic Acids Res 2012;40:11450−62.

[14] Zhu L, Zhong J, Jia X, Liu G, Kang Y, Dong M, et al. Precision methylome characterization of *Mycobacterium tuberculosis complex* (MTBC) using PacBio single-molecule real-time (SMRT) technology. Nucleic Acids Res 2016;44:730−43.

[15] Cota I, Bunk B, Sproer C, Overmann J, Konig C, Casadesus J. OxyR-dependent formation of DNA methylation patterns in OpvABOFF and OpvABON cell lineages of *Salmonella enterica*. Nucleic Acids Res 2016;44:3595−609.

[16] Cohen NR, Ross CA, Jain S, Shapiro RS, Gutierrez A, Belenky P, et al. A role for the bacterial GATC methylome in antibiotic stress survival. Nat Genet 2016;48:581−6.

[17] Wolanski M, Donczew R, Zawilak-Pawlik A, Zakrzewska-Czerwinska J. oriC-encoded instructions for the initiation of bacterial chromosome replication. Front Microbiol 2014;5:735.

[18] Kozdon JB, Melfi MD, Luong K, Clark TA, Boitano M, Wang S, et al. Global methylation state at base-pair resolution of the Caulobacter genome throughout the cell cycle. Proc Natl Acad Sci U S A 2013;110:E4658−67.

[19] Lobner-Olesen A, Hansen FG, Rasmussen KV, Martin B, Kuempel PL. The initiation cascade for chromosome replication in wild-type and Dam methyltransferase deficient *Escherichia coli* cells. EMBO J 1994;13:1856−62.

[20] Lahue RS, Su SS, Modrich P. Requirement for d(GATC) sequences in *Escherichia coli* mutHLS mismatch correction. Proc Natl Acad Sci U S A 1987;84:1482−6.

[21] Blow MJ, Clark TA, Daum CG, Deutschbauer AM, Fomenkov A, Fries R, et al. The Epigenomic Landscape of Prokaryotes. PLoS Genet 2016;12:e1005854.

[22] Lee CR, Lee JH, Park KS, Kim YB, Jeong BC, Lee SH. Global dissemination of

carbapenemase-producing *Klebsiella pneumoniae*: epidemiology, genetic context, treatment options, and detection methods. Front Microbiol 2016;7:895.

[23] Darling AE, Mau B, Perna NT. progressiveMauve: multiple genome alignment with gene gain, loss and rearrangement. PLoS One 2010;5:e11147.

[24] Roberts RJ, Vincze T, Posfai J, Macelis D. REBASE--a database for DNA restriction and modification: enzymes, genes and genomes. Nucleic Acids Res 2015;43:D298−9.

[25] de Jong A, Pietersma H, Cordes M, Kuipers OP, Kok J. PePPER: a webserver for prediction of prokaryote promoter elements and regulons. BMC Genomics 2012;13:299.

[26] Luo H, Gao F, Lin Y. Evolutionary conservation analysis between the essential and nonessential genes in bacterial genomes. Sci Rep 2015;5:13210.

[27] Waldron DE, Owen P, Dorman CJ. Competitive interaction of the OxyR DNA-binding protein and the Dam methylase at the antigen 43 gene regulatory region in *Escherichia coli*. Molecular Microbiology 2002;44:509−20.

[28] Campbell JL, Kleckner N. *E. coli* oriC and the dnaA gene promoter are sequestered from *dam* methyltransferase following the passage of the chromosomal replication fork. Cell 1990;62:967−79.

[29] Chang SW, Chao WR, Ruan A, Wang PH, Lin JC, Han CP. A promising hypothesis of c-KIT methylation/ expression paradox in c-KIT (+) squamous cell carcinoma of uterine cervix-CTCF transcriptional repressor regulates c-KIT proto-oncogene expression. Diagn Pathol 2015;10:207.

[30] Nievera C, Torgue JJ, Grimwade JE, Leonard AC. SeqA blocking of DnaA-oriC interactions ensures staged assembly of the *E. coli* pre-RC. Mol Cell 2006;24:581−92.

[31] Coffin SR, Reich NO. Modulation of Escherichia coli DNA methyltransferase activity by biologically derived GATC-flanking sequences. J Biol Chem 2008;283:20106−16.

[32] Pollak AJ, Chin AT, Brown FL, Reich NO. DNA looping provides for "intersegmental hopping" by proteins: a mechanism for long-range site localization. J Mol Biol 2014;426:3539−52.

[33] Barel I, Naughton B, Reich NO, Brown FLH. Specificity versus processivity in the sequential modification of DNA: a study of DNA adenine methyltransferase.

J Phys Chem B 2018;122:1112–20.

[34] Zheng Y, Sweet SM, Popovic R, Martinez-Garcia E, Tipton JD, Thomas PM, et
al. Total kinetic analysis reveals how combinatorial methylation patterns are
established on lysines 27 and 36 of histone H3. Proc Natl Acad Sci U S A
2012;109:13549–54.

[35] Wodarz D, Boland CR, Goel A, Komarova NL. Methylation kinetics and CpG-
island methylator phenotype status in colorectal cancer cell lines. Biol Direct
2013;8:14.

[36] Hojfeldt JW, Laugesen A, Willumsen BM, Damhofer H, Hedehus L, Tvardovskiy
A, et al. Accurate H3K27 methylation can be established de novo by SUZ12-
directed PRC2. Nat Struct Mol Biol 2018;25:225–32.

[37] Siu LK, Yeh K-M, Lin J-C, Fung C-P, Chang F-Y. *Klebsiella pneumoniae* liver
abscess: a new invasive syndrome. The Lancet Infectious Diseases
2012;12:881–7.

[38] English AC, Richards S, Han Y, Wang M, Vee V, Qu J, et al. Mind the gap:
upgrading genomes with Pacific Biosciences RS long-read sequencing
technology. PLoS One 2012;7:e47768.

[39] Brettin T, Davis JJ, Disz T, Edwards RA, Gerdes S, Olsen GJ, et al. RASTtk: a
modular and extensible implementation of the RAST algorithm for building
custom annotation pipelines and annotating batches of genomes. Sci Rep
2015;5:8365.

[40] Loytynoja A, Goldman N. webPRANK: a phylogeny-aware multiple sequence
aligner with interactive alignment browser. BMC Bioinformatics 2010;11:579.

[41] Page AJ, Cummins CA, Hunt M, Wong VK, Reuter S, Holden MT, et al. Roary:
rapid large-scale prokaryote pan genome analysis. Bioinformatics 2015;31:3691-
3.

[42] Price MN, Dehal PS, Arkin AP. FastTree 2--approximately maximum-likelihood
trees for large alignments. PLoS One 2010;5:e9490.

[43] He Z, Zhang H, Gao S, Lercher MJ, Chen WH, Hu S. Evolview v2: an online
visualization and management tool for customized and annotated phylogenetic
trees. Nucleic Acids Res 2016;44:W236–41.

[44] Ma W, Noble WS, Bailey TL. Motif-based analysis of large nucleotide data sets
using MEME-ChIP. Nat Protoc 2014;9:1428–50.

[45] Bolger AM, Lohse M, Usadel B. Trimmomatic: a flexible trimmer for Illumina
sequence data. Bioinformatics 2014;30:2114–20.

[46] Krueger F, Andrews SR. Bismark: a flexible aligner and methylation caller for
Bisulfite-Seq applications. Bioinformatics 2011;27:1571–2.

[47] Katoh K, Standley DM. MAFFT multiple sequence alignment software version 7:
improvements in performance and usability. Mol Biol Evol 2013;30:772–80.

[48] Zhang Z, Xiao J, Wu J, Zhang H, Liu G, Wang X, et al. ParaAT: a parallel tool
for constructing multiple protein-coding DNA alignments. Biochem Biophys Res
Commun 2012;419:779–81.

[49] Cooper GM, Stone EA, Asimenos G, Program NCS, Green ED, Batzoglou S, et
al. Distribution and intensity of constraint in mammalian genomic sequence.
Genome Res 2005;15:901–13.

 Figure legends

 Figure 1 Circos plots displaying the general genomic information of 14 *K.*

 ***pneumoniae* strains (NTUH-K2044; 11492; 11420; 11454; 11311; 12208; 11311; 23;**

 11021; N201205880; 309074; 11305; 13190; 283747 and 721005)

 The circles are as follows (from outside to inside): (1) A physical map scaled in

 megabases (Mb) from the base 1, which is the start of putative replication origin; (2)

 Coding sequences transcribed in a clockwise direction; (3) Coding sequences

 transcribed in a counterclockwise direction; (4) G+C content in a window of 2 kb

 sliding (red and green indicates G+C content that is higher and lower than average,

 respectively); (5) GC (G-C/G+C) skew in a window of 2 kb sliding (orange and green

 indicates GC skew above and below zero, respectively); (6) GA<u>T</u>C motifs; (7)

 CCW<u>G</u>G motifs. The other circles inside indicate the other 13 methylation motifs.

 Figure 2 The 15 MTase genes and corresponding methylated sequence motifs

 for the 14 *K. pneumoniae* strains

 15 MTase genes and corresponding methylated sequence motifs in 14 *K. pneumoniae*

 strains. Hyper-virulence strains and low-virulence strains were separately labeled in

 red and green letters. The large circles represents chromosome genome , and small

 circles around represents plasmid genome. Only the MTases with enzymatic activity

 are shown in this figure.

 Figure 3 Density distribution of the GATC/CCWGG motifs on the NTUH-

 K2044 genome and random generated genome

 The orange histograms show the density distribution of GA<u>T</u>C/CCW<u>G</u>G motifs on

 the random generated genome, which follow Poisson distribution with $\lambda = 3.73/2.89$

 (total number of GA<u>T</u>C/CCWGG motifs$\times$1000 per genome size). The green

 histograms show the density distribution of GA<u>T</u>C/CCW<u>G</u>G on the NTUH-K2044

 genome, which also follow Poisson distributions with $\lambda = 5.64/3.62$.

 Figure 4 Distribution of GATC/CCWGG motifs with different methylation

 patterns in GR and IGR regions

 A. Bar plots showing the ratio of GA<u>T</u>C/CCW<u>G</u>G motifs with different methylation

 patterns in GRs and IGRs of 14 *K. pneumonia* strains. The blue, yellow and green

 bars indicate the ratios of methylated, hemi-methylated and un-methylated motifs in

 GRs and IGRss. **B**. Box plots showing the methylated levels of GA<u>T</u>C/CCW<u>G</u>G

 motifs in GRs and IGRs (red and blue boxes). **C**. Frequency distribution of the

methylated (top) and hemi/un-methylated (bottom) GA**T**C/C**C**W**G**G motifs in GRs

and IGRs of 14 *K. pneumonia* strains. The dotted grey lines represent the position of

the start and stop codons.

Figure 5   Dynamic methylation analysis of GA<u>T</u>C and CCW<u>G</u>G motifs of NTUH-K2044 and 11492

A. Growth curves of *K. pneumonia* NTUH-K2044 (red) and 11492 (blue). X-axis represents the growth time, including lag phase, exponential phase, transition-to-stationary phase and stationary phase. Y-axis represents the logarithms of normalized O.D. value. At 1, 4 and 24 h, aliquots of bacterial cultures were collected and sequenced using Pacbio and Illumina platforms. The doubling time (tD) of the two strains are also labelled in the plot. **B**. Genome-wide methylation level versus genome position for the two motifs (GA**T**C and C**C**W**G**G) at exponential phase. The bold fitting lines approximate the average methylation levels across the genomes (5 kb window size). **C**. Schematic diagram showing the process of DNA replication in exponential phase. (1) The origin replication complex binds to oriC region. (2) The first round of replication will be initiated when original replication complex binds to oriC region. (3) The second round of replication begins before completion of the first round replication. (4) Six replication forks are generated in one bacterium. (5-6) The first round of replication completes, followed by a new cell replication cycle. **D**. Fitting of the genomic coverages with the mathematical model of replication. The genomic coverage plots of 11492 (upper) and NTUH-K2044 (lower) strains are shown in grey; the mathematically fitting curves are shown in red. The equation for fitting plot is also shown in the figure.

Figure 6   Comparison of the methylation time of GACT motifs in different genome regions

A. Box plot shows the methylation time of GA**T**C motifs in *oriC* and other regions. **B**. Box plot shows the methylation time of GA**T**C and C**C**W**G**G motifs in GR and IGR regions. ** $P < 0.01$, *** $P < 0.001$.

Tables

Table 1   General genomic information of the 14 *K. pneumoniae* strains

Table 2   Three types of modification patterns of GA<u>T</u>C motif from the 14 *K.*

***pneumoniae* strains**

Table 3   Modification patterns of the 12 motifs with 6mA among the 14 *K.*

***pneumoniae* strains**

Table 4   Modification patterns of the two motifs with 5mC among the 14 *K.*

***pneumoniae* strains**

Table 5   15 methylated motifs and corresponding DNA MTases from the 14 *K.*

***pneumoniae* strains**

Table 6   Average methylation time per motif in the two *K. pneumoniae* strains

(11492 and NTUH-K2044)

plus-strand genes
minus-strand genes
High GC content
Low GC content
(G-C)/(G+C)>0
(G-C)/(G+C)<0
GATC
CCWGG
MTCGAK
AGGAAG
GRACRAC
CTAN5GTAA
TTCAN7TTC
RTACN5GGC
CATCN6TTYG
CCAN7TCAC
CCAYN7TTYG
AGCN5CTTC
CAGN6TCAA
GGCAN8TCG
CCAGN7RTTC

Table 1 General genomic information of the 14 *K. pneumoniae* strains

Strains No.	Chromosome/ plasmid	Genomic size (Mb)	GC content (%)	Gene number	Average gene size (bp)	Coding region (%)	tRNAs	rRNAs
NTUH-K2044	NUTH-K2044_chr	5.25	57.68	5050	917	90.09	86	25
	pNUTH-K2044-1	0.22	50.17	243	701	83.28	0	0
11492	11492_chr	5.25	57.45	5017	922	88.43	87	25
	p11492-1	0.19	50.37	202	724	82.72	0	0
11420	11420_chr	5.44	57.46	5264	912	88.35	88	25
	p11420-1	0.23	50.10	257	688	83.43	0	0
	p11420-2	0.08	50.56	101	605	84.45	0	0
11454	11454_chr	5.20	57.60	4990	917	88.35	86	25
	p11454-1	0.21	50.67	225	717	83.74	0	0
12208	12208_chr	5.21	57.59	5067	910	89.42	82	25
	p12208-1	0.33	47.27	353	709	83.79	0	0
	p12208-2	0.21	49.69	232	698	82.90	0	0
11311	11311_chr	5.27	57.56	5048	918	88.26	86	25
	p11311-1	0.20	49.93	209	733	82.74	0	0
23	23_chr	5.29	57.54	5120	910	88.31	85	25
	p23-1	0.20	49.93	212	730	82.76	0	0
	p23-2	0.11	49.46	106	831	87.78	2	0

Strains No.	Chromosome/ plasmid	Genomic size (Mb)	GC content (%)	Gene number	Average gene size (bp)	Coding region (%)	tRNAs	rRNAs
11305	11305_chr	5.32	57.46	5186	901	88.37	88	25
	p11305-1	0.06	53.29	60	690	75.98	0	0
N201205880	N201205880_chr	5.21	57.50	5106	901	88.85	87	25
	pN201205880-1	0.23	51.08	240	758	82.03	0	0
	pN201205880-2	0.15	52.33	146	755	88.63	0	0
	pN201205880-3	0.11	54.85	134	661	83.30	0	0
	pN201205880-4	0.11	49.31	108	815	88.07	1	0
	pN201205880-5	0.08	50.29	88	733	84.69	0	0
	pN201205880-6	0.08	54.00	96	596	78.08	0	0
	pN201205880-7	0.07	52.85	83	657	86.33	0	0
	pN201205880-8	0.06	52.45	69	609	82.19	0	0
309074	309074_chr	5.31	57.46	5181	907	90.23	89	25
	p309074-1	0.19	53.44	197	735	83.60	0	0
	p309074-2	0.07	51.83	85	668	83.78	0	0
	p309074-3	0.05	49.03	65	667	85.67	0	0
13190	13190_chr	5.26	57.50	5172	897	89.34	88	25
	p13190-1	0.31	47.26	316	703	84.09	0	0
	p13190-2	0.08	53.86	85	649	79.42	0	0

Strains No.	Chromosome/plasmid	Genomic size (Mb)	GC content (%)	Gene number	Average gene size (bp)	Coding region (%)	tRNAs	rRNAs
	p13190-3	0.04	45.72	50	674	86.66	0	0
283747	283747_chr	5.50	57.37	5441	891	90.07	85	25
	p283747-1	0.15	54.40	192	574	78.82	0	0
721005	721005_chr	5.48	57.38	5468	886	89.57	85	25
	p721005-1	0.27	46.48	283	716	82.85	0	0
	p721005-2	0.16	53.92	199	627	82.14	0	0
	p721005-3	0.06	51.76	77	631	88.59	0	0
11021	11021_chr	5.45	57.46	5386	893	89.67	85	25
	p11021-1	0.24	51.69	271	682	82.43	0	0
	p11021-2	0.17	53.48	217	605	81.85	0	0

Table 2 Three types of modification patterns of GATC motif from the 14 *K. pneumoniae* strains

Strain No.	Motif [a]	Total motif number [b]	Methylated motif (%)	Hemi-methylated motif (%)	Un-methylated motif (%)
NTUH-K2044	GATC	30,727	30,151 (98.13%)	551 (1.79%)	25 (0.08%)
11492	GATC	30,316	30,271 (99.85%)	31 (0.10%)	14 (0.05%)
11420	GATC	31,847	26,835 (84.26%)	4669 (14.66%)	343 (1.08%)
11454	GATC	30,278	28,046 (92.63%)	2142 (7.07%)	90 (0.30%)
12208	GATC	31,512	30,695 (97.41%)	790 (2.51%)	27 (0.09%)
11311	GATC	30,623	29,520 (96.4%)	1076 (3.51%)	27 (0.09%)
23	GATC	31,275	26,523 (84.81%)	4423 (14.14%)	329 (1.05%)
11305	GATC	30,369	29,222 (96.22%)	1122 (3.69%)	25 (0.08%)
N201205880	GATC	33,171	28,007 (84.43%)	4852 (14.63%)	312 (0.94%)
309074	GATC	31,208	30,384 (97.36%)	782 (2.51%)	42 (0.13%)
13190	GATC	31,419	28,766 (91.56%)	2515 (8.00%)	138 (0.44%)
283747	GATC	31,335	30,156 (96.24%)	1126 (3.59%)	53 (0.17%)
721005	GATC	32,242	31,521 (97.76%)	687 (2.13%)	34 (0.11%)
11021	GATC	32,080	27,907 (86.99%)	3968 (12.37%)	205 (0.64%)

Note: [a] The methylated nucleotide in the motif is shown as bold letter. The underlined letter represents the thymine pairing with the methylated adenine on the complementary strand. [b] The motif number includes the ones on the plus and minus strands of chromosomes and plasmids.

Table 3 Modification patterns of the 12 motifs with 6mA among the 14 *K. pneumoniae* strains

Strain No.	Motif [a]	Total motif number [b]	Methylated motif (%)	Hemi-methylated motif (%)	Un-methylated motif (%)
NTUH-K2044	GR**A**CRAC*	2104	2060 (97.91%)	/	44 (2.09%)
11420	RT**A**CN$_5$GGC*	1275	1156 (90.67%)	/	119 (9.33%)
	TTCAN$_7$TTC*	853	598 (70.11%)	231 (27.08%)	24 (2.81%)
11454	**A**GCN$_5$CTTC	991	930 (93.84%)	61 (6.16%)	/
12208	AGG**A**AG*	2864	2848 (99.30%)	/	16 (0.7%)
11311#	CC**A**YN$_7$TTYG*	667	608 (91.15%)	58 (8.70%)	1 (0.15%)
23#	CC**A**YN$_7$TTYG*	687	534 (77.73%)	145 (21.11%)	8 (1.16%)
11305^	CC**A**GN$_7$RTTC	342	325 (95.03%)	17 (4.97%)	/
N201205880	CC**A**N$_7$TCAC*	543	433 (79.74%)	102 (18.78%)	8 (1.47%)
309074	CT**A**N$_5$GTAA	165	164 (99.39%)	1 (0.61%)	/
	C**A**GN$_6$TCAA*	439	413 (94.08%)	26 (5.92%)	/
	C**A**TCN$_6$TTYG	634	589 (92.90%)	45 (7.10%)	/
13190^	CC**A**GN$_7$RTTC	379	338 (89.18%)	40 (10.55%)	1 (0.26%)
	GGC**A**N$_8$TCG	1065	874 (82.07%)	184 (17.28%)	7 (0.66%)

Note: [a] The methylated nucleotide in the motif is shown as bold letter. The underlined letter represents the thymine pairing with the methylated adenine on the complementary strand. Degenerate bases used in our recognition sequences are listed in the following: R = G or A, Y = C or T, M = A or C, K = G or T, S = G or C, W = A or T, B = not A (C or G or T), D = not C (A or G or T), H = not G (A or C or T), V = not T (A or C or G), N = A or C or G or T. [b] The motif number includes the ones on the plus and minus strands of

chromosomes and plasimds. [*] Newly reported motifs. [#] Both strain 11311 and 23 possess motif CC**A**YN7<u>T</u>TYG. ^ Both strain 11305 and 13190 possess motif CC**A**GN7R<u>T</u>TC.

Table 4 Modification patterns of the two motifs with 5mC among the 14 *K. pneumoniae* strains

Strain No.	Motif [a]	Total motif number [b]	Methylated motif (%)	Hemi-methylated motif (%)	Un-methylated motif (%)
NTUH-K2044	CCW<u>G</u>G	19,530	16,117 (82.52%)	1756 (8.99%)	1657 (8.49%)
	MT**C**<u>G</u>AK[*]	5410	2339 (43.23%)	270 (4.99%)	2801 (51.78%)
11492	CCW<u>G</u>G	19,284	19,015 (98.61%)	86 (0.45%)	183 (0.94%)
	MT**C**<u>G</u>AK[*]	5386	2368 (43.97%)	258 (4.79%)	2760 (51.24%)
11420	CCW<u>G</u>G	20,175	18,828 (93.32%)	839 (4.16%)	508 (2.52%)
11454	CCW<u>G</u>G	19,314	14,983 (77.58%)	2367 (12.26%)	1964 (10.17%)
12208	CCW<u>G</u>G	20108	19,874 (98.84%)	35(0.17%)	199 (0.99%)
11311	CCW<u>G</u>G	19,668	19,126 (97.24%)	299 (1.52%)	243 (1.24%)
23	CCW<u>G</u>G	19,941	18,668 (93.62%)	963 (0.48%)	310 (0.16%)
11305	CCW<u>G</u>G	19,460	18,964 (97.45%)	261 (1.34%)	235 (1.21%)
N201205880	CCWGG	21,115	20,867 (98.83%)	38 (0.18%)	210 (0.99%)
309074	CCWGG	19,966	17,464 (87.47%)	1460 (7.31%)	1042 (5.22%)
13190	CCWGG	20,060	19,149 (95.46%)	694 (3.46%)	217 (1.08%)
283747	CCWGG	20,156	19,141 (94.96%)	574 (2.85%)	441 (2.19%)
721005	CCWGG	20,832	20,470 (98.26%)	145 (0.70%)	217 (1.04%)
11021	CCWGG	20,736	18,352 (88.50%)	1358 (6.55%)	1026 (4.95%)

Note: [a] The methylated nucleotide in the motif is shown as bold letter. The underlined letter represents the guanine pairing with the methylated cytosine on the complementary strand. Degenerate bases used in our recognition sequences are listed in the following: R =

G or A, Y = C or T, M = A or C, K = G or T, S = G or C, W = A or T, B = not A (C or G or T), D = not C (A or G or T), H = not G (A or C or T), V = not T (A or C or G), N = A or C or G or T. [b] The motif number includes the ones on the plus and minus strands of chromosomes and plasimds. [*] Newly reported motifs.

Table 5 15 methylated motifs and corresponding DNA MTases from the 14 *K. pneumoniae* strains

Motif [a]	Modification	R-M/orphan[b]	MTase	Comments
G**A**TC	6mA	orphan	Dam	Known
RT**A**CN$_5$GGC	6mA	RM-I	KamA	New
TTC**A**N$_7$TTC	6mA	RM-I	KamB	New
GR**A**CRAC	6mA	RM-II	KamC	New
AGCN$_5$CTTC	6mA	RM-I	M.KpnGH01II	Known
AGGA**A**G	6mA	RM-II	KamD	New
CC**A**YN$_7$TTYG	6mA	RM-I	KamE	New
CC**A**N$_7$TCAC	6mA	RM-I	KamF	New
CC**A**GN$_7$RTTC	6mA	RM-I	M.KpnAATI	Known
CT**A**N$_5$GTAA	6mA	RM-I	M.Kpn35657I	Known
C**A**GN$_6$TCAA	6mA	RM-I	KamG	New
C**A**TCN$_6$TTYG	6mA	RM-I	M.Kpn39795II	Known
GGC**A**N$_8$TCG	6mA	RM-I	M.KpnAATIV	Known
C**C**WGG	5mC	orphan	Dcm	Known
MT**CG**AK	5mC	RM-II	KcmA	New

Note: [a] The methylated nucleotide in the motif is shown as bold letter. The underlined letter represents the guanine pairing with the methylated cytosine on the complementary strand. Degenerate bases used in the recognition sequences are listed in the following: R = G or A, Y = C or T, M = A or C, K = G or T, S = G or C, W = A or T, B = not A (C or G or T), D = not C (A or G or T), H = not G (A or C or T), V = not T (A or C or G), N = A or C or G or T. [b] The MTase prediction was based on the sequence alignment with REBASE database (http://rebase.neb.com/rebase/rebase.html). The predicted MTases were further classified as Type I, Type II or orphan MTases according to the annotation information.

Table 6 Average methylation time per motif in the two *K. pneumoniae* strains (11492 and NTUH-K2044)

Motif sequence	No. of motif (%)	Methylation time (mean ± sd, min)
GA<u>T</u>C	59424 (100%)	3.5 ± 0.5
SGA<u>T</u>CS (fast)	17041 (28.7%)	3.4 ± 0.5**
WGA<u>T</u>CW (slow)	9803 (16.5%)	3.8 ± 0.5***
CCW<u>G</u>G	35516 (100%)	9.2 ± 4.9
SCCW<u>G</u>GS (slow)	13573 (38.2%)	9.6 ± 4.9***
WCCW<u>G</u>GW (fast)	5800 (16.3%)	8.2 ± 4.5***
MTC<u>G</u>AK	10467 (100%)	4.5 ± 0.5
GR<u>A</u>CRAC	2002 (100%)	3.5 ± 0.4

Note: Degenerate bases used in our recognition sequences are listed in the following: R = G or A, Y = C or T, M = A or C, K = G or T, S = G or C, W = A or T, B = not A (C or G or T), D = not C (A or G or T), H = not G (A or C or T), V = not T (A or C or G), N = A or C or G or T. *** means that the methylation times of the motifs with certain flanking sequences show the significant difference ($P < 0.001$) comparing with the average methylation time of total motifs.