Prevalence of Hypervirulence-Associated Pathogenicity Loci Among *Klebsiella*

pneumoniae Bloodstream Isolates at a United States Hospital

Travis J. Kochan[1], Natalia Khalatyan[1], Bettina H. Cheung[1], Sophie H. Nozick[1],

Egon A. Ozer[2], Alan R. Hauser[1,2]

[1]Department of Microbiology-Immunology, Northwestern University, Feinberg

School of Medicine, Chicago, IL, USA

[2]Division of Infectious Diseases, Department of Medicine, Northwestern

University, Feinberg School of Medicine, Chicago, IL, USA

Abstract

Klebsiella pneumoniae is a major threat to human health worldwide. "Classical" *K.*

pneumoniae (cKp) strains commonly cause multidrug-resistant (MDR) infections in

debilitated patients residing in hospitals and long-term care facilities. In contrast,

hypervirulent *K. pneumoniae* (hvKp) strains may cause invasive, community-acquired

infections in otherwise healthy individuals. These highly virulent strains were first

identified in Taiwan in 1986 and have since disseminated across the globe. The

prevalence of hvKp has been well described in Southeast Asia, but is not well

understood in the United States. In this work, 141 consecutive *K. pneumoniae* blood

stream isolates were collected at Northwestern Memorial Hospital from 2015-2017. To

screen this collection for hvKp, isolates were tested for the hypermucoviscous

phenotype and the presence of aerobactin and salmochelin biosynthesis genes,

biomarkers that have been used to distinguish hvKp from cKp. These biomarkers were

identified in 12 (8.5%), 6 (4.3%), and 5 (3.5%) isolates, respectively. Whole genome

sequencing was performed on the six isolates that contained aerobactin biosynthesis

genes and demonstrated that these genes were contained on large plasmids. Five

isolates contained plasmids nearly identical to previously described hvKp virulence

plasmids while one was similar to a plasmid from a swine isolate of *K. pneumoniae*.

These findings indicate that several non-clonal hvKp-like strains are present in the U.S.

Introduction

Klebsiella pneumoniae (Kp) is a leading cause of nosocomial infections

worldwide including pneumonia, bloodstream, and urinary tract infections (1). Kp can

cause severe, secondary pneumonia in patients with respiratory viral infections and was

recently identified as a leading cause of secondary pneumonia in patients with the novel

coronavirus SARS-CoV-2 (2). Kp pathogenicity is due to a variety of virulence factors,

including siderophores, fimbriae, and a polysaccharide capsule of various types. The

polysaccharide capsule is the major virulence factor for Kp as it allows this bacterium to

evade host immune defenses, such as complement-mediated phagocytosis (3, 4). The

most common type of Kp are referred to as classical strains (cKp), which infect

debilitated patients residing in hospitals and long-term care facilities, (5). cKp strains are

increasingly resistant to antibiotics, including carbapenems (6-11). Genes conferring

antimicrobial resistance are often carried by conjugative plasmids allowing for extensive

dissemination among cKp strains. Thus, it is not surprising that the IDSA, WHO, and the

CDC have each deemed Kp as a serious public health priority in need of new

therapeutic development (12-14). As such, multidrug-resistant (MDR) cKp strains have

been the focus of research within the United States.

A second type of Kp, referred to as hypervirulent Kp (hvKp), was first identified in

Taiwan in 1986 as a common cause of pyogenic liver abscesses (PLA) in young,

otherwise healthy individuals living in the community (15-22). These strains are now

common in Asia, but little is known about their prevalence in the U.S. Case reports

document the presence of hvKp infections and PLA in North America, but only a few

surveillance studies have been performed (21, 24-26). hvKp strains frequently express a

highly mucoid capsule, leading to a "hypermucoviscous" colony phenotype, and are

considerably more virulent in mouse models than cKp strains (23). Their increased

virulence is usually attributed to the presence of one of a number of large, non-

conjugative, "hypervirulence" plasmids (27-29). The limitations imposed by the hyper-

mucoid capsule on genetic exchange and the non-conjugative nature of these virulence

plasmids may restrict the hypervirulence phenotype to few sequence types (e.g. ST23,

ST86, ST66) and capsule types (e.g. K1, K2). The best characterized hypervirulence

plasmids are pK2044 and KP52.145pII (Fig 1A, B) (29, 30). In addition, we previously

identified a unique plasmid with hvKp features, pTK421_2 (Fig 1C), in a bloodstream

isolate from Northwestern Memorial Hospital (NMH), suggesting considerable diversity

among the hypervirulence plasmids of hvKp (31). Each hypervirulence plasmid contains

two distinct pathogenicity loci (PAL-1 and PAL-2) (Fig 1D). PAL-1 usually consists of a

mucoid regulator gene (*rmpA2*), the aerobactin receptor gene (*iutA*), and aerobactin

biosynthesis genes (*iucABCD*). PAL-2 usually consists of a distinct mucoid regulator

operon (*rmpADC*), the salmochelin receptor gene (*iroN*), and salmochelin biosynthesis

genes (*iroBD*) (Fig 1D). Plasmids KP52.145pII and pTK421_2 contain an incomplete

PAL-1 that lacks *rmpA2* (Fig 1D). The mucoid regulators *rmpA* and *rmpA2* play a major

role in virulence by increasing capsule production and hypermucoviscosity (28), but the

roles of the different siderophores (aerobactin, salmochelin, and yersiniabactin) in hvKp

virulence remain controversial (30, 32). A recent report suggested that aerobactin was

the only siderophore required for the hypervirulent phenotype of the ST86-K2 capsule

type strain hvKp1 (32). In contrast, another report showed that only a triple siderophore

mutant (yersiniabactin, aerobactin, and salmochelin) of the ST23-K1 strain NTUH-K2044

was attenuated in the mouse intraperitoneal model (30). In addition, strain ATCC43816

(also called KPPR1) is hypervirulent in mice (4, 33) but is otherwise atypical in that it

lacks a hypervirulence plasmid and PAL-1 but contains a version of PAL-2 integrated

into a chromosomal integrative and conjugative element (ICEKp1, Fig 1D). In KPPR1,

yersiniabactin promoted disease in the lung and salmochelin promoted dissemination to

tissues (34). Thus, it is clear that the mucoid regulators and siderophores play roles in

conferring the hypervirulent phenotype, but significant strain-to-strain variability of

virulence factors exists.

A microbiological definition of hvKp has remained elusive. Initially,

hypermucoviscosity was used as a proxy for the hvKp phenotype. However, several

reports have described hypermucoviscous Kp strains that are not hypervirulent in mice

and lack pathogenicity loci usually associated with hvKp (35-37). As a result, genetic

biomarkers have been used to define hvKp, but these definitions have varied from study-

to-study. Recently, Russo and colleagues systematically examined the accuracy of using

rmpA, *rmpA2*, *peg-344*, *iucA*, and *iroB,* genes found on virulence plasmids, as diagnostic

biomarkers for hvKp (23, 38). In their cohort of 175 isolates, the presence of *iucA* or *iroB*

or a hypermucoviscosity phenotype distinguished hvKp from cKp with an accuracy of

96%, 97%, and 90%, respectively. Here, we use these biomarkers to describe the

prevalence of hvKp at NMH in Chicago, Illinois. From 2015 – 2017, we collected 141

consecutive Kp bloodstream isolates and tested them for these biomarkers by

polymerase chain reaction and the string test. We identified twelve hypermucoviscous

isolates and six isolates that contain hypervirulence pathogenicity loci located on large

hvKp-like plasmids.

Results

Identification of hvKp using diagnostic biomarkers

To assess the prevalence of hvKp-like strains at NMH, we collected 141

consecutive Kp isolates from patients with positive blood cultures from 2015–2017.

Given that hvKp strains are thought to disseminate through the bloodstream, we

hypothesized that bloodstream isolates would be enriched for hvKp. We screened all

141 isolates for the following previously published diagnostic biomarkers of hvKp: *iucA,*

iroB, and the hypermucoviscous phenotype. The presence of *iucA* and *iroB* was

evaluated by PCR. Among NMH bloodstream isolates, six (4.3%) were positive for *iucA*

and five (3.5%) were positive for *iroB* (Table 1). Five isolates were positive for both, and

a single isolate was *iucA+* and *iroB-* (Table 1). The collection was screened for

hypermucoviscosity by both string test and centrifugation. Twelve (6.7%) and 24 (13.5%)

isolates, respectively, were positive for hypermucoviscosity by these two tests (Table 1).

In addition, isolates were tested for tellurite resistance, a phenotype commonly encoded

by pK2044 plasmids (Fig 1). Six isolates (4.3%) were able to grow on tellurite, including

three isolates that were positive for *iroB* and *iucA* (Table 1). This suggested that these

three isolates may contain pK2044-like plasmids while the other double-positive (*iroB+,*

iucA+) isolates may contain KP52.145pII-like plasmids or unique hypervirulence

plasmids. Collectively, these findings indicate that Kp strains with features of hvKp are

circulating at NMH and suggest that these strains may contain multiples types of hvKp-

associated pathogenicity loci.

Analysis of hvKp Plasmid Content

Since hvKp pathogenicity loci are commonly encoded on a large virulence

plasmid, we next focused on the six isolates that contained either salmochelin or

aerobactin biosynthesis genes. To determine whether hvKp-associated genes of these

six isolates were carried on large plasmids similar to those found in characterized hvKp

isolates, complete genome sequences were obtained using both Nanopore and Illumina

platforms (Table 2). Whole-genome sequencing revealed that each of these isolates

harbored large plasmids with some hvKp pathogenicity loci. Three isolates (KpN8,

KpN89, and KpN115) contained plasmids with near complete alignment with pK2044,

including PAL-1, PAL-2, and tellurite resistance genes (Fig 2A). KpN49 and KpN165

harbored plasmids with poor overall alignment with plasmid pK2044, (Fig 2B) but with

nearly complete alignment with KP52.145pII (Fig 3). Similar to KP52.145pII, these two

plasmids contained a version of PAL-1 with *iucABCD* and *iutA* but lacking *rmpA2* (Fig 1).

The remaining isolate, KpN23, harbored a conjugative virulence plasmid containing only

the aerobactin biosynthesis genes and receptor (Fig 3, 4A and 4B). pKpN23_1 was

highly homologous to a plasmid harbored by a swine isolate collected in Thailand in

2016, KpCTRSRTH01_p2, but had minimal alignment with pK2044 or KP52.145pII. We

had previously identified pTK421_2, a Kp plasmid with PAL-1, PAL-2, and conjugation

genes (Fig 1). This plasmid was also distinct from pK2044 or KP52.145pII (Fig 2B & 3).

We therefore aligned pKPN23_1 to pTK421_2 but found that these plasmids shared

relatively little alignment beyond PAL-1 (Fig 4B). Overall, among the 141 bloodstream

isolates, we identified six isolates that cumulatively contained three distinct plasmids with

pathogenicity loci associated with hypervirulent Kp.

Discussion

Case reports indicate that hvKp strains exist in the U.S., but their prevalence is

unclear and has only been investigated in two surveillance studies (24, 25). In this work,

we determined the prevalence of hvKp at NMH by identifying isolates using diagnostic

criteria of hvKp. The three criteria we tested were *iroB*, *iucA,* and hypermucoviscosity.

We identified five isolates (3.5%) that met all three of these examined diagnostic criteria

for hvKp; each contained a large plasmid nearly identical to previously described hvKp

plasmids. We identified one additional isolate that contained *iucA* on a large plasmid but

was not hypermucoviscous and lacked *iroB*. Currently, it is unclear if isolates positive for

only a single biomarker are highly virulent and cause the types of infections

characteristic of hvKp. Further studies are warranted to understand the level of virulence

in strains such as this one with incomplete pathogenicity loci. In addition, we identified

12 isolates that were hypermucoviscous by string test and 24 that were

hypermucoviscous by centrifugation. It is unclear why some isolates were

hypermucoviscous by centrifugation but not the string test. These data agree with

previous reports suggesting that the string test is less sensitive than centrifugation for

measuring hypermucoviscosity and that hypermucoviscosity does not accurately predict

hypervirulence (37, 39). However, all strains that contained *rmpA* were positive by both

string and centrifugation tests (Table 2).

Since hvKp pathogenicity loci are normally encoded on a large plasmid, we

performed long-read sequencing of the six isolates that contained *iucA*. Whole genome

sequencing confirmed that each of these isolates harbored a large plasmid. Five of

these isolates harbored a plasmid highly similar to one of the two previously described

hvKp virulence plasmids, pK2044 or KP52.145pII, and the other, KpN23, harbored a

unique plasmid containing only aerobactin. Taken together, these data suggest that

multiple clones of hvKp isolates are circulating at NMH.

Our study agrees with the two published U.S. surveillance studies, which

describe the hvKp prevalence in the U.S. as 3.5%-6.25% of Kp infections. In 2010, 64

Kp strains were analyzed from two hospitals in the greater Houston area (25). They

identified four strains (6.25%) containing hvKp pathogenicity-associated genes, based

on the presence of *rmpA* and K1 capsule type by PCR amplification. In 2018, a larger

study in New York City characterized Kp isolates from 462 patients (24). This study

identified 17 (3.7%) isolates by PCR amplification that contained the hvKp biomarkers

rmpA and *iucA*. This is much lower than the prevalence of hvKp isolates in China in

2013 (37.8%, defined as *iuc+*), where hvKp is endemic (40). A recent genomic

surveillance study of seven countries in South and Southeast Asia analyzed the

sequences of 365 bloodstream isolates collected from 2010-2017 (41). Sequencing

revealed that 26% (95/365) were *iuc+*, 17% (63/365) were *iuc+* and either *rmpA+* or

rmpA2+, and an additional 0.8% (3/365) were *iro+* and *rmpA+*. Of the *iuc+* isolates,

13.7% (13/95) contained carbapenamase genes, but only 1 was positive for *iuc*, *rmpA/2*

and a carbapenamase gene. Our results together with these reports indicate that while

hvKp-like isolates are circulating within the United States, they remain relatively rare

compared to Southeastern Asia.

The presence of hvKp in U.S. hospitals has important clinical implications. In

patients, hvKp infections are associated with invasive infections most commonly

characterized by the development of PLA. Extrahepatic complications such as

endogenous endophthalmitis, necrotizing fasciitis, and meningitis have also been

reported (42). These invasive infections are associated with significant mortality and

morbidity; patients with PLA take longer to recover and 9-18% have recurrent PLA (43,

44). In addition, one study showed that 10/14 patients with endogenous endophthalmitis

progressed to blindness (45). Interestingly, while hvKp isolates cause severe invasive

infections, several studies reported no difference in mortality following bloodstream

infection with hvKp or cKp (46, 47). This is likely related to the fact that community

acquired hvKp infections tend to occur in young healthy patients, and outcome of

bloodstream infection is largely dependent on the time it takes to initiate appropriate

antimicrobial therapy. Delay in effective antimicrobial therapy, which is more likely to

occur in infections cause by multidrug-resistant bacteria, leads to poor outcomes in

bloodstream infection (48, 49). None of the six hvKp-like isolates identified at NMH

contained antimicrobial resistance genes. Thus far, antimicrobial resistant hvKp

infections are exceedingly rare in the United States. However, antimicrobial resistant

hvKp infections are on the rise in Asia, likely due to the increase in hospital acquired

hvKp infections (46, 50, 51). While this study is limited to a single institution, the findings

presented here describe the prevalence and diversity of hvKp plasmids at a major U.S.

academic medical center and call attention to the need for further studies evaluating the

risk and clinical outcomes of these diverse hvKp isolates.

Materials and Methods

Bacterial isolates and growth conditions

The isolates used in this study were collected from patients with a positive blood culture

from 2015-2017 and are described in Table S1. These isolates were identified as Kp on

a VITEK-2 (52).. Kp was grown at 37°C in Lysogeny broth (LB) or on LB agar plates. For

screening of tellurite resistance, Kp was grown at 37°C on LB agar plates supplemented

with 3 μg/mL potassium tellurite (Sigma-Aldrich, USA).

Screening for *iroB* and *iucA*

Aerobactin and salmochelin biosynthesis genes were detected by polymerase chain

reaction using custom designed primers:

iroB Fwd- ATTTCCGCCGCTACCTCTTCAG; iroB Rev-

TGGGCTATTGTATCCTGTGCTATCTCTG; iucA Fwd-

GTTGTGGTCGTTCAGATAACAGAAAAAGTT; iucA Rev-

AATTCTCATTATCATGACGAGTGAGATCTGT

Hypermucoviscosity testing

A string test was used to assess the hypermucoviscosity of each Kp isolate as described

previously (53). Isolates were grown overnight at 37°C on LB agar. A single colony was

lifted with a loop to evaluate the formation of a viscous string between the loop and the

colony. A positive string test was defined as a string length ≥ 5 mm. Hypermucoviscosity

was also assessed by centrifugation of overnight cultures (5 mL LB) at 3,220 rcf for 10

minutes (39). Hypermucoviscous isolates were identified by the persistence of turbidity.

Whole-Genome Sequencing

Genomic DNA (gDNA) was prepared from a single colony cultured overnight at 37°C in

LB using the Maxwell 16 system (Promega Corp., Madison, WI). Nanopore sequencing

was performed as described previously (54, 55). Briefly, libraries were prepared from

gDNA using the ligation sequencing kit (SQK-LSK109, Oxford Nanopore, UK). Libraries

were sequenced on the MinION using a FLO-MIN106 flow cell and base calling and

demultiplexing of sequence reads was performed using Guppy v3.4.5. Libraries for

Illumina sequencing were prepared using Nextera XT library kits (Illumina, Inc., San

Diego, CA) and sequenced on an Illumina NextSeq instrument to generate paired-end

150-bp reads. Hybrid assembly and circularization of Nanopore and Illumina reads were

performed using Flye v2.6. Nanopore sequencing errors were corrected by aligning

Illumina reads to the assembly using BWA v0.7.17 and correcting the errors with serial

rounds of Pilon v1.23. Annotation was performed using the NCBI Prokaryotic Genome

Annotation Pipeline.

Molecular typing and identification of virulence genes

Assembled whole genome sequences were analyzed with the bioinformatics tools

Kleborate and Kaptive to evaluate multi locus sequence type, capsule locus typing, and

virulence gene content (56, 57). Plasmids identified by whole genome sequencing were

aligned using BLAST Ring Image generator (BRIG) with the alignment threshold set at

85% identity (58).

Bibliography

1. Podschun R, Ullmann U. 1998. *Klebsiella spp.* as nosocomial pathogens: epidemiology, taxonomy, typing methods, and pathogenicity factors. Clin Microbiol Rev 11:589-603.

2. Dhesi Z, Enne VI, Brealey D, Livermore DM, High J, Russell C, Colles A, Kandil H, Mack D, Martin D, Page V, Parker R, Roulston K, Singh S, Wey E, Swart AM, Stirling S, Barber JA, O'Grady J, Gant VA. 2020. Organisms causing secondary pneumonias in COVID-19 patients at 5 UK ICUs as detected with the FilmArray test doi:doi: https://doi.org/10.1101/2020.06.22.20131573, medRvix.

3. Paczosa MK, Mecsas J. 2016. *Klebsiella pneumoniae*: Going on the Offense with a Strong Defense. Microbiol Mol Biol Rev 80:629-61.

4. Lawlor MS, Hsu J, Rick PD, Miller VL. 2005. Identification of *Klebsiella pneumoniae* virulence determinants using an intranasal infection model. Mol Microbiol 58:1054-73.

5. Russo TA, Marr CM. 2019. Hypervirulent *Klebsiella pneumoniae*. Clin Microbiol Rev 32.

6. Marcade G, Brisse S, Bialek S, Marcon E, Leflon-Guibout V, Passet V, Moreau R, Nicolas-Chanoine MH. 2013. The emergence of multidrug-resistant *Klebsiella pneumoniae* of international clones ST13, ST16, ST35, ST48 and ST101 in a teaching hospital in the Paris region. Epidemiol Infect 141:1705-12.

7. Meng X, Yang J, Duan J, Liu S, Huang X, Wen X, Huang X, Fu C, Li J, Dou Q, Liu Y, Wang J, Yan Q, Zou M, Liu W, Peng Z, Chen L, Li C, Wu A. 2019. Assessing Molecular Epidemiology of Carbapenem-resistant *Klebsiella pneumoniae* (CR-Kp) with MLST and MALDI-TOF in Central China. Sci Rep 9:2271.

8. Roulston KJ, Bharucha T, Turton JF, Hopkins KL, Mack DJF. 2018. A case of NDM-carbapenemase-producing hypervirulent *Klebsiella pneumoniae* sequence type 23 from the UK. JMM Case Rep 5:e005130.

9. Shen D, Ma G, Li C, Jia X, Qin C, Yang T, Wang L, Jiang X, Ding N, Zhang X, Yue L, Yin Z, Zeng L, Zhao Y, Zhou D, Chen F. 2019. Emergence of a Multidrug-Resistant Hypervirulent *Klebsiella pneumoniae* Sequence Type 23 Strain with a Rare bla CTX-M-24-Harboring Virulence Plasmid. Antimicrob Agents Chemother 63.

10. Shu L, Dong N, Lu J, Zheng Z, Hu J, Zeng W, Sun Q, Chan EW, Zhou H, Hu F, Chen S, Zhang R. 2019. Emergence of OXA-232 Carbapenemase-Producing *Klebsiella pneumoniae* That Carries a pLVPK-Like Virulence Plasmid among Elderly Patients in China. Antimicrob Agents Chemother 63.

11. Lester CH, Olsen SS, Jakobsen L, Arpi M, Fuursted K, Hansen DS, Heltberg O, Holm A, Hojbjerg T, Jensen KT, Johansen HK, Justesen US, Kemp M, Knudsen JD, Roder B, Frimodt-Moller N, Hammerum AM. 2011. Emergence of extended-spectrum beta-lactamase (ESBL)-producing *Klebsiella pneumoniae* in Danish hospitals; this is in part explained by spread of two CTX-M-15 clones with multilocus sequence types 15 and 16 in Zealand. Int J Antimicrob Agents 38:180-2.

298   12.   Tacconelli E, Magrini N. 2017. Global priority list of antibiotic-resistant
299   bacteria to guide research, discovery, and development of new antibiotics.
300   https://www.who.int/medicines/publications/WHO-PPL-
301   Short_Summary_25Feb-ET_NM_WHO.pdf. Accessed October 2.
302   13.   Talbot GH, Bradley J, Edwards JE, Jr., Gilbert D, Scheld M, Bartlett JG. 2006.
303   Bad bugs need drugs: an update on the development pipeline from the
304   Antimicrobial Availability Task Force of the Infectious Diseases Society of
305   America. Clin Infect Dis 42:657-668.
306   14.   CDC. 2013. Antibiotic resistance threats in the United States, 2013.
307   http://www.cdc.gov/drugresistance/threat-report-2013/. Accessed
308   September 18.
309   15.   Liu YC, Cheng DL, Lin CL. 1986. *Klebsiella pneumoniae* liver abscess
310   associated with septic endophthalmitis. Arch Intern Med 146:1913-6.
311   16.   Fang FC, Sandler N, Libby SJ. 2005. Liver abscess caused by magA+ *Klebsiella*
312   *pneumoniae* in North America. J Clin Microbiol 43:991-2.
313   17.   Lam MMC, Wyres KL, Duchene S, Wick RR, Judd LM, Gan YH, Hoh CH,
314   Archuleta S, Molton JS, Kalimuddin S, Koh TH, Passet V, Brisse S, Holt KE.
315   2018. Population genomics of hypervirulent *Klebsiella pneumoniae* clonal-
316   group 23 reveals early emergence and rapid global dissemination. Nat
317   Commun 9:2703.
318   18.   Turton JF, Englender H, Gabriel SN, Turton SE, Kaufmann ME, Pitt TL. 2007.
319   Genetically similar isolates of *Klebsiella pneumoniae* serotype K1 causing
320   liver abscesses in three continents. J Med Microbiol 56:593-7.
321   19.   Siu LK, Fung CP, Chang FY, Lee N, Yeh KM, Koh TH, Ip M. 2011. Molecular
322   typing and virulence analysis of serotype K1 *Klebsiella pneumoniae* strains
323   isolated from liver abscess patients and stool samples from noninfectious
324   subjects in Hong Kong, Singapore, and Taiwan. J Clin Microbiol 49:3761-5.
325   20.   Vila A, Cassata A, Pagella H, Amadio C, Yeh KM, Chang FY, Siu LK. 2011.
326   Appearance of *Klebsiella pneumoniae* liver abscess syndrome in Argentina:
327   case report and review of molecular mechanisms of pathogenesis. Open
328   Microbiol J 5:107-13.
329   21.   Pastagia M, Arumugam V. 2008. *Klebsiella pneumoniae* liver abscesses in a
330   public hospital in Queens, New York. Travel Med Infect Dis 6:228-33.
331   22.   Rahimian J, Wilson T, Oram V, Holzman RS. 2004. Pyogenic liver abscess:
332   recent trends in etiology and mortality. Clin Infect Dis 39:1654-9.
333   23.   Russo TA, Olson R, Fang CT, Stoesser N, Miller M, MacDonald U, Hutson A,
334   Barker JH, La Hoz RM, Johnson JR. 2018. Identification of Biomarkers for
335   Differentiation of Hypervirulent *Klebsiella pneumoniae* from Classical *K.*
336   *pneumoniae.* J Clin Microbiol 56.
337   24.   Parrott AM, Shi J, Aaron J, Green DA, Whittier S, Wu F. 2020. Detection of
338   multiple hypervirulent *Klebsiella pneumoniae* strains in a New York City
339   hospital through screening of virulence genes. Clin Microbiol Infect
340   doi:10.1016/j.cmi.2020.05.012.
341   25.   Chou A, Nuila RE, Franco LM, Stager CE, Atmar RL, Zechiedrich L. 2016.
342   Prevalence of hypervirulent *Klebsiella pneumoniae*-associated genes rmpA

and magA in two tertiary hospitals in Houston, TX, USA. J Med Microbiol
65:1047-8.
26.   Peirano G, Pitout JD, Laupland KB, Meatherall B, Gregson DB. 2013.
Population-based surveillance for hypermucoviscosity *Klebsiella pneumoniae*
causing community-acquired bacteremia in Calgary, Alberta. Can J Infect Dis
Med Microbiol 24:e61-4.
27.   Chen YT, Chang HY, Lai YC, Pan CC, Tsai SF, Peng HL. 2004. Sequencing and
analysis of the large virulence plasmid pLVPK of *Klebsiella pneumoniae* CG43.
Gene 337:189-98.
28.   Nassif X, Fournier JM, Arondel J, Sansonetti PJ. 1989. Mucoid phenotype of
Klebsiella pneumoniae is a plasmid-encoded virulence factor. Infect Immun
57:546-52.
29.   Wu KM, Li LH, Yan JJ, Tsao N, Liao TL, Tsai HC, Fung CP, Chen HJ, Liu YM,
Wang JT, Fang CT, Chang SC, Shu HY, Liu TT, Chen YT, Shiau YR, Lauderdale
TL, Su IJ, Kirby R, Tsai SF. 2009. Genome sequencing and comparative
analysis of *Klebsiella pneumoniae* NTUH-K2044, a strain causing liver abscess
and meningitis. J Bacteriol 191:4492-501.
30.   Hsieh PF, Lin TL, Lee CZ, Tsai SF, Wang JT. 2008. Serum-induced iron-
acquisition systems and TonB contribute to virulence in *Klebsiella*
pneumoniae causing primary pyogenic liver abscess. J Infect Dis 197:1717-
27.
31.   Kochan TJ, Ozer EA, Pincus NB, Fitzpatrick MA, Hauser AR. 2020. Complete
Genome Sequence of *Klebsiella pneumoniae* Strain TK421, a Conjugative
Hypervirulent Isolate. Microbiol Resour Announc 9.
32.   Russo TA, Olson R, MacDonald U, Beanan J, Davidson BA. 2015. Aerobactin,
but not yersiniabactin, salmochelin, or enterobactin, enables the
growth/survival of hypervirulent (hypermucoviscous) *Klebsiella pneumoniae*
ex vivo and in vivo. Infect Immun 83:3325-33.
33.   Xiong H, Carter RA, Leiner IM, Tang YW, Chen L, Kreiswirth BN, Pamer EG.
2015. Distinct Contributions of Neutrophils and CCR2+ Monocytes to
Pulmonary Clearance of Different *Klebsiella pneumoniae* Strains. Infect
Immun 83:3418-27.
34.   Bachman MA, Oyler JE, Burns SH, Caza M, Lepine F, Dozois CM, Weiser JN.
2011. *Klebsiella pneumoniae* yersiniabactin promotes respiratory tract
infection through evasion of lipocalin 2. Infect Immun 79:3309-16.
35.   Fang CT, Chuang YP, Shun CT, Chang SC, Wang JT. 2004. A novel virulence
gene in *Klebsiella pneumoniae* strains causing primary liver abscess and
septic metastatic complications. J Exp Med 199:697-705.
36.   Lee CH, Liu JW, Su LH, Chien CC, Li CC, Yang KD. 2010. Hypermucoviscosity
associated with *Klebsiella pneumoniae*-mediated invasive syndrome: a
prospective cross-sectional study in Taiwan. Int J Infect Dis 14:e688-92.
37.   Krapp F, Morris AR, Ozer EA, Hauser AR. 2017. Virulence Characteristics of
Carbapenem-Resistant *Klebsiella pneumoniae* Strains from Patients with
Necrotizing Skin and Soft Tissue Infections. Sci Rep 7:13533.
38.   Harada S, Doi Y. 2018. Hypervirulent *Klebsiella pneumoniae*: a Call for
Consensus Definition and International Collaboration. J Clin Microbiol 56.

39. Walker KA, Miller VL. 2020. The intersection of capsule gene expression, hypermucoviscosity and hypervirulence in *Klebsiella pneumoniae*. Curr Opin Microbiol 54:95-102.

40. Zhang Y, Zhao C, Wang Q, Wang X, Chen H, Li H, Zhang F, Li S, Wang R, Wang H. 2016. High Prevalence of Hypervirulent *Klebsiella pneumoniae* Infection in China: Geographic Distribution, Clinical Characteristics, and Antimicrobial Resistance. Antimicrob Agents Chemother 60:6115-20.

41. Wyres KL, Nguyen TNT, Lam MMC, Judd LM, van Vinh Chau N, Dance DAB, Ip M, Karkey A, Ling CL, Miliya T, Newton PN, Lan NPH, Sengduangphachanh A, Turner P, Veeraraghavan B, Vinh PV, Vongsouvath M, Thomson NR, Baker S, Holt KE. 2020. Genomic surveillance for hypervirulence and multi-drug resistance in invasive *Klebsiella pneumoniae* from South and Southeast Asia. Genome Med 12:11.

42. Shon AS, Bajwa RP, Russo TA. 2013. Hypervirulent (hypermucoviscous) *Klebsiella pneumoniae*: a new and dangerous breed. Virulence 4:107-18.

43. Wang H, Ren Y, Chang Z, Liu Z. 2020. The increased recurrence rate of liver abscess caused by extended-spectrum beta-lactamase-producing *Klebsiella pneumoniae*. Eur J Clin Microbiol Infect Dis 39:1315-1320.

44. Siu LK, Yeh KM, Lin JC, Fung CP, Chang FY. 2012. *Klebsiella pneumoniae* liver abscess: a new invasive syndrome. Lancet Infect Dis 12:881-7.

45. Chou FF, Kou HK. 1996. Endogenous endophthalmitis associated with pyogenic hepatic abscess. J Am Coll Surg 182:33-6.

46. Li J, Ren J, Wang W, Wang G, Gu G, Wu X, Wang Y, Huang M, Li J. 2018. Risk factors and clinical outcomes of hypervirulent *Klebsiella pneumoniae* induced bloodstream infections. Eur J Clin Microbiol Infect Dis 37:679-689.

47. Kim D, Park BY, Choi MH, Yoon EJ, Lee H, Lee KJ, Park YS, Shin JH, Uh Y, Shin KS, Shin JH, Kim YA, Jeong SH. 2019. Antimicrobial resistance and virulence factors of *Klebsiella pneumoniae* affecting 30 day mortality in patients with bloodstream infection. J Antimicrob Chemother 74:190-199.

48. Lambregts MMC, Wijnakker R, Bernards AT, Visser LG, Cessie SL, Boer MGJ. 2020. Mortality after Delay of Adequate Empiric Antimicrobial Treatment of Bloodstream Infection. J Clin Med 9.

49. Weiss SL, Fitzgerald JC, Balamuth F, Alpern ER, Lavelle J, Chilutti M, Grundmeier R, Nadkarni VM, Thomas NJ. 2014. Delayed antimicrobial therapy increases mortality and organ dysfunction duration in pediatric sepsis. Crit Care Med 42:2409-17.

50. Karlsson M, Stanton RA, Ansari U, McAllister G, Chan MY, Sula E, Grass JE, Duffy N, Anacker ML, Witwer ML, Rasheed JK, Elkins CA, Halpin AL. 2019. Identification of a Carbapenemase-Producing Hypervirulent *Klebsiella pneumoniae* Isolate in the United States. Antimicrob Agents Chemother 63.

51. Liu C, Du P, Xiao N, Ji F, Russo TA, Guo J. 2020. Hypervirulent *Klebsiella pneumoniae* is emerging as an increasingly prevalent *K. pneumoniae* pathotype responsible for nosocomial and healthcare-associated infections in Beijing, China. Virulence 11:1215-1224.

52. Leverstein-van Hall MA, Fluit AC, Paauw A, Box AT, Brisse S, Verhoef J. 2002. Evaluation of the Etest ESBL and the BD Phoenix, VITEK 1, and VITEK 2

automated instruments for detection of extended-spectrum beta-lactamases
in multiresistant *Escherichia coli* and *Klebsiella spp.* J Clin Microbiol 40:3703-
11.

53.   Hadano Y. 2013. String test. BMJ Case Rep 2013.

54.   Cherny KE, Ozer EA, Kochan TJ, Kociolek LK. 2020. Complete Genome
Sequence of *Clostridium innocuum* Strain ATCC 14501. Microbiol Resour
Announc 9.

55.   Cherny KE, Ozer EA, Kochan TJ, Johnson S, Kociolek LK. 2020. Complete
Genome Sequence of *Clostridium innocuum* Strain LC-LUMC-CI-001, Isolated
from a Patient with Recurrent Antibiotic-Associated Diarrhea. Microbiol
Resour Announc 9.

56.   Lam MMC, Wyres KL, Judd LM, Wick RR, Jenney A, Brisse S, Holt KE. 2018.
Tracking key virulence loci encoding aerobactin and salmochelin
siderophore synthesis in *Klebsiella pneumoniae.* Genome Med 10:77.

57.   Wick RR, Heinz E, Holt KE, Wyres KL. 2018. Kaptive Web: User-Friendly
Capsule and Lipopolysaccharide Serotype Prediction for *Klebsiella* Genomes.
J Clin Microbiol 56.

58.   Alikhan NF, Petty NK, Ben Zakour NL, Beatson SA. 2011. BLAST Ring Image
Generator (BRIG): simple prokaryote genome comparisons. BMC Genomics
12:402.

Figure Legends

Figure 1. Virulence plasmids associated with hypervirulence in *Klebsiella pneumoniae.* Genomic elements and plasmid architecture are depicted for pK2044 **(A)**, KP52.145pII **(B)** and pTK421 **(C)**. PAL-1 and PAL-2 contents and organization are depicted for pK2044, KP52.145pII, pTK421_2, and ICEKp1 **(D)**. (Drawings not to scale.)

Figure 2. The bloodstream isolates KpN115, KpN8, and KpN89 harbor

pK2044-like plasmids. DNA sequences for plasmids identified by nanopore

sequencing were aligned with either pK2044 **(A)**, or KP52.145pII **(B)** using blast

ring image generator (BRIG). A sequence identity threshold of 85% was used.

Aerobactin biosynthesis genes are indicated with green arrows, salmochelin with

orange arrows, *rmpA* with blue arrows, and *rmpA2* with red arrows. Plasmid and

PALoc alignments are shown in the indicated colors.

Figure 3. The bloodstream isolates KpN49_2 and KpN165_1 harbor

KP52.145pII-like plasmids. DNA sequences for plasmids identified by Nanopore

sequencing were aligned with KP52.145pII using blast ring image generator

(BRIG). A sequence identity threshold of 85% was used. Aerobactin biosynthesis

genes are indicated with green arrows, salmochelin with orange arrows, and

rmpA with blue arrows. Plasmid and PALoc alignments are shown in the

indicated colors.

Figure 4. A unique plasmid containing aerobactin biosynthesis genes was

identified in strain KpN23. DNA sequences for plasmid KpCTRSRTH01_p2

and plasmids from NMH bloodstream isolates **(A)** or pK2044, KP52.145pII, and

pTK421_2 **(B)** were aligned with pKpN23_1 using blast ring image generator

(BRIG). A sequence identity threshold of 85% was used. Aerobactin biosynthesis

genes are indicated with green arrows. Plasmid and PALoc alignments are

shown in the indicated colors.

Table 1. Prevalence of hypervirulence features in blood stream isolates

	Isolates n = 141
	Positive
Hypervirulence Marker	n(%)
iroB	5 (3.5)
iucA	6 (4.3)
iroB/iucA	5 (3.5)
String Test	12 (8.5)
Centrifugation Test	24 (17.0)
Tellurite Resistance	6 (4.3)

Table 2. Characteristics of bloodstream isolates containing hvKp pathogenicity loci

Strain	ST	K-Locus	*iuc*	*iro*	hmv*	*rmpA/2*	Plasmid
KPN8	ST23	KL1	*iuc 1*	*iro 1*	+	*rmpA, rmpA2*	pK2044
KPN23	ST881	KL2	*iuc 3*	-	-	-	pKPN23_1
KPN49	ST66	KL2	*iuc 2*	*iro 2*	+	*rmpA*	KP52.145pII
KPN89	ST23	KL1	*iuc 1*	*iro 1*	+	*rmpA, rmpA2*	pK2044
KPN115	ST23	KL1	*iuc 1*	*iro 1*	+	*rmpA, rmpA2*	pK2044
KPN165	ST380	KL2	*iuc 2*	*iro 2*	+	*rmpA*	KP52.145pII

*hmv = hupermucoviscous; ST = Sequence Type