Neural mechanisms underlying the temporal control of sequential saccade planning in the frontal eye fields.

Debaleena Basu[1*], Naveen Sendhilnathan[2], Aditya Murthy[1]

[1]Centre for Neuroscience, Indian Institute of Science, Bengaluru, Karnataka 560012, India

[2]Department of Neuroscience, Columbia University in the City of New York, NY 10027, USA

*Corresponding author: Dr. Debaleena Basu; e-mail: basu.debaleena@gmail.com

Abbreviated title: Temporal control of sequential saccades

Number of pages: 42;

Number of figures: 6,

Number of supplementary figures: 6

Summary:

Sequences of saccadic eye movements are instrumental in navigating our visual
environment. While neural activity has been shown to ramp up to a threshold before single
saccades, the neural underpinnings of multiple saccades is unknown. To understand the
neural control of rapid saccade sequences, we recorded from the frontal eye field (FEF) of
macaque monkeys while they performed a sequential saccade task. We show that concurrent
planning of two saccade plans brings forth processing bottlenecks, specifically by decreasing
the growth rate and increasing the threshold of saccade-related ramping activity. The rate
disruption affected both saccade plans, and a computational model wherein activity related to
the two saccade plans bilaterally and asymmetrically inhibited each other, predicted the
behavioral and neural results observed experimentally. Borrowing from models in
psychology, our results demonstrate a capacity-sharing mechanism of processing bottlenecks,
wherein multiple saccade plans in a sequence, compete for the processing capacity by
perturbation of the saccade-related ramping activity. Finally, we show that in contrast to
movement related neurons, visual activity in FEF neurons is not affected by the presence of
multiple saccade targets, indicating that for perceptually simple tasks, inhibition amongst
movement-related neurons mainly instantiates capacity sharing. Taken together, we show
how psychology-inspired models of capacity sharing can be mapped onto neural responses to
understand the control of rapid saccade sequences.

Keywords: motor sequences, oculomotor control, electrophysiology, processing bottlenecks

Introduction:

Saccadic eye movements shift the fovea from one point to another, serially sampling our visual surroundings, and aiding consequent behavior. Proper planning and execution of saccade sequences is essential for performing everyday tasks such as reading. Despite extensive research on the neural basis of planning individual saccades, the neural mechanisms underlying the sequencing of multiple saccades remain largely unknown. Previous research has shown that sequential saccades can be processed in parallel (Basu and Murthy, 2020; Becker and Jürgens, 1979; Bhutani et al., 2012; Bhutani et al., 2013; McPeek et al., 2003; McPeek and Keller, 2002; McPeek et al., 2000; Minken et al., 1993; Phillips and Segraves, 2010; Port and Wurtz, 2003; Ray et al., 2004; Sharika et al., 2008; Shen and Paré, 2014; Tian et al., 2000; Wu et al., 2013). Sequential saccade studies have shown that as the temporal gap between the targets (TSD; target step delay) decreases, the latency of the response to the second stimulus increases markedly, as if the brain inherently cannot process two simple decisions at the same time (Pashler, 1994; Marois and Ivanoff, 2005; Ray et al., 2012; Ray et al., 2004; Ruthruff et al., 2001). The bottlenecks associated with parallel programming of multiple saccade plans form the basis of this study.

Various theoretical frameworks have been proposed to explain how closely spaced action plans interfere with each other. Single-channel bottleneck models propose that a central, decision-making stage constitutes the bottleneck, wherein the central stages of multiple plans can only proceed serially and cannot be 'co-active' (Pashler, 1994; Ruthruff et al., 2001; Welford, 1967; Welford, 1952). For a sequence of two saccades, the first plan is likely to reach the central stage first, and thus the saccade 2 plan must 'wait' till central processing of the first is over (**Fig. 1A**). In contrast, capacity-sharing models argue that the decision-making stages of both plans can proceed in parallel, albeit with differential rates. The concept of the brain's 'capacity' corresponds to the brain's general information processing capabilities (Broadbent, 1971; Gopher and Navon, 1980; Kahneman, 1973; McLeod, 1977), independent of task type. The capacity-sharing models predict that because of its temporal precedence, the first saccade plan will get the major share of the capacity and the second saccade plan will get a smaller fraction, thus delaying the onset of the second response (**Fig. 1B**; Navon and Miller, 2002; Tombu and Jolicœur, 2003).

The neural mechanisms of processing bottlenecks in sequential saccade planning are not known. To investigate the neural architecture of saccade-related bottlenecks, we recorded neural activity from the frontal eye field (FEF) of macaque monkeys performing a sequential

saccade task. FEF is a good candidate region to study the neural imprints of processing
bottlenecks since it is a higher-order control center for goal-directed saccadic planning
(Sendhilnathan et al., 2021; Sendhilnathan et al., 2017, 2020). Further, the activity of FEF
movement neurons follow the dynamics of accumulator models and resemble the central
capacity-limited stage observed in computational models of dual-task studies (Hanes and
Schall, 1996; Ray et al., 2012; Sigman and Dehaene, 2005). Finally, FEF movement neurons
can encode two saccade plans in parallel (Basu and Murthy, 2020), and thus, any limitations
arising during the concurrent programming of saccades may be found in the activity of
movement-related neurons in the FEF. Our results show that FEF movement neurons
constitute a bottleneck locus—the processing of saccadic sequences is slowed down by
reducing the speed of activity growth or by increasing movement activation threshold. Such
adjustments were observed for both the first and second saccade plans, indicating that a
capacity-sharing mechanism might underlie temporal delays seen during the sequencing of
multiple actions.
89

Results:

Two monkeys, a *Macaca radiata* (J) and a *Macaca mulatta* (G) performed a
sequential saccade 'FOLLOW' task (**Fig. S1**; see methods), where the majority (70%) of the
trials were 'step trials' in which they had to perform a rapid sequence of saccades to two
targets in the order of their presentation. The remaining 30% of the trials were 'no-step'
trials, wherein a single visual target was presented, and the monkeys had to make a single
saccade to it. The two types of trials were randomly interleaved. The temporal gap (target
step delay or TSD) between the first and second target onsets in step trials was randomly
chosen among 17 ms, 83 ms, and 150 ms (Basu and Murthy, 2020).
99

Behavioral evidence of processing bottlenecks during sequential saccades
In the scheme of single-channel bottleneck models, the second plan shows the
hallmark of processing bottlenecks: increase in latencies with decrease in TSD, whilst the
saccade 1 latencies (RT1) stay unaffected (**Fig 1C** left). However, unlike the single-channel
bottleneck model, where plan 1 may be assumed to get 100% of the capacity, the capacity-
sharing model predict that the latencies of the first saccade (RT1) will also increase as it only
gets a part of the full available capacity (**Fig 1C** right).

107   To ensure that the behavioral data are matched to the neural data, we analyzed trials in

which saccades were made into the response field (RF; see methods). That is, for RT1, the

first saccade was made into the RF, and for RT2, the second saccade was made into the RF.

Both RT1 and RT2 slowed down as the TSD decreased, indicating a capacity-sharing

mechanism (**Fig. 1D**; RT1: Kruskal-Wallis, χ^2 (2, 240) = 17.85, p < .001, η^2 = 0.07; RT2:

Kruskal-Wallis, χ^2 (2, 233) = 158.37, p < .001, η^2 = 0.67). While the effect on RT1 was

typically much smaller than that on RT2, the increases in saccade latencies with decreasing

TSD corroborated with previously well-established evidence of processing bottlenecks in

concurrent action planning. Our behavioral data, thus, supports the presence of a capacity-

sharing bottleneck as opposed to the single-channel bottleneck as the first saccade plan does

not stay unaffected.

Movement-related activity during single saccades:

120   Previous work has shown that the pattern of activity of FEF movement neurons are

correlated with stochastic accumulation, which is widely used in computational models of

saccadic reaction times (Boucher et al., 2007; Hanes and Schall, 1996; Ratcliff et al., 2007;

Woodman et al., 2008) and are directly linked to saccade initiation times (Huerta et al., 1986;

Langer and Kaneko, 1990; Segraves, 1992). Since reaction time lengthening is the main

behavioral evidence of processing bottlenecks, movement neurons are well projected to carry

neural correlates of the same. To confirm whether movement-related activity in FEF adheres

to an accumulation-to-threshold model of reaction time, we first studied the no-step single-

saccade trials. We divided these trials into fast, medium, and slow reaction time groups and

we measured the parameters of accumulator models from the movement activity (**Fig. S2A**).

The reaction time grouping was obtained by partitioning reaction times in each session using

the mean reaction time of that session (see methods). The main parameters of accumulator

models, i.e., baseline, onset, growth rate, and threshold activity were measured for the three-

reaction time conditions (fast, medium, slow), for each neuron (**Fig. S2C-F;** see methods).

Consistent with the earlier studies (Hanes and Schall, 1996), adjustments in the rate of growth

of activity of the movement neuron population predicted reaction times in the no-step trials:

across the movement neuron population, the slope of the best fitting line for growth rate

variation in the reaction time groups was significantly different from zero (Z_{rate} = -4.27, p <

.001; **Fig. S2E**). Further, the slopes for the growth rate were negative, indicating that fast

reaction times were preceded by a steeper rate of growth of movement activity and vice

versa. While the growth rate varied with reaction time, the threshold did not ($Z_{threshold}$ = -0.98,

p = .323; **Fig. S2F**), corroborating with the established reaction time models of accumulation
to a fixed threshold. The slope distributions of other accumulator measures like baseline, and
onset, were not statistically significant from zero ($Z_{baseline}$ = -2.04, p = 0.05; Z_{onset} = 1.92, p =
0.054).

Processing bottlenecks underlie the representation of sequential saccades:

Using a computational model, Sigman and Dehaene (2005) had shown that evidence
accumulation, representing a central decision process, constituted a bottleneck in dual-tasks,
while the perceptual stage and the execution stage did not. Based on the mapping between
accumulator models and movement neuron dynamics, four possible hypotheses (**Fig. S2A**)
can explain how the activity of FEF movement neurons coding for the second saccade might
bring about the systematic increase of the latency of the second saccade (RT2) with decrease
in TSD that characterizes processing bottlenecks. The lengthening of reaction time may be
due to (1) lowering of the baseline firing rate with shorter TSDs (2) delaying of the onset of
the activity related to the second saccade with shorter TSDs (3) reduced growth rate of the
activity with shorter TSDs (4) and an increase of the saccade threshold firing rate with larger
TSDs.

Fig. 2 schematically shows the possible modulations of the accumulation process in
the planning stage (P) and the corresponding movement neuron activity. The accumulation
process is represented as a noisy integrator accumulating visual evidence till it reaches the
threshold. In the single-channel bottleneck model, RT1 is unaffected, and thus the dynamics
of the integrator and the corresponding neural activity will be unchanged across the three
TSDs (**Fig. 2A**). For RT2, the single-channel bottleneck model posits a postponement of the
central stage, thus the onset of the accumulating process and of the neural activity will get
delayed as the overlap between the two saccade plans increases from long to short TSD (**Fig.**
2B). According to the capacity-sharing model, the first and second saccade plans can proceed
in parallel; thus, there is no 'waiting period' for the accumulation process of the second
plan—the onset of neural activity will be similar across TSDs for both first and second
saccade. However, since both motor plans share the limited processing capacity, the central
stages of both plans will be lengthened. This may be brought about by a decrease in the rate
of integration from long to short TSD, or an increase in the decision threshold. At the level of
neural activity, the rate of ramping up of movement-related activity may slow down, or the
threshold firing rate for saccade onset may increase to account for the increase in saccade
latencies with decrease in TSD. Critically, the rate and/or threshold modulation will be

present for both saccade plans according to the capacity-sharing model, although the effect
may be lesser for the first plan as the corresponding increase in RT1 is also less (**Fig. 2C &**
Fig. 2D). While we have presented polarized scenarios for the two bottleneck models, it is
possible that at the population level, there would a combination of the factors mentioned.

To assess which of the above possibilities explain the increase in RT2, we analyzed
the neural activity in trials where the second saccade was made into the RF (RFin trials; see
methods) for all three TSDs (**Fig. 3A**; see **Fig S3** for single neuron example). Across the
population, the rate of neural activity growth slowed down from long to short TSD, and the
activity ramped up to a higher firing rate threshold. We measured each of the four
accumulator parameters: baseline, onset, rate, and threshold (averaged across trials of the
same TSD) for the three TSD conditions, for each neuron (**Fig. 3B;** see methods) using linear
regression. The slopes from all the movement neurons were compared using a Wilcoxon
signed-rank test. Across the movement neuron population, the slopes of the rate and the
threshold, as a function of TSD were significantly different from zero ($Z_{rate} = 4.27$, $p < .001$;
$Z_{threshold} = -2.67$, $p < 0.01$; **Fig. 3B**). Further, the slopes for the rate of activity growth were
positive, indicating that the rate of activity grew faster at longer TSDs, where presumably the
effect of processing bottlenecks was the least among the three TSD conditions. Threshold
slopes were significantly negative, indicating that as the TSD increased, the threshold
required for initiation of the second saccade was reduced at the population level. However,
the slope distributions of other accumulator measures like baseline, and onset, were not
statistically significant from zero ($Z_{baseline} = -0.62$, $p = 0.53$; $Z_{onset} = 0.86$, $p = 0.17$). Thus,
processing bottlenecks at the level of FEF movement neurons were characterized by
multifaceted adjustments in the rate and threshold of the activity related to S2.

While the classical evidence of processing bottlenecks is indexed by the increase in
RT2, RT1 may also be affected according to the capacity-sharing scheme of processing
bottlenecks (**Fig. 1B**). We tested whether movement-related activity encoding first saccade
remained unchanged as would be expected in the single-channel bottleneck scheme, or
changed systematically, across TSDs as the capacity-sharing model predicted. To address this
issue, we performed the same analyses as before but for the condition in which the first
saccade was made into the RF (RFout trials; **Fig 3C**; see **Fig S3** for single neuron example).
At the population level, rate perturbation occurred with decrease in TSD in the first plan,
mirroring the modulation observed for the second plan (Wilcoxon signed-rank test for slopes
of rates, $Z_{rate} = 3.62$, $p < .001$; **Fig 3D**). However, unlike the second plan, threshold activity

did not show a significantly decreasing relation with TSD ($Z_{threshold}$ = 1.16, p = 0.25;). Slope

distributions of other accumulator measures like baseline, and onset, were not statistically

significant from zero ($Z_{baseline}$ = 0.85, p = 0.39; Z_{onset} = 1.04, p = 0.29; **Fig 3D**). Thus, rate

perturbation constituted a major mechanism through which the ramping up of activity of FEF

movement neurons was controlled during parallel planning of sequential saccades.

State space dynamics and inhibitory control may enable capacity sharing during
sequential saccade planning

To gain deeper insights into neural mechanisms underlying capacity sharing, we

studied the population dynamics underlying the trajectory of neural activity in FEF. First, we

visualized this by performing a principal component analysis (PCA) separately for the

population neural activity (for saccades into RF) aligned to target 1 and target 2 onsets for

each of the three TSDs (**Fig 4A**). PCA is a commonly used unsupervised learning algorithm

to extract the latent information from the data (see methods). This method allows us to look

at the high dimensional FEF population neural activity in a much lower dimension that

captures the maximum variance of the population. At least ~ 7-8 PCs were required to

explain >99% of the variance for any of the six conditions (three TSDs, two plans; although

there was a trend of fewer PCs explaining more variance as TSD increased). However, the

top three PCs explain >90% variance. Therefore, we visualized a 'state-space trajectory' by

plotting the top three PCs versus one another (**Fig 4B**). Each point on the trajectory indicates

the neural state at each time point.

If planning for the first and the second saccades are processed in parallel but compete

for the same shared space due to limited capacity (according to the capacity sharing model),

we should expect the neural trajectories to span different subspaces at shorter TSDs and span

the same subspace at higher TSDs. That is, in the lowest possible TSD, we should expect the

two subspaces to be completely orthogonal (no overlap) and as the TSD increases and

approaches the reaction time of the first saccade, the subspaces can begin to overlap.

Therefore, in our case with the lowest TSD being 17 ms, we should expect a low degree of

overlap and at TSD = 150 ms (~RT1), we should expect a high degree of overlap. In contrast,

the single-channel bottleneck hypothesis predicts that the subspaces corresponding to the

planning of the first and the second saccades would completely overlap, since the plan 2

would be completely dormant until plan 1 is completed.

We found that the neural trajectories significantly differed between the planning of
the first and the second saccades for the shortest TSD but became more similar as the TSD
increased (**Fig 5D**). We quantified the degree of overlap between the subspaces spanned by
these neural trajectories (see methods). At the shortest TSD, the magnitude of overlap
between the signals for planning of the first and the second saccades was 47% and this
increased as the TSD increased from medium (84%) to long TSDs (92%; **Fig 4C**)**,** aligned
more with the predictions of the capacity sharing model. This result also held true for all
saccade directions (**Fig S4**). We also confirmed that these differences were related to the
TSDs and not to differences in saccade kinematics, which were similar across TSDs for the
first and the second saccades (**Fig S5**).

Next, we investigated the mechanism behind the differences in the neural subspace
overlap among different TSDs. We performed two sets of simulations (see methods; **S5A-F**)
using the accumulator framework (**Fig S5H**). For each of the two sets, we simulated 40
neurons with 900 trials per neuron (with three types of TSD trials) using a firing rate model
to approximately match the statistical power of our experimental dataset (see methods). We
constructed an inhibition function such that the magnitude of the inhibition inversely varied
with TSD (see methods; **Fig S5G**).

In the first set of simulations, we introduced a unilateral inhibition (**Fig 5A;** see
methods). Here, the activities for plan 2 were temporally shifted by plan 1 following the
inhibition curve as a function of TSD. The resulting simulated neural activities (**Fig 5B**)
resembled the predictions of a single-channel bottleneck model (**Fig 2A-B**). Very few (~3)
PCs explained >99% of the variance. The state-space neural trajectories were not
significantly different between planning of the first and the second saccades for any of the
TSDs (**Fig 5C**) as the subspace overlap was 98% between any pair of plans (**Fig 5D**), as
expected from the single-channel bottleneck model.

In the next set of simulations, we introduced bilateral, asymmetric mutual inhibition
(see methods; **Fig 5E**). That is plan 1 temporally shifted plan 2 just like before but plan 2
reduced the magnitude of peak firing of plan 1. Hence the nature of inhibition is both bilateral
and asymmetric. The simulations of this model (**Fig 5F**) resembled the neural data (**Fig 3A,
C**) and the predictions of a capacity sharing bottleneck model (**Fig 2C-D**). Here, ~ 7-8 PCs
were required to explain >99% of the variance for the shortest TSD and fewer (~5-6) PCs
were required to explain >99% of the variance for the longest TSD. The neural trajectories
significantly differed for short TSD but were similar for longer TSDs (**Fig 5G**) and the
degree of subspace overlap between the two plans increased with TSD, consistent with the

structure present in the neural data (**Fig 5H**) resembling the experimental data (**Fig 4C**), as
expected from the capacity sharing model.

FEF visually-related neurons do not show processing bottlenecks

Previous studies have reported a separation between the visual and motor processing
of FEF neurons with only motor processing affecting reaction time in perceptually simple
tasks (Sato et al., 2001; Thompson et al., 1997; Woodman et al., 2008). Thus, it is plausible
that the responses of visual neurons are not gated by inhibitory bottlenecks. This notion was
tested by analyzing target-related activity in purely visual (**Fig 6A**) and visuomovement
neurons (**Fig 6B**).

We analyzed the average target-related response in the 200 ms window following
target onset for each neuron to identify signatures of processing bottlenecks. If target
selection is capacity-limited, then presumably, neural responses encoding saccade targets
appearing in close succession will be inhibited, either due to single-channel bottleneck (only
second target response gets affected) or due to capacity sharing (both first and second target
responses get affected). In contrast to movement-related activity, the average activity in the
target-related period did not vary with TSD (Kruskalwallis: χ^2 (2, 129) = 0.47, p = .79 (first
saccade); χ^2 (2, 124) = 0.06, p = .97 (S2)) for both saccade plans, suggesting that the visual
processing stage is pre-bottleneck, at least of a perceptually simple task like the FOLLOW
task.

Discussion:

In this study, we explored the limits of parallel processing involved in saccade
sequences. Processing bottlenecks were found within FEF, the mechanisms being rate
perturbation and threshold modulation in the movement neuron population. Additionally, we
found evidence of processing bottlenecks for both motor plans for the first and the second
saccades, suggesting that the associated bottleneck could be a consequence of capacity
sharing between co-activated movement plans. The notion of such shared and limited
processing was also revealed in the state space dynamics of FEF movement activity, which
showed a potential role for inhibitory control that gated access of concurrent motor plans to a
planning subspace. Our analysis of visual activity did not reflect any consistent modulation

that could be considered a significant bottleneck. The major results are discussed and
interpreted in the following sections.

Processing bottlenecks in sequential saccade planning

Processing bottlenecks and parallel programming represent functionally antithetical
processes, and yet both are essential for optimal saccadic behavior. While parallel
programming allows for rapid execution of a saccade sequence, processing bottlenecks are
likely to arise to check unbridled parallel programming of motor plans, as failure to control it
might lead to errors like averaged saccades or incorrect order of execution of a saccade
sequence (Bhutani et al., 2012; Coëffé and O'regan, 1987; Findlay, 1982; Ray et al., 2012;
Viviani and Swensson, 1982; Zambarbieri et al., 1987). In the context of the current study,
we tested whether a single-channel bottleneck (Pashler, 1994) or a capacity-sharing
bottleneck (Kahneman, 1973) best explained our reaction time data since behavioral evidence
of both the models have been found in dual-task paradigms (Arnell and Duncan, 2002; Navon
and Miller, 2002; Pashler, 1994). In our data, we found evidence of increase in both RT1 and
RT2 with TSD, ruling out the single-channel bottleneck model being the exclusive
framework underlying bottlenecks in sequential saccades. Our neural data also suggested a
capacity-sharing mechanism of bottlenecks: the onset of saccade-related activity did not vary
with TSD as predicted by the single-channel bottleneck hypothesis (see **Fig. 2**), and both
saccade plans showed consistent activity modulations with TSD. A reduction in the rate of
accumulation and an increase in the threshold activity level were seen for the second saccade
plan. In contrast, only changes in the slope of the activity corresponding to the first saccade
were observed, which may account for the more subtle changes in RT1.

Inhibitory control underlying processing bottlenecks

We tested whether mutually inhibitory accumulators encoding distinct saccade plans
can mimic capacity sharing, wherein both the saccadic eye movements are executed with
delays, especially for the second saccade. Modelling such a response required two important
conditions: the first condition required that the inhibition be asymmetric, being greater for the
first saccade plan than the second saccade plan, which manifest as greater capacity and faster
information processing for the former compared to the latter. Such an asymmetry is a natural
consequence of the temporal delay allowing for greater activity in the first saccade to inhibit
the second saccade; the second condition required an inhibitory kernel that decreased with
target step delay, such that inhibition from the first accumulator to be greater at shorter delays

despite being the level of activity in the accumulator being lesser compared to what it would

be at larger target step delays. Such an inhibitory kernel is necessary to match the observed

behavioral data of greater second saccade reaction times as well as the neural data which

showed greater interference for the second saccade motor plans at the shorter TSDs.

Interestingly, using a dynamical systems approach under the assumption of stationarity of

noise across trials (Elsayed et al., 2016), this model of inhibitory control could be also shown

to act as a "queuing" mechanism, in which non-orthogonal neural spaces can simultaneously

allow parallel processing but yet temporarily slow the processing of the second saccade. We

believe that the ability of such inhibition to reconfigure the neural space may reflect the

nonlinear effects of inhibition on the pattern of activity representing accumulator activity that

underlie the saccades.

The simplest and most parsimonious explanation for the location of such a bottleneck

is at the level of FEF via bilateral mutual inhibition (Ray et al., 2009) of competing motor

plans developing in the FEF. This type of inhibitory gating can be brought about by

inhibitory interneurons within the FEF (Markram et al., 2004; Somogyi, 1977). Although

such a form of inhibition is intuitive and can be readily implemented within the proposed

frameworks described for decision-making circuits (Bogacz et al., 2006; Ratcliff and Smith,

2004), implementing an inhibitory kernel that decreases with increasing TSD cannot be easily

implemented in a straightforward manner by mutually inhibitory accumulators. Furthermore,

using an identical task, our previous work has shown that the basal ganglia is causally

involved in the conversion of parallel movement plans into sequential behavior (Bhutani et

al., 2013). Inactivation of the basal ganglia in monkeys with muscimol or impairment of the

basal ganglia in Parkinson's disease patients resulted in a significantly greater extent of

saccadic errors that develop due to unchecked parallel programming leading to a 'collision'

of saccade plans. The results of both these studies can be reconciled by the fact that FEF and

basal ganglia share a closed connection through the cortico-BG-thalamo-cortical loop

wherein the thalamus, a major relay center, receives projections from BG output nuclei, and

in turn projects to multiple cortical regions, including the FEF, which are again routed to the

input nuclei of basal ganglia (Alexander et al., 1986; Middleton and Strick, 2000; Parent and

Hazrati, 1995a, b). Thus , the origin of the bottleneck could also be in the well-established

inhibitory control circuitry of the basal ganglia (Hikosaka et al., 2000) and then re-routed to

the FEF through the basal ganglia-thalamo-cortical loop (Goldman-Rakic and Porrino, 1985),

which then manifests it various adjustments of movement-related neuronal activity.

Neural representations of processing bottlenecks within FEF

Our data show robust signatures of processing bottlenecks involving rate and
threshold adjustments of FEF movement neurons contributing to the observed processing
bottlenecks. Interestingly, similar adjustments of rate have been observed in FEF movement-
related neurons when monkeys slow their reaction times to improve their accuracy (Heitz and
Schall, 2012), consistent with movement-related activity reflecting a developing motor plan
that can be adjusted by strategic requirements of the task. However, in contrast to
speed/accuracy adjustments, we did find systematic increases in threshold for the second
saccade with shorter TSDs that together with decreases in accumulation rate, contribute to the
lengthening of reaction times for the second saccade. Interestingly, similar changes in growth
rate for both the first and second saccade, particularly at shorter TSDs, were also observed in
our model of mutually inhibiting accumulators but without any changes in threshold (**Fig 5**),
raising the possibility that these changes may involve additional processes such as
adjustments in the excitability of superior colliculus neurons from the basal ganglia (Lo and
Wang, 2006; Wurtz and Hikosaka, 1986) that were not modelled here.

In contrast to the movement neurons, the activity of visual neurons displayed little
evidence of active inhibitory control, suggesting that they are 'pre-bottleneck'. This is not
surprising since many studies have reported a separation between the visual and motor
processing of FEF neurons with only motor processing affecting reaction time in perceptually
simple tasks, thus it is plausible that the responses of visual neurons are not gated by
inhibitory bottlenecks for our task. However, it can be speculated that in a more perceptually
challenging task, manifestations of processing bottlenecks would show up in the activity of
visual responses as well. Thereby, it can be concluded that movement neurons, which are
thought to be functionally downstream of visual neurons (Woodman et al., 2008), are
subjected to a greater degree of inhibitory control, possibly due to its direct role in saccade
initiation. A similar result was observed in the countermanding (Hanes et al., 1998) and
redirect tasks (Murthy et al., 2009), where movement-related neurons showed the strongest
evidence of inhibitory control that reflected the monkeys' abilities to withhold or change
saccade plans. Thus, movement-related activity would fall under the 'post-/peri- bottleneck'
category while visually-related activity would be 'pre-bottleneck', at least for perceptually
simple tasks.

References

Alexander, G.E., DeLong, M.R., and Strick, P.L. (1986). Parallel organization of functionally
segregated circuits linking basal ganglia and cortex. Annual review of neuroscience *9*, 357-
381.
409

410 Arnell, K.M., and Duncan, J. (2002). Separate and shared sources of dual-task cost in
411 stimulus identification and response selection. Cognitive psychology *44*, 105-147.
412

413 Basu, D., and Murthy, A. (2020). Parallel programming of saccades in the macaque frontal
414 eye field: are sequential motor plans coactivated? Journal of neurophysiology *123*, 107-119.
415

416 Becker, W., and Jürgens, R. (1979). An analysis of the saccadic system by means of double
417 step stimuli. Vision research *19*, 967-983.
418

419 Bhutani, N., Ray, S., and Murthy, A. (2012). Is saccade averaging determined by visual
420 processing or movement planning? Journal of neurophysiology *108*, 3161-3171.
421

422 Bhutani, N., Sureshbabu, R., Farooqui, A.A., Behari, M., Goyal, V., and Murthy, A. (2013).
423 Queuing of concurrent movement plans by basal ganglia. Journal of Neuroscience *33*, 9985-
424 9997.
425

426 Bogacz, R., Brown, E., Moehlis, J., Holmes, P., and Cohen, J.D. (2006). The physics of
427 optimal decision making: a formal analysis of models of performance in two-alternative
428 forced-choice tasks. Psychological review *113*, 700.
429

430 Boucher, L., Palmeri, T.J., Logan, G.D., and Schall, J.D. (2007). Inhibitory control in mind
431 and brain: an interactive race model of countermanding saccades. Psychological review *114*,
432 376.
433

434 Broadbent, D.E. (1971). Decision and stress.
435

436 Bruce, C.J., and Goldberg, M.E. (1985). Primate frontal eye fields. I. Single neurons
437 discharging before saccades. Journal of neurophysiology *53*, 603-635.
438

439 Churchland, M.M., Cunningham, J.P., Kaufman, M.T., Foster, J.D., Nuyujukian, P., Ryu, S.I.,
440 and Shenoy, K.V. (2012). Neural population dynamics during reaching. Nature *487*, 51-56.
441

Coëffé, C., and O'regan, J.K. (1987). Reducing the influence of non-target stimuli on

saccade accuracy: Predictability and latency effects. Vision research *27*, 227-240.

Ding, L., and Gold, J.I. (2012). Separate, causal roles of the caudate in saccadic choice and

execution in a perceptual decision task. Neuron *75*, 865-874.

Elsayed, G.F., Lara, A.H., Kaufman, M.T., Churchland, M.M., and Cunningham, J.P. (2016).

Reorganization between preparatory and movement population responses in motor cortex.

Nature communications *7*, 1-15.

Findlay, J.M. (1982). Global visual processing for saccadic eye movements. Vision research

22, 1033-1045.

Gold, J.I., and Shadlen, M.N. (2007). The neural basis of decision making. Annual review of

neuroscience *30*.

Goldman-Rakic, P.S., and Porrino, L.J. (1985). The primate mediodorsal (MD) nucleus and

its projection to the frontal lobe. Journal of Comparative Neurology *242*, 535-560.

Gopher, D., and Navon, D. (1980). How is performance limited: Testing the notion of central

capacity. Acta psychologica *46*, 161-180.

Hanes, D.P., Patterson, W.F., and Schall, J.D. (1998). Role of frontal eye fields in

countermanding saccades: visual, movement, and fixation activity. Journal of

neurophysiology *79*, 817-834.

Hanes, D.P., and Schall, J.D. (1996). Neural control of voluntary movement initiation.

Science *274*, 427-430.

Heitz, R.P., and Schall, J.D. (2012). Neural mechanisms of speed-accuracy tradeoff. Neuron

76, 616-628.

Hikosaka, O., Takikawa, Y., and Kawagoe, R. (2000). Role of the basal ganglia in the control

of purposive saccadic eye movements. Physiological reviews *80*, 953-978.

Huerta, M.F., Krubitzer, L.A., and Kaas, J.H. (1986). Frontal eye field as defined by
intracortical microstimulation in squirrel monkeys, owl monkeys, and macaque monkeys: I.
Subcortical connections. Journal of Comparative Neurology *253*, 415-439.
480

481   Jagadisan, U.K., and Gandhi, N.J. (2016). Disruption of fixation reveals latent sensorimotor
482   processes in the superior colliculus. Journal of Neuroscience *36*, 6129-6140.
483

484   Kahneman, D. (1973). Attention and effort, Vol 1063 (Citeseer).
485

486   Langer, T.P., and Kaneko, C.R. (1990). Brainstem afferents to the oculomotor omnipause
487   neurons in monkey. Journal of Comparative Neurology *295*, 413-427.
488

489   Lo, C.-C., and Wang, X.-J. (2006). Cortico–basal ganglia circuit mechanism for a decision
490   threshold in reaction time tasks. Nature neuroscience *9*, 956-963.
491

492   Markram, H., Toledo-Rodriguez, M., Wang, Y., Gupta, A., Silberberg, G., and Wu, C. (2004).
493   Interneurons of the neocortical inhibitory system. Nature reviews neuroscience *5*, 793-807.
494

495   Marois, R., and Ivanoff, J. (2005). Capacity limits of information processing in the brain.
496   Trends in cognitive sciences *9*, 296-305.
497

498   McLeod, P. (1977). A dual task response modality effect: Support for multiprocessor models
499   of attention. Quarterly Journal of Experimental Psychology *29*, 651-667.
500

501   McPeek, R.M., Han, J.H., and Keller, E.L. (2003). Competition between saccade goals in the
502   superior colliculus produces saccade curvature. Journal of neurophysiology *89*, 2577-2590.
503

504   McPeek, R.M., and Keller, E.L. (2002). Superior colliculus activity related to concurrent
505   processing of saccade goals in a visual search task. Journal of Neurophysiology *87*, 1805-
506   1815.
507

508   McPeek, R.M., Skavenski, A.A., and Nakayama, K. (2000). Concurrent processing of
509   saccades in visual search. Vision research *40*, 2499-2516.
510

511   Middleton, F.A., and Strick, P.L. (2000). Basal ganglia and cerebellar loops: motor and
512   cognitive circuits. Brain research reviews *31*, 236-250.
513

Minken, A., Van Opstal, A., and Van Gisbergen, J. (1993). Three-dimensional analysis of
strongly curved saccades elicited by double-step stimuli. Experimental Brain Research *93*,
521-533.

Murthy, A., Ray, S., Shorter, S.M., Priddy, E.G., Schall, J.D., and Thompson, K.G. (2007).
Frontal eye field contributions to rapid corrective saccades. Journal of Neurophysiology *97*,
1457-1469.

Murthy, A., Ray, S., Shorter, S.M., Schall, J.D., and Thompson, K.G. (2009). Neural control
of visual search by frontal eye field: effects of unexpected target displacement on visual
selection and saccade preparation. Journal of Neurophysiology *101*, 2485-2506.

Navon, D., and Miller, J. (2002). Queuing or sharing? A critical evaluation of the single-
bottleneck notion. Cognitive psychology *44*, 193-251.

Nosofsky, R.M., and Palmeri, T.J. (1997). Comparing exemplar-retrieval and decision-bound
models of speeded perceptual classification. Perception & Psychophysics *59*, 1027-1048.

Parent, A., and Hazrati, L.-N. (1995a). Functional anatomy of the basal ganglia. I. The
cortico-basal ganglia-thalamo-cortical loop. Brain research reviews *20*, 91-127.

Parent, A., and Hazrati, L.-N. (1995b). Functional anatomy of the basal ganglia. II. The place
of subthalamic nucleus and external pallidium in basal ganglia circuitry. Brain research
reviews *20*, 128-154.

Pashler, H. (1994). Dual-task interference in simple tasks: data and theory. Psychological
bulletin *116*, 220.

Phillips, A.N., and Segraves, M.A. (2010). Predictive activity in macaque frontal eye field
neurons during natural scene searching. Journal of neurophysiology *103*, 1238-1252.

Port, N.L., and Wurtz, R.H. (2003). Sequential activity of simultaneously recorded neurons in
the superior colliculus during curved saccades. Journal of neurophysiology *90*, 1887-1903.

Purcell, B.A., Heitz, R.P., Cohen, J.Y., and Schall, J.D. (2012). Response variability of frontal
eye field neurons modulates with sensory input and saccade preparation but not visual
search salience. Journal of neurophysiology *108*, 2737-2750.

Purcell, B.A., Heitz, R.P., Cohen, J.Y., Schall, J.D., Logan, G.D., and Palmeri, T.J. (2010).

Neurally constrained modeling of perceptual decision making. Psychological review *117*,

1113.

Ramakrishnan, A., and Murthy, A. (2013). Brain mechanisms controlling decision making

and motor planning. In Progress in brain research (Elsevier), pp. 321-345.

Ratcliff, R., Hasegawa, Y.T., Hasegawa, R.P., Smith, P.L., and Segraves, M.A. (2007). Dual

diffusion model for single-cell recording data from the superior colliculus in a brightness-

discrimination task. Journal of neurophysiology *97*, 1756-1774.

Ratcliff, R., and Rouder, J.N. (1998). Modeling response times for two-choice decisions.

Psychological science *9*, 347-356.

Ratcliff, R., and Smith, P.L. (2004). A comparison of sequential sampling models for two-

choice reaction time. Psychological review *111*, 333.

Ray, S., Bhutani, N., and Murthy, A. (2012). Mutual inhibition and capacity sharing during

parallel preparation of serial eye movements. Journal of vision *12*, 17-17.

Ray, S., Pouget, P., and Schall, J.D. (2009). Functional distinction between visuomovement

and movement neurons in macaque frontal eye field during saccade countermanding.

Journal of Neurophysiology *102*, 3091-3100.

Ray, S., Schall, J.D., and Murthy, A. (2004). Programming of double-step saccade

sequences: modulation by cognitive control. Vision research *44*, 2707-2718.

Ruthruff, E., Pashler, H.E., and Klaassen, A. (2001). Processing bottlenecks in dual-task

performance: Structural limitation or strategic postponement? Psychonomic bulletin & review

8, 73-80.

Sato, T., Murthy, A., Thompson, K.G., and Schall, J.D. (2001). Search efficiency but not

response interference affects visual selection in frontal eye field. Neuron *30*, 583-591.

Segraves, M.A. (1992). Activity of monkey frontal eye field neurons projecting to oculomotor

regions of the pons. Journal of Neurophysiology *68*, 1967-1985.

Segraves, M.A., and Goldberg, M.E. (1987). Functional properties of corticotectal neurons in
the monkey's frontal eye field. Journal of Neurophysiology *58*, 1387-1419.

Sendhilnathan, N., Basu, D., Goldberg, M.E., Schall, J.D., and Murthy, A. (2021). Neural
correlates of goal-directed and non–goal-directed movements. Proceedings of the National
Academy of Sciences *118*.

Sendhilnathan, N., Basu, D., and Murthy, A. (2017). Simultaneous analysis of the LFP and
spiking activity reveals essential components of a visuomotor transformation in the frontal
eye field. Proceedings of the National Academy of Sciences *114*, 6370-6375.

Sendhilnathan, N., Basu, D., and Murthy, A. (2020). Assessing within-trial and across-trial
neural variability in macaque frontal eye fields and their relation to behaviour. European
Journal of Neuroscience.

Sharika, K., Ramakrishnan, A., and Murthy, A. (2008). Control of predictive error correction
during a saccadic double-step task. Journal of neurophysiology *100*, 2757-2770.

Shen, K., and Paré, M. (2014). Predictive saccade target selection in superior colliculus
during visual search. Journal of Neuroscience *34*, 5640-5648.

Sigman, M., and Dehaene, S. (2005). Parsing a cognitive task: a characterization of the
mind's bottleneck. PLoS biology *3*, e37.

Smith, P.L., and Van Zandt, T. (2000). Time-dependent Poisson counter models of response
latency in simple judgment. British Journal of Mathematical and Statistical Psychology *53*,
293-315.

Sommer, M.A., and Wurtz, R.H. (2000). Composition and topographic organization of signals
sent from the frontal eye field to the superior colliculus. Journal of Neurophysiology *83*,
1979-2001.

Somogyi, P. (1977). A specific 'axo-axonal'interneuron in the visual cortex of the rat. Brain
Res *136*, 345-350.

Thompson, K.G., Bichot, N.P., and Schall, J.D. (1997). Dissociation of visual discrimination
from saccade programming in macaque frontal eye field. Journal of neurophysiology *77*,
1046-1050.

Thompson, K.G., Hanes, D.P., Bichot, N.P., and Schall, J.D. (1996). Perceptual and motor
processing stages identified in the activity of macaque frontal eye field neurons during visual
search. Journal of neurophysiology *76*, 4040-4055.

Tian, J., Schlag, J., and Schlag-Rey, M. (2000). Testing quasi-visual neurons in the
monkey's frontal eye field with the triple-step paradigm. Experimental brain research *130*,
433-440.

Tombu, M., and Jolicœur, P. (2003). A central capacity sharing model of dual-task
performance. Journal of Experimental Psychology: Human Perception and Performance *29*,
3.

Usher, M., and McClelland, J.L. (2001). The time course of perceptual choice: the leaky,
competing accumulator model. Psychological review *108*, 550.

Viviani, P., and Swensson, R.G. (1982). Saccadic eye movements to peripherally
discriminated visual targets. Journal of Experimental Psychology: Human Perception and
Performance *8*, 113.

Welford, A. (1967). Single-channel operation in the brain. Acta psychologica *27*, 5-22.

Welford, A.T. (1952). The psychological refractory period and the timing of high-speed
performance-a review and a theory. British Journal of Psychology *43*, 2.

Westheimer, G. (1954). Eye movement responses to a horizontally moving visual stimulus.
AMA archives of ophthalmology *52*, 932-941.

Wheeless, L.L., Boynton, R.M., and Cohen, G.H. (1966). Eye-movement responses to step
and pulse-step stimuli. JOSA *56*, 956-960.

Woodman, G.F., Kang, M.-S., Thompson, K., and Schall, J.D. (2008). The effect of visual
search efficiency on response preparation: neurophysiological evidence for discrete flow.
Psychological science *19*, 128-136.

Wu, E.X., Gilani, S.O., van Boxtel, J.J., Amihai, I., Chua, F.K., and Yen, S.-C. (2013).

Parallel programming of saccades during natural scene viewing: Evidence from eye

movement positions. Journal of vision *13*, 17-17.

Wurtz, R.H., and Hikosaka, O. (1986). Role of the basal ganglia in the initiation of saccadic

eye movements. In Progress in brain research (Elsevier), pp. 175-190.

Zambarbieri, D., Schmid, R., and Ventre, J. (1987). Saccadic eye movements to predictable

visual and auditory targets. In Eye Movements from Physiology to Cognition (Elsevier), pp.

131-140.

Zylberberg, A., Ouellette, B., Sigman, M., and Roelfsema, P.R. (2012). Decision making

during the psychological refractory period. Current biology *22*, 1795-1799.

METHODS:

678
679   **KEY RESOURCES TABLE**

REAGENT OR RESOURCE	SOURCE	IDENTIFIER
Experimental models: Organisms/Strains		
Rhesus macaque (*Macaca mulatta*)		**N/A**
Bonnet macaque (*Macaca radiata*)		**N/A**
Software and Algorithms		
MATLAB	**Mathworks**	**https://www.mathworks.com/products/matlab.html**
Blackrock	**Blackrock Microsytstems**	**https://www.blackrockmicro.com/**
ISCAN eye tracking system	**ISCAN**	**http://iscaninc.com/**
Others		
Tungsten microelectrode	**FHC**	**https://www.fh-co.com/product/metal-microelectrodes/**

680
681
682   **CONTACT FOR REAGENT AND RESOURCE SHARING**
683   Further information and requests for resources and reagents should be directed to and
684   will be fulfilled by the Lead Contact, Debaleena Basu (basu.debaleena@gmail.com).

685
686
687   **EXPERIMENTAL MODEL AND SUBJECT DETAILS**
688   The detailed methods pertaining to this dataset has been published in a previous study
689   (Basu & Murthy 2020; Sendhilnathan et al., 2021). A brief overview is given below.

Single-unit recordings were done from two adult monkeys (J, male *Macaca radiata*, and G, female *Macaca mulatta*). The animals were cared for in accordance with the animal ethics guidelines of the Committee for the Purpose of Control and Supervision of Experiments on Animals (CPCSEA), Government of India, and the Institutional Animal Ethics Committee (IAEC) of the Indian Institute of Science (IISc.).

Surgical Procedures

Each monkey underwent two surgeries: first, to implant a titanium headpost for the purpose of head-fixation during experiments, and second to make an MRI-guided craniotomy over the FEF and implant a recording chamber (Crist instruments, USA). Training or recording sessions were conducted only after the monkeys completed surgical recovery.

METHOD DETAILS

Behavioral tasks:

Monkeys were trained on two oculomotor tasks: the memory-guided (MG) saccade task and the FOLLOW saccade task. Trials in the MG task began with a red fixation point ($0.6° \times 0.6°$) which was presented in the center of a screen. After a variable fixation period (~300 ms), a gray target stimulus ($1° \times 1°$) was presented peripherally. Post-appearance, the target disappeared after 100 ms; however, the monkeys were required to fixate for a delay period of around 1000 ms. The fixation spot was extinguished after the delay period, following which the monkeys had to make a saccade to the remembered target location. Correct trials were reinforced with juice rewards. The delay period served to aid the classification of FEF neurons by isolating the stimulus-related (visual) and saccade-related (motor) epochs.

The FOLLOW task (**Fig S1**) is a modified version of the double-step task (Becker and Jürgens, 1979; Westheimer, 1954; Wheeless et al., 1966), where single saccade no-step trials (30%) were randomly interleaved with sequential saccade step trials (70%). Trials started with fixation, following which a green saccade target ($1° \times 1°$) was presented in one of the

six possible peripheral locations (eccentricity 12º). The fixation spot was removed at target

onset. In no-step trials, the monkeys had to execute a single, correct saccade to the target. In

step trials a red second target ($1° \times 1°$) appeared after target 1, signaling the monkey to make

ordered sequential saccades. Step trials comprised two targets, the first one being same as in

the no-step trials. After a variable time delay (target step delay (TSD): 17 ms, 83 ms, or 150

ms), a red target was displayed. Monkeys had to make an additional second saccade from

target 1 to target 2 to get rewarded in step trials.

Response field (RF) identification was done using the MG task. The RF center, and

the two flanking positions were set as 'RFin' positions and the three diametrically opposite

positions were considered 'RFout' positions. No-step targets and the first target of step trials

could appear at any one of the six RFin and RFout locations. The second target in step trials

was presented in any one of three positions diametrically opposite to the location of the first

target. Based on this scheme, RFin trials refer to trials in which the second target or target 2

was presented in the RF while target 1 was outside RF. RFout trials are those in which target

1 was inside RF and target 2 was outside. Neural activity in RFin trials would mainly encode

the second target or second saccade, while RFout trials would represent the first target or the

saccade.

Recording setup and procedures:

The tasks were designed and displayed using TEMPO and VIDEOSYNC software

(Reflective computing, St. Louis, MO, USA). A Sony Bravia LCD monitor (42 inches, 60 Hz

refresh rate; 640×480 resolution) was used to show the task stimuli to the monkeys. An

infrared eye tracker (ISCAN, Woburn, MA USA) was used to track the pupils throughout the

recording session.

Neural recordings were undertaken using tungsten microelectrodes (FHC, Bowdoin,

ME, USA, impedance 2 - 4 MΩ). A Cerebus data acquisition system (Blackrock

Microsystems, Salt Lake City, UT, USA), which was synchronized to the TEMPO software,

was used to sample and store neuronal data at 30,000 Hz. In each recording session, the MG

task was used to identify and classify FEF neurons. After RF and cell type was identified, the

FOLLOW task was started.

QUANTIFICATION AND STATISTICAL ANALYSIS

Data Analysis:

The collected neural data was sorted offline using the in-built spike-sorting tool of
Cerebus system (Blackrock Microsystems). Saccades were detected from eye position data
using a 30°/s velocity threshold. Analysis of the data was done using MATLAB (MathWorks,
Natick, MA, USA). This study used only trials with correct responses, with at least eight
trials per condition as the inclusion criteria. The final dataset for this study comprised 84 FEF
neurons. A filter mimicking an excitatory post-synaptic potential (EPSP) was used to
convolve spike data (Murthy et al., 2007).

The classification of FEF neurons was done using the MG task. The delay period
separated the visual epoch (90-180 ms after target onset) from the movement epoch (80 ms
window preceding saccade onset). Visual neurons were identified if the activity was
increased in the visual epoch compared to baseline (300-100 ms preceding target onset),
movement neuron displayed higher activity in the movement epoch, and visuomovement
neurons showed increased activity in both the epochs. A visuo-motor index (VMI) was used
to validate the cell classification (Murthy et al., 2007).

$$VMI = \frac{VA - MA}{VA + MA}$$

769   where $VA = mean\ firing\ rate\ above\ baseline\ in\ the\ visual\ epoch$
         $MA = mean\ firing\ rate\ above\ baseline\ in\ the\ movement\ epoch$

With the range being from +1 to -1, visual neurons had positive VMIs, while
movement neurons had negative VMIs. Visuomovement neurons yielded intermediate VMIs.
Activity in the single saccade trials of the FOLLOW task was also taken in account for proper
cell classification, as changing task contexts have been shown to influence neuronal activity
profiles (Jagadisan and Gandhi, 2016).

Accumulator parameters:

Taking cue from accumulator models and previous studies (Woodman et al., 2008),
four main parameters were calculated: (1) Baseline firing rate; (2) Onset of firing rate
increase; (3) Threshold activity required for saccade initiation; (4) Rate of growth of activity
from onset to threshold. These measures of accumulator dynamics were calculated separately
for FOLLOW step trials in which the first saccade went into the RF (RFout) and those in
which the second saccade was towards the RF (RFin). Since correct FOLLOW step trials
always had a sequence of two saccades stepping from an RF-in position to an RF-out position

or vice versa, the activity in the RFin trials was a mix of RFout and RFin activities. To

specifically analyze activity that contributed only to the second saccades made into the RF,

the mean RF-out activity of no-step trials was used as a reference and subtracted from the

mean activity in RFin step-trials. For single neurons, the parameter calculations were made

from non-normalized, differential activity for RFin trials.

790   For FOLLOW no-step trials, trials in which the saccade was into the response field

were used for accumulator parameter calculation. Trials in each session were grouped into

fast, medium, or slow reaction time trials based on the average reaction time of that session,

i.e. trials with reaction time less than 30 ms below mean reaction time were considered as fast

trials, those with reaction time more than 30 ms above mean reaction time were slow trials,

and trials around the mean reaction time ($\pm$ 10 ms) were medium reaction time trials.

Accumulator parameters were then calculated for the three-reaction time groups.

797   Baseline activity was measured as the average of the differential activity in the RFin

condition in the 100 ms before the appearance of the first FOLLOW target. Onset was

defined as the time point when FEF activity first exceeded 2 SDs above baseline, provided

that the differential activity ultimately reached 4 SDs and was maintained above 2 SDs for at

least 50 ms for the second saccade plan and 20 ms for the first saccade plan. Threshold

activation was the average firing rate in the RF-in condition in the interval from 10 to 20 ms

before saccade initiation (Hanes and Schall, 1996). Rate of activity growth was measured by

subtracting the threshold-activity level from the onset-activity level and dividing by the time

interval between onset and threshold. This measure was robust against fluctuations in the

rise-profile. To better understand non-linear rise profiles, the rate was also measured by

piecewise regression fits using a sliding window of width 40 ms (for RFin trials) from onset

to threshold and calculating the slopes and intercepts at each point. For population analyses,

difference SDFs of each session were normalized to the peak average activity in the TSD =

17 ms group and for each saccade plan i.e. activity related to second saccade plan was

normalized with respect to TSD 17 activity for second saccades going into the RF and vice

versa for the first saccade plan. In the case of no-step trials, the SDFs were normalized to the

peak activity of the fast trials in each session.

Principal Components Analysis (PCA):

816   For analyses based on dimensionality reduction, we performed two steps of data

preprocessing before further analyses. First, we 'soft normalized' the neural responses (r) for

each neuron, i, by dividing the neural activity for each neuron (r_i) by its range ($|r_i| =$

$r_i/(range(r_i)))$ (Churchland et al., 2012). Soft normalization preserves the structure of

inter-neuronal variation while normalizing the population response so that neurons with

strong responses could be reduced to approximately unity range, but neurons with weak

responses could be reduced to less than unity range. Second, we mean-centered the responses

of each neuron by subtracting the mean activity of a given neuron across all conditions ($\overline{r_i}$)

from the neural response ($|r_i| - \overline{r_i}$).

To identify the signals that best represent the population activity of neurons, we

performed principal component analysis (PCA), a common unsupervised learning algorithm,

on the data. To do this, we constructed two matrices P_1 and P_2 of size $t \times n$ where t is the

time and n is the number of neurons with population response for the first and the second

saccade plans, respectively. We applied PCA to P_1 and P_2 yielding W_1 and W_2 respectively,

which are $n \times k$ matrix each, of principal components.

We used a metric to index the degree to which the population response occupied

different neural dimensions on trials with different TSDs. To compute this 'subspace overlap'

between the first and the second saccade plans for each TSD, we first defined the variance

captured as $V(P, W) = 1 - \frac{\|P - PWW^T\|}{\|P\|}$, where the operator $\|X\|$ means the Frobenius norm of

the matrix X. Then, the subspace overlap was given by: $\frac{V(P_2 W_1)}{V(P_2 W_2)}$.

Subspace overlap should be equal to one if the population responses occupied the

same dimensions (i.e., are spanned by the same PCs) on both the saccade plans. And the

subspace overlap should be equal to zero if the population responses occupied mutually

orthogonal dimensions on both the saccade plans.

Neural Simulations:

We simulated 40 motor neurons with 900 trials (with three types of TSD trials) using

a firing rate model (**Fig S5A**) to approximately match the statistical power of our

experimental dataset. We defined the spatial properties of each neuron (tuning curve) through

a cosine function centered on one of the 8 positions which was randomly chosen and called it

the neuron's RF:

$$tuning(\theta) = r\cos(w\theta + \phi)$$

848   where r is the peak firing rate, w defines the width of the tuning curve, θ is a set of 8 target

849   positions and ϕ is the displacement (**Fig S5B**). We then defined the temporal properties of

850   the neurons (**Fig S5C**) using a skewed Gaussian distribution response kernel:

$$kernel\ (x) = \ 2\frac{1}{\sqrt{2\pi}}\ e^{\frac{-(x-m)^2}{2c}}\ \int_{-\infty}^{\alpha x} \frac{1}{\sqrt{2\pi}}\ e^{\frac{-t^2}{2c}}\ dt$$

$$= \frac{1}{\sqrt{2\pi}}\ e^{\frac{-x^2}{2c}}\ \left[1 + erf\left(\frac{\alpha x}{\sqrt{2}}\right)\right]$$

where $c = 8000$ controls the full width at half maximum of the distribution, $erf(x)$ is the

error function and α is the shape parameter that controls the shape of the distribution. To

simulate a noisy accumulator (**Fig S5H**), we sampled $m \sim \mathcal{N}(0,40)$ and $\alpha \sim -$

$|\mathcal{N}(0,0.01)|$. Note that similar results can be obtained using a response kernel resembling a

Poisson post-synaptic potential function:

$$kernerl\ (x) = \ \left[1 - e^{\frac{-x}{\tau_g}}\right] \times \left[e^{\frac{-x}{\tau_d}}\right]$$

where τ_g controls the rate of increase and τ_d controls the rate of decay.

We further injected a small noise to the system by convolving the response kernel with

Gaussian noise of $\mu = 0$ and $\sigma^2 = 1$. For each neuron, we multiplied the tuning curve and

the convolved kernel to get the spatiotemporal firing rate for that neuron (**Fig S5D**). Saccade

onsets were taken as the time when the normalized activity reached a fixed threshold of 1

unit.

For the simulated data with asymmetric bilateral inhibition, we followed the above

steps (for both plans 1 and 2) until the activity for plan 1 reached y% of its peak response.

This 'y' is given by an inhibition function (**Fig S5G**) which was constructed by dividing the

neural response by the response in no-step trials. To simulate the inhibitory effect of plan 1

on plan 2, we temporally shifted the response for plan 2, to until after the time of first saccade

onset (t_{s1}), after plan 1 reached y% of its peak response and then interpolated the data in

between using a cubic spline function.

$$r_2(t) = \begin{cases} r_2(t), & \text{if } t < T \\ r_2(t + (t_{s1} - T)), & \text{if } t \geq t_{s1} \end{cases}$$

878   where $T = arg\left(y \times \max(r_1(t))\right)$

880   To simulate the effect of plan 2 on plan 1, we first normalized plan 2's response from
881   0 to 1, then multiplied it by the same inhibition factor, y and then subtracted it from plan 1's
882   response.

$$r_1(t) = r_1(t) - y\left(\frac{r_2(t) - \min(r_2(t))}{\max(r_2(t)) - \min(r_2(t))}\right)$$

Therefore, the nature of inhibition was asymmetric and, in this way, plans 1 and 2
have the highest bilateral inhibition effect on each other for the shortest TSD and the least
effect for the longest TSD. We estimated all the above hyper-parameters such that the
simulated data closely resembled the experimental data.

For the simulated data with unilateral inhibition, we followed the same steps as above
with the exception of the last step. That is, we modeled the effect of plan 1 on plan 2 by
temporally shifting the plan 2 as described above but we did not account for the effect of plan
2 on plan 1.

After simulating the data, we followed the same data pre-processing step before
dimensionality reduction similar to the experimental data.

Statistical testing:

A two-sided Wilcoxon signed-rank test to analyze a single sample set of data. For
group comparisons, the non-parametric Kruskal-Wallis test was used. Trials were considered
to be independent observations as the TSDs on each trial was chosen randomly. All the
results are presented as mean ($\pm$ standard error of mean, SEM) and all tests are performed at a
significance level of $\alpha = 0.05$ unless otherwise mentioned.

Data and code availability:

All data is available in the main text or the supplementary materials. Raw data and codes are
available upon reasonable request.
905

906   **Acknowledgments:**
907   We thank S. Sengupta for helping with behavioral training and Dr. A. Gopal P.A. for helping
908   with data collection.
909
910   **Funding:**
911   This work was supported by a D.B.T.-I.I.Sc (Department of Biotechnology, Government of
912   India – Indian Institute of Science) partnership grant given to A.M. D.B was supported by a
913   graduate fellowship from the Ministry of Human Resource Development (MHRD),
914   Government of India, through the Indian Institute of Science.
915
916   **Author Contributions:** D.B. Performed research, Analyzed data, Wrote the paper. N.S.
917   Analyzed data, Wrote the paper. A.M. Designed research, Wrote the paper.
918
919   **Competing interests:** Authors declare no competing interests.
920
921   **Disclosure:** The authors declare no competing financial interests
922

**Figure 1. Behavioral predictions for processing bottlenecks during the planning of
sequential saccades**

A. Single-channel bottleneck framework. Each task is made up of three stages. The visual stage (V) can be
carried on in parallel with stages of another task, but the central planning stage, P, can only proceed singly. In a
two-saccade sequence, the stages of the first saccade plan proceed to completion unabated leading to its
execution (E). For the second plan however, if the second target closely follows the first (low TSD condition),
the central planning stage, P2, is postponed till P1 is complete. Such a postponement does not occur in the long
TSD condition, where the two saccade plans are well-separated, thereby leading to an increase of RT2 from long
to short TSD.

B. Capacity-sharing bottleneck framework. In this framework, the P stages of multiple plans can proceed in
parallel and access the brain's limited processing capacity simultaneously. In the low TSD condition, P1 and P2
concurrently 'share' the capacity, resulting in slower progress of the saccade plans. This leads to lengthening of
both RT1 and RT2 in the low TSD condition, the effect on RT2 being greater as the second saccade plan gets a
smaller share of the central capacity.

C. Predictions of reaction time vs TSD for single-channel bottleneck framework (left) and capacity-sharing
bottleneck framework (right). RT2 increases with decrease in TSD for both frameworks, whereas RT1 increase
is predicted only by the capacity-sharing model.

D. Behavioral data for reaction time vs TSD. Data shows trials in which the first (for RT1) or second (for RT2)
saccade was into the response field. Both reaction times increased significantly with decrease in TSD.

Figure 2. Neural activity predictions for processing bottlenecks during the planning of sequential saccades

A. Hypothesized neural activity for saccade plan 1 in the single-channel bottleneck framework: Bottom: After the first visual target is presented (vertical broken line; T1), there is an initial visual processing stage (V1) which is often of constant duration for all plans. The planning stage (P1) for the first saccade is represented as a noisy integrator, accumulating activity till the motor threshold (horizontal solid line) is reached and the saccade is executed (E1). Top panel: The corresponding neural activity is shown as the ramping up of FEF movement neuron activity till saccade onset (S1). The activities corresponding to three different TSDs are shown in three different colors.

B. Hypothesized neural activity for saccade plan 2 in the single-channel bottleneck framework. The onset of the accumulation process and the ramping up of the neural activity will shift later with decrease in TSD to account for RT2 elongation (same format as A; T2: onset of second target; V2: visual processing stage for target 2, P2: planning stage for saccade 2; E2: execution stage for plan 2; S2: onset of second saccade).

C. Hypothesized neural activity for saccade plan 1 in the capacity-sharing bottleneck framework. The onsets of the integrators and the movement neuron activity do not change with TSD on account of parallel programming of the two saccade plans. Increase in saccade latencies at shorter TSDs maybe brought about by a decrease in the growth rate from long to short TSD. Same format as **A**.

D. Hypothesized neural activity for saccade plan 2 in the capacity-sharing bottleneck framework. Same as C, with the addition of threshold modulation and a greater degree of rate adjustment with TSD to constitute the larger increase in RT2 from long to short TSD. Same format as **A**.

Figure 3. Activity for FEF movement neurons during sequential saccades

A. Top: Population activity of FEF movement neurons when the second saccade went into the response field, aligned on second target onset (T2). Mean saccade onset times (S2) for the short, medium, and long TSD conditions are shown as vertical, colored lines with s.e.m error bars. Right: same as left but activity aligned to the second saccade onset. Shading indicates mean ± SEM.

B. Population histogram of slopes of each measure of accumulator dynamics (baseline, onset, growth rate and threshold) as a function of TSD for FEF movement neurons. Asterisks denote cases where the distribution of movement neuron slopes was significantly different from zero (Wilcoxon signed-rank test, *** $p < .001$, ** $p < .01$).

C. Same as **A** but for saccade 1

D. Same as **B** but for saccade 1

1025
1026

Figure 4: Extent of subspace sharing explains processing bottlenecks during the planning of sequential saccades

A. Normalized mean population neural responses aligned to target 1 (T1) and target 2 (T2) for short, medium and long TSD trials (n.u. = normalized unit); same as **Fig 3A** and **3C.**

B. Cumulative percent variance explained by the first 10 PCs for target (T1) (black) and target 2 (T2) (color) related responses for short (left), medium (center) and long (right) TSD trials.

C. Subspace overlap between a pair of conditions. S, M and L indicate short, medium and long TSDs, and 1 and 2 indicate the saccade plan number.

1056

Figure 5: Only simulations with bilateral, asymmetric inhibition capture the empirical data's population dynamics

A. Schematic of the simulation with bilateral inhibition. Top row: simulated neural activity for the first saccade plan and bottom row: simulated neural activity for the second saccade plan (see **Fig S6** and methods; n.u. = normalized unit).

B. Normalized mean population neural responses, for data simulated with bilateral inhibition, aligned to target 1 and target 2 for short, medium and long TSD trials.

C. First three PCs plotted against each other for target 1 (black) and target 2 (color) related responses for short (left), medium (center) and long (right) TSD trials. Filled circle markers indicate the starts of the respective trajectories.

D. Subspace overlap between a pair of conditions. S, M and L indicate short, medium and long TSDs, and 1 and 2 indicate the plan number.

E. Same as **A,** but for simulation with unilateral inhibition (see methods).

F. Same as **B**, but for simulated data with unilateral inhibition.

G. Same as **C**, but for simulated data with unilateral inhibition.

H. Same as **D**, but for simulated data with unilateral inhibition.

Figure 6. Processing bottlenecks in FEF V and VM neurons

A. A representative FEF visual neuron aligned to target onset (T) and saccade onset (S) in a memory-guided task for saccades into the RF (yellow) and saccades out of RF (into aRF; black).

B. A representative FEF vismov neuron aligned to target onset (T) and saccade onset (S) in a memory-guided task for saccades into the RF (yellow) and saccades out of RF (into aRF; black).

C. FEF visual and vismov neuron population activity encoding for different TSD conditions (short, medium, long), aligned to target 2 onset. Mean saccade onset times for the TSD conditions are shown as vertical colored lines with s.e.m error bars. Inset: Kruskal–Wallis box plots for average activity in the visual epoch (gray shaded area) for the three TSDs. Shading indicates mean ± SEM.

D. Same as **C** but for neural activity aligned to target 1.

1119
1120   **Figure S1. Schematic of the FOLLOW task** (*related to Fig 1*)
1121
1122   **A.** A representative no-step trial. The trial starts with the appearance of a central fixation point (FP), followed by
1123   the presentation of the green target (T1) at any one of the six possible peripheral locations. The monkey had to
1124   make a single saccade (S1) to the target to get a juice reward. In the representative framework, the processes
1125   leading to the culmination of a saccade are simplified to consist of three stages: visual encoding of stimuli(V),
1126   central planning (P), and saccade execution (E). RT refers to the reaction time.
1127
1128   **B.** A representative step trial. Similar to no-step trials, a step trial started with central fixation, after which a
1129   green target (T1) appeared. A second red target (T2) was then presented after T1. A variable delay separated the
1130   first and the second target onsets (target step delay; TSD). The monkey had to make a sequence of two saccades
1131   (S1, S2) to the two targets in order of their appearance to get rewarded in step trials. The abbreviations used are
1132   the same as in **A**, but for two saccades.
1133

Figure S2. Population activity of FEF movement neurons in no-step trials (*related to Fig 3*)

A. Schematic of possible adjustments of neural activity related to processing bottlenecks. The lengthening of the reaction time may be due to four possible adjustments according to accumulator dynamics. Each schematic shows a noise-free, simplistic accumulation process, starting from the baseline and reaching up to the threshold for saccade initiation. Representative reaction time distributions are plotted above each of the schematics. Lowering of baseline activity, delaying the onset of activity, slowing down the rate of growth, and increasing the threshold level can either singly or in combination, bring about increased reaction times.

B. Left: FEF movement neuron population activity encoding for different reaction times (short: green, medium: gray, long: red), aligned to target onset (vertical broken line) . Mean saccade onset times for all the reaction time conditions are shown as vertical colored lines with s.e.m error bars. Right: same as left but activity aligned to saccade onset (vertical broken line). Shading indicates mean ± SEM.

C. Illustration of the measurement of accumulator parameter: baseline. Left: Schematic of a spike density function of a representative neuron aligned to target onset showing the baseline value. Middle: The baseline was calculated for short, medium and long reaction times for a representative neuron and the slope of the best fit was calculated. Right: Histogram of distribution of slopes measured this way for all the neurons was compared with zero (vertical broken line).

D. Same as **C** but for measurement of onset.

E. Same as **C** but for measurement of growth rate.

F. Same as **C** but for measurement of threshold.

Figure S3: Single neuron example for processing bottlenecks *(related to Fig 3)*

A. A representative FEF movement neuron aligned to target onset (T) and saccade onset (S) in a memory guided task for saccades into the RF and saccades out of RF.
B. Left: Neural activity, a representative neuron encoding for different TSD conditions (short, medium, long TSD), aligned to target 2 onset. Right: same as left but activity aligned to S2 onset.
C. Same as **B** but for neural activity aligned to target 1 (left) and saccade 1 (right).

Figure S4: Population dynamics for all target positions (*related to Fig 4*)

A. Soft-normalized mean population neural responses towards all six target positions (3 towards RF, shown in warm colors and 3 out of RF, shown in cold colors; see inset for colors), aligned to target 1 (left) and target 2 (right) for short TSD trials (n.u. = normalized unit).
B. Same as **A**, but for medium TSD trials.
C. Same as **A**, but for long TSD trials.
D. First three PCs for each target position shown in **A**, plotted against each other for target 1 (left) and target 2 (right) related responses for short TSD trials. Filled circle markers indicate the starts of the respective trajectories.
E. Same as **D**, but for medium TSD trials shown in **B**.
F. Same as **D**, but for long TSD trials shown in **C**.

1216

Figure S5: Saccade kinematics with different TSDs (*related to Fig 4*)

A. Saccade 1 trajectories, towards RF (filled, broken circle), during short, medium and long TSD trials from a representative session. The central cross denotes the fixation point.

B. Mean saccade 1 velocity profiles for short, medium and long TSD trials (thick lines) superimposed on saccade 1 velocity profiles from individual trials (thin lines). Shading indicates mean ± SEM.

C. Peak saccade 1 velocities from each session for short, medium and long TSD trials. P values: short-medium: 0.81, ranksum test; medium-long: 0.60, ranksum test.

D. Saccade 1 main sequence for short (left), medium (center) and long (right) TSD trials. The overlaid contours represent the density of data.

E. Saccade 2 trajectories, towards RF (filled, broken circle), during short, medium and long TSD trials from the same representative session as **A**.

F. Same as **B**, but for saccade 2.

G. Same as **C**, but for saccade 2. P values: short-medium: 0.92, ranksum test; medium-long: 0.80, ranksum test.

H. Same as **D**, but for saccade 2.

1239

Figure S6: Data simulation process, reaction time and neural activity profiles (*related to Fig 5*)

- **A.** Schematic illustration of the simulation.
- **B.** Tuning curves (peak firing rates as a function of target positions) for all the simulated neurons. One representative neuron's tuning curve is highlighted in purple.
- **C.** Response kernels (activity as a function of time) for all the simulated neurons. Same representative neuron's stimulus kernel is highlighted in purple.
- **D.** Multiplication of tuning curves and stimulus kernels and addition of noise results in the simulated neuron's activity for each of the 8 target positions. Activity towards the 8 target positions is shown for the same representative neuron as above.
- **E.** Distribution of RF positions (argmax(tuning curve)) of the simulated neurons shows a uniform spread of RFs across the simulated neurons.
- **F.** Distribution of peak firing rate (max(tuning curve)) of the simulated neurons shows a unimodal distribution with a mean ~ 37 sp/s.
- **G.** Probability of inhibition as a function of TSD
- **H.** Bottom: Example simulation for one session, for RF position, following the simulation pipeline shown in A. Top: reaction time distribution obtained from the simulation.
- **I.** Top: The same reaction time distribution from H, but uniformly divided into fast, medium and slow reaction times. Bottom: Average neural activity from the simulation in H, divided into fast, medium and slow reaction time conditions as explained above.
- **J.** Same FEF neural data from **Fig S2B** (divided into fast, medium and slow reaction time).
- **K.** Top: reaction time distributions of two monkeys for short, medium and long TSD trials for plan 1 (left) and plan 2 (right). Bottom: Same as **Fig 3C;** right: Same as **Fig 3A.**
- **L.** Top: reaction time distributions from the simulations with bilateral inhibition in the same format as above. Bottom: Same as **Fig 5F.**
- **M.** Top: reaction time distributions from the simulations with unilateral inhibition in the same format as above. Bottom: Same as **Fig 5B.**