Social network architecture and the tempo of cumulative cultural

evolution

Mauricio Cantor[1,2,†,*], Michael Chimento[3,6,7,†], Simeon Q. Smeele[3,4,†], Peng He[5,6,7,8], Danai

Papageorgiou[5,6,7,8], Lucy M. Aplin[3,6,^,*], Damien R. Farine[5,6,7,8,^]

[1] Department for the Ecology of Animal Societies, Max Planck Institute of Animal Behavior, Konstanz,

Germany

[2] Departamento de Ecologia e Zoologia, Universidade Federal de Santa Catarina, Florianópolis, Brazil

[3] Cognitive & Cultural Ecology Research Group, Max Planck Institute of Animal Behavior, Radolfzell,

Germany

[4] Department of Human Behavior, Ecology and Culture, Max Planck Institute for Evolutionary

Anthropology, Leipzig, Germany

[5] Department of Collective Behaviour, Max Planck Institute of Animal Behavior, Konstanz, Germany

[6] Department of Biology, University of Konstanz, Konstanz, Germany

[7] Centre for the Advanced Study of Collective Behaviour, University of Konstanz, Konstanz, Germany

[8] Department of Evolutionary Biology and Environmental Science, University of Zurich, Switzerland

^Joint senior authors; †Joint first authors, arranged alphabetically

* Corresponding Authors:

Mauricio Cantor. Department for the Ecology of Animal Societies, Max Planck Institute of Animal

Behavior, Am Obstberg 1, Radolfzell 78315, Germany; Email**:** mcantor@ab.mpg.de

Lucy M. Aplin. Cognitive & Cultural Ecology Research Group, Max Planck Institute of Animal

Behavior, Am Obstberg 1, Radolfzell 78315, Germany, Phone: +49 7732 1501-13.; Email:

laplin@ab.mpg.de

Abstract

The ability to build upon previous knowledge—cumulative cultural evolution—is a hallmark of human societies. While cumulative cultural evolution depends on the interaction between social systems, cognition and the environment, there is increasing evidence that cumulative cultural evolution is facilitated by larger and more structured societies. However, such effects may be interlinked with patterns of social wiring, thus the relative importance of social network architecture as an additional factor shaping cumulative cultural evolution remains unclear. By simulating innovation and diffusion of cultural traits in populations with stereotyped social structures, we disentangle the relative contributions of network architecture from those of population size and connectivity. We demonstrate that while more structured networks, such as those found in multilevel societies, can promote the recombination of cultural traits into high-value products, they also hinder spread and make products more likely to go extinct. We find that transmission mechanisms are therefore critical in determining the outcomes of cumulative cultural evolution. Our results highlight the complex interaction between population size, structure and transmission mechanisms, with important implications for future research.

Keywords

Cultural evolution | Cultural complexity | Multilevel societies | Small-world networks | Social structure

Background

Cumulative cultural evolution (CCE)—where iterative innovations and social transmission generate cultural accumulation over time [1-3]—is key to the human's ecological success and worldwide distribution [4,5]. While CCE fundamentally depends on the interplay between cognition and social learning mechanisms [1], it is increasingly clear that demography can modulate the rate of cultural evolution [6-9]. Large population sizes [10,11], greater population turnover, and more densely

connected societies [3,12] can all provide greater innovative potential, more learning models, faster

diffusion, and reduced extinction risk of useful innovations [7,13-15]. For example, increasing

population density as well as migration of hunter-gatherers during the upper Paleolithic transition led

to the explosion of culture that forms the basis of modern human societies [8,15]. Yet, it remains

unclear how variation in the wiring of these social connections shape the tempo of cumulative cultural

evolution.

Network architecture—here defined as a social structure with a characteristic set of properties—can

shape transmission of behaviours, thus setting the tempo of CCE—here defined as the rate of cultural

recombination events. For example, architectures with low network connectivity (i.e. density;

proportion of realised connections), high clustering (tendency of connected individuals to share the

same social neighbours) and high modularity (tendency of the network to contain sets of individuals

more connected to each other than with others) slow down the spread of information across

populations [16-18]. Slower spread can then potentially favour greater cultural diversity by allowing

multiple cultural lineages to arise in populations before any one lineage dominates [19,20]. While

previous work has largely focused on how new behaviours spread through a social network [16,17] to

establish cultures [20,21] and how cultural traits can generate a feedback shaping network structure

[18,22], more recently it has been argued that emergent network properties could affect CCE [12,14,19]

by shaping how new traits are produced, recombined, and maintained [14]. For example, partial

connectivity facilitates the emergence of multiple cultural lineages in parallel [20], which is required

for achieving cultural accumulation, but partially-connected networks suffer from cultural loss if

connectivity is too low for new innovations to spread [14]. By contrast, full connectivity facilitates

rapid spread of new innovations, but can prevent the accumulation of alternative cultural traits

[12,14]. However, within a given level of connectivity, how connections are structured—the social

network architecture—could also impact CCE by influencing how fast and widely information can

spread.

Because network architecture can shape the effect of connectivity on diffusion dynamics [23], those

architectures that balance the ability for cultural accumulation together with the recombination of

different cultural traits should have a selective advantage in facilitating CCE [19]. Multilevel societies,

such as those in modern hunter-gatherers, feature high clustering and nested modularity. These

network properties are expected to favour CCE by allowing coexistence of multiple cultural traits in

different parts of the network, and for different cultural lineages to come into contact to allow

combinations from lineages to produce new traits [19]. Multilevel societies have been demonstrated to

accelerate CCE when compared to fully connected networks [19]. However, when considering their

potential for facilitating CCE, multilevel and fully-connected networks represent possible endpoints

along a continuum of possible architectures. Here, we ask how a range of social network architectures

can affect the tempo of CCE within a given population size and number, or density, of social

connections within that population. Our approach allows us to explicitly disentangle the relative

contribution of network architecture from those of connectivity and population size.

Materials and Methods

Overview

We first generated social networks with six different architectures—random, small-world, lattice,

modular, modular lattice, and multilevel—capturing different levels and combinations of clustering

and modularity (**Fig. 1A**). We expressed these network architectures in populations with different

sizes and densities of connections (average degree), where all individuals in the network had the same

degree. We then build two agent-based models to explore how network architecture affects

cumulative culture evolution. Briefly, our models allow innovations of cultural products to take place

along two cultural lineages, with the knowledge of new products being spread through social

connections via two transmission mechanics: either one-to-many or one-to-one diffusion. Once a high

level of product diversity has been reached in both lineages, agents can recombine each lineage's

products into one with a final higher payoff product (hereafter 'recombination'). Finally, we compared

the performance of agents arranged in the different network architecture in terms of time to cultural

recombination (i.e. tempo), time to diffusion, and the diversity of cultural traits.

Social network architectures

We generated networks in which nodes representing individuals were linked by binary social

relationships to represent the following six stereotypical social structures. We generated: (i) small-

world networks, using the Watts-Strogatz model [23] with node degree K links; (ii) random networks,

by randomly connecting nodes ensuring all nodes had the same degree K; (iii) lattices, by placing

nodes on a grid and connecting each to its K nearest neighbours; (iv) modular networks, by assigning

nodes into nine modules, randomly connecting each to $K-1$ nodes from the same module and one node

from another module; (v) modular lattices, as per modular networks, but where the connections

within modules were lattices; and (vi) multilevel networks, as per modular lattices, but assigning

nodes into three sets of three modules, and connecting each to $K-2$ nodes within their module, one

node from each module from within its set and one node from a module outside of it.

Each network architecture differs in clustering and modularity, but we standardize their connectivity

(degree) and population size. For each architecture, we generated an ensemble of networks with

different sizes ($N \in \{64,144,324\}$ nodes), and densities of connections (in which $K \in \{8,12,18,24,30\}$

average links). We used these population sizes because they allowed us to partition the network into

equally-sized clusters composed of equally-sized groups in which all individuals had the same degree,

and in which connectivity was greater within groups and within clusters than between groups and

between clusters. Although all networks had comparable sizes and densities (i.e. the proportion links),

the six architectures varied in levels and combinations of clustering coefficient and modularity.

Clustering, C, informs the tendency of connected nodes to share the same connections with other

nodes, while modularity, Q, informs the tendency of the nodes to be organized into cohesive subsets

that are more connected to each other than to the rest of the network [24].

Cultural evolution simulations

We constructed two agent-based models to simulate cultural evolution on the different types network

architectures. The two models differed in the mechanics of information transmission: one-to-many

versus one-to-one diffusion pathways. Our first agent-based model (model 1) followed Migliano et al.

[19]. All agents were initialized with an inventory of three items from each of two lineages. In each

simulation round (epoch), each focal agent was selected once, at random, and a partner randomly

chosen from its social network connections. These agents combined one or two items from their

inventory in proportion to their value into a triad of items. If this triad was a valid product, knowledge

of that product was learned, spread immediately to all their network connections (one-to-many

diffusion), and subsequently became available as an ingredient for making new products. Simulations

finished once a recombination product (a triad that recombines specific products from both lineages)

was first innovated. We ran 5,000 simulations for each of the network architecture types, sizes and

densities of connections, recording time to achieve the recombination product (in epochs) and tracking

the diversity of cultural innovations over time. An epoch was one simulation round in which each

agent was selected once as a focal agent in random order.

Our second agent-based model (model 2) extended the first by changing the transmission mechanic

and altering the set of valid combinations such that the model can run past the first innovation of

either recombination product. Transmission of valid products now occurred between dyads of agents

(one-to-one diffusion) prior to choosing items from their inventory, in contrast to the broadcast style of

diffusion in model 1. Secondly, if a triad contained either recombination products, the final product

was that recombination product. In the case where both recombination products were present in the

triad, one was chosen as the final product at random. This allowed us to track the diffusion of

recombination products beyond their innovation. We also ran 5,000 simulations for the same

parameter space of model 1, recording time (in epochs) to cultural lineage recombination, as well as

time to diffusion to the majority of the network (i.e. the latency to more than $(N/2) + 1$ nodes having

a high value product).

Data analyses

We compared the performance of agents organized in the different social network architectures in

terms of the time to recombination, time to diffusion, and the diversity of cultural lineages over time.

To compare time to recombination, we used time-to-event (survival) analyses [25] where time to

recombination was a function of network architecture and connectivity. Our simulations yielded time-

to-event data, so we used the following methods from survival analysis, which are suitable for such

data. For each population size, we used the Cumulative Incidence Function to estimate the proportion

of simulations in which agents reached the recombination of each cultural lineage's products into a

final high-payoff product (the 'event'). We used the non-parametric Kaplan-Meier product limit

estimator to estimate the 'survival function' from this time-to-event data; since we represented the

time intervals based on observed recombination events from 5,000 simulations from model 1, we also

calculated the 95% confidence intervals (using the Greenwood estimator). To measure variance in time

to recombination across population sizes, we measured the quartile coefficient of dispersion

(QCD=(Q3-Q1)/(Q3+Q1)), as this variable is not normally distributed and QCD offers a more robust

measure.

While statistics are not typically performed on data from agent-based models since the posterior is

directly sampled, we wanted to quantify the relative contributions of architecture, size and

connectivity without the cumbersome descriptions of the entire distribution (distributions can be

readily seen in **Fig. 2**). For both models 1 and 2, we created three sets of generalized linear models

(GLMs) that predicted logged time to recombination (in epochs). Time to recombination was logged to

account for non-normality of residuals, and to make comparisons more fair by bringing the mean

closer to the median of the distribution. All models used log link function, as the data was non-linear

conditional on predictors, even after the log transformation. Also, the log link function allowed the

presentation of exponentiated coefficients, which simplify the comparison to the reference (here, the

random networks at the GLM intercept). The first set of GLMs used a full interaction structure to

partition the relative contributions of architecture, size and connectivity to average time to

recombination (Supplementary Material, **Table S1**), excluding fully connected networks. To then

compare fully connected networks to all other networks, we built a GLM using architecture and

population size as predictors in a full interaction structure (Supplementary Material, **Table S2**).

Connectivity was excluded as a predictor, as all fully connected networks only have 1 possible degree

($K=N-1$). Finally, to compare differences between architectures more precisely, we subset data by

connectivity and population size and performed a GLM with only architecture as a predictor for each

subset, again excluding fully connected networks. We performed all data analyses in R [26], using

'survival' [27] and 'survminer' [28] packages.

Data availability

We implemented the agent-based models in R and Python. The code to generate the social networks,

perform the simulations, replicate the data, the statistical analyses and the figures are available at [29].

The simulated data used in this manuscript can be retrieved at [45].

Results

When comparing the time to cultural recombination across population sizes and levels of connectivity,

partially connected networks consistently outperform fully connected networks [12,14,19]. A

generalized linear model indicated that fully connected networks were, on average, 65% slower (GLM,

$\exp(\beta)$=1.652, t=59.208, p<.001; **Table S1**) compared to the least structured network architecture of the

same size (random, N=64 taken as the intercept), with similar decreases in performance independent

of size. Further, we also confirm [7] that larger populations take less time (about 40% less) to reach

recombination (GLM, $\exp(\beta)$=0.618, t=-68.481, p<.001) compared to networks of the same architecture

and connectivity (**Figs. 1B,2, Table S2**). Larger partially-connected network architectures were less

variable in their time to recombination (Quartile Coefficient of Dispersion: QCD=0.688 for N=64;

QCD=0.444 for N=144; QCD=0.273 for N=324, **Fig. 2A**). We also found that time to recombination was

optimized at intermediate densities of connections, confirming that intermediate levels of connectivity

can favour CCE [14], and revealing that the optimal level of connectivity varied with population size

(**Fig. 1B**). In the smallest population (N=64), sparse networks outperformed the others, but this was

reversed in the largest population (N=328) (**Fig. 1B**). However, differences in time to recombination

were generally small (**Fig. 1B, 2A**).

Architectures favouring cumulative cultural evolution under some conditions disfavour it under other

conditions

Under the one-to-many diffusion mechanism (model 1), multilevel, modular, and modular lattice

architectures had relatively shorter times to recombination in smaller populations with greater

connectivity and in larger populations with less connectivity (**Fig. 3A**). However, such network

architectures performed worse than lattice, small world, and random architectures in smaller

populations with less connectivity and in larger populations with greater connectivity (**Fig. 3A**).

Multilevel, modular, and modular lattice architectures were optimal at lower and higher levels of

connectivity in medium-sized populations (N=144, **Fig. 3A**), although connectivity generally had a

lesser impact for these architectures relative to random, small-world, and lattice architectures (**Fig. 1B**).

Surprisingly, multilevel performed the worst in seven out of the 11 size and connectivity combinations

(**Fig. 3A**) despite having the highest clustering and modularity—properties that have been predicted

to favour CCE [19]. The modular lattice architecture (which had similar modularity and clustering to

multilevel architecture) performed best in the other four combinations (**Fig. 3A**). Thus, no one network

architecture proved optimal, with those favoured under some conditions being disfavoured under

other conditions.

The large variation in the time to recombination (**Fig. 2**) within a given combination of network

architecture, population size, and density of connections suggests that the outcomes of a simulation

were predominantly driven by stochastic events. The impact of such stochasticity is best revealed by

the bimodal outcome for partially connected networks, which arises most often in smaller populations

(**Fig. 2A**). This bimodality occurs because there are fewer independent innovation events when there

are fewer individuals, which increases the chance that cultural products all emerge from the same

lineage and, therefore, that this single lineage spreads to the whole population before the other lineage

is innovated. Tracking the diversity of products over time (**Figs. 4, S1**) highlights how the stochasticity

in early events can affect cultural diversity, and therefore the outcomes of CCE, even within the same

network architecture. Overall, measuring the tempo of CCE under one-to-many diffusion (model 1)

revealed differences in the best performing architecture across population sizes and levels of

connectivity (**Figs. 2A, 3A**); however, these between-architecture differences were small (range =

0.886-1.145; **Fig. 3A**), compared to the large variance in the time to recombination within architecture

(**Fig. 2A**).

The same architectures that promote diversity also restrict transmissions of novel products

To identify the relative contribution of diffusion mechanisms to CCE, we extended model 1 by

implementing a one-to-one diffusion mechanism (model 2). Whereas model 1 represents an extreme

scenario where information spreads instantaneously to all the contacts of a focal agent, model 2 tests

another extreme in which information about discoveries spread only to a single contact at a time.

Interestingly, when employing such one-to-one diffusion dynamics, fully connected networks were

only estimated to be 3% slower to recombination compared to networks of the same population size

(GLM, $\exp(\beta)$=1.033, t=6.808, p<.001; **Table S1**). Again, larger populations had a significantly shorter

average time to recombination compared to smaller networks of the same architecture and degree

(GLM, exp(β)=0.595, t=-109.094, p<.001; **Fig. 2B, Table S2**). However, under one-to-one diffusion, the

relative times to recombination of different architectures was generally more consistent than under

one-to-many diffusion, both in their median times to recombination (**Fig. 2B**) and in their relative

performance under a given population size and level of connectivity (**Fig. 3B**). Within a given

population size, multilevel architecture typically had the shortest times to recombination when

networks had greater connectivity, but there was almost no difference in performance among

architectures when connectivity was low (**Fig. 3B**). Thus, in contrast to one-to-many, one-to-one

diffusion increased the tempo for architectures with greater modularity and clustering (modular,

modular lattice, multilevel) relative to the other architectures.

Model 2 also tracked the time for the recombination product to diffuse to the majority of the

population, something which model 1 was not designed to track. The time from recombination to

diffusion was shortest in fully connected networks, and increased with population size (**Fig. 2C**).

When evaluating performance from the start of the simulations until the time to diffusion, population

size caused the most variation in outcomes, compared to network architecture or connectivity (**Table

S2**). In small populations, the contribution of the final diffusion was relatively small compared to the

time to recombination, meaning that the best performing networks in achieving recombination also

performed best overall (**Fig. 3C**). By contrast, in larger populations, the performance of modular and

clustered network architectures (modular, modular lattice, and multilevel) all performed worse: they

were the slowest at reaching final diffusion (**Fig. 3C**) despite typically reaching recombination the

fastest (**Fig. 3B**). These differences, however, remain minor relative to the variance in outcomes within

each set of conditions (architecture, population size and connectivity).

Discussion

We revisit recent empirical and *in silico* experiments in humans to tease apart the contributions of

different candidate social structures to the tempo of cumulative cultural evolution. Our results suggest

that it is unlikely that one specific social network architecture consistently promotes cumulative

cultural evolution across all population sizes, densities of social connections, or diffusion mechanisms.

Rather, the relationship is nuanced; the broad distribution of outcomes from our two models indicate

that the best performing architecture under some conditions can be the worst performing architecture

under others. Further, the outcome of any diffusion mechanism is as likely to be affected by stochastic

processes as by the architecture of the networks itself. While not at odds with previous work showing

that multilevel societies can accelerate cumulative cultural evolution [19], our results suggest that a

range of other partially connected architectures could equally increase the tempo of cumulative

cultural evolution.

The fact that alternative architectures can have similar outcomes in terms of CCE has important

consequences for how the social structure of societies and CCE are framed in future discussions, and

where future research is directed. Current thinking is that complex, highly structured societies, such as

multilevel societies, might precede recombinatory CCE in the timeline of human evolution, or that the

benefits accrued from cultural evolution [22] or CCE [19] might co-evolve with clustered and modular

network structures. However, our results suggest that simple patterns of spatial distribution (e.g. a

lattice social network caused by distributed resources) could lead to largely equivalent effects on CCE.

It follows that we might expect to find recombinatory CCE even before the evolution of complex

societies. Indeed, evidence that simple, lattice-like social structures [30] can provide a substrate for

recombinatorial culture might be provided by the combinatorial, spatially variable song structure of

territorial passerine birds [31-35], which several authors have proposed to be a simple form of CCE

[36,37].

Population size has been suggested as another major demographic factor affecting rates of CCE [3,7-9].

Our findings align with this previous research, with our simulations showing that larger populations

always have a higher rate of cultural accumulation. Population size also interacted with connectivity

(which we modelled as a fixed network degree, i.e. the number of individuals' social connections [22]),

with changes in connectivity having a more pronounced effect in smaller populations. This outcome is

likely to arise because an increase in one unit of mean degree corresponds to a greater increase in

network connectivity in smaller populations (more rapidly pushing the network towards becoming

fully connected). However, in our simulations we did not vary the distribution in connectivity among

individuals, which has previously been shown to impact the properties of information cascades [38]

and differences among groups in behaviours such as cooperation [39]. Skewed degree distributions,

where some nodes are much more connected than others, could allow independent lineages to arise in

peripheral nodes and for highly connected 'hubs' to combine the products from these lineages, thereby

facilitating CCE. Thus, variation in how much or how little individuals are connected, independently

of other factors (mean connectivity, population size, and network properties), is an important

dimension for future studies on CCE to consider.

Fully connected networks have been commonly used to evaluate the performance of a transmission

network with a given set of characteristics [3,19]. However, a fully-connected social network

representing a human population of any reasonable size would represent an unrealistically high level

of connectivity [40], even in the fractal-like human social networks [41]. In addition, our simulations

demonstrate that the contribution of large differences in connectivity to the rates of CCE outweighs

any effects of network architecture, at least when information is broadcast (i.e. a one-to-many diffusion

mechanism). Thus, we suggest that fully connected networks are uninformative null models for

testing the influence of social structure on the tempo of CCE. Instead, random networks of similar

sizes and densities of connections as the network of interest would provide a more robust benchmark

[see also 42]. Our results suggest, however, that any effect of network architecture on increased rate of

CCE inferred from noisy field data would likely be indistinguishable from the null expectancy, as

variation within architecture greatly exceeded that of between architectures.

One major remaining question is how the diversity of cultural lineages might affect CCE across

different network architectures. Our simulations were built on previous empirical and theoretical

work that considered two cultural lineages [12,19]. Increasing the number of lineages could potentially

reduce or increase the time to recombination, depending on how the recombinations have to be made

and on their pay-offs. For example, recombinations from any two of three or more lineages would

likely emerge faster, as the probability that any two lineages survive is higher. By contrast,

recombinations requiring products from three lineages could take substantially longer, especially in

more connected networks (or with one-to-many diffusion dynamics), since all three would have to

become established. However, adding more lineages would be unlikely to reduce the influence of

stochasticity in any given diffusion. This is especially the case in highly connected networks or one-to-

many diffusion dynamics, because it would not stop one lineage from initially dominating (when a

single invention spreads throughout the population, before another lineage is invented, see **Fig. 4**).

Rather, the tendency for one lineage to dominate would be more heavily impacted by the payoff

structure associated with incremental improvements to products. While we do not believe that

increasing the number of lineages would substantially change the primary findings of this study, a

promising avenue for further research would be a more in-depth exploration of how exposure to

multiple cultural lineages may shape the tempo and mode of CCE.

The evolutionary benefits of CCE not only rely on cultural accumulation, but also on the ability for

new cultural traits to spread through populations. When we extended simulations to examine the time

from recombination to the diffusion of the final higher payoff product, our results suggested that the

network architecture hypothesized to improve time to recombination (multilevel) paradoxically

inhibited diffusion most. These findings complement and extend previous studies demonstrating that

populations with partially-connected network structures can suffer from cultural loss when

connectivity becomes too low for new innovations to spread [12], and that cultural evolution is more

profoundly impacted by the rates of information loss and transmission than differences in social

network architecture [42]. Further, we show that the relative performance of network architectures can

change dramatically when considering performance in terms of acquisition of behaviour by the

majority of individuals in a population, as opposed to the time when a single individual has reached

cultural recombination, especially in larger populations. For example, while populations in a

multilevel social network architecture consistently reached recombination faster than those organized

as random networks under one-to-one diffusion, the multilevel architecture then restricted the final

spread of higher-value cultural traits. These results therefore suggest that multiple dimensions of

performance—including every step from innovations to the final acquisition of higher-valued traits—

may need to be considered when studying the role of social structure in shaping CCE and vice-versa.

Our work reinforces the need for studies of CCE to explicitly consider how network structure interacts

with transmission mechanisms to form a realised transmission network. We show that a very

restrictive transmission dynamic (one-to-one) mitigates the effect of network connectivity on CCE by

generating a partially connected transmission network within an otherwise fully-connected social

network. The consequences of transmission dynamics on CCE were demonstrated, for instance, by

Migliano et al. [19] who found that CCE was faster in simulations where transmission was limited to

kin-based connections (i.e. reduced connectivity). Under one-to-one diffusion, independent lineages

can develop in fully-connected networks because new information is not immediately accessible to all,

leading to more comparable performance between fully and partially connected networks. Thus, when

simulating CCE, it is important to match the transmission dynamics with the time scale of the model.

One-to-many diffusion can be realistic when each epoch represents one generation (e.g. the innovation

of a new medicine [19] could take tens or hundreds of epochs to reach high recombinatory levels),

while the one-to-one diffusion might be more realistic when cultural traits are simpler to recombine.

The production and innovation frequency, as well as transmission biases, may further vary between

species, populations, tasks and contexts. Together with network structure, innovation frequency and

transmission biases may fundamentally alter the transmission dynamics—for example, conformity

overrides pay-off biases [21,43] and homophily reduces social connectivity [18,44]—fuelling

evolutionary feedbacks between network structure and cultural evolution [22]. Both factors will

therefore alter the resulting transmission networks, potentially restricting spread of new cultural traits

and slowing recombinatory CCE. More than highlighting the intricate, yet nuanced, interplay between

demography and cultural transmission, our work strengthens our emerging understanding that

realised connectivity, rather than network architecture, is important for cumulative cultural evolution

[2,3].

Competing interests

The authors have no competing interests with this study

Author Contributions

DRF, DP conceived the idea. CMC, SQS, DRF developed and ran the models and simulations. MC,

CMC analysed the data. MC, LMA, DRF, CMC drafted the initial manuscripts, and all authors

contributed to writing and editing the final article.

Acknowledgments

We thank the Social Evolutionary Ecology Lab and the Cognitive and Cultural Ecology Lab at the Max

Planck Institute of Animal Behavior for the discussion that inspired this manuscript, and the

anonymous reviewers for the insightful comments.

Funding

This work was supported by the Max Planck Society, and the Advanced Centre for Collective

Behaviour, funded by the Deutsche Forschungsgemeinschaft (DFG, German Research Foundation)

under Germany's Excellence Strategy (EXC 2117–422037984). DRF and DP were funded by a grant

from the European Research Council (ERC) under the European Union's Horizon 2020 research and

innovation programme (grant agreement No. 850859), awarded to DRF. DRF was also funded by an

Eccellenza Professorship Grant of the Swiss National Science Foundation (Grant Number

PCEFP3_187058). LMA was funded by a Max Planck Independent Group Leader Fellowship. PH was

funded by a scholarship from the China Scholarship Council (No. 201706100183). M.C. was funded by

CAPES (88881.170254/2018-01) and the Max Planck Society. MCh, DP and SQS received funding from

the IMPRS for Organismal Biology.

References

1. M. Tomasello, The human adaptation for culture. *Annual Rev. Anthrop.* **28(1),** 509-529 (1999).

2. A. Mesoudi, A. Thornton, What is cumulative cultural evolution? *Proc. R. Soc. B.* **285(1880),** 20180712

(2018).

3. M. Derex, A. Mesoudi, A Cumulative cultural evolution within evolving population structures.

Trends Cogn. Sci. 24 (8): 654-667

4. R. Boyd, P.J. Richerson, *The origin and evolution of cultures*. Oxford University Press, Oxford (2005).

5. R. Boyd, P.J. Richerson, J. Henrich, The cultural niche: Why social learning is essential for human

adaptation. *Proc. Natl. Acad. Sci.* **108**, 10918–10925 (2011).

6. S. Shennan, Demography and cultural innovation: a model and its implications for the emergence of

modern human culture. *Cambridge Arch. J.* **11(1)**:5-16 (2001).

7. J. Henrich, Demography and cultural evolution: How adaptive cultural processes can produce

maladaptive losses—the Tasmanian case. *Am. Antiq.* **69(2)**, 197–214 (2004).

8. A. Powell, S. Shennan, M.G. Thomas, Late Pleistocene demography and the appearance of modern

human behavior. *Science* **324(5932)**, 1298–1301 (2009).

9. M. Derex, M.-P. Beugin, B. Godelle, M. Raymond, Experimental evidence for the influence of group

size on cultural complexity. *Nature* **503(7476),** 389–391 (2013).

10. N. Fay, N. de Kleine, B. Walker, C.A. Caldwell, Increasing population size can inhibit cumulative

cultural evolution. *Proc. Natl. Acad. Sci.* **116(14)**, 6726-6731 (2019).

11. J. P. Martens, Scenarios where increased population size can enhance cumulative cultural evolution

are likely common. *Proc. Natl. Acad. Sci.* **116(35),** 17160-17160 (2019).

12. M. Derex, R. Boyd, Partial connectivity increases cultural accumulation within groups. *Proc. Natl.*

Acad. Sci. **113(11)**, 2982-2987 (2016).

13. R. Baldini, Revisiting the effect of population size on cumulative cultural evolution. *J. Cogn. Cult.*

15(3-4), 320-336 (2015).

14. M. Derex, C. Perreault, R. Boyd, Divide and conquer: intermediate levels of population

fragmentation maximize cultural accumulation. *Phil. Trans. R. Soc. B* **373(1743)**, 20170062 (2018).

15. J. Henrich, R. Boyd, M. Derex, M.A Kline, A. Mesoudi, M. Muthukrishna, A.T. Powell, S.J.

Shennan, M.G. Thomas, Understanding cumulative cultural evolution. *Proc. Natl. Acad. Sci.* **113(44)**,

E6724-E6725 (2016).

16. B. Voelkl, R. Nöe, The influence of social structure on the propagation of social information in

artificial primate groups: a graph-based simulation approach. *J. Theor. Biol.* **252(1)**, 77-86 (2008)

17. C.L. Nunn, P.H. Thrall, K. Bartz, T. Dasgupta, C. Boesch, Do transmission mechanisms or social

systems drive cultural dynamics in socially structured populations? *An. Behav.* **77(6)**,*1515-1524*

(2009).

18. M. Cantor, H. Whitehead, The interplay between social networks and culture: theoretically and

among whales and dolphins. *Phil. Trans. R. Soc. B* **368(1618)**, 20120340 (2013).

19. A.B. Migliano, F. Battiston, S. Viguier, A.E. Page, M. Dyble, R. Schlaepfer, D. Smith, L. Astete, M.

Ngales, J. Gomez-Gardenes, V. Latora, Hunter-gatherer multilevel sociality accelerates cumulative

cultural evolution. *Science Adv.* **6(9)**, p.eaax5913 (2020).

20. H. Whitehead, D. Lusseau, Animal social networks as substrate for cultural behavioural diversity.

J. Theor. Biol. **294**,19-28 (2012).

21. L.M. Aplin, D.R. Farine, J. Morand-Ferron, A. Cockburn, A. Thornton, B.C. Sheldon,

Experimentally induced innovations lead to persistent culture via conformity in wild birds. *Nature*

518(7540), 538-541 (2015).

22. M., Smolla, E. Akçay, Cultural selection shapes network structure. *Sci. Adv*. **5(8)**, eaaw0609 (2019).

23. D.J. Watts, S.H. Strogatz, Collective dynamics of 'small-world' networks. *Nature* **393(6684)**, 440-442

(1998)

24. M. Newman, *Networks*. Oxford University Press (2018). New York, NY, USA

25. R.G. Miller Jr, *Survival analysis* (Vol. 66). John Wiley & Sons (1981). New York, NY, USA

26. R Core Team. R: A language and environment for statistical computing. R Foundation for

Statistical Computing, Vienna, Austria. URL https://www.R-project.org/ (2019).

27. T. Therneau, A Package for Survival Analysis in R. R package version 3.2-3, https://CRAN.R-

project.org/package=survival (2020)

28. A. Kassambara, M. Kosinski, P. Biecek, Survminer: Drawing Survival Curves using 'ggplot2'. R

package version 0.4.7. https://CRAN.R-project.org/package=survminer (2020)

29. M. Cantor, M. Chimento, S.Q. Smeele, P. He, D. Papageorgiou, L.M. Aplin, D.R. Farine. Social

network architecture and the tempo of cumulative cultural evolution. Github repository. (2020).

https://github.com/simeonqs/Social_network_architecture_and_the_tempo_of_cumulative_cultu

ral_evolution

30. T.A. Wilkin, D. Garant, A.G. Gosler, B.C. Sheldon, Density effects on life-history traits in a wild

population of the great tit *Parus major*: analyses of long-term data with GIS techniques. *J. Anim.*

Ecol., **75**(2), 604-615 (2006)

31. J.P. Hailman, M.S. Ficken. Combinatorial animal communication with computable syntax: chick-a-

dee calling qualifies as 'language' by structural linguistics. *An. Behav.* **34**, 1899–1901 (1986).

32. L.L. Bloomfield, I. Charrier, C.B. Sturdy, Note types and coding in parid vocalizations. II: The

chick-a-dee call of the Mountain Chickadee (*Poecile gambeli*). *Can. J. Zool.* **82**, 780–793 (2004).

33. D.E. Kroodsma, Winter Wren singing behavior: a pinnacle of song complexity. *Condor* **82**, 357–365

(1980).

34. D.W. Leger, First Documentation of combinatorial song syntax in a suboscine passerine species.

Condor **107(4)**, 765–774 (2005).

35. C.L. Branch, V.V. Pravosudov, Mountain chickadees from different elevations sing different songs:

acoustic adaptation, temporal drift or signal of local adaptation? *R. Soc. Open Sci* **2.4,** 150019 (2015).

36. L. M. Aplin, Culture and cultural evolution in birds: a review of the evidence. *Anim. Behav.* **147**,

179-187 (2019).

37. A. Whiten, Cultural evolution in animals. *Ann. Rev. Ecol. Evol. System.* **50,** 27-48 (2019).

38. D.J. Watts, A simple model of global cascades on random networks. *Proc. Natl. Acad. Sci.* **99(9)**,

5766-5771 (2002).

39. C.L. Apicella, F.W. Marlowe, J.H. Fowler, N.A. Christakis, Social networks and cooperation in

hunter-gatherers. *Nature* **481(7382)**, 497-501 (2012).

40. B.J. West, G.F. Massari, G. Culbreth, R. Failla, M. Bologna, R.I.M. Dunbar, P. Grigolini, Relating size

and functionality in human social networks through complexity. *Proc. Natl. Acad. Sci.* **117(31)**,

18355-18358 (2020).

41. R.I.M. Dunbar, Structure and function in human and primate social networks: implications for

diffusion, network stability and health. *Proc. R. Soc. A* **476**:20200446 (2020).

42. S., Yeaman, A. Schick, L. Lehmann, Social network architecture and the maintenance of deleterious

cultural traits. *J R Soc Interface* **9(70)**, 848-858 (2012).

43. K.K. Denton, Y. Ram, U. Liberman, M.W. Feldman, Cultural evolution of conformity and

anticonformity. *Proc. Natl. Acad. Sci* 117(24), 13603-13614 (2020)

www.pnas.org/cgi/doi/10.1073/pnas.2004102117

44. M. McPherson, L. Smith-Lovin, J.M. Cook, Birds of a feather: Homophily in social networks.

Annual Review of Sociology, **27(1)**, 415-444 (2001).

45. M. Cantor, M. Chimento, S.Q. Smeele, P. He, D. Papageorgiou, L.M. Aplin, D.R. Farine. Data from:

Social network architecture and the tempo of cumulative cultural evolution, Dryad, Dataset (2021)

doi: https://doi.org/10.5061/dryad.3r2280gff

Figures

Figure 1. Social network architectures, and the time to recombination for each architecture across population sizes and levels of connectivity using model 1. (A) Network architectures vary in clustering and modularity: Random (unclustered C=0.03, non-modular Q=0.24), small-world (clustered C=0.52, medium-modular Q=0.63), lattice (clustered C=0.45, medium-modular Q=0.54), modular (unclustered C=0.23, modular Q=0.82), modular lattices (clustered C=0.41, modular Q=0.81), multilevel (clustered C=0.42, modular Q=0.83). Each binary network depicts populations with the same number of individuals (here, N=324 nodes) that have the same number of social connections (here, degree K=12 links per node; density

D=0.037) but are wired differently. **(B)** Cumulative incidence of recombination events (y-axis) as a stepwise

function over time (x-axis, log epochs) for small (N=64), medium (N=144), and large population sizes

(N=324). The line shading represents the amount of network connectivity (node degree K, where the lighter

the shade, the smaller the degree (K∈{8,12} for N=64; K∈{8,12,18,24} for N=144; K∈{8,12,18,24,30} for

N=324). Vertical dashed lines indicate the median of time to recombination (S(t)≤0.5) per network

connectivity, across architectures. The time to reach recombination was truncated to 100 epochs for better

visualization. Curves were calculated based on 5,000 simulations.

Figure 2. Time to recombination and time from recombination to diffusion across network architectures

with varying sizes but a fixed degree. Comparison of the performance across the range of network

architectures of the same degree (here K=12 links per node) and fully connected networks of the same size

(N=64, N=144, and N=324 nodes, K=63, 143, and 323 respectively). **(A)** Time to recombination (log epochs)

from 5,000 simulations with model 1 that uses a broadcast (one-to-many) diffusion dynamic. **(B)** Time to

recombination (log epochs) from 5,000 simulations with model 2 that uses a dyadic (one-to-one) diffusion

dynamic. **(C)** Difference between the time to recombination and the time to diffusion, where time to

diffusion corresponds to the latency until the majority of the individuals in the population has information

about the final higher-payoff product, from 5,000 simulations using model 2 (one-to-one diffusion). All

ridges were plotted with the same bandwidth (0.18).

Figure 3. Relative performance of network architectures within each of the 11 combinations of

population size and level of connectivity used in the simulations. Each row of each table reports the

coefficient estimate of GLMs of network architecture (column) in function of the time to recombination

while maintaining degree (row) and network size (box) constant. Higher coefficients (red colours) represent

a poorer performance (longer latency to recombination) while lower coefficients (blue colours) represent

network architectures that perform better (shorter latency to recombination) for that combination of

population size and level of connectivity (using random networks as the reference architecture in the GLM

intercept). The relative performance of each architecture is shown for **(A)** time to recombination under a

one-to-many diffusion mechanism (model 1), **(B)** time to recombination under a one-to-one diffusion

mechanism (model 2), and **(C)** total time to diffusion (from simulation start until the majority of the

population has information about the final higher-payoff product) under a one-to-one diffusion mechanism

(model 2).

Figure 4. Cultural product diversity across time in a fully connected social network and in a highly

structured network architecture. The distributions of time to recombination and cultural product diversity

illustrates how early stochasticity can affect cultural diversity, even within the same network architecture,

and therefore can shape the outcomes of cumulative cultural evolution. **(A)** Time to recombination (epochs)

from 5,000 simulations with one-to-many diffusion dynamics (model 1) in multilevel and fully-connected

networks (with N=64 and K=12) highlights distinct cultural trajectories among the highly structured

networks (note the bimodal distribution). Following panels show cultural diversity over time from one

simulation taken from the **(B)** single mode of the fully-connected network, and the **(C)** first (*) and **(D)**

second (**) modes of the distribution of results from the multilevel architecture. Cultural diversity (y-axis)

represents the proportion of the population with one of the possible products over time: a combination of

two inventory items (2nd stage; dotted lines), a valid combination triad of items (3rd stage; dashed lines),

and the final higher payoff product, i.e. a triad recombining products from the two lineages (recombination;

thick full lines). These products could come from two independent cultural lineages: lineage A (top row)

and lineage B. For better visualization, the distribution of single inventory items (1st stage product) was

omitted (but see **Fig. S1**).

Supporting Material for:
Social network architecture and the tempo of cumulative cultural evolution
581
582 Mauricio Cantor*, Michael Chimento, Simeon Q. Smeele, Peng He, Danai Papageorgiou, Lucy M.
583 Aplin*, Damien R. Farine
584
Corresponding Authors:
* Mauricio Cantor. Department for the Ecology of Animal Societies, Max Planck Institute of Animal
Behavior, Am Obstberg 1, Radolfzell 78315, Germany;
* Lucy M. Aplin. Cognitive & Cultural Ecology Research Group, Max Planck Institute of Animal Behavior,
Am Obstberg 1, Radolfzell 78315, Germany, Phone: +49 7732 1501-13.
Emails: mcantor@ab.mpg.de, laplin@ab.mpg.de

Table S1. Generalized linear models (GLMs) for agent-based model 1 (with one-to-many
diffusion mechanism) and model 2 (one-to-one diffusion), in which time to recombination (time
to recombination, measured as log(epoch+1)) is predicted by an interaction between population
size (N= number of nodes) and social network architecture. Values represent the model
coefficients (and standard errors, SE), which are exponentiated for easy reference against the
constant, as these GLMs used a log link function. Network connectivity was not included, as this
contains the same information as population size for fully connected networks. Intercepts
represents random network architectures of population size N=64. *t*-values and *p*-values are also
presented, where ***p<0.01.
600

Dependent variable: log(epoch + 1)	Model 1			Model 2		
	β (SE)	t-value	p-value	β (SE)	t-value	p-value
Intercept	3.425 (0.003)	1,328.07	0.000***	4.205 (0.002)	2,442.73	0.000***
architecture (small-world)	0.999 (0.004)	273.691	0.000***	1.003 (0.002)	412.456	0.000***
architecture (lattice)	1.010 (0.004)	278.152	0.000***	1.013 (0.002)	418.671	0.000***
architecture (modular)	0.970 (0.004)	261.819	0.000***	0.994 (0.002)	407.023	0.000***
architecture (modular lattice)	0.972 (0.004)	262.838	0.000***	0.991 (0.002)	405.079	0.000***
architecture (multilevel)	0.977 (0.004)	264.828	0.000***	0.982 (0.002)	399.537	0.000***
architecture (full)	1.652 (0.003)	486.607	0.000***	1.033 (0.003)	353.642	0.000***
population size (N=144)	0.715 (0.004)	197.009	0.000***	0.759 (0.002)	322.453	0.000***
population size (N=324)	0.512 (0.004)	125.054	0.000***	0.587 (0.003)	231.672	0.000***
architecture (small-world) : population size (N=144)	0.999 (0.005)	194.533	0.000***	0.999 (0.003)	300.603	0.000***
architecture (lattice) : population size (N=144)	0.996 (0.005)	194.851	0.000***	0.989 (0.003)	298.247	0.000***
architecture (modular) : population size (N=144)	1.013 (0.005)	195.078	0.000***	0.996 (0.003)	298.061	0.000***
architecture (modular lattice) : population size (N=144)	0.999 (0.005)	191.92	0.000***	1.003 (0.003)	300.253	0.000***
architecture (multilevel) : population size (N=144)	1.024 (0.005)	198.357	0.000***	1.011 (0.003)	301.777	0.000***
architecture (full) : population size (N=144)	1.206 (0.005)	243.188	0.000***	0.994 (0.005)	217.448	0.000***
architecture (small-world) : population size (N=324)	1.005 (0.006)	173.584	0.000***	0.997 (0.004)	278.573	0.000***
architecture (lattice) : population size (N=324)	1.012 (0.006)	176.012	0.000***	0.986 (0.004)	276.2	0.000***
architecture (modular) : population size (N=324)	1.059 (0.006)	182.989	0.000***	1.001 (0.004)	278.842	0.000***
architecture (modular lattice) : population size (N=324)	1.036 (0.006)	178.259	0.000***	1.006 (0.004)	280.226	0.000***
architecture (multilevel) : population size (N=324)	1.100 (0.006)	192.975	0.000***	1.015 (0.004)	282.066	0.000***
architecture (full) : population size (N=324)	1.406 (0.006)	252.195	0.000***	0.980 (0.005)	182.571	0.000***
Observations	345,000			345,000		
Log Likelihood	-446,718.40			-378,097.40		
Akaike Information Criterion	893,478.70			756,236.80		

Table S2. Generalized linear models (GLMs) for agent-based model 1 (with one-to-many
diffusion mechanism) and model 2 (one-to-one diffusion), in which time to recombination (time
to recombination, measured as log(epoch+1)) is predicted by an interaction between population
size (N = number of nodes), network architecture, and network connectivity (i.e. average network
degree, K). Coefficients are exponentiated for easy reference to the intercept, as these GLMs used
a log link function. Full networks were excluded. Intercepts represent random network
architectures of population size *N*=64, average degree of *K*=8. *t*-values and *p*-values are also
presented, where ***p<0.01.
609

Dependent variable: log(epoch + 1)	Model 1			Model 2		
	β (SE)	t-value	p-value	β (SE)	t-value	p-value
Intercept	3.196 (0.004)	827.046	0.000***	4.168 (0.002)	1,707.64	0.000***
architecture (small-world)	0.999 (0.005)	182.779	0.000***	1.009 (0.003)	293.735	0.000***
architecture (lattice)	1.010 (0.005)	185.703	0.000***	1.011 (0.003)	294.487	0.000***
architecture (modular)	1.005 (0.005)	184.396	0.000***	1.006 (0.003)	292.469	0.000***
architecture (modular lattice)	1.011 (0.005)	186.113	0.000***	1.004 (0.003)	291.433	0.000***
architecture (multilevel)	1.020 (0.005)	188.461	0.000***	0.993 (0.003)	286.534	0.000***
population size (N=144)	0.724 (0.007)	109.84	0.000***	0.756 (0.004)	186.603	0.000***
population size (N=324)	0.618 (0.007)	84.05	0.000***	0.595 (0.005)	124.502	0.000***
connectivity (K=12)	1.143 (0.005)	222.751	0.000***	1.018 (0.003)	297.42	0.000***
connectivity (K=18)	0.847 (0.010)	87.507	0.000***	0.996 (0.006)	171.202	0.000***
connectivity (K=24)	0.831 (0.010)	84.886	0.000***	0.995 (0.006)	171.007	0.000***
connectivity (K=30)	0.856 (0.010)	89.049	0.000***	0.992 (0.006)	170.253	0.000***
architecture (small-world) : population size (N=144)	0.997 (0.009)	106.845	0.000***	0.992 (0.006)	173.547	0.000***
architecture (lattice) : population size (N=144)	1.007 (0.009)	108.852	0.000***	0.989 (0.006)	173.083	0.000***
architecture (modular) : population size (N=144)	0.973 (0.009)	103.674	0.000***	0.998 (0.006)	174.678	0.000***
architecture (modular lattice) : population size (N=144)	0.942 (0.009)	99.622	0.000***	1.008 (0.006)	176.732	0.000***
architecture (multilevel) : population size (N=144)	0.967 (0.009)	103.559	0.000***	1.014 (0.006)	177.167	0.000***
architecture (small-world) : population size (N=324)	1.004 (0.010)	96.642	0.000***	0.987 (0.007)	146.186	0.000***
architecture (lattice) : population size (N=324)	1.021 (0.010)	99.355	0.000***	0.988 (0.007)	146.499	0.000***
architecture (modular) : population size (N=324)	0.930 (0.011)	87.286	0.000***	0.993 (0.007)	147.195	0.000***
architecture (modular lattice) : population size (N=324)	0.876 (0.011)	80.546	0.000***	0.996 (0.007)	147.486	0.000***
architecture (multilevel) : population size (N=324)	0.955 (0.010)	91.24	0.000***	1.005 (0.007)	148.566	0.000***

architecture (small-world) : connectivity (K=12)	0.999 (0.007)	137.563	0.000***	0.987 (0.005)	204.283	0.000***
architecture (lattice) : connectivity (K=12)	1.000 (0.007)	138.353	0.000***	1.004 (0.005)	208.69	0.000***
architecture (modular) : connectivity (K=12)	0.934 (0.007)	127.039	0.000***	0.976 (0.005)	200.973	0.000***
architecture (modular lattice) : connectivity (K=12)	0.928 (0.007)	126.334	0.000***	0.974 (0.005)	200.323	0.000***
architecture (multilevel) : connectivity (K=12)	0.922 (0.007)	125.835	0.000***	0.978 (0.005)	200.263	0.000***
architecture (small-world) : connectivity (K=18)	1.002 (0.014)	73.377	0.000***	1.003 (0.008)	121.873	0.000***
architecture (lattice) : connectivity (K=18)	0.996 (0.013)	73.791	0.000***	0.997 (0.008)	121.036	0.000***
architecture (modular) : connectivity (K=18)	1.131 (0.014)	82.63	0.000***	0.988 (0.008)	119.744	0.000***
architecture (modular lattice) : connectivity (K=18)	1.170 (0.014)	83.951	0.000***	0.996 (0.008)	120.896	0.000***
architecture (multilevel) : connectivity (K=18)	1.130 (0.013)	84.295	0.000***	0.998 (0.008)	121.107	0.000***
architecture (small-world) : connectivity (K=24)	1.005 (0.014)	72.791	0.000***	1.002 (0.008)	121.564	0.000***
architecture (lattice) : connectivity (K=24)	0.992 (0.014)	72.609	0.000***	1.003 (0.008)	121.974	0.000***
architecture (modular) : connectivity (K=24)	1.163 (0.014)	84.717	0.000***	0.991 (0.008)	120.13	0.000***
architecture (modular lattice) : connectivity (K=24)	1.231 (0.014)	88.561	0.000***	0.996 (0.008)	120.82	0.000***
architecture (multilevel) : connectivity (K=24)	1.172 (0.013)	87.205	0.000***	0.995 (0.008)	120.561	0.000***
architecture (small-world) : connectivity (K=30)	0.999 (0.014)	73.539	0.000***	1.006 (0.008)	122.03	0.000***
architecture (lattice) : connectivity (K=30)	0.977 (0.013)	72.431	0.000***	1.000 (0.008)	121.265	0.000***
architecture (modular) : connectivity (K=30)	1.168 (0.013)	86.595	0.000***	0.996 (0.008)	120.651	0.000***
architecture (modular lattice) : connectivity (K=30)	1.237 (0.014)	90.541	0.000***	0.992 (0.008)	120.044	0.000***
architecture (multilevel) : connectivity (K=30)	1.175 (0.013)	89.005	0.000***	0.999 (0.008)	121.122	0.000***
population size (N=144): connectivity (K=12)	0.864 (0.009)	94.311	0.000***	0.992 (0.006)	174.263	0.000***
population size (N=324): connectivity (K=12)	0.796 (0.011)	74.925	0.000***	0.975 (0.007)	144.303	0.000***
population size (N=144): connectivity (K=18)	1.251 (0.012)	102.992	0.000***	1.023 (0.007)	138.758	0.000***
population size (N=144): connectivity (K=24)	1.426 (0.012)	118.629	0.000***	1.030 (0.007)	139.805	0.000***
architecture (small-world) : population size (N=144) : connectivity (K=12)	1.000 (0.013)	77.063	0.000***	1.014 (0.008)	126.148	0.000***
architecture (lattice) : population size (N=144) : connectivity (K=12)	0.989 (0.013)	76.711	0.000***	0.998 (0.008)	124.302	0.000***
architecture (modular) : population size (N=144) : connectivity (K=12)	1.101 (0.013)	84.372	0.000***	1.017 (0.008)	126.196	0.000***
architecture (modular lattice) : population size (N=144) : connectivity (K=12)	1.134 (0.013)	86.46	0.000***	1.016 (0.008)	126.215	0.000***
architecture (multilevel) : population size (N=144) : connectivity (K=12)	1.133 (0.013)	87.411	0.000***	1.016 (0.008)	125.951	0.000***

architecture (small-world) : population size (N=324) : connectivity (K=12)	0.999 (0.015)	66.585	0.000***	1.017 (0.010)	106.402	0.000***
architecture (lattice) : population size (N=324) : connectivity (K=12)	0.989 (0.015)	66.504	0.000***	0.995 (0.010)	104.209	0.000***
architecture (modular) : population size (N=324) : connectivity (K=12)	1.125 (0.015)	73.525	0.000***	1.027 (0.010)	107.365	0.000***
architecture (modular lattice) : population size (N=324) : connectivity (K=12)	1.160 (0.016)	74.596	0.000***	1.030 (0.010)	107.742	0.000***
architecture (multilevel) : population size (N=324) : connectivity (K=12)	1.154 (0.015)	76.922	0.000***	1.024 (0.010)	106.886	0.000***
architecture (small-world) : population size (N=144) : connectivity (K=18)	1.005 (0.017)	58.571	0.000***	1.000 (0.010)	95.91	0.000***
architecture (lattice) : population size (N=144) : connectivity (K=18)	0.999 (0.017)	58.796	0.000***	1.006 (0.010)	96.475	0.000***
architecture (modular) : population size (N=144) : connectivity (K=18)	0.907 (0.017)	52.727	0.000***	0.991 (0.010)	94.773	0.000***
architecture (modular lattice) : population size (N=144) : connectivity (K=18)	0.902 (0.017)	51.672	0.000***	0.977 (0.010)	93.62	0.000***
architecture (multilevel) : population size (N=144) : connectivity (K=18)	0.921 (0.017)	54.341	0.000***	0.982 (0.010)	94.073	0.000***
architecture (small-world) : population size (N=144) : connectivity (K=24)	0.994 (0.017)	58.554	0.000***	0.997 (0.010)	95.705	0.000***
architecture (lattice) : population size (N=144) : connectivity (K=24)	0.981 (0.017)	58.226	0.000***	0.998 (0.010)	95.864	0.000***
architecture (modular) : population size (N=144) : connectivity (K=24)	0.835 (0.017)	49.062	0.000***	0.981 (0.010)	93.886	0.000***
architecture (modular lattice) : population size (N=144) : connectivity (K=24)	0.794 (0.017)	46.1	0.000***	0.973 (0.010)	93.235	0.000***
architecture (multilevel) : population size (N=144) : connectivity (K=24)	0.836 (0.017)	49.902	0.000***	0.977 (0.010)	93.492	0.000***
Observations	330,000			330,000		
Log Likelihood	-423,477.80			-359,565.20		
Akaike Information Criterion	847,087.60			719,262.30		

610
611

Fully connected

Random

Small-world

Lattice

Modular

Modular lattice

Multilevel

lineage A   lineage B

Proportion of population

Time (epoch)

—— Item   ▪ ▪ Dyad   ▬ ▬ Triad   ▬▬▬ Recombination

Figure S1. Cultural trait diversity across social network architectures and proposed null
models. Comparison of the diversity of cultural traits along the network architecture spectrum,
across populations of same size (N=64 nodes) and connectivity (average degree K=12 links) in
relative to a fully connected network (top) of the same size (N=64, K=63). Cultural innovation
diversity across network architectures was measured as the proportion of the population with
one of the four possible traits over time (y-axis): a single inventory item (1st stage; thin lines), a
combination of two items (2nd stage; dotted lines), a valid combination triad of items (3rd stage;
dashed lines), and the final higher payoff product, i.e. a triad recombining products from the two
lineages (recombination; thick full lines). The inventory items came from two independent
cultural lineages: A (black lines) and B (red lines).
(red lines).