Title: Granzyme B skews CD4[+] T cell differentiation resulting in increased intestinal

pathogenicity

Authors: Kristen L. Hoek,[a] Michael J. Greer,[b*] Kathleen G. McClanahan,[a*] Ali Nazmi,[a] M. Blanca

Piazuelo,[c,e] Kshipra Singh,[c,e] Keith T. Wilson,[a,c,e,f,g] and Danyvid Olivares-Villagómez.[a,e,g,h]

*These two authors contributed equally to this report

Affiliations: [a]Department of Pathology, Microbiology and Immunology, Vanderbilt University

Medical Center, A5301 Medical Center North, Nashville, Tennessee 37232, USA; [b]Department of

Biomedical Informatics, Vanderbilt University, Nashville, Tennessee, 37232, USA; [c]Division of

Gastroenterology, Hepatology and Nutrition, Department of Medicine, Vanderbilt University

Medical Center, Nashville, Tennessee 37232, USA; [e]Center for Mucosal Inflammation and

Cancer, Vanderbilt University Medical Center, Nashville, Tennessee 37232; [f]Veterans Affairs

Tennessee Valley Healthcare System, Nashville, Tennessee 37232; [g]Vanderbilt Institute for

Infection, Immunology and Inflammation, Vanderbilt University Medical Center, A5301 Medical

Center North, Nashville, Tennessee 37232, USA. [h]ORCID, 0000-0002-1158-8976

Corresponding author: Danyvid Olivares-Villagómez. Address: Vanderbilt University Medical

center, A5211 Medical Center North, Nashville Tennessee 37232-0001. Fax number: 615-343-

7392. Telephone number: 615-936-0134. email: danyvid.olivares-villagomez@vumc.org.

Abstract: CD4$^+$ T cell activation and differentiation are important events that set the stage for

proper immune responses. Many factors are involved in the activation and differentiation of T

cells, and these events are tightly controlled to prevent unwanted and/or exacerbated immune

responses that may harm the host. It has been well documented that granzyme B, a potent serine

protease involved in cell-mediated cytotoxicity, is readily expressed by certain CD4$^+$ T cells, such

as regulatory T cells and CD4$^+$CD8$\alpha\alpha^+$ intestinal intraepithelial lymphocytes, both of which

display cytotoxicity associated with granzyme B. However, because not all CD4$^+$ T cells

expressing granzyme B are cytotoxic, additional roles for this protease in CD4$^+$ T cell biology

remain unknown. Here, using a combination of *in vivo* and *in vitro* approaches, we report that

granzyme B-deficient CD4$^+$ T cells display increased IL-17 production. In the adoptive transfer

model of intestinal inflammation, granzyme B-deficient CD4$^+$ T cells triggered a more rapid

disease onset than their WT counterparts, and presented a differential transcription profile. Similar

results were also observed in granzyme B-deficient mice infected with *Citrobacter rodentium*. Our

results suggest that granzyme B modulates CD4$^+$ T cell differentiation, providing a new

perspective into the biology of this enzyme.

Introduction

Granzymes are a family of serine proteases expressed by many immune cell populations.

There are five different granzymes in humans (A, B, H, K and M) and 11 in mice (A, B, C, D, E,

F, G, K, L, M and N), which differ in function and substrate specificity.[1] Most research has focused

on granzymes A and B due to their hallmark role in CD8[+] T and NK cell-mediated cytotoxicity

against tumor cells or virus-infected cells.[2] However, mounting evidence indicates that granzymes

are also involved in other processes, including inflammatory responses,[3-5] remodeling of the

extracellular matrix,[6] and cardiovascular disorders.[7]

Activated CD4[+] T cells express granzyme B, and some of these cells exhibit lytic activity

similar to CD8[+] T cells.[8, 9] Antigen-specific CD4[+] T cell lytic activity mediated by granzyme B

can be triggered by viral and bacterial infections.[10-12] Similarly, some tumor-infiltrating CD4[+] T

cells also present anti-tumor cytotoxicity mediated by granzyme B.[13] In addition, one of the

mechanisms by which regulatory CD4[+] T cells in humans and mice control immune responses is

through granzyme B-mediated cell death.[14-18]

After activation, CD4[+] T cells can differentiate into a diverse set of T helper (Th) cells

depending on environmental cues and transcription factors, which ultimately leads Th cells to

produce a specific cytokine profile.[19] In certain populations of differentiated CD4[+] T cells,

granzyme B controls CD4[+] T cell immune responses by initiating activation-induced cell death.[20]

Downregulation of granzyme B, for example when Th2 cells are cultured in the presence of

vasoactive intestinal peptide, reduces Th2 cell death.[20] Moreover, granzyme B expression in Th2

cells is enhanced by IL-10R signaling, which results in cell death.[21] These studies clearly indicate

that granzyme B is involved in controlling some immune CD4[+] T cells responses. However, it is

unclear whether this enzyme is actively involved during the process of Th differentiation.

In this report, we present data indicating that granzyme B is expressed during CD4$^+$ T cell

differentiation. Cells lacking granzyme B show a different cytokine profile than granzyme B-

competent cells, primarily presenting increased IL-17 production. Moreover, using the adoptive

transfer model of intestinal inflammation, we show that granzyme B-deficient CD4$^+$ T cells

activated *in vivo* have a different gene expression profile and display increased pathogenicity

relative to WT cells. Similarly, we present evidence indicating that granzyme B deficiency

predisposes mice to higher susceptibility to *Citrobacter rodentium* infection. These results indicate

that granzyme B is required for proper CD4$^+$ T cell differentiation, and in its absence, CD4$^+$ T cells

may acquire an aberrant phenotype characterized by increased IL-17 production and pathogenicity.

Results

Th0 and Th1, but not Th17 CD4$^+$ T cells express granzyme B during differentiation

To determine the role of granzyme B in T cell activation and differentiation, we cultured

naïve CD4$^+$ T cells in Th0, Th1 and Th17 differentiation conditions. One and three days after

culture, RNA was extracted to determine granzyme B expression relative to naïve CD4$^+$ T cells.

Under Th0 conditions, expression of granzyme B at day 1 did not increase above the levels

observed for naïve CD4$^+$ T cells (Figure 1a, top), but presented an almost two-fold increase at day

3 (Figure 1a, bottom). Th1 cells expressed a 6-fold increase in granzyme B mRNA over naïve T

cells at day 1, decreasing to 2-fold at day 3. On the other hand, CD4$^+$ T cells under Th17

differentiation conditions did not induce granzyme B mRNA expression at either time point

(Figure 1a).

Around 15% of Th0 and Th1 cells expressed granzyme B intracellularly at day 5 (Figure

1b), whereas this enzyme was not detected in Th17 cells, confirming the results for granzyme B

mRNA expression. Granzyme B in Th1 cells was not detected at 1 day after differentiation but it

was conspicuously expressed at day 3 and day 5 (Figure 1c). On the other hand, Th17 cells did not

express granzyme B at any of the time points analyzed (Figure 1c). Because Th1 and Th17 cultures

contain different cytokine/antibody cocktails, and granzyme B is expressed in the former but not

in the latter helper cells, we investigated what factors present in these culture conditions induce or

repress granzyme B expression. For this purpose, we activated CD4$^+$ T cells under Th0 conditions

(anti-CD3/CD28, IL-2, and irradiated spleen cells), in the presence of individual factors present in

Th1 and Th17 differentiation cocktails (Figure 1d). IL-2 alone induced ~7.5% of CD4$^+$ T cells to

express granzyme B, which was not significantly different from cells cultured without IL-2 or in

the presence of IL-12, IL-1β, and IL-23, these last two cytokines involved in Th17 differentiation.

However, Th17-inducing cytokines IL-6 and TGFβ, as well as the blocking anti-IFNγ antibody

significantly ablated granzyme B expression. These results indicate that CD4$^+$ T cell activation

induces granzyme B activation, whereas some cytokines and anti-IFNγ inhibit its expression.

Because most of our differentiation cultures included spleen cells as a source of APC, we

determined whether these cells influence granzyme B expression during CD4$^+$ T cell activation.

As shown in Figure 1e, CD4$^+$ T cell activation in the absence of APC (but including IL-2), resulted

in less than 2% of CD4$^+$ T cells expressing granzyme B, whereas irradiated spleen APC

significantly increased granzyme B expression.

Overall, these results indicate that granzyme B is expressed during Th0 and Th1

differentiation, with irradiated APC being an important expression-enhancing factor. On the other

hand, Th17 differentiation conditions ablate granzyme B expression. This effect can be attributed

to IL-6, TGFβ, and anti-IFNγ.

As shown in Figure 1b, during Th0 and Th1 differentiation, two populations of CD4$^+$ T

cells can be distinguished: WT-gzmB^{pos} and WT-gzmB^{neg}. To determine whether these

populations present different cytokine profiles, we determined the frequencies of IFNγ and IL-17-

producing cells in each of these subpopulations. Around half of Th0 WT-gzmB^{pos} expressed IFNγ,

whereas only around 5% of the WT-gzmB^{neg} CD4$^+$ T cells were positive for IFNγ (Figure 1f). Th0

cells derived from granzyme B-deficient (Gzmb$^{-/-}$) mice presented decreased levels of IFNγ-

producing cells in comparison to WT-gzmB^{pos} cells, but significantly more IFNγ$^+$ cells than WT-

gzmB^{neg} cells (Figure 1f). On the other hand, IL-17 production was not significantly different

among Th0 cells from any of the CD4$^+$ T cell populations.

Th1 differentiation induced high levels of IFNγ$^+$ cells in all CD4$^+$ T cell populations

analyzed; however, WT-gzmB^{pos} cells presented the highest levels of IFNγ$^+$ cells in comparison

to WT-gzmB^{neg} or cells derived from Gzmb$^{-/-}$ mice (Figure 1g). Interestingly, under Th1

conditions, cells from Gzmb$^{-/-}$ mice presented significantly greater levels of IL-17$^+$ cells when

compared to WT-gzmB^{pos} and WT-gzmB^{neg} cells (Figure 1g). These results suggest that granzyme

B may be involved in preventing IL-17 production when CD4$^+$ T cells are cultured under Th1

conditions.

Mesenteric lymph node APC enhance granzyme B and IL-17 expression

As shown in the previous section, addition of irradiated spleen APC to the cultures

enhanced granzyme B expression by activated CD4$^+$ T cells under Th0 conditions (Figure 1e).

Because APC activity varies between lymphoid organs, we investigated whether APC from spleen

and mesenteric lymph nodes (MLN) induced differential granzyme B expression, and if so,

whether IFNγ and IL-17 expression is affected. Spleen APC induced greater granzyme B

expression than their MLN counterparts in WT CD4$^+$ T cells (Supp. Figure 1a). IFNγ production

between WT-gzmB^{pos} CD4$^+$ T cells was similar regardless of the source of irradiated APC, which

was also observed between WT-gzmB^{neg} CD4$^+$ T cells (Supp. Figure 1b). However, irradiated

APC from MLN induced higher IL-17 production than spleen APC in the WT-gzmB^{pos} CD4$^+$ T

cell population (Supp. Figure 1c). Granzyme B-deficient CD4$^+$IFNγ$^+$ T cells were similar between

cultures containing spleen or MLN APC (Supplemental Fig. 1d). However, increased IL-17

production was also observed in granzyme B-deficient CD4$^+$ T cells differentiated in the presence

of MLN APC (Supp. Figure 1e).

Overall, these results indicate that APC present in distinct immunological compartments

stimulate differential granzyme B expression. Moreover, APC that induce CD4$^+$ T cells to express

less granzyme B also seem to promote increased IL-17 production (particularly in the WT-gzmB^{pos}

and granzyme B-deficient CD4$^+$ T cell populations), suggesting an inverse relationship between

granzyme B expression and IL-17 production.

***In vivo* activation of granzyme B-deficient CD4$^+$ T cells results in increased pathogenicity**

To further understand the role of granzyme B in CD4$^+$ T cell differentiation, we adoptively

transferred naïve CD4$^+$CD45RB^{hi} T cells from WT or *Gzmb$^{-/-}$* mice into *Rag2$^{-/-}$* recipient mice.

Because it is well-established that transfer of naïve CD4$^+$ T cells into immunocompromised

recipients results in chronic intestinal inflammation, [22, 23] this set up allows us to: 1) determine

whether granzyme B-deficiency alters CD4$^+$ T cell pathogenicity, and 2) whether absence of

granzyme B influences the cytokine profile of CD4$^+$ T cells activated *in vivo* (discussed in the next

section).

In our animal colony, transfer of naïve CD4$^+$ T cells from WT mice typically results in

disease symptoms by 6 weeks after transfer. However, when naïve CD4$^+$ T cells from *Gzmb*$^{-/-}$ mice

were transferred into *Rag2*$^{-/-}$ mice, weight loss was evident starting at 14 days after transfer, and

by day 21 these mice had lost around 20% of their original weight (Figure 2a). At this point, the

experiments were terminated to comply with our institution's animal welfare regulations. In

contrast, at day 21, *Rag2*$^{-/-}$ mice that received WT cells were either gaining or maintaining their

weight (Figure 2a). These results suggest that absence of granzyme B resulted in more pathogenic

T cells. To determine whether this was a CD4$^+$ T cell intrinsic effect, we adoptively transferred

naïve CD4$^+$ T cells from WT or *Gzmb*$^{-/-}$ donor mice into *Rag2*$^{-/-}$ *Gzmb*$^{-/-}$ mice. Donor CD4$^+$ T cells

from WT mice did not cause early weight loss in the double knock-out recipient mice, while CD4$^+$

T cells deficient in granzyme B caused similar weight loss as in *Rag2*$^{-/-}$ recipient mice (Figure 2a).

These results indicate a granzyme B-intrinsic effect in CD4$^+$ T cell pathogenesis. Weight loss in

mice receiving granzyme B-deficient cells was accompanied by other signs of disease, such as

diarrhea, pilo-erection, and hunching (Figure 2b). As expected, colon pathology based on immune

cell infiltration, loss of goblet cells, epithelial damage and hyperplasia was higher in mice

receiving CD4$^+$ T cells from *Gzmb*$^{-/-}$ donors (Figure 2c). Finally, granzyme B-deficient CD4$^+$ T

cells reconstituted the recipient mouse colons better than WT cells, regardless of the recipient mice

(Figure 2d).

In summary, our data indicate that absence of granzyme B in CD4$^+$ T cells activated *in vivo*

results in accelerated pathogenicity and increased cell reconstitution.

Granzyme B-deficient CD4$^+$ T cells present increased IL-17 production *in vivo*.

To determine whether granzyme B deficiency results in differential CD4$^+$ T cell cytokine

profiles *in vivo*, we adoptively transferred naïve CD4$^+$CD45RB^{hi} T cells from WT or *Gzmb$^{-/-}$* mice

into *Rag2$^{-/-}$* recipient mice. Three weeks after transfer, donor-derived cells were recovered from

the MLN and lamina propria and analyzed for IFNγ and IL-17 expression. As shown in Figure 3,

IFNγ was the predominant cytokine produced by donor CD4$^+$ T cells derived from WT and *Gzmb$^-$*

$^{/-}$ mice. In cells recovered from the MLN, there were similar levels of IFNγ production between

WT and *Gzmb$^{-/-}$* CD4$^+$ T cells (Figure 3a). However, CD4$^+$ T cells from *Gzmb$^{-/-}$* donor mice

presented significantly increased IL-17 production in comparison to WT CD4$^+$ T cells (Figure 3a).

A similar pattern was observed in donor-derived cells recovered from the lamina propria (Figure

3b).

In summary, *in vivo* activated CD4$^+$ T cells deficient in granzyme B presented increased

IL-17 production. These results suggest that absence of granzyme B results in skewed *in vivo*

differentiation.

Granzyme B deficient CD4$^+$ T cells present a distinct gene expression profile

Because granzyme B-deficient CD4$^+$ T cells had increased IL-17 production when

activated *in vivo*, we investigated whether these cells presented a distinct gene expression profile

in comparison to *in vivo* activated CD4$^+$ T cells from WT mice. For this purpose, CD4$^+$CD45RB^{hi}

T cells from WT and *Gzmb$^{-/-}$* donor mice were adoptively transferred into *Rag2$^{-/-}$* mice. Three

weeks after transfer, cells were recovered from the MLN, their RNA extracted and gene expression

profile determined by RNAseq. As shown in Figure 4a, WT and granzyme B-deficient CD4$^+$ T

cells presented significantly different gene expression profiles. We also performed gene set

enrichment analysis to determine whether WT and granzyme B-deficient CD4$^+$ T cells recovered

from recipient mice segregated in specific groups. As indicated in Figure 4b, GSEA indicated that

these two CD4[+] T cell populations presented distinct differential gene expression within several

GSEA groups (Figure 4b). These results indicate that granzyme B-deficient CD4[+] T cells, in the

adoptive transfer system, follow a distinct differentiation pathway compared to WT CD4[+] T cells.

We validated the results generated from the RNAseq analysis by comparing the expression

of selected genes in naïve CD4[+] T cells (Figure 4c, top), and donor-derived cells recovered 21d

post-adoptive transfer (Figure 4c, bottom). Hallmark genes involved in Th17 responses, such as

Rorc and *Il17a* were similarly expressed in naïve CD4[+] T cells, regardless of the granzyme B

status, but their expression had significantly increased in recovered granzyme B-deficient CD4[+] T

cells. *Ddit4*, a gene important for optimal T cell proliferation,[24] is highly expressed in T cells from

multiple sclerosis patients, and is important in Th17 differentiation.[25] *Ddit4* expression was 3-fold

higher in activated CD4[+] T cells derived from *Gzmb*[-/-] mice in comparison to WT CD4[+] T cells

(Figure 4c). Granzyme B-deficient CD4[+] T cells presented a 2-fold increase in *Il7r* expression

(Figure 4c), which mediates IL-7 signaling, a well-known lymphopoietic cytokine also expressed

in T cells undergoing homeostatic expansion.[26] Plexin D1, encoded by the *Plxnd1* gene, binds

semaphorins[27] and is critical for directing thymocyte migration.[28] Although the role of *Plxnd1* is

not well known in mature T cell biology, granzyme B-deficient CD4[+] T cells presented

approximately a 4-fold increase in expression. Interestingly, donor CD4[+] T cells derived from

Gzmb[-/-] mice presented decreased *Il10ra* expression in comparison to WT counterparts (Figure

4c).

Overall, these results show that in the adoptive transfer model of CD4[+] T cell activation,

granzyme B deficiency results in a distinct gene expression profile, characterized by genes relevant

to Th17 differentiation and T cell proliferation.

Granzyme B-deficient CD4$^+$ T cells possess better reconstitution capabilities

As indicated in Figure 2d, granzyme B-deficient CD4$^+$ T cells showed greater

reconstitution in the colon than WT CD4$^+$ T cells when transferred into Rag-2$^{-/-}$ recipient mice.

However, this observation could be due to inflammation-driven proliferation, which was more

prevalent in *Rag2$^{-/-}$* mice receiving CD4$^+$ T cells from *Gzmb$^{-/-}$* mice. To further determine the

reconstitution potential of CD4$^+$ T cells derived from WT and *Gzmb$^{-/-}$* mice, we performed

competitive adoptive T cell transfer experiments. For this purpose, naïve CD4$^+$ T cells from WT

CD45.1 and *Gzmb$^{-/-}$* CD45.2 mice (1x10^5 each) were adoptively co-transferred into the same

recipient mice. Of the total donor-derived cells obtained from the MLN, granzyme B-deficient

CD4$^+$ T cells presented ~60% reconstitution, whereas WT-derived cells showed ~40%

reconstitution (Figure 5a). Although the IFNγ profile of co-transferred cells was similar to cells

transferred alone, co-transferred WT cells presented increased IL-17 production in comparison to

WT CD4$^+$ T cells transferred alone (Figure 5b). These results raise the possibility that granzyme

B-deficient CD4$^+$ T cells influenced the behavior of granzyme-competent CD4$^+$ T cells. *Rag2$^{-/-}$*

mice co-transferred with WT and GzB$^{-/-}$ CD4$^+$ T cells displayed similar weight loss (Figure 5c)

and colon pathology (Figure 5d) to mice transferred only with CD4$^+$ T cells from *Gzmb$^{-/-}$* mice,

suggesting that the increased pathogenicity of the latter cells was maintained even in the presence

of granzyme-competent CD4$^+$ T cells.

Granzyme B-deficient mice present normal cellularity and IFNγ /IL-17 production

Since *in vivo* activated granzyme B-deficient CD4$^+$ T cells possess differential gene

expression and cytokine profiles, it is possible that these differences are also observed in naïve

cells. To investigate this possibility, we enumerated the CD4$^+$ T cell cellularity of spleen, MLN,

and IEL compartments from WT and *Gzmb*$^{-/-}$ mice. As indicated in Figure 6a, frequencies and

numbers of TCRβ$^+$CD4$^+$ T cells were similar among the two groups of mice. *Ex vivo* IFNγ and IL-

17 production by non-stimulated (ns) or stimulated (s) naïve CD4$^+$ T cells derived from the MLN

and IEL showed no distinguishable difference between cells from WT and *Gzmb*$^{-/-}$ mice (Figure

6b).

In summary, naïve WT and *Gzmb*$^{-/-}$ animals appear to have similar CD4$^+$ T cell cellularity

and basal IFNγ and IL-17 production in peripheral lymphoid organs and the intestines.

Granzyme B deficiency results in increased disease severity during *Citrobacter rodentium*

infection

As demonstrated in the previous section, naïve *Gzmb*$^{-/-}$ mice have a normal CD4$^+$ T cell

compartment compared to WT mice. However, because we observed that activated T cells from

Gzmb$^{-/-}$ mice became more pathogenic and had a distinct gene expression profile in the adoptive

transfer model of colitis, we investigated whether *Gzmb*$^{-/-}$ mice respond properly to antigenic

stimulus. For this purpose, we infected WT and *Gzmb*$^{-/-}$ mice with the extracellular bacterium

Citrobacter rodentium, which preferentially colonizes the colon of mice and induces an IL-17/IL-

22 immune response. After infection, WT mice maintained similar weight throughout the course

of the experiment; however, starting at 9 days post-infection, *Gzmb*$^{-/-}$ mice lost weight, reaching

approximately ~20% loss of the starting weight by day 12 (Figure 7a). *Gzmb*$^{-/-}$ mice also presented

other signs of disease, such as diarrhea, hunched posture, and pilo-erection, which were mostly

absent in WT mice (Figure 7b). Despite similar colon bacterial burden (Figure 7c) *Gzmb*$^{-/-}$ mice

developed higher pathology characterized by increased infiltration and epithelial injury than WT

mice (Figure 7d). Analysis of cytokine production showed that in the spleen, MLN and LP of

infected $Gzmb^{-/-}$ mice, there was a significant increase in CD4$^+$ T cells expressing IL-17, while

IFNγ production was increased only in the MLN (Figure 7e). Because innate immune cells are

important for the response against *C. rodentium*, we investigated whether the absence of granzyme

B in innate immune cells was responsible for the observed increased pathology in infected $Gzmb^{-/-}$

mice. For this purpose, $Rag-2^{-/-}$ and $Rag-2^{-/-}Gzmb^{-/-}$ mice were infected with *C. rodentium* and

monitored for 14 days. Both groups of mice maintained similar weights throughout the course of

the experiment (Supplemental Figure 2a), indicating that the effect caused by granzyme B-

deficiency is most likely associated with adaptive immune cells. To further confirm these results,

we treated $Rag-2^{-/-}$ and $Rag-2^{-/-}Gzmb^{-/-}$ mice with anti-CD40 antibodies, which induce acute

intestinal inflammation in the absence of T or B cells.[29] As shown in Supplemental Figure 2b, $Rag-$

$2^{-/-}$ and $Rag-2^{-/-}Gzmb^{-/-}$ mice presented similar weight loss throughout the course of the experiment.

Overall, these results recapitulate what was observed in the adoptive transfer model of

colitis, and indicate that the absence of granzyme B increased pathogenicity, which is associated

with adaptive immune responses.

Discussion

Granzyme B expression in CD4$^+$ T cells is well-documented, and has been primarily

studied in the context of granzyme B/perforin-dependent regulatory T cell suppressor activity.[30]

However, as our data indicate, granzyme B expression appears to be a common feature among a

significant fraction of activated CD4$^+$ T cells. Interestingly, not all differentiation conditions

induced granzyme B expression. For example, Th0 and Th1 cells rapidly express granzyme B

starting at 1-day post activation, whereas Th17 cells do not express this enzyme at any time after

activation. In the former conditions, expression of granzyme B is primarily driven by the presence

of irradiated splenocytes, indicating the possibility that cell-to-cell contact is important for

granzyme B expression.

Although none of the cytokines involved in Th1 differentiation induced granzyme B

expression above the levels observed with the addition of irradiated splenocytes, anti-IFNγ and

some cytokines present in Th17 cocktails, such as IL-6 and TGFβ prevented the expression of the

enzyme, which suggests that restriction of granzyme B expression is necessary for Th17

polarization. Studies in cytotoxic CD8$^+$ T cells have shown that activation of these cells with anti-

CD3 in the presence of IL-6 induces granzyme B expression, but not in the presence of TGF-β.[31]

Here we show that in CD4$^+$ T cell differentiation, IL-6 does not enhance the expression of

granzyme B, indicating that CD8$^+$ and CD4$^+$ T cells possess different mechanisms for modulating

granzyme B production.

Granzyme B expression may not only be limited during the initial CD4$^+$ T cell priming and

differentiation. For example, TGFβ is known for its role in regulatory T cell differentiation, and

some of these cells require granzyme B for their suppressor functions. However, as our results

indicate, TGFβ ablates expression of this enzyme, suggesting that regulatory T cells must acquire

granzyme B expression during post-priming events. Similarly, a fraction of effector CD4$^+$ T cells

that migrate into the intestinal intraepithelial lymphocyte compartment acquire granzyme B

expression after transcriptional reprogramming.[32]

One of the most intriguing questions is: Why do CD4$^+$ T cells express granzyme B during

Th0/Th1 activation and differentiation? During *in vitro* Th0/Th1 differentiation WT-gzmB^{pos} and

WT-gzmB^{neg} cells have a distinct IFNγ profile, suggesting that these cells may represent different

lineages, and the activity of granzyme B influences their outcome. This is better exemplified when

analyzing granzyme B-deficient CD4$^+$ T cells, where *in vivo* and *in vitro* activation skew them

towards an IL-17-producing phenotype, with a distinct gene expression signature. We postulate

that granzyme B-deficient IL-17$^+$ cells represent a lineage of cells that underwent aberrant

differentiation. Therefore, expression of granzyme B during Th0/Th1 activation serves as a

checkpoint for proper CD4$^+$ T cell differentiation, preventing IL-17 production and increased

pathology. These observations have important significance because it has been reported that allelic

variants of granzyme B correlate with autoimmune disorders,[33-35] raising the possibility that in

certain CD4$^+$ T cell-mediated disorders, lack of proper granzyme B function increases the

probability of improper differentiation and increased pathogenesis potential. If this idea is correct,

then WT-gzmB^{pos} CD4$^+$ T cells may represent cells in which the differentiation signals have the

potential to skew cells into an unwanted phenotype, and granzyme B expression in these cells

ensures the correct differentiation process.

Granzyme B activity in lymphocyte development and/or differentiation has been

previously reported.[36] These authors showed that granzyme A and B are important for the

development of a subset of TCR^{neg} intestinal intraepithelial lymphocytes known for the expression

of intracellular CD3γ. Development of these cells requires granzyme B to cleave and inactivate

the intracellular domain of Notch1. This raises the interesting possibility that during CD4$^+$ T cell

differentiation, granzyme B prevents unwanted phenotypes by disrupting signals coming from the

environment, such those provided by Notch and its ligands, which are known to influence CD4$^+$ T

cell differentiation.[37] Our observation that granzyme B expression was increased in the presence

of irradiated splenocytes, supports this hypothesis and we are currently investigating what signals

derived from APC enhance granzyme B expression.

It has been reported that granzyme B-deficient CD4$^+$CD25$^-$ T cells used in murine models

of graft versus host disease (GVHD), induced faster disease onset with increased lethality, in

comparison to their WT counterparts.[38] Although this group did not provide a mechanistic

explanation for the increased pathogenicity observed, they suggested that increased proliferation

and decreased cell death may be involved. According to our data, we believe that in the GVHD

model, differential proliferation and survival of granzyme B-deficient CD4$^+$ T cells are the result

of aberrant CD4$^+$ T cell differentiation, which allows escape of highly pathogenic clones. Further

investigations in the GVDH model are needed to test this hypothesis.

In summary, granzyme B has been known for its role in cell mediated-cytotoxicity, and

while many groups have reported potential extracellular roles for this enzyme,[39] here we present

in vitro and *in vivo* evidence supporting a novel intrinsic function for granzyme B during CD4$^+$ T

cell differentiation. Although many questions remain to be answered regarding how granzyme B

activity is involved in this process, our results provide a strong foundation for a better

understanding of the function of this enzyme in CD4$^+$ T cell biology.

Methods

Mice. C57BL/6J and C57BL/6J.CD45.1 mice were originally purchased from The Jackson

Laboratory (000664, and 002014 respectively) and have been maintained and acclimated in our

colony for several years. Granzyme B (*Gzmb*)$^{-/-}$ mice were kindly provided by Dr. Xuefang Cao.

CD57BL/6J *Rag2*$^{-/-}$ mice were kindly provided by Dr. Luc Van Kaer. *Rag2*$^{-/-}$*Gzmb*$^{-/-}$ were

generated in our colony. All mice were between 6 to 10 weeks of age at the time of

experimentation. Male and female mice were used for all experiments, except for *Citrobacter*

rodentium infection where 7-week-old females were used. Mice were maintained in accordance

with the Institutional Animal Care and Use Committee at Vanderbilt University.

Reagents and flow cytometry. Fluorochrome-coupled anti-mouse CD4 (GK1.5), CD8α (53-6.7),

CD45RB (C363.16A), granzyme B (NGZB), IFNγ (XMG1.2), IL-17a (TC11-18H10), TCRβ

(H57-597) and ghost viability dyes were purchased from ThermoFisher, BD Bioscences or Tonbo.

Biotinylated anti-CD8α (53-1.7) and CD19 (1D3) antibodies were purchased from Tonbo. Anti-

CD40 antibody was purchased from BioXcell. Surface cell staining was performed following

conventional techniques. For intracellular cytokine staining, cells were stimulated with PMA and

ionomycin in the presence of Golgi Stop (BD Biosciences) for 4hr prior to staining. Extracellular

markers were stained, cells were fixed briefly with 2% paraformaldehyde, followed by

permeabilization and intracellular staining using the BD Cytofix/Cytoperm kit, according to

manufacturer's instructions. All stained samples were acquired using BD FACS Canto II, 3- or 4-

Laser Fortessa, or 5-Laser LSR II flow cytometers (BD Biosciences). Data were analyzed using

FlowJo software (Tree Star).

Lymphocyte isolation. Spleen and MLN lymphocytes were isolated by conventional means. IEL

and LP cells were isolated by mechanical disruption as previously reported.[40] Briefly, after

flushing the intestinal contents with cold HBSS and removing excess mucus, the intestines were

cut into small pieces (~1 cm long) and shaken for 45 minutes at 37°C in HBSS supplemented with

5% fetal bovine serum and 2 mM EDTA. Supernatants were recovered and cells isolated using a

discontinuous 40/70% Percoll (General Electric) gradient. To obtain lamina propria lymphocytes,

intestinal tissue was recovered and digested with collagenase (187.5 U/ml, Sigma) and DNase I

(0.6 U /ml, Sigma). Cells were isolated using a discontinuous 40/70% Percoll gradient.

In vitro *CD4 T cell activation/differentiation.* Naïve CD4 T cells were isolated from the spleens of

the indicated mice by magnetic sorting using the Miltenyi or StemCell naïve CD4$^+$ T cell isolation

kits according to the manufacturer's instructions. Splenocytes or MLN cells used as APC were

incubated for 1hr at 37°C, non-adherent cells were removed, adherent cells were washed and

resuspended in complete RPMI, and irradiated with 7 Gy (700 rads). 5x10^5 CD4$^+$ T cells with or

without 5x10^5 irradiated APC were incubated in the presence of plate bound anti-CD3 (5μg/ml)

and soluble anti-CD28 (2.5μg/ml), for the indicated times, under the following conditions: Th0

(10ng/mL 1L-2), Th1 (10ng/mL 1L-2, 20ng/mL IL-12, and 10μg/mL anti-IL-4) , and Th17

(5ng/mL TGFβ, 20ng/mL IL-6, 10ng/mL IL-1β, 10ng/mL IL-23, 10μg/mL anti-IL-4, and

10μg/mL anti-IFNγ).

Adoptive transfer of naïve CD4 T cells. Splenocytes from WT or *Gzmb*$^{-/-}$ mice were depleted of

CD19$^+$ and CD8α$^+$ cells using magnetic bead sorting (Miltenyi). Enriched cells were stained with

antibodies directed against TCRβ, CD4 and CD45RB, and naïve CD4$^+$ T cells were sorted as

CD4$^+$CD45RB^{hi} using a FACSAria III at the Flow Cytometry Shared Resource at VUMC. 1x10^5

cells were adoptively transferred into the indicated recipient mice. Starting weight was determined

prior to injection. Mice were monitored and weighed weekly. At the end time point, mice were

sacrificed and donor-derived cells were isolated from the indicated organs or tissues. Colon

histopathology was performed in a blind fashion by a GI pathologist (MBP), following established

parameters.[41] For competitive transfer experiments, 1x10^5 CD4$^+$CD45RB^{hi} cells from both WT-

CD45.1 and *Gzmb*[-/-] CD45.2 mice were adoptively co-transferred into *Rag2*[-/-] mice and analyzed

as indicated above.

Transcription profile analysis. For gene expression analysis, TCRβ⁺CD4⁺ T cells were purified by

flow cytometry from a pool of MLN cells comprised of 1 male and 1 female animal with similar

disease progression 21 days after adoptive transfer. RNA was isolated using the RNeasy micro kit

(Qiagen). cDNA library preparation and RNAseq was performed by the VANTAGE Core at

VUMC on an Illumina NovaSeq 6000 (2 x 150 base pair, paired-end reads). The tool Salmon[42]

was used for quantifying the expression of RNA transcripts. The R project software along with the

edgeR method[43] was used for differential expression analysis. For gene set enrichment analysis

(GSEA), RNAseq data was ranked according to the t-test statistic. The gene sets curated (C2), GO

(C5), immunological signature collection (C7) and hallmarks of cancer (H) of the Molecular

Signatures Database (MSigDB) were used for enrichment analysis. GSEA enrichment plots were

generated using the GSEA software[44] from the Broad Institute with 1000 permutations.

Real time PCR. For *in vitro* analysis, RNA was extracted from MACS-enriched, differentiated

CD4⁺ T cells at day 1 and 3 of culture. For *in vivo* analysis, RNA was isolated from flow cytometry

purified donor-derived TCRβ⁺CD4⁺ T cells as described above. cDNA was synthesized using the

RT2 First Strand kit (Qiagen). Real time PCR was performed using RT2 SYBR Green Mastermix

(Qiagen) on an Applied BioSystems Proflex PCR cycler. All primers were purchased from Qiagen.

Citrobacter rodentium *infection.* Seven-week old female WT and *Gzmb*[-/-] animals were infected

with *C. rodentium* as previously described[45]. Briefly, mice were infected with 5x10⁸ CFU

exponentially grown bacteria by oral gavage. Starting weight was determined prior to gavage.

Mice were monitored and weighed daily for 14d. At the end time point, cells were isolated from

the indicated organs or tissues. Cellularity was determined, intracellular cytokine staining was

performed, and colon histopathology was performed in a blind fashion by a GI pathologist (MBP)

as previously described.[46]

Anti-CD40 disease induction. *Rag-2*[-/-] and *Rag-2*[-/-]*Gzmb*[-/-] mice were injected i.p. with

150μg/mouse of anti-CD40 (BioXcell). Mice were monitored and weighed for daily for 7 days.

Acknowledgments

This work was supported by NIH grant R01DK111671 (D.O-V.); National Library of Medicine

T15 LM00745 (M.J.G.); Vanderbilt Training in Cellular, Biochemical and Molecular Sciences

Training Program, T32 GM008554-25 (K.G.M.); Vanderbilt Training in Gastroenterology T32

grant NIH T32 DK007673 (A.N.); and the Digestive Disease Research Center at Vanderbilt

University Medical Center, NIH grant P30DK058404 (M.B.P., K.T.W., and D.O-V.). We thank

the Flow Cytometry Shared Resource and the Vanderbilt Technologies for Advanced Genomics

(VANTAGE) for their invaluable support.

Author contributions: K.L.H., and D.O-V. designed research; K.L.H., M.J.G., K.M., A.N., and

K.S. performed research; K.L.H., M.J.G., M.B.P. and D.O-V. analyzed data; K.S. and K.T.W.

provided expertise with *Citrobacter rodentium* infections; K.L.H. and D.O-V. wrote the

manuscript.

Competing interests: The authors declare no competing interests.

References

1.   Anthony DA, Andrews DM, Watt SV, Trapani JA, Smyth MJ. Functional dissection
of the granzyme family: cell death and inflammation. *Immunol Rev* 2010; **235**(1)**:**
73-92.

458
459   2.   Chowdhury D, Lieberman J. Death by a thousand cuts: granzyme pathways of
460   programmed cell death. *Annu Rev Immunol* 2008; **26:** 389-420.

461
462   3.   Afonina IS, Cullen SP, Martin SJ. Cytotoxic and non-cytotoxic roles of the CTL/NK
463   protease granzyme B. *Immunol Rev* 2010; **235**(1)**:** 105-116.

464
465   4.   Wensink AC, Hack CE, Bovenschen N. Granzymes regulate proinflammatory
466   cytokine responses. *J Immunol* 2015; **194**(2)**:** 491-497.

467
468   5.   Kumar AA, Delgado AG, Piazuelo MB, Van Kaer L, Olivares-Villagomez D. Innate
469   CD8alphaalpha(+) lymphocytes enhance anti-CD40 antibody-mediated colitis in
470   mice. *Immun Inflamm Dis* 2017; **5**(2)**:** 109-123.

471
472   6.   Parkinson LG, Toro A, Zhao H, Brown K, Tebbutt SJ, Granville DJ. Granzyme B
473   mediates both direct and indirect cleavage of extracellular matrix in skin after
474   chronic low-dose ultraviolet light irradiation. *Aging Cell* 2015; **14**(1)**:** 67-77.

475
476   7.   Chamberlain CM, Ang LS, Boivin WA, Cooper DM, Williams SJ, Zhao H *et al.*
477   Perforin-independent extracellular granzyme B activity contributes to abdominal
478   aortic aneurysm. *Am J Pathol* 2010; **176**(2)**:** 1038-1049.

479
480   8.   Lancki DW, Hsieh CS, Fitch FW. Mechanisms of lysis by cytotoxic T lymphocyte
481   clones. Lytic activity and gene expression in cloned antigen-specific CD4+ and
482   CD8+ T lymphocytes. *J Immunol* 1991; **146**(9)**:** 3242-3249.

483
484   9.   Ebnet K, Chluba-de Tapia J, Hurtenbach U, Kramer MD, Simon MM. In vivo primed
485   mouse T cells selectively express T cell-specific serine proteinase-1 and the
486   proteinase-like molecules granzyme B and C. *Int Immunol* 1991; **3**(1)**:** 9-19.

487
488   10.   Sun Q, Burton RL, Pollok KE, Emanuel DJ, Lucas KG. CD4(+) Epstein-Barr virus-
489   specific cytotoxic T-lymphocytes from human umbilical cord blood. *Cell Immunol*
490   1999; **195**(2)**:** 81-88.

491
492   11.   Canaday DH, Wilkinson RJ, Li Q, Harding CV, Silver RF, Boom WH. CD4(+) and
493   CD8(+) T cells kill intracellular Mycobacterium tuberculosis by a perforin and
494   Fas/Fas ligand-independent mechanism. *J Immunol* 2001; **167**(5)**:** 2734-2742.

12. Yasukawa M, Ohminami H, Yakushijin Y, Arai J, Hasegawa A, Ishida Y *et al.* Fas-independent cytotoxicity mediated by human CD4+ CTL directed against herpes simplex virus-infected cells. *J Immunol* 1999; **162**(10)**:** 6100-6106.

13. Dorothee G, Vergnon I, Menez J, Echchakir H, Grunenwald D, Kubin M *et al.* Tumor-infiltrating CD4+ T lymphocytes express APO2 ligand (APO2L)/TRAIL upon specific stimulation with autologous lung carcinoma cells: role of IFN-alpha on APO2L/TRAIL expression and -mediated cytotoxicity. *J Immunol* 2002; **169**(2)**:** 809-817.

14. Grossman WJ, Verbsky JW, Tollefsen BL, Kemper C, Atkinson JP, Ley TJ. Differential expression of granzymes A and B in human cytotoxic lymphocyte subsets and T regulatory cells. *Blood* 2004; **104**(9)**:** 2840-2848.

15. Koonpaew S, Shen S, Flowers L, Zhang W. LAT-mediated signaling in CD4+CD25+ regulatory T cell development. *J Exp Med* 2006; **203**(1)**:** 119-129.

16. Kawamura K, Kadowaki N, Kitawaki T, Uchiyama T. Virus-stimulated plasmacytoid dendritic cells induce CD4+ cytotoxic regulatory T cells. *Blood* 2006; **107**(3)**:** 1031-1038.

17. Price JD, Schaumburg J, Sandin C, Atkinson JP, Lindahl G, Kemper C. Induction of a regulatory phenotype in human CD4+ T cells by streptococcal M protein. *J Immunol* 2005; **175**(2)**:** 677-684.

18. Gondek DC, Lu LF, Quezada SA, Sakaguchi S, Noelle RJ. Cutting edge: contact-mediated suppression by CD4+CD25+ regulatory cells involves a granzyme B-dependent, perforin-independent mechanism. *J Immunol* 2005; **174**(4)**:** 1783-1786.

19. Zhu J, Yamane H, Paul WE. Differentiation of effector CD4 T cell populations (*). *Annu Rev Immunol* 2010; **28:** 445-489.

20. Sharma V, Delgado M, Ganea D. Granzyme B, a new player in activation-induced cell death, is down-regulated by vasoactive intestinal peptide in Th2 but not Th1 effectors. *J Immunol* 2006; **176**(1)**:** 97-110.

21. Coomes SM, Kannan Y, Pelly VS, Entwistle LJ, Guidi R, Perez-Lloret J *et al.* CD4(+) Th2 cells are directly regulated by IL-10 during allergic airway inflammation. *Mucosal immunology* 2017; **10**(1)**:** 150-161.

22. Powrie F, Correa-Oliveira R, Mauze S, Coffman RL. Regulatory interactions between CD45RBhigh and CD45RBlow CD4+ T cells are important for the balance between protective and pathogenic cell-mediated immunity. *J Exp Med* 1994; **179**(2)**:** 589-600. PMC2191378.

542   23.   Powrie F, Leach MW, Mauze S, Menon S, Caddle LB, Coffman RL. Inhibition of
543   Th1 responses prevents inflammatory bowel disease in scid mice reconstituted
544   with CD45RBhi CD4+ T cells. *Immunity* 1994; **1**(7)**:** 553-562.
545

546   24.   Reuschel EL, Wang J, Shivers DK, Muthumani K, Weiner DB, Ma Z *et al.* REDD1
547   Is Essential for Optimal T Cell Proliferation and Survival. *PLoS One* 2015; **10**(8)**:**
548   e0136323.
549

550   25.   Zhang F, Liu G, Li D, Wei C, Hao J. DDIT4 and Associated lncDDIT4 Modulate
551   Th17 Differentiation through the DDIT4/TSC/mTOR Pathway. *J Immunol* 2018;
552   **200**(5)**:** 1618-1626.
553

554   26.   Fry TJ, Mackall CL. The many faces of IL-7: from lymphopoiesis to peripheral T
555   cell maintenance. *J Immunol* 2005; **174**(11)**:** 6571-6576.
556

557   27.   Carvalheiro T, Rafael-Vidal C, Malvar-Fernandez B, Lopes AP, Pego-Reigosa JM,
558   Radstake T *et al.* Semaphorin4A-Plexin D1 Axis Induces Th2 and Th17 While
559   Represses Th1 Skewing in an Autocrine Manner. *Int J Mol Sci* 2020; **21**(18).
560

561   28.   Choi YI, Duke-Cohan JS, Ahmed WB, Handley MA, Mann F, Epstein JA *et al.*
562   PlexinD1 glycoprotein controls migration of positively selected thymocytes into the
563   medulla. *Immunity* 2008; **29**(6)**:** 888-898.
564

565   29.   Uhlig HH, McKenzie BS, Hue S, Thompson C, Joyce-Shaikh B, Stepankova R *et*
566   *al.* Differential activity of IL-12 and IL-23 in mucosal and systemic innate immune
567   pathology. *Immunity* 2006; **25**(2)**:** 309-318.
568

569   30.   Cao X, Cai SF, Fehniger TA, Song J, Collins LI, Piwnica-Worms DR *et al.*
570   Granzyme B and perforin are important for regulatory T cell-mediated suppression
571   of tumor clearance. *Immunity* 2007; **27**(4)**:** 635-646.
572

573   31.   Huber M, Heink S, Grothe H, Guralnik A, Reinhard K, Elflein K *et al.* A Th17-like
574   developmental process leads to CD8(+) Tc17 cells with reduced cytotoxic activity.
575   *Eur J Immunol* 2009; **39**(7)**:** 1716-1725.
576

577   32.   Mucida D, Husain MM, Muroi S, van Wijk F, Shinnakasu R, Naoe Y *et al.*
578   Transcriptional reprogramming of mature CD4(+) helper T cells generates distinct
579   MHC class II-restricted cytotoxic T lymphocytes. *Nat Immunol* 2013; **14**(3)**:** 281-
580   289.
581

582   33.   Jin Y, Birlea SA, Fain PR, Gowan K, Riccardi SL, Holland PJ *et al.* Variant of TYR
583   and autoimmunity susceptibility loci in generalized vitiligo. *The New England*
584   *journal of medicine* 2010; **362**(18)**:** 1686-1697.
585

586 34. Li S, Yao W, Pan Q, Tang X, Zhao S, Wang W *et al.* Association analysis revealed
587 one susceptibility locus for vitiligo with immune-related diseases in the Chinese
588 Han population. *Immunogenetics* 2015; **67**(7)**:** 347-354.
589

590 35. Shen C, Gao J, Sheng Y, Dou J, Zhou F, Zheng X *et al.* Genetic Susceptibility to
591 Vitiligo: GWAS Approaches for Identifying Vitiligo Susceptibility Genes and Loci.
592 *Front Genet* 2016; **7:** 3.
593

594 36. Ettersperger J, Montcuquet N, Malamut G, Guegan N, Lopez-Lastra S, Gayraud S
595 *et al.* Interleukin-15-Dependent T-Cell-like Innate Intraepithelial Lymphocytes
596 Develop in the Intestine and Transform into Lymphomas in Celiac Disease.
597 *Immunity* 2016; **45**(3)**:** 610-625.
598

599 37. Amsen D, Helbig C, Backer RA. Notch in T Cell Differentiation: All Things
600 Considered. *Trends Immunol* 2015; **36**(12)**:** 802-814.
601

602 38. Du W, Leigh ND, Bian G, O'Neill RE, Mei L, Qiu J *et al.* Granzyme B-Mediated
603 Activation-Induced Death of CD4+ T Cells Inhibits Murine Acute Graft-versus-Host
604 Disease. *J Immunol* 2015; **195**(9)**:** 4514-4523.
605

606 39. Boivin WA, Cooper DM, Hiebert PR, Granville DJ. Intracellular versus extracellular
607 granzyme B in immunity and disease: challenging the dogma. *Laboratory*
608 *investigation; a journal of technical methods and pathology* 2009; **89**(11)**:** 1195-
609 1220.
610

611 40. Nazmi A, Greer MJ, Hoek KL, Piazuelo MB, Weitkamp J, Olivares-Villagomez D.
612 Osteopontin and iCD8alpha cells promote intestinal intraepithelial lymphocyte
613 homeostasis. *J Immunol* 2020; **In press**.
614

615 41. Aranda R, Sydora BC, McAllister PL, Binder SW, Yang HY, Targan SR *et al.*
616 Analysis of intestinal lymphocytes in mouse colitis mediated by transfer of CD4+,
617 CD45RBhigh T cells to SCID recipients. *J Immunol* 1997; **158**(7)**:** 3464-3473.
618

619 42. Patro R, Duggal G, Love MI, Irizarry RA, Kingsford C. Salmon provides fast and
620 bias-aware quantification of transcript expression. *Nature methods* 2017; **14**(4)**:**
621 417-419.
622

623 43. Robinson MD, McCarthy DJ, Smyth GK. edgeR: a Bioconductor package for
624 differential expression analysis of digital gene expression data. *Bioinformatics*
625 2010; **26**(1)**:** 139-140.
626

627 44. Subramanian A, Tamayo P, Mootha VK, Mukherjee S, Ebert BL, Gillette MA *et al.*
628 Gene set enrichment analysis: a knowledge-based approach for interpreting
629 genome-wide expression profiles. *Proc Natl Acad Sci U S A* 2005; **102**(43)**:** 15545-
630 15550.
631

632   45.   Olivares-Villagomez D, Algood HM, Singh K, Parekh VV, Ryan KE, Piazuelo MB
633   *et al.* Intestinal epithelial cells modulate CD4 T cell responses via the thymus
634   leukemia antigen. *J Immunol* 2011; **187**(8)**:** 4051-4060.
635
636   46.   Singh K, Gobert AP, Coburn LA, Barry DP, Allaman M, Asim M *et al.* Dietary
637   Arginine Regulates Severity of Experimental Colitis and Affects the Colonic
638   Microbiome. *Front Cell Infect Microbiol* 2019; **9:** 66.
639
640
641

Figure legends

Figure 1. Granzyme B expression during T cell differentiation. (a) Naïve CD4$^+$ T cells from WT mice were cultured in the presence of plate-bound anti-CD3 and soluble CD28 under the indicated differentiation conditions, and in the absence of irradiated splenocytes. At day 1 (top) and 3 (bottom), cells were recovered, RNA extracted and granzyme B mRNA expression determined. (b) Intracellular granzyme B in cells cultured as in (a), but in the presence of 5x10^5 irradiated splenocytes. Left, representative dot plot; live, TCRβ$^+$CD4$^+$ cells are displayed. Right, data summary. Each dot represents an individual sample. (c) Granzyme B expression of cells cultured in Th1 and Th17 conditions in the presence of irradiated splenocytes at different time points. (d) Cells cultured in Th0 conditions in the presence of the indicated individual cytokines, harvested 5 days after culture. (e) Cells cultured as indicated in (d) in the presence or absence of irradiated APC from spleen. Intracellular IFNγ and IL-17 staining of representative samples cultured in (f) Th0 or (g) Th1 conditions in the presence of irradiated splenocytes. Top, intracellular IFNγ and IL-17 staining representative samples; bottom, data summary. Each dot represents an individual sample. Data are representative of at least 3 independent experiments, performed in triplicate. n=6. For (e), ***P<0.001, Student's t test; for (g) and (h), *P<0.05; **P<0.01; ***P<0.001; ****P<0.001, One-way ANOVA.

Supplemental Figure 1. Granzyme B and cytokine expression in the presence of 5x10^5 irradiated spleen or MLN APC. Naïve CD4$^+$ T cells from WT mice were cultured in the presence of plate-bound anti-CD3 and soluble anti-CD28 under Th1 differentiation conditions. (a) Granzyme B expression 5 days after culture. (b) IFNγ and (c) IL-17 production from the same cultures. (d) IFNγ and (e) IL-17 expression from granzyme B-deficient CD4$^+$ T cells under similar cultures as in (a). Each dot represents an individual sample. Data are representative of at least 2 independent experiments, performed in triplicate. n=3-6. *P<0.05; **P<0.01; ***P<0.001. For (a) and (e), Student's t-test; for (b) and (c), One-way ANOVA.

Figure 2. *In vivo* activated granzyme B-deficient CD4$^+$ T cells present increased pathogenicity. 1x10^5 CD4$^+$CD45RB^{hi} T cells from WT or *Gzmb$^{-/-}$* donor mice were adoptively transferred into *Rag2$^{-/-}$* or *Rag2$^{-/-}$Gzmb$^{-/-}$* recipient mice. (a) Mice were weighed weekly, and (b) monitored for

signs of disease. Twenty-one days after transfer, (c) colon pathology was scored (left,

representative micrographs, 100X magnification; right, data summary), and (d) donor-cell

reconstitution of the IEL compartment in the colon was determined. For (a), dotted lines

represent untreated *Rag2*$^{-/-}$ and *Rag-2*$^{-/-}$*Gzmb*$^{-/-}$ mice. Each dot represents an individual mouse.

Data are representative of at least 3 independent experiments. n=8-10. For (a), ***P<0.001;

****P<0.0001; Student's t-test for each time point. For (b), # compares WT > *Rag2*$^{-/-}$ to *Gzmb*$^{-/-}$

> *Rag2*$^{-/-}$ and WT > *Gzmb*$^{-/-}$*Rag2*$^{-/-}$ at 14 days, and ## compares WT > *Rag2*$^{-/-}$ to *Gzmb*$^{-/-}$ > *Rag2*$^{-}$

$^{/-}$ and WT > *Gzmb*$^{-/-}$*Rag2*$^{-/-}$ at 21 days; Two-way ANOVA. For (c) and (d), *P<0.05;

***P<0.001; One-way ANOVA.

Figure 3. Granzyme B-deficient CD4$^+$ T cells present skewed IL-17 differentiation *in vivo*.

1x10^5 CD4$^+$CD45RB^{hi} T cells from WT or *Gzmb*$^{-/-}$ donor mice were adoptively transferred into

Rag2$^{-/-}$ recipient mice. Three weeks after transfer, donor-derived cells were recovered from (a)

MLN, and (b) lamina propria, and their IFNγ/IL-17 profile was determined by intracellular

staining. Dot plots indicate a representative sample. Live, TCRβ$^+$CD4$^+$ cells are displayed.

Graphs represent the summary. Each dot indicates an individual mouse. For MLN, n=11-13; for

lamina propria, n=7-11. Data are representative of at least 3 independent experiments. **P<0.01;

****P<0.001. Student's t-Test.

Figure 4. Granzyme B-deficient CD4$^+$ T cells activated *in vivo* present a distinct gene expression

profile. 1x10^5 CD4$^+$CD45RB^{hi} T cells from WT or *Gzmb*$^{-/-}$ donor mice were adoptively

transferred into *Rag2*$^{-/-}$ recipient mice. Three weeks after transfer, donor-derived cells were

recovered from the MLN, sorted for live TCRβ$^+$CD4$^+$, their RNA isolated and sequenced. (a)

Representative heat map. (b) Representative gene set enrichment analysis. (c) Real-time PCR

validation of selected genes. n=6-7. *P<0.05; **P<0.01. Student's t-Test.

Figure 5. Granzyme B-deficient CD4$^+$ T cells present greater *in vivo* reconstitution.

1x10^5 CD4$^+$CD45RB^{hi} T cells from WT (CD45.1) and *Gzmb*$^{-/-}$ (CD45.2) donor mice were co-

transferred into the same *Rag2*$^{-/-}$ recipient mice. As controls, cells from the same donors were

independently transferred into *Rag2*$^{-/-}$ recipient mice. Three weeks after transfer, (a) MLN

reconstitution (right, representative dot plot displaying live, TCRβ$^+$CD4$^+$ cells; left, summary),

and (b) IFNγ/IL-17 production was determined. (c) *Rag2*[-/-] mice that received CD4[+] T cells from

both WT/*Gzmb*[-/-] or only *Gzmb*[-/-] mice were weighed weekly. (d) Colon pathology was

determined 3 weeks post transfer. Each dot represents an individual mouse. For (a) to (c), n=8-

10; for (e) to (g), n=5-11. Data are representative of at least 2 independent experiments.

**P<0.01; Student's t-Test.

Figure 6. Naïve granzyme B-deficient mice present a normal CD4[+] T cell compartment. (a)

CD4[+] T cell frequencies and cellularity from spleen, MLN, and IEL compartment were

determined by flow cytometry. (b) *Ex vivo* IFNγ and IL-17 production from CD4[+] T cells derived

from naïve WT and *Gzmb*[-/-] mice. Cells were non-stimulated (ns) or stimulated (s) for 4 h with

PMA/ionomycin. Each dot represents an individual mouse. For (a), spleen and MLN, n=6-10; for

IEL, n=12-13. For (b), n=6.

Figure 7. Granzyme B-deficient mice present severe disease after *C. rodentium* infection.

Indicated mice were orogastrically infected with *C. rodentium* ($5x10^8$ CFU/mouse) and

monitored for (a) weight change, and (b) signs of disease. At 14 days post infection, colons were

dissected, (c) colonization/g of colon was determined, and (d) pathology scored as indicated in

the Methods section; left, representative micrographs (200X magnification); right, data

summary. (e) CD4[+] T cell IFNγ and IL-17 expression in the indicated organs was determined.

Cells were non-stimulated (ns) or stimulated (s) for 4 h with PMA/ionomycin. Each dot

represents an individual mouse. For (a-d), n=7-9; for (e), 3-5. Data are representative of at least 3

independent experiments. For (a): **P<0.01; ***P<0.01; ****P<0.001; Student's t-Test for each

time point. For (b): ****P<0.001; Student's t-Test for each time point. For (d) and (e): *P<0.05;

**P<0.01; One-way ANOVA.

Supplemental Figure 2. (a) Absence of adaptive immune cells rescues granzyme B-deficient

mice from severe disease. The indicated mice were orogastrically infected with *C. rodentium*

($5x10^8$ CFU/mouse) and monitored for weight change for 14 days. Data are representative of at

least 2 independent experiments. n=3-4 for sham groups. n=5-7 for infected mice. (b) Granzyme

B deficiency does not alter disease progression in the anti-CD40 model of colitis. The indicated

mice were treated i.p. with 150 μg/mouse and weighed daily. Data are representative of at least 3

independent experiments. n=2 for sham groups. n=14 for treated group.

744
745

Figure 2

a

TNF family | Chemokines | Cell survival | Adhesion

WT

Gzmb⁻/⁻

Transcription factors | Innate response | Cytokine receptors | Ag presentation

WT

Gzmb⁻/⁻

b

Enrichement score (ES)

IL4 and IL12 response | Naive vs memory | Unstimulated vs null | T conventional vs Treg

c

Pre-transfer

Ddit4 | *IL7r* | *IL10ra* | *IL17a* | *Rorc* | *Plxnd1*

Fold increase — WT, *Gzmb⁻/⁻*

Post-transfer

Fold increase — WT, *Gzmb⁻/⁻*

** | * | * | ** | * | **

Figure 4

Figure 5

a

b

Figure 6

Figure 7