Title: Granzyme B prevents aberrant IL-17 production and intestinal pathogenicity in CD4[+] T cells

Authors: Kristen L. Hoek, PhD [a]; Michael J. Greer, PhD [b*]; Kathleen G. McClanahan,[a*] Ali Nazmi

[a], PhD; M. Blanca Piazuelo, MD[c,e]; Kshipra Singh, PhD[c,e]; Keith T. Wilson, MD[a,c,e,f,g]; and

Danyvid Olivares-Villagómez, PhD[a,e,g,h].

*These two authors contributed equally to this report

Affiliations: [a]Department of Pathology, Microbiology and Immunology, Vanderbilt University

Medical Center, A5301 Medical Center North, Nashville, Tennessee 37232, USA; [b]Department of

Biomedical Informatics, Vanderbilt University, Nashville, Tennessee, 37232, USA; [c]Division of

Gastroenterology, Hepatology and Nutrition, Department of Medicine, Vanderbilt University

Medical Center, Nashville, Tennessee 37232, USA; [e]Center for Mucosal Inflammation and

Cancer, Vanderbilt University Medical Center, Nashville, Tennessee 37232; [f]Veterans Affairs

Tennessee Valley Healthcare System, Nashville, Tennessee 37232; [g]Vanderbilt Institute for

Infection, Immunology and Inflammation, Vanderbilt University Medical Center, A5301 Medical

Center North, Nashville, Tennessee 37232, USA. [h]ORCID, 0000-0002-1158-8976

Corresponding author: Danyvid Olivares-Villagómez. Address: Vanderbilt University Medical

center, A5211 Medical Center North, Nashville Tennessee 37232-0001. Fax number: 615-343-

7392. Telephone number: 615-936-0134. email: danyvid.olivares-villagomez@vumc.org.

Conflict of Interest: No conflict

Author contributions: K.L.H., and D.O-V. designed research; K.L.H., M.J.G., K.M., A.N., and

K.S. performed research; K.L.H., M.J.G., M.B.P. and D.O-V. analyzed data; K.S. and K.T.W.

provided expertise with *Citrobacter rodentium* infections; K.L.H. and D.O-V. wrote the

manuscript.

Acknowledgments

This work was supported by NIH grant R01DK111671 (D.O-V.); National Library of Medicine

T15 LM00745 (M.J.G.); Vanderbilt Training in Cellular, Biochemical and Molecular Sciences

Training Program, T32 GM008554-25 (K.G.M.); Vanderbilt Training in Gastroenterology T32

grant NIH T32 DK007673 (A.N.); and the Digestive Disease Research Center at Vanderbilt

University Medical Center, NIH grant P30DK058404 (M.B.P., K.T.W., and D.O-V.). Flow

cytometry experiments were performed in the VMC Flow Cytometry Shared Resource, supported

by the Vanderbilt Ingram cancer Center (P30 CA068485) and the Vanderbilt Digestive Disease

Research Center (DK058404). We acknowledge the Translational Pathology Shared Resource

supported by NCI/NIH cancer Center Support Grant 5P30 CA68485-19 and the Mouse Metabolic

Phenotyping Center Grant 2 U24 DK059637-16. We also thank the Vanderbilt Technologies for

Advanced Genomics (VANTAGE) for their invaluable support.

Key words: Granzyme B; CD4$^+$ T cells; T cell differentiation; IL-17; intestinal inflammation.

Abstract: CD4[+] T cell activation and differentiation are important events that set the stage for

proper immune responses. Many factors are involved in the activation and differentiation of T

cells, and these events are tightly controlled to prevent unwanted and/or exacerbated immune

responses that may harm the host. It has been well documented that granzyme B, a potent serine

protease involved in cell-mediated cytotoxicity, is readily expressed by certain CD4[+] T cells, such

as regulatory T cells and CD4[+]CD8$\alpha\alpha$[+] intestinal intraepithelial lymphocytes, both of which

display cytotoxicity associated with granzyme B. However, because not all CD4[+] T cells

expressing granzyme B are cytotoxic, additional roles for this protease in CD4[+] T cell biology

remain unknown. Here, using a combination of *in vivo* and *in vitro* approaches, we report that

granzyme B-deficient CD4[+] T cells display increased IL-17 production. In the adoptive transfer

model of intestinal inflammation, granzyme B-deficient CD4[+] T cells triggered a more rapid

disease onset than their WT counterparts, and presented a differential transcription profile. Similar

results were also observed in granzyme B-deficient mice infected with *Citrobacter rodentium*. Our

results suggest that granzyme B modulates CD4[+] T cell differentiation, providing a new

perspective into the biology of this enzyme.

Introduction

Granzymes are a family of serine proteases expressed by many immune cell populations.

There are five different granzymes in humans (A, B, H, K and M) and 11 in mice (A, B, C, D, E,

F, G, K, L, M and N), which differ in function and substrate specificity.[1] Most research has focused

on granzymes A and B due to their hallmark role in CD8[+] T and NK cell-mediated cytotoxicity

against tumor cells or virus-infected cells.[2] However, mounting evidence indicates that granzymes

are also involved in other processes, including inflammatory responses,[3-5] remodeling of the

extracellular matrix,[6] and cardiovascular disorders.[7]

Activated CD4[+] T cells express granzyme B, and some of these cells exhibit lytic activity

similar to CD8[+] T cells.[8, 9] Antigen-specific CD4[+] T cell lytic activity mediated by granzyme B

can be triggered by viral and bacterial infections.[10-12] Similarly, some tumor-infiltrating CD4[+] T

cells also present anti-tumor cytotoxicity mediated by granzyme B.[13] In addition, one of the

mechanisms by which regulatory CD4[+] T cells in humans and mice control immune responses is

through granzyme B-mediated cell death.[14-18]

After activation, CD4[+] T cells can differentiate into a diverse set of T helper (Th) cells

depending on environmental cues and transcription factors, which ultimately leads Th cells to

produce a specific cytokine profile.[19] In certain populations of differentiated CD4[+] T cells,

granzyme B controls CD4[+] T cell immune responses by initiating activation-induced cell death.[20]

Downregulation of granzyme B, for example when Th2 cells are cultured in the presence of

vasoactive intestinal peptide, reduces Th2 cell death.[20] Moreover, granzyme B expression in Th2

cells is enhanced by IL-10R signaling, which results in cell death.[21] These studies clearly indicate

that granzyme B is involved in controlling some immune CD4[+] T cell responses. However, it is

unclear whether this enzyme is actively involved during the process of Th differentiation.

In this report, we present data indicating that granzyme B is expressed during CD4$^+$ T cell

differentiation. Cells lacking granzyme B show a different cytokine profile than granzyme B-

competent cells, primarily presenting increased IL-17 production. Moreover, using the adoptive

transfer model of intestinal inflammation, we show that granzyme B-deficient CD4$^+$ T cells

activated *in vivo* have a different gene expression profile and display increased pathogenicity

relative to WT cells. Similarly, we present evidence indicating that granzyme B deficiency

predisposes mice to higher susceptibility to *Citrobacter rodentium* infection. These results indicate

that granzyme B is required for proper CD4$^+$ T cell differentiation, and in its absence, CD4$^+$ T cells

may acquire an aberrant phenotype characterized by increased IL-17 production and pathogenicity.

Results

Th0 and Th1, but not Th17 CD4$^+$ T cells express granzyme B during differentiation

To determine the role of granzyme B in T cell activation and differentiation, we cultured

naïve CD4$^+$ T cells in Th0, Th1 and Th17 differentiation conditions. One and three days after

culture, RNA was extracted to determine granzyme B expression relative to naïve CD4$^+$ T cells.

Under Th0 conditions, expression of granzyme B at day 1 displayed a slight increase above the

levels observed for naïve CD4$^+$ T cells (Figure 1a, top), but presented an almost three-fold increase

at day 3 (Figure 1a, bottom). Th1 cells expressed a 3-fold increase in granzyme B mRNA over

naïve T cells at day 1, which was maintained until day 3. On the other hand, CD4$^+$ T cells under

Th17 differentiation conditions did not induce granzyme B mRNA expression at either time point

(Figure 1a).

Around 13% of Th0 and Th1 cells expressed granzyme B intracellularly at day 5 (Figure

1b; see Supplemental Figure 1a for gating strategy), whereas this enzyme was not detected in Th17

cells, confirming the results for granzyme B mRNA expression. Although granzyme B mRNA was

detected at day 1 in Th1 cells, granzyme B protein was not, but the protein was conspicuously

expressed at day 3 and day 5 (Figure 1c). These results suggest a lag between granzyme B mRNA

transcription and translation. A similar granzyme B protein expression profile was detected for

Th0 cells (Figure 1c). On the other hand, Th17 cells did not express granzyme B at any of the time

points analyzed (Figure 1c). Similarly, Th2 cells did not express granzyme B (Supplemental Figure

2b).

Because Th1 and Th17 cultures contain different cytokine/antibody cocktails, and

granzyme B is expressed in the former, but not in the latter, we investigated what factors present

in these culture conditions induce or repress granzyme B expression. For this purpose, we activated

CD4$^+$ T cells under Th0 conditions (anti-CD3/CD28, IL-2, and irradiated spleen cells), in the

presence of individual factors included in Th1 and Th17 differentiation cocktails (Figure 1d). IL-

2 alone induced ~7.5% of CD4$^+$ T cells to express granzyme B, which was not significantly

different from cells activated without IL-2 (background levels), or in the presence of IL-12, IL-1β,

and IL-23, the last two of which are involved in Th17 differentiation. However, the Th17-inducing

cytokines IL-6 and TGFβ, as well as the blocking anti-IFNγ antibody, significantly ablated

granzyme B expression. These results indicate that CD4$^+$ T cell activation induces granzyme B

activation, whereas some cytokines and anti-IFNγ inhibit its expression.

Because most of our differentiation cultures included spleen cells as a source of APC, we

determined whether these cells influence granzyme B expression during CD4$^+$ T cell activation.

As shown in Figure 1e, CD4$^+$ T cell activation in the absence of APC (but including IL-2), resulted

in less than 2% of CD4$^+$ T cells expressing granzyme B, whereas irradiated spleen APC

significantly increased granzyme B expression.

Overall, these results indicate that granzyme B is expressed during Th0 and Th1

differentiation, with irradiated APC being an important expression-enhancing factor. On the other

hand, Th17 and Th2 differentiation conditions ablate granzyme B expression. This effect can be

attributed to IL-6, TGFβ, and anti-IFNγ.

As shown in Figure 1b, during Th0 and Th1 differentiation, two populations of CD4$^+$ T

cells can be distinguished: WT-gzmB^{pos} and WT-gzmB^{neg}. To determine whether these

populations present different cytokine profiles, we determined the frequencies of IFNγ- and IL-

17-producing cells in each of these subpopulations. Around half of Th0 WT-gzmB^{pos} expressed

IFNγ, whereas only around 5% of the WT-gzmB^{neg} CD4$^+$ T cells were positive for IFNγ (Figure

1f). Th0 cells derived from granzyme B-deficient (*Gzmb$^{-/-}$*) mice presented decreased levels of

IFNγ-producing cells in comparison to WT-gzmB^{pos} cells, but significantly more IFNγ$^+$ cells than

WT-gzmB^{neg} cells (Figure 1f). On the other hand, IL-17 production was not significantly different

among Th0 cells from any of the CD4$^+$ T cell populations.

Th1 differentiation induced high levels of IFNγ$^+$ cells in all CD4$^+$ T cell populations

analyzed; however, WT-gzmB^{pos} cells presented the highest levels of IFNγ$^+$ cells in comparison

to WT-gzmB^{neg} or cells derived from *Gzmb$^{-/-}$* mice (Figure 1g). Interestingly, under Th1

conditions, cells from *Gzmb$^{-/-}$* mice presented significantly greater levels of IL-17$^+$ cells when

compared to WT-gzmB^{pos} and WT-gzmB^{neg} cells (Figure 1g). These results suggest that granzyme

B may be involved in preventing IL-17 production when CD4$^+$ T cells are cultured under Th1

conditions.

CD4$^+$ T cells cultured in Th2 differentiation conditions, which contains anti-IFNγ, lack

granzyme B expression (Supplemental Figure 1b); therefore, as expected, promiscuous cytokine

expression in granzyme B-deficient CD4$^+$ T cells was not observed in Th2 differentiation

conditions (Supplemental Figure 1c).

***In vivo* activation of granzyme B-deficient CD4$^+$ T cells results in increased pathogenicity**

To further understand the role of granzyme B in CD4$^+$ T cell differentiation, we adoptively

transferred naïve CD4$^+$CD45RB^{hi} T cells from WT or *Gzmb$^{-/-}$* mice into *Rag2$^{-/-}$* recipient mice.

Because it is well-established that transfer of naïve CD4$^+$ T cells into immunocompromised

recipients results in chronic intestinal inflammation, [22, 23] this set up allows us to: 1) determine

whether granzyme B-deficiency alters CD4$^+$ T cell pathogenicity, and 2) whether absence of

granzyme B influences the cytokine profile of CD4$^+$ T cells activated *in vivo* (discussed in the next

section).

In our animal colony, transfer of naïve CD4$^+$ T cells from WT mice typically results in

disease symptoms by 6 weeks after transfer. However, when naïve CD4$^+$ T cells from *Gzmb$^{-/-}$* mice

were transferred into *Rag2$^{-/-}$* mice, weight loss was evident starting at 14 days after transfer, and

by day 21 these mice had lost around 20% of their original weight (Figure 2a). At this point, the

experiments were terminated to comply with our institution's animal welfare regulations. In

contrast, at day 21, *Rag2$^{-/-}$* mice that received WT cells were either gaining or maintaining their

weight (Figure 2a). Weight loss was accompanied by increased disease severity characterized by

diarrhea, pilo-erection, and hunching (Figure 2b); increased intestinal pathology based on immune

cell infiltration, loss of goblet cells, epithelial damage and hyperplasia (Figure 2c); and donor-

derived CD4$^+$ T cell reconstitution (Figure 2d). To confirm that the pathogenicity observed in

granzyme B-deficient CD4$^+$ T cells is an intrinsic effect, we adoptively transferred naïve CD4$^+$ T

cells from littermate *Gzmb$^{+/-}$* and *Gzmb$^{-/-}$* mice into *Rag2$^{-/-}$* recipient mice. Although disease

development was delayed for almost a week, *Gzmb*^{-/-} mice displayed earlier disease onset and

increased colon pathology compared to *Gzmb*^{+/-} littermate controls (Supplemental Figure 2a),

indicating that the pathogenicity observed is intrinsic to CD4$^+$ T cells.

To investigate whether the recipient mice's granzyme B status is important for disease

development, we adoptively transferred naïve CD4$^+$ T cells from WT or *Gzmb*^{-/-} donor mice into

Rag2^{-/-} *Gzmb*^{-/-} recipient mice. Donor CD4$^+$ T cells from WT mice did not cause early weight

loss in the double knock-out recipient mice, while CD4$^+$ T cells deficient in granzyme B caused

similar weight loss as in *Rag2*^{-/-} recipient mice (Supplemental Figure 2b and Figure 2a). Weight

loss in mice receiving granzyme B-deficient CD4$^+$ T cells was accompanied by increased disease

severity (Supplemental Figure 2c), colon pathology (Supplemental Figure 3d), and donor-derived

CD4$^+$ T cell reconstitution (Supplemental Figure 3e). Therefore, the granzyme B status in the

recipient mice is irrelevant for the intrinsic pathogenicity of granzyme B-deficient CD4$^+$ T cells.

In summary, our data indicate that absence of granzyme B in CD4$^+$ T cells activated *in vivo*

results in accelerated pathogenicity and increased cell reconstitution.

Granzyme B-deficient CD4$^+$ T cells present increased IL-17 production *in vivo*.

To determine whether granzyme B deficiency results in differential CD4$^+$ T cell cytokine

profiles *in vivo*, we adoptively transferred naïve CD4$^+$CD45RB^{hi} T cells from WT or *Gzmb*^{-/-} mice

into *Rag2*^{-/-} recipient mice. Three weeks after transfer, donor-derived cells were recovered from

the MLN and lamina propria and analyzed for IFNγ and IL-17 expression. As shown in Figure 3,

IFNγ was the predominant cytokine produced by donor CD4$^+$ T cells derived from WT and *Gzmb*^{-/-}

mice, with similar levels between WT and *Gzmb*^{-/-} CD4$^+$ T cells (Figure 3a and 3b). However,

CD4$^+$ T cells from *Gzmb*^{-/-} donor mice presented significantly increased IL-17 production in

comparison to WT CD4$^+$ T cells (Figure 3a and 3b). It has been reported that pathogenesis in colitis

requires the development of Th17 cells expressing IFNγ.[24] As shown in Figure 3a and 3b, the

frequency of IL-17$^+$IFNγ$^+$ donor-derived CD4$^+$ T cells was similar among both groups in the MLN

and lamina propria, suggesting that these cells may not be responsible for the accelerated colitis

development observed. Albeit less pronounced, cytokine profiles were also similar when littermate

Gzmb$^{+/-}$ and *Gzmb$^{-/-}$* donor mice were used (Supplemental Figure 3).

In summary, *in vivo* activated CD4$^+$ T cells deficient in granzyme B presented increased

IL-17 production. These results suggest that absence of granzyme B results in skewed *in vivo*

differentiation.

Granzyme B deficient CD4$^+$ T cells present a distinct gene expression profile

Because granzyme B-deficient CD4$^+$ T cells had increased IL-17 production when

activated *in vivo*, we investigated whether these cells presented a distinct gene expression profile

in comparison to *in vivo* activated CD4$^+$ T cells from WT mice. For this purpose, CD4$^+$CD45RB^{hi}

T cells from WT and *Gzmb$^{-/-}$* donor mice were adoptively transferred into *Rag2$^{-/-}$* mice. Three

weeks after transfer, CD4$^+$ T cells were recovered from the MLN, their RNA extracted and gene

expression profile determined by RNAseq. As shown in Figure 4a, WT and granzyme B-deficient

CD4$^+$ T cells presented significantly different gene expression profiles. Examples of genes

encoding transcription factors or molecules associated with gene expression upregulated in CD4$^+$

T cells derived from *Gzmb$^{-/-}$* mice include Special AT-rich Binding protein (*Satb1*), a chromatin

remodeling protein involved in T cell development and implicated in Th2 and regulatory T cell

responses;[25] Krüppel-like transcription factor 13 (*Klf13*), which has been implicated in T cell

survival; [26] nuclear receptor subfamily 1, group D, member 1 (*Nr1d1*), which in human T cells is

important for circadian cycles;[27] and the Th17 master transcription factor *Rorc*. We also performed

gene set enrichment analysis to determine whether WT and granzyme B-deficient CD4+ T cells

recovered from recipient mice segregated in specific groups. As indicated in Figure 4b, GSEA

indicated that these two CD4+ T cell populations presented distinct differential gene expression

within several GSEA groups (Figure 4b). These results indicate that granzyme B-deficient CD4+

T cells, in the adoptive transfer system, follow a distinct differentiation pathway compared to WT

CD4+ T cells.

We validated the results generated from the RNAseq analysis by comparing the expression

of selected genes in naïve CD4+ T cells (Figure 4c, top), and donor-derived cells recovered 21d

post-adoptive transfer (Figure 4c, bottom). Hallmark genes involved in Th17 responses, such as

Rorc and *Il17a*, were similarly expressed in naïve CD4+ T cells, regardless of the granzyme B

status, but their expression was significantly increased in recovered granzyme B-deficient CD4+ T

cells. *Ddit4*, a gene important for optimal T cell proliferation,[28] is highly expressed in T cells from

multiple sclerosis patients, and is important in Th17 differentiation.[29] *Ddit4* expression was 3-fold

higher in activated CD4+ T cells derived from *Gzmb*[-/-] mice in comparison to WT CD4+ T cells

(Figure 4c). Granzyme B-deficient CD4+ T cells presented a 2-fold increase in *Il7r* expression

(Figure 4c), which mediates IL-7 signaling, a well-known lymphopoietic cytokine also expressed

in T cells undergoing homeostatic expansion.[30] Plexin D1, encoded by the *Plxnd1* gene, binds

semaphorins[31] and is critical for directing thymocyte migration.[32] Although the role of *Plxnd1* is

not well known in mature T cell biology, granzyme B-deficient CD4+ T cells presented

approximately a 4-fold increase in expression. Interestingly, donor CD4+ T cells derived from

Gzmb[-/-] mice presented decreased *Il10ra* expression in comparison to WT counterparts (Figure

4c).

In terms of IFNγ expression, the total count read average from the RNAseq analysis for

this gene were 2940.7±1332 and 2609.3±1348.9 (P=0.66) for cells derived from WT and *Gzmb*$^{-/-}$

CD4$^+$ T cells, respectively, indicating that IFNγ expression is similar among the two groups. These

results confirm what we observed with intracellular staining for this cytokine (Figure 3).

Overall, these results show that in the adoptive transfer model of CD4$^+$ T cell activation,

granzyme B deficiency results in a distinct gene expression profile, which is characterized by genes

relevant to Th17 differentiation and T cell proliferation.

Granzyme B-deficient CD4$^+$ T cells possess better reconstitution capability

As indicated in Figure 2d, granzyme B-deficient CD4$^+$ T cells showed greater

reconstitution in the colon than WT CD4$^+$ T cells when transferred into Rag-2$^{-/-}$ recipient mice.

However, this observation could be due to inflammation-driven proliferation, which was more

prevalent in *Rag2*$^{-/-}$ mice receiving CD4$^+$ T cells from *Gzmb*$^{-/-}$ mice. To further determine the

reconstitution potential of CD4$^+$ T cells derived from WT and *Gzmb*$^{-/-}$ mice, we performed

competitive adoptive T cell transfer experiments. For this purpose, naïve CD4$^+$ T cells from WT

CD45.1 and *Gzmb*$^{-/-}$ CD45.2 mice (1x10^5 each) were adoptively co-transferred into the same *Rag2*$^-$

$^{/-}$ recipient mice. Of the total donor-derived cells obtained from the MLN, granzyme B-deficient

CD4$^+$ T cells presented ~60% reconstitution, whereas WT-derived cells showed ~40%

reconstitution (Figure 5a). To determine whether increased cell reconstitution by granzyme B-

deficient CD4$^+$ T cells correlated with increased cell division, we stained the cells for Ki67 as a

surrogate marker for cell division. As shown in Figure 5b, there was higher percentage of Ki67$^+$

cells in granzyme-B deficient donor-derived cells. To further confirm these results, we cultured

donor-derived cells without any stimulation, and measured their division potential by CFSE

dilution after 3 days. CD4$^+$ T cells from *Gzmb$^{-/-}$* mice showed a slight increase in cell division

compared to cells derived from WT mice (Figure 5c).

We also analyzed the cytokine profile of the cells derived from the competitive adoptive

CD4$^+$ T cell approach. Although the IFNγ profile of co-transferred cells was similar to cells

transferred alone, co-transferred WT cells presented increased IL-17 production in comparison to

WT CD4$^+$ T cells transferred alone (Figure 5d, left and middle graphs). These results raise the

possibility that granzyme B-deficient CD4$^+$ T cells influenced the behavior of granzyme B-

competent CD4$^+$ T cells. Interestingly, co-transferred granzyme B-deficient CD4$^+$ T cells

displayed a higher percentage of IFNγ$^+$IL-17$^+$ cells in comparison to WT cells (Figure 5d, right

graph). *Rag2$^{-/-}$* mice co-transferred with WT and GzB$^{-/-}$ CD4$^+$ T cells displayed similar weight

loss (Figure 5e) and colon pathology (Figure 5f) to mice transferred only with CD4$^+$ T cells from

Gzmb$^{-/-}$ mice, suggesting that the increased pathogenicity of the latter cells was maintained even

in the presence of granzyme-competent CD4$^+$ T cells.

Granzyme B-deficient mice present normal cellularity and IFNγ /IL-17 production

Since *in vivo* activated granzyme B-deficient CD4$^+$ T cells possess differential gene

expression and cytokine profiles, it is possible that these differences are also observed in naïve

cells. To investigate this possibility, we enumerated the CD4$^+$ T cell cellularity of spleen, MLN,

and IEL compartments from WT and *Gzmb$^{-/-}$* mice. As indicated in Supplemental Figure 4a,

frequencies and numbers of TCRβ$^+$CD4$^+$ T cells were similar among the two groups of mice. *Ex*

vivo IFNγ and IL-17 production by non-stimulated (ns) or stimulated (s) naïve CD4$^+$T cells derived

from the MLN and IEL showed no distinguishable difference between cells from WT and *Gzmb$^{-/-}$*

mice (Supplemental Figure 4b).

In summary, naïve WT and $Gzmb^{-/-}$ animals appear to have similar CD4$^+$ T cell cellularity

and basal IFNγ and IL-17 production in peripheral lymphoid organs and the intestines.

Granzyme B deficiency results in increased disease severity during *Citrobacter rodentium*

infection

As demonstrated in the previous section, naïve $Gzmb^{-/-}$ mice have a normal CD4$^+$ T cell

compartment compared to WT mice. However, because we observed that activated T cells from

$Gzmb^{-/-}$ mice became more pathogenic and had a distinct gene expression profile in the adoptive

transfer model of colitis, we investigated whether $Gzmb^{-/-}$ mice respond properly to antigenic

stimulus. For this purpose, we infected WT and $Gzmb^{-/-}$ mice with the extracellular bacterium

Citrobacter rodentium, which preferentially colonizes the colon of mice and induces an IL-17/IL-

22-based immune response. After infection, WT mice maintained similar weight throughout the

course of the experiment; however, starting at 9 days post-infection, $Gzmb^{-/-}$ mice lost weight,

reaching approximately ~20% loss of the starting weight by day 12 (Figure 6a). $Gzmb^{-/-}$ mice also

presented other signs of disease, such as diarrhea, hunched posture, and pilo-erection, which were

mostly absent in WT mice (Figure 6b). Despite similar colonic bacterial burden (Figure 6c), $Gzmb^{-/-}$

mice developed greater pathology characterized by increased infiltration and epithelial injury

than WT mice (Figure 6d). Analysis of cytokine production showed that in the spleen and LP of

infected $Gzmb^{-/-}$ mice, there was a significant increase in CD4$^+$ T cells expressing IL-17, while

IFNγ production was only increased in the MLN (Figure 6e).

Because innate immune cells are important for the response against *C. rodentium*, we

investigated whether the absence of granzyme B in innate immune cells was responsible for the

observed increased pathology in infected $Gzmb^{-/-}$ mice. For this purpose, $Rag-2^{-/-}$ and $Rag-2^{-/-}$

Gzmb[-/-] mice were infected with *C. rodentium* and monitored for 14 days. Both groups of mice

maintained similar weights throughout the course of the experiment (Supplemental Figure 5a),

indicating that the effect caused by granzyme B-deficiency is most likely associated with adaptive

immune cells. To further confirm these results, we treated *Rag-2*[-/-] and *Rag-2*[-/-]*Gzmb*[-/-] mice with

anti-CD40 antibodies, which induce acute intestinal inflammation in the absence of T or B cells.[33]

As shown in Supplemental Figure 5b, *Rag-2*[-/-] and *Rag-2*[-/-]*Gzmb*[-/-] mice presented similar weight

loss throughout the course of the experiment.

Interestingly, when *Gzmb*[+/-] and *Gzmb*[-/-] littermate mice were infected with *C. rodentium*,

both groups displayed similar weight curves and colon colonization (Supplemental Figure 5c and

5d). However, there was a slight trend for increased IFNγ[+]IL-17[+] CD4[+] T cells in the spleen of

Gzmb[-/-] mice (Supplemental Figure 5e).

Discussion

Granzyme B expression in CD4[+] T cells is well-documented, and has been primarily

studied in the context of granzyme B/perforin-dependent regulatory T cell suppressor activity.[34]

However, as our data indicate, granzyme B expression appears to be a common feature among a

significant fraction of activated CD4[+] T cells. Interestingly, not all differentiation conditions

induced granzyme B expression. For example, Th0 and Th1 cells rapidly express granzyme B

starting at 1-day post activation, whereas Th17 cells do not express this enzyme at any time after

activation. In the former conditions, expression of granzyme B is primarily driven by the presence

of irradiated splenocytes, indicating the possibility that cell-to-cell contact is important for

granzyme B expression.

Although none of the cytokines involved in Th1 differentiation induced granzyme B

expression above the levels observed with the addition of irradiated splenocytes, anti-IFNγ and

some cytokines present in the Th17 cocktail, such as IL-6 and TGFβ prevented the expression of

this enzyme, which suggests that restriction of granzyme B expression is necessary for Th17

polarization. Studies in cytotoxic CD8$^+$ T cells have shown that activation of these cells with anti-

CD3 in the presence of IL-6, but not in the presence of TGFβ, induces granzyme B expression.[35]

Here we show that in CD4$^+$ T cell differentiation, IL-6 does not enhance the expression of

granzyme B, indicating that CD8$^+$ and CD4$^+$ T cells possess different mechanisms for modulating

granzyme B production.

Granzyme B expression may not only be limited during the initial CD4$^+$ T cell priming and

differentiation. For example, TGFβ is known for its role in regulatory T cell differentiation, and

some of these cells require granzyme B for their suppressor functions. However, as our results

indicate, TGFβ ablates expression of this enzyme, suggesting that regulatory T cells must acquire

granzyme B expression during post-priming events. Similarly, a fraction of effector CD4$^+$ T cells

that migrate into the intestinal intraepithelial lymphocyte compartment acquire granzyme B

expression after transcriptional reprogramming.[36]

One of the most intriguing questions is: Why do CD4$^+$ T cells express granzyme B during

Th0/Th1 activation and differentiation? During *in vitro* Th0/Th1 differentiation, WT-gzmB^{pos} and

WT-gzmB^{neg} cells have distinct IFNγ profiles, suggesting that these cells may represent different

lineages and that the activity of granzyme B influences their outcome. This is better exemplified

when analyzing granzyme B-deficient CD4$^+$ T cells, where *in vivo* and *in vitro* activation skew the

cells towards an IL-17-producing phenotype, with a distinct gene expression signature. We

postulate that granzyme B-deficient IL-17$^+$ cells represent a lineage of cells that underwent

aberrant differentiation. Therefore, expression of granzyme B during Th0/Th1 activation serves as

a checkpoint for proper CD4$^+$ T cell differentiation, preventing IL-17 production and increased

pathology. These observations have important significance because it has been reported that allelic

variants of granzyme B correlate with autoimmune disorders,[37-39] raising the possibility that in

certain CD4$^+$ T cell-mediated disorders, lack of proper granzyme B function increases the

probability of improper differentiation and increased pathogenesis potential. If this idea is correct,

then WT-gzmB^{pos} CD4$^+$ T cells may represent cells in which the differentiation signals have the

potential to skew cells into an unwanted phenotype, and granzyme B expression in these cells

ensures the correct differentiation process.

We have shown that adoptive transfer of CD4$^+$ T cells derived from non-littermate (Figure

2) and littermate (Supplemental Figure 2a) mice resulted in increased pathogenicity and IL-17

production, which argues that granzyme B deficiency has an intrinsic effect in CD4$^+$ T cells.

However, the host's environment also influences the pathogenicity and IL-17 production of

granzyme B-deficient CD4$^+$ T cells. For example, infection of $Gzmb^{-/-}$ mice with $C.\ rodentium$

resulted in more severe disease with increased IL-17 production relative to WT mice (Figure 6);

however, this effect was not observed in infected $Gzmb^{+/-}$ and $Gzmb^{-/-}$ littermate mice

(Supplemental Figure 5c-5e). This observation suggests that granzyme B deficiency in CD4$^+$ T

cells is necessary but not sufficient for their associated increased IL-17 production and

pathogenicity. One potential factor that may be responsible for controlling the pathogenicity of

granzyme B-deficient CD4$^+$ T cells may be the microbiota present in granzyme B-competent mice.

We are currently investigating this interesting hypothesis.

Granzyme B activity in lymphocyte development and/or differentiation has been

previously reported.[40] These authors showed that granzyme A and B are important for the

development of a subset of TCR[neg] intestinal intraepithelial lymphocytes known for the expression

of intracellular CD3γ. Development of these cells requires granzyme B to cleave and inactivate

the intracellular domain of Notch1. This raises the interesting possibility that during CD4[+] T cell

differentiation, granzyme B prevents unwanted phenotypes by disrupting signals coming from the

environment, such those provided by Notch and its ligands, which are known to influence CD4[+] T

cell differentiation.[41] Our observation that granzyme B expression was increased in the presence

of irradiated splenocytes supports this hypothesis. We are currently investigating what signals

derived from APC enhance granzyme B expression.

It has been reported that granzyme B-deficient CD4[+]CD25[-] T cells used in murine models

of graft versus host disease (GVHD) induced faster disease onset with increased lethality in

comparison to their WT counterparts.[42] Although this group did not provide a mechanistic

explanation for the increased pathogenicity observed, they suggested that increased proliferation

and decreased cell death may be involved. According to our data, we believe that in the GVHD

model, differential proliferation and survival of granzyme B-deficient CD4[+] T cells are the result

of aberrant CD4[+] T cell differentiation, which allows escape of highly pathogenic clones. Further

investigations in the GVHD model are needed to test this hypothesis.

In summary, granzyme B has been known for its role in cell mediated-cytotoxicity, and

while many groups have reported potential extracellular roles for this enzyme,[43] here we present

in vitro and *in vivo* evidence supporting a novel intrinsic function for granzyme B during CD4[+] T

cell differentiation. Although many questions remain to be answered regarding how granzyme B

is involved in this process, our results provide a strong foundation for a better understanding of

the function of this enzyme in CD4[+] T cell biology.

Methods

Mice. C57BL/6J and C57BL/6J.CD45.1 mice were originally purchased from The Jackson Laboratory (000664, and 002014 respectively) and have been maintained and acclimated in our colony for several years. Granzyme B (*Gzmb*)$^{-/-}$ mice were kindly provided by Dr. Xuefang Cao. Littermate mice were generated by crossing *Gzmb*$^{-/-}$ with WT mice; F1 males (*Gzmb*$^{+/-}$) were subsequently crossed to *Gzmb*$^{-/-}$ female mice to obtain *Gzmb*$^{+/-}$ and *Gzmb*$^{-/-}$ mice. CD57BL/6J *Rag2*$^{-/-}$ mice were kindly provided by Dr. Luc Van Kaer. *Rag2*$^{-/-}$*Gzmb*$^{-/-}$ were generated in our colony. All mice were between 6 to 10 weeks of age at the time of experimentation. Male and female mice were used for all experiments. Mice were maintained in accordance with the Institutional Animal Care and Use Committee at Vanderbilt University.

Reagents and flow cytometry. Fluorochrome-coupled anti-mouse CD4 (GK1.5), CD8α (53-6.7), CD45RB (C363.16A), granzyme B (NGZB), IFNγ (XMG1.2), IL-17a (TC11-18H10), Ki67 (solA15), TCRβ (H57-597) and ghost viability dyes were purchased from ThermoFisher, BD Bioscences or Tonbo. Biotinylated anti-CD8α (53-1.7) and CD19 (1D3) antibodies were purchased from Tonbo. The Invitrogen Vybrant CFDA SE cell tracer kit was purchased from ThermoFisher Scientific. Anti-CD40 antibody was purchased from BioXcell. Surface cell staining was performed following conventional techniques. For intracellular cytokine staining, cells were stimulated with PMA and ionomycin in the presence of Golgi Stop (BD Biosciences) for 4hr prior to staining. Extracellular markers were stained, cells were fixed briefly with 2% paraformaldehyde, followed by permeabilization and intracellular staining. For intracellular cytokine and granzyme B staining, the BD Cytofix/Cytoperm kit was used according to the manufacturer's instructions. For intracellular Ki67 staining, the eBioscience transcription factor staining buffer set was used

according to the manufacturer's instructions. All stained samples were acquired using BD FACS

Canto II, 3- or 4-Laser Fortessa, or 5-Laser LSR II flow cytometers (BD Bioscences). Data were

analyzed using FlowJo software (Tree Star).

Lymphocyte isolation. Spleen and MLN lymphocytes were isolated by conventional means. IEL

and LP cells were isolated by mechanical disruption as previously reported.[44] Briefly, after

flushing the intestinal contents with cold HBSS and removing excess mucus, the intestines were

cut into small pieces (~1 cm long) and shaken for 45 minutes at 37°C in HBSS supplemented with

5% fetal bovine serum and 2 mM EDTA. Supernatants were recovered and cells isolated using a

discontinuous 40/70% Percoll (General Electric) gradient. To obtain lamina propria lymphocytes,

intestinal tissue was recovered and digested with collagenase (187.5 U/ml, Sigma) and DNase I

(0.6 U /ml, Sigma). Cells were isolated using a discontinuous 40/70% Percoll gradient.

In vitro *CD4 T cell activation/differentiation.* Naïve CD4 T cells were isolated from the spleens of

the indicated mice by magnetic sorting using the Miltenyi or StemCell Technologies naïve CD4$^+$

T cell isolation kits according to the manufacturer's instructions. Splenocytes or MLN cells used

as APC were incubated for 1hr at 37°C, non-adherent cells were removed, adherent cells were

washed and resuspended in complete RPMI, and irradiated with 7 Gy (700 rads). 5×10^5 CD4$^+$ T

cells with or without 5×10^5 irradiated APC were incubated in the presence of plate bound anti-

CD3 (5μg/ml) and soluble anti-CD28 (2.5μg/ml), for the indicated times, under the following

conditions: Th0 (10ng/mL IL-2), Th1 (10ng/mL IL-2, 20ng/mL IL-12, and 10μg/mL anti-IL-4) ,

Th17 (5ng/mL TGFβ, 20ng/mL IL-6, 10ng/mL IL-1β, 10ng/mL IL-23, 10μg/mL anti-IL-4, and

10μg/mL anti-IFNγ), and Th2 (10ng/mL IL-4, 2μg/mL anti-IFNγ, and 2μg/mL anti IL-12).

Adoptive transfer of naïve CD4 T cells. Splenocytes from WT or *Gzmb*[-/-] mice were depleted of

CD19[+] and CD8α[+] cells using magnetic bead sorting (Miltenyi). Enriched cells were stained with

antibodies directed against TCRβ, CD4 and CD45RB, and naïve CD4[+] T cells were sorted as

CD4[+]CD45RB[hi] using a FACSAria III at the Flow Cytometry Shared Resource at VUMC. 1×10^5

cells were adoptively transferred into the indicated recipient mice. Starting weight was determined

prior to injection. Mice were monitored and weighed weekly. At the end time point, mice were

sacrificed and donor-derived cells were isolated from the indicated organs or tissues. Colon

histopathology was performed in a blind fashion by a GI pathologist (MBP), following established

parameters.[45] For competitive transfer experiments, 1×10^5 CD4[+]CD45RB[hi] cells from both WT-

CD45.1 and *Gzmb*[-/-] CD45.2 mice were adoptively co-transferred into *Rag2*[-/-] mice and analyzed

as indicated above.

Transcription profile analysis. For gene expression analysis, TCRβ[+]CD4[+] T cells were purified by

flow cytometry from a pool of MLN cells comprised of 1 male and 1 female animal with similar

disease progression 21 days after adoptive transfer. RNA was isolated using the RNeasy micro kit

(Qiagen). cDNA library preparation and RNAseq was performed by the VANTAGE Core at

VUMC on an Illumina NovaSeq 6000 (2 x 150 base pair, paired-end reads). The tool Salmon[46]

was used for quantifying the expression of RNA transcripts. The R project software along with the

edgeR method[47] was used for differential expression analysis. For gene set enrichment analysis

(GSEA), RNAseq data was ranked according to the t-test statistic. The gene sets curated (C2), GO

(C5), immunological signature collection (C7) and hallmarks of cancer (H) of the Molecular

Signatures Database (MSigDB) were used for enrichment analysis. GSEA enrichment plots were

generated using the GSEA software[48] from the Broad Institute with 1000 permutations.

Real time PCR. For *in vitro* analysis, RNA was extracted from MACS-enriched, differentiated

CD4$^+$ T cells at day 1 and 3 of culture. For *in vivo* analysis, RNA was isolated from flow cytometry

purified donor-derived TCRβ$^+$CD4$^+$ T cells as described above. cDNA was synthesized using the

RT2 First Strand kit (Qiagen). Real time PCR was performed using RT2 SYBR Green Mastermix

(Qiagen) on an Applied BioSystems Proflex PCR cycler. All primers were purchased from Qiagen.

Citrobacter rodentium *infection*. Seven-week old female WT and *Gzmb$^{-/-}$* or *Gzmb$^{+/-}$* and *Gzmb$^{-/-}$*

littermate animals were infected with *C. rodentium* as previously described[49]. Briefly, mice were

infected with 5x10^8 CFU exponentially grown bacteria by oral gavage. Starting weight was

determined prior to gavage. Mice were monitored and weighed daily for 14d. At the end time

point, cells were isolated from the indicated organs or tissues. Cellularity was determined,

intracellular cytokine staining was performed, and colon histopathology was performed in a blind

fashion by a GI pathologist (MBP) as previously described.[50]

Anti-CD40 disease induction. *Rag-2$^{-/-}$* and *Rag-2$^{-/-}$Gzmb$^{-/-}$* mice were injected i.p. with

150µg/mouse of anti-CD40 (BioXcell). Mice were monitored and weighed for daily for 7 days.

References

1.   Anthony DA, Andrews DM, Watt SV, Trapani JA, Smyth MJ. Functional dissection of the
499   granzyme family: cell death and inflammation. *Immunol Rev* 2010; **235**(1)**:** 73-92.
500

501 2. Chowdhury D, Lieberman J. Death by a thousand cuts: granzyme pathways of programmed
502   cell death. *Annu Rev Immunol* 2008; **26:** 389-420.
503

504 3. Afonina IS, Cullen SP, Martin SJ. Cytotoxic and non-cytotoxic roles of the CTL/NK
505   protease granzyme B. *Immunol Rev* 2010; **235**(1)**:** 105-116.
506

507 4. Wensink AC, Hack CE, Bovenschen N. Granzymes regulate proinflammatory cytokine
508   responses. *J Immunol* 2015; **194**(2)**:** 491-497.
509

510 5. Kumar AA, Delgado AG, Piazuelo MB, Van Kaer L, Olivares-Villagomez D. Innate
511   CD8alphaalpha(+) lymphocytes enhance anti-CD40 antibody-mediated colitis in mice.
512   *Immun Inflamm Dis* 2017; **5**(2)**:** 109-123.
513

514 6. Parkinson LG, Toro A, Zhao H, Brown K, Tebbutt SJ, Granville DJ. Granzyme B mediates
515   both direct and indirect cleavage of extracellular matrix in skin after chronic low-dose
516   ultraviolet light irradiation. *Aging Cell* 2015; **14**(1)**:** 67-77.
517

518 7. Chamberlain CM, Ang LS, Boivin WA, Cooper DM, Williams SJ, Zhao H *et al.* Perforin-
519   independent extracellular granzyme B activity contributes to abdominal aortic aneurysm.
520   *Am J Pathol* 2010; **176**(2)**:** 1038-1049.
521

522 8. Lancki DW, Hsieh CS, Fitch FW. Mechanisms of lysis by cytotoxic T lymphocyte clones.
523   Lytic activity and gene expression in cloned antigen-specific CD4+ and CD8+ T
524   lymphocytes. *J Immunol* 1991; **146**(9)**:** 3242-3249.
525

526 9. Ebnet K, Chluba-de Tapia J, Hurtenbach U, Kramer MD, Simon MM. In vivo primed
527   mouse T cells selectively express T cell-specific serine proteinase-1 and the proteinase-
528   like molecules granzyme B and C. *Int Immunol* 1991; **3**(1)**:** 9-19.
529

530 10. Sun Q, Burton RL, Pollok KE, Emanuel DJ, Lucas KG. CD4(+) Epstein-Barr virus-specific
531   cytotoxic T-lymphocytes from human umbilical cord blood. *Cell Immunol* 1999; **195**(2)**:**
532   81-88.
533

534 11. Canaday DH, Wilkinson RJ, Li Q, Harding CV, Silver RF, Boom WH. CD4(+) and CD8(+)
535   T cells kill intracellular Mycobacterium tuberculosis by a perforin and Fas/Fas ligand-
536   independent mechanism. *J Immunol* 2001; **167**(5)**:** 2734-2742.
537

538 12. Yasukawa M, Ohminami H, Yakushijin Y, Arai J, Hasegawa A, Ishida Y *et al.* Fas-
539   independent cytotoxicity mediated by human CD4+ CTL directed against herpes simplex
540   virus-infected cells. *J Immunol* 1999; **162**(10)**:** 6100-6106.
541

542 13. Dorothee G, Vergnon I, Menez J, Echchakir H, Grunenwald D, Kubin M *et al.* Tumor-
543   infiltrating CD4+ T lymphocytes express APO2 ligand (APO2L)/TRAIL upon specific
544   stimulation with autologous lung carcinoma cells: role of IFN-alpha on APO2L/TRAIL
545   expression and -mediated cytotoxicity. *J Immunol* 2002; **169**(2)**:** 809-817.
546

547   14.   Grossman WJ, Verbsky JW, Tollefsen BL, Kemper C, Atkinson JP, Ley TJ. Differential
548   expression of granzymes A and B in human cytotoxic lymphocyte subsets and T regulatory
549   cells. *Blood* 2004; **104**(9)**:** 2840-2848.
550

551   15.   Koonpaew S, Shen S, Flowers L, Zhang W. LAT-mediated signaling in CD4+CD25+
552   regulatory T cell development. *J Exp Med* 2006; **203**(1)**:** 119-129.
553

554   16.   Kawamura K, Kadowaki N, Kitawaki T, Uchiyama T. Virus-stimulated plasmacytoid
555   dendritic cells induce CD4+ cytotoxic regulatory T cells. *Blood* 2006; **107**(3)**:** 1031-1038.
556

557   17.   Price JD, Schaumburg J, Sandin C, Atkinson JP, Lindahl G, Kemper C. Induction of a
558   regulatory phenotype in human CD4+ T cells by streptococcal M protein. *J Immunol* 2005;
559   **175**(2)**:** 677-684.
560

561   18.   Gondek DC, Lu LF, Quezada SA, Sakaguchi S, Noelle RJ. Cutting edge: contact-mediated
562   suppression by CD4+CD25+ regulatory cells involves a granzyme B-dependent, perforin-
563   independent mechanism. *J Immunol* 2005; **174**(4)**:** 1783-1786.
564

565   19.   Zhu J, Yamane H, Paul WE. Differentiation of effector CD4 T cell populations (*). *Annu
566   Rev Immunol* 2010; **28:** 445-489.
567

568   20.   Sharma V, Delgado M, Ganea D. Granzyme B, a new player in activation-induced cell
569   death, is down-regulated by vasoactive intestinal peptide in Th2 but not Th1 effectors. *J
570   Immunol* 2006; **176**(1)**:** 97-110.
571

572   21.   Coomes SM, Kannan Y, Pelly VS, Entwistle LJ, Guidi R, Perez-Lloret J *et al.* CD4(+) Th2
573   cells are directly regulated by IL-10 during allergic airway inflammation. *Mucosal
574   immunology* 2017; **10**(1)**:** 150-161.
575

576   22.   Powrie F, Correa-Oliveira R, Mauze S, Coffman RL. Regulatory interactions between
577   CD45RBhigh and CD45RBlow CD4+ T cells are important for the balance between
578   protective and pathogenic cell-mediated immunity. *J Exp Med* 1994; **179**(2)**:** 589-600.
579   PMC2191378.
580

581   23.   Powrie F, Leach MW, Mauze S, Menon S, Caddle LB, Coffman RL. Inhibition of Th1
582   responses prevents inflammatory bowel disease in scid mice reconstituted with CD45RBhi
583   CD4+ T cells. *Immunity* 1994; **1**(7)**:** 553-562.
584

585   24.   Harbour SN, Maynard CL, Zindl CL, Schoeb TR, Weaver CT. Th17 cells give rise to Th1
586   cells that are required for the pathogenesis of colitis. *Proc Natl Acad Sci U S A* 2015;
587   **112**(22)**:** 7061-7066.
588

589   25.   Zelenka T, Spilianakis C. SATB1-mediated chromatin landscape in T cells. *Nucleus* 2020;
590   **11**(1)**:** 117-131.
591

592   26.   Zhou M, McPherson L, Feng D, Song A, Dong C, Lyu SC *et al.* Kruppel-like transcription
593   factor 13 regulates T lymphocyte survival in vivo. *J Immunol* 2007; **178**(9)**:** 5496-5504.
594

595   27.   Bollinger T, Leutz A, Leliavski A, Skrum L, Kovac J, Bonacina L *et al.* Circadian clocks
596   in mouse and human CD4+ T cells. *PLoS One* 2011; **6**(12)**:** e29801.
597

598   28.   Reuschel EL, Wang J, Shivers DK, Muthumani K, Weiner DB, Ma Z *et al.* REDD1 Is
599   Essential for Optimal T Cell Proliferation and Survival. *PLoS One* 2015; **10**(8)**:** e0136323.
600

601   29.   Zhang F, Liu G, Li D, Wei C, Hao J. DDIT4 and Associated lncDDIT4 Modulate Th17
602   Differentiation through the DDIT4/TSC/mTOR Pathway. *J Immunol* 2018; **200**(5)**:** 1618-
603   1626.
604

605   30.   Fry TJ, Mackall CL. The many faces of IL-7: from lymphopoiesis to peripheral T cell
606   maintenance. *J Immunol* 2005; **174**(11)**:** 6571-6576.
607

608   31.   Carvalheiro T, Rafael-Vidal C, Malvar-Fernandez B, Lopes AP, Pego-Reigosa JM,
609   Radstake T *et al.* Semaphorin4A-Plexin D1 Axis Induces Th2 and Th17 While Represses
610   Th1 Skewing in an Autocrine Manner. *Int J Mol Sci* 2020; **21**(18).
611

612   32.   Choi YI, Duke-Cohan JS, Ahmed WB, Handley MA, Mann F, Epstein JA *et al.* PlexinD1
613   glycoprotein controls migration of positively selected thymocytes into the medulla.
614   *Immunity* 2008; **29**(6)**:** 888-898.
615

616   33.   Uhlig HH, McKenzie BS, Hue S, Thompson C, Joyce-Shaikh B, Stepankova R *et al.*
617   Differential activity of IL-12 and IL-23 in mucosal and systemic innate immune pathology.
618   *Immunity* 2006; **25**(2)**:** 309-318.
619

620   34.   Cao X, Cai SF, Fehniger TA, Song J, Collins LI, Piwnica-Worms DR *et al.* Granzyme B
621   and perforin are important for regulatory T cell-mediated suppression of tumor clearance.
622   *Immunity* 2007; **27**(4)**:** 635-646.
623

624   35.   Huber M, Heink S, Grothe H, Guralnik A, Reinhard K, Elflein K *et al.* A Th17-like
625   developmental process leads to CD8(+) Tc17 cells with reduced cytotoxic activity. *Eur J
626   Immunol* 2009; **39**(7)**:** 1716-1725.
627

628   36.   Mucida D, Husain MM, Muroi S, van Wijk F, Shinnakasu R, Naoe Y *et al.* Transcriptional
629   reprogramming of mature CD4(+) helper T cells generates distinct MHC class II-restricted
630   cytotoxic T lymphocytes. *Nat Immunol* 2013; **14**(3)**:** 281-289.
631

632   37.   Jin Y, Birlea SA, Fain PR, Gowan K, Riccardi SL, Holland PJ *et al.* Variant of TYR and
633   autoimmunity susceptibility loci in generalized vitiligo. *The New England journal of
634   medicine* 2010; **362**(18)**:** 1686-1697.
635

636   38.   Li S, Yao W, Pan Q, Tang X, Zhao S, Wang W *et al.* Association analysis revealed one
637   susceptibility locus for vitiligo with immune-related diseases in the Chinese Han
638   population. *Immunogenetics* 2015; **67**(7)**:** 347-354.
639

640   39.   Shen C, Gao J, Sheng Y, Dou J, Zhou F, Zheng X *et al.* Genetic Susceptibility to Vitiligo:
641   GWAS Approaches for Identifying Vitiligo Susceptibility Genes and Loci. *Front Genet*
642   2016; **7:** 3.
643

644   40.   Ettersperger J, Montcuquet N, Malamut G, Guegan N, Lopez-Lastra S, Gayraud S *et al.*
645   Interleukin-15-Dependent T-Cell-like Innate Intraepithelial Lymphocytes Develop in the
646   Intestine and Transform into Lymphomas in Celiac Disease. *Immunity* 2016; **45**(3)**:** 610-
647   625.
648

649   41.   Amsen D, Helbig C, Backer RA. Notch in T Cell Differentiation: All Things Considered.
650   *Trends Immunol* 2015; **36**(12)**:** 802-814.
651

652   42.   Du W, Leigh ND, Bian G, O'Neill RE, Mei L, Qiu J *et al.* Granzyme B-Mediated
653   Activation-Induced Death of CD4+ T Cells Inhibits Murine Acute Graft-versus-Host
654   Disease. *J Immunol* 2015; **195**(9)**:** 4514-4523.
655

656   43.   Boivin WA, Cooper DM, Hiebert PR, Granville DJ. Intracellular versus extracellular
657   granzyme B in immunity and disease: challenging the dogma. *Laboratory investigation; a*
658   *journal of technical methods and pathology* 2009; **89**(11)**:** 1195-1220.
659

660   44.   Nazmi A, Greer MJ, Hoek KL, Piazuelo MB, Weitkamp J, Olivares-Villagomez D.
661   Osteopontin and iCD8alpha cells promote intestinal intraepithelial lymphocyte
662   homeostasis. *J Immunol* 2020; **In press**.
663

664   45.   Aranda R, Sydora BC, McAllister PL, Binder SW, Yang HY, Targan SR *et al.* Analysis of
665   intestinal lymphocytes in mouse colitis mediated by transfer of CD4+, CD45RBhigh T
666   cells to SCID recipients. *J Immunol* 1997; **158**(7)**:** 3464-3473.
667

668   46.   Patro R, Duggal G, Love MI, Irizarry RA, Kingsford C. Salmon provides fast and bias-
669   aware quantification of transcript expression. *Nature methods* 2017; **14**(4)**:** 417-419.
670

671   47.   Robinson MD, McCarthy DJ, Smyth GK. edgeR: a Bioconductor package for differential
672   expression analysis of digital gene expression data. *Bioinformatics* 2010; **26**(1)**:** 139-140.
673

674   48.   Subramanian A, Tamayo P, Mootha VK, Mukherjee S, Ebert BL, Gillette MA *et al.* Gene
675   set enrichment analysis: a knowledge-based approach for interpreting genome-wide
676   expression profiles. *Proc Natl Acad Sci U S A* 2005; **102**(43)**:** 15545-15550.
677

678   49.   Olivares-Villagomez D, Algood HM, Singh K, Parekh VV, Ryan KE, Piazuelo MB *et al.*
679   Intestinal epithelial cells modulate CD4 T cell responses via the thymus leukemia antigen.
680   *J Immunol* 2011; **187**(8)**:** 4051-4060.
681

682   50.   Singh K, Gobert AP, Coburn LA, Barry DP, Allaman M, Asim M *et al.* Dietary Arginine
683   Regulates Severity of Experimental Colitis and Affects the Colonic Microbiome. *Front*
684   *Cell Infect Microbiol* 2019; **9:** 66.

Figure legends

Figure 1. Granzyme B expression during T cell differentiation. (a) Naïve CD4+ T cells from WT

mice were cultured in the presence of plate-bound anti-CD3 and soluble anti-CD28 under the

indicated differentiation conditions, and in the absence of irradiated splenocytes. At day 1 (top)

and 3 (bottom), cells were recovered, RNA extracted and granzyme B mRNA expression

determined. Naïve CD4+ T cells without stimulation were used as reference. (b) Intracellular

granzyme B in cells cultured as in (a), but in the presence of irradiated splenocytes. Left,

representative dot plot (see Supplemental Figure 1a for gating strategy). Right, data summary. (c)

Granzyme B expression in cells cultured in Th0, Th1 and Th17 conditions in the presence of

irradiated splenocytes at different time points. (d) Cells cultured in Th0 conditions in the

presence of irradiated splenocytes and the indicated individual cytokines, harvested 5 days after

culture. (e) Cells cultured with IL-2, with or without irradiated splenocytes (APC). (f) and (g),

intracellular IFNγ and IL-17 staining of WT and *Gzmb*-/- CD4+ T cells cultured in Th0 or Th1

conditions, respectively, in the presence of irradiated splenocytes. Top, intracellular IFNγ and

IL-17 staining of representative concatenated plots; bottom, data summary. Each dot represents

an individual sample. Data are representative of at least 3 independent experiments, performed in

triplicate. n=6. For (d), (f) and (g), *P<0.05; **P<0.01; ***P<0.001; ****P<0.001, One-way

ANOVA; For (e), ***P<0.001, Student's t test.

Figure 2. *In vivo* activated granzyme B-deficient CD4+ T cells display increased pathogenicity.

1×10^5 CD4+CD45RBhi T cells from WT or *Gzmb*-/- donor mice were adoptively transferred into

Rag2-/- recipient mice. (a) Mice were weighed weekly, and (b) monitored for signs of disease.

Twenty-one days after transfer, (c) colon pathology was scored (left, representative micrographs;

right, data summary), and (d) donor-cell reconstitution of the IEL compartment in the colon was

determined. For (a), dotted lines represent untreated *Rag2*-/- mice. Each dot represents an

individual mouse. Data are representative of at least 3 independent experiments. n=8-10.

**P<0.01; ***P<0.001; ****P<0.0001; Student's t-test.

Figure 3. Granzyme B-deficient CD4+ T cells display skewed IL-17 differentiation *in vivo*.

1×10^5 CD4+CD45RBhi T cells from WT or *Gzmb*-/- donor mice were adoptively transferred into

Rag2-/- recipient mice. Three weeks after transfer, donor-derived cells were recovered from (a)

MLN, and (b) lamina propria, and their IFNγ/IL-17 profile was determined by intracellular

staining. Dot plots indicate a representative sample. Live, TCRβ⁺CD4⁺ cells are displayed.

Graphs represent the summary. Each dot indicates an individual mouse. For MLN, n=11-13; for

lamina propria, n=7-11. Data are representative of at least 3 independent experiments. **P<0.01;

****P<0.001. Student's t-Test.

Figure 4. Granzyme B-deficient CD4⁺ T cells activated *in vivo* display a distinct gene expression

profile. 1x10⁵ CD4⁺CD45RB^hi T cells from WT or *Gzmb-/-* donor mice were adoptively

transferred into *Rag2-/-* recipient mice. Three weeks after transfer, donor-derived cells were

recovered from the MLN, sorted for live TCRβ⁺CD4⁺, their RNA isolated and sequenced. (a)

Representative heat map. (b) Representative gene set enrichment analysis. The green line on the

plots indicate the enrichment score; the black lines show where the genes related to the pathway

are located in the ranking; the legend at the bottom indicates whether the expression of the genes

correlate more with CD4⁺ T cells derived from WT or *Gzmb-/-* mice. (c) Real-time PCR

validation of selected genes. For day 0, n=2; for day 21, n=3-6. *P<0.05; **P<0.01. Student's t-

Test.

Figure 5. Granzyme B-deficient CD4⁺ T cells present greater *in vivo* reconstitution.

1x10⁵ CD4⁺CD45RB^hi T cells from WT (CD45.1) and *Gzmb-/-* (CD45.2) donor mice were co-

transferred into the same *Rag2-/-* recipient mice. As controls, cells from the same donors were

independently transferred into *Rag2-/-* recipient mice. Three weeks after transfer, (a) MLN

reconstitution (right, representative dot plot displaying live, TCRβ⁺CD4⁺ cells; left, summary),

and (b) Ki67 staining were determined. (c) *In vitro* proliferation potential measured by CFSE

dilution. Histograms indicate representative samples. Gray, WT; Red, *Gzmb-/-*. The gates indicate

CFSE-negative cells; graph represents summary. (d) IFNγ/IL-17 production was determined

from cells as in (a); graphs represent single IFNγ⁺ or IL-17⁺ (left and middle, respectively), and

IFNγ⁺IL-17⁺ cells (right). (e) *Rag2-/-* mice that received CD4⁺ T cells from both WT/*Gzmb-/-* or

only *Gzmb-/-* mice were weighed weekly. (f) Colon pathology was determined 3 weeks post

transfer. For (a) n=11; (b) n=9; (c) n=5; (d) n=4-9; (e), n=5-9. Each dot represents an individual

mouse. Data are representative of at least 2 independent experiments. *P<0.05; **P<0.01;

***P<0.001; ****P<0.0001. For (a) and (b) Student's t-Test; for (d) One-Way ANOVA.

Figure 6. Granzyme B-deficient mice present severe disease after *C. rodentium* infection.

Indicated mice were orogastrically infected with *C. rodentium* ($5x10^8$ CFU/mouse) and

monitored for (a) weight change, and (b) signs of disease. At 14 days post infection, colons were

dissected, (c) colonization/g of colon was determined, and (d) pathology scored as indicated in

the Methods section; left, representative micrographs (200X magnification); right, data

summary. (e) CD4$^+$ T cell IFNγ and IL-17 expression in the indicated organs was determined.

Cells were non-stimulated (ns) or stimulated (s) for 4 h with PMA/ionomycin. Each dot

represents an individual mouse. For (a-d), n=7-9; for (e), 3-5. Data are representative of at least 3

independent experiments. For (a): **P<0.01; ***P<0.01; ****P<0.001; Student's t-Test. For (b):

****P<0.001; Student's t-Test. For (d) and (e): *P<0.05; **P<0.01; One-way ANOVA.

Figure 1

167x246mm (240 x 240 DPI)

Figure 2

Figure 2

156x210mm (240 x 240 DPI)

Figure 3

Figure 3

149x117mm (240 x 240 DPI)

Figure 4

Figure 4

185x283mm (240 x 240 DPI)

Figure 5

Figure 5

157x209mm (240 x 240 DPI)

Figure 6

164x238mm (240 x 240 DPI)

Supplemental Figure 1. (a) Gating strategy for Figure 1. Plots are from representative WT

CD4$^+$ T cells cultured in Th0 conditions. Cells from *Gzmb*$^{-/-}$ mice presented similar gating

profiles (not shown). (b) Granzyme B expression in differentiated Th1 and Th2 cells from the

indicated mice; left, representative dot plots; right, summary. n=3. (c) IL-17 and IL-4 expression

in Th2 differentiated cells from the indicated mice; left, representative dot plots; right, summary.

n=3.

Supplemental Figure 2. *In vivo* activated granzyme B-deficient CD4$^+$ T cells display increased pathogenicity. (a) 1×10^5 CD4$^+$CD45RB^{hi} T cells from *Gzmb$^{+/-}$* or *Gzmb$^{-/-}$* littermate donor mice were transferred into *Rag2$^{-/-}$* recipient mice and their weights monitored weekly (left), and colon pathology scored (right). (b-e) 1×10^5 CD4$^+$CD45RB^{hi} T cells from WT or *Gzmb$^{-/-}$* donor mice were adoptively transferred into *Rag2$^{-/-}$Gzmb$^{-/-}$* recipient mice; (b) mice were weighed weekly, and (c) monitored for signs of disease. Twenty-one days after transfer, (d) colon pathology was scored (left, representative micrographs; right, data summary), and (e) donor-cell reconstitution of the IEL compartment in the colon was determined. For (a), dotted lines represent untreated *Rag2$^{-/-}$Gzmb$^{-/-}$* mice. Each dot represents an individual mouse. Data are representative of at least 3 independent experiments. n=8-9. *P<0.05; **P<0.01; ***P<0.001; ****P<0.0001; Student's t-test.

Supplemental

Figure 3

Supplemental Figure 3. Granzyme B-deficient CD4$^+$ T cells present skewed IL-17 differentiation *in vivo.* 1x10^5 CD4$^+$CD45RB^{hi} T cells from littermate *Gzmb$^{+/-}$* or *Gzmb$^{-/-}$* donor mice were adoptively transferred into *Rag2$^{-/-}$* recipient mice. Three weeks after transfer, donor-derived cells were recovered from MLN and their IFNγ/IL-17 profile was determined by intracellular staining. Dot plots indicate a representative sample. Live, TCRβ$^+$CD4$^+$ cells are displayed. Graphs represent the summary. n=3. Each dot indicates an individual mouse. Student's t-Test.

a

b

Supplemental Figure 4

Supplemental Figure 4. Naïve granzyme B-deficient mice display a normal CD4$^+$ T cell compartment. (a) CD4$^+$ T cell frequencies and cellularity from spleen, MLN, and IEL compartment were determined by flow cytometry. (b) *Ex vivo* single cytokine positive CD4$^+$ T cells derived from naïve WT and *Gzmb*$^{-/-}$ mice. Cells were non-stimulated (ns) or stimulated (s) for 4 h with PMA/ionomycin. Each dot represents an individual mouse. For (a), spleen and MLN, n=6-10; for IEL, n=12-13. For (b), n=6.

Supplemental Figure 5. (a) Absence of adaptive immune cells rescues granzyme B-deficient mice from severe disease. The indicated mice were orogastrically infected with *C. rodentium* (5×10^8 CFU/mouse) and monitored for weight change for 14 days. Data are representative of at least 2 independent experiments. n=3-4 for sham groups. n=5-7 for infected mice. (b) Granzyme B deficiency does not alter disease progression in the anti-CD40 model of colitis. The indicated mice were treated i.p. with 150 µg/mouse and weighed daily. Data are representative of at least 3 independent experiments. n=2 for sham groups. n=14 for treated group. (c-e) Littermate *Gzmb*$^{+/-}$ and *Gzmb*$^{-/-}$ mice were infected with *C. rodentium* (5×10^8 CF/mouse). (c) Animals were monitored daily for weight change. At 14 days post infection, (d) colon colonization and (e) IFNγ and IL-17 production were determined in the indicated organs. For (c and d), n=11-12; for (e), n=3-5. Student's t-Test. *$P<0.05$.