1

Composition Wheels: Visualizing dissolved organic matter using common composition metrics across a variety of Canadian Ecozones

Pieter J. K. Aukes[1,2]¶*

Sherry L. Schiff[1]¶

[1] Department of Earth & Environmental Studies, University of Waterloo, Waterloo, Ontario,

Canada

[2] Geography & Environmental Studies, Wilfrid Laurier University, Waterloo, Ontario, Canada

* Corresponding author

E-mail: paukes@uwaterloo.ca (PJKA)

¶These authors contributed equally to this work.

Abstract

Dissolved organic matter (DOM) is a ubiquitous component of aquatic systems, impacting aquatic health and drinking water quality. These impacts depend on the mixture of organic molecules that comprise DOM. Changing climates are altering both the amount and character of DOM being transported from the terrestrial system into adjacent surface waters, yet DOM composition is not monitored as often as overall concentration. Many DOM characterization methods exist, confounding comparison of DOM composition across different studies. The objective of this research is to determine which parameters in a suite of relatively simple and common DOM characterization techniques explain the most variability in DOM composition from surface and groundwater sites. Further, we create a simple visualization tool to easily compare compositional differences in DOM. A large number of water samples (n=250) was analyzed from six Canadian ecozones for DOM concentration, ultraviolet-visible light absorbance, molecular size, and elemental ratios. Principal component analyses was used to identify quasi-independent DOM compositional parameters that explained the highest variability in the dataset: spectral slope, specific-UV absorbance at 255nm, humic substances fraction, and dissolved organic carbon to dissolved organic nitrogen ratio. A 'Composition Wheel' was created by plotting these four parameters as a polygon. Our results find similarities in DOM composition irrespective of site differences in vegetation and climate. Further, two main end-member Composition Wheel shapes were revealed that correspond to DOM in organic-rich groundwaters and DOM influenced by photodegradation. The Composition Wheel approach uses easily visualized differences in polygon shape to quantify how DOM evolves by natural processes along the aquatic continuum and to track sources and degradation of DOM.

Introduction

Dissolved organic matter (DOM) is a pervasive component of aquatic environments and

an important determinant of overall water quality and ecosystem function. For example, DOM

dictates light, thermal, and pH regimes within lakes [1], complexes with and mobilizes metals

[2], and acts as an important redox constituent for biogeochemical reactions [3]. Further, DOM

affects drinking water quality through taste, odour, and colour [4], consumption of added

oxidizing chemicals, and reactions with chlorine during drinking water treatment processes to

form carcinogenic disinfection by-products [5]. The overall reactivity of DOM is determined by

its mixture of thousands of organic molecules with differing structural and chemical

characteristics. Increased DOM concentrations have been observed in surface waters across the

United Kingdom, Europe, and North America [1,6] and linked with declines to water

transparency [1,7], that may, in turn have significant effects upon future drinking water treatment

options [8,9]. Yet, little information is found on the processes that dictate DOM composition

and its impact on the surrounding environment [10]. Hence, quantifying changes to both the

amount and composition of DOM across temporal and spatial scales allows for a better

understanding of future changes to DOM and its influence upon water quality.

Both concentration and composition of DOM are used to identify the source, quality, and

fate of DOM within the environment. The overall DOM concentration is operationally

quantified by the concentration of carbon in molecules passing through a chosen filter size

(generally between 0.2 and 0.7 μm), whereas DOM composition refers to the mixture of organic

molecules that comprise it. Differences in DOM composition have been used to quantify

hydrologic mixing and changes to redox potential [11]. Various options are available to

characterize DOM composition, but many are expensive, analyze only a subfraction of the DOM,

require complicated analysis, or are not widely accessible. For instance, Fourier transform ion

cyclotron resonance mass spectrometry (FT-ICR-MS) analyses provides insightful and novel

information on DOM composition [12,13] but is not readily available. Alternatively, a number

of common and simple measures can provide compositional information on bulk DOM.

Ultraviolet and visible light absorbance are used to estimate aromaticity and molecular size of

light-absorbing DOM [14–16], while fluorescence provides information on the amount of humic

and proteinaceous DOM [17,18]. Simple elemental ratios, such as dissolved organic carbon to

dissolved organic nitrogen, indicate DOM lability [19], whereas comparison of various

molecular weight groupings can identify differences in DOM sources or processes (i.e. large

humic substances from soil-derived sources versus low-molecular weight components from

enhanced degradation) [20]. Thus, various characterization techniques can provide a holistic

representation of the components that comprise DOM.

Dissolved organic matter is the net product of varying sources and degrees of processing

at a point within the watershed. Quantifying changes to DOM composition along a hydrologic

continuum can provide information on dominant sources or processes that influence DOM

evolution [21]. For instance, DOM sampled from organic-rich substrates typically has high

concentrations of large molecular weight, aromatic DOM, while enhanced processing in

groundwaters shifts DOM towards smaller components [22–24]. Soil-derived DOM

characteristics dominate in headwater streams across various climatic regimes, as shown by

higher UV-absorbing properties and higher molecular weight components [18,21]. Conversely,

exposure to sunlight and *in-situ* DOM production within surface waters produces DOM with

lower UV-absorbing, low-molecular weight components [15]. Once in the aquatic system,

differences in DOM composition, or intrinsic controls, are thought to dictate DOM fate more-so

than extrinsic controls such as temperature, nutrients, and sunlight exposure [25]. Soil-derived

aliphatic components of DOM are preferentially lost as water moves from subsurface to surface

waters, at which point aromatic, high-nominal oxidation state DOM becomes further degraded

along the fluvial network to more aliphatic, low nominal oxidation state DOM [12,25].

Increased degradation and processing of DOM along the hydrologic continuum reduces its

chemodiversity, with persistence of specific components linked to the original DOM

composition [25–27]. Hence, compositional measures can be used to better understand how

DOM composition varies, and the potential sources or processes that have resulted in the specific

mixture of DOM at that point of sampling.

Recent progress in the understanding of DOM and carbon cycling has led to recognition

of the importance of characterizing DOM composition along with concentration. Although

characterization techniques have improved [28], the final synthesis and discussion of various

compositional DOM measures has generally relied on a matrix of scatter plots, requiring the user

to jump between graphs and build the comparison themselves. Given the direct impact of DOM

composition on water treatment and aquatic health, there is a need to relay compositional

information in an efficient and understandable manner to inform policy decisions. This can be

difficult if numerous graphs are needed in order to compare DOM differences, all with units and

scales specific to the method, especially if the target audience has little knowledge of carbon

chemistry or the characterization techniques.

Clear communication of science and its relevance to society is increasing in importance

for both informing the general public and supporting evidence-based policy decisions. Effective

data visualization can reduce the cognitive load required to understand and retain new

knowledge [29,30], creating an accessibility to science that enhances scientific literacy [31–33].

Recently, scientists and graphic designers have collaborated to utilize large datasets and translate

data-driven conclusions using novel, creative techniques [34,35]. In terms of DOM composition,

graphics have not moved beyond various scatter plots. Further, comparison of different DOM

composition metrics have generally focussed on single hydrological environments (i.e. solely

lakes [26] or rivers [36]) or on one or two measures across spatial scales or varying

environments. Given the complex mixture of organic compounds, and that different measures of

DOM composition record different attributes, the combined use of several complimentary

measures may provide a better characterization of DOM [37] and enhance public outreach

through presentation as a simple, effective image.

The objectives of this study are to select the best suite of readily accessible DOM

characterization techniques, and to create a simple, effective visual tool that can be used to

illustrate and discuss differences in DOM composition across environmental gradients. These

objectives will be accomplished in three parts: 1) determine which broadly-used measures of

DOM composition capture the most variability within a dataset of surface and groundwater sites

in various Canadian ecozones (groupings of similar areas of biodiversity [38]), 2) create a

visualization tool to allow facile comparison of DOM from different samples based upon a set of

quasi-independent composition measures, and 3) use the tool to explore DOM composition

resulting from differences in hydrological setting and climate across a range of northern

environments.

Methods

Sites & sampling

Locations were selected where DOM was expected to differ due to differences in

surrounding watershed characteristics (e.g. land use, wetland coverage, climate, and vegetation).

Various surface and groundwater samples were collected between June to October from the

Northwest Territories (Yellowknife, Wekweètì, Daring Lake), high arctic Nunavut (Lake Hazen

Watershed), and Ontario (IISD-Experimental Lakes Area) (Fig 1; S1 Table). Surface water

samples were collected 0.25 m below the surface from lakes, rivers, creeks, and ponds.

Groundwater samples from northern locations (Yellowknife, Wekweètì, Daring Lake, Lake

Hazen) were collected from the deepest extent of the mid-summer active-layer, just above the

permafrost boundary (0.1 to 0.5 m below surface). Additional samples were collected from

Turkey Lakes Watershed (ON), Nottawasaga River Watershed (ON), Grand River (ON),

Mackenzie River (NT), and Black Brook Watershed (NB) to expand the ranges of individual

composition measures but only overall DOM concentration and select composition metrics are

available at these sites (Fig 1; S1 Table). A total of 250 samples were collected and analysed in

this study.

Surface water samples were collected using a 60 mL syringe and filtered in-field using

0.45 μm syringe-tip filters (Whatman GD/X 45mm) into 40 mL acid-washed, pre-rinsed glass

vials. Groundwater samples were collected using a peristaltic pump with an attached 0.45 μm

syringe-tip filter. Vials and filters were pre-rinsed with filtered sample water before collection.

Samples were kept cool (<4ºC) and in the dark until analysis at the University of Waterloo

within three weeks of collection.

DOM quantity & composition analyses

Dissolved organic carbon and total nitrogen concentrations were measured using a

Shimadzu Total Organic Carbon (TOC-L) Combustion Analyzer with TNM-1 module.

Dissolved organic nitrogen (DON) was calculated as the difference between total dissolved

nitrogen concentration and the sum of inorganic nitrogen species (nitrate, nitrite, and

ammonium). Inorganic nitrogen species were measured using SmartChem 200 Automated

Chemistry Analyzer (Unity Scientific, MA United States). The DOC:DON ratio was calculated

using molar concentrations of DOC (M_C) and DON (M_N).

Absorbance was measured in a 1 cm cuvette using a Cary 100 UV-VIS

Spectrophotometer (Agilent, CA United States) at 5 nm increments from 200 to 800 nm.

Deionized water was used to zero the instrument and run intermittently during analyses to correct

for baseline drift. The Naperian absorption coefficient (a; m^{-1}) was calculated using:

$$a_\lambda = \frac{ln(10) \times A_\lambda}{L}$$

where A is the baseline-corrected absorbance at wavelength λ and L is the cell length (m). A

suite of absorbance characteristics were then calculated (Table 1).

Molecular-size based fractions of DOM were determined using a size exclusion

chromatography technique (Liquid Chromatography – Organic Carbon Detection, LC-OCD) at

the University of Waterloo. Detailed instrument setup and analysis is described elsewhere [20].

Briefly, the sample was injected through a size-exclusion column (SEC; Toyopearl HW-50S,

Tosoh Bioscience) that separated DOM based on hydrodynamic radii into five hydrophilic

fractions (from largest to smallest): biopolymers (BP; polysaccharides or proteins), humic

substances fraction (HSF; humic and fulvic acid fraction), building blocks (BB; lower weight

humic substances), low molecular weight neutrals (LMWN; aldehydes, small organic materials),

and LMW-acids (LMWA; saturated mono-protic acids). A portion of the sample by-passes the

SEC for determination of the overall DOC concentration, here referred to as DOM concentration

in mg C/L. A number average molecular weight was derived only for the HSF based on elution

time. Duplicates run at six concentrations yield a precision for the LC-OCD of <0.1 mg C/L for

all fractions. Concentrations of each fraction were calculated using specialized software

(ChromCALC, DOC-Labor, Germany) that integrated chromatograms from the LC-OCD.

Statistical analyses & Composition Wheel design

184   Samples from sites with multiple sampling events were averaged to create a single value

185   per site. Data were analysed using unconstrained ordination analysis via principal components

186   analysis (PCA) on a subset of samples that contained all composition measures listed in Table 1

187   (subset n=130). Data were scaled before PCA and analysed using R Statistical Software [42].

188 The Composition Wheel (CW) is a polygon drawn from axes of various composition

189 measures that are independent of DOM concentration in order to focus solely on differences in

190 DOM composition. Composition Wheel parameters were chosen based on the highest

191 contribution of variables explaining the first two principal component axes (S1 Fig). Further,

192 independent measures of DOM composition were preferentially chosen to minimize overlap in

193 information between similar techniques. Each CW axis corresponds to a specific parameter. For

194 each axis, the individual value for each sample is normalized as a value between the maximum

195 and minimum encountered for that parameter within the dataset. All R code and data used to

196 create the DOM CW, manuscript figures, and statistical analyses can be found at

197 https://github.com/paukes/DOM-Comp-Wheel.

Results

DOM concentration & composition

200 DOM concentrations ranged from 0.1 to 273 mg C/L, with highest mean values in boreal

201 groundwater, pond, and creek samples (Fig 2; S1 File). The highest DOM concentrations were

202 found in groundwater environments in Yellowknife, while the lowest concentrations were found

203 in high arctic environments (Fig 2). High arctic seeps, rivers, and lakes also generally had the

204 lowest average DOC:DON values, but higher average specific ultra-violet absorbance at 255 nm

205 (SUVA), spectral slope ($S_{275-295}$), and HSF values than other locations. SUVA and spectral slope

206 values covered the known range as described broadly in the literature (SUVA: 1.1 to 21

207 L/(mg·m); spectral slope: 0.005 to 0.032 nm^{-1}). Highest SUVA values (>11 L/(mg·m)) were

208 found at two sites: the anoxic bottom of a boreal lake and in the high arctic groundwater, but

209 absorbance may be affected by high iron concentrations [14,43]. These two sites have unusually

high iron concentrations and were not included further in the discussion. Values of DOC:DON

ranged between 9 to 124, and were lowest in rivers and high arctic samples. Humic substances

fraction (HSF) ranged from 14% to 85% and on average were lowest in lakes. Overall,

groundwater samples contained the highest DOC:DON and HSF values, and lowest $S_{275-295}$.

Hence, both DOM concentration and composition varied across geographic scales and

hydrological environments.

PCA on DOM composition measures

The first three principal component axes explained 66% of the variance in DOM

composition using the measures contained in the dataset, with PC1 and PC2 accounting for 54%

of the variability (Fig 3). Comparison of the first two principal components (PC) yield four

distinct groupings of strongly-contributing measures: I) SUVA, SAC_{420}, SAC_{350}; II) HSF, HS

MW; III) $S_{275-295}$, $E_2:E_3$, and S_R; and IV) BB, LMWN, and BP. Groups I and II were negatively

associated to groups III and IV. Highest contributions to PC1 and PC2 axes were HSF, SAC_{350},

SAC_{420}, SUVA, and $S_{275-295}$. Variables with contribution to PC1 and PC2 lower than 2% were

$E_4:E_6$, DOC:DON, $S_{350-400}$, and LMWA (Fig. S1). Absorbance parameters normalized to DOM

concentrations (SAC_{350}, SAC_{420}, and SUVA) all plotted closely to each other and trended

positively with measures of HSF, HSF molecular weight, and DOC:DON. Absorbance

techniques plotted perpendicular to LC-OCD size fractions (Fig 3).

Based on contribution to the first two PC axes, we selected four DOM composition

measures to define DOM composition. Further, these selected measures are quasi-independent

as they are based on different analytical principles or attributes (SUVA, $S_{275-295}$: two different

absorbance aspects; HSF: size-exclusion chromatography; DOC:DON: stoichiometry).

Although the absorbance at different wavelengths within a sample are closely related [21], they

can be used to provide different information on DOM components as one characterizes the

amount of UV-absorbing components normalized to DOM concentration at a specific

wavelength (SUVA) while the other is associated with the shape of the absorbance spectra and

varies with the distribution of DOM molecular weights ($S_{275-295}$). These four measures include

three high-contributing variables to PC1 and PC2 axes (HSF, SUVA, and $S_{275-295}$). The length of

the CW axis, and thus CW size, is dictated by the range of each compositional measure

encountered in the entire dataset. Composition Wheels are then used as a basis for comparing

DOM.

Shapes of Composition Wheels

Different environments are characterized by CW different shapes. Ponds and lakes tend

to have lower values of SUVA and DOC:DON and higher $S_{275-295}$, resulting in a triangle shape

elongated to the top-right corner (Fig 4). Groundwater DOM samples form a trapezoidal shape

due to higher HSF and SUVA values than other sites. Differences in shape are found within

locations between different hydrologic settings whereas similarities in shape are found within

hydrological setting across locations. For instance, Yellowknife pond sites are similar in shape

to groundwater DOM, while the creek, lake, and river shapes reflect DOM with higher $S_{275-295}$,

lower DOC:DON, and lower HSF. Representation of different measures using these shapes

allows for a facile comparison of DOM composition among samples.

Shapes also help to visualize and highlight the most extremes in DOM composition for

each location. The spread of compositions is not evenly distributed between such extreme

shapes at any given site but sampling was not targeted *a priori* based on shape. Most surface

water sites contained a range in DOM composition between these extremes, forming either

shapes similar to groundwater DOM or thin, elongated triangles in the $S_{275-295}$ direction. For this

reason, extremes were highlighted in sites with many different shapes to help identify a

continuum of possible DOM sources or mixtures (Fig 4). Further, some sites did not conform to

these end-members, as seen by the ELA creek sample with high DOC:DON, allowing easy

identification of anomalous data or sites that may warrant further investigation.

Discussion

Comparison of DOM composition measures

Ranges in individual measures of DOM composition are large at both low and high

concentrations of DOM (Fig 4, S2 Table) making it difficult to consistently separate sites or

hydrological environments across different measures. For instance, accounting for both DOM

concentration and DOC:DON may help differentiate some locations, but the range in DOC:DON

values across moderate DOM concentrations covers most sites (Fig 2). Hence, new techniques

that simultaneously consider different aspects of DOM composition, and are independent of

concentration, are needed to compare variations in DOM composition across sample types and

environments.

Different characterization techniques provide different information on DOM. In

particular, the perpendicular relationship between absorbing and non-absorbing parameters

observed among the first two PC axes indicate a range in different DOM properties are captured.

For instance, all ELA creeks contain high and similar HSF values, yet we can use absorbance

techniques based on molecular weight or UV-absorbing capability (via SUVA; Fig 2) to

differentiate between these samples. However, correlations between size-based and absorbance

parameters do not mean causation. For example, higher SUVA values are not consistently

related to HSF proportions even though this is often assumed based on the higher aromaticity

and UV-absorbing capabilities of humic material [14,45]. Daring Lake creeks have high SUVA

but low HSF. Although absorbance parameters are easily measured and helpful in comparing

different DOM compositions, they do not accurately reflect non-absorbing components (e.g.

stoichiometry or functional groups) that could differ among samples.

Effects of DOM Processes and CW End Members

Similar DOM CW shapes were found within hydrologic settings even though samples

were collected from sites with varying climates and vegetation, spanning areas of boreal shield

watersheds (IISD-ELA) to high arctic (Lake Hazen). Further, the range in values across the four

CW measures compared well with DOM from other studies (i.e. similar DOM concentrations

and SUVA values to rivers in the United States [16,44] and Canadian boreal lakes [46])

indicating that DOM composition and CW shape may not be unique to its locale. The similarity

in CW shapes across different locations may result from analogous drivers of DOM fate. For

instance, the proportion of biodegradable DOM was found to be a function of DOM composition

and nutrients, rather than ecosystem or region-specific characteristics [47], suggesting that

differences in the quality of DOM may not result from location alone. Hence, CWs can be used

to amalgamate different compositional measures and quantify similarities in DOM sources and

processing across a range of geomorphic, climate, and vegetation environments.

Composition Wheels can be further used to determine the degree of DOM degradation and

facilitate comparisons of change along the aquatic continuum. This requires knowledge of the

effect of processing, as well as the end-members (e.g. original sources of DOM). Microbial

degradation of DOM induces a shift towards higher SUVA and lower DOC:DON (Fig 5; [48–51]

but not $S_{275-295}$, resulting in a distinctive effect of microbial DOM transformation on the CW.

Photolytic processing of DOM results in an increase in $S_{275-295}$ values and a general decrease in

the other three DOM quality measures [52–54]. Further, photolytic degradation affects all four

compositional axes, whereas microbial degradation only affects two axes (Fig 5), indicating that

certain measures of DOM composition respond differently to different DOM degradation

processes. This has important implications for studies using single-characterization techniques

as some parameters may not always faithfully serve as useful surrogates for others. For instance,

microbial degradation changes SUVA but not HSF (Fig 5), even though SUVA has been linked

to DOM aromaticity in humic substances [14]. Thus, comparative differences in a CW can help

determine the dominant processes that produce the observed DOM composition of a sample.

Further, knowing the degree DOM composition changes with photolysis allows CW to be used

in a quantitative manner to quantify the amount of degradation. Although the quantitative

response of DOM sources to photo- or microbial degradation is not yet sufficiently known across

different environments, in areas where these rates of change to individual measures have been

measured, CWs can be used in a quantitative manner to assess the extent and major processes

contributing to DOM loss.

Laboratory experiments are useful to assess DOM source composition prior to extensive

biotic or abiotic processing. Although this study did not conduct analysis on leachates, various

studies have used similar DOM composition measures to identify DOM leachates. Leachates

(organic matter steeped in de-ionized water) result in either low molecular weight, low aromatic

DOM from mosses and coniferous sources [22,55] or large, humic, aromatic DOM components

from other plant material and deciduous leaf litter [37,57,58]. DOM produced from aquatic

(primary production) consists of low molecular weight, high protein and nitrogen content, and

low UV-absorbing components and is thought to be rapidly consumed [37,56,59,60]. These

parameters can also be represented by a CW (Figure 5b – using a range in values based on what

is presented in literature), and provide a shape to compare DOM origins from aquatic primary

production versus terrestrial plants.

Two dominant DOM compositional end-member shapes are evident in our dataset:

groundwater DOM sources and DOM altered by photolysis. Other shapes are intermediate

between these two distinctive end-members. Degradation in groundwater in organic-rich

substrates allows for the accumulation of large, aromatic components of DOM (high HSF,

SUVA, and low $S_{275-295}$; [61]), resulting in a large trapezoid shape CW. This trapezoid shape is

consistent for groundwater DOM across most sites, supporting previous groundwater studies that

attribute the narrow DOM compositional range to groundwater processing [23]. In contrast, a

'kite-like' shape results from the effects of photolysis (Fig 4), resulting in DOM with lower

molecular weight and less UV-absorbing components [15]. Further, photolysis end-member

CWs are similar in shape to the end-products of photolytic experiments (Fig 5; [51]), and are

clearly seen in surface waters across all sites (Fig 4). Hence, these end-member shapes may

constrain the possible range in CW shapes that can be found within aquatic environments.

The groundwater/photolysis end-member categorization frames our conceptual model (Fig 6)

that quantifies how the two end-member CW shapes evolve depending on the aquatic-terrestrial

hydrologic connection (Fig 6; pathway A-b) or exposure to sunlight (Fig 6: pathway A-B). For

instance, creeks with little to no processing of groundwater-derived DOM can be identified by

CW shapes similar to groundwater DOM, as seen in ELA and some Daring creeks (Fig 4). ELA

lakes have a wide gradient of CW shape and thus overall DOM quality (Fig 4). The use of CW

to pair process-based knowledge of DOM composition with hydrologic transport provides a

framework that can be used identify key knowledge gaps (i.e. rates of degradation, kinetics based

on composition, relative organic source contributions, water residence time) and quantify the

relationship between DOM sources and degree of processing within the environment.

Certain creeks and rivers from YK or WK do not fit into the end-member categories as they

contain shapes with larger right sides (higher $S_{275-295}$ and SUVA than other samples), indicating

other processes may be important in determining DOM composition. This may be a result of our

sampling of mostly oligotrophic systems that are dominated by terrestrial carbon. We

hypothesize that these different shapes represent surface water systems with greater *in-situ*, or

autochthonous, contributions of DOM, generally characterized by higher amounts of proteins

and smaller molecular weight components [62–64]. Thus, although our dataset was not

comprehensive in terms of capturing the full range of DOM sources, the CW-informed

conceptual model provides an initial framework to build the factors leading to a specific DOM

composition.

Adaptability of DOM Composition Wheels

Expressing DOM composition with only four concentration-independent parameters

excludes advantages of other techniques not used in this study. However, agreement between

multiple parameters allows for selection of surrogates for the CW. LC-OCD is not as widely

available but many studies have traditionally used resins [16,65,66] or other size-exclusion

columns [67,68] to characterize DOM. Further, LC-OCD fractions of humics have been well

correlated to measures of ^{13}C-NMR and fluorescence measures such as HIX [12,69,70].

Although fluorescence was not used in this study, strong associations between certain

fluorescence parameters and absorbance or molecular-weight groupings could be used to replace

LC-OCD defined fractions, such as using PARAFAC modelling to discern components most

similar to HSF [12,70] (Fig 7). These fluorescence parameters are also independent measures of

DOM composition and could be readily substituted into the CW. Elemental ratios of DOC:DON

are generally positively correlated with humic-like fluorescence and negatively correlated to

protein-like components [71]. Other absorbance indices such as slope ratio and E_2:E_3 (Table 1)

are potential surrogates for $S_{275\text{-}295}$ (Fig 2), providing various parameters that can be substituted

for one another when comparing studies where only one of these measures is provided.

Measurements of SUVA could be substituted with SAC_{420}, SAC_{350}, or E_4:E_6 [25,71]. However,

although SUVA has been correlated with HIX and to fluorescence component C3 [25,71], the

opposite has also been found [70]. Composition Wheels can easily incorporate advanced

techniques, such as FT-ICR-MS, that can provide information on the molecular heterogeneity of

DOM, identification of either terrestrial-like (via condensed and polycyclic aromatics or

polyphenolic compounds) or *in situ* DOM sources (higher aliphatic and peptide-like

compounds), or to identify differences in H:C and O:C due to microbial or photodegradation

[12,13,72]. Associations between different characterization techniques allow for mixing of

techniques and comparison of different variables, indicating a wide-range of applicability of CW

within environmental sciences.

The CW visualization method provides an efficient communication tool not only among

scientists, but also between scientists and other stakeholders concerned with water quality,

including community members. Shapes from CWs show clearly how DOM differs, whereas

changes in the numerical value of multiple metrics are only easily grasped by those with

familiarity with the methods and environmental ranges. Further work could include pairing

different shapes of DOM with key DOM roles such as disinfection demand and by-product

formation, metal mobility, and mercury bioaccumulation. By reducing the complexity of

independent DOM measures and creating an easily comparable shape, differences in DOM

composition can be easily grasped and communicated to larger audiences.

Acknowledgements

We thank Michael English, Roy Judas, and Jordan Reid for field assistance in collecting samples

in the Northwest Territories; Paul Dainard, Kyra St. Pierre, Igor Lehnherr, Vince St. Louis,

Jessica Serbu, Maria Cavacao, Victoria Wisniewski, and Michael English for field assistance at

Lake Hazen, NU; Will Robertson for assistance at Long Point, ON; Environmental

Geochemistry Lab, University of Waterloo for sampling the Grand River, ON; John Spoelstra for

sampling Black Brook Watershed, Nottawasaga River Watershed, and Turkey Lakes Watershed.

We also thank the Community of Wekweètì and the Wek'èezhìi Land & Water Board for their

support, and the Community-Based Monitoring Program and Department of Environment and

Natural Resources, Government of Northwest Territories, for collection of samples from the

Mackenzie River, NT. We would like thank Richard J. Elgood for logistical, field, and technical

help, Monica Tudorancea assistance with the LC-OCD analysis, and Mackenzie Schultz for help

with Figure 1. We are grateful for the high arctic logistic support provided by Parks Canada, and

funding from the Polar Continental Shelf Program (PCSP), Northwest Territories Cumulative

Impact Monitoring Program, Northern Scientific Training Program, and the National Science

and Engineering Research Council of Canada. We thank Igor Lehnherr, Susan Ziegler, Chris

Parsons, Roland Hall, and three anonymous reviewers for their comments on earlier drafts of this

manuscript. PJKA also benefitted from scholarships from the Garfield Weston Foundation, TD

Friends of the Environment, and the University of Waterloo. Data used in this manuscript can be

found in the publicly-available Wilfrid Laurier University Scholars Portal Dataverse

(https://doi.org/10.5683/SP2/NG6D02), while all R code used for statistical analyses, figure

creation, and a Composition Wheel template is available from https://github.com/paukes/DOM-

Comp-Wheel.

References

1.   Strock KE, Theodore N, Gawley WG, Ellsworth AC, Saros JE. Increasing dissolved
organic carbon concentrations in northern boreal lakes: Implications for lake water
transparency and thermal structure. J Geophys Res Biogeosciences. 2017;122: 1022–1035.
doi:10.1002/2017JG003767
2.   Baken S, Degryse F, Verheyen L, Merckx R, Smolders E. Metal complexation properties
of freshwater dissolved organic matter are explained by its aromaticity and by
anthropogenic ligands. Environ Sci Technol. 2011;45: 2584–2590. doi:10.1021/es103532a
3.   Bauer M, Heitmann T, Macalady DL, Blodau C. Electron transfer capacities and reaction
kinetics of peat dissolved organic matter. Environ Sci Technol. 2007;41: 139–145.
doi:10.1021/es061323j
4.   Matilainen A, Gjessing ET, Lahtinen T, Hed L, Bhatnagar A, Sillanpää M. An overview
of the methods used in the characterisation of natural organic matter (NOM) in relation to
drinking water treatment. Chemosphere. 2011;83: 1431–1442.
doi:10.1016/j.chemosphere.2011.01.018
5.   Krasner SW, Weinberg HS, Richardson SD, Pastor SJ, Chinn R, Sclimenti MJ, et al.
Occurrence of a New Generation of Disinfection Byproducts. Environ Sci Technol.
2006;40: 7175–7185. doi:10.1021/es060353j
6.   Evans CD, Monteith DT, Cooper DM. Long-term increases in surface water dissolved
organic carbon: observations, possible causes and environmental impacts. Environ Pollut.
2005;137: 55–71. doi:10.1016/j.envpol.2004.12.031
7.   Weyhenmeyer GA, Prairie YT, Tranvik LJ. Browning of boreal freshwaters coupled to
carbon-iron interactions along the aquatic continuum. PLoS One. 2014;9: e88104.
doi:10.1371/journal.pone.0088104
8.   Worrall F, Burt TP. Changes in DOC treatability: Indications of compositional changes in
DOC trends. J Hydrol. 2009;366: 1–8. doi:10.1016/j.jhydrol.2008.11.044
9.   Ritson JP, Graham NJD, Templeton MR, Clark JM, Gough R, Freeman C. The impact of
climate change on the treatability of dissolved organic matter (DOM) in upland water
supplies: A UK perspective. Sci Total Environ. 2014;473–474: 714–730.
doi:10.1016/j.scitotenv.2013.12.095
10.   Creed IF, Bergström AK, Trick CG, Grimm NB, Hessen DO, Karlsson J, et al. Global
change-driven effects on dissolved organic matter composition: Implications for food
webs of northern lakes. Glob Chang Biol. 2018;24: 3692–3714. doi:10.1111/gcb.14129
11.   Larsen LG, Aiken GR, Harvey JW, Noe GB, Crimaldi JP. Using fluorescence
spectroscopy to trace seasonal DOM dynamics, disturbance effects, and hydrologic
transport in the Florida Everglades. J Geophys Res. 2010;115: 1–14.
doi:10.1029/2009JG001140
12.   Hutchins RHS, Aukes PJK, Schiff SL, Dittmar T, Prairie YT, del Giorgio PA. The
Optical, Chemical, and Molecular Dissolved Organic Matter Succession Along a Boreal
Soil-Stream-River Continuum. J Geophys Res Biogeosciences. 2017;122: 2892–2908.

456   doi:10.1002/2017JG004094

457   13.   Kellerman AM, Guillemette F, Podgorski DC, Aiken GR, Butler KD, Spencer RGM.
458   Unifying Concepts Linking Dissolved Organic Matter Composition to Persistence in
459   Aquatic Ecosystems. Environ Sci Technol. 2018;52: 2538–2548.
460   doi:10.1021/acs.est.7b05513

461   14.   Weishaar JL, Aiken GR, Bergamaschi BA, Fram MS, Fujii R, Mopper K. Evaluation of
462   specific ultraviolet absorbance as an indicator of the chemical composition and reactivity
463   of dissolved organic carbon. Environ Sci Technol. 2003;37: 4702–4708.
464   doi:10.1021/es030360x

465   15.   Helms JR, Stubbins A, Ritchie JD, Minor EC, Kieber DJ, Mopper K. Absorption spectral
466   slopes and slope ratios as indicators of molecular weight, source, and photobleaching of
467   chromophoric dissolved organic matter. Limnol Oceanogr. 2008;53: 955–969.
468   doi:10.4319/lo.2008.53.3.0955

469   16.   Spencer RGM, Butler KD, Aiken GR. Dissolved organic carbon and chromophoric
470   dissolved organic matter properties of rivers in the USA. J Geophys Res Biogeosciences.
471   2012;117. doi:10.1029/2011JG001928

472   17.   Stedmon C a, Markager S, Bro R. Tracing dissolved organic matter in aquatic
473   environments using a new approach to fluorescence spectroscopy. Mar Chem. 2003;82:
474   239–254. doi:10.1016/S0304-4203(03)00072-0

475   18.   Jaffé R, Yamashita Y, Maie N, Cooper WT, Dittmar T, Dodds WK, et al. Dissolved
476   Organic Matter in Headwater Streams: Compositional Variability across Climatic Regions
477   of North America. Geochim Cosmochim Acta. 2012;94: 95–108.
478   doi:10.1016/j.gca.2012.06.031

479   19.   Hunt AP, Parry JD, Hamilton-Taylor J. Further evidence of elemental composition as an
480   indicator of the bioavailability of humic substances to bacteria. Limnol Oceanogr.
481   2000;45: 237–241.

482   20.   Huber SA, Balz A, Abert M, Pronk W. Characterisation of aquatic humic and non-humic
483   matter with size-exclusion chromatography – organic carbon detection – organic nitrogen
484   detection (LC-OCD-OND). Water Res. 2011;45: 879–885.
485   doi:10.1016/j.watres.2010.09.023

486   21.   Massicotte P, Asmala E, Stedmon C, Markager S. Global distribution of dissolved organic
487   matter along the aquatic continuum: Across rivers, lakes and oceans. Sci Total Environ.
488   2017;609: 180–191. doi:10.1016/j.scitotenv.2017.07.076

489   22.   Wickland KP, Neff JC, Aiken GR. Dissolved Organic Carbon in Alaskan Boreal Forest:
490   Sources, Chemical Characteristics, and Biodegradability. Ecosystems. 2007;10: 1323–
491   1340. doi:10.1007/s10021-007-9101-4

492   23.   Shen Y, Chapelle FH, Strom EW, Benner R. Origins and bioavailability of dissolved
493   organic matter in groundwater. Biogeochemistry. 2015;122: 61–78. doi:10.1007/s10533-
494   014-0029-4

495   24.   Aukes PJK, Schiff SL, Robertson WD. Evolution of Dissolved Organic Matter Along a
496   Septic System Plume: Evidence of Sustained Biogeochemical Activity in the Groundwater
497   Zone. J Geophys Res Biogeosciences. 2019;124: 1389–1400. doi:10.1029/2018JG004758

498   25.   Kellerman AM, Kothawala DN, Dittmar T, Tranvik LJ. Persistence of dissolved organic
499   matter in lakes related to its molecular characteristics. Nat Geosci. 2015;8.
500   doi:10.1038/ngeo2440

501   26.   Kellerman AM, Dittmar T, Kothawala DN, Tranvik LJ. Chemodiversity of dissolved

502   organic matter in lakes driven by climate and hydrology. Nat Commun. 2014;5: 1–8.
503   doi:10.1038/ncomms4804

504 27. Creed IF, Mcknight DM, Pellerin BA, Green MB, Bergamaschi BA, Aiken GR, et al. The
505   river as a chemostat : fresh perspectives on dissolved organic matter flowing down the
506   river continuum. Can J Fish Aquat Sci. 2015;14: 1–14.

507 28. McCallister SL, Ishikawa NF, Kothawala DN. Biogeochemical tools for characterizing
508   organic carbon in inland aquatic ecosystems. Limnol Oceanogr Lett. 2018;3: 444–457.
509   doi:10.1002/lol2.10097

510 29. Pinker S. Theory of Graph Comprehension. In: Freedle R, editor. Artificial intelligence
511   and the future of testing. Lawrence Erlbaum Associates, Inc.; 1990. pp. 73–126.

512 30. Zhu B, Watts SA. Visualization of network concepts: The impact of working memory
513   capacity differences. Inf Syst Res. 2010;21: 327–344. doi:10.1287/isre.1080.0215

514 31. Gordin DN, Pea RD. Prospects for Scientific Visualization as an Educational Technology.
515   J Learn Sci. 1995;4: 249–279.

516 32. Pauwels L. An integrated model for conceptualising visual competence in scientific
517   research and communication. Vis Stud. 2008;23: 147–161.
518   doi:10.1080/14725860802276305

519 33. Martin LJ, Turnquist A, Groot B, Huang SYM, Kok E, Thoma B, et al. Exploring the Role
520   of Infographics for Summarizing Medical Literature. Heal Prof Educ. 2019;5: 48–57.
521   doi:10.1016/j.hpe.2018.03.005

522 34. Shamoun-Baranes J, Farnsworth A, Aelterman B, Alves JA, Azijn K, Bernstein G, et al.
523   Innovative visualizations shed light on avian nocturnal migration. PLoS One. 2016;11: 1–
524   15. doi:10.1371/journal.pone.0160106

525 35. Lupi G, Accurat. Beautiful Reasons - Engaging Aesthetics for Data Narratives [Internet].
526   2016 [cited 18 Jan 2020]. Available: https://medium.com/accurat-studio/beautiful-reasons-
527   c1c6926ab7d7

528 36. Lundin EJ, Klaminder J, Bastviken D, Olid C, Hansson S V., Karlsson J. Large difference
529   in carbon emission - Burial balances between boreal and arctic lakes. Sci Rep. 2015;5: 1–
530   7. doi:10.1038/srep14248

531 37. Hansen AM, Kraus TEC, Pellerin BA, Fleck JA, Downing BD, Bergamaschi BA. Optical
532   properties of dissolved organic matter (DOM): Effects of biological and photolytic
533   degradation. Limnol Oceanogr. 2016;61: 1015–1032. doi:10.1002/lno.10270

534 38. Marshall IB, Schut PH, Ballard M. A National Ecological Framework for Canada:
535   Attribute Data [Internet]. Ottawa, ON, Canada: Ecosystem Stratification Working Group,
536   Agriculture and Agri-Food Canada & Environment Canada; 1999. Available:
537   http://sis.agr.gc.ca/cansis/nsdb/ecostrat/1999report/index.html

538 39. Dahlén J, Bertilsson S, Pettersson C. Effects of UV-A radiation on dissolved organic
539   matter in humic surface waters. Environ Int. 1996;22: 501–506.

540 40. Chin Y-P, Aiken GR, O'Loughlin E. Molecular Weight, Polydispersity, and Spectroscopic
541   Properties of Aquatic Humic Substances. Environ Sci Technol. 1994;28: 1853–1858.
542   doi:10.1021/es00060a015

543 41. Curtis PJ, Schindler DW. Hydrologic control of dissolved organic matter in low-order
544   Precambrian Shield lakes. Biogeochemistry. 1997;36: 125–138.
545   doi:10.1023/A:1005787913638

546 42. R Core Team. R: A language and environment for statistical computing. [Internet].
547   Vienna, Austria: R Foundation for Statistical Computing; 2020. Available: http://www.r-

548   project.org

549 43. Poulin B a, Ryan JN, Aiken GR. Effects of Iron on Optical Properties of Dissolved
550   Organic Matter. Environ Sci Technol. 2014; 1–20. doi:10.1021/es502670r

551 44. Jaffé R, McKnight D, Maie N, Cory RM, McDowell WH, Campbell JL. Spatial and
552   temporal variations in DOM composition in ecosystems: The importance of long-term
553   monitoring of optical properties. J Geophys Res. 2008;113: 1–15.
554   doi:10.1029/2008JG000683

555 45. Hua G, Reckhow DA, Abusallout I. Correlation between SUVA and DBP formation
556   during chlorination and chloramination of NOM fractions from different sources.
557   Chemosphere. 2015;130: 82–89. doi:10.1016/j.chemosphere.2015.03.039

558 46. Mueller KK, Fortin C, Campbell PGC. Spatial Variation in the Optical Properties of
559   Dissolved Organic Matter (DOM) in Lakes on the Canadian Precambrian Shield and
560   Links to Watershed Characteristics. Aquat Geochemistry. 2012;18: 21–44.
561   doi:10.1007/s10498-011-9147-y

562 47. Lapierre J-F, del Giorgio PA. Partial coupling and differential regulation of biologically
563   and photochemically labile dissolved organic carbon across boreal aquatic networks.
564   Biogeosciences. 2014;11: 5969–5985. doi:10.5194/bg-11-5969-2014

565 48. Koehler B, Von Wachenfeldt E, Kothawala DN, Tranvik LJ. Reactivity continuum of
566   dissolved organic carbon decomposition in lake water. J Geophys Res Biogeosciences.
567   2012;117: 1–14. doi:10.1029/2011JG001793

568 49. McDowell WH, Zsolnay A, Aitkenhead-Peterson J, Gregorich EG, Jones DL, Jödemann
569   D, et al. A comparison of methods to determine the biodegradable dissolved organic
570   carbon from different terrestrial sources. Soil Biol Biochem. 2006;38: 1933–1942.
571   doi:10.1016/j.soilbio.2005.12.018

572 50. Kalbitz K, Schwesig D, Schmerwitz J, Kaiser K, Haumaier L, Glaser B, et al. Changes in
573   properties of soil-derived dissolved organic matter induced by biodegradation. Soil Biol
574   Biochem. 2003;35: 1129–1142. doi:10.1016/S0038-0717(03)00165-2

575 51. Aukes PJK. Dissolved Organic Matter in the Canadian Arctic & Sub-Arctic: Importance
576   of DOM Quality & Quantity in a Warming Climate. University of Waterloo. 2019.

577 52. Winter AR, Fish TAE, Playle RC, Smith DS, Curtis PJ. Photodegradation of natural
578   organic matter from diverse freshwater sources. Aquat Toxicol. 2007;84: 215–222.
579   doi:10.1016/j.aquatox.2007.04.014

580 53. Franke D, Hamilton MW, Ziegler SE. Variation in the photochemical lability of dissolved
581   organic matter in a large boreal watershed. Aquat Sci. 2012;74: 751–768.
582   doi:10.1007/s00027-012-0258-3

583 54. Mann PJ, Davydova a., Zimov NS, Spencer RGM, Davydov S, Bulygina E, et al.
584   Controls on the composition and lability of dissolved organic matter in Siberia's Kolyma
585   River basin. J Geophys Res. 2012;117: G01028. doi:10.1029/2011JG001798

586 55. Sun L, Perdue EM, Meyer JL, Weis J. Use of elemental composition to predict
587   bioavailability dissolved organic matter in a Georgia river. Limnol Oceanogr. 1997;42:
588   714–721. doi:10.4319/lo.1997.42.4.0714

589 56. Catalán N, Obrador B, Felip M, Pretus JL. Higher reactivity of allochthonous vs.
590   autochthonous DOC sources in a shallow lake. Aquat Sci. 2013;75: 581–593.
591   doi:10.1007/s00027-013-0302-y

592 57. Lee M-H, Hur J. Photodegradation-Induced Changes in the Characteristics of Dissolved
593   Organic Matter with Different Sources and Their Effects on Disinfection By-Product

Formation Potential. CLEAN - Soil, Air, Water. 2014;42: 552–560.
doi:10.1002/clen.201200685

58. Singh S, Inamdar S, Mitchell MJ, McHale P. Seasonal pattern of dissolved organic matter
(DOM) in watershed sources : influence of hydrologic flow paths and autumn leaf fall.
Biogeochemistry. 2014;118: 321–337. doi:10.1007/s10533-013-9934-1

59. Pivokonsky M, Safarikova J, Baresova M, Pivokonska L, Kopecka I. A comparison of the
character of algal extracellular versus cellular organic matter produced by
cyanobacterium, diatom and green alga. Water Res. 2014;51: 37–46.
doi:10.1016/j.watres.2013.12.022

60. McKnight DM, Andrews ED, Spaulding SA, Aiken GR. Aquatic fulvic acids in algal-rich
antarctic ponds. Limnol Oceanogr. 1994;39: 1972–1979. doi:10.4319/lo.1994.39.8.1972

61. Wauthy M, Rautio M, Christoffersen KS, Forsström L, Laurion I, Mariash HL, et al.
Increasing dominance of terrigenous organic matter in circumpolar freshwaters due to
permafrost thaw. Limnol Oceanogr Lett. 2018; doi:10.1002/lol2.10063

62. Tank SE, Lesack LFW, Gareis J a. L, Osburn CL, Hesslein RH. Multiple tracers
demonstrate distinct sources of dissolved organic matter to lakes of the Mackenzie Delta,
western Canadian Arctic. Limnol Oceanogr. 2011;56: 1297–1309.
doi:10.4319/lo.2011.56.4.1297

63. Roiha T, Peura S, Cusson M, Rautio M. Allochthonous carbon is a major regulator to
bacterial growth and community composition in subarctic freshwaters. Sci Rep. 2016;6:
34456. doi:10.1038/srep34456

64. Hood E, Fellman JB, Spencer RGM, Hernes PJ, Edwards R, D'Amore D V., et al.
Glaciers as a source of ancient and labile organic matter to the marine environment.
Nature. 2009;462: 1044–7. doi:10.1038/nature08580

65. Cleveland CC, Neff JC, Townsend AR, Hood E. Composition, Dynamics, and Fate of
Leached Dissolved Organic Matter in Terrestrial Ecosystems: Results from a
Decomposition Experiment. Ecosystems. 2004;7: 275–285. doi:10.1007/s10021-003-
0236-7

66. Bourbonniere R. Distribution patterns of dissolved organic matter fractions in natural
waters from eastern Canada. Org Geochem. 1989;14: 97–107. doi:10.1016/0146-
6380(89)90023-5

67. Her N, Amy GL, Foss D, Cho J, Yoon Y, Kosenka P. Optimization of method for
detecting and characterizing NOM by HPLC-size exclusion chromatography with UV and
on-line DOC detection. Environ Sci Technol. 2002;36: 1069–76. Available:
http://www.ncbi.nlm.nih.gov/pubmed/11917993

68. Kawasaki N, Matsushige K, Komatsu K, Kohzu A, Nara FW, Ogishi F, et al. Fast and
precise method for HPLC-size exclusion chromatography with UV and TOC (NDIR)
detection: importance of multiple detectors to evaluate the characteristics of dissolved
organic matter. Water Res. 2011;45: 6240–6248. doi:10.1016/j.watres.2011.09.021

69. Romera-Castillo C, Chen M, Yamashita Y, Jaffé R. Fluorescence characteristics of size-
fractionated dissolved organic matter: Implications for a molecular assembly based
structure? Water Res. 2014;55C: 40–51. doi:10.1016/j.watres.2014.02.017

70. Bodmer P, Heinz M, Pusch M, Singer GA, Premke K. Carbon dynamics and their link to
dissolved organic matter quality across contrasting stream ecosystems. Sci Total Environ.
2016;553: 574–586. doi:10.1016/j.scitotenv.2016.02.095

71. Kothawala DN, Stedmon CA, Müller RA, Weyhenmeyer GA, Köhler SJ, Tranvik LJ.

Controls of dissolved organic matter quality: Evidence from a large-scale boreal lake
survey. Glob Chang Biol. 2014;20: 1101–1114. doi:10.1111/gcb.12488
72.   Ward CP, Nalven SG, Crump BC, Kling GW, Cory RM. Photochemical alteration of
organic carbon draining permafrost soils shifts microbial metabolic pathways and
stimulates respiration. Nat Commun. 2017; 1–8. doi:10.1038/s41467-017-00759-2
645
646

Tables

Table 1. Dissolved organic matter composition as described by chemical, absorbance, and molecular-size based measures used in this study.

Measure (*common abbreviation*)	Equation	Unit	Characteristic	Reference
Chemical				
DOC:DON	$M_C \div M_N$	-	Stoichiometry	[19]
Absorbance				
E$_2$:E$_3$	$A_{255} \div A_{365}$	-	Inversely related to molecular size	[39]
E$_4$:E$_6$	$A_{465} \div A_{665}$	-	Humic molecular weight and size	[40]
SAC$_{350}$	$\dfrac{ln(10) \times A_{420}/L}{[DOC]}$	$\dfrac{cm^2}{mg\text{-}C} \div 10^3$	Specific absorbance at 350 nm and attributed to colour	[41]
SAC$_{420}$	$\dfrac{ln(10) \times A_{420}/L}{[DOC]}$	$\dfrac{L}{mg \times m}$	Specific absorbance at 420 nm	
SUVA	$\dfrac{ln(10) \times A_{255}/L}{[DOC]}$	$\dfrac{L}{mg \times m}$	Correlated to degree of aromaticity	[14]
S$_{275\text{-}295}$	$a_\lambda = a_{\lambda_{275}} e^{-S(\lambda - 275)}$	nm^{-1}	Inversely related to MW (where S is the slope)	[15]
S$_{350\text{-}400}$	$a_\lambda = a_{\lambda_{350}} e^{-S(\lambda - 350)}$	nm^{-1}	Used with S$_{275\text{-}295}$ to calculated S$_R$	[15]
S$_R$	$\dfrac{S_{275-295}}{S_{350-400}}$	-	Inversely related to overall molecular weight; integrative indicator of light-absorbing DOM history	[15]
Size Exclusion Chromatography				
Biopolymers (BP)	-	% of DOM	Polysaccharides, proteins	[20]
Humic substances (HSF)	-	% of DOM	Humic or fulvic-like components	[20]
Building Blocks (BB)	-	% of DOM	Degraded HS	[20]
Low-molecular weight neutrals (LMWN)	-	% of DOM	Aldehydes, small organic materials	[20]
Low-molecular weight acids (LMWA)	-	% of DOM	Saturated mono-protic acids	[20]
HS molecular weight (HS MW)	-	g / mol	Nominal average molecular weight of HS	[20]

Figure

Fig 1. Locations of sampling sites and ecozones. River locations (Grand River, ON; Mackenzie River, NT) are labelled at the mouth of the river.

659
Fig 2. Compositional measures versus total DOM concentration. Measures include A)
SUVA, specific ultraviolet absorbance at 255 nm, B) $S_{275-295}$, spectral slope between 275 to
295nm, C) DOC:DON, and D) HSF; humic substances fraction. Colours represent geographical
sampling locations (HZ: Lake Hazen Watershed, NU; DL: Daring Lake, NT; WK: Wekweètì,
NT; YK: Yellowknife, NT; MR: Mackenzie River, NT; ELA: IISD-Experimental Lakes Area,
ON; TLW: Turkey Lakes Watershed, ON; NW: Nottawasaga River Watershed, ON; GR: Grand
River, ON; LP: Long Point, ON; BBK: Black Brook Watershed, ON) while shapes represent
hydrologic environments. Light grey circles represent two other published DOM characterization
studies conducted at similar scales [16,44].
669

670
Fig 3. Principal component analyses for samples from different ecozones. Grey dots
represent individual samples, while vectors represent absorbance (red), elemental (green), and
LC-OCD (purple) compositional measures.

Fig 4. Composition wheels for DOM from different hydrologic settings within each geographical site. Axes are a numerical value normalized relative to the maximum and minimum encountered within the dataset for each parameter. Parameters for each axis found in bottom left. The orientation of the composition wheel (CW) axes are arbitrary. Different samples from the same hydrological and geographic setting are plotted within the same CW. Two 'end member' compositions are highlighted in sites with many samples to help visualize the continuum of DOM mixtures: groundwater-like DOM (low DOC:DON, high SUVA and HSF) or photolyzed DOM (high $S_{275\text{-}295}$).

Fig 5. Changes to a Composition Wheel resulting from photolytic and microbial degradation (left) and differences in terrestrial versus aquatic DOM sources (right).
Changes to dissolved organic matter (DOM) (grey) with photolytic (orange; 12 to 18 day exposure to sunlight) or microbial (purple; 30-day experiment) degradation based on experimental incubations of natural DOM samples [51] (left). The mean decrease in DOC concentration was 18% during photolysis and 10% during microbial degradation experiments (n=11, 9; respectively). Hypothetical differences in DOM sources (right) are based on various leachate experiments in literature [22,37,55–59].

696
Fig 6. Conceptual model of DOM evolution along the aquatic continuum using
Composition Wheels. A conceptual model of a few select processes and sources of dissolved
organic matter (DOM) along the aquatic continuum. Two main hydrologic flowpaths are
illustrated (A-B-C and A-b-c) and originate from the same source 'A'. However, each flowpath
undergoes different processing within the environment. Arrows represent the transport or
addition of DOM among groundwaters (dark purple) and surface waters (light purple), while
lowercase text highlights some of the specific processes and sources encountered. The wetland
'b' represents an area of organic matter degradation and DOM accumulation within organic-rich
groundwaters, resulting in the transport of large, aromatic DOM components into surface water
'c'. Conversely, DOM is rapidly transported down the river 'B' and into a large lake 'C',
resulting in much higher exposure to sunlight. The use of Composition Wheels to illustrate
differences in sources and processing is shown on the bottom panel, comparing DOM from an
organic-rich source with longer exposure to sunlight (A-B-C) versus the same DOM with less
sunlight exposure and greater microbial degradation (A-b-c).
711
712

714 **Fig 7. Surrogate parameters for the DOM Composition Wheel.**

Supporting information

S1 Fig. Contribution of each variable within the PCA.

719 **S1 Table. Environmental description for all sampling sites.** Included are the number of total
720 samples taken for DOM concentration with at least one compositional measure (n_{DOC}), and
721 number of samples used from that site in the PCA analysis (n_{PCA}).
722

SURFACE WATERS

Name	Location	n_{DOC}	n_{PCA}	Description
IISD-Experimental Lakes Area, ON (ELA)	49° 39' 40"N, 93° 43' 48"W Ontario, Canada	**74**	**45**	Boreal forest, underlain by Precambrian bedrock with discontinuous surficial layer of sandy-gravel till. Sampled from 2010 to 2012. Ecozone mean annual temperature: 0.7 C; Ecozone total annual precipitation: 825 mm.
Lakes		*49*	*36*	
Creeks		*25*	*9*	
Grand River, ON (GR)	43° 30' 41"N, 80° 29' 43"W Ontario, Canada	**39**	**-**	Surrounding land predominately agricultural and flows past six wastewater treatment plants. Sampled from 6 consecutive locations along a 90km stretch every two months from 2011 to 2012. Ecozone mean annual temperature: 6.5 C; Ecozone total annual precipitation: 940 mm.
Yellowknife, NT (YK)		**23**	**20**	Samples from the Taiga Shield underlain by discontinuous permafrost. Surface waters are surrounded by bedrock and peat plateaux around Yellowknife. Sampled in July or October between 2013 and 2017. Ecozone mean annual temperature: -4.8 C; Ecozone total annual precipitation: 565 mm.
Lakes	62° 27' 14"N, 114° 22' 18"W Northwest Territories, Canada	*2*	*2*	
Ponds		*8*	*7*	
Rivers		*10*	*9*	
Creek		*3*	*2*	
Mackenzie River, NT (MK)	63° 14' 17"N, 123° 34' 0"W Northwest Territories, Canada	**13**	**-**	Samples taken by the Community Based Monitoring network along the Mackenzie River in July and August of 2015. Samples ranged from WHERE to Inuvik. River flows through Taiga Shield and Taiga Plains.
Rivers		*13*	*-*	
Wekweètì, NT (WK)	64° 11' 24"N, 114° 11' 10"W Northwest Territories, Canada	**11**	**11**	Situated in the Taiga Shield, below treeline, continuous permafrost. Samples taken in October of 2015 and 2016. Ecozone mean annual temperature: -4.8 C; Ecozone total annual precipitation: 565 mm.
Lakes		*9*	*9*	
Creeks		*2*	*2*	
Daring Lake, NT (DL)	64° 31' 29"N, 111° 40' 24"W Northwest Territories, Canada	**19**	**19**	Found in the Southern Arctic above treeline, continuous permafrost. Ecozone mean annual temperature: -9.8 C; Ecozone total annual precipitation: 313 mm.
Lakes		*7*	*7*	
Ponds		*1*	*1*	
Creeks		*11*	*11*	
Lake Hazen, NU (LH)		**160**	**12**	
Lakes	81° 49' 30"N, 71° 19' 26"W Nunavut, Canada	*38*	*2*	
Ponds		*18*	*4*	Tundra located in the high arctic; Lake Hazen Watershed is considered a local polar oasis.
Creeks		*41*	*1*	
Rivers		*32*	*0*	
Seeps		*31*	*5*	

GROUND WATERS

Name	Location	n	nPCA	Description
Turkey Lakes Watershed, ON (TLW)	47° 2' 54"N, 84° 24' 25"W Ontario, Canada	**16**	**-**	Relatively un-impacted watershed in the Great Lakes-St. Lawrence forest region. Area consists of Precambrian bedrock and surficial glacial deposits of glaciofluvial outwash. Samples collected from depths ranging between 0.90 -

				6.89m below surface. Ecozone mean annual temperature: 0.7 C; Ecozone total annual precipitation: 825 mm.
IISD-Experimental Lakes Area, ON (ELA)	49° 39' 40"N, 93° 43' 48"W Ontario, Canada	**17**	**-**	Piezometers constructed in transect along a wetland, ranging from 0.70 - 3.85m below surface. Ecozone mean annual temperature: 0.7 C; Ecozone total annual precipitation: 825 mm.
Nottawasaga Aquifer, ON (NW)	44° 7' 26"N, 79° 49' 12"W Ontario, Canada	**6**	**-**	Surficial deposits of glaciolacustrine deposits in an agriculturally-impacted aquifer. Samples collected from single multi-level piezometer within an unconfined surficial sand aquifer at depths of 4.35m, 5.13m, 6.68m, 9.90m, and 11.3m below surface. Ecozone mean annual temperature: 6.5 C; Ecozone total annual precipitation: 940 mm.
Black Brook Watershed, NB (BBK)	47° 6' 11"N, 67° 45' 40"W New Brunswick, Canada	**15**	**-**	Site is an agriculturally-impacted aquifer, sampled during the summer of 2012. Surficial deposits of till and small deposits of glacial outwash. Samples taken from twelve domestic wells and three multi-level piezometers (6.1 - 30m below surface). Ecozone mean annual temperature: 4.6 C; Ecozone total annual precipitation: 1185 mm.
Long Point, ON (LP)	42° 34' 46"N, 80° 22' 57"W Ontario, Canada	**23**	**-**	Unconfined sand aquifer atop of a clay aquitard. Piezometers range from 1 to 4m below surface. Sampling of groundwater containing a septic plume.
Yellowknife, NT (YK)	62° 27' 14"N, 114° 22' 18"W Northwest Territories, Canada	**33**	**16**	Samples from the Taiga Shield underlain by discontinuous permafrost. Surface waters are surrounded by bedrock and peat plateaux around Yellowknife. Sampled in July or October between 2013 and 2017. Ecozone total annual precipitation: 565 mm.
Wekweètì, NT (WK)	64° 11' 24"N, 114° 11' 10"W Northwest Territories, Canada	**1**	**1**	Situated in the Taiga Shield, below treeline, continuous permafrost. Samples taken in October of 2015 and 2016. Ecozone total annual precipitation: 565 mm.
Daring Lake, NT (DL)	64° 31' 29"N, 111° 40' 24"W Northwest Territories, Canada	**4**	**4**	Found in the Southern Arctic above treeline, continuous permafrost. Ecozone mean annual temperature: -9.8 C; Ecozone total annual precipitation: 313 mm.
Lake Hazen, NU (LH)	81° 49' 30"N, 71° 19' 26"W Nunavut, Canada	**17**	**2**	Samples taken from piezometer installed at deepest extent of active-layer (~0.25m). Location was in a subcatchment wetland containing organic-rich soil. Flow direction was through wetland into nearby lake. Organic-rich layer underlain by silt or clay material.

S2 Table. DOM Composition versus Concentration Statistics. Results from a linear model of dissolved organic matter (DOM) composition (SUVA, slope between 275-295nm, DOC:DON,

726   and humic substances fraction) as predicted by the overall DOM concentration (mg C/L; log
727   transformed).
728

Parameter	Linear Regression to log(DOM)	
	R^2	p-value
SUVA	0.10	<0.01
S275	0.01	0.61
DOC:DON	0.33	<0.01
HSF	0.09	<0.01