TITLE

Visualizing subcellular structures in neurons with expansion microscopy

AUTHORS

Logan A. Campbell[1]*, Katy E. Pannoni[1]*, Niesha A. Savory[2], Dinesh Lal[3], Shannon Farris[1-4#]

AFFILIATIONS

[1]Center for Neurobiology Research, Fralin Biomedical Research Institute, Virginia Tech
Carilion, Roanoke, VA, 24016, USA.
[2]School of Neuroscience, Virginia Tech, Blacksburg, VA, 24016, USA.
[3]Virginia Tech Carilion School of Medicine, Roanoke, VA, 24016, USA.
[4]Department of Biomedical Sciences & Pathobiology, Virginia-Maryland College of Veterinary
Medicine, Virginia Tech, Blacksburg, VA, 24061, USA.

*Authors contributed equally
#Lead Contact Farrissl@vtc.vt.edu

Abstract Word Count 174
Introduction, Results, Discussion Word Count 3,573
Figures 6
Tables 3

ABSTRACT

Protein expansion microscopy (proExM) is a powerful technique that crosslinks proteins to a
swellable hydrogel to physically expand and optically clear biological samples. The resulting
increased resolution (~70 nm) and physical separation of labeled proteins make it an attractive
tool for studying the localization of subcellular organelles in densely packed tissues, such as the
brain. However, the digestion and expansion process greatly reduces fluorescence signals
making it necessary to optimize ExM conditions per sample for specific end goals. Here we
describe a proExM workflow optimized for resolving subcellular organelles (mitochondria and
the Golgi apparatus) and reporter-labeled spines in fixed mouse brain tissue. By directly
comparing proExM staining and digestion protocols, we found that immunostaining before
proExM and using a Proteinase K based digestion for 8 hours consistently resulted in the best
fluorescence signal to resolve subcellular organelles while maintaining sufficient reporter
labeling to visualize spines and trace individual neurons. With these methods, we more
accurately quantified mitochondria size and number and better visualized Golgi ultrastructure in
reconstructed CA2 neurons of the hippocampus.

Figure 1: ExM workflow for visualizing subcellular organelles in fixed brain tissue. Schematic of immunostaining and proExM workflows tested to optimize visualization of organelles in genetically-labeled neurons.

INTRODUCTION

Protein retention expansion microscopy (proExM) is a powerful tool that crosslinks proteins to a

swellable hydrogel to optically clear and physically expand tissues up to ~4-fold in volume[1,2].

Because expansion physically separates crosslinked moieties, this technology is particularly

useful for visualizing subcellular structures in densely packed tissues, such as the brain.

However, one consequence of tissue expansion is a decrease in the fluorescence intensity of

labeled proteins primarily due to the digestion process and the dilution of fluorescence signal per

unit volume. Various ExM protocols have described different methods for improving

fluorescence retention, primarily by modifying fixation, crosslinking, and/or digestion conditions

to preserve protein epitopes[2–8]. One common ExM protocol uses a strong protease-based

digestion (Proteinase K[2]), but other gentler proteases have also been used (LysC[2,3]), as well as a

combination of heat and detergents in place of proteases (e.g. autoclave in an alkaline buffer[2–5]).

Immunostaining can also be done before or after ExM to boost fluorescence[2,3] (Fig. 1),

although results are often dependent on the protein epitope and the quality of antibody staining.

To improve antibody staining in brain tissue, antigen retrieval is often performed prior to

immunostaining via boiling tissue in water or heating tissue in a citrate buffer (pH 6) to expose

protein epitopes and reduce nonspecific staining. Here we set out to compare ExM

immunostaining and digestion conditions for fluorescently labeled subcellular organelles in

perfused brain sections using antibodies that either require or do not require antigen retrieval

(COX4-labeling of mitochondria and GOLGA5-labeling of Golgi, respectively). Visualizing the

spatial organization of organelles within compartmentalized cells, such as neurons, is important

for understanding their function, thus we paid particular attention to conditions that maintained

fluorescence signal from genetically-encoded reporters (i.e. enhanced green fluorescent protein,

EGFP, and tdTomato). We found that performing IHC, with or without antigen retrieval, prior to

ExM with Proteinase K digestion for 8 hours best preserves fluorescence signal for co-
visualizing subcellular organelles and neuronal morphology, including spines. Further, we report
our optimized conditions for antibodies against widely used reporters and conclude with protocol
considerations for achieving specific end goals.
69
RESULTS
Proteinase K digestion time impacts fluorescence intensity and expansion factor
Sufficient digestion or homogenization (also referred to as hydrolysis in some protocols) is
required to prevent sample distortion during expansion and is highly dependent on digestion
conditions, including time, temperature, pH, buffer composition and enzyme quality[1,3,9]. Varying
digestion time impacts fluorescence intensity[1] and how much the tissue expands in water, or the
expansion factor[5]. To determine the effect of digestion time on the fluorescence intensity of the
Amigo2-EGFP reporter line, which predominantly labels hippocampal area CA2 neurons, we
performed a time course of enzymatic digestion with Proteinase K as described in the proExM
protocol[3]. Initially, we performed the time course on 40-micron vibratome cut sections from
perfused adult Amigo2-EGFP mouse brains. Hydrogels were digested for 2, 4, 8 or 16 hours at
room temperature. Unfortunately, there was insufficient EGFP signal remaining at the 8 and 16
hour timepoints to directly compare hydrogels across conditions (data not shown). We then
repeated the experiment on sections immunostained for GFP prior to ExM and imaged the
resulting gels with identical acquisition parameters to directly compare fluorescence intensities
and expansion factors (Fig. 2). Importantly, we measured micro expansion factors, or the degree
to which cell soma areas expanded versus the commonly reported macro expansion factors
calculated by measuring how much the hydrogel expands. We found that average fluorescence
intensities diminished as the length of digestion time increased (one-way ANOVA; F-stat: 20.96,
p-value: 8.17E-14. N = two animals, 1-2 sections per animal per time point; 159 total cells, 32 $\pm$
4 cells per time point, Fig. 2AB). Further, we found that average expansion factors increased as
length of digestion time increased (one-way ANOVA; F-stat: 21.07, p-value: 4.26E-11, Fig.
2CD), resulting in an inverse relationship between fluorescence intensity and expansion factor
(Fig. 2E). Interestingly, we did not detect significant decreases in fluorescence intensities from
pairwise comparisons between the 8-hr digestion and the 2- or 4-hr digestions (p=0.90 and 0.55,
respectively, Tukey's post hoc test, Fig. 2D), despite significant increases in expansion factors
between the same comparisons (p=0.001 for each). These data indicate that 8-hr digestion retains
the most fluorescence for a sizable expansion factor (~3) that is not significantly different from
the overnight expansion factor (p=0.18). However, we note that the fluorescence intensities
reported here are corrected for background, and shorter digestion times have greater fluorescence
background signal compared with longer digestion times (see Table 1). Regardless of digestion
time, we were able to successfully acquire robust fluorescent images at each time point,
including overnight (Fig. 2F), as long as immunolabeling was performed prior to ExM.

Figure 2: The effect of Proteinase K digestion time on fluorescence intensity and expansion factor.
A) Representative single z-section images of unexpanded (i) and expanded GFP+ CA2 neurons after 2-hour (ii), 4-hour (iii), 8-hour (iv), or overnight (v) digestion with Proteinase K. Image acquisition parameters were identical for each condition, except for the unexpanded condition.
B) GFP fluorescence intensity measured in the cell soma after the different digestion times. Fluorescence was corrected for background and $\log_2$ transformed. Line represents the median. Digestion time had a significant effect on GFP fluorescence (one-way ANOVA; F-stat: 20.96, p-value: 8.17E-14, 32 ± 4 cells per time point from two animals, 159 total cells).
C) Expansion factor of the expanded cell somas in B. Expansion factors were calculated using cell soma areas relative to the average unexpanded cell soma area. Digestion time had a significant effect on expansion factor (one-way ANOVA; F-stat: 21.07, p-value: 4.26E-11).
D) Matrix of p-values visualizing the results of pairwise Tukey's post hoc tests comparing mean fluorescence (upper diagonal) or expansion factor (lower diagonal) at each digestion time point. Matrix is color coded by p-value (red = significant; grey = not significant; α=0.05).
E) Effect of digestion time on fluorescence intensity (cyan; left axis) and expansion factor (orange; right axis). Plot shows the mean and 95% confidence intervals from **B** and **C**.
F) Unexpanded (i) and overnight expanded (ii) max-projection 10X confocal images of GFP+ CA2 cells. Images are the same as in **A**(i) and (v). Imaging parameters were optimized separately to obtain the best image of both. Scale bars = (**A**, **F**) 50 µm.

Cells expand equivalently, independent of section depth

Next we considered if the time-dependent effect of Proteinase K on expansion factor and

fluorescence intensity is impacted by tissue depth. We reasoned that cells near the surface may

have greater access to Proteinase K and/or fluorescently-labeled antibodies compared with cells

deeper within tissue that may impact expansion factor and/or fluorescence intensity, respectively.

To test this, we compared cell soma areas binned by Z-section depth (adjusted by hydrogel

thickness) across digestion time points.

Figure 3: No appreciable effect of tissue depth on expansion factor.
A) Average cell soma area (in μm^2) by Z section depth for unexpanded control and digestion conditions. Cell soma area was measured in the Z section with the widest soma diameter. Because of the overall increase in depth with expansion, cells were binned by Z section to achieve ~6 bins per time point. N = 32 ± 4 cells per time point.
B) Background subtracted average cell soma fluorescence by Z section depth. Binning of the data is the same as in A. The data are not $\log_2$ transformed as in Fig 2.
C) Representative confocal images of GFP+ CA2 cell somas after 2-hour (left panel) or overnight (right panel) digestion at three Z-section depths. Cell soma size and fluorescence is variable in the Z dimension in unexpanded conditions, which does not appreciably change under difference digestion conditions. Error bars show the standard deviation. Scale bar is 50μm.

We did not detect a systematic difference in cell soma area across z-section depth (Fig. 3),
indicating that Proteinase K equivalently digests 40-micron thick tissue, at least by the 2-hr time
point. For some time points (e.g. 4- and 8-hrs) fluorescence intensity appeared brighter near the
surface compared with deeper sections (Fig. 3B). However, we note this is likely due to optical
limits, as fluorescence intensity was not brighter at the far-end surface that has equivalent access
to Proteinase K and antibody solutions. We saw similar results with 100-micron thick tissue
(data not shown).
120

Immunolabeled organelles are best resolved with IHC prior to ExM
Protease digestion can decrease fluorescence intensity by impacting fluorescently-labeled
antibodies and/or target antigen availability. Thus, immunohistochemistry (IHC) labeling before
or after ExM (further referred to as IHC-pre and IHC-post, respectively) can be affected, but it is
unknown if they equally affect the fluorescence intensity of antibodies that require antigen
retrieval. Protease-free digestion protocols (e.g. detergent plus heat created in an autoclave liquid
cycle) have been shown to effectively digest hydrogels, and avoids protease-dependent depletion
of fluorescence intensity[3]. However, these protocols have not been tested with antibodies that
require antigen retrieval. In Figure 4, we directly compared COX4-mitochondria labeling after
protease (Proteinase K) or autoclave (i.e. mild digestion) digestion ExM protocols[3]. We further
compared these digestion protocols with IHC performed pre (Fig. 4AB) or post ExM (Fig. 4CD).
Our optimized protocol for COX4 immunostaining on unexpanded sections (Fig. 4E) requires
antigen retrieval (boiling) prior to IHC, thus an antigen retrieval step was included for sections
bound for Proteinase K digestion.
135

We detected COX4-labeled mitochondria using both Proteinase K and autoclave digestion
ExM protocols (Fig. 4), however Proteinase K digested gels performed better with IHC-pre, and
autoclave digested gels performed better with IHC-post. Compared with unexpanded COX4-
labeled mitochondria, expanded COX4-labeled mitochondria were on average slightly greater in
number per cell (ExM: 117.5 ± 5.6, No ExM: 95.8 ± 4.4) and smaller in size (mitochondria size:
ExM 0.001 ± 0.0, No ExM 0.003 ± 0.0) after normalizing to cell soma area (total mitochondria
area: ExM 422.5 ± 93.7, No ExM 123.8 ± 2.4) (Fig. 4G). To account for potential anisotropic
expansion of cell somas, we compared the ratio of nuclear area (measured via DAPI) to the cell
soma area (measured via reporter labeling) and found them to be similar on average with and
without ExM (ratio: ExM 0.357 ± 0.00, No ExM 0.347 ± 0.01; nuclear area: ExM $1,383.1 \pm$
93.5, No ExM 154.5 ± 7.8; soma area: ExM $3,869.5 \pm 259.2$, No ExM 443.3 ± 11.8), indicating
that the decrease in percent cytoplasmic area of mitochondria (Fig. 4I) is due to better
individually resolved expanded mitochondria.
149

In regards to reporter labeling, RFP fluorescence only fared well when IHC was done prior to
ExM, regardless of digestion method. Thus, for epitopes that require antigen retrieval and/or
reporter labeling, IHC-pre-ExM is the preferred method of choice. Note that antigen retrieval

diminishes reporter labeling (RFP and GFP) and explains the difference in RFP intensity
between autoclave (Fig. 4Aii) and Proteinase K (Fig. 4Bii) digested gels. If only mitochondria
labeling is required, the autoclave digestion protocol performs well with IHC-post. The inferior
staining of COX4 with IHC-pre with autoclave digestion compared with IHC-pre with Proteinase
K digestion may be due to a lack of antigen retrieval in the former that is achieved by autoclave
digestion.

Figure 4: ProExM pipeline comparison for visualizing mitochondria.
A) 20X Max-projection image of a section immunostained for COX4 (i) and RFP (ii) pre-ExM with autoclave digestion. Images in panels A-D were acquired with identical imaging parameters to directly compare conditions. Brightness and contrast were adjusted to the same extent across the entire image.
B) 20X Max-projection image of a boiled section immunostained for COX4 (i) and RFP (ii) pre-ExM with overnight Proteinase K digestion.
C) 20X Max-projection image of a section immunostained for COX4 (i) and RFP (ii) post-ExM with autoclave digestion.
D) 20X Max-projection image of a boiled section immunostained for COX4 (i) and RFP (ii) post-ExM with overnight Proteinase K digestion
E) 40X Max-projection image from a subset of Z sections from a boiled, unexpanded section immunostained for COX4 and RFP. Inset is a single Z section from the indicated representative cell (white asterisk) with mitochondria outlined as quantified in **G-I**. Imaging parameters were optimized per condition in **E-F**.
F) 20X Max-projection image from a subset of Z sections from a boiled section immunostained for COX4 and GFP pre-ExM with overnight Proteinase K digestion as in **B**.
G) The average number of mitochondria per cell in unexpanded (**E**) and expanded (**F**) sections.
H) Mitochondria area per cell (normalized by soma area). Line reflects median and whiskers are twice the inner quartile range.
I) The average percent cytoplasm area (soma area - nuclear area) containing mitochondria. Note that ExM better resolves densely packed mitochondria, resulting in greater number, smaller sized, and fewer percent area per cell soma compared with non-ExM.
Scale bars: 50 µm (**A,E,F**) and 5 µm (insets).

160
We next compared the Proteinase K and autoclave ExM protocols with GOLGA5-
immunolabeling of Golgi apparatus (Fig. 5), which does not require antigen retrieval. Consistent
with our COX4 results, GOLGA5-labeled Golgi were detected with either Proteinase K or
autoclave digested protocols. Golgi in expanded sections were well resolved and revealed
complex Golgi structure compared with Golgi in unexpanded sections (Fig. 5F). In contrast to
our COX4 results, GOLGA5-labeling fared well in both IHC-pre-ExM digestion conditions,
likely due to robust GOLGA5-labeling in unboiled sections. Similar to COX4, GOLGA5-
labeling post-ExM with Proteinase K digestion was unsuccessful. IHC-pre with autoclave

digestion was not tested. As with RFP reporter labeling, GFP reporter labeling also fared well
with IHC-pre and autoclave digestion, albeit at lower intensities than Proteinase K digestion.
Thus, GFP and RFP fluorescence retention are comparable when antibody-labeled prior to ExM
and they retain more fluorescence with Proteinase K digestion compared with autoclave
digestion. It is important to note that we detect qualitative differences in the ability of different
reporter antibodies to detect spines, with and without ExM (see Table 3).
175

Figure 5: ProExM pipeline comparison for visualizing Golgi.
A) 20X Max-projection image of a section immunostained for GOLGA5 (i) and GFP (iii) pre-ExM with autoclave digestion. (ii) Insets are zoomed images of the dotted box in (i). Images in panels **A**-**C** were acquired with identical imaging parameters to directly compare conditions. Brightness and contrast were adjusted to the same extent across the entire image.
B) 20X Max-projection image of a section immunostained for GOLGA5 (i) and GFP (iii) pre-ExM with overnight Proteinase K digestion.
C) 20X Max-projection image of a section immunostained for GOLGA5 (i) and GFP (iii) post-ExM with overnight Proteinase K digestion.
D) 40X Max-projection of a subset of Z sections from an unexpanded section immunostained for GOLGA5 (i) and RFP (iii). Imaging parameters were optimized per condition in **D**-**E**.
E) 20X Max-projection of a subset of Z sections from image in (**B**) immunostained pre-ExM with overnight Proteinase K digestion. Note the increase in Golgi complexity that is resolved with autoclave or Proteinase K digested gels compared with unexpanded sections, although fluorescence intensity was better retained with Proteinase K digestion.
F) Line plot profile showing the fluorescence intensity of unexpanded Golgi (cyan) and Golgi after IHC-pre-ExM with Proteinase K digestion (magenta). Measured Golgi are insets in **D**iii and **E**.
Scale bars: 50μm (**A**i, **D**i, **E**) and 10μm (**A**ii, **D**ii).

176
177

Dendritic spines are best resolved in expanded GFP or RFP-immunolabeled tissue
In addition to resolving subcellular organelles such as mitochondria and Golgi apparatus, the
ability to resolve fine dendritic structures such as spines allows us to address questions about
function and plasticity at synapses. Thus, we set out to find the optimal combination of ExM
conditions for subcellular organelles and reporter antibodies to resolve dendritic spines in our
EGFP and tdTomato reporter mice. Table 3 compares spines in unexpanded and expanded tissue,
with or without immunolabeling for the reporter protein, and with or without antigen retrieval by
boiling or citrate. Expanded samples were immunostained pre-ExM (if at all) and digested
overnight with Proteinase K, as described in the methods. To boost the signal of the reporter

protein, we tested two antibodies against each EGFP and tdTomato reporters (anti-GFP and anti-

RFP, respectively). The ability to resolve dendritic spines in each condition was qualitatively

assessed by multiple investigators (not blinded to condition), and indicated by the number of + in

the table (from + to ++++). A greater number of + indicates better discrimination of spines and

"-" indicates dendritic spines could not be resolved for a given condition. Some conditions in the

table have yet to be tested as indicated where applicable.

Figure 6: Resolving dendritic spines with ExM.
A) 20X with 3X zoom max-projection image (i) of a section immunostained for RFP pre-ExM with overnight Proteinase K digestion. This mouse received 2 tamoxifen injections. (ii) Insets are 2X zoomed images of the dotted box in (i). All images were acquired with parameters optimal per condition. Brightness and contrast were adjusted per condition across the entire image.
B) 40X with 2X zoom max-projection image of an unexpanded tdTomato section. This mouse received 2 tamoxifen injections.
C) 20X with 2X zoom max-projection image of a section immunostained for GFP pre-ExM with overnight Proteinase K digestion.
D) 40X with 2X zoom max-projection image from an unexpanded section immunostained for GFP. Note the increased background labeling and density of dendritic branches in the unexpanded images, making dendrites more difficult to trace compared with expanded images.
Scale bars: 25μm (**Ai**, **Bi**, **Ci**, **Di**) and 5μm (**Aii**, **Bii**, **Cii**, **Dii**).

Figure 6 shows representative images of dendritic spines for each of the four conditions tested. In

general, spines were easier to resolve in tdTomato reporter mice (top panel) than EGFP reporter

mice (bottom panel), even with the same ExM and IHC conditions. However, both reporter lines

benefited from the addition of IHC to boost the reporter signal. Without an IHC boost, neither

reporter line had resolvable spines after ExM (see Table 3), primarily due to a loss of

fluorescence. We saw the best resolution of spines in expanded tdTomato+ tissue that was
stained with the rabbit anti-RFP without any antigen retrieval (Fig. 6A; "Am2-
icre;tdTomato/RFP-rabbit" in Table 3). We noted that the chicken RFP antibody did not label
spines as well as the rabbit RFP antibody ("Am2-icre;tdTomato/RFP-chicken", Table 3),
indicating that not all reporter antibodies are equal when it comes to labeling spines. The IHC-
pre-ExM condition increased the resolution of tdTomato+ dendritic spines compared with
unexpanded and unlabeled tdTomato+ dendritic spines (Fig. 6B). While spines were still visible
on tdTomato+ CA2 dendrites without immunostaining or expansion (Fig. 6Bii), we were not able
to reliably resolve spines on unexpanded or expanded EGFP+ dendrites without immunostaining
("am2-EGFP", Table 3). With the addition of IHC, however, spines could be resolved on GFP+
dendrites in both expanded (Fig. 6C) and unexpanded tissue (Fig. 6D). Compared to unexpanded
tissue with the same IHC conditions, the ExM protocol increases the resolution of spines by
increasing their physical size and separation from nearby dendritic branches and reducing
background fluorescence (compare Fig. 6C and D) enabling spine morphometric analyses on
individually traced neurons.
216
DISCUSSION
ProExM is a powerful tool that increases the resolution of conventional fluorescence microscopy
to ~70 nm and can be performed with tools available in a standard molecular biology
laboratory[10,11]. Because a fully expanded hydrogel is mostly water, the optically clear sample is
well suited to resolve densely packed organelles and tissues. However, the digestion and
expansion process greatly reduces fluorescence retention making it necessary to optimize ExM
conditions per sample for specific end goals. Here we described a proExM workflow optimized
for resolving subcellular organelles (mitochondria and the Golgi apparatus) and spines in fixed
mouse brain tissue. We reliably found that immunostaining before proExM (IHC-pre-ExM) and
using a Proteinase K based digestion for 8 hours resulted in the best fluorescence signal to
resolve subcellular organelles while maintaining sufficient reporter labeling to visualize spines
and trace individual neurons. With these methods, we were able to more accurately quantify
mitochondria size and number and better visualize Golgi ultrastructure in reconstructed CA2
neurons of the hippocampus.
231
Several groups have optimized expansion protocols to visualize subcellular organelles across
different sample types, including various cell lines[1,5,8,12–15], rat liver[16], clinical specimens[17],
fungi[6,18], songbird[19] and drosophila[20,21]. Others have used ExM to visualize subcellular
structures, including mitochondria[1,2,22–24] and/or spines[2,19,22] in brain tissue, but few have
systematically analyzed how fluorescence intensities and expansion factors compare across
protocols or with unexpanded measurements. This is critically important as several groups have
noted discrepancies in micro versus macro expansion factors in other sample types[7,13,16,25–27],
including dissimilarities in expansion factors of different subcellular organelles[13] or of
subcellular organelles across neighboring cells and tissues[16]. Others, however, have reported
minimal to no differences in micro vs macro expansion factors[6,18]. We did not measure macro

expansion factors here, but our average micro expansion factors are lower than the commonly
reported 4-4.5X macro expansion factor for proExM, which is consistent with other reports using
micro expansion factor measurements[13,14] (but see also ref[25]), reinforcing the notion that each
sample type needs to be independently optimized and validated for ExM.

To determine the optimal digestion time for fluorescence retention in fixed brain sections, we
performed a digestion time course and found the greatest drop in fluorescence after the overnight
digestion. There was no significant drop in fluorescence between 2, 4 or 8 hours of digestion.
However, there was a significant increase in expansion factor during this time period. Expansion
factor begins to plateau after 8 hours of digestion, and while there is a slight increase in
expansion factor in the overnight condition it is not statistically significant. Therefore, under the
conditions used here, a digestion time of 8 hours is ideal for achieving a robust expansion (~3X)
without further loss of fluorescence seen with overnight digestion. At all digestion time points,
the expansion factor did not systematically vary by depth, indicating that 2 hours in Proteinase K
is sufficient for uniform penetration and isotropic expansion in the Z dimension of 40 micron
brain sections, as previously reported for thicker sections and longer digestion times[10].

In regards to labeling subcellular organelles in fixed brain sections, we were able to visualize
expanded mitochondria with a COX4 antibody and expanded Golgi apparatus with a GOLGA5
antibody, using either the proteinase K digestion or the mild autoclave digestion. In our hands,
the IHC-pre ExM with proteinase K digestion outperformed the other conditions based on
fluorescence signal retention for both immunostained organelles and reporter proteins. While
EGFP and tdTomato have been reported to have different percent fluorescence retention after
ExM[2], they perform comparably when antibody-labeled prior to ExM, as recently reported in
cultured cells[28]. They also retain more fluorescence with Proteinase K digestion compared with
autoclave digestion. However, if the goal is only to visualize mitochondria, the mild autoclave
digestion with IHC-post ExM also produced good COX4 staining, as seen for other mitochondria
immunostains, like TOMM20[2]. The GOLGA5 antibody performed decently using IHC-pre ExM
and autoclave digestion, which the COX4 antibody did not, likely due to COX4 immunolabeling
requiring antigen retrieval. Neither antibody performed well with IHC-post ExM and Proteinase
K digestion.

Despite diminished fluorescence, ExM afforded better resolution for quantification of
subcellular organelles compared to unexpanded organelles. We quantified the number and size of
expanded mitochondria and found that the expanded mitochondria were smaller and more
numerous than unexpanded mitochondria. This presumably is indicative of tightly packed
mitochondria in the unexpanded samples being lumped together that can be separately resolved
with expansion. Here, we normalized subcellular measurements (i.e. mitochondria size) to within
cell nuclear and cytoplasmic areas. Another study found discrepancies in cell soma vs nuclear
expansion factors[16], but here the ratio of nuclear area to cytoplasmic area remained constant

between unexpanded and expanded states, indicating isotropic expansion, perhaps due to our
longer digestion times.

The fine details of the Golgi cisternae were also better resolved after expansion, whereas
without expansion the Golgi were smoothened and much of the details lost. For the goal of
visualizing reporter-labeled dendritic spines, we found that the addition of IHC-pre ExM was
necessary for the resolution of both EGFP+ and tdTomato+ spines. Dendritic spines were best
resolved in IHC-pre ExM with Proteinase K digestion with either GFP or RFP antibodies. The
ability to label dendritic spines in ExM was antibody dependent, with some antibodies against
the same reporter faring better than others under the same IHC conditions.

Our lab has begun applying the described proExM methods to answer open questions
involving subcellular organelles in neurons. One such open question is whether there is Golgi
present in dendrites, or if the Golgi is limited to neuronal cell bodies. It is known that local
translation of RNA occurs in dendrites[29]; however, there is conflicting evidence of the existence
of Golgi apparatus in the dendrites, which would normally process newly transcribed membrane
bound proteins. Using combined GOLGA5-labeling of Golgi apparatus and reporter neuron
labeling with the proExM protocol described here, we do not detect GOLGA5 staining outside
the cell soma or very proximal dendrites, consistent with reports that canonical Golgi markers
are retained in the soma and not present in distal dendrites[24,31,32].
297
298 *Considerations for subcellular imaging of expanded samples*
299 The benefits of expansion come at several costs, including diminished signal concentration,
300 hydrogel mechanical integrity and movement, and increased imaging volume and time[5]. Here we
301 comment on the proposed workarounds[2,3,11] for these issues that have or have not worked well
302 for subcellular imaging of expanded brain sections.
303
Expansion microscopy substantially increases the thickness of a sample, which limits its ability
to be imaged with standard high magnification microscope objectives with limited working
distances. It can be expected that the gel thickness will be equivalent to 4-fold the depth of the
gelation chamber. Further, keeping the tissue in plane during gel chamber assembly is difficult,
often resulting in increased sample z-distance. Minimizing tissue thickness (40 microns versus
100 microns) and using a single coverslip for the gelation chamber helped minimize gel
thickness without sacrificing ability to reconstruct neurons.
311
Loss of fluorescence due to the digestion of antibodies or epitopes and dilution of fluorescent
molecules per unit volume can result in low contrast samples not suitable for high resolution
imaging even with overexpressed fluorescent reporter proteins. We found that performing IHC
beforehand and limiting the Proteinase K digestion to 8 hours largely negated this issue. When
the fluorescence signal is insufficient, imaging in low concentrations of PBS (0.5X PBS instead
of 0.0001X PBS or water) substantially improved the contrast by increasing the concentration of

fluorescent molecules per area. This will decrease the expansion factor, but in our hands, even a
2-3-fold expansion in optically clear tissue produces better resolved images of subcellular
structures than unexpanded tissue.
321
Hydrogel movement during imaging is a common issue for us. Because gels expand 4-fold in x,
y and z compared with unexpanded brain sections, tile scans are required to reconstruct entire
neurons even at 10x (see Fig. 2Fii for a 10x single field of view of expanded neurons). Tile
scanning increases the length of time required to image a gel and thus worsens gel shift. Gel shift
was most noticeable while imaging on an upright microscope equipped with a water immersion
lens, since the gel can easily shift when submerged. Gel shift was less noticeable when imaging
either on an inverted scope or on an upright scope with air objectives (if the gel was dried for 30
minutes to adhere to the glass bottom plate), but these options negatively impact objective
working distance or resolution, respectively. To minimize gel shift during upright imaging with a
water dipping lens, we applied the following techniques to stabilize the gel in the imaging
chamber (50 mm WillCo Well). Following full expansion in water, we surrounded the gel with
2% agarose in the imaging chamber. This noticeably reduced gel movement during acquisition of
single images but shift was still detected during longer tile scans. We also tested re-embedding
the gelated sample in an unexpandable gel and covalently linking it to the glass imaging dish[2].
This completely eliminated movement of the gel and allowed us to take long tile scans on an
upright or inverted microscope. Unexpectedly, however, this also seemed to dampen the
fluorescence signal, which was irreversible.
339
340 **Table 1: Digestion time course.** Table showing mean expansion factor and fluorescence
341 intensities (FI) for each digestion time point and the unexpanded control. Also included is the
342 average background fluorescence intensity ("BG FI"), the normalized $\log_2$ transformed
343 fluorescence, and the number of cells ("N Cells") for each digestion condition.

Digest Time	Exp Factor	Mean FI	BG FI	Log₂ Norm FI	N Cells
no ExM	1.0 (±0.02)	69.0 (±5.39)	7.24 (±0.63)	5.795 (±0.12)	31
2 hour	2.6 (±0.08)	32.91 (±1.93)	3.56 (±0.4)	4.78 (±0.10)	29
4 hour	2.6 (±0.06)	42.79 (±4.77)	4.21 (±0.24)	5.024 (±0.14)	33
8 hour	3.0 (±0.08)	32.43 (±2.43)	2.34 (±0.01)	4.742 (±0.13)	35
overnight	3.2 (±0.05)	22.84 (±2.72)	2.28 (±0.03)	4.025 (±0.18)	31

344
345
346 **Table 2: Antibodies and conditions.** Table showing primary and secondary antibody conditions
347 used for expansion microscopy.

Primary Antibody	Host	Clonality	Supplier	Catalogue #	Antigen retrieval	Concentration	Incubation
GFP	Chicken	Polyclonal IgY	Abcam	ab13970	Not Required	1:500	72+ hrs at RT

GFP Booster Atto 488	Alpaca	Monoclonal VHH	Chromotek	gba488-100	Not Required	1:200	72+ hrs at RT
RFP	Rabbit	Polyclonal IgG	Rockland	600-401-379	Not Requred	1:500	72+ hrs at RT
RFP	Chicken	Polyclonal IgY	Rockland	600-901-379	Not Required	1:250	72+hrs at RT
GOLGA5/Golgin-84	Rabbit	Polyclonal IgG	Abcam	ab224040	Not Required	1:500	72+ hrs at RT
COX4	Rabbit	Polyclonal IgG	Synaptic Systems	298 003	5 min at 100C	1:500	72+hrs at RT
ERGIC-53/P58	Rabbit	Polyclonal IgG	Sigma Aldrich	E1031	Citrate at 80C	1:600	72+ hrs at RT
ST3GAL5	Rabbit	Polyclonal IgG	Sigma Aldrich	AV46358	Citrate at 80C	1:250	72+ hrs at RT
Secondary Antibody							
Goat anti Mouse IgG (H+L) Alexa 488	Goat	Polyclonal IgG	Invitrogen	A11029	NA	1:500	48+ hrs at RT
Goat anti Mouse IgG (H+L) Alexa 546	Goat	Polyclonal IgG	Invitrogen	A11030	NA	1:500	48+ hrs at RT
Goat anti Rabbit IgG (H+L) Alexa 488	Goat	Polyclonal IgG	Invitrogen	A11034	NA	1:500	48+ hrs at RT
Goat anti Rabbit IgG (H+L) Alexa 546	Goat	Polyclonal IgG	Invitrogen	A11035	NA	1:500	48+ hrs at RT
Goat anti Chicken IgY (H+L) Alexa 488	Goat	Polyclonal IgY	Invitrogen	A11039	NA	1:500	48+ hrs at RT
Goat anti Chicken IgY (H+L) Alexa 546	Goat	Polyclonal IgY	Invitrogen	A11040	NA	1:500	48+ hrs at RT

348
349 **Table 3: Detection of reporter labeled spines**

Tissue/Antibody	ExM (protK)	IHC (pre-ExM)	Spines
Am2-GFP	unexpanded	no IHC	+
		ag-no IHC	-
	expanded	no IHC	-
		ag-no IHC	-
Am2-GFP/GFP-chicken	unexpanded	IHC	++
		ag-IHC	+
	expanded	IHC	+++
		ag-IHC	++
Am2-GFP/GFP booster-alpaca	unexpanded	IHC	+
		ag-IHC	-
	expanded	IHC	-
		ag-IHC	nt
Am2-icre;tdTomato	unexpanded	no IHC	++

		ag-no IHC	+
	expanded	no IHC	-
		ag-no IHC	nt
Am2-icre;tdTomato/RFP-chicken	unexpanded	IHC	+++
		ag-IHC	+
	expanded	IHC	++
		ag-IHC	+
Am2-icre;tdTomato/RFP-rabbit	unexpanded	IHC	++++
		ag-IHC	nt
	expanded	IHC	++++
		ag-IHC	nt
- not detected; + rare SLM only; ++ OK; +++ Good; ++++ Excellent; nt Not tested			

350
351

352 **METHODS**
353
Animals
Adult male and female Amigo2-EGFP (RRID:MMRRC 033018-UCD, bred for at least 10
generations onto C57BL/6J background) or Amigo2-icreERT2;RosaTdTomato transgenic mice
were used. Amigo2-icreERT2;RosaTdTomato mice were generated by crossing Amigo2-
icreERT2 mice[33] to Ai14 mice (Jax #007914). Amigo2-icreER;ROSA-TdTomato mice were
given 2 or 3 daily intraperitoneal injections of tamoxifen (Sigma T5648, 100mg/kg freshly
dissolved daily in 100% ethanol then diluted 10-fold in corn oil and heated at 60C for 1 hour
until in solution). Mice were group-housed under a 12:12 light/dark cycle with access to food
and water *ad libitum*. All procedures were approved by the Animal Care and Use Committee of
Virginia Tech.
364
Immunofluorescence
Mice were anesthetized with 150mg/kg fatal plus solution and perfused with ice-cold 4%
paraformaldehyde. Amigo2-icreERT2;RosaTdTomato mice were perfused one week post
tamoxifen injections. Brains were dissected and post-fixed for at least 24 hours before sectioning
40 μm thick sections in the horizontal plane on a vibratome (Leica VT1000S). Sections to be
stained with COX4 underwent antigen retrieval by boiling free floating sections in 1.7ml tubes
for 3 min in nanopure water. All sections were washed in PBS and blocked for at least 1 hour in
5% Normal Goat Serum (NGS)/0.3% Triton-100x. See Table 3 for primary and secondary
antibodies and conditions. Antibodies were diluted in blocking solution and sections were
incubated for 72+ hours at room temperature (RT). After several rinses in PBS-T (0.3% Triton-
100x), sections were incubated in secondary antibodies for 48 hours at RT. Prior to imaging,
unexpanded sections were washed in PBS-T and mounted under Vectashield fluorescence media
with 4′,6-diamidino-2-phenylindole (DAPI) (Vector Laboratories).
378
Protein expansion microscopy solution preparation
Solutions were prepared as described by Asano et. al 2018[3]. Anchoring stock solution was
prepared by dissolving Acryloyl-X, SE (ThermoFisher A20770) in DMSO (1:100 w/v) and
stored at -20C. Monomer solution components were prepared by dissolving Sodium Acrylate in
npH20 (33% w/v, Sigma 408220), Acrylamide in npH20 (50% w/v, Sigma A9099), N, N'-
Methylenebisacylamide in npH20 (2% w/v, Sigma M7279). Monomer working solution was
prepared by adding 2.25mL of 33% SA solution (8.6% w/v), 0.5 mL of 50% Acrylamide
solution (2.5% w/v), 0.75 mL of 2% N,N-Methylenebisacrylamide solution (0.15% w/v), 4 mL
of 5M NaCl (11.7% w/v), and 1 mL of 10X PBS. Inhibitor stock was prepared by dissolving 4-
Hydroxy-TEMPO (0.5% w/v, Sigma 176141) in npH2O and initiator stock was made by
dissolving Ammonium persulfate in npH2O (10% w/v, Sigma 248614). Accelerator solution was
prepared by diluting TEMED in npH2O (10% v/v, Sigma T7024) immediately before use. All
solutions except the TEMED accelerator were prepared before use and stored at -20C.

Protein expansion microscopy

4X protein expansion microscopy (proExM) was carried out on horizontal mouse brain sections containing dorsal hippocampus as described in Asano et al 2018[3]. Sections were incubated overnight in Acryloyl-X stock/PBS (1:100, ThermoFisher, A20770) at RT in the dark. Following incubation, the slices were washed twice with PBS for 15 minutes each at RT. The gelation solution was prepared by adding 384 uL of monomer solution, 8 uL 4-Hydroxy-TEMPO inhibitor (1:200 w/v, Sigma Aldrich, 176141), 8uL TEMED accelerator (10% v/v, Sigma Aldrich, T7024), and lastly 8uL of APS initiator (10% w/v, Sigma Aldrich, 248614) for each section. Sections were then incubated in the gelation solution for 30-45 minutes at 4C in the dark. Gelated sections were placed on gelation chambers constructed from microscope slides with coverslips as spacers. Our gelation chambers produce gels with the thickness of two type No. 1.5 coverslips (~0.3mm thick). The chambers were filled with gelation solution and allowed to incubate at 37 C for 2 hours in a humidified container. Following gelation incubation, the gelation chamber was deconstructed to uncover the gelated brain section. To remove the gel from the chamber, digestion solution without Proteinase K was applied and a coverslip was used to gently remove the sample. Digestion solution containing Proteinase K (8U/mL, New England BioLabs, P8107S) was applied to gels and allowed to digest for 2-16 hours (see Results) at room temperature. Upon completion of digestion, gels were stained with DAPI (Sigma Aldrich, D9542, 1:10,000 in PBS) for 10 minutes at room temperature with shaking. The gels were then washed twice for 10 minutes with npH2O to remove excess DAPI and fully expand the gel.

Stabilizing ExM Gels with Agarose

To prevent movement during imaging, gels were fully expanded in water or 0.001X PBS in WillCo wells (HBSB-5040) and reversibly immobilized by applying liquid 2% agarose around and on top of the gel (in areas not containing tissue). Following application, the gel embedded with agarose was placed at 4C for at least 15 minutes to allow the agarose to fully solidify prior to imaging.

Re-embedding and Linking Gels to Imaging Dish

Re-embedding and covalently linking gels to a WillCo well allowed long-term imaging without gel shifting as described by Tillberg et al 2016[2]. First, the gel was completely expanded in water and then incubated in a non-expanding re-embedding solution (3% w/v acrylamide, 0.15% w/v N,N-methylenebisacylamide, 0.05% w/v APS, 0.05% w/v TEMED, and 5mM Tris). The gels were incubated with shaking at room temperature for 20 minutes. The gel was then transferred to a Bind-Silane treated WillCo Well plate and covered with a coverslip. Fresh re-embedding solution was then lightly applied surrounding the sample, which was then incubated at 37C for 1.5-2hrs without shaking. Once the re-embedding solution gelated, the coverslip was removed from the covalently linked gel and could be imaged.

Bind Silane Treatment of Imaging Dishes
Immediately before use, imaging dishes were treated with a bind silane solution as described by
Tillberg et al 2016[2]. Before bind silane treatment, imaging dishes were briefly washed with
npH2O, 100% Ethanol, and then allowed to dry. Bind silane solution (80% v/v EtOH, 2% v/v
acetic acid, 0.05% v/v Bind-Silane) was then applied with shaking for 5 minutes. The dish was
then washed with 100% EtOH and allowed to dry before usage.
438
Digestion Time Course Experiment
To assess the effect of digestion time on tissue expansion and fluorescence retention, we ran a
digestion time course experiment by varying the amount of time the ExM gels were in digestion
solution-- either 2 hours, 4 hours, 8 hours or 16 hours (overnight). The brains of two Amigo2-
EGFP mice (One male and one female, 21-23 weeks old) were processed as above. During a
pilot run, we noted that samples with digestion > 2 hours lost the majority of EGFP fluorescence,
making it impossible to acquire images with identical parameters for direct comparison.
Therefore, to boost the EGFP signal, approximately ten sections per brain (two sections per
condition per mouse) were first immunostained in a 24-well plate as described in the
immunohistochemistry methods, with a primary chicken antibody against EGFP (Abcam,
ab13970; 1:500 concentration) at RT for 3 days, followed by a secondary antibody (Invitrogen
Alexa-488, A11039; 1:500 concentration) incubation for 48 hours. As a control, a few sections
were set aside after immunostaining to mount on slides without expanding.
452
Sections processed for ExM were anchored in Acryloyl-X overnight, washed twice with 1X PBS
and incubated in gelation solution for 30 min at 4C the following day (see "Protein Expansion
Methods" for more detail). The gels were incubated in a humid environment at 37C for 2 hours
to set, and then carefully removed from the chamber and placed in a digestion solution with 8
U/mL of Proteinase K (New England Bio, Cat # P8107S) for the designated period of time. All
gels in this experiment were digested on a shaker at room temperature. At the end of the
digestion time, the digestion solution was replaced with 1X PBS several times to wash the gels.
Gels were stored in 1X PBS in the dark at 4C until imaging, at which point the PBS was replaced
with npH2O to fully expand the gels. Gels were imaged at 10X (EC-Plan-Neofluar lens; 10x; 0.3
NA; 5.2mm WD) taking $10\mu m$ steps for distance on a Zeiss 710 confocal microscope with the
same image acquisition parameters, as described in more detail under "image acquisition".
464
Digestion Time Course Analysis
The ExM images from the time course experiment were analyzed with the image processing
program Fiji (v.+03..30, NIH). Before analysis, images were flattened across the Z dimension
with the *"Z Project"* function to get an idea of how many GFP positive cells were present. This
flattened image was only used for the selection of cells. Cells were excluded from analysis if
they met any of the following criteria: 1) Incomplete cell (i.e. on the image border); 2) Any cell
overlapping in Z with a cell already analyzed; 3) Cells with an ambiguous border or that were

difficult to differentiate. Given these exclusion criteria, an average of 32 cells were analyzed for
each time point across both animals. For some time points, cells were included from multiple
hippocampal sections from the same animal.

For the fluorescence analysis, each cell body meeting the inclusion criteria was manually traced
with the freehand selection tool in a single z section at the widest point of the cell. Once the cells
were traced, the *"measure"* tool in Fiji was used to measure the mean intensity and the area
within each individual cell ROI. To subtract background signal, one cell ROI was selected (at
random) and moved to a location without EGFP signal and the background mean intensity was
measured. The background mean intensity value was subtracted from the mean fluorescence
intensity of each cell from the same image, resulting in a normalized mean fluorescence. The
area of the cell body ROIs were used to calculate expansion factor by comparing it to the average
area of 37 unexpanded cell bodies, which was calculated to be 300 μm^2. To calculate the linear
expansion factor for each cell, we used the below formula:

$$Linear\ Expansion\ Factor = \frac{\sqrt{(cell\ area)}}{\sqrt{(avg\ non\ expanded\ cell\ area)}}$$

487
488
A total of 159 cells were included in the analyses (average of 32 +/- 4 cells per time point from
two brains). The mean fluorescence and expansion factor measurements were imported into
Python (v 3.7, installed with the Anaconda distribution), where the average mean fluorescence
and average linear expansion factor were calculated for each time point, including the
unexpanded control (see Fig. 2). The mean fluorescence data was not normally distributed
(Shapiro-Wilk test for normality; p < 0.0001), so the mean fluorescence data was log_2
transformed and 11 cell means were removed as outliers, pre-defined as two standard deviations
from the mean. These same outliers were also removed from the expansion factor data; however,
because this data was normally distributed (Shapiro-Wilk; p = 0.235), it was not transformed.
One-way ANOVAs were run using an ordinary least squares model with the StatsModels
package (ols, ANOVA_lm, statsmodels v0.11.1) comparing fluorescence intensity and linear
expansion factors across digestion time points. After overall significance was reached, pairwise
post-hoc Tukey's multiple comparison tests (pairwise_tukeyhsd, statsmodels v0.11.1) were
performed to determine which digestion time points were significantly different (see Fig. 2D).

Analysis of Expanded and Unexpanded Mitochondria
To determine if the ProExM protocol affects our ability to resolve and quantify mitochondria, we
analyzed the number and size of expanded mitochondria compared to unexpanded mitochondria.
Tissue from an adult male EGFP reporter mouse was immunostained with COX4 to label
mitochondria and expanded using our optimized ExM protocol with overnight digestion in
Proteinase K (see Expansion Microscopy methods). An adult male tdTomato reporter mouse was
similarly processed and immunostained with COX4 but not expanded. Confocal images were

taken of both samples for image analysis in FIJI. Note that the COX4 signal was imaged in
different color channels for the expanded and unexpanded samples (546nm vs 488nm,
respectively).
514
Confocal images were imported into Fiji and converted to 8-bit for the analysis. An intensity
threshold was chosen separately for each image which best represented the signal in the original
image. Using the reporter label as a guide, four cells were analyzed from the expanded sample
and five cells from the unexpanded sample. Cells were chosen using the same criteria as for the
digestion time course analysis. An ROI for each cell was drawn by fitting an oval to the cell
body using the oval selection tool, rotating if necessary, and the signal outside of the cell's ROI
was removed for the analysis. The "*nucleus counter*" plug-in from the FIJI "Cookbook"
microscopy analysis collection was used to segment mitochondria within the ROI of each cell
analyzed. The size threshold used for the expanded sample was adjusted for expansion factor,
which was calculated using the ratio of the cell body diameters in the expanded images
compared to the unexpanded images. The calculated expansion factor for the expanded images
was 3x in the X and Y dimensions-- an expansion factor of 9x in total area.
527
Once the mitochondria were properly segmented, FIJI's "*measure*" feature was used to measure
the number, area and intensity of each individual mitochondrial ROI. A custom Python code was
written to calculate and plot the averages of the mitochondria number, size and total area for the
expanded cells and the unexpanded cells. To account for the effect of expansion on size, the area
of each mitochondria was divided by the area of the soma to get a normalized mitochondria area.
The mitochondrial coverage was calculated by dividing the total summed mitochondrial area in
the cell by the area of the cytoplasm (*soma area – nucleus area*) and converting it to a
percentage.
536
Image acquisition
Images were acquired on an upright Zeiss 710 confocal microscope equipped with a motorized
stage, 488/561/633 laser lines, and 20X/1.0 NA water immersion or 20X/0.4 NA air lenses. Gels
were expanded by washing with 0.001X PBS three times and by incubating overnight at RT in
0.001X PBS. Gels were immobilized in 50mm glass bottom wells (WillCo Wells, HBSB-5040)
by applying 2% agarose to the edges of the gels. Gels were then imaged in a fully expanded state
in or 0.001X PBS.
544
ExM image processing
Czi files were imported into Fiji, individual channel images were split and adjusted for
brightness and contrast equivalently across conditions, then images were viewed with the volume
viewer plugin (v. 2.01.2). If images showed considerable shift or too many neurons overlapped, a
subset stack was created without the offending Z sections. Mode was set to max-projection and
interpolation was set to tricubic smooth with z-aspect and sampling optimized per image

(typically 0.5-2.0 for each parameter). Snapshots were taken in the XY plane at 1.0 and 2.0 scale
in grayscale and 1D. The resulting snapshot images were imported into Photoshop (v. 21.2) and
converted to 300 dpi. Any further brightness and contrast edits done in photoshop were minimal
and applied equivalently to all comparable images in the figure.
555

556   **Statistical analyses**
557   Statistical analyses were done in python (v 3.7) with an alpha of 0.05 considered significant.
558

Acknowledgements
Research reported in this publication was supported by the National Institute of Mental Health of
the NIH under award R00MH109626 to S.F. We thank the members of the Dudek lab for
feedback and critically reading this manuscript, as well as the Virginia Tech animal care staff for
their support.

Author Contributions
Conceptualization, S.F.; Methodology, L.A.C., K.E.P., N.A.S., D.L., S.F.; Formal Analysis,
L.A.C., K.E.P., N.A.S., S.F.; Investigation, L.A.C., K.E.P., N.A.S., D.L., S.F.; Writing –
Original Draft, L.A.C., K.E.P., S.F.; Writing – Review & Editing, L.A.C., K.E.P., S.F.;
Visualization, L.A.C., K.E.P., S.F.; Supervision, S.F.; Funding Acquisition, S.F.
570

571   **Declaration of Interests**
572   The authors declare no conflict of interest or competing financial interest.
573
574

FIGURE LEGENDS

Fig. 1: ExM workflow for visualizing subcellular organelles in fixed brain tissue.
Schematic of immunostaining and proExM workflows tested to optimize visualization of
organelles in genetically-labeled neurons.

**Fig. 2: The effect of Proteinase K digestion time on fluorescence intensity and expansion
factor.**
A) Representative single z-section images of unexpanded (i) and expanded GFP+ CA2 neurons
after 2-hour (ii), 4-hour (iii), 8-hour (iv), or overnight (v) digestion with Proteinase K. Image
acquisition parameters were identical for each condition, except for the unexpanded condition.
B) GFP fluorescence intensity measured in the cell soma after the different digestion times.
Fluorescence was corrected for background and $\log_2$ transformed. Line represents the median.
Digestion time had a significant effect on GFP fluorescence (one-way ANOVA; F-stat: 20.96, p-
value: 8.17E-14, 32 ± 4 cells per time point from two animals, 159 total cells).
C) Expansion factor of the expanded cell somas in **B**. Expansion factors were calculated using
cell soma areas relative to the average unexpanded cell soma area. Line represents the median.
Digestion time had a significant effect on expansion factor (one-way ANOVA; F-stat: 21.07, p-
value: 4.26E-11).
D) Matrix of p-values visualizing the results of pairwise Tukey's post hoc tests comparing mean
fluorescence (upper diagonal) or expansion factor (lower diagonal) at each digestion time point.
Matrix is color coded by p-value (red = significant; grey = not significant; α=0.05).
E) Effect of digestion time on fluorescence intensity (cyan; left axis) and expansion factor
(orange; right axis). Plot shows the mean and 95% confidence intervals from **B** and **C**.
F) Unexpanded (i) and overnight expanded (ii) max-projection 10X confocal images of GFP+
CA2 cells. Images are the same as in **A**(i) and (v). Imaging parameters were optimized separately
to obtain the best image of both.
Scale bars = (**A**, **F**) 50 μm.

Fig. 3: No appreciable effect of tissue depth on expansion factor.
A) Average cell soma area (in μm^2) by Z section depth for unexpanded control and digestion
conditions. Cell soma area was measured in the Z section with the widest soma diameter.
Because of the overall increase in depth with expansion, cells were binned by Z section to
achieve ~6 bins per time point. N = 32 ± 4 cells per time point.
B) Background subtracted average cell soma fluorescence by Z section depth. Binning of the
data is the same as in A. The data are not $\log_2$ transformed as in Fig. 2.
C) Representative confocal images of GFP+ CA2 cell somas after 2-hour (left panel) or
overnight (right panel) digestion at three Z-section depths. Cell soma size and fluorescence is
variable in the Z dimension in unexpanded conditions, which does not appreciably change under
difference digestion conditions.

615   Error bars show the standard deviation. Scale bar is 50μm.
616
Fig. 4: ProExM pipeline comparison for visualizing mitochondria.
A) 20X Max-projection image of a section immunostained for COX4 (i) and RFP (iii) pre-ExM
with autoclave digestion. Images in panels A-D were acquired with identical imaging parameters
to directly compare conditions. Brightness and contrast were adjusted to the same extent across
the entire image.
B) 20X Max-projection image of a boiled section immunostained for COX4 (i) and RFP (ii) pre-
ExM with overnight Proteinase K digestion.
C) 20X Max-projection image of a section immunostained for COX4 (i) and RFP (ii) post-ExM
with autoclave digestion.
D) 20X Max-projection image of a boiled section immunostained for COX4 (i) and RFP (ii)
post-ExM with overnight Proteinase K digestion
E) 40X Max-projection image from a subset of Z sections from a boiled, unexpanded section
immunostained for COX4 and RFP. Inset is a single Z section from the indicated representative
cell (white asterisk) with mitochondria outlined as quantified in G-I. Imaging parameters were
optimized per condition in E-F.
F) 20X Max-projection image from a subset of Z sections from a boiled section immunostained
for COX4 and GFP pre-ExM with overnight Proteinase K digestion as in **B.**
G) The average number of mitochondria per cell in unexpanded **(E)** and expanded **(F)** sections.
H) Mitochondria area per cell (normalized by soma area). Line reflects median and whiskers are
twice the inner quartile range.
I) The average percent cytoplasm area (soma area - nuclear area) containing mitochondria. Note
that ExM better resolves densely packed mitochondria, resulting in greater number, smaller
sized, and fewer percent area per cell soma compared with no-ExM.
Scale bars: 50 μm (**A,E,F**) and 5 μm (insets).
641
Fig. 5: ProExM pipeline comparison for visualizing Golgi.
A) 20X Max-projection image of a section immunostained for GOLGA5 (i) and GFP (iii) pre-
ExM with autoclave digestion. (ii) Insets are zoomed images of the dotted box in (i). Images in
panels **A-C** were acquired with identical imaging parameters to directly compare conditions.
Brightness and contrast were adjusted to the same extent across the entire image.
B) 20X Max-projection image of a section immunostained for GOLGA5 (i) and GFP (iii) pre-
ExM with overnight Proteinase K digestion.
C) 20X Max-projection image of a section immunostained for GOLGA5 (i) and GFP (iii) post-
ExM with overnight Proteinase K digestion.
D) 40X Max-projection of a subset of Z sections from an unexpanded section immunostained for
GOLGA5 (i) and RFP (iii). Imaging parameters were optimized per condition in **D-E.**
E) 20X Max-projection of a subset of Z sections from image in (**B**) immunostained pre-ExM
with overnight Proteinase K digestion. Note the increase in Golgi complexity that is resolved

with autoclave or Proteinase K digested gels compared with unexpanded sections, although
fluorescence intensity was better retained with Proteinase K digestion.
F) Line plot profile showing the fluorescence intensity of unexpanded Golgi (cyan) and Golgi
after IHC-pre-ExM with Proteinase K digestion (magenta). Measured Golgi are inset in **Diii** and
E.
Scale bars: 50µm (**Ai, Di, E**) and 10µm (**Aii, Dii**).
661
Fig. 6: Resolving dendritic spines with ExM.
A) 20X with 3X zoom max-projection image (i) of a section immunostained for RFP pre-ExM
with overnight Proteinase K digestion. This mouse received 2 tamoxifen injections. (ii) Insets are
2X zoomed images of the dotted box in (i). Images were acquired with imaging parameters
optimal per condition. Brightness and contrast were adjusted per condition across the entire
image.
B) 40X with 2X zoom max-projection image of an unexpanded tdTomato section. This mouse
received 2 tamoxifen injections.
C) 20X with 2X zoom max-projection image of a section immunostained for GFP pre-ExM with
overnight Proteinase K digestion.
D) 40X with 2X zoom max-projection image from an unexpanded section immunostained for
GFP. Note the increased background labeling and density of dendritic branches in the
unexpanded images, making dendrites more difficult to trace compared with expanded images.
Scale bars: 25µm (Ai, Bi, Ci, Di) and 5µm (Aii, Bii, Cii, Dii).
676

REFERENCES

1. Chozinski TJ, Halpern AR, Okawa H, Kim H-J, Tremel GJ, Wong ROL, Vaughan JC. Expansion microscopy with conventional antibodies and fluorescent proteins. Nature Methods. Nature Publishing Group; 2016 Jun;13(6):485–488.

2. Tillberg PW, Chen F, Piatkevich KD, Zhao Y, Yu C-CJ, English BP, Gao L, Martorell A, Suk H-J, Yoshida F, DeGennaro EM, Roossien DH, Gong G, Seneviratne U, Tannenbaum SR, Desimone R, Cai D, Boyden ES. Protein-retention expansion microscopy of cells and tissues labeled using standard fluorescent proteins and antibodies. Nat Biotechnol. 2016;34(9):987–992. PMCID: PMC5068827

3. Asano SM, Gao R, Wassie AT, Tillberg PW, Chen F, Boyden ES. Expansion Microscopy: Protocols for Imaging Proteins and RNA in Cells and Tissues. Curr Protoc Cell Biol. 2018;80(1):e56. PMCID: PMC6158110

4. Ku T, Swaney J, Park J-Y, Albanese A, Murray E, Cho JH, Park Y-G, Mangena V, Chen J, Chung K. Multiplexed and scalable super-resolution imaging of three-dimensional protein localization in size-adjustable tissues. Nat Biotechnol. 2016;34(9):973–981. PMCID: PMC5070610

5. Park H-E, Choi D, Park JS, Sim C, Park S, Kang S, Yim H, Lee M, Kim J, Pac J, Rhee K, Lee J, Lee Y, Lee Y, Kim S-Y. Scalable and Isotropic Expansion of Tissues with Simply Tunable Expansion Ratio. Adv Sci (Weinh). 2019 Nov;6(22):1901673. PMCID: PMC6864509

6. Gambarotto D, Zwettler FU, Le Guennec M, Schmidt-Cernohorska M, Fortun D, Borgers S, Heine J, Schloetel J-G, Reuss M, Unser M, Boyden ES, Sauer M, Hamel V, Guichard P. Imaging cellular ultrastructures using expansion microscopy (U-ExM). Nat Methods. 2019;16(1):71–74. PMCID: PMC6314451

7. Katoh Y, Chiba S, Nakayama K. Practical method for superresolution imaging of primary cilia and centrioles by expansion microscopy using an amplibody for fluorescence signal amplification. Mol Biol Cell. 2020 15;31(20):2195–2206. PMCID: PMC7550703

8. Shi X, Li Q, Dai Z, Tran AA, Feng S, Ramirez AD, Lin Z, Wang X, Chow TT, Seiple IB, Huang B. Label-retention expansion microscopy. bioRxiv. Cold Spring Harbor Laboratory; 2019 Jul 2;687954.

9. Truckenbrodt S, Sommer C, Rizzoli SO, Danzl JG. A practical guide to optimization in X10 expansion microscopy. Nature Protocols. Nature Publishing Group; 2019 Mar;14(3):832–863.

10. Chen F, Tillberg PW, Boyden ES. Expansion microscopy. Science. 2015 Jan 30;347(6221):543–548. PMID: 25592419

11. Wassie AT, Zhao Y, Boyden ES. Expansion microscopy: principles and uses in biological research. Nat Methods. 2019;16(1):33–41. PMCID: PMC6373868

12. Kunz TC, Götz R, Gao S, Sauer M, Kozjak-Pavlovic V. Using Expansion Microscopy to
Visualize and Characterize the Morphology of Mitochondrial Cristae. Front Cell Dev Biol.
2020;8:617. PMCID: PMC7373753

13. Büttner M, Lagerholm CB, Waithe D, Galiani S, Schliebs W, Erdmann R, Eggeling C,
Reglinski K. Challenges of Using Expansion Microscopy for Super-resolved Imaging of
Cellular Organelles. Chembiochem. 2020 Oct 13; PMID: 33049107

14. Pesce L, Cozzolino M, Lanzanò L, Diaspro A, Bianchini P. Measuring expansion from
macro- to nanoscale using NPC as intrinsic reporter. J Biophotonics.
2019;12(8):e201900018. PMCID: PMC7065622

15. Sun D, Fan X, Zhang H, Huang Z, Tang Q, Li W, Bai J, Lei X, Chen X. Click-ExM enables
expansion microscopy for all biomolecules. bioRxiv. Cold Spring Harbor Laboratory; 2020
Mar 20;2020.03.19.998039.

16. Pernal SP, Liyanaarachchi A, Gatti DL, Formosa B, Pulvender R, Kuhn ER, Ramos R, Naik
AR, George K, Arslanturk S, Taatjes DJ, Jena BP. Nanoscale imaging using differential
expansion microscopy. Histochem Cell Biol. 2020 Jun;153(6):469–480. PMID: 32193594

17. Zhao Y, Bucur O, Irshad H, Chen F, Weins A, Stancu AL, Oh E-Y, DiStasio M, Torous V,
Glass B, Stillman IE, Schnitt SJ, Beck AH, Boyden ES. Nanoscale imaging of clinical
specimens using pathology-optimized expansion microscopy. Nat Biotechnol. 2017
Aug;35(8):757–764. PMCID: PMC5548617

18. Götz R, Panzer S, Trinks N, Eilts J, Wagener J, Turrà D, Di Pietro A, Sauer M, Terpitz U.
Expansion Microscopy for Cell Biology Analysis in Fungi. Front Microbiol. 2020;11:574.
PMCID: PMC7147297

19. Düring DN, Rocha MD, Dittrich F, Gahr M, Hahnloser RHR. Expansion Light Sheet
Microscopy Resolves Subcellular Structures in Large Portions of the Songbird Brain. Front
Neuroanat. 2019;13:2. PMCID: PMC6365838

20. Jiang N, Kim H-J, Chozinski TJ, Azpurua JE, Eaton BA, Vaughan JC, Parrish JZ.
Superresolution imaging of Drosophila tissues using expansion microscopy. Mol Biol Cell.
2018 15;29(12):1413–1421. PMCID: PMC6014096

21. Cahoon CK, Yu Z, Wang Y, Guo F, Unruh JR, Slaughter BD, Hawley RS. Superresolution
expansion microscopy reveals the three-dimensional organization of the Drosophila
synaptonemal complex. Proc Natl Acad Sci U S A. 2017 15;114(33):E6857–E6866. PMCID:
PMC5565445

22. Gao R, Asano SM, Upadhyayula S, Pisarev I, Milkie DE, Liu T-L, Singh V, Graves A,
Huynh GH, Zhao Y, Bogovic J, Colonell J, Ott CM, Zugates C, Tappan S, Rodriguez A,
Mosaliganti KR, Sheu S-H, Pasolli HA, Pang S, Xu CS, Megason SG, Hess H, Lippincott-
Schwartz J, Hantman A, Rubin GM, Kirchhausen T, Saalfeld S, Aso Y, Boyden ES, Betzig
E. Cortical column and whole-brain imaging with molecular contrast and nanoscale
resolution. Science. 2019 Jan 18;363(6424).

23. Fecher C, Trovò L, Müller SA, Snaidero N, Wettmarshausen J, Heink S, Ortiz O, Wagner I,
Kühn R, Hartmann J, Karl RM, Konnerth A, Korn T, Wurst W, Merkler D, Lichtenthaler SF,
Perocchi F, Misgeld T. Cell-type-specific profiling of brain mitochondria reveals functional
and molecular diversity. Nature Neuroscience. 2019 Oct;22(10):1731–1742.

24. Karagiannis ED, Kang JS, Shin TW, Emenari A, Asano S, Lin L, Costa EK, Consortium
IGC, Marblestone AH, Kasthuri N, Boyden ES. Expansion Microscopy of Lipid Membranes.
bioRxiv. Cold Spring Harbor Laboratory; 2019 Nov 4;829903.

25. Martínez GF, Gazal NG, Quassollo G, Szalai AM, Cid-Pellitero ED, Durcan TM, Fon EA,
Bisbal M, Stefani FD, Unsain N. Quantitative expansion microscopy for the characterization
of the spectrin periodic skeleton of axons using fluorescence microscopy. Sci Rep. 2020
19;10(1):2917. PMCID: PMC7031372

26. Scheible MB, Tinnefeld P. Quantifying Expansion Microscopy with DNA Origami
Expansion Nanorulers. bioRxiv. Cold Spring Harbor Laboratory; 2018 Feb 14;265405.

27. Vanheusden M, Vitale R, Camacho R, Janssen KPF, Acke A, Rocha S, Hofkens J.
Fluorescence Photobleaching as an Intrinsic Tool to Quantify the 3D Expansion Factor of
Biological Samples in Expansion Microscopy. ACS Omega. 2020 Mar 17;5(12):6792–6799.
PMCID: PMC7114699

28. Min K, Cho I, Choi M, Chang J-B. Multiplexed expansion microscopy of the brain through
fluorophore screening. Methods. 2020 01;174:3–10. PMID: 31326595

29. Steward O, Schuman EM. Protein synthesis at synaptic sites on dendrites. Annu Rev
Neurosci. 2001;24:299–325. PMID: 11283313

30. Pierce JP, Mayer T, McCarthy JB. Evidence for a satellite secretory pathway in neuronal
dendritic spines. Curr Biol. 2001 Mar 6;11(5):351–355. PMID: 11267872

31. Bowen AB, Bourke AM, Hiester BG, Hanus C, Kennedy MJ. Golgi-independent secretory
trafficking through recycling endosomes in neuronal dendrites and spines. Elife. 2017 06;6.
PMCID: PMC5624785

32. Kennedy MJ, Ehlers MD. ORGANELLES AND TRAFFICKING MACHINERY FOR
POSTSYNAPTIC PLASTICITY. Annual Review of Neuroscience. 2006 Jul 21;29(1):325–
362.

33. Alexander GM, Brown LY, Farris S, Lustberg D, Pantazis C, Gloss B, Plummer NW, Jensen
P, Dudek SM. CA2 neuronal activity controls hippocampal low gamma and ripple
oscillations. Elife. 2018 02;7. PMCID: PMC6251629

34. Schindelin J, Arganda-Carreras I, Frise E, Kaynig V, Longair M, Pietzsch T, Preibisch S,
Rueden C, Saalfeld S, Schmid B, Tinevez J-Y, White DJ, Hartenstein V, Eliceiri K,
Tomancak P, Cardona A. Fiji: an open-source platform for biological-image analysis. Nat
Methods. 2012 Jun 28;9(7):676–682. PMCID: PMC3855844

Figure 1: ExM workflow for visualizing subcellular organelles in fixed brain tissue.
Schematic of immunostaining and proExM workflows tested to optimize visualization of organelles in genetically-labeled neurons.

Figure 2: The effect of Proteinase K digestion time on fluorescence intensity and expansion factor.

A) Representative single z-section images of unexpanded (i) and expanded GFP+ CA2 neurons after 2-hour (ii), 4-hour (iii), 8-hour (iv), or overnight (v) digestion with Proteinase K. Image acquisition parameters were identical for each condition, except for the unexpanded condition.

B) GFP fluorescence intensity measured in the cell soma after the different digestion times. Fluorescence was corrected for background and $\log_2$ transformed. Line represents the median. Digestion time had a significant effect on GFP fluorescence (one-way ANOVA; F-stat: 20.96, p-value: 8.17E-14, 32 ± 4 cells per time point from two animals, 159 total cells).

C) Expansion factor of the expanded cell somas in B. Expansion factors were calculated using cell soma areas relative to the average unexpanded cell soma area. Digestion time had a significant effect on expansion factor (one-way ANOVA; F-stat: 21.07, p-value: 4.26E-11).

D) Matrix of p-values visualizing the results of pairwise Tukey's post hoc tests comparing mean fluorescence (upper diagonal) or expansion factor (lower diagonal) at each digestion time point. Matrix is color coded by p-value (red = significant; grey = not significant; α=0.05).

E) Effect of digestion time on fluorescence intensity (cyan; left axis) and expansion factor (orange; right axis). Plot shows the mean and 95% confidence intervals from **B** and **C**.

F) Unexpanded (i) and overnight expanded (ii) max-projection 10X confocal images of GFP+ CA2 cells. Images are the same as in **A**(i) and (v). Imaging parameters were optimized separately to obtain the best image of both. Scale bars = (**A**, **F**) 50 µm.

Figure 3: No appreciable effect of tissue depth on expansion factor.
A) Average cell soma area (in μm²) by Z section depth for unexpanded control and digestion conditions. Cell soma area was measured in the Z section with the widest soma diameter. Because of the overall increase in depth with expansion, cells were binned by Z section to achieve ~6 bins per time point. N = 32 ± 4 cells per time point.
B) Background subtracted average cell soma fluorescence by Z section depth. Binning of the data is the same as in A. The data are not $\log_2$ transformed as in Fig 2.
C) Representative confocal images of GFP+ CA2 cell somas after 2-hour (left panel) or overnight (right panel) digestion at three Z-section depths. Cell soma size and fluorescence is variable in the Z dimension in unexpanded conditions, which does not appreciably change under difference digestion conditions. Error bars show the standard deviation. Scale bar is 50μm.

Figure 4: ProExM pipeline comparison for visualizing mitochondria.

A) 20X Max-projection image of a section immunostained for COX4 (i) and RFP (ii) pre-ExM with autoclave digestion. Images in panels A-D were acquired with identical imaging parameters to directly compare conditions. Brightness and contrast were adjusted to the same extent across the entire image.

B) 20X Max-projection image of a boiled section immunostained for COX4 (i) and RFP (ii) pre-ExM with overnight Proteinase K digestion.

C) 20X Max-projection image of a section immunostained for COX4 and RFP (ii) post-ExM with autoclave digestion.

D) 20X Max-projection image of a boiled section immunostained for COX4 and RFP (ii) post-ExM with overnight Proteinase K digestion

E) 40X Max-projection image from a subset of Z sections from a boiled, unexpanded section immunostained for COX4 and RFP. Inset is a single Z section from the indicated representative cell (white asterisk) with mitochondria outlined as quantified in **G-I**. Imaging parameters were optimized per condition in **E-F**.

F) 20X Max-projection image from a subset of Z sections from a boiled section immunostained for COX4 and GFP pre-ExM with overnight Proteinase K digestion as in **B**.

G) The average number of mitochondria per cell in unexpanded (**E**) and expanded (**F**) sections.

H) Mitochondria area per cell (normalized by soma area). Line reflects median and whiskers are twice the inner quartile range.

I) The average percent cytoplasm area (soma area - nuclear area) containing mitochondria. Note that ExM better resolves densely packed mitochondria, resulting in greater number, smaller sized, and fewer percent area per cell soma compared with non-ExM.

Scale bars: 50 μm (**A,E,F**) and 5 μm (insets).

Figure 5: ProExM pipeline comparison for visualizing Golgi.

A) 20X Max-projection image of a section immunostained for GOLGA5 (i) and GFP (iii) pre-ExM with autoclave digestion. (ii) Insets are zoomed images of the dotted box in (i). Images in panels A-C were acquired with identical imaging parameters to directly compare conditions. Brightness and contrast were adjusted to the same extent across the entire image.

B) 20X Max-projection of a section immunostained for GOLGA5 (i) and GFP (iii) pre-ExM with overnight Proteinase K digestion.

C) 20X Max-projection image of a section immunostained for GOLGA5 (i) and GFP (iii) post-ExM with overnight Proteinase K digestion.

D) 40X Max-projection of a subset of Z sections from an unexpanded section immunostained for GOLGA5 (i) and RFP (iii). Imaging parameters were optimized per condition in D-E.

E) 20X Max-projection of a subset of Z sections from image in (B) immunostained pre-ExM with overnight Proteinase K digestion. Note the increase in Golgi complexity that is resolved with autoclave or Proteinase K digested gels compared with unexpanded sections, although fluorescence intensity was better retained with Proteinase K digestion.

F) Line plot profile showing the fluorescence intensity of unexpanded Golgi (cyan) and Golgi after IHC-pre-ExM with Proteinase K digestion (magenta). Measured Golgi are insets in Diii and E.

Scale bars: 50µm (Ai, Di, E) and 10µm (Aii, Dii).

Figure 6: Resolving dendritic spines with ExM.
A) 20X with 3X zoom max-projection image (i) of a section immunostained for RFP pre-ExM with overnight Proteinase K digestion. This mouse received 2 tamoxifen injections. (ii) Insets are 2X zoomed images of the dotted box in (i). All images were acquired with parameters optimal per condition. Brightness and contrast were adjusted per condition across the entire image.
B) 40X with 2X zoom max-projection image of an unexpanded tdTomato section. This mouse received 2 tamoxifen injections.
C) 20X with 2X zoom max-projection image of a section immunostained for GFP pre-ExM with overnight Proteinase K digestion. This mouse received 2 tamoxifen injections.
D) 40X with 2X zoom max-projection image from an unexpanded section immunostained for GFP. Note the increased background labeling and density of dendritic branches in the unexpanded images, making dendrites more difficult to trace compared with expanded images.
Scale bars: 25µm (**Ai**, **Bi**, **Ci**, **Di**) and 5µm (**Aii**, **Bii**, **Cii**, **Dii**).