Networks of marine protected areas under fluctuating connectivity: The importance of within-MPA dynamics

Ridouan Bani
Tianna Peller
Justin Marleau
Marie-Josée Fortin
Frédéric Guichard

Abstract The design of marine protected areas (MPAs) has been optimized under assumptions of spatially and temporally homogeneous larval dispersal, despite complex spatiotemporal patterns displayed by ocean currents. Here we studied the effect of dispersal variability on the effectiveness of MPA networks across scales. We adopted a nested approach integrating the dynamics of both within and among MPA connectivity into a stochastic metapopulation model and derived stability over MPA networks by partitioning within and among MPA contributions in relation to the spatial resolution of within-MPA connectivity. We applied this framework over a range of dispersal traits (spawning time and pelagic larval duration) and MPA network configurations, based on simulated biophysical connectivity along the northeast Pacific coast. Our re-

R. Bani
Department of Biology, McGill University, 1205 Docteur Penfield, Montréal, Québec H3A 1B1, Canada;
E-mail: ridouan.bani@mail.mcgill.ca

T. Peller
Department of Biology, McGill University, 1205 Docteur Penfield, Montréal, Québec H3A 1B1, Canada;

J. Marleau
Department of Biology, McGill University, 1205 Docteur Penfield, Montréal, Québec H3A 1B1, Canada;

M.-J. Fortin
Department of Ecology & Evolutionary Biology, University of Toronto, 25 Willcocks Street, Toronto, Ontario M5S 3B2, Canada;

F. Guichard
Department of Biology, McGill University, 1205 Docteur Penfield, Montréal, Québec H3A 1B1, Canada;

sults show how within-MPA dynamics affect predictions based on parameters of MPA networks such as MPA size, spacing, and pelagic larval duration. Increasing withn-MPA spatial resolution predicted increasing population stability independently of other network properties. Our analysis also resolved the role of pelagic larval duration for scaling up within-MPA dynamics to MPA networks: species with short larval duration led to increasing network stability with MPA spacing while the opposite was observed for species with long larval duration. Our study stresses the importance of integrating fluctuating larval connectivity, both within and among MPAs, and more specifically suggest the benefit of small and nearby MPAs under increasing ocean variability.

Keywords Marine protected areas · spatiotemporal dispersal · metapopulation persistence · size & spacing

Introduction

Networks of marine protected area (MPA) are designed to allow for synergistic interactions among individual MPAs across spatial scales and for the protection of biological diversity, and of natural and associated cultural resources (IUCN 1994; WCPA/IUCN 2007). Effective MPAs network planning can allow populations to grow and persist within MPAs and contribute larval recruitment (Harrison et al. 2012; Baetscher et al. 2019) as well as spillover through adult movement (Kellner et al. 2008; Moffitt et al. 2009; Goñi et al. 2010; Kough et al. 2019) to unprotected areas. Metapopulation theories concerning MPA networks associate these principles with four proxies: location (e.g. representation, nurseries, special features), size (e.g. richness, adult movement, home range), spacing (e.g. connectivity), and/or number (e.g. replication). Effective MPA network planning is equivalent to understanding how location, size, spacing and number of MPAs contribute to species reproductive (Goodyear 1993) and dispersal (Moffitt et al. 2011; Green et al. 2015) abilities. Many studies concerned with principles for designing MPAs networks that address social, economic and biological criteria, but few have set these principles in the context of variability in environmental conditions and biological traits (McLeod et al. 2009; Magris et al. 2014). MPA network design needs to reflect the complexity of species' niches, especially the spatial configuration and environmental and biological spatiotemporal variability (Caswell et al. 1997; Caswell 2001; Lande 2002; Tuljapurkar and Caswell 2012; Tuljapurkar 2013).

Theoretical studies concerned with the effects of MPA size and spacing on species persistence show that MPA size and the fraction of coastline protection required for species persistence should be proportional to mean dispersal distance (Botsford et al. 2019). However, others have drawn attention to the impact dispersal heterogeneity can have on species persistence in MPAs (Kaplan 2006; Aiken et al. 2007; Kaplan et al. 2009; White et al. 2010; Aiken and Navarrete 2011). Recent studies have indicated the importance of larval dispersal patterns, especially the spatiotemporal variability that can arise from

hydrodynamic processes, climatic variability, and from developmental and behavioral changes (Munday et al. 2008; Siegel et al. 2008; Mitarai et al. 2009; Munday et al. 2009; Lett et al. 2010; Magris et al. 2014). Current MPA network theories typically ignore such variability by describing larval transport and development with dispersal kernels (Gaylord et al. 2006; Aiken et al. 2007; Woodson et al. 2012; Kerr et al. 2015; Le Corre et al. 2019). Spatiotemporal fluctuations and aggregation in recruitment events can be observed over a broad range of spatial scales, including scales that are smaller than the size of an MPA. This can lead to a network of larval connectivity within individual MPAs in addition to among-MPA connectivity.

Spatiotemporal variability in dispersal can produce inflating or deflating effects on growth and density (Watson et al. 2012; Williams and Hastings 2013) which may affect stability of metapopulation in MPA networks (Bani et al. 2019) and can be used in MPA network planning especially in extreme conditions when average larval recruitment is low. These inflating or deflating effects are purely a product of dispersal fluctuations and depend on the temporal pattern of larval exchange among populations, a feature that dispersal kernels fail to address when modeling dispersal dynamics. When MPA network connectivity is assumed static and larval recruitment fails to offset post-settlement mortality, the metapopulation defined by the MPA network will go extinct. Positive growth and stable dynamics can still be achieved when fluctuations in larval recruitment are autocorrelated (Williams and Hastings 2013) and spatially asynchronous (Watson et al. 2012; Bani et al. 2019). Recent theory applied to simulated biophysical connectivity (Bani et al. 2019), predicts the species density in MPA networks can undergo these inflating or deflating effects with changes in the configuration of MPA networks (spatial heterogeneity in larval recruitment, spacing, proportion of protected area). However, there is currently no theory of how these effects of spatiotemporal patterns of connectivity can scale up from within population dynamics to whole metapopulations. Such theory is needed to inform the optimal configuration of MPA networks under increasing climate and ocean variability.

Here, we adopt a metapopulation model (Bani et al. 2019) that captures the dynamics of a single marine species subject to density regulation and occupying MPAs interconnected by time varying larval dispersal. We predict the growth and stability of the metapopulation in a MPA network from the stochastic and stage-structured estimates of stationary distribution. We derive an explicit expression for metapopulation stability as a contribution of scale-dependent effect of within- and among- MPA fluctuations in larval recruitment. We then apply our framework to the Pacific northeast coastal system, where we vary the spacing, and proportion of protected area by either random or structured MPA assignment along the coast.

1 Metapopulation dynamics within MPA networks

We study populations dynamics of a single marine species characterized by an adult sedentary phase and a dispersive larval phase. The populations are distributed over an MPAs network noted $\mathcal{M}$ ($\mathcal{M} = \{U, V, ...\}$ is the network of MPAs, U, and V are two individual MPA within the network $\mathcal{M}$, and $|\mathcal{M}|$ is the number of MPAs). The population occupying the MPAs ($\{U, V, ...\}$) are interconnected by time-varying larval dispersal. We adopted a stochastic and density-dependent model that assumes densities are near carrying capacity. Our stochastic framework captures and predict species metapopulation growth and stability conditions when dispersal varies in time and space.

The population $N_U(t)$ within each MPA at time changes due to three processes: (1) adult individuals die with a per-capita rate m_a, (2) a time constant self-recruitment with a per-capita rate of $\bar{I}_{UU}$ and (3) time varying external-recruitment from other MPAs $\sum_{V \in \mathcal{M}; V \neq U} I_{VU}(t) N_V(t)$. We used a time averaged value for local recruitment (local-retention) because larvae that recruit directly back to their natal population are not exposed to ocean current variability compared to larvae that travel greater distances. We also assume that the intensity of each process varied with local population size to account for competition among conspecifics at both recruitment and adult stages. Competition can be for space or for other resources, and is implemented as a logistic function with carrying-capacity K_U specific to each MPA.

$$\frac{dN_U(t)}{dt} = (r_U N_U(t) + \sum_{V \in \mathcal{M}; V \neq U} I_{VU}(t) N_V(t))(1 - \frac{N_U(t)}{K_U}) \qquad (1)$$

The local growth is equal to the adult mortality rate subtracted from the self-recruitment rate ($r_U = \bar{I}_{UU} - m_a$). The time varying per capita immigration rate $I_{VU}(t) = \delta f C_{VU}(t)$ is a function of $C_{VU}(t)$: the probability or proportion of individuals dispersing from V to U at time t, f and δ are adult fecundity and larval survival rates respectively and are spatially uniform over the metapopulation. It is important to note that the loss of larvae to unprotected areas may be of importance for fisheries purposes, however, our model only tracks the dynamics of population' individuals within MPAs and we assume no survival of larvae ($\delta = 0$) outside MPAs.

We convert the density-dependent model in eq. (1) to a stochastic and multivariate Markovian process that converges to a stationary distribution when its deterministic counterpart reaches the stable equilibrium ($K_U, K_V, ...$). If we assume that all the MPAs have similar relative carrying capacity (i.e. $k_U = \frac{1}{|\mathcal{M}|}$), the mean and variance of the stationary distribution of regional relative abundance are as follows (Bani et al. 2019):

$$\mu_{\sum_U n_U} = \sum k_U = 1 \qquad (2)$$

$$\sigma^2_{\sum_U n_U} = \frac{m_a R_{\mathcal{M}}}{2|\mathcal{M}|^2} \sum_{U \in \mathcal{M}} \frac{\sum_{V \in \mathcal{M}; V \neq U} \sum_{W \in \mathcal{M}; W \neq U} cov(C_{VU}(t), C_{WU}(t))}{\sum_{V=1}^s \bar{C}_{VU} - \frac{1}{R_{\mathcal{M}}}} = \sum_{U \in \mathcal{M}} \sigma^2_{U \in \mathcal{M}}$$

(3)

1.1 Metapopulation stability

We can consider the variation in total relative abundance over the MPA network around carrying capacity as a measure of population stability defined as:

$$Stability = \frac{1}{CV} = \frac{\mu_{\sum n_U}}{\sigma_{\sum n_U}} = \sqrt{\frac{1}{\sum_{U \in \mathcal{M}} \sigma^2_{U \in \mathcal{M}}}}$$

(4)

The contribution of each MPA to MPAs network' metapopulation stability measured in terms of the variance contribution of each MPA to metapopulation variance, which is represented by the variance of the stationary distribution $\sigma^2_{\sum_U n_U}$ (eq. 3). For instance, the contribution of MPA U is:

$$\sigma^2_{U \in \mathcal{M}} = \frac{m_a R_{\mathcal{M}}}{2|\mathcal{M}|^2} \times \frac{\sum_{V \in \mathcal{M}; V \neq U} \sum_{W \in \mathcal{M}; W \neq U} cov(C_{VU}(t), C_{WU}(t))}{\sum_{V=1}^s \bar{C}_{VU} - \frac{1}{R_{\mathcal{M}}}}$$

(5)

Where $R_{\mathcal{M}}$ is the net reproductive rate of network $\mathcal{M}$ calculated such that the largest eigenvalue (λ_d) of the average projection matrix of the density-independent (a.k.a. average connectivity matrix - diagonal adult mortality matrix) version of model (eq. 1) to be equal to 1. The variance contribution (eq. 5) assumes spatially homogeneous larval recruitment within MPAs. We now develop an expression for the variance contribution of each MPA to MPAs network' stability for the case where within-MPA larval recruitment can be heterogeneous.

2 Spatial resolution of larval recruitment within MPAs

In addition to the temporal variability in immigration terms ($I_{VU}(t)$), we also consider the case of spatially heterogeneous larval recruitment within individual MPAs. In such case, we divide each MPA U into smaller patches (P_i) such that $U = \{P_1, ..., P_u\}$, and that within each patch P_i larval recruitment is assumed to be homogeneous. However, we assume that larval recruitment can differ among the patches within-MPA.

The connectivity between MPAs represent a spatially averaged connectivity between within MPAs patches P_i. For example, the connectivity from MPA $V = \{P_1, ..., P_v\}$ to $U = \{P_1, ..., P_u\}$ at time t is:

$$C_{VU}(t) = \sum_{P_j \in V} \frac{d_j}{\sum_{P_v \in V} d_v} \sum_{P_i \in U} C_{ji}(t)$$

(6)

The connectivity $C_{ji}(t)$ is the proportion of the total number of larvae d_j released from patch $P_j \in V$ arriving to patch $P_i \in U$.

The size of patches P_i corresponds to the level of larval recruitment spatial resolution within each MPA U: when MPA U has a spatially homogeneous larval recruitment then the size of patches P_i within which larval recruitment is assumed homogeneous should be larger than the MPA U size (*i.e.* there exists a patch P_u such that $U \subset P_u$), otherwise the size of patch P_i should be smaller than the MPA U size for the MPA U to have spatially heterogeneous larval recruitment (i.e. $\exists U \in \mathcal{M}$ such that $U = \bigcup_{i=1,\ldots,u} P_i$ and $u > 1$).

When patches P_i are smaller than the size of individual MPAs, larval connectivity can become nested within individual MPAs. The within-MPA spatial resolution in larval recruitment can affect how individual MPAs contribute to metapopulation growth and stability. In the following section, we derive an explicit relationship that partitions the contribution of MPA U to metapopulation stability in relation to the spatial resolution of larval recruitment within MPA U.

2.1 Partitioning MPA and within-MPA contributions to population stability

When larval recruitment is spatially heterogeneous such that patches P_i are smaller than the size of individual MPAs (*i.e.* $\exists U \in \mathcal{M}$ such that $U = \bigcup_{i=1,\ldots,u} P_i$ and $u > 1$), individual MPAs become themselves networks of their within patches P_i. Thus, individual MPAs are subnetworks within the MPA network and their variance contribution to the MPA network metapopulation stability depends on within-MPA dynamics. We can derive the nested MPA contribution to metapopulation stability across the whole MPA network starting from MPA U contribution expression with spatially homogeneous larval recruitment ($\sigma^2_{U \in \mathcal{M}}$ in eq. 5) and by integrating between-MPA connectivity as a function of within-MPA patches P_i connectivity (eq. 6) (see appendix B for details).

$$\sigma^2_{U \in \mathcal{M}} = \sum_{i \in U} \left[(\frac{|\Omega|}{|\mathcal{M}| \times |U|})^2 (\frac{R_{\mathcal{M}}}{R_{\Omega}})^2 \omega_{i \in \Omega} \sigma^2_{i \in \Omega} - (\frac{1}{|\mathcal{M}|})^2 (\frac{R_{\mathcal{M}}}{R_U})^2 \omega_{i \in U} \sigma^2_{i \in U} \right]$$

(7)

The population within patch P_i contributes

$$\sigma^2_{i \in \Omega} = \frac{m_a R_{\Omega}}{2|\Omega|^2} \times \frac{\sum_{q \in \Omega; q \neq i} \sum_{q \in \Omega; p \neq i} cov(C_{qi}(t), C_{pi}(t))}{\sum_{j \in \Omega} \bar{C}_{ji} - \frac{1}{R_{\Omega}}}$$

to whole ($\Omega = \bigcup_{U \in \mathcal{M}} \{P_i | P_i \in U\}$) metapopulation variance, and

$$\sigma^2_{i \in U} = \frac{m_a R_U}{2|U|^2} \times \frac{\sum_{k \in U; k \neq i} \sum_{l \in U; l \neq i} cov(C_{ki}(t), C_{li}(t))}{\sum_{j \in U} \bar{C}_{ji} - \frac{1}{R_U}}$$

to within MPA U ($\{P_i|P_i \in U\}$) metapopulations variance (Fig. 1). Both contributions are weighted by

$$\omega_{i \in \Omega} = \frac{\sum_{j \in \Omega} \bar{I}_{ji} - m_a}{\sum_{V \in \mathcal{M}} \bar{I}_{VU} - m_a}$$

and

$$\omega_{i \in U} = \frac{\sum_{j \in U} \bar{I}_{ji} - m_a}{\sum_{V \in \mathcal{M}} \bar{I}_{VU} - m_a}$$

which are patch P_i larval recruitment relative to adult mortality when recruitment is from the whole Ω or from within-MPA U metapopulations, respectively. $R_{\mathcal{M}}$, R_{Ω}, and R_U are the net reproductive rates of MPAs network $\mathcal{M}$ when larval recruitment is spatially homogeneous within MPAs, of whole metapopulation Ω across patches P_i (regardless of the MPA they belong to), and of patches within-MPA U.

It is important to note that while $\sigma^2_{i \in \Omega}$ measures variance contribution of each P_i population to stability of metapopulation within all MPAs (Ω), $\sigma^2_{i \in U}$ measures the contribution of the same population to stability of metapopulation within-MPA (U) only (Fig. 1). The right-hand side of eq. (7) is composed of two parts, the first part measures the weighted contribution of each patch to the metapopulation of all (Ω) regardless of what MPA they belong two and the second part measures its contribution to the metapopulation within the MPA it belongs to.

We verify that the relationship in eq. (7) holds for two boundary conditions when larval recruitment is homogenous within-MPA (i.e. $\{\forall U \in \mathcal{M}, \exists P_j | U \subset P_j\}$) as well as for the case of a single MPA. When larval recruitment is homogenous within-MPA, $\sigma^2_{i \in U} = 0$, $\frac{R_{\mathcal{M}}}{R_{\Omega}} = 1$, and $\omega_{i \in \Omega} = 1$ and MPA variance contribution in eq. (7) reduces to the case when larval recruitment is homogenous within-MPA in eq. (5). Moreover, in the case of a single MPA, $\sigma^2_{i \in \Omega} = \sigma^2_{i \in U}$, $\frac{R_{\mathcal{M}}}{R_{\Omega}} = \frac{R_{\mathcal{M}}}{R_U}$, and $\frac{|\Omega|}{|\mathcal{M}| \times |U|} = \frac{1}{|\mathcal{M}|} = 1$ such that $\sigma^2_{U \in \mathcal{M}} = 0$ regardless of the level of within-MPA spatial resolution. In this case, single MPA dynamics with homogeneous larval recruitment are governed by local growth rate ($r_U = \bar{I}_{UU} - m_a$) and since self-recruitment is optimized to balance adult mortality, the population grows deterministically to reach carrying (i.e. $\sigma^2_{U \in \mathcal{M}} = 0$).

By using the nested network contribution of MPAs to variance $\sigma^2_{U \in \mathcal{M}}$ given by eq. (7) in our measure of stability (eq. 4), the stability of a network of nested MPAs becomes:

$$Stability = \sqrt{\frac{1}{\sum_{U \in \mathcal{M}} \sum_{i \in U} \left[(\frac{|\Omega|}{|\mathcal{M}||U|})^2 (\frac{R_{\mathcal{M}}}{R_{\Omega}})^2 \omega_{i \in \Omega} \sigma^2_{i \in \Omega} - (\frac{1}{|\mathcal{M}|})^2 (\frac{R_{\mathcal{M}}}{R_U})^2 \omega_{i \in U} \sigma^2_{i \in U} \right]}}$$

(8)

Fig. 1 An illustration of two MPAs (top panel: $\mathcal{M} = \{U, V\}$)) with equal size when larval recruitment is well mixed within both MPAs and showing two connectivity pathways: a self-recruitment back to MPA U and external recruitment from MPA V. When larval recruitment is spatially heterogeneous, MPAs become sub-networks (middle panel) and the both connectivity arrows shown for well-mixed MPAs (top panel) are replaced with multiple connectivity arrows between patches. Spatial heterogeneity in larval recruitment can generate within-MPA dynamics (bottom panel).

3 Case study along northeast Pacific coasts

3.1 biophysical connectivity

We apply our model to a case study along the coast of British Columbia, Canada and Washington State, USA. To quantify the dynamic connectivity, a solution from a three-dimensional hydrodynamic model ROMS (Regional Ocean Modeling System) output is used to generate neutrally buoyant particle trajectories along the coast. The ROMS model uses 3 km horizontal resolution and a 31 vertical resolution levels. To hind-cast ocean currents over the 10-year period from 1998 to 2007, Masson and Fine (2012) forced the ROMS output with tides, 3-hourly winds from the North American Regional Reanalysis (NARR; Mesinger et al. (2006)), heat fluxes, freshwater runoff (Morrison et al. 2012), and lateral boundary salinities and temperature from Simple Ocean Data Assimilation (SODA; Carton and Giese (2008)).

444 circular release sites (5 km radius) with centers located on the coastline are distributed uniformly throughout the nearshore (Fig. 3-a). 133 million passive Lagrangian particles (LTRANS: North et al. (2011)) were released from the 444 patches for a period of 10 years between Jan 1, 1998 and Aug 26, 2007. The passive particles were released biweekly starting Jan 1 to Aug 26 every year and then tracked (Latitude, Longitude) weekly for a period

Fig. 2 The North-East Pacific coast (British Columbia, Canada and Washington, U.S.) and the distribution of (a) 10 km sites along the coastline, and aggregated patches of (b) 2-times, (c) 4-times, and (d) 8-times the size of circular patches. The patches: 1-time, 2-times, 4-times, and 8-times the size of the site with 10 km diameter are the four levels of spatial heterogeneity in larval recruitment within-MPA. We used equation (6) to calculate the connectivity between the patches using the connectivity between the sites.

of 18 weeks (120 days). We refer to the release day by spawning time and the duration spent after release as pelagic larval duration, so we can assume hypothetical species represented by pairs of spawning time and pelagic larval duration. In total we had 18 spawning times ($\{Jan\ 01,\ Jun\ 15,...,\ Aug\ 12,\ Aug\ 26\}$) and 18 pelagic larval durations ($\{1, 8, ..., 113, 120\}$) and a total of $18 \times 18 = 324$ hypothetical species. We assume that our hypothetical species reproduce during the one-day spawning time window every year and that all larvae reach competency to settle exactly after the pelagic larval duration. Connectivity among the 444 circular patches are probability of moving from a patch to another and are calculated as the proportion of particles that are

within the 5 km radius of each patches divided by the number of particles released from the source patch. The connectivity times series are calculated for each pair of spawning time, pelagic larval duration and year.

3.2 MPA network configuration: MPA size, MPA spacing and within-MPA connectivity

We studied the effects of within-MPA connectivity and of its spatial resolution on MPA persistence defined as the minimal net reproductive rate (minimum NRR) and stability ($\frac{1}{CV}$). We assumed that and individual MPA can be formed by aggregating neighboring sites of 5 km radius (Fig. 2a). We first assessed the effect of spatial resolution on local retention and minimal net reproductive rate for the case of single MPAs by aggregating contiguous sites into a single patch forming a single MPA of increasing size, up to 100% of the total 444 sites. MPAs of each size were replicated by varying the location of the starting site along the coastline.

We implemented MPA networks and controlled for MPA spacing by limiting the size of individual MPAs to an area equivalent to 8 contiguous sites (80 km), with a total of 49 MPAs (Fig. 2d). We varied the proportion of protected area (10%, 20%, 30%, and 40%) by randomly sampling (5, 10, 15, and 20) out of 49 potential MPAs. We further divided the sampled MPAs into 5 categories of MPA spacing (76 km, 163 km, 250 km, 337 km, and 425 km), resulting in the same number of MPA network replicates for all distance categories. The MPA spacing was calculated as an average distance between MPAs as follows ($X_U^T = (X_{U,lon}, X_{U,lat})$ is the center of MPA U):

$$spacing = \frac{2}{|\mathcal{M}|(|\mathcal{M}| - 1)} \sum_{U > V \in M} \sqrt{(X_U - X_V)^T \times (X_U - X_V)}, \quad (9)$$

For example, if we assume a network of 5 MPAs, the MPA spacing is calculated as follows:

$$\frac{1}{\frac{5 \times (5-1)}{2}} \times \begin{bmatrix} 1 & 1 & 1 & 1 & 1 \end{bmatrix} \times \begin{bmatrix} 0 & d_{12} & d_{13} & d_{14} & d_{15} \\ 0 & 0 & d_{23} & d_{24} & d_{25} \\ 0 & 0 & 0 & d_{34} & d_{35} \\ 0 & 0 & 0 & 0 & d_{45} \\ 0 & 0 & 0 & 0 & 0 \end{bmatrix} \times \begin{bmatrix} 1 \\ 1 \\ 1 \\ 1 \\ 1 \end{bmatrix}$$

Where $d_{UV} = \sqrt{(X_{U,lon} - X_{V,lon})^2 + (X_{U,lat} - X_{V,lat})^2}$ is the distance between MPA U and V. When applied to the case study, the MPA spacing range span between 32 km and 469 km and 76 km, 163 km, 250 km, 337 km, and 425 km are representative values of the 5 different distance groups.

To control for within-MPA spatial resolution (*i.e.* size of patches P_i), we aggregated sites within-MPAs from its highest level of spatial resolution corresponding to a single site (no aggregation) to patches of increasing size corresponding to 3 lower spatial resolution levels represented by patch length: 10

km (1-site in Fig. 2a), 20 km (2-site aggregation in Fig. 2b), 40 km (4-site aggregation in Fig. 2c), and 80 km (8-site aggregation in Fig. 2d). Within-MPA, larval recruitment is spatially-homogeneous when the spatial resolution is as coarse as MPA size. Within-MPA spatial resolution increases with decreasing patch length.

4 Results

4.1 Metapopulation stability in MPA network with within-MPA connectivity

When we consider within-MPA connectivity as a nested subnetwork, each MPA contribution to stability ($\frac{1}{CV}$) can be partitioned into its within-patch contribution to within- and between-MPA networks, both governed by dynamic connectivity within and between MPAs. Our results show how stability of overall MPA networks is affected by dynamic connectivity when within-MPA spatial resolution increased.

Within-MPA spatial resolution covaries with temporal variations in connectivity, and more specifically with its statistical moments including the mean, variance, covariance, and realized connections. Within-MPA spatial resolution decreased the mean and increased the variance of larval recruitment (Fig. 3), both changes resulted in reducing overall stability of MPA networks with increasing within-MPA spatial resolution when proportion of protected area is at 10% (Fig. 4). Moreover, since the mean and variance of connectivity decay exponentially with pelagic larval duration, we observed stronger response of short pelagic larval durations compared to longer pelagic larval duration (Fig. 3), and stability decreased minimally for longer compared to short pelagic larval duration (10% of protected area in Fig. 4).

The response of metapopulation stability to MPA spacing within MPA networks varied with within-MPA spatial resolution, proportion of protected area, and pelagic larval duration. When the proportion of protected area is 10%, MPA spacing increased (decreased) network stability for short (long) pelagic larval durations regardless of the level of within-MPA spatial resolution. Increasing the proportion of protected area from 10% to 40%, increased (decreased) stability at shorter MPA spacing for high (low) spatial resolution especially with long pelagic larval durations.

The decreasing metapopulation stability in MPA networks with increasing within-MPA spatial resolution results from the combination of between- (U; Fig.2) and within- (Ω; Fig.2) MPA responses. These responses can be captured by the within-MPA population contributions to both within- ($\sigma^2_{i \in U}$) and between-MPA ($\sigma^2_{i \in \Omega}$) networks (eq. 7).

Metapopulation stability in MPA networks decreased with increasing MPA contribution ($\sigma^2_{U \in \mathcal{M}}$), which itself increased with increasing difference between weighted variance contribution of populations to between- ($\omega_{i \in \Omega} \sigma^2_{i \in \Omega}$) and within- ($\omega_{i \in U} \sigma^2_{i \in U}$) MPA networks. Larval recruitment when recruitment is from MPAs ($\omega_{i \in \Omega}$) and within-MPA ($\omega_{i \in U}$) networks and variance contribu-

Fig. 3 Response of statistical components of dynamic connectivity to pelagic larval duration and spatial spatial resolution within MPAs: (a) mean, (b) variance, (c) covariance, and (d) percentage of realized connections. For each pelagic larval duration, points represent the mean over MPA replicates and spawning times.

tion to between-MPA ($\sigma^2_{i \in \Omega}$) varied little compared to how variance contribution to within-MPA ($\sigma^2_{i \in U}$) varied with within-MPA spatial resolution when the proportion of protected area is low (10% in Figs. B1 and B2). Within-MPA dynamics is what reduces MPA network stability: the higher the within-MPA spatial resolution, the lower the variance contribution of populations to within-compare to between-MPA networks (Fig. B1). The decrease of population contribution to the within-MPA variance translated into a stable within-MPA sub-networks and unstable between-MPA networks with increasing within-MPA spatial resolution.

5 Discussion

We studied the impact of connectivity within and between MPAs on metapopulation persistence in MPA network in relation to population growth from low density and to population stability at carrying capacity. Theories of MPA network design based on metapopulation dynamics typically focus on the study

Fig. 4 Mean MPA Network stability ($1/CV$) as a function of MPA spacing and within-MPA spatial resolution (dot size) for different pelagic larval durations (panel columns) and proportions of protected area (panel rows). Solid lines show best quadratic fit and error bars represent standard deviation over MPA replicates and spawning times

of population growth at low densities, mostly ignoring density-dependent processes (Burgess et al. 2014). However, the establishment of MPA networks not only aim to enable populations at low densities to recover, but also to promote the long-term stability of populations at higher abundances (Carr et al. 2017). When populations are near carrying capacity, their dynamics can be density-dependent (e.g. Hixon et al. 2012). Density-dependent dynamics involve mortality due intraspecific competition on both adults and new recruits, and both are affected by fluctuating larval dispersal/connectivity within and between MPAs. In general, our findings show connectivity dynamics within MPAs can promote the persistence of populations at both low and high densities. Specifically, our findings predict a lower minimum net reproductive effort

(NRR) is required for growth of a populations at low densities and more stable dynamics of populations near carrying capacity.

In addition to the common assumption of low-density dynamics, most current MPA theories rely on temporally and spatially averaged dispersal (Burgess et al. 2014). Our work extends this current framework to account for spatially and temporally fluctuating connectivity interacting with density-dependent (meta)population growth across life stages. Differential responses to spatiotemporal dispersal of low and high density populations found in our study highlight the roles played by multiple statistical components of dynamic connectivity and their interaction with ecological processes such as intraspecific competition. At low densities, average larval recruitment is the most important predictor of population persistence. However, at high abundance, the temporal average, variance and covariance of larval recruitment interact to govern the stability of fluctuations around carrying capacity. This suggests that variance and covariance in recruitment can impact the long-term maintenance of (meta)populations in MPA networks. Recruitment is widely recognized as being spatially and temporally variable (Caffey 1985; Cowen and Sponaugle 2009) and our work suggests that the explicit consideration of this variability is important for understanding population dynamics.

Our study support results from other studies that incorporate spatiotemporal larval dispersal in MPA planning with the goal of designing networks that can be robust to environmental variations and climate change (e.g. Carr et al. 2017; Coleman et al. 2017). Similar to recent studies, we show the importance of pelagic larval traits, in particular pelagic larval duration (PLD), for predicting temporal patterns of dynamic connectivity (Bani et al. 2019). By explicitly considering larval dispersal within and between MPAs, we reveal how PLD interacts with the spatial structure and resolution of connectivity networks. For instance, larval recruitment for species with short PLD show strong spatial variation compared to species with long PLD (Fig. **??**). This mapping between the pelagic traits and spatiotemporal properties of larval dispersal could predict the multiple and interacting effects of climate change on physical ocean currents and pelagic traits (Bani et al. in review).

MPAs can self-persist when local-retention is high enough to compensate for both larval post-settlement and adult mortality or persist as a network when at least one MPA is self-persistent and well connected to all other MPAs (Botsford et al. 2001; Armsworth 2002). Metapopulation persistence in MPA networks has commonly been formulated as a problem of MPA spacing and distance (Botsford et al. 2001; Kaplan et al. 2006; Moffitt et al. 2009). Studies have suggested that MPA size should be increased and spacing decreased in response to environmental variability, including variability that directly affect larval dispersal (Airamé et al. 2003; Shanks et al. 2003; Salm et al. 2006; Green et al. 2007; McLeod et al. 2009). Increasing size and decreasing spacing are both expected to increase the average larval recruitment regardless of environmental variability. However, these predictions are based on the assumption that larval recruitment is proportional to MPA size. This is clearly not the case in our results (Fig. 3c), which show that local retention in MPAs follows

a bimodal - first decreasing then increasing - relationship with MPA size, especially for short larval duration (Fig. **??**). It is more likely that increasing MPA size will decrease local retention for species with short PLD. The chances of recruiting back to a natal population decreases with MPA size within the MPA network, and MPA self-persistence thus decreases with MPA size.

We found that population growth and stability in MPA networks exposed to fluctuating connectivity are strongly dependent on within-MPA connectivity. Local retention under the assumption of well-mixed recruitment results from spatially-averaged self- and external larval recruitment within each MPA. By considering within-MPA dynamics, it is possible for an individual population within an MPA to persist when the spatially-averaged recruitment is not enough to compensate for post-settlement and adult mortality (Fig. **??**b). In that context, the minimal NRR can reveal the interaction between within-MPA connectivity and MPA size for recovery when at low abundance: if all MPAs are self-persistent independently of distance, the minimal NRR should remain the same regardless of MPA distance. Otherwise, MPAs are dependent on external recruitment and within-MPA connectivity for network persistence. Despite the strong focus on connectivity in MPA theory, connectivity within MPAs has rarely been discussed. Yet, marine populations are commonly patchily distributed across relatively small spatial scales (Sale et al. 2006) and empirical evidence has shown that recruits into one population can come from nearby patches (Planes et al. 2009; Berumen et al. 2012). Our findings suggest that larval connectivity within MPAs can promote population persistence and stability.

Across our range of PLD values, we found that networks with a few large MPAs have greater self-recruitment than external recruitment, but networks with many small MPAs have greater external than self recruitment (Fig. 4d). Minimal NRR required for persistence in our smallest MPAs within networks set at a small proportion of protected area (¡10%) is smaller than in large MPAs. These same small MPAs are more self-persistent and export more larvae, thus subsidizing non-self-persistent MPAs and contributing to MPA network persistence. These results suggest that external recruitment from small MPAs allows reaching low minimum NRR with less overall proportion of protected area compared to large MPAs that rely on self-recruitment. When connectivity within MPAs is considered, small and adjacent sites within MPAs reduce minimal NRR variations among species with different PLD (Fig. **??**b) compared to large MPAs with well-mixed dispersal that are sensitive to PLD (Fig. **??**).

When population abundance is near carrying capacity, large MPAs are predicted to benefit the stability of MPAs networks when within-MPA connectivity is ignored. On the other hand, within-MPA aggregation of larval recruitment spatially decouples the dynamics of an individual MPA from the network, which destabilizes the MPA network. However, this increase in network-level variability resulting from within-MPA dynamics is alleviated by reducing both MPA size and spacing. For example, with 20% of the coastline protected, MPA network stability is robust to changes in within-MPA spatial resolution when

MPAs are small and close to each other, but network stability becomes sensitive to within-MPA spatial resolution with increasing MPA spacing (Fig. 4).

MPA theory has long highlighted a perceived tradeoff between MPA spacing and size for achieving fisheries and conservation objectives (Hastings and Botsford 2003; Gaines et al. 2010): conservation objectives are predicted to be best achieved with large MPAs that are self-sustaining while fisheries objectives are best served by small MPAs that maximize spillover effects and larval export (e.g. Botsford et al. 2009). This tradeoff has led to the suggested implementation of MPA networks with moderately sized and spaced MPAs (Gaines et al. 2010). Our cross-scale analysis incorporating within-MPA connectivity predicts that many small and nearby MPAs can maintain high population abundance and greater network-level stability, and thus achieve conservation goals for sedentary species with dispersive larval stages, while simultaneously providing spillover effects and larval export to achieve fisheries goals. Our study suggests spatiotemporal fluctuations in connectivity can impact MPA network design through the consideration of within-MPA connectivity. The importance of such a cross-scale theory of MPA networks is likely to grow with ongoing climate change and associated increase in environmental variability.

Acknowledgments

References

IUCN. Guidelines for protected area management categories, 1994.

WCPA/IUCN. Establishing networks of marine protected areas: A guide for developing national and regional capacity for building mpa networks. 2007.

Hugo B Harrison, David H Williamson, Richard D Evans, Glenn R Almany, Simon R Thorrold, Garry R Russ, Kevin A Feldheim, Lynne Van Herwerden, Serge Planes, Maya Srinivasan, et al. Larval export from marine reserves and the recruitment benefit for fish and fisheries. *Current biology*, 22(11): 1023–1028, 2012.

Diana S Baetscher, Eric C Anderson, Elizabeth A Gilbert-Horvath, Daniel P Malone, Emily T Saarman, Mark H Carr, and John Carlos Garza. Dispersal of a nearshore marine fish connects marine reserves and adjacent fished areas along an open coast. *Molecular ecology*, 28(7):1611–1623, 2019.

Julie B Kellner, Roger M Nisbet, and Steven D Gaines. Spillover from marine reserves related to mechanisms of population regulation. *Theoretical Ecology*, 1(2):117–127, 2008.

Elizabeth A Moffitt, Louis W Botsford, David M Kaplan, and Michael R O'Farrell. Marine reserve networks for species that move within a home range. *Ecological applications*, 19(7):1835–1847, 2009.

Raquel Goñi, Ray Hilborn, David Díaz, Sandra Mallol, and Sara Adlerstein. Net contribution of spillover from a marine reserve to fishery catches. *Marine Ecology Progress Series*, 400:233–243, 2010.

Andrew S Kough, Carolyn A Belak, Claire B Paris, Agnessa Lundy, Heather Cronin, Gaya Gnanalingam, Sam Hagedorn, Rachel Skubel, Amanda C Weiler, and Allan W Stoner. Ecological spillover from a marine protected area replenishes an over-exploited population across an island chain. *Conservation Science and Practice*, 1(3):e17, 2019.

C Phillip Goodyear. Spawning stock biomass per recruit in fisheries management: foundation and current use. *Canadian Special Publication of Fisheries and Aquatic Sciences.*, 120:67–81, 1993.

Elizabeth A Moffitt, J Wilson White, and Louis W Botsford. The utility and limitations of size and spacing guidelines for designing marine protected area (MPA) networks. *Biological Conservation*, 144(1):306–318, 2011.

Alison L Green, Aileen P Maypa, Glenn R Almany, Kevin L Rhodes, Rebecca Weeks, Rene A Abesamis, Mary G Gleason, Peter J Mumby, and Alan T White. Larval dispersal and movement patterns of coral reef fishes, and implications for marine reserve network design. *Biological Reviews*, 90(4): 1215–1247, 2015.

Elizabeth McLeod, Rodney Salm, Alison Green, and Jeanine Almany. Designing marine protected area networks to address the impacts of climate change. *Frontiers in Ecology and the Environment*, 7(7):362–370, 2009.

Rafael A Magris, Robert L Pressey, Rebecca Weeks, and Natalie C Ban. Integrating connectivity and climate change into marine conservation planning. *Biological Conservation*, 170:207–221, 2014.

Hal Caswell, Roger M Nisbet, André M de Roos, and Shripad Tuljapurkar. Structured-population models: many methods, a few basic concepts. In *Structured-population models in marine, terrestrial, and freshwater systems*, pages 3–17. Springer, 1997.

H Caswell. Construction, analysis, and interpretation. *Sunderland: Sinauer*, 585, 2001.

Russell Lande. Incorporating stochasticity in population viability analysis. *Population viability analysis. University of Chicago Press, Chicago*, pages 18–40, 2002.

Shripad Tuljapurkar and Hal Caswell. *Structured-population models in marine, terrestrial, and freshwater systems*, volume 18. Springer Science & Business Media, 2012.

Shripad Tuljapurkar. *Population dynamics in variable environments*, volume 85. Springer Science & Business Media, 2013.

Louis W Botsford, J Wilson White, and Alan Hastings. *Population Dynamics for Conservation*. Oxford University Press, 2019.

David M Kaplan. Alongshore advection and marine reserves: consequences for modeling and management. *Marine Ecology Progress Series*, 309:11–24, 2006.

Christopher M Aiken, Sergio A Navarrete, Manuel I Castillo, and Juan Carlos Castilla. Along-shore larval dispersal kernels in a numerical ocean model of the central Chilean coast. *Marine Ecology Progress Series*, 339:13–24, 2007.

David M Kaplan, Louis W Botsford, Michael R O'Farrell, Steven D Gaines, and Salvador Jorgensen. Model-based assessment of persistence in proposed marine protected area designs. *Ecological Applications*, 19(2):433–448, 2009.

J Wilson White, Louis W Botsford, Alan Hastings, and John L Largier. Population persistence in marine reserve networks: incorporating spatial heterogeneities in larval dispersal. *Marine Ecology Progress Series*, 398:49–67, 2010.

Christopher M Aiken and Sergio A Navarrete. Environmental fluctuations and asymmetrical dispersal: generalized stability theory for studying metapopulation persistence and marine protected areas. *Marine Ecology Progress Series*, 428:77–88, 2011.

Philip L Munday, Geoffrey P Jones, Morgan S Pratchett, and Ashley J Williams. Climate change and the future for coral reef fishes. *Fish and Fisheries*, 9(3):261–285, 2008.

DA Siegel, S Mitarai, CJ Costello, SD Gaines, BE Kendall, RR Warner, and KB Winters. The stochastic nature of larval connectivity among nearshore marine populations. *Proceedings of the National Academy of Sciences*, 105 (26):8974–8979, 2008.

S Mitarai, DA Siegel, JR Watson, C Dong, and JC McWilliams. Quantifying connectivity in the coastal ocean with application to the southern california bight. *Journal of Geophysical Research: Oceans (1978–2012)*, 114(C10), 2009.

PL Munday, JM Leis, JM Lough, CB Paris, MJ Kingsford, ML Berumen, and Jonathan Lambrechts. Climate change and coral reef connectivity. *Coral*

Reefs, 28(2):379–395, 2009.

Christophe Lett, Sakina-Dorothée Ayata, Martin Huret, and Jean-Olivier Irisson. Biophysical modelling to investigate the effects of climate change on marine population dispersal and connectivity. *Progress in Oceanography*, 87 (1):106–113, 2010.

Brian Gaylord, Daniel C Reed, Peter T Raimondi, and Libe Washburn. Macroalgal spore dispersal in coastal environments: mechanistic insights revealed by theory and experiment. *Ecological Monographs*, 76(4):481–502, 2006.

CB Woodson, MA McManus, JA Tyburczy, JA Barth, L Washburn, JE Caselle, MH Carr, DP Malone, PT Raimondi, BA Menge, et al. Coastal fronts set recruitment and connectivity patterns across multiple taxa. *Limnology and Oceanography*, 57(2):582–596, 2012.

Kecia A Kerr, Anabell Cornejo, Frédéric Guichard, Augusto C Crespi Abril, and Rachel Collin. Planktonic predation risk: effects of diel state, season and prey life history stage. *Journal of Plankton Research*, 37(2):452–461, 2015.

Nicolas Le Corre, Pierre Pepin, Guoqi Han, Zhimin Ma, and Paul VR Snelgrove. Assessing connectivity patterns among management units of the Newfoundland and Labrador shrimp population. *Fisheries Oceanography*, 28(2):183–202, 2019.

James R Watson, Bruce E Kendall, David A Siegel, and Satoshi Mitarai. Changing seascapes, stochastic connectivity, and marine metapopulation dynamics. *The American Naturalist*, 180(1):99–112, 2012.

Paul David Williams and Alan Hastings. Stochastic dispersal and population persistence in marine organisms. *The American Naturalist*, 182(2):271–282, 2013.

Ridouan Bani, Marie-Josée Fortin, Rémi M Daigle, and Frédéric Guichard. Dispersal traits interact with dynamic connectivity to affect metapopulation growth and stability. *Theoretical Ecology*, 12(1):111–127, 2019.

Diane Masson and Isaac Fine. Modeling seasonal to interannual ocean variability of coastal British Columbia. *Journal of Geophysical Research: Oceans (1978–2012)*, 117(C10), 2012.

Fedor Mesinger, Geoff DiMego, Eugenia Kalnay, Kenneth Mitchell, Perry C Shafran, Wesley Ebisuzaki, Dušan Jović, Jack Woollen, Eric Rogers, Ernesto H Berbery, et al. North american regional reanalysis. *Bulletin of the American Meteorological Society*, 87(3):343–360, 2006.

J Morrison, MGG Foreman, and D Masson. A method for estimating monthly freshwater discharge affecting British Columbia coastal waters. *Atmosphere-Ocean*, 50(1):1–8, 2012.

James A Carton and Benjamin S Giese. A reanalysis of ocean climate using Simple Ocean Data Assimilation (SODA). *Monthly Weather Review*, 136 (8):2999–3017, 2008.

Elizabeth W North, E Eric Adams, Zachary Schlag, Christopher R Sherwood, Ruoying He, Kyung Hoon Hyun, and Scott A Socolofsky. Simulating oil droplet dispersal from the deepwater horizon spill with a Lagrangian ap-

proach. *Geophysical Monograph Series*, 195:217–226, 2011.

Scott C Burgess, Kerry J Nickols, Chris D Griesemer, Lewis AK Barnett, Allison G Dedrick, Erin V Satterthwaite, Lauren Yamane, Steven G Morgan, J Wilson White, and Louis W Botsford. Beyond connectivity: how empirical methods can quantify population persistence to improve marine protected-area design. *Ecological Applications*, 24(2):257–270, 2014.

Mark H Carr, Sarah P Robinson, Charles Wahle, Gary Davis, Stephen Kroll, Samantha Murray, Ervin Joe Schumacker, and Margaret Williams. The central importance of ecological spatial connectivity to effective coastal marine protected areas and to meeting the challenges of climate change in the marine environment. *Aquatic Conservation: Marine and Freshwater Ecosystems*, 27:6–29, 2017.

Mark A Hixon, Todd W Anderson, Kevin L Buch, Darren W Johnson, J Brock McLeod, and Christopher D Stallings. Density dependence and population regulation in marine fish: a large-scale, long-term field manipulation. *Ecological Monographs*, 82(4):467–489, 2012.

Hugh M Caffey. Spatial and temporal variation in settlement and recruitment of intertidal barnacles. *Ecological Monographs*, 55(3):313–332, 1985.

Robert K Cowen and Su Sponaugle. Larval dispersal and marine population connectivity. *Annual review of marine science*, 1:443–466, 2009.

Melinda A Coleman, Paulina Cetina-Heredia, Moninya Roughan, Ming Feng, Erik van Sebille, and Brendan P Kelaher. Anticipating changes to future connectivity within a network of marine protected areas. *Global Change Biology*, 23(9):3533–3542, 2017.

Louis W Botsford, Alan Hastings, and Steven D Gaines. Dependence of sustainability on the configuration of marine reserves and larval dispersal distance. *Ecology Letters*, 4(2):144–150, 2001.

Paul R Armsworth. Recruitment limitation, population regulation, and larval connectivity in reef fish metapopulations. *Ecology*, 83(4):1092–1104, 2002.

David M Kaplan, Louis W Botsford, and Salvador Jorgensen. Dispersal per recruit: an efficient method for assessing sustainability in marine reserve networks. *Ecological Applications*, 16(6):2248–2263, 2006.

Satie Airamé, Jenifer E Dugan, Kevin D Lafferty, Heather Leslie, Deborah A McArdle, and Robert R Warner. Applying ecological criteria to marine reserve design: a case study from the California Channel Islands. *Ecological applications*, 13(sp1):170–184, 2003.

Alan L Shanks, Brian A Grantham, and Mark H Carr. Propagule dispersal distance and the size and spacing of marine reserves. *Ecological applications*, 13(sp1):159–169, 2003.

Rodney V Salm, Terry Done, and Elizabeth McLeod. Marine protected area planning in a changing climate. *Coral reefs and climate change: science and management. American Geophysical Union, Washington, DC*, pages 207–221, 2006.

Alison Green, Paul Lokani, Stuart Sheppard, Jeanine Almany, Stephen Keu, Joseph Aitsi, J Warku Karvon, Richard Hamilton, and Geoff Lipsett-Moore. Scientific design of a resilient network of marine protected areas, kimbe bay,

west new britain, papua new guinea. *TNC Pacific Island Countries Report*, 2(07), 2007.

Peter F Sale, Ilkka Hanski, and Jacob P Kritzer. The merging of metapopulation theory and marine ecology: establishing the historical context. In *Marine metapopulations*, pages 3–28. Elsevier, 2006.

Serge Planes, Geoffrey P Jones, and Simon R Thorrold. Larval dispersal connects fish populations in a network of marine protected areas. *Proceedings of the National Academy of Sciences*, 106(14):5693–5697, 2009.

Michael L Berumen, Glenn R Almany, Serge Planes, Geoffrey P Jones, Pablo Saenz-Agudelo, and Simon R Thorrold. Persistence of self-recruitment and patterns of larval connectivity in a marine protected area network. *Ecology and evolution*, 2(2):444–452, 2012.

Alan Hastings and Louis W Botsford. Comparing designs of marine reserves for fisheries and for biodiversity. *Ecological Applications*, 13(sp1):65–70, 2003.

Steven D Gaines, Crow White, Mark H Carr, and Stephen R Palumbi. Designing marine reserve networks for both conservation and fisheries management. *Proceedings of the National Academy of Sciences*, 107(43):18286–18293, 2010.

Louis W Botsford, Daniel R Brumbaugh, Churchill Grimes, Julie B Kellner, John Largier, Michael R O'Farrell, Stephen Ralston, Elaine Soulanille, and Vidar Wespestad. Connectivity, sustainability, and yield: bridging the gap between conventional fisheries management and marine protected areas. *Reviews in Fish Biology and Fisheries*, 19(1):69–95, 2009.

Appendix A: Partitioning MPAs variance contribution into within-MPA patches variance contribution

A marine protected area contributes ($\sigma^2_{U \in \mathcal{M}}$) with the expression as follow:

$$\sigma^2_{U \in \mathcal{M}} = \frac{m_a R_\mathcal{M}}{2|\mathcal{M}|^2} \times \frac{\sum_{V \in \mathcal{M}; U \neq U} \sum_{W \in \mathcal{M}; W \neq U} Cov(C_{VU}(t), C_{WU}(t))}{\sum_{V \in \mathcal{M}} \overline{C_{VU}} - \frac{1}{R_\mathcal{M}}}$$

If we replace the connectivity between the MPAs ($C_{VU}(t)$, $C_{WU}(t)$, and $\overline{C_{VU}}$) by their expression as a function of their within patches (Eq. 6):

$$\sigma^2_{U \in \mathcal{M}} = \frac{m_a R_\mathcal{M}}{2|\mathcal{M}|^2} \times \frac{\sum_{V \in \mathcal{M}; U \neq U} \sum_{W \in \mathcal{M}; W \neq U} Cov(\sum_{j \in V} (\frac{d_j}{\sum_{v \in V} d_v}) \sum_{i \in U} C_{ji}(t), \sum_{k \in W} (\frac{d_k}{\sum_{w \in W} d_W}) \sum_{l \in U} C_{kl}(t))}{\sum_{V \in \mathcal{M}} \sum_{j \in V} (\frac{d_j}{\sum_{v \in V} d_v}) \sum_{i \in U} \overline{C_{ji}}(t) - \frac{1}{R_\mathcal{M}}}$$

$$= \frac{m_a R_\mathcal{M}}{2|\mathcal{M}|^2} \times \frac{\sum_{V \in \mathcal{M}; U \neq U} \sum_{j \in V} (\frac{d_j}{\sum_{v \in V} d_v}) \sum_{W \in \mathcal{M}; W \neq U} \sum_{k \in W} \frac{d_k}{\sum_{w \in W} d_W} \sum_{i, l \in U} Cov(C_{ji}(t), C_{kl}(t))}{\sum_{V \in \mathcal{M}} \sum_{j \in V} (\frac{d_j}{\sum_{v \in V} d_v}) \sum_{i \in U} \overline{C_{ji}}(t) - \frac{1}{R_\mathcal{M}}}$$

$$= \frac{m_a R_\mathcal{M}}{2|\mathcal{M}|^2} \times [\frac{\sum_{V \in \mathcal{M}} \sum_{j \in V} (\frac{d_j}{\sum_{v \in V} d_v}) \sum_{W \in \mathcal{M}} \sum_{k \in W} \frac{d_k}{\sum_{w \in W} d_W} \sum_{i \in U} Cov(C_{ji}(t), C_{ki}(t))}{\sum_{V \in \mathcal{M}} \sum_{j \in V} (\frac{d_j}{\sum_{v \in V} d_v}) \sum_{i \in U} \overline{C_{ji}}(t) - \frac{1}{R_\mathcal{M}}} -$$

$$\frac{\sum_{k, l \in U} \sum_{i \in U} (\frac{d_j}{\sum_{u \in U} d_u}) \times (\frac{d_j}{\sum_{u \in U} d_u}) Cov(C_{ji}(t), C_{ki}(t))}{\sum_{V \in \mathcal{M}} \sum_{j \in V} (\frac{d_j}{\sum_{v \in V} d_v}) \sum_{i \in U} \overline{C_{ji}}(t) - \frac{1}{R_\mathcal{M}}}]$$

For simplicity and clarity, we assume that the patches do have similar size with equal net larval reproduction $d_i = d_i$, then $\frac{d_j}{\sum_{v \in V} d_v} = \frac{1}{|V|}$:

$$\sigma^2_{U \in \mathcal{M}} = \frac{m_a R_\mathcal{M}}{2|\mathcal{M}|^2} \times [\frac{\sum_{V \in \mathcal{M}} \sum_{j \in V} (\frac{1}{|V|}) \sum_{W \in \mathcal{M}} \sum_{k \in W} \frac{1}{|W|} \sum_{i \in U} Cov(C_{ji}(t), C_{ki}(t))}{\sum_{V \in \mathcal{M}} \sum_{j \in V} (\frac{1}{|V|}) \sum_{i \in U} \overline{C_{ji}}(t) - \frac{1}{R_\mathcal{M}}} -$$

$$\frac{\sum_{k, l \in U} \sum_{i \in U} (\frac{1}{|U|}) \times (\frac{1}{|U|}) Cov(C_{ji}(t), C_{ki}(t))}{\sum_{V \in \mathcal{M}} \sum_{j \in V} (\frac{1}{|V|}) \sum_{i \in U} \overline{C_{ji}}(t) - \frac{1}{R_\mathcal{M}}}]$$

$$= \frac{m_a R_\mathcal{M}}{2|\mathcal{M}|^2} \times [\sum_{i \in U} \frac{\sum_{V \in \mathcal{M}} \sum_{j \in V} (\frac{1}{|V|}) \sum_{W \in \mathcal{M}} \sum_{k \in W} \frac{1}{|W|} Cov(C_{ji}(t), C_{ki}(t))}{\sum_{V \in \mathcal{M}} \sum_{j \in V} (\frac{1}{|V|}) \sum_{i \in U} \overline{C_{ji}}(t) - \frac{1}{R_\mathcal{M}}} -$$

$$\sum_{i \in U} \frac{\sum_{k, l \in U} (\frac{1}{|U|})(\frac{1}{|U|}) Cov(C_{ji}(t), C_{ki}(t))}{\sum_{V \in \mathcal{M}} \sum_{j \in V} (\frac{1}{|V|}) \sum_{i \in U} \overline{C_{ji}}(t) - \frac{1}{R_\mathcal{M}}}]$$

$$\sigma^2_{U\in\mathcal{M}} = \frac{m_a R_\mathcal{M}}{2|\mathcal{M}|^2} \times [\sum_{i\in U} \frac{\sum_{V\in\mathcal{M}}\sum_{j\in V}(\frac{1}{|V|})\sum_{W\in\mathcal{M}}\sum_{k\in W}\frac{1}{|W|}Cov(C_{ji}(t),C_{ki}(t))}{\sum_{V\in\mathcal{M}}\sum_{j\in V}(\frac{1}{|V|})\sum_{i\in U}\overline{C_{ji}}(t) - \frac{1}{R_\mathcal{M}}} -$$

$$\sum_{i\in U} \frac{\sum_{k,l\in U}(\frac{1}{|U|})(\frac{1}{|U|})Cov(C_{ji}(t),C_{ki}(t))}{\sum_{V\in\mathcal{M}}\sum_{j\in V}(\frac{1}{|V|})\sum_{i\in U}\overline{C_{ji}}(t) - \frac{1}{R_\mathcal{M}}}]$$

$$= \frac{m_a R_\mathcal{M}}{2|\mathcal{M}|^2} \times \sum_{i\in U}[\frac{\sum_{V\in\mathcal{M}}\sum_{j\in V}(\frac{1}{|V|})\sum_{W\in\mathcal{M}}\sum_{k\in W}\frac{1}{|W|}Cov(C_{ji}(t),C_{ki}(t))}{\sum_{V\in\mathcal{M}}\sum_{j\in V}(\frac{1}{|V|})\sum_{i\in U}\overline{C_{ji}}(t) - \frac{1}{R_\mathcal{M}}} -$$

$$\frac{\sum_{k,l\in U}(\frac{1}{|U|})(\frac{1}{|U|})Cov(C_{ji}(t),C_{ki}(t))}{\sum_{V\in\mathcal{M}}\sum_{j\in V}(\frac{1}{|V|})\sum_{i\in U}\overline{C_{ji}}(t) - \frac{1}{R_\mathcal{M}}}]$$

$$= \frac{m_a R_\mathcal{M}}{2|\mathcal{M}|^2} \times \sum_{i\in U}[\frac{\sum_{j\in\Omega}\overline{C_{ji}} - \frac{1}{R_\Omega}}{\sum_{V\in\mathcal{M}}\sum_{j\in V}(\frac{1}{|V|})\sum_{i\in U}\overline{C_{ji}}(t) - \frac{1}{R_\mathcal{M}}} \times$$

$$\frac{\sum_{V\in\mathcal{M}}\sum_{j\in V}(\frac{1}{|V|})\sum_{W\in\mathcal{M}}\sum_{k\in W}\frac{1}{|W|}Cov(C_{ji}(t),C_{ki}(t))}{\sum_{j\in\Omega}\overline{C_{ji}} - \frac{1}{R_\Omega}} -$$

$$\frac{\sum_{j\in U}\overline{C_{ji}} - \frac{1}{R_U}}{\sum_{V\in\mathcal{M}}\sum_{j\in V}(\frac{1}{|V|})\sum_{i\in U}\overline{C_{ji}}(t) - \frac{1}{R_\mathcal{M}}} \times \frac{\sum_{k,l\in U}(\frac{1}{|U|})(\frac{1}{|U|})Cov(C_{ji}(t),C_{ki}(t))}{\sum_{j\in U}\overline{C_{ji}} - \frac{1}{R_U}}]$$

If we assume that MPAs nestedness is similar such that $\{|U| = |V|, \forall U, V \in \mathcal{M}\}$, we can further simpify the expression into:

$$\sigma^2_{U\in\mathcal{M}} = \frac{m_a R_\mathcal{M}}{2|\mathcal{M}|^2} \times \sum_{i\in U}[\frac{\sum_{j\in\Omega}\overline{C_{ji}} - \frac{1}{R_\Omega}}{\sum_{V\in\mathcal{M}}\frac{1}{|V|}\sum_{j\in V;i\in U}\overline{C_{ji}}(t) - \frac{1}{R_\mathcal{M}}} \times$$

$$\frac{1}{|U|^2} \times \frac{\sum_{j\in\Omega,j\neq i}\sum_{k\in\Omega,k\neq i}Cov(C_{ji}(t),C_{ki}(t))}{\sum_{j\in\Omega}\overline{C_{ji}} - \frac{1}{R_\Omega}} -$$

$$\frac{\sum_{j\in U}\overline{C_{ji}} - \frac{1}{R_U}}{\sum_{V\in\mathcal{M}}\sum_{j\in V}(\frac{1}{|V|})\sum_{i\in U}\overline{C_{ji}}(t) - \frac{1}{R_\mathcal{M}}} \times \frac{1}{|U|^2} \times \frac{\sum_{k,l\in U}Cov(C_{ji}(t),C_{ki}(t))}{\sum_{j\in U}\overline{C_{ji}} - \frac{1}{R_U}}]$$

$$
= \sum_{i \in U} [\frac{m_a R_{\mathcal{M}}}{2|\mathcal{M}|^2} \times \frac{\sum_{j \in \Omega} \overline{C_{ji}} - \frac{1}{R_\Omega}}{\sum_{V \in \mathcal{M}} \frac{1}{|V|} \sum_{j \in V; i \in U} \overline{C_{ji}}(t) - \frac{1}{R_{\mathcal{M}}}} \times
$$

$$
\frac{1}{|U|^2} \times \frac{\sum_{j \in \Omega, j \neq i} \sum_{k \in \Omega, k \neq i} Cov(C_{ji}(t), C_{ki}(t))}{\sum_{j \in \Omega} \overline{C_{ji}} - \frac{1}{R_\Omega}} -
$$

$$
\frac{m_a R_{\mathcal{M}}}{2|\mathcal{M}|^2} \times \frac{\sum_{j \in U} \overline{C_{ji}} - \frac{1}{R_U}}{\frac{1}{|U|} \times \sum_{V \in \mathcal{M}} \sum_{j \in V; i \in U} \overline{C_{ji}}(t) - \frac{1}{R_{\mathcal{M}}}} \times
$$

$$
\frac{1}{|U|^2} \times \frac{\sum_{k,l \in U} Cov(C_{ji}(t), C_{ki}(t))}{\sum_{j \in U} \overline{C_{ji}} - \frac{1}{R_U}}]
$$

$$
\sigma^2_{U \in \mathcal{M}} = \sum_{i \in U} [\frac{|\Omega|^2 R_{\mathcal{M}}}{|\mathcal{M}|^2 R_\Omega} \times \frac{\sum_{j \in \Omega} \overline{C_{ji}} - \frac{1}{R_\Omega}}{\sum_{V \in \mathcal{M}} \frac{1}{|V|} \sum_{j \in V; i \in U} \overline{C_{ji}}(t) - \frac{1}{R_{\mathcal{M}}}} \times
$$

$$
\frac{1}{|U|^2} \times \frac{m_a R_\Omega}{2|\Omega|^2} \times \frac{\sum_{j \in \Omega, j \neq i} \sum_{k \in \Omega, k \neq i} Cov(C_{ji}(t), C_{ki}(t))}{\sum_{j \in \Omega} \overline{C_{ji}} - \frac{1}{R_\Omega}} -
$$

$$
\frac{R_{\mathcal{M}}}{|\mathcal{M}|^2 R_U} \times \frac{\sum_{j \in U} \overline{C_{ji}} - \frac{1}{R_U}}{\sum_{V \in \mathcal{M}} \frac{1}{|V|} \sum_{j \in V; i \in U} \overline{C_{ji}}(t) - \frac{1}{R_{\mathcal{M}}}} \times
$$

$$
\frac{m_a R_U}{2|U|^2} \times \frac{\sum_{k,l \in U} Cov(C_{ji}(t), C_{ki}(t))}{\sum_{j \in U} \overline{C_{ji}} - \frac{1}{R_U}}]
$$

$$
= \sum_{i \in U} [\frac{|\Omega|^2}{|\mathcal{M}|^2 |U|^2} \times \frac{R_{\mathcal{M}}}{R_\Omega} \times \frac{\frac{1}{R_\Omega}}{\frac{1}{R_{\mathcal{M}}}} \times (\frac{\sum_{j \in \Omega} \delta f \overline{C_{ji}} - m_a}{\sum_{V \in \mathcal{M}} \frac{1}{|V|} \sum_{j \in V; i \in U} \delta f \overline{C_{ji}}(t) - m_a}) \times
$$

$$
(\frac{m_a R_\Omega}{2|\Omega|^2} \times \frac{\sum_{j \in \Omega, j \neq i} \sum_{k \in \Omega, k \neq i} Cov(C_{ji}(t), C_{ki}(t))}{\sum_{j \in \Omega} \overline{C_{ji}} - \frac{1}{R_\Omega}}) -
$$

$$
\frac{1}{|\mathcal{M}|^2} \times \frac{R_{\mathcal{M}}}{R_U} \times \frac{\frac{1}{R_U}}{\frac{1}{R_{\mathcal{M}}}} \times (\frac{\sum_{j \in U} \delta f \overline{C_{ji}} - m_a}{\sum_{V \in \mathcal{M}} \frac{1}{|V|} \sum_{j \in V; i \in U} \delta f \overline{C_{ji}}(t) - m_a}) \times
$$

$$
(\frac{m_a R_U}{2|U|^2} \times \frac{\sum_{k,l \in U} Cov(C_{ji}(t), C_{ki}(t))}{\sum_{j \in U} \overline{C_{ji}} - \frac{1}{R_U}})]
$$

Then:

$$\sigma^2_{U\in\mathcal{M}} = \sum_{i\in U}[(\frac{|\Omega|}{|\mathcal{M}||U|})^2 \times (\frac{R_\mathcal{M}}{R_\Omega})^2 \times (\frac{\sum_{j\in\Omega}\overline{I_{ji}} - m_a}{\sum_{V\in\mathcal{M}}\frac{1}{|V|}\sum_{j\in V;i\in U}\overline{I_{ji}}(t) - m_a})\times$$

$$(\frac{m_a R_\Omega}{2|\Omega|^2} \times \frac{\sum_{j\in\Omega,j\neq i}\sum_{k\in\Omega,k\neq i}Cov(C_{ji}(t),C_{ki}(t))}{\sum_{j\in\Omega}\overline{C_{ji}} - \frac{1}{R_\Omega}})-$$

$$\frac{1}{|\mathcal{M}|^2} \times (\frac{R_\mathcal{M}}{R_U})^2 \times (\frac{\sum_{j\in U}\overline{I_{ji}} - m_a}{\sum_{V\in\mathcal{M}}\frac{1}{|V|}\sum_{j\in V;i\in U}\overline{I_{ji}}(t) - m_a})\times$$

$$(\frac{m_a R_U}{2|U|^2} \times \frac{\sum_{k,l\in U}Cov(C_{ji}(t),C_{ki}(t))}{\sum_{j\in U}\overline{C_{ji}} - \frac{1}{R_U}})]$$

Therefore, if we let:

$$\sigma^2_{i\in\Omega} = \frac{m_a R_\Omega}{2|\mathcal{M}|^2} \times \frac{\sum_{j\in\Omega,j\neq i}\sum_{k\in\Omega,k\neq i}Cov(C_{ji}(t),C_{ki}(t))}{\sum_{j\in\Omega}\overline{C_{ji}} - \frac{1}{R_\Omega}}$$

$$\sigma^2_{i\in\Omega} = \frac{m_a R_U}{2|U|^2} \times \frac{\sum_{k,l\in U}Cov(C_{ji}(t),C_{ki}(t))}{\sum_{j\in U}\overline{C_{ji}} - \frac{1}{R_U}})$$

and:

$$\omega_{i\in\Omega} = \frac{\sum_{j\in\Omega}\overline{I_{ji}} - m_a}{\sum_{V\in\mathcal{M}}\frac{1}{|V|}\sum_{j\in V;i\in U}\overline{I_{ji}}(t) - m_a}$$

$$\omega_{i\in U} = \frac{\sum_{j\in U}\overline{I_{ji}} - m_a}{\sum_{V\in\mathcal{M}}\frac{1}{|V|}\sum_{j\in V;i\in U}\overline{I_{ji}}(t) - m_a}$$

Thus, we can write the contribution in variance of MPA U to network stability as:

$$\sigma^2_{U\in\mathcal{M}} = \sum_{i\in U}[(\frac{|\Omega|}{|\mathcal{M}||U|})^2 \times (\frac{R_\mathcal{M}}{R_\Omega})^2 \times \omega_{i\in\Omega} \times \sigma^2_{i\in\Omega} - \frac{1}{|\mathcal{M}|^2} \times (\frac{R_\mathcal{M}}{R_U})^2 \times \omega_{i\in U} \times \sigma^2_{i\in U}]$$

Fig. B1 The variance contribution of patches within-MPA to whole ($\sigma^2_{i \in \Omega}$) and within-MPA ($\sigma^2_{i \in U}$) networks stability and how they vary with MPA spacing and spatial heterogeneity within-MPA (dots size: tiny 10 km, small 20 km, medium 40 km, and large 80 km) for different pelagic larval durations (columns: short 8 days, intermediate 64 days, and long 120 days) and proportion of protected areas (rows: 10%, 20%, 30%, and 40%). The lines are best quadratic fit, and points and error bars represent averages and standard deviations of different MPA replicates and spawning times. The within-MPA patch contribution to overall network ($\sigma^2_{i \in \Omega}$) varied little with spatial heterogeneity within-MPA compared their variance contribution to within-MPA ($\sigma^2_{i \in U}$) network stability regardless of MPA spacing especially at low proportion of protected areas (10%) and short pelagic larval duration.

Appendix B: Supplementary Figures

Fig. B2 Relative larval recruitment when it is from whole ($\omega_{i \in \Omega}$) and within-MPA ($\omega_{i \in U}$) networks (proportion in y-axis) and how they vary with MPA spacing (x-axis) and spatial heterogeneity within-MPA (dots size: tiny 10 km, small 20 km, medium 40 km, and large 80 km) for different pelagic larval durations (columns: short 8 days, intermediate 64 days, and long 120 days) and proportion of protected areas (rows: 10%, 20%, 30%, and 40%). The lines are best quadratic fit, and points and error bars represent averages and standard deviations of different MPAs replicates and spawning times, respectively. Larval recruitment from whole network varied little with spatial heterogeneity within-MPA compared larval recruitment from within-MPA network regardless of MPA spacing especially at low proportion of protected areas (10%) and short pelagic larval duration.