TITLE

Commensal Microbiota Regulates Skin Barrier Function And Repair Via Signaling

Through The Aryl Hydrocarbon Receptor

AUTHORS

Aayushi Uberoi[1], Casey Bartow-McKenney[1], Qi Zheng[1], Laurice Flowers[1], Amy Campbell[1],

Simon A.B. Knight[1], Neal Chan[1], Monica Wei[1], Victoria Lovins[1], Julia Bugayev[1], Joseph

Horwinski[1], Charles Bradley[1], Jason Meyer[2], Debra Crumrine[2], Carrie Hayes Sutter[3], Peter

Elias[2], Elizabeth Mauldin[1], Thomas R. Sutter[3]*, Elizabeth A. Grice[1]*

AFFILIATIONS

[1]University of Pennsylvania, Perelman School of Medicine, Department of Dermatology,

Philadelphia, Pennsylvania, USA.

[2]San Francisco Veterans Affairs Medical Center, Dermatology Service and University of

California San Francisco; Department of Dermatology, San Francisco, California

[3] Department of Biological Sciences; W. Harry Feinstone Center for Genomic Research,

University of Memphis, Memphis, Tennessee, USA.

CONTACT INFORMATION

*Address for correspondence:

egrice@pennmedicine.upenn.edu

tsutter@memphis.edu

SUMMARY

The epidermis forms a barrier that defends the body from desiccation and entry of harmful substances, while sensing and integrating environmental signals. The tightly orchestrated cellular changes required for the proper formation and maintenance of this epidermal barrier occur in the context of the skin microbiome. Using germ free mice, we demonstrate the microbiota is necessary for proper differentiation and repair of the epidermal barrier. These effects were mediated by the aryl hydrocarbon receptor (AHR) in keratinocytes, a xenobiotic receptor also implicated in epidermal differentiation. Murine skin lacking keratinocyte AHR was more susceptible to barrier damage and infection, during steady state and epicutaneous sensitization. Colonization with a defined consortium of human skin isolates restored barrier competence in an AHR-dependent manner. We reveal a fundamental mechanism whereby the microbiota regulates skin barrier formation and repair, with far-reaching implications for the numerous skin disorders characterized by epidermal barrier dysfunction.

KEYWORDS

Skin; barrier; microbiome; epidermis; aryl hydrocarbon receptor; keratinocyte; microbiota; commensal

MAIN TEXT

INTRODUCTION

The skin is the primary barrier between the human body and the environment and functions to prevent desiccation and entry of foreign and/or harmful substances. The barrier properties of the skin reside in the epidermis, a semi-permeable stratified epithelium that is formed as a result of keratinocyte terminal differentiation. Though continuously exposed to xenobiotic toxins, physical insults, and pathogenic microbes, the epidermis is also associated with diverse commensal microbial communities that are critical players in regulating skin physiology [1]. These microbial communities, collectively referred to as skin microbiota, are specialized to thrive in the unique nutrient and environmental conditions of this organ. The skin microbiome is topographically diverse, temporally complex, and distinct from other organs [2, 3]. How the commensal microbiota influences development of skin's barrier function is undefined, as are the molecular mechanisms that mediate these interactions.

The barrier function of the skin may be conceptualized as four intertwined "levels" consisting of microbial, immune, chemical, and physical barriers [4]. The skin microbiome itself provides a barrier to pathogenic micro-organisms via a variety of different mechanisms e.g. production of proteases, antimicrobial peptides and antibiotics, and interference with quorum sensing [5]. This outermost microbial barrier also interacts with and mediates other functional levels of the cutaneous barrier. Skin microbiota play a fundamental role in the induction, training, and function of the skin immune barrier in part through the release of antimicrobial peptides, short-chain fatty acids, and polyamines [6]. Neonatal colonization by microbiota has long-lasting impacts on adult immune barrier as commensal skin microbes' prime immune cells to differentiate between commensal versus pathogenic bacterium [7]. Bacterial lipases can hydrolyze lipids resulting in production of free fatty acids that impact the acidic surface pH of the skin, which dictates the chemical barrier of the skin [8, 9]. While studies in gnotobiotic mice suggest that epidermal differentiation and barrier genes are microbially regulated [10],

mechanistic roles for the skin microbiota in development, regeneration, and function of the

physical barrier are not well defined.

The epidermal permeability barrier (EPB) comprises of the stratum corneum and a

complex system of tight junctions and adhesion complexes and their associated cytoskeletal

networks that mediate cell-cell adhesion to create a mechanical barrier between the

environment and underlying tissue [11]. Actively dividing keratinocytes in the stratum basale

commit to terminal differentiation and move progressively into suprabasal layers, i.e. stratum

spinosum, the stratum granulosum and eventually the stratum corneum [12]. In the stratum

corneum, keratinocytes become flattened and denucleated (which are then called corneocytes),

and plasma membranes are replaced with cornified envelopes. Lamellar bodies secrete their

lipid-rich contents into the intercellular space between the corneocytes and are subsequently

processed into barrier-providing lipid lamellae. Altogether, corneocytes, lipids and a complex

network of trans-membrane proteins, provide a highly hydrophobic EPB against the

environment. Microbial influences on this process of epidermal differentiation and EPB

formation are not well understood, nor are the mechanisms whereby the EPB senses and

responds to changes in the microbiota.

The sensing of xenobiotics, or compounds foreign to a living organism, is critical for

barrier defense and homeostasis in the skin [13]. Keratinocytes function as sentinels that sense

and respond to external stimuli [14]. Activation of xenobiotic receptors in keratinocytes induces

expression of detoxification enzymes and membrane transporters that promote elimination of

toxic compounds[15]. Accumulating evidence suggests that roles for xenobiotic receptors

extend to cellular processes beyond xenobiotic metabolism that include cellular proliferation,

tissue repair, and immune responses [13]. Microbes produce a plethora of small molecules and

secondary metabolites, which are hypothesized to mediate their interactions with the host

toward a mutualistic relationship [16]. The diversity of molecular signals produced by skin

microbes, and how keratinocytes decipher and respond to them, remain largely unexplored.

The aryl hydrocarbon receptor (AHR) is a xenobiotic receptor that has emerged as a

critical player in EPB development, function and integrity. Activation of the AHR, a ligand-

activated transcription factor of the basic, helix-loop-helix motif-containing Per-ARNT-Sim family

[17], induces a variety of epidermal differentiation and barrier genes, accelerates terminal

differentiation, and increases stratum corneum thickness [18-22]. The AHR can be activated by

halogenated and non-halogenated aromatic hydrocarbons, including dioxins such as 2,3,7,8-

Tetrachlorodibenzo-p-dioxin (TCDD), polychlorinated biphenyls (PCBs) and polycyclic aromatic

hydrocarbons (PAHs); clinically used drugs, food-derived molecules, endobiotics, and bacterial

metabolites [23-26]. Microbial regulation of the AHR in the context of the skin barrier remains

poorly understood, as well as the consequences of perturbing the commensal microbiota with

respect to EPB function and defense.

In addition to congenital barrier deficiencies, epidermal barrier dysfunction is a hallmark

of inflammatory skin diseases, including atopic dermatitis and psoriasis, and predisposes skin to

infections [27-29]. Additionally, epicutaneous sensitization, as a result of epidermal barrier

dysfunction, may lead to atopic and allergic disease [30].Thus, there is an urgent scientific and

clinical need to define the mechanistic basis by which the commensal microbiota regulate

homeostatic barrier function, as such mechanisms provide new targets for prevention and/or

intervention in skin barrier deficiencies.

Here, we investigated the role of commensal microbiota in regulation of permeability

barrier homeostasis of skin. We found that commensal microbes are necessary for normal

epidermal differentiation, EPB function, and repair. These effects were mediated by microbial

signaling through the keratinocyte AHR. Murine skin lacking keratinocyte AHR signaling

displayed increased barrier permeability, enhanced susceptibility to infection by *S. aureus,* and

increased pathology in a model of atopic dermatitis. We show that topical colonization with a

defined consortium of human skin commensals improves EPB function in murine germ-free skin

and models of barrier dysfunction. Our findings reveal a fundamental role for the commensal

skin microbiota in regulating the physical integrity and repair of the skin barrier, provides

mechanistic insights into microbial-skin crosstalk, and uncovers therapeutic targets for

improving skin barrier function.

RESULTS

Epithelial development and differentiation programs are impaired in germ free skin

To characterize microbially-mediated regulation of homeostatic epithelial gene

expression programs, we performed RNA-seq on epidermal sheets isolated from dorsal skin of

C57BL/6 mice of 3 different colonization states (n=8 mice each. **Figure 1A**): Specific pathogen

free (SPF) mice that were conventionally raised in presence of microbiota, germ free (GF) mice

born and raised in sterile gnotobiotic isolators, and a third group of mice that were born GF and

then colonized (COL) with SPF microbiota for 2 weeks. Within 2 weeks, COL mice were

colonized with microbiota from SPF mice as validated by 16S ribosomal RNA (rRNA) gene

sequencing (**Figure 1B**). We identified differentially expressed genes (DEGs) between

colonization states by training negative-binomial linear models using DESeq2 R package and 3-

way comparisons. This analysis revealed 6396, 427, and 3232 DEGs for SPF vs. GF, COL vs.

GF and SPF vs. COL comparisons, respectively (**Figure 1C, Table S1**). We focused on the 396

shared DEGs of SPF and COL epidermis when compared to GF epidermis (**Figure 1D**). We

reasoned that this subset of DEGs meet the criteria of being induced and sustained by microbial

colonization, suggesting homeostatic control. The 396 DEGs were significantly enriched for

biological functions such as skin development, keratinocyte differentiation, and epidermis

development (**Figure 1E**). This result suggests the microbiota plays an important role in

epithelial barrier formation.

To further examine DEGs involved in epithelial barrier function, we manually curated

genes involved in different facets of epithelial barrier: keratinization, cornified envelope

formation, adherence and gap junction, basement membrane function, barrier development,

differentiation and intercellular lipid-lamellae processing (**Table S2)**. Focusing on the SPF vs.

GF subset of DEGs, multiple genes across each of these categories were expressed at lower

levels in GF mice (**Figure 1F**, **Tables S3, S4**). In particular, genes critical for cornified envelope

formation [e.g. *involucrin (Ivl)*, *envoplakin (Evpl)*] and its desquamation, [e.g. *Kallikrien-related*

peptidases 5, 7(Klk5, 7)] were downregulated in GF skin. We hypothesized that such significant

and widespread differences in gene expression would result in structural differences between

GF and SPF skin. However, consistent with prior reports [10, 31] we did not notice any overt

differences between the epithelial organization of GF vs SPF mice by traditional

histopathological examination (**Figure S1**). Analysis of skin ultrastructure by electron

microscopy showed that overall, SPF mice had a greater number of individual layers within the

stratum corneum than GF mice (**Figure S1**). Immunofluorescence-based analysis of molecular

biomarkers showed decreased expression of loricrin and cytokeratin-10 in skin of GF mice

(**Figure 1G**) that are implicated in barrier integrity [32-34]. Among genes that were

downregulated in GF skin were tight and adherens junctions family members (**Figure 1F**) such

as *tight junction protein 3 (Tjp3), desmogleins 1a-b (Dsg1),* and *claudin-1 (Cldn1)* that are

critical players involved in skin barrier formation (reviewed in [35]). Remarkably, tight junction

integrity appeared compromised in the suprabasal epithelium as evident by downregulated and

diffused expression of *Dsg1* in GF mouse epidermis (**Figure 1G)**. Disruption of *Dsg1* is

associated with improper formation of desmosomes in suprabasal epithelia and has been

associated with skin barrier impairment [36-38]. Together, these findings suggest the

hypothesis that skin barrier formation requires the commensal microbiota.

Commensal microbiota promotes skin barrier function and repair

A fully functioning stratum corneum closely controls the water concentration gradient in

the skin such that passive diffusion of water occurs from inner layers towards the outside.

Barrier disruption compromises the ability of the stratum corneum to maintain this water

concentration gradient and results in increased transepidermal water loss (TEWL), measured

using a sensor for water vapor flow density [39, 40]. Low TEWL values are indicative of intact

skin and increased TEWL is associated with a disrupted barrier (**Figure 2A, Panel i**). Under

basal conditions, GF mice had slightly increased TEWL compared to SPF mice corroborating

the findings from ultrastructure analysis (**Figure S2**). Skin barrier of GF mice was perturbed

more readily with tape stripping than SPF skin (**Figure S2**), consistent with fewer layers of

stratum corneum in GF mice. After comparable insults following tape-strip injury (TEWL 15-20

g/m^2/h), SPF mice more rapidly repaired their barrier compared to GF mice when measured

over a period of 24 hours (**Figure 2B**). We observed similar delays in barrier recovery in GF

Rag1$^{-/-}$ mice (that lack mature T and B-cells) compared to age-matched SPF *Rag1$^{-/-}$* mice

(**Figure 2C, Figure S2**) suggesting that microbially-regulated adaptive immune responses are

not responsible for the delayed barrier repair phenotype. Skin from GF *Rag1$^{-/-}$* mice also

showed decreased expression of genes involved in terminal differentiation and formation of

transmembrane junctions compared to SPF *Rag1$^{-/-}$* mice (**Figure S2)** as seen in wild-type mice.

182   Basal keratinocytes undergo a spatiotemporal and highly controlled differentiation

program dependent on intracellular calcium flux to establish and maintain barrier [41]. In

presence of high calcium, primary epidermal keratinocytes can differentiate *in vitro* to express

genes involved in formation of cornified envelope [42]. We derived murine primary epidermal

keratinocytes from GF and SPF skin, respectively, and exposed them to high calcium containing

medium (**Figure 2D**). Expression of genes involved in terminal differentiation [*cytokeratin-10*

(Krt10), *involucrin (Ivl)*] and formation of transmembrane junctions [*Corneodesmosin (Cdsn),*

Desmocollin-1(Dsc1), Desmoglein-1a (Dsg1a)] were reduced in GF keratinocytes compared to

SPF keratinocytes (**Figure 2E, Figure S2**). Additionally, GF keratinocytes had decreased

transepithelial electrical resistance (TEER) which is indicative of decreased transmembrane

junction strength (**Figure 2F, Figure S2**).

193   To further examine the implications of perturbing the microbiota on skin barrier function

we used an antibiotic depletion model (**Figure 2G**). Prior studies had shown that antibiotics

traditionally used to disrupt gut microbiota in mice were not sufficient to disrupt skin microbiota

in mice [43]. We developed a new regimen consisting of antibiotics (Metronidazole,

Sulfamethoxazole, Trimethoprim, Cephalexin and Enrofloxacin) that are administered orally in

hospitals and veterinary clinics to target skin bacteria [44-47] and were able to inhibit prominent

murine skin commensal *Staphylococcus xylosus* [48, 49]. Oral administration of antibiotics for

two weeks diminished microbial burden on skin as observed by both quantitative cultures and

16S rRNA gene sequencing (**Figure 2H, I**) but did not significantly affect microbial burden in the

gut (**Figure S3).** Antibiotic-treated mice were delayed in barrier repair compared to control mice

that were treated with vehicle (**Figure 2J**). Together, these data confirm a role for commensal

microbes in promoting skin barrier function and repair.

Aryl hydrocarbon receptor pathway is attenuated in germ free skin

The sensing of external physiological and chemical signals is critical for barrier defense

and homeostasis in the skin [13]. Therefore, we hypothesized that xenobiotic receptors that act

as epithelial sensors and relay microbial signals would be among those genetic pathways

dysregulated in GF epidermis. Previous studies have identified at least 304 xenobiotic

processing genes (XPGs) in mice, which encode the enzymes, transporters, and transcription

factors required to metabolize xenobiotics [50]. We found that 52/304 XPGs were differentially

expressed (*P<0.*05) in skin of SPF and GF mice, and the majority were upregulated in SPF

mice (n=43/52; **Table S5**). The pregnane X receptor (PXR*, NR1I2*), constitutive androstane

receptor (CAR*, NR1I3*), peroxisome proliferator-activated receptor-alpha (PPARa) and aryl

hydrocarbon receptor (AHR) are key transcription factors that regulate xenobiotic processing in

skin [51]. Of these, only the AHR gene was differentially expressed and was upregulated in SPF

skin compared to GF skin. Canonically, after ligand binding, the AHR translocates to the cell

nucleus and binds DNA at xenobiotic responsive elements (XRE), to regulate transcription of

target genes [52]. Expression of key downstream target genes, i.e. cytochrome-p450 *Cyp1a1*

and molecular chaperones *Hsp90aa1* and *Hsp90ab1* that respond to AHR activation, were also

downregulated in GF murine epidermis (**Figure 3A**). Primary keratinocytes derived from GF skin

were also impaired in expression of these genes (**Figure S2**). Overall, microbiota-mediated

upregulation of *AHR* was consistent with changes in its canonical pathway, suggesting that

regulation of xenobiotic processing genes in the skin may be mediated through the *AHR*.

Since the AHR can activate multiple signaling pathways [53], we explored the impact of

microbial colonization on expression of genes reported to be a part of canonical and non-

canonical AHR signaling networks (**Figure 3A**). In this context, we saw upregulation of several

genes in SPF mice, including those involved in cell proliferation, differentiation, and

inflammation (e.g. *Hes1, Jun* and *Tgfb1).* Together these findings suggest the AHR as a

potential mechanism by which skin microbes modulate epithelial barrier integrity.

Treatment with an AHR agonist improves barrier function and recovery in germ free mice

We hypothesized that if the GF skin barrier phenotype we observed was due to

attenuated AHR signaling, then activation via AHR ligand would improve barrier recovery. We

treated adult GF mice topically with AHR ligand 6-Formylindolo[3,2-b] carbazole (FICZ), daily for

2 weeks at a low dose (100µg/kg) (**Figure 3B**), a regimen shown to induce expression of

Cyp1a1 [54]. FICZ is a tryptophan photoproduct and is a well-characterized AHR ligand in skin

[55, 56]. At the end of treatment, we compared the rate of barrier recovery in tape-stripped GF

mice that were either treated with FICZ or vehicle (**Figure 3C**). We observed that FICZ

significantly accelerated barrier recovery in GF mice. In parallel, we also treated SPF mice with

FICZ at the same dose and observed that FICZ accelerated early stage barrier recovery

(**Figure3D**). FICZ treatment activated AHR signaling in the treated group as *CYP1A1*

expression was induced in all FICZ treated mice in comparison to the untreated group (**Figure**

3E). Additionally, FICZ treatment increased expression of genes implicated in barrier repair

(**Figure 3E**) in GF mice. AHR upregulation increases expression of genes involved in epidermal

differentiation and formation of gap junctions [57]. In primary human keratinocytes, FICZ

increased epithelial resistance as measured by TEER (**Figure 3F, Figure S4**), which was

diminished by co-treatment with an AHR inhibitor (**Figure 3G**). Overall, activation of the AHR in

GF skin improves epidermal barrier recovery, supporting a mechanistic role for the commensal

microbiota in homeostatic regulation of the AHR.

Mice deficient in epithelial AHR have a defective skin barrier and are more susceptible to

infection

AHR is expressed in a variety of cell types in the skin, but abrogating AHR function in

keratinocytes has been suggested to impact skin barrier [58]. To confirm the role of keratinocyte

AHR in skin barrier function, we generated mice where the *floxed Ahr* allele ($Ahr^{f/f}$) was

conditionally knocked out in the skin epithelia using a *Cre* driven by the keratin-14 promoter

($K14^{Cre/+}Ahr^{f/f}$). While littermates that retained AHR function ($Ahr^{f/f}$) repaired their barrier within

24 hours following tape-strip disruption, barrier repair was significantly diminished in

$K14^{Cre/+}Ahr^{f/f}$ skin (**Figure 4A**). Additionally, keratinocytes derived from $K14^{Cre/+}Ahr^{f/f}$ mice

showed increased TEER compared to $Ahr^{f/f}$ keratinocytes (**Figure 4B**).

Patients with diseases of skin barrier impairment such as atopic dermatitis (AD) are

highly susceptible to colonization and infection by pathogens including *S. aureus* [59]. To test if

AHR-dependent barrier impairment leads to increased infection, we topically applied *S. aureus*

to tape-stripped skin [60] and quantified bacteria following 48 hours. We observed enhanced

infection of *S. aureus* and increased overall bacterial burden on tape stripped skin (**Figure 4C**)

of $K14^{Cre/+}Ahr^{f/f}$ mice compared to AHR sufficient controls (**Figure 4D**). Since barrier dysfunction

is a hallmark in the development of AD [61], and mice lacking AHR are impaired in barrier

repair, we hypothesized that these mice will be more prone to barrier damage, infection, and

atopic disease. We adapted a mouse model of AD (**Figure 4E**) induced by repeated

epicutaneous sensitization of tape-stripped skin with ovalbumin (OVA) [30]. In this model,

$K14^{Cre/+}Ahr^{f/f}$ skin was exacerbated in disease pathology, with enhanced TEWL (**Figure 4F**) and

increased susceptibility to *S. aureus* infection compared to $Ahr^{f/f}$ mice (**Figure 4G**). Together,

these data demonstrate that AHR function in keratinocytes is essential for barrier function, and

loss of AHR can lead to enhanced barrier damage in the setting of inflammatory skin disease

and facilitate bacterial entry.

Human skin microbial consortium restores barrier repair and function via the AHR

In the gastrointestinal tract, the AHR is activated by the gut microbiota [62] to enhance

intestinal barrier integrity by inducing tight junction proteins in intestinal epithelial cells [63].

However, the relation between skin commensals and cutaneous AHR-dependent barrier

regulation is unexplored. Given our observations that the commensal microbiota directly impacts

barrier repair (**Figure 1, 2**) and regulates the AHR genetic pathway (**Figure 3**) in murine skin,

we hypothesized that topical association of GF skin with human skin commensals would

activate the AHR and restore skin barrier repair. To test this hypothesis, we first curated a

collection of cultured skin microbes that were abundant and prevalent on healthy human skin.

Referred to as Flowers' Flora, the collection consists of members of Firmicutes phylum i.e.

Staphylococcus epidermidis, S. warneri, S. hemolyticus and members of Actinobacteria phylum

i.e. *Micrococcus luteus, Corynebacterium aurimucosum* (**Figure 5A**)*.* These skin microbes

activated AHR in keratinocytes as determined by way of a reporter assay consisting of the AHR

reporter element conjugated with *Cyp1a1* (**Figure 5B**). Flowers' Flora colonized murine GF skin

as determined by bacterial culture swabs and species-specific qPCR analysis (**Figure 5C, D**).

After two weeks of colonizing with this defined consortium of human skin commensals (**Figure**

5E) barrier recovery function in GF skin was restored (**Figure 5F**). Skin of colonized mice

showed elevated expression of differentiation genes as well as *Cyp1a1* (**Figure 5G**) compared

to GF skin. Upon terminal differentiation, keratinocytes derived from colonized mice were

enhanced in TEER compared to GF controls (**Figure 5H**). Colonization of $K14^{Cre/+}Ahr^{f/f}$ mice

with Flowers' Flora did not improve barrier repair (**Figure 5I**), further supporting an AHR-

dependent mechanism. Finally, to determine if Flowers' Flora could prevent barrier damage in

AD-like disease, we pre-colonized wild-type SPF murine skin prior to short-term epicutaneous

sensitization with OVA (**Figure 5J**). Pre-colonization with Flowers' Flora significantly improved

barrier recovery compared to skin that was not pre-colonized (**Figure 5K**). Together, these

findings indicate that skin commensal microbes signal through the AHR to maintain homeostatic

control of epidermal barrier integrity, and suggest new targets for preventing and/or treating

epidermal barrier dysfunction.

DISCUSSION

The skin microbiome provides the first level of barrier defense to the human body. While

commensal skin microbes have demonstrated effects on immune and chemical barriers of the

skin, their regulation of the physical barrier is not well-defined. Proper epithelial differentiation

and cornification is essential for formation of the epithelial barrier [11, 64, 65]. By using

comparative transcriptomics of gnotobiotic mice, we identified genes involved in skin

development, differentiation and barrier function as top candidates for microbial regulation.

Using multiple models of microbial depletion, loss of commensal microbiota impaired the barrier

repair function of the skin. We mechanistically linked the xenobiotic receptor AHR in mediating

microbial signals to the keratinocyte to boost epithelial differentiation and adhesion, and thus

integrity of the EPB. The absence of such signaling resulted in exacerbated pathology and

increased bacterial infection in models of barrier disruption and epicutaneous sensitization.

Finally, we showed that topical association with a consortium of human skin commensals

restores skin barrier function and integrity, an effect that was dependent on the keratinocyte

AHR.

Our findings parallel those in the gastrointestinal tract, where gut commensals have

been demonstrated to regulate intestinal barrier formation by modulating epithelial turnover [66]

and controlling mucus production [67]. However, unlike the simple mucosal epithelium that

provides the intestinal barrier, the skin is composed of a multi-layered stratified squamous

epithelium that terminally differentiates. Such complexity requires tightly orchestrated signals to

balance differentiation with proliferation. We show here that skin commensals produce signals

that directly regulate epithelial stratification through the xenobiotic sensor AHR. Future studies

will be required to address the identity of the microbial metabolites that interact spatially within

the complex architecture of the skin.

Our studies focused on the keratinocyte AHR and its role in forming the EPB, but the

AHR can also be expressed by epidermal Langerhans cells, innate and adaptive immune cells,

and dermal cells. Previous studies suggest that repair of the EPB is not dependent on AHR

derived from Langerhans cells [58]. However, during epicutaneous sensitization, Langerhans

specific loss of AHR led to decreased Langerhans cells number and function and dysregulated

T cell responses [68]. Thus, AHR likely represents a sensor for Langerhans cell activation as

part of the immunological barrier. Further studies will need to better define the cell-type

specificity of AHR signaling to the different levels of barrier function, including the immune

barrier.

Through depletion models and topical association with defined microbial consortia, we

demonstrate the necessity and sufficiency, respectively, of commensal microbes in epithelial

barrier function. We note that while colonization of germ-free mice with commensal microbes for

two-weeks (COL) resulted in restoration of key epithelial differentiation signals, there were still

differences between gene expression profiles of COL mice and SPF mice raised in the

presence of microbes. It has been shown that skin-resident immune cells rely on imprinting by

early life bacterial exposures to regulate acute wound healing in adult mouse skin [69].

Therefore, it is possible that certain skin development genes rely on early-life exposure to the

microbiota for complete restoration of epithelial gene expression profiles.

Our data supports the hypothesis that sensing of microbial signals by the xenobiotic

receptor AHR are crucial for self-renewal required of the epidermis. Topical application of coal

tar is one of the oldest therapies for atopic dermatitis and has been shown to activate AHR to

induce epithelial differentiation [70]. Recently, the natural product derived small molecule,

Tapinarof, was found to bind and activate the AHR to moderate inflammatory responses in

atopic dermatitis and psoriasis [71]. In support of the hypothesis that AHR mediates microbial

signals to promote barrier function, we found that treatment with the potent and selective AHR

ligand FICZ at a low dose was able to restore epithelial barrier repair in GF mice and induced

epithelial differentiation. However, the role of AHR in skin barrier regulation may be highly

context dependent. In murine models, exposure to pollutants can lead to hyperactivation of AHR

that results in skin barrier damage and inflammation, which mirrors the phenotype of mice that

constitutively express AHR in the keratinocyte [72, 73]. Thus, the balance in the specificity and

quantity of AHR ligand, from endogenous and environment sources, is likely a key factor in

modulating downstream signaling and impact on the skin, and requires further investigation.

AHR has now been recognized as an intracellular pattern recognition receptor that can

identify and metabolize bacterial pigmented virulence factors, and promote antibacterial defense

responses [74]. Our studies show that loss of AHR in skin led to increased susceptibility to the

skin pathogen *S. aureus.* It remains to be determined whether increased susceptibility was due

to impaired physical barrier, impaired antimicrobial barrier, or both. For example, the microbiota-

induced antimicrobial protein RELMα protects against skin infection in a Vitamin A dependent

manner [75]. Since there is interaction between AHR and retinoic acid signaling pathways [76,

77], this may represent a mechanism by which the skin microbiota mediates the antimicrobial

barrier. The differential roles for pathogens and commensals in regulating AHR and promoting

downstream effects are undefined, though are critical when considering diseases of barrier

impairment or wounding, which are often complicated by *S. aureus* colonization and/or infection.

In summary, our findings show a role for skin microbiota in regulating epithelial

differentiation and barrier function in stratified epithelia through AHR. These studies show that

skin microbiome directly impacts development of the epidermal physical barrier. Future studies

that address how microbial communities interact with each other to influence xenobiotic signals

in homeostatic versus disease states will help leverage how personalized microbiota-based

therapies can be used to improve the skin barrier.

FIGURE LEGENDS

Figure 1. Commensal microbiota regulates epithelial barrier genes. (A) Three

groups of mice were employed, specific pathogen free (SPF), germ free (GF), and germ-free

mice colonized (COL) with SPF microbiota for 2 weeks. **(B)** Skin microbiota composition

determined by 16S rRNA gene sequencing. Y-axis indicates absolute read counts of most

abundant phylum (by relative abundance in the dataset) for each mouse (x-axis). **(C)** RNA-seq

workflow. **(D)** Overlap of differentially expressed genes when comparing groups of gnotobiotic

mice. **(E)** Shown in white are the number of genes that were further analyzed for uniquely

enriched gene ontology biological process terms for aforementioned DEGs. Shown on the y-

axis are the uniquely enriched terms, with *p*-values indicated on the x-axis. *P*-values are based

on Fisher's exact test and FDR-adjusted under dependency using the "BY" method. **(F)** To

examine DEGs involved in epithelial barrier function, we manually curated genes involved in

different facets of epithelial barrier: keratinization, cornified envelope formation, adherence and

gap junction, basement membrane function, barrier development, differentiation and intercellular

lipid-lamellae processing. Shown here is a snapshot of key genes that were differentially

expressed in the SPF vs GF subset (*p<0.001*). Horizontal bars represent the Log2 fold-change

comparison (genes upregulated in SPF: log2FC>0, downregulated in SPF: log2FC<0).

Error bars represent standard error estimate for the log2 fold-change computed using the

DESeq2 package. **(G)** Tail-skin from SPF and GF mice. Immunofluorescence-based detection

of differentiation markers (i) loricrin (red) and (ii) keratin-10 (purple) and adhesion marker (iii)

desmoglein-1(grey). Nuclei are counter-stained with Hoechst stain (blue). Images were taken at

constant light exposure (100ms for proteins of interest and 10ms for DAPI channel) and then

overlaid for representation. White dashed-line indicates boundary separating epithelial-stromal

compartments. Scale bar (10µm) is indicated in white (bottom-right). For quantification 10-12

random images were taken in a blinded fashion for each mouse tissue (n=6) at constant light

exposure and processed through ImageJ. For each image integrated density of signal was

normalized to Hoechst stain signal from the same area. Each dot corresponds to average

normalized signal across 10-12 images for each mouse. Asterisk indicates statistical

significance (p<0.05, T test, two-sided). See also Table S1, S2, S3, S4 and Figure S1.

Figure 2. Commensal microbiota promotes skin barrier repair function. (A) Schematic

depicts (i) principle of measuring transepidermal water loss (TEWL) to assess barrier repair

function in adult mice (6-8 weeks old). (ii) Experimental design for assessing barrier recovery.

Dorsal skin of mice was tape-stripped to achieve comparable insults and TEWL was measured

up to 24 hours post-tape stripping. Effect of colonization of microbes was assessed by

comparing age-matched germ-free (GF) and specific pathogen-free (SPF) mice (n=4 mice per

group) in **(B)** wild-type C57/BL6 mice [ANCOVA, $F_{(1,69)} = 50.649$, ***$P<0.001$*)] and **(C)** $Rag1^{-/-}$

mice [ANCOVA, $F_{(1,53)} = 188.1$, ***$P<0.001$*)]. **(D)** Primary mouse keratinocytes were derived

wild-type GF (n=4) and SPF (n=4) C57/BL6 mice and grown in 5% FBS and 1.6mM Ca^{2+} for

three days to induce terminal differentiation. **(E)** Expression of genes involved in differentiation

[Involucrin *(Ivl),* cytokeratin-10 *(Krt10)*] and adherence [Corneodesmosin *(Cdsn)*, Desmocollin-1

(Dsc1), Desmoglein-1a *(Dsg1a)*] was assessed by qRT-PCR. Cycle thresholds were normalized

to housekeeping genes (*Rplp2*, *Sptbn1* and *18S rRNA*) and normalized relative to Cq values of

SPF condition. Each square represents average readings from keratinocytes (n= 4 technical

replicates) derived from an individual mouse (n=4 mice per group). *$P<0.01$, ** $P<0.001$ by T-

test adjusted by Bonferroni correction. **(F)** Primary keratinocytes were grown on transwells in

5% FBS and 1.6mM Ca^{2+}. Epithelial adhesion was assessed by measuring transepithelial

electrical resistance (TEER) at indicated time points. Data from one experiment is represented

for visualization (See Figure S2). One dot represents average TEER readings from technical

replicates (n=3) derived from one individual mouse (n=4 mice per group). ***$P<0.001$ by two-

way ANOVA adjusted for multiple experiments. **(G)** To decrease skin microbial burden, wild-

type SPF mice were treated with antibiotic cocktail (n=5) or vehicle (n=6) for two weeks. **(H)** To

determine microbial burden, mice were swabbed 14 days after treatment and colony forming

units (CFU) were determined. **(I)** Genomic DNA was extracted from swabs collected at baseline

(Day-0) and after one week of treatment (Day 7). V1-V3 region was amplified and analyzed by

16S rRNA gene sequencing and overall abundance i.e. total operational taxonomic unit (OTUs)

counts belonging to different phyla in each sample are depicted**.** Phyla with total read count

<1000 are grouped into 'Other'. (See Figure S3) **(J)** At the end of two weeks the two groups of

mice were tape stripped to achieve comparable insults and TEWL was measured and plotted

against time [ANCOVA, *F (1,41) =26.315, ***P<0.001)*]. TEWL/TEER vs time readings were

fitted by linear modeling (in **B, C, F** and **J**) and significance was assessed by ANCOVA analysis.

Modeling parameters (adjusted R^2 and *F*-statistics) are indicated on top-right for each plot. Span

indicated by shaded area represents 95% CI. Temperature and humidity conditions during

TEWL measurement are indicated for each experiment. Also see Figures S2 and S3.

Figure 3. Activation of aryl hydrocarbon receptor (AHR) signaling in skin rescues barrier

dysfunction in germ free mice. (A) Differentially expressed genes (DEGs) in SPF vs GF mice,

as identified by RNAseq, were mapped onto the AHR pathway. Log_2FC of significant DEGs

(*P<0.01)* in SPF mice are represented in colored boxes according to the legend. **(B)** Schematic

illustrating experimental design. Age matched 6-week-old, GF and SPF (n=5 mice/group)

C57/BL6 mice, respectively, were treated daily with 6-formylindolo[3,2-b] carbazole (FICZ) at

100µg/kg or vehicle for 2 weeks. At end of treatment dorsal skin of mice was tape-stripped to

achieve comparable insults (~20g/m^2/h) and TEWL was measured up to 24 hours post-tape

stripping to assess barrier recovery. TEWL vs time readings were fitted by linear modeling and

covariance was assessed by ANCOVA. Barrier recovery was compared in **(C)** GF mice [*F (1,47)*

=21.9, ***P<0.001)] and **(D)** SPF mice *[F(1,57)=2.98, *P=0.0492]* that were either treated with

FICZ or vehicle. **(E)** Expression of genes [*Ahr, Cdsn, Cyp1a1, Cyp1b1, Dsg1a, Ivl* and Krt14]

was assessed by qRT-PCR in GF mouse skin treated with FICZ or vehicle (4 mice per group).

Cycle thresholds were normalized to housekeeping genes (*Rplp2*, *Sptbn1* and *18s rRNA*) and

normalized relative to Cq values of FICZ treatment. **P<0.05,* by T-test adjusted by Bonferroni

correction. **(F)** Primary human keratinocytes grown on transwells (in 5% FBS and 1.6mM Ca^{2+})

in presence of FICZ (0nM, 10nM and 100nM) for three days and transepithelial electrical

resistance (TEER) was measured. Cells from different donors are represented by different

symbol. See Figure S4. **(G)** Primary human keratinocytes (in 5% FBS and 1.6mM Ca^{2+}) were

treated as indicated with FICZ and/or AHR inhibitor at 100nM doses. TEER values at the end of

three days of treatment are reported. ****P<0.001* by T-test for panels F and G.

Figure 4. Loss of aryl hydrocarbon receptor (AHR) in keratinocytes impairs skin barrier in

mice. (A) *Ahr* floxed *($Ahr^{f/f}$)* allele was knocked out in mice using Cre driven by a keratin-14

promoter *($K14^{Cre/+}Ahr^{f/f}$)*. Dorsal skin was tape-stripped to achieve comparable insults (20-25

$g/m^2/h$) and TEWL recovery curves were compared between *$Ahr^{f/f}$* (n=5) and *$K14^{Cre/+} Ahr^{f/f}$* (n=6)

mice [ANCOVA, *F (1,96) =131.34, ***P<0.001)*]. **(B)** Primary mouse keratinocytes were derived

from mice and polarized in 5% FBS and 1.6mM Ca^{2+} for three days and transepithelial electrical

resistance (TEER) was measured (***P<0.005,* T-test). **(C)** *$Ahr^{f/f}$* (n=5) and *$K14^{Cre/+} Ahr^{f/f}$* (n=7)

mice were tape-stripped (TEWL=20 $g/m^2/h$) and 10^7 CFU *S. aureus* containing tdTomato was

applied to back skin. **(D)** 48 hours post-infection tissue was collected, weighed, homogenized

and plated. *S. aureus* (visible as red colored colonies) and total bacterial colonies were counted

(***P<0.005,* T-test). **(E)** Model for atopic dermatitis induced by repeated epicutaneous

sensitization of tape-stripped skin with ovalbumin (OVA) or vehicle (PBS) [30] was implemented.

Mice were tape stripped at the beginning of the experiment (TEWL~20 $g/m^2/h$) and OVA was

applied daily for 7 days, 3 times with rest for 2 weeks between each treatment. At the end of

final treatment, mice were tape-stripped (TEWL~40 $g/m^2/h$) and 24 hours later **(F)** TEWL levels

were assessed and **(G)** *S. aureus* was applied to back skin, *S. aureus* CFUs were determined

(as described in E). Statistical significance in panels G and H were assessed using a 2-way

ANOVA (***P<0.005*, ****P<0.0005*). (See Figure S4)

Figure 5. Commensal microbes curated from human skin restores skin barrier function in

germ-free mice via AHR activation. (A) Curation of bacteria for Flowers' Flora (FF)

consortium. **(B)** HaCaT cells (10^6 cells/well) were co- transfected with plasmids containing a

Firefly luciferase reporter conjugated to *Cyp1a*- AHR response element and a *Renilla* luciferase

as transfection control. Transfected cells were treated with indicated bacteria at indicated

multiplicity of infection (MOI) and luminescence was measured. *Firefly* luciferase activities were

normalized to *Renilla* luciferase levels and relative response compared to 10nM FICZ treatment

(positive control) was computed. **(C)** Germ-free mice were colonized with FF daily for two

weeks. Mice were swabbed at indicated days and CFUs were enumerated by plating on blood

agar plates. **(D)** To determine whether individual bacteria of FF colonized skin, genomic DNA

was extracted from skin swabs collected from mice at day 14 and qPCR analysis was

conducted using species-specific primers for each bacterium and percentage composition

relative to total 16S rRNA was determined. **(E)** Two-weeks post colonization mice (n=5

mice/group) that were either germ-free (GF) or colonized with FF were tape stripped (TEWL

~20-30 g/m^2/h) and **(F)** barrier recovery was assessed by TEWL (ANCOVA, *F(1,157)=181.25,*

P<0.0001). **(G)** Expression of genes [*Ahr, Cdsn, Cyp1a1, Cyp1b1, Dsg1a, Ivl* and Krt14] was

assessed by qRT-PCR in mouse skin treated with FF or vehicle (GF). Cycle thresholds were

normalized to housekeeping genes (*Rplp2*, *Sptbn1* and *18s rRNA*) and normalized relative to

Cq values of FF treatment. **P<0.05* and ***P<0.005* by T-test adjusted with Bonferroni

correction. **(H)** Primary mouse keratinocytes were derived and polarized in 5% FBS and 1.6mM

Ca^{2+} for three days and transepithelial electrical resistance (TEER) was measured (***P<0.005,*

T-test). **(I)** To test if improved barrier recovery via FF is through AHR, *K14$^{Cre/+}$Ahr$^{f/f}$* (n=6) were

pre-colonized as shown in Fig. 4F and compared to *K14$^{Cre/+}$Ahr$^{f/f}$* (n=3) that were treated with

Control (Ctrl) [ANCOVA, *F(1,61)=0.1191, P=0.73115*]. Additionally, *Ahr$^{f/f}$* mice that were

colonized (n=4) and untreated (n=3) were included in comparisons. **(J)** To test if pre-

colonization with FF could improve barrier recovery in an epicutaneous sensitization model, we

pre-colonized C57BL6/J mice (n=5) with FF for 2-weeks. At the end of colonization, mice were

tape-stripped and treated with Ovalbumin for 1-week. Mice were subjected to comparable

insults (TEWL~40g/m^2/h) and barrier recovery kinetics were compared in FF colonized versus

control (non-colonized) mice. FF colonized mice showed improved barrier recovery compared to

control mice [ANOVA, **P<0.01*]. (See Figures S5).

MATERIALS AND METHODS

RESOURCE AVAILABILITY

Lead contact

Further information and requests for resources and reagents should be directed to and will be

fulfilled by the corresponding author, Elizabeth Grice (egrice@pennmedicine.upenn.edu).

EXPERIMENTAL MODEL AND SUBJECT DETAILS

Animal models and husbandry conditions

All mouse experiments were conducted under protocols approved by the University of

Pennsylvania Institutional Animal Care and Use Committee (Protocol #804065). Age-matched

6-8 weeks old mice were used for all experiments. The following strains of mice were used in

these studies: C57BL6/J (JAX stock #000664), Rag1 KO (JAX stock #002216)[78], Ahr^{fx} (JAX

stock #006203)[79] and K14cre (JAX stock #018964)[80].

Germ-free studies were conducted in the Penn Gnotobiotic Mouse facility. Indicated

strains were bred and maintained as germ-free (GF) in flexible vinyl isolators at the Penn

Gnotobiotic Mouse facility housed in the University of Pennsylvania, School of Veterinary

Medicine. Mice were housed as 3-5 mice per cage, until they were euthanized for tissue

harvest. Aggressive mice or those that showed scratching wounds were not used in the studies.

SPF counterparts were purchased from Jackson Laboratories and allowed to acclimatize in the

facility for one week prior to beginning any experiments. One week prior to beginning of an

experiment, mice (GF or SPF) were transferred to hermetically-sealed cages with individually

filtered-positive airflow. The mice were maintained in these cages for the duration of the study.

This allowed similar housing conditions for both GF and SPF mice for consistent TEWL

readings. All mice were given autoclaved bedding, water and irradiated chow (5021

Autoclavable Mouse Breeder Diet, LabDiet®). Mouse handling was conducted in a laminar flow

cabinet through double layer protective gloves. Germ-free status of mice was confirmed by

weekly bacterial checks by the germ-free facility. At the end of each experiment, mice were

autopsied and germ-free status was confirmed by enlarged cecum. Additionally, skin swabs,

fecal pellet, and bedding samples were cultured by on blood-agar plates.

Studies in wild-type and AHR KO mice were conducted in Clinical Research Building

vivaria at the University of Pennsylvania. C57BL6/J mice were maintained and bred by lab

personnel. To generate $K14^{Cre/+}Ahr^{f/f}$ mice $Ahr^{f/f}$ mice were crossed with $K14^{Cre/+}$ mice to

generate $K14^{Cre/+}Ahr^{f/+}$ F1 mice, $K14^{Cre/+}Ahr^{f/+}$ F1 mice were backcrossed to $Ahr^{f/f}$ mice to obtain

experimental mice ($K14^{Cre/+}Ahr^{f/f}$) and litter-mate controls ($Ahr^{f/f}$) lacking *Cre*. To ensure

robustness of studies, mice were randomly housed. Genotyping protocol: $Ahr^{f/f}$ status was

determined using PCR primers oIMR6075-Reverse (5' CAG TGG GAA TAA GGC AAG AGT GA

3') and oIMR6076-Forward (5' GGT ACA AGT GCA CAT GCC TGC 3') and resolving on a 5%

polyacrylamide gel. *Cre* allele was determined using Generic Cre protocol (Protocol #22392,

Jax Lab, Version 1.3) and resolved on a 2.5% Agarose gel.

Primary keratinocytes from adult mouse skin

Keratinocytes were derived from mouse tail or ear skin as described previously with

slight modifications [81]. Following euthanasia mouse ears and/or tail were excised. With the

help of forceps, ears were split into dorsal and ventral halves. To peel the tail skin from the

bone, a scalpel was used to cut along the ventral axis from base of tail to tip. The exposed tail

bone was peeled off using blunt-tip forceps. The resultant skin was cut into $0.75cm^2$ pieces.

Skin obtained from ears and tails were incubated dermis side down in 6-well dishes and floated

in ice-cold dispase (1mg/ml) in 1X PBS overnight at 4^oC. Epidermal sheets were separated by

lifting the epidermis using forceps. The separated epidermal sheets were cut into tiny pieces

using forceps and scissors and incubated in 60mm untreated culture dish containing 2ml of

0.25% Trypsin-EDTA at 37^oC, 5% CO_2 for 15 minutes. At this point, 5ml suspension media

559 (DMEM+10% FBS+ P/S) was added to the dish and the skin pieces were pipetted vigorously

560 using a 10ml pipette to obtain a single cell suspension. Cell suspension was centrifuged at 150g

561 for 5 minutes at 4°C and supernatant was removed. The cell pellet was suspended in 10ml

562 suspension media and passed through 100µm cell strainer. Cell suspension was centrifuged at

563 150g for 15 minutes at 4°C, supernatant was removed and cells were suspended in 1ml

564 suspension media and plated in collagen-coated 60mm dishes at 0.5 mouse equivalents (i.e. 5

565 million cells/ml) in plating media (low $Ca2^+$ KSFM+ growth supplements+ 5% dialyzed FBS+4%

566 DMEM) containing 10µM ROCK inhibitor (abcam120129) to prevent differentiation as described

567 previously[82]. Typical cell counts were 2-5 x 10^6 cells per mouse (cell count) and viability

568 greater than 70% (cell viability) as determined by Trypan blue exclusion assay. Dishes were

569 collagen coated by incubating 1.5ml collagen solution (50µg/ml collagen in 0.02N acetic acid)

570 for 1 hour at 37 °C, 5% CO_2 or overnight at room temperature. After incubation plates were

571 rinsed 3 times with 1X PBS. Cells were remained undisturbed for 48 hours and then passaged

572 for different experiments. Cells were maintained at 37°C in an atmosphere of 5% CO2 with

573 humidity.

574 ***Human keratinocyte cultures***

575 <u>Primary cultures of human keratinocytes</u> were obtained from neonatal foreskins through

576 the Penn Dermatology Skin Biology and Diseases Resource-based Center: Skin Translational

577 Research Core (STaR) Core B (visit: https://dermatology.upenn.edu/sbdrc/core-b/). Each

578 experiment was conducted with at least three donors (as indicated in text). All experiments were

579 conducted with cells at passage number less than 4. Briefly, cell suspensions were generated

580 using dispase and trypsin, and the cells were cultured in a keratinocyte growth media [50%

581 Medium 154, M154500 (Life Technologies), 50% Keratinocyte SFM, 17005042 (Life

582 Technologies), 1% HKGS supplement, S0015 (Life Technologies), 1% Antibiotic/Antimycotic,

583 15240062 (Invitrogen)]. For routine passaging, cells were split when they were less than 70%

584 confluent. Cells were washed with 1X PBS and trypsinized with 0.25% Trypsin-EDTA for 5

minutes, trypsin was inactivated using trypsin inhibitor (R007100, Thermo Fisher Scientific) and

cell suspension was centrifuged. Following removal of supernatant, cell pellet was suspended in

culture media and seeded as per experimental design. Cells were maintained at 37°C in an

atmosphere of 5% CO2 with humidity.

Immortalized human keratinocyte HaCaT cells [83] were used for AHR reporter assay.

HaCaT cells were verified for lack of mycoplasma contamination by ATCC. Experiments were

conducted on cells that were at passage numbers between 26-36. For routine cell culture,

HaCaTs were maintained in DMEM high glucose (11965092, Thermo Fisher Scientific)

supplemented with 1% Sodium Pyruvate, 5% FBS, 1% Antibiotic/Antimycotic, 15240062

(Invitrogen) and 1% Non-essential amino acids). For routine passaging, cells were split when

they were less than 70% confluent. Cells were washed with 1X PBS and trypsinized with 0.25%

Trypsin-EDTA for 5 minutes, trypsin was quenched with DMEM containing 5% FBS. Cells were

maintained at 37°C in an atmosphere of 5% CO2 with humidity.

Microbial strains

The Flowers' Flora Consortium, consisted of *Staphylococcus epidermidis* (EGM 2-01)*,*

Staphylococcus hemolyticus (EGM 2-08)*, Staphylococcus warneri* (EGM 2-09)*, Micrococcus*

luteus (EGM 2-04) and *Corynebacterium aurimucosum* (EGM 2-02)*,* that had been isolated from

healthy human skin and maintained in the Grice lab culture repository (number in parenthesis

indicates identifier code in Grice lab culture collection). The *S. aureus* strain (AH3926) used in

these studies was generously provided by Dr. Alexander Horswill (University of Colorado,

Anschutz Medical Campus). *S. aureus* AH3926 consists of tdTomato stably integrated into *S.*

aureus LAC (AH1263) and it's construction has been described in detail previously [60].

Culturing conditions: All strains were cultured on solid blood agar plates at room temperature for

24-48 hours. For liquid cultures, all species (except *M. luteus*) were inoculated in tryptic soy

broth and grown by shaking at 100rpm at 37°C. *M. luteus* was inoculated in nutrient broth.

METHOD DETAILS

RNA-sequencing of murine epithelia and analysis

Mice were shaved, and skin was collected from dorsal region. The fat layer was scraped off using a scalpel and then the skin was floated in dispase (1mg/ml) in 1X PBS overnight at 37^{o}C for 1 hour in order to separate the epidermis from the dermis. The epidermis was stored in RNAlater. Mouse epidermis that had been stored in RNA-Later (Thermo-Fisher) was blotted dry and approximately 20 mg of tissue was placed in a Lysing Matrix A tube (MP Bio) with 600 µl RLT buffer (Qiagen) containing 2-mercaptoethanol. The tissue was homogenized with three, 1 min bursts of bead beating in a FastPrep 24 (MP Bio). The lysate was centrifuged (14,000 x g, 3 min) and the supernatant was transferred to a new tube to which 1 volume of 70 % ethanol was added. RNA was purified using a RNeasy Tissue Kit (Qiagen), as per manufacturer's guidelines. RNA was quantified on a Qubit and RNA-integrity was assessed using BioAnalyser according to manufacturer's instructions. 1µg RNA was used to construct RNA-seq libraries using the stranded-TruSeq RNA Sample Prep Kit (Illumina), spiked with phiX and sequenced on the Illumina NextSeq-500 Platform in 3 runs of 1x75 reads. The three runs were aggregated and then analyzed and aligned against the mouse genome [Genome Reference Consortium Mouse Build 38 patch release 5 (GRCm38.p5)] using AlignerBoost [84] and STAR 2.5.3[85]. Gene counts were fitted into a negative binomial model where both the gnotobiotic condition (SPF, GF or COL) and sex of the mouse were included using the DESeq2 [86] R package . Pairwise DEGs between conditions were obtained by setting corresponding "contrasts" for each pairwise comparison and filtering genes with FDP adjusted p-values less than 0.1. To identify enriched Gene Ontology (GO) terms, all annotated GO terms for aforementioned DEGs were retrieved using the ENSEMBL biomaRt R package[87], and significant enriched GO terms were identified using the topGO R package with the FDR-adjusted p-values < 0.1 under dependency [88].Uniquely enriched GO terms were selected by grouping similar GO terms using the online GO visualization tool REVIGO[89] with default (medium) similarity settings.

16S rRNA Gene Sequencing

Sample collection

Mice were swabbed prior to shaving with sterile foam-tipped applicators (Puritan) as

described previously [49]. The swabs were snap frozen and stored at −80°C immediately

following collection. Bacterial DNA was extracted from swabs as described [90]. In brief, swabs

were incubated for one hour at 37°C with shaking in 300µL yeast cell lysis solution (from

Epicentre MasterPure Yeast DNA Purification kit) and 10,000 units of ReadyLyse Lysozyme

solution (Epicentre). Samples were subjected to bead beating for ten minutes at maximum

speed on a vortex mixer with 0.5 mm glass beads (MoBio), followed by a 30-minute incubation

at 65°C with shaking. Protein precipitation reagent (Epicentre) was added and samples were

spun at maximum speed. The supernatant was removed, mixed with isopropanol and applied to

a column from the PureLink Genomic DNA Mini Kit (Invitrogen). Instructions for the Invitrogen

PureLink kit were followed exactly, and DNA was eluted in 50 mL elution buffer (Invitrogen). At

each sampling event, swab control samples that never came into contact with the skin were

collected, prepared and sequenced exactly as the experimental samples. No significant

background contamination from either reagent and/or collection procedures was recovered.

Sequencing and analysis

Amplification of the 16S rRNA gene V1−V3 region was performed as described

previously [90]. Sequencing was performed at the PennCHOP microbiome core on the Illumina

MiSeq using 300 bp paired-end chemistry. The mock community control (MCC; obtained from

BEI Resources, NIAID, NIH as part of the Human Microbiome Project: Genomic DNA from

Microbial Mock Community B (Even, Low Concentration), v5.1L, for 16S rRNA Gene

Sequencing, HM-782D) was sequenced in parallel. Sequencing of the V1-V3 region was

performed using 300 bp paired-end chemistry. Sequences were preprocessed and quality

filtered prior to analysis, including size filtering to 460-600 nucleotides. HmmUFOtu was used

for sequence alignment and phylogeny-based OTU clustering as described previously [91].

Statistical analysis and visualization was performed using the phyloseq package [92] in the R

statistical computing environment.

Barrier recovery

Barrier was assessed as described [93, 94] by using noninvasive probe

(Courage+Khazaka, Cologne, Germany) to measure transepidermal water loss (TEWL) by

diffusion (Tewameter ®, TM300) according to the manufacturer's instructions. The dorsal flanks

of mice were shaved 24 hours prior to beginning of barrier analyses. Basal epidermal

permeability barrier function was assessed 24 hours after shaving. Barrier was disrupted by

tape stripping (3M Scotch High Performance Packaging Tape, 2"X800") to achieve comparable

insults between experimental and control groups as indicated for each experiment. Mice were

anesthetized using isoflurane during TEWL measurements. TEWL measurements were

averaged at 1-second intervals for a 30 second period. Indoor ambient temperature and mean

relative humidity were recorded for each experiment. For barrier recovery assessment, TEWL

was measured by placing probe at the same location on the dorsal flank of the mouse each

time. For consistency, the same person made all TEWL measurements.

Antibiotic treatment of mice

An antibiotic cocktail consisting of Metronidazole (1g/L), Sulfamethoxazole (0.8g/L),

Trimethoprim (0.16g/L), Cephalexin (4g/L) and Baytril (0.025g/L) dissolved in drinking water

containing Splenda (1 packet/250ml) as sweetener was provided to the mice for two weeks. To

ensure decreased microbial burden, cages were changed 3 times a week for antibiotic treated

mice as described previously [49]. Control cages were changed once a week, as per

conventional policies to ensure microbial biodiversity.

Differentiation assay

Confluent cells were trypsinized (0.25% Trypsin EDTA) and plated at 10^4 cells/well in 12

well collagen coated dishes without ROCK inhibitor. Cells were allowed to grow to confluency

(typically 2-3 days) and then media was switched to cornification medium i.e. traditional E-media

without EGF (3 parts DMEM+ 1 part DMEM/F12 + 5% FBS + cholera toxin + insulin + adenine +

hydrocortizone + antibiotics) for three days, to induce polarization and transmembrane junction

formation in keratinocytes [95]. At the end of three days cells were scraped off and processed

for qPCR analyses.

Transepithelial electrical resistance (TEER) measurements

Keratinocytes were trypsinized (0.25% Trypsin EDTA) and plated at 10^4 cells/well on

collagen coated 12mm transwells with 0.4µm pore (Sigma-Aldrich; CLS3460). Twenty-four

hours post-plating cells were 100% confluent and media was switched to KSFM containing 5%

FBS and 1.6mM Ca^{2+}. Every 24 hours, transepithelial electrical resistance (TEER) was

measured using an epithelial volt/ohm meter (EVOM) using a STX2 manual electrode. To

measure TEER, three readings were taken per chamber to cover three different areas of the

transwell membrane longitudinally over three days. The electrode was cleaned with 0.5%

bleach followed by 70% ethanol between each transwell. Readings are reported in ohms-cm^2.

FICZ and AHR inhibitor treatments: For experiments described in Figure 4, primary human

keratinocytes were seeded on transwells and grown in presence of 100nM each of AHR ligand

6-formylindolo[3,2-b] carbazole (FICZ) (Sigma-Aldrich, #SML1489) and/or AHR inhibitor (Sigma

Aldrich, #CH-223191) throughout the course of the experiments.

Tissue preparation and immunofluorescence analysis

Murine skin tissue was collected and fixed in 4% paraformaldehyde and embedded in

paraffin and sectioned at 6µm, as described previously[10]. Immunofluorescence protocols are

described in detail at dx.doi.org/10.17504/protocols.io.k95cz86 [96]. Briefly, tissue sections were

deparaffinized with xylenes and rehydrated with graded ethanol. Heat-induced antigen retrieval

was performed in 0.01M citrate buffer, pH 6.0 and blocked in 10% normal goat serum.

Antibodies against the following proteins were used at indicated dilution: Cytokeratin-10 1:1000

(Biolegend, #905401), Desmoglein 1a 1:200 (Abcam, #ab124798), Loricrin 1:500 (Biolegend,

#905101). Alexafluor 594 conjugated goat-anti rabbit 1:1000 (ThermoFisher Scientific,

#A32740) was used as secondary antibodies. Tissue was counterstained with Hoechst stain.

Wide-field fluorescent images were acquired using by means of a 20X lens objective on a Leica

DM6000 Widefield Fluorescence Microscope at the University of Pennsylvania, School of

Veterinary Medicine Imaging core. For purpose of quantification 10-12 random images were

taken in a blinded fashion at constant light exposure of 100 miliseconds for the Alexafluor 594

channel and 10 seconds for the Hoechst Stain. Images were processed using ImageJ software

version 10.2 (NIH, Bethesda, MD). Krt10, Dsg1a, and Loricrin levels of each image were

calculated by the integrated density of the signal, normalized to the Hoechst stain signal from

the same area. For statistical analysis, each stain was calculated by taking average levels in

each corresponding to 10-12 images per mouse and a two-sided T-test was used to determine

the significance of signal differences between groups.

Electron Microscopy

The ultrastructural analysis by EM was performed as described previously [97] at the VA

Medical Center and Department of Dermatology at the University of California, San Francisco,

United States. Skin samples were fixed in 2% glutaraldehyde and 2% paraformaldehyde and

post-fixed in reduced ruthenium tetroxide before epoxy embedding. The samples were cut on a

Leica Ultracut E microtome (Leica microsystems, Wetzlar, Germany) and imaged on a JEOL

100CX transmission electron microscope (JEOL, Tokyo, Japan) using a Gatan digital camera.

For quantification, the thickness of the cornified envelope was measured in at least 25 randomly

selected positions in 5 random high-powered electron micrographs of the mid stratum corneum

from three mice of each colonization state. The observer recording these measurements was

blinded to the groups.

AHR reporter assay

Immortalized human keratinocyte HaCaT cells [83] were seeded in 96 well plates at 10^4

cells/well in 100µl KSFM media supplemented with supplements. Twenty-four hours later, when

cells were 80-90% confluent, each well was co-transfected with 20ng of *Renilla* luciferase DNA

pGL4.74 (Promega) and 180ng of *Firefly* luciferase reporter plasmids: pGL4.23 which has

xenobiotic response element (XRE) corresponding to *Cyp1A1* activity. Construction of plasmid

is described previously[20]. Transient transfections of HaCaT cells was performed with

Fugene® HD (Promega, #E2311) at 3:1 Fugene transfection reagent: DNA ratio according to

manufacturer instructions. Twelve hours post-transfection, media containing transfection

complexes was removed and replaced with media containing either indicated bacterial strains to

represent MOI=0.1, 1 and 10, peptidoglycan (10ng) or FICZ (10nM). Twenty-four hours post-

infection, dual luciferase readings (*Renilla* and *Firefly*) were read using Dual-Luciferase®

reporter assay system (Promega, #E1910) on the BioTek Synergy HT fluorescence plate

reader. Background correction was performed by subtracting readings of empty wells from

observed readings and *Firefly* luciferase activity was normalized to *Renilla* luciferase. Relative

response ratio to compare *CYP1A1* induction by each bacterial species to treatment with known

AHR ligand FICZ (10nM) was computed as follows: RRR= [(experimental sample ratio)-

(negative control ratio)]/ [(positive control ratio)-(negative control ratio)][98]. Experimental

sample refers to cells treated with indicated bacteria or peptidoglycan; negative control refers to

unstimulated cells and positive control refers to cells stimulated with 10nM FICZ.

cDNA synthesis for qPCR analyses

Mouse skin or epidermis that had been stored in RNA-Later (Thermo-Fisher) was blotted

dry and approximately 20 mg of tissue was placed in a Lysing Matrix A tube (MP Bio) with 600

µl RLT buffer (Qiagen) containing 2-mercaptoethanol. The tissue was homogenized with three,

1 min bursts of bead beating in a FastPrep 24 (MP Bio). The lysate was centrifuged (14,000 x

g, 3 min) and the supernatant was transferred to a new tube to which 1 volume of 70 % ethanol

was added.   RNA was purified using a RNeasy Tissue Kit (Qiagen). Traces of DNA were

removed with DNase-1 and the RNA was stored at -80°C.RNA was quantified (Qubit) and 10 ng

RNA was used as a template for Superscript III (Invitrogen) reverse transcription with random

hexamer primers. Following treatment with RNase H, the cDNA was stored at -20°C.

Colonizing mice with human skin commensals

For liquid cultures of human skin commensals, *S. epidermidis, S. warneri, S. hemolyticus* and *C. aurimucosum* were inoculated in tryptic soy broth (TSB) and *M. luteus* was inoculated in nutrient broth media, respectively and grown by shaking at 200rpm for 16 hours at 37^{o}C. Cultures were centrifuged, media was removed and bacterial pellets were suspended in PBS to obtain 10^9 CFU/ml. Equal amount of each bacteria (10^9 CFU) was combined in 5ml PBS and inoculated in bedding of mouse cages daily for 2 weeks. For each inoculum, a fresh batch culture was grown overnight.

Mouse treatment with AHR ligand FICZ

Mice were treated with a low dose (100µg/kg) of AHR ligand 6-Formylindolo[3,2-b] carbazole (FICZ) (Sigma-Aldrich, #SML1489), that has been shown to be sufficient to induce expression of *Cyp1a1*[54]. A stock solution (4.5mg/ml) of FICZ was made in DMSO and diluted in 50% Acetone. Mice were shaved 24 hours prior to treatment and FICZ was applied topically by directly pipetting onto shaved skin, daily for 2 weeks (**Figure 3B)**.

***S. aureus* skin infection protocol**

Epicutaneous infection: Infection protocol for epicutaneous infection with *S. aureus* has been described previously [99] and was implemented with slight modifications. Briefly, mice were anesthetized with isoflurane, tape stripped (TEWL=20 g/m^2/h) and 24 hours later, 10^7 CFU *S. aureus* in 100µl was applied to back skin and spread using a swab. The inoculum was allowed to dry for 10 minutes and mice were returned to their cages.

CFU enumeration: Forty-eight hours post-infection, mice were euthanized and approximately 1cm^2 infected skin area was collected and weighed and transferred to tubes containing 300µl 1X PBS. Tissue was homogenized by bead beating for twenty minutes at maximum speed on a vortex mixer with 0.5 mm ceramic beads and CFUs were enumerated by serial dilution on blood agar plates after overnight incubation at 37^{o}C. Both *S. aureus* and total bacterial counts were determined and normalized to weight of tissues. *S. aureus* colonies were

792   visible as red colonies due to stable expression of tDTomato and could be distinguished from

793   total bacteria.

Epicutaneous sensitization with ovalbumin

Procedures to induce barrier defects that mimic atopic dermatitis by repeated

epicutaneous sensitization by ovalbumin (OVA) [30, 100] followed by infection with *S. aureus*

[101] have been described previously. The dorsal skin of mice was shaved and TEWL was

assessed to give a baseline reading. To measure and compare barrier repair between wild-type

and AhR null mice, we standardized the amount of initial barrier disruption to give identical

TEWL values. To achieve this, mice were tape stripped to give a reading of TEWL ~20g/m^2/h.

Twenty-four hours post tape-stripping, the mice were treated daily for 7 days with 100 μg OVA

(Sigma Aldrich, # A5503) suspended in 100 μl PBS was applied onto backs of mice and spread

using a skin swab, and allowed to dry for 2 minutes. For inducing atopic dermatitis-like

condition, the 7-day OVA treatment regime was repeated twice more, with 2 weeks rest

between subsequent treatments. At the end of treatment mice were mice were tape stripped to

give a reading of TEWL ~40g/m^2/h and *S. aureus* was applied as described earlier or barrier

recovery was assessed.

QUANTIFICATION AND STATISTICAL ANALYSIS

Data visualization and statistics.

All statistical analysis was performed using functions built into the R statistical environment

(RStudio Version 1.3.1056). Data was visualized using ggplot2 [102] package and GraphPad

Prism version 8.0.0 for Mac OS X, GraphPad Software, San Diego, California USA,

www.graphpad.com. TEWL/TEER vs time readings were fitted by linear modeling function in R

statistical package and visualized using ggplot2 package. Significance was assessed by

ANCOVA analysis. Fit parameters (adjusted R^2 and *F*-statistics) are indicated for each plot.

Span indicated by shaded area represents 95% CI. Gene expression analysis from qPCR was

conducted as per guidelines described previously[103]. Cycle thresholds were normalized to

housekeeping genes (Rplp2, Sptbn1 and 18s rRNA) and normalized relative to quantitative

cycle (Cq) values of control. Heatmaps for qPCR analysis were made using Morpheus heat map

viewer from Broad Institute (https://software.broadinstitute.org/morpheus). Each square

represents average normalized readings (n= 3 technical replicates). Bonferrroni correction was

used to adjust for multiple comparisons. The AHR pathway was originally downloaded from

wikipathways (https://www.wikipathways.org/index.php/Pathway:WP2873), then modified by

highlighting DEGs identified in our analysis using a customized Perl script. The log2 fold-change

values between SPF and GF mice were used to determine the color hue (red/blue: up/down in

SPF vs. GF) and saturation of the highlighted DEGs.

KEY RESOURCES TABLE

SUPPLEMENTARY MATERIALS

Table S1. Related to Figure 1.

Results from differential expression analysis of RNA-seq data for genes depicted in Fig. 1F. As

a result, we identified 6,396, 427, and 3,232 DEGs for SPF vs. GF, COL vs. GF and SPF vs.

COL comparisons, respectively. DEGs defined as those with FDR adjusted p-values < 0.1. In

Sheet SPF vs GF (upregulatedin SPF: $log2FC > 0$, downregulated in SPF: $log2FC < 0$); in

sheet COLvsGF (upregulated in COL: $log2FC > 0$, downregulated in COL: $log2FC < 0$); in

sheet SPFvsCOL (upregulated in SPF: $log2FC > 0$, downregulated in SPF: $log2FC < 0$).

Table S2. Related to Figure 1. Contains list of genes manually curated based on literature

analyses that have been implicated in barrier function. Genes are listed under the following

categories based on their functions: adherence junction formation, lipid-lamellae formation,

keratin network, differentiation, skin barrier development, formation of cornified envelope and

basement membrane.

Table S3. Related to Figure 1. Contains the key genes involved in skin barrier function that

were differentially expressed between SPF and GF murine skin and were used to generate

graph depicted in Figure 1F.

Table S4. Related to Figure 1. Contains list of differentially expressed barrier genes involved in

skin barrier function that were differentially expressed between SPF and GF murine skin by

using genes described in Table S2 as reference.

Figure S1. Related to Figure 1. Structural analysis of murine dorsal skin from SPF and GF

mice. **(i)** Histopathological analysis of Hematoxylin and Eosin stained tissue did not demonstrate

overt differences between age-matched GF and SPF mice. **(ii)** Electron Microscopy (EM) was

performed on skin and number of layers of cornified envelope were counted. *** indicates

P<0.01, T-test.

Figure S2. Related to Figure 2. Assessment of **(A)** TEWL and **(B)** pH at baseline in age-

matched age-matched germ-free (GF) and specific pathogen-free (SPF) C57BL6/J mice (n=4

mice per group). **(C)** Shown is indicated TEWL corresponding to number of tape strips in GF vs

SPF mice. **(D)** GF and SPF *Rag1-/-* mice were swabbed colony forming units (CFU) were

determined and **(E)** TEWL was measured at baseline. **(F)** Dorsal skin from GF and SPF *Rag1-/-*

mice was collected and gene expression was analyzed by qRT-PCR. Cycle thresholds were

normalized to housekeeping genes (*Rplp2*, *Sptbn1* and *18s rRNA*) and normalized relative to

Cq values of SPF condition. Each square represents average readings from three technical

replicates from each mouse. **P<0.01*, ** *P<0.001* by T-test adjusted by Bonferrroni correction.

(G) Primary keratinocytes were derived from germ free mice and grown on transwells in basal

medium before switching to cornification media containing 5% FBS and 1.6mM Ca^{2+}. Shown

here is expression of genes between GF and SPF mice before and after calcium switch.

Expression of genes involved in differentiation [Involucrin *(Ivl),* cytokeratin-10 *(Krt10)*] and

adherence [Corneodesmosin *(Cdsn)*, Desmocollin-1 *(Dsc1)*, Desmoglein-1a *(Dsg1a)*] and AHR

pathway (*Cyp1a1, Cyp1b1, Hsp90ab1*) was assessed by qRT-PCR. Cycle thresholds were

normalized to housekeeping genes (*Rplp2*, *Sptbn1* and *18s rRNA*) and normalized relative to

Cq values of SPF condition. Each square represents average readings from technical replicates

(n= 4 technical replicates) derived from an individual mouse (n=4 mice per group). **P<0.01*, **

P<0.001 by T-test adjusted by Bonferrroni correction. **(H)** Primary keratinocytes grown on

transwells were grown in 5% FBS and 1.6mM Ca^{2+} epithelial adhesion was assessed by

measuring transepithelial electrical resistance (TEER) at indicated time points. Data from three

individual experiments are shown. One dot represents average TEER readings from technical

replicates (n=3) derived from one individual mouse (n=4 mice per group).

Figure S3. Related to Figure 2. To decrease skin microbial burden, wild-type SPF mice were

treated with antibiotic cocktail (n=5) or vehicle (n=6) for two weeks. Genomic DNA was

extracted from skin swabs and fecal samples collected at baseline (Day 0 i.e. D0) and after one

week of treatment (Day 7 i.e. D7). V1-V3 region was amplified and analyzed by 16S rRNA gene

sequencing. **(A)** Microbiota composition of each sample shown as the relative abundance of the

top 15 most abundant genus of the entire dataset**. (B)** Fecal pellets collected at end of treatment

(i.e. 14 days later) were weighed, homogenized and plated on blood agar plates and CFUs were

enumerated. **(C)** Within-sample alpha diversity for samples from groups shown as the Shannon

index. The within sample Shannon index was calculated using the Phyloseq R package. **(D)**

Principle Coordinates Analysis (PCoA) plot of between microbiota sample beta-diversity for

different sample types (skin or fecal) and antibiotic treatment (control or antibiotic). Ctrl: control

treatment, Abx: Antibiotic treatment. Treatment timepoint was color labeled. The Bray-Curtis

dissimilarity was used for the between-sample beta-diversity metric.

Figure S4. Related to Figure 4. TEWL dynamics during the course of ovalbumin sensitization

treatment regime. **(A)** Schematic depicts sequence of manipulations to mice during OVA

sensitization. Mice were tape stripped at the beginning of the experiment (TEWL~20 $g/m^2/h$)

and OVA was applied daily for 7 days, 3 times with rest for 2 weeks between each treatment. At

the end of final treatment, mice were tape-stripped (TEWL~40 $g/m^2/h$) and *S. aureus* was

applied to back skin. **(B)** TEWL readings were taken at indicated time points in **(i)** $K14^{Cre/+}Ahr^{f/f}$

and **(ii)** $Ahr^{f/f}$ mice treated with OVA or vehicle (PBS)*.* Endpoint readings are reported in Figure

4. **(C)** Total bacterial counts per gram of tissue are reported for $K14^{Cre/+}Ahr^{f/f}$ (KO) and $Ahr^{f/f}$

(WT) mice for indicated comparisons.

Figure S5. Related to Figure 5. TEWL dynamics during the course of short-term ovalbumin

sensitization treatment regime after Flowers' Flora colonization **(A)** Schematic depicts sequence

of manipulations to C57BL6/J mice. Mice were pre-colonized with Flowers' Flora daily for 2

weeks. At the end of pre-colonization, mice were tape stripped at the beginning of the

experiment (TEWL~20 $g/m^2/h$) and OVA was applied daily for 7 days. At the end of final

treatment, mice were tape-stripped (TEWL~40 $g/m^2/h$) and barrier recovery was assessed. **(B)**

TEWL readings were taken at indicated time points in **(i)** mice pre-colonized with Flowers' Flora

and **(ii)** those treated with vehicle (PBS)*.*

ACKNOWLEDGEMENTS

We thank Dr. Alex Horswill (University of Colorado) for the kind gift of the *S. aureus* strain; Dr.

Jorge Henao-Mejia (University of Pennsylvania) for the kind gift of $Ahr^{f/f}$ mice; Dr. John Seykora

(University of Pennsylvania) for the kind gift of HaCaT keratinocytes; Dmytro Kobuley,

Gnotobiotic Mouse Facility (University of Pennsylvania) and current and former members of the

Grice lab and the Department of Dermatology for critical discussion and review of the work.

FUNDING

This work was funded by grants from the National Institutes of Health, National Institute of

Environmental and Health Sciences (R56-ES-030218 to EAG/TRS, R01-ES-017014 to TRS),

National Institute of Arthritis, Musculoskeletal, and Skin Diseases (R01-AR-006663 and R00-

AR-060873 to EAG), the Burroughs Wellcome Fund PATH Award (EAG), the Linda Pechenik

Montague Investigator Award (EAG), and the Dermatology Foundation Sun Pharma Research

Award (EAG). This research was also supported by the Resource Cores of the Penn Skin

Biology and Disease Resource-based Center funded by P30-AR-069589 (*PI: George*

Cotsarelis, M.D.). CBM and LF were supported by the Penn Dermatology Research Training

Grant, T32-AR-007465, from the National Institute of Arthritis, Musculoskeletal, and Skin

Diseases.

REFERENCES

1. Flowers, L. and E.A. Grice, *The Skin Microbiota: Balancing Risk and Reward.* Cell Host Microbe, 2020. **28**(2): p. 190-200.
2. Byrd, A.L., Y. Belkaid, and J.A. Segre, *The human skin microbiome.* Nat Rev Microbiol, 2018. **16**(3): p. 143-155.
3. Grice, E.A. and J.A. Segre, *The skin microbiome.* Nat Rev Microbiol, 2011. **9**(4): p. 244-53.
4. Eyerich, S., et al., *Cutaneous Barriers and Skin Immunity: Differentiating A Connected Network.* Trends Immunol, 2018. **39**(4): p. 315-327.
5. Parlet, C.P., M.M. Brown, and A.R. Horswill, *Commensal Staphylococci Influence Staphylococcus aureus Skin Colonization and Disease.* Trends Microbiol, 2019. **27**(6): p. 497-507.
6. Belkaid, Y. and J.A. Segre, *Dialogue between skin microbiota and immunity.* Science, 2014. **346**(6212): p. 954-9.
7. Leech, J.M., et al., *Toxin-Triggered Interleukin-1 Receptor Signaling Enables Early-Life Discrimination of Pathogenic versus Commensal Skin Bacteria.* Cell Host Microbe, 2019. **26**(6): p. 795-809 e5.
8. Elias, P.M., *The skin barrier as an innate immune element.* Semin Immunopathol, 2007. **29**(1): p. 3-14.
9. Schmid-Wendtner, M.H. and H.C. Korting, *The pH of the skin surface and its impact on the barrier function.* Skin Pharmacol Physiol, 2006. **19**(6): p. 296-302.
10. Meisel, J.S., et al., *Commensal microbiota modulate gene expression in the skin.* Microbiome, 2018. **6**(1): p. 20.
11. Simpson, C.L., D.M. Patel, and K.J. Green, *Deconstructing the skin: cytoarchitectural determinants of epidermal morphogenesis.* Nat Rev Mol Cell Biol, 2011. **12**(9): p. 565-80.
12. Egawa, G. and K. Kabashima, *Multifactorial skin barrier deficiency and atopic dermatitis: Essential topics to prevent the atopic march.* J Allergy Clin Immunol, 2016. **138**(2): p. 350-358 e1.
13. Mackowiak, B., et al., *The Roles of Xenobiotic Receptors: Beyond Chemical Disposition.* Drug Metab Dispos, 2018. **46**(9): p. 1361-1371.
14. Nestle, F.O., et al., *Skin immune sentinels in health and disease.* Nat Rev Immunol, 2009. **9**(10): p. 679-91.
15. Kazem, S., E.C. Linssen, and S. Gibbs, *Skin metabolism phase I and phase II enzymes in native and reconstructed human skin: a short review.* Drug Discov Today, 2019. **24**(9): p. 1899-1910.

16.　Donia, M.S. and M.A. Fischbach, *HUMAN MICROBIOTA. Small molecules from*
the human microbiota. Science, 2015. **349**(6246): p. 1254766.
17.　Burbach, K.M., A. Poland, and C.A. Bradfield, *Cloning of the Ah-receptor cDNA*
reveals a distinctive ligand-activated transcription factor. Proc Natl Acad Sci U S
A, 1992. **89**(17): p. 8185-9.
18.　Esser, C., et al., *Functions of the aryl hydrocarbon receptor in the skin.* Semin
Immunopathol, 2013. **35**(6): p. 677-91.
19.　Furue, M., et al., *Role of AhR/ARNT system in skin homeostasis.* Arch Dermatol
Res, 2014. **306**(9): p. 769-79.
20.　Sutter, C.H., et al., *2,3,7,8-Tetrachlorodibenzo-p-dioxin increases the expression*
of genes in the human epidermal differentiation complex and accelerates
epidermal barrier formation. Toxicol Sci, 2011. **124**(1): p. 128-37.
21.　Sutter, C.H., et al., *EGF receptor signaling blocks aryl hydrocarbon receptor-*
mediated transcription and cell differentiation in human epidermal keratinocytes.
Proc Natl Acad Sci U S A, 2009. **106**(11): p. 4266-71.
22.　Kennedy, L.H., et al., *2,3,7,8-Tetrachlorodibenzo-p-dioxin-mediated production of*
reactive oxygen species is an essential step in the mechanism of action to
accelerate human keratinocyte differentiation. Toxicol Sci, 2013. **132**(1): p. 235-
49.
23.　Stevens, E.A., J.D. Mezrich, and C.A. Bradfield, *The aryl hydrocarbon receptor: a*
perspective on potential roles in the immune system. Immunology, 2009. **127**(3):
p. 299-311.
24.　Nguyen, L.P. and C.A. Bradfield, *The search for endogenous activators of the*
aryl hydrocarbon receptor. Chem Res Toxicol, 2008. **21**(1): p. 102-16.
25.　McIntosh, B.E., J.B. Hogenesch, and C.A. Bradfield, *Mammalian Per-Arnt-Sim*
proteins in environmental adaptation. Annu Rev Physiol, 2010. **72**: p. 625-45.
26.　Esser, C., *The Aryl Hydrocarbon Receptor in Immunity: Tools and Potential.*
Methods Mol Biol, 2016. **1371**: p. 239-57.
27.　Kubo, A., K. Nagao, and M. Amagai, *Epidermal barrier dysfunction and*
cutaneous sensitization in atopic diseases. J Clin Invest, 2012. **122**(2): p. 440-7.
28.　Roberson, E.D. and A.M. Bowcock, *Psoriasis genetics: breaking the barrier.*
Trends Genet, 2010. **26**(9): p. 415-23.
29.　Elias, P.M., *Primary role of barrier dysfunction in the pathogenesis of atopic*
dermatitis. Exp Dermatol, 2018. **27**(8): p. 847-851.
30.　Spergel, J.M., et al., *Epicutaneous sensitization with protein antigen induces*
localized allergic dermatitis and hyperresponsiveness to methacholine after
single exposure to aerosolized antigen in mice. J Clin Invest, 1998. **101**(8): p.
1614-22.
31.　Yockey, L.J., et al., *The absence of a microbiota enhances TSLP expression in*
mice with defective skin barrier but does not affect the severity of their allergic
inflammation. J Invest Dermatol, 2013. **133**(12): p. 2714-2721.
32.　Jensen, J.M., et al., *Impaired cutaneous permeability barrier function, skin*
hydration, and sphingomyelinase activity in keratin 10 deficient mice. J Invest
Dermatol, 2000. **115**(4): p. 708-13.

33. Schmuth, M., et al., *Structural and functional consequences of loricrin mutations in human loricrin keratoderma (Vohwinkel syndrome with ichthyosis).* J Invest Dermatol, 2004. **122**(4): p. 909-22.

34. Chamcheu, J.C., et al., *Keratin gene mutations in disorders of human skin and its appendages.* Arch Biochem Biophys, 2011. **508**(2): p. 123-37.

35. Brandner, J.M., et al., *Epidermal tight junctions in health and disease.* Tissue Barriers, 2015. **3**(1-2): p. e974451.

36. Broussard, J.A., J.L. Koetsier, and K.J. Green, *Desmosomes pattern cell mechanics to govern epidermal tissue form and function.* bioRxiv, 2020: p. 2020.01.21.914176.

37. Samuelov, L., et al., *Desmoglein 1 deficiency results in severe dermatitis, multiple allergies and metabolic wasting.* Nat Genet, 2013. **45**(10): p. 1244-1248.

38. Elias, P.M., et al., *Desmoglein isoform distribution affects stratum corneum structure and function.* J Cell Biol, 2001. **153**(2): p. 243-9.

39. Grubauer, G., P.M. Elias, and K.R. Feingold, *Transepidermal water loss: the signal for recovery of barrier structure and function.* J Lipid Res, 1989. **30**(3): p. 323-33.

40. Fluhr, J.W., K.R. Feingold, and P.M. Elias, *Transepidermal water loss reflects permeability barrier status: validation in human and rodent in vivo and ex vivo models.* Exp Dermatol, 2006. **15**(7): p. 483-92.

41. Tellkamp, F., S. Vorhagen, and C.M. Niessen, *Epidermal polarity genes in health and disease.* Cold Spring Harb Perspect Med, 2014. **4**(12): p. a015255.

42. Hennings, H., et al., *Calcium regulation of growth and differentiation of mouse epidermal cells in culture.* Cell, 1980. **19**(1): p. 245-54.

43. Naik, S., et al., *Compartmentalized control of skin immunity by resident commensals.* Science, 2012. **337**(6098): p. 1115-9.

44. Bowen, A.C., et al., *Sulfamethoxazole-Trimethoprim (Cotrimoxazole) for Skin and Soft Tissue Infections Including Impetigo, Cellulitis, and Abscess.* Open Forum Infect Dis, 2017. **4**(4): p. ofx232.

45. Lofmark, S., C. Edlund, and C.E. Nord, *Metronidazole is still the drug of choice for treatment of anaerobic infections.* Clin Infect Dis, 2010. **50 Suppl 1**: p. S16-23.

46. Beco, L., et al., *Suggested guidelines for using systemic antimicrobials in bacterial skin infections: part 2-- antimicrobial choice, treatment regimens and compliance.* Vet Rec, 2013. **172**(6): p. 156-60.

47. Derrick, C.W., Jr. and K. Reilly, *The role of cephalexin in the treatment of skin and soft-tissue infections.* Postgrad Med J, 1983. **59 Suppl 5**: p. 43-6.

48. Nagase, N., et al., *Isolation and species distribution of staphylococci from animal and human skin.* J Vet Med Sci, 2002. **64**(3): p. 245-50.

49. SanMiguel, A.J., et al., *Topical Antimicrobial Treatments Can Elicit Shifts to Resident Skin Bacterial Communities and Reduce Colonization by Staphylococcus aureus Competitors.* Antimicrob Agents Chemother, 2017. **61**(9).

50. Fu, Z.D., et al., *RNA Sequencing Quantification of Xenobiotic-Processing Genes in Various Sections of the Intestine in Comparison to the Liver of Male Mice.* Drug Metab Dispos, 2016. **44**(6): p. 842-56.

51. Minzaghi, D., P. Pavel, and S. Dubrac, *Xenobiotic Receptors and Their Mates in*
Atopic Dermatitis. Int J Mol Sci, 2019. **20**(17).
52. Wright, E.J., et al., *Canonical and non-canonical aryl hydrocarbon receptor*
signaling pathways. Curr Opin Toxicol, 2017. **2**: p. 87-92.
53. Larigot, L., et al., *AhR signaling pathways and regulatory functions.* Biochim
Open, 2018. **7**: p. 1-9.
54. Ehrlich, A.K., et al., *TCDD, FICZ, and Other High Affinity AhR Ligands Dose-*
Dependently Determine the Fate of CD4+ T Cell Differentiation. Toxicol Sci,
2018. **161**(2): p. 310-320.
55. Fritsche, E., et al., *Lightening up the UV response by identification of the*
arylhydrocarbon receptor as a cytoplasmatic target for ultraviolet B radiation.
Proc Natl Acad Sci U S A, 2007. **104**(21): p. 8851-6.
56. Syed, D.N. and H. Mukhtar, *FICZ: A Messenger of Light in Human Skin.* J Invest
Dermatol, 2015. **135**(6): p. 1478-1481.
57. van den Bogaard, E.H., et al., *Genetic and pharmacological analysis identifies a*
physiological role for the AHR in epidermal differentiation. J Invest Dermatol,
2015. **135**(5): p. 1320-1328.
58. Haas, K., et al., *Aryl Hydrocarbon Receptor in Keratinocytes Is Essential for*
Murine Skin Barrier Integrity. J Invest Dermatol, 2016. **136**(11): p. 2260-2269.
59. Alexander, H., et al., *The role of bacterial skin infections in atopic dermatitis:*
expert statement and review from the International Eczema Council Skin
Infection Group. Br J Dermatol, 2020. **182**(6): p. 1331-1342.
60. Miller, R.J., et al., *Development of a Staphylococcus aureus reporter strain with*
click beetle red luciferase for enhanced in vivo imaging of experimental
bacteremia and mixed infections. Sci Rep, 2019. **9**(1): p. 16663.
61. Cork, M.J., et al., *Epidermal barrier dysfunction in atopic dermatitis.* J Invest
Dermatol, 2009. **129**(8): p. 1892-908.
62. Rooks, M.G. and W.S. Garrett, *Gut microbiota, metabolites and host immunity.*
Nat Rev Immunol, 2016. **16**(6): p. 341-52.
63. Singh, R., et al., *Enhancement of the gut barrier integrity by a microbial*
metabolite through the Nrf2 pathway. Nat Commun, 2019. **10**(1): p. 89.
64. Wikramanayake, T.C., O. Stojadinovic, and M. Tomic-Canic, *Epidermal*
Differentiation in Barrier Maintenance and Wound Healing. Adv Wound Care
(New Rochelle), 2014. **3**(3): p. 272-280.
65. Matsui, T. and M. Amagai, *Dissecting the formation, structure and barrier*
function of the stratum corneum. Int Immunol, 2015. **27**(6): p. 269-80.
66. Kaiko, G.E., et al., *The Colonic Crypt Protects Stem Cells from Microbiota-*
Derived Metabolites. Cell, 2016. **165**(7): p. 1708-1720.
67. Johansson, M.E., et al., *Normalization of Host Intestinal Mucus Layers Requires*
Long-Term Microbial Colonization. Cell Host Microbe, 2015. **18**(5): p. 582-92.
68. Hong, C.H., et al., *Selective AhR knockout in langerin-expressing cells abates*
Langerhans cells and polarizes Th2/Tr1 in epicutaneous protein sensitization.
Proc Natl Acad Sci U S A, 2020. **117**(23): p. 12980-12990.
69. Constantinides, M.G., et al., *MAIT cells are imprinted by the microbiota in early*
life and promote tissue repair. Science, 2019. **366**(6464).

| 1096 | 70. | van den Bogaard, E.H., et al., *Coal tar induces AHR-dependent skin barrier repair in atopic dermatitis.* J Clin Invest, 2013. **123**(2): p. 917-27. |

70.   van den Bogaard, E.H., et al., *Coal tar induces AHR-dependent skin barrier*
repair in atopic dermatitis. J Clin Invest, 2013. **123**(2): p. 917-27.

71.   Smith, S.H., et al., *Tapinarof Is a Natural AhR Agonist that Resolves Skin*
Inflammation in Mice and Humans. J Invest Dermatol, 2017. **137**(10): p. 2110-
2119.

72.   Tauchi, M., et al., *Constitutive expression of aryl hydrocarbon receptor in*
keratinocytes causes inflammatory skin lesions. Mol Cell Biol, 2005. **25**(21): p.
9360-8.

73.   Hidaka, T., et al., *The aryl hydrocarbon receptor AhR links atopic dermatitis and*
air pollution via induction of the neurotrophic factor artemin. Nat Immunol, 2017.
18(1): p. 64-73.

74.   Moura-Alves, P., et al., *AhR sensing of bacterial pigments regulates antibacterial*
defence. Nature, 2014. **512**(7515): p. 387-92.

75.   Harris, T.A., et al., *Resistin-like Molecule alpha Provides Vitamin-A-Dependent*
Antimicrobial Protection in the Skin. Cell Host Microbe, 2019. **25**(6): p. 777-788
e8.

76.   Murphy, K.A., L. Quadro, and L.A. White, *The intersection between the aryl*
hydrocarbon receptor (AhR)- and retinoic acid-signaling pathways. Vitam Horm,
2007. **75**: p. 33-67.

77.   Andreola, F., et al., *Aryl hydrocarbon receptor knockout mice (AHR-/-) exhibit*
liver retinoid accumulation and reduced retinoic acid metabolism. Cancer Res,
1997. **57**(14): p. 2835-8.

78.   Mombaerts, P., et al., *RAG-1-deficient mice have no mature B and T*
lymphocytes. Cell, 1992. **68**(5): p. 869-77.

79.   Walisser, J.A., et al., *Aryl hydrocarbon receptor-dependent liver development*
and hepatotoxicity are mediated by different cell types. Proc Natl Acad Sci U S A,
2005. **102**(49): p. 17858-63.

80.   Dassule, H.R., et al., *Sonic hedgehog regulates growth and morphogenesis of*
the tooth. Development, 2000. **127**(22): p. 4775-85.

81.   Lichti, U., J. Anders, and S.H. Yuspa, *Isolation and short-term culture of primary*
keratinocytes, hair follicle populations and dermal cells from newborn mice and
keratinocytes from adult mice for in vitro analysis and for grafting to
immunodeficient mice. Nat Protoc, 2008. **3**(5): p. 799-810.

82.   Chapman, S., et al., *The effect of Rho kinase inhibition on long-term keratinocyte*
proliferation is rapid and conditional. Stem Cell Res Ther, 2014. **5**(2): p. 60.

83.   Boukamp, P., et al., *Normal keratinization in a spontaneously immortalized*
aneuploid human keratinocyte cell line. J Cell Biol, 1988. **106**(3): p. 761-71.

84.   Zheng, Q. and E.A. Grice, *AlignerBoost: A Generalized Software Toolkit for*
Boosting Next-Gen Sequencing Mapping Accuracy Using a Bayesian-Based
Mapping Quality Framework. PLoS Comput Biol, 2016. **12**(10): p. e1005096.

85.   Dobin, A., et al., *STAR: ultrafast universal RNA-seq aligner.* Bioinformatics, 2013.
29(1): p. 15-21.

86.   Love, M.I., W. Huber, and S. Anders, *Moderated estimation of fold change and*
dispersion for RNA-seq data with DESeq2. Genome Biol, 2014. **15**(12): p. 550.

87. Durinck, S., et al., *Mapping identifiers for the integration of genomic datasets with the R/Bioconductor package biomaRt.* Nature Protocols, 2009. **4**(8): p. 1184-1191.

88. Benjamini, Y. and D. Yekutieli, *The control of the false discovery rate in multiple testing under dependency.* Annals of Statistics, 2001. **29**(4): p. 1165-1188.

89. Supek, F., et al., *REVIGO Summarizes and Visualizes Long Lists of Gene Ontology Terms.* Plos One, 2011. **6**(7).

90. Meisel, J.S., et al., *Skin Microbiome Surveys Are Strongly Influenced by Experimental Design.* J Invest Dermatol, 2016. **136**(5): p. 947-956.

91. Zheng, Q., et al., *HmmUFOtu: An HMM and phylogenetic placement based ultra-fast taxonomic assignment and OTU picking tool for microbiome amplicon sequencing studies.* Genome Biol, 2018. **19**(1): p. 82.

92. McMurdie, P.J. and S. Holmes, *phyloseq: an R package for reproducible interactive analysis and graphics of microbiome census data.* PLoS One, 2013. **8**(4): p. e61217.

93. Man, G., et al., *Topical hesperidin enhances epidermal function in an aged murine model.* J Invest Dermatol, 2015. **135**(4): p. 1184-1187.

94. Bradley, C.W., et al., *Longitudinal Evaluation of the Skin Microbiome and Association with Microenvironment and Treatment in Canine Atopic Dermatitis.* J Invest Dermatol, 2016. **136**(6): p. 1182-1190.

95. Simpson, C.L., et al., *Plakoglobin rescues adhesive defects induced by ectodomain truncation of the desmosomal cadherin desmoglein 1: implications for exfoliative toxin-mediated skin blistering.* Am J Pathol, 2010. **177**(6): p. 2921-37.

96. Uberoi, A. *Standard protocol for performing immunofluorescence or immunohistochemistry on paraffin embedded tissue.* 2017.

97. Crumrine, D., et al., *Mutations in Recessive Congenital Ichthyoses Illuminate the Origin and Functions of the Corneocyte Lipid Envelope.* J Invest Dermatol, 2019. **139**(4): p. 760-768.

98. Eggers, C.H., B.; Lewis, S.; Strayer, C.; Landreman, A. *Designing a Bioluminescent Reporter Assay: Normalization Options.* 2016; Available from: https://www.promega.com/resources/pubhub/bioluminescent-reporter-assay-normalization/.

99. Wanke, I., et al., *Staphylococcus aureus skin colonization is promoted by barrier disruption and leads to local inflammation.* Exp Dermatol, 2013. **22**(2): p. 153-5.

100. Jin, H., et al., *Animal models of atopic dermatitis.* J Invest Dermatol, 2009. **129**(1): p. 31-40.

101. Nakatsuji, T., et al., *Staphylococcus aureus Exploits Epidermal Barrier Defects in Atopic Dermatitis to Trigger Cytokine Expression.* J Invest Dermatol, 2016. **136**(11): p. 2192-2200.

102. Wickham, H. *ggplot2: Elegant Graphics for Data Analysis.*

. 2016; Available from: https://ggplot2.tidyverse.org.

103. Taylor, S.C., et al., *The Ultimate qPCR Experiment: Producing Publication Quality, Reproducible Data the First Time.* Trends Biotechnol, 2019. **37**(7): p. 761-774.