Distinct Contributions of Nonpredictive and Predictive Peripheral Cues to Visual

Conscious Perception: an MEG study

Alfredo Spagna[1,2,*], Dimitri J. Bayle[3], Zaira Romeo[4], Lydia Yahia-Cherif[2], Ana B. Chica[5],

Paolo Bartolomeo[2,*]

[1] Department of Psychology, Columbia University in the City of New York, NY, USA, 10027

[2] Sorbonne Université, Inserm U 1127, CNRS UMR 7225, Paris Brain Institute, ICM, Hôpital

de la Pitié-Salpêtrière, 75013 Paris, France

[3] Licae Lab, Université Paris Nanterre, Nanterre, France

[4] Department of General Psychology, University of Padova, Padova 35131, Italy

[5] Department of Experimental Psychology; Mind, Brain, and Behavior Research Center

(CIMCYC), University of Granada, Granada, Spain, 18071

* Corresponding authors

Alfredo Spagna, Ph. D.

Department of Psychology,

Columbia University in the City of New York,

NY, USA, 10027

as5559@columbia.edu

Paolo Bartolomeo, M.D., Ph.D.

Sorbonne Université, Inserm U 1127,

CNRS UMR 7225, Paris Brain Institute,

ICM, Hôpital de la Pitié-Salpêtrière,

75013 Paris, France

paolo.bartolomeo@icm-institute.org

Number of pages: 39

Number of Figures (6) and Tables (1)

Number of words for abstract (208/250), introduction (648/650), discussion (1227/1500)

Conflict of interest statement (yes)

Author Contributions (yes)

Funding (yes)

Abstract

Do we need attention to become aware of an external event? We used

magnetoencephalography (MEG) in human participants to assess the effects of nonpredictive

and predictive supra-threshold peripheral visual cues on the conscious perception of near-

threshold Gabor patches. Both nonpredictive and predictive valid cues increased the number

of detected targets, and shifted the response criterion towards a more liberal decision.

Predictive cues unexpectedly induced a greater sensitivity (d') for invalid trials than for valid

trials. With nonpredictive cues, seen targets were associated with right-lateralized

frontoparietal feed-forward and feedback sweeps. For seen targets there was increased

connectivity among visual regions, and between these areas and the inferior parietal lobes and

the anterior insular cortices (AIC), bilaterally. Valid predictive cues interacted with conscious

target detection, with greater activation of areas mostly located in the left hemisphere,

especially in the frontoparietal network and temporoparietal junction, and induced an

increased connectivity between the right AIC and areas of the visual ventral stream in the

seen condition only. Thus, neural activity induced by nonpredictive and predictive spatial

cues can enhance conscious visual perception through distinct mechanisms, mostly relying on

frontoparietal activity in the right or left hemisphere, respectively. Connectivity involving the

AIC participates in shaping the interaction between attention and conscious visual perception.

51   **Significance Statement**

Do we need to pay attention to external objects in order to become aware of them?

Characterizing the spatiotemporal dynamics of attentional effects on visual perception is

critical to understand how humans process information coming from relevant aspects of their

environment. Participants detected near-threshold visual targets preceded by supra-threshold

spatial cues with varying degrees of predictivity, while their brain activity was recorded using

magnetoencephalography. Results demonstrated that spatial cues, especially when predictive,

biased participants' conscious perception through an early recruitment of frontoparietal

regions. This work highlights an interactive pattern between spatial attention and

consciousness, as shown by the effects of attention-related regions on visual sensory cortices

bilaterally, consistent with the hypothesis that attention is a pathway to conscious perception.

Introduction

The relationship between attention and consciousness remains debated. Some argue

that conscious perception cannot occur without attention (Posner, 1994; Dehaene et al., 2006;

Mashour et al., 2020); others propose that attention and consciousness are dissociable

processes (Wyart and Tallon-Baudry, 2008; Tallon-Baudry, 2012). In spite of the growing

effort devoted to characterizing the neural substrates supporting visual conscious perception,

this quest is still open, possibly because attention is a heterogeneous psychological construct

consisting of independent but interactive functions, which may differentially influence

conscious visual perception.

Studies investigating how attention modulates conscious perception mainly focused on

visuospatial orienting, which enhances the processing of stimuli appearing in a specific region

of the visual field (Posner, 1980). Nonpredictive visual cues, which are not informative about

the future location of the target, exogenously capture attention (Fan et al., 2002; Funes et al.,

2007; Chica et al., 2013b; Spagna et al., 2015). For short stimulus-onset asynchronies (SOAs)

a benefit in performance often occurs when the cue and the target appear on the same side

(so-called "valid" trials), compared to when they occur on opposite sides ("invalid" trials).

More endogenous, or goal-driven forms of orienting occur with predictive cues that correctly

indicate the location of the upcoming target in most trials. Neuroimaging studies have shown

partially distinct neural processes underlying exogenous and endogenous orienting (Corbetta

and Shulman, 2002; Fan et al., 2005; Chica et al., 2013b; Xuan et al., 2016), but the

spatiotemporal dynamics of these mechanisms remain unclear, possibly due to insufficient

temporal resolution.

Longstanding evidence indicates that nonpredictive orienting cues improve conscious

perception (Chica et al., 2011b; Sergent et al., 2013). However, the effects of endogenous

orienting remain more controversial (Ling and Carrasco, 2006; Koch and Tsuchiya, 2007;

Kentridge et al., 2008; Liu et al., 2009; Van Boxtel et al., 2010; Tsuchiya and Koch, 2014,

2016; Botta et al., 2017; Vernet et al., 2019). Peripheral predictive cues engage a mix of

exogenous and endogenous orienting mechanisms, because they initially capture attention

exogenously, but as time passes attention is endogenously kept at the cued location. This

endogenous persistence is apparently strategic, because it is more likely that the target appears

at the cued location (but see Bartolomeo et al., 2007). A series of studies (Chica et al., 2011b;

Chica et al., 2013b; Botta et al., 2017) showed that exogenous attention affects perceptual

consciousness, and that the maximal attentional effect on conscious perception is observed

when attention is first exogenously captured, and then endogenously maintained at the target

location. Characterizing the spatial and temporal dynamics of the neural substrates supporting

possible interactions between attentional orienting and visual conscious processing could

reduce the tension between competing, but not necessarily mutually exclusive, theoretical

frameworks.

Here, we addressed two questions: (1) What are the spatiotemporal dynamics of the

effects of peripheral cues on visual conscious processing? (2) How does cue predictivity

modulate these effects? We recorded magnetoencephalography, capitalizing on its unique

capacity to characterize a wide range of neural dynamics (Baillet, 2017), while participants

performed a version of a Posner-type cueing paradigm (Chica et al., 2014b) with supra-

threshold peripheral cues and near-threshold Gabor targets. In different experiments, spatial

cues were either nonpredictive or predictive of the site of occurrence of targets. This setting

enabled us to examine the effects of cues on conscious visual perception, in terms of

behavioral effects, neural activity, and brain connectivity.

The gateway hypothesis would be supported by findings of increased reportability of

the target stimuli under conditions of increased attention (i.e., valid cues) (Posner, 1994),

perhaps with larger effects for predictive than for nonpredictive cues (Chica et al., 2011b),

and a possible involvement of frontoparietal attention networks in the interaction between

attention and consciousness (Chica et al., 2013c). The absence of cue validity effects on

conscious perception would instead favor the cumulative influence hypothesis (Wyart and

Tallon-Baudry, 2008; Tallon-Baudry, 2012), together with an absence of interaction between

attention and consciousness in frontoparietal activations.

Materials and Methods

Participants. To estimate the sample size required to detect a difference in d' for valid and

invalid cue trials, we conducted a statistical power analysis using the parameters estimated in

our previous study (Chica et al., 2011b). With alpha = 0.05, an expected power of 0.80 and an

effect size of 0.81, the projected sample size needed was of n = 15 (two-tailed). We also

conducted a statistical power analysis to estimate the sample size required to detect a

difference in *criterion* for valid and invalid cue trials in the experiment with predictive cues.

With alpha = 0.05, an expected power of 0.80, and the effect size of 1.53 as shown in

previous research (Chica et al., 2011b), the projected sample size needed was of n = 6 (two-

tailed).

In total, 37 participants were recruited across two experiments. Eighteen participants

completed the experiment with nonpredictive cues (age = 24 ± 3.13 years; age range = 22-33

years; 6M), and nineteen participants completed the experiment with predictive cues (age =

24 ± 3.79 years; age range = 20-32 years; 7M). Five participants had to be excluded from data

analysis of the predictive cue experiment, due to issues in the data quality of MEG recordings.

All participants reported normal or corrected-to-normal vision, and gave written informed

consent before participation. The study was approved by the INSERM and by the Institutional

Review Boards of Paris Ile de France (CPP 1).

Stimuli and Procedure. The tasks were compiled and run using E-Prime software (RRID:

SCR_009567; Psychology Software Tools, Pittsburgh, PA) on a Windows XP desktop

computer. All stimuli were presented on a gray background at the center of a black projection

screen using a PROPixx projector (resolution, 1050×1400 pixels; refresh rate, 60 Hz) located

outside the shielded recording room. **Fig 1a.** shows a schematic representation of the

sequence of events in an invalid trial. **Fig 1b.** illustrates size of the stimuli used in our tasks.

The display consisted of three black boxes ($3.6° \times 4.9°$ of visual angle) presented on a gray

background; the central one was presented at the center of the screen and contained a fixation

point (a black cross) at its center. The other two boxes were located 6° of visual angle to the

left and right side and 4° of visual angle below the central box, respectively, a setting created

to maximize MEG responses from early visual areas (Portin et al., 1999).

146   Participants sat in the MEG recording room, with the screen being positioned

approximately 80 cm away from their eyes, and performed two tasks sequentially: 1) a

discrimination task, which required to identify the orientation of a Gabor patch (spatial

frequency: 5 cycles per degree of visual angle; diameter: 2.5° of visual angle; orientation:

chosen among 12 equally spaced between 0 and 180°, vertical and horizontal orientations

being excluded) that was presented for 16ms in either the box to the left or to the right side of

the display. After a 484ms delay, participants were asked to press a button on a response box

with 3 vertical buttons to indicate the orientation of the grating among two possibilities

presented vertically on the screen, distant by 3° from each other. Participants pressed the

upper response button with their index finger to choose the upper orientation or the middle

response button with their middle finger to choose the lower orientation. The location of the

correct orientation was randomized. After the participants' response, or after 3s without

response, 2) a detection task was presented, which required to press one of the three buttons

of the response box to indicate whether the target was absent, or whether it had been

presented in the left or right box. Two arrow-like stimuli (>>>>>> or <<<<<<) were

presented above and below the fixation cross, their respective position being randomized

across trial, and the word "absent" presented under the arrow-like stimuli. For trials in which

participants reported to have seen a stimulus, they then pressed the upper or lower response

button (with their index or middle finger, respectively) to indicate the visual hemifield of the

target presentation. For trials in which participants reported to have not seen the stimulus,

they pressed the lower response button with their ring finger. After the participants' response,

or after 3s without response, the next trial began after a variable delay of 1 to 1.5 s.

Before the recording session, participants were briefly instructed about the goal of the

study and were then shown instructions on the screen. Each participant underwent a

calibration session (mean duration, 6 min), during which the target contrast was manipulated

in order to estimate the individual threshold for which the percentage of consciously

perceived target was 50%. The calibration session consisted of two randomly inter-leaved

psychophysical staircases (one-up / one-down), theoretically converging toward a detection

rate of 50%. During the calibration session, participants were engaged in the same paradigm

as described previously, except that the contrast of the stimuli was varied from trial to trial

depending on their previous seen – unseen report in the corresponding staircase. Threshold

contrasts were estimated separately for the valid and invalid locations. The calibration session

was followed by eight recording sessions (mean duration, 8 min per session).

Each trial started with a fixation display, whose duration varied randomly between

1,000ms and 1,500ms. In the target-present trials, a cue occurred 300ms before targets, in the

form of a black dot with a 1° diameter, and presented for 50ms near the external upper corner

of one of the two peripheral boxes. Such a small cue was used in order to avoid possible

phenomena of perceptual interference with the subsequent, near-threshold target.

Experimental designs and statistical analyses. In the nonpredictive cue experiment,

each of the eight MEG recording sessions consisted of 110 trials, including 88 stimulus-

present trials (in which stimuli at threshold contrast were presented either in the left or right

lower visual quadrants) and 22 stimulus-absent trials (in which no stimulus was presented).

The total number of trials was 880, with 50% valid cue trials (352 trials), 50% invalid cue

trials (352 trials), and 176 catch trials. Fifty percent of the targets were presented at the cued

location (valid cue condition); the other 50% was presented at the uncued location (invalid

cue condition) (**Fig. 1c**). Trials within a recording session were presented in a different

randomized order for each subject.

In the experiment with predictive cues, parameters of stimulus size and timing of

presentation were the same as those used in the nonpredictive cue experiment, except that the

total number of trials on each of the eight MEG recording sessions consisted of 784 trials,

with 67% valid cue trials (448 trials), 33% invalid cue trials (224 trials), and 112 catch trials

(**Fig. 1d**).

Figure 1 a. Schematic representation of the sequence of events in an invalid-cue trial. **b.** Size of the stimuli and

exact location of presentation on the screen. The experiment with nonpredictive cues (*NonPredCue*) and the

experiment with predictive cues (*PredCue*) shared the same sequence of event and size of stimuli; **c.** in the

NonPredCue experiment 50% of cues were valid and 50% were invalid; **d.** in the *PredCue* experiment 67% of

203   cues were valid and 33% were invalid.

MEG recordings. Continuous MEG recordings were conducted at the CENIR

(http://www.cenir.org) with an ELEKTA Neuromag TRIUX® machine (204 planar

gradiometers and 102 magnetometers) located in a magnetically shielded room with a

sampling frequency rate of 1kHz and a bandwidth ranging from of 0.01 to 300 Hz. The

recordings were then MaxFiltered (v2.2) (Taulu and Simola, 2006) to attenuate environmental

noise, Signal Space Separation (SSS) was then implemented, automatic detection of bad

channels was conducted, data were filtered (1 to 250 Hz), and resampled at a rate of 250Hz,

and then converted in the Fieldtrip structure (RRID: SCR_004849;

http://www.fieldtriptoolbox.org/) (Oostenveld et al., 2011) to conduct further preprocessing

and analytic steps. Cardiac activity (electrocardiogram – ECG), vertical and horizontal EOG

signals were also recorded together with the electrophysiological data. The exact timing of the

presentation of the stimuli onset was corrected in accordance to the signal received from a

photodiode located in the MEG room, in order to adjust to the delay produced by the refresh

rate of the projector.

Preprocessing and Artifact Rejection. Additional preprocessing steps were conducted using

Fieldtrip and included an initial visual inspection of the recordings conducted by two of the

authors (D.J.B. and Z.R.) to exclude segments with artifacts and ensure data quality control.

Electroculogram (EOG) recordings from both vertical and horizontal sensors were then used

to reject trials in which eye movements (beyond 3º) occurred. Rejection thresholds for both

horizontal and vertical EOG traces was set to $\pm$ .66V, corresponding to a deviation greater

than 3º of visual angle (and with the target at 6º of visual angle). Trials with excessive eye

movements and eye blinks (~10.52% of trials) were rejected offline from the MEG traces

according to the 3º threshold mentioned above. Signal from the photodiode was used to

discard 1) trials with a delay between the trigger and the photodiode greater than 300ms; 2)

trials with a delay between the cue and the target greater than 827ms; 3) trials in which the

delay between the trigger of the cue and the photodiode was greater than 40ms or smaller than

30ms, for a total of ~1% of the trials. Last, trials contaminated by muscular activity (jump or

movement) were rejected manually upon visual inspection (~15%). For the nonpredictive cue

experiment, out of the 15,840 trials acquired, 8517 trials were analyzed (*right visual field*:

seen invalid = 1033; seen valid = 1284; unseen invalid = 1056; unseen valid = 882; *left visual*

field: seen invalid = 1191; seen valid 1176; unseen invalid = 956; unseen valid = 939). For the

predictive cue experiment, out of the 10,796 trials acquired in total, 7454 trials were analyzed

(*right visual field*: seen invalid = 396; seen valid = 1588; unseen invalid = 812; unseen valid =

1379; *left visual field*: seen invalid = 639; seen valid 1144; unseen invalid = 576; unseen valid

= 1379) (see **Table S1** for a subject-by-subject breakdown of the number of trials in each

condition remaining after artifact rejection).

Event-Related Magnetic Fields. Data from 102 neuromag channels was analyzed in this study.

A Matlab® script was used to separate the MEG continuous recordings into 2300ms-long

epochs (ranging from -1000 before the cue and 1300ms after the cue), and epochs from the

eight experimental conditions from each participant were then imported into Brainstorm

(Tadel et al., 2011). For each condition, event-related magnetic fields were then averaged

(weighted) along their entire length (2300ms).

Source reconstruction. Signal amplitude from the 15,000 cortical elemental dipoles

underlying the signals measured by the sensors were then estimated from the epochs using the

weighted minimum norm estimation (wMNE) imaging method as implemented in Brainstorm

(Tadel et al., 2011), which first identifies a current source density image fitting the data

through the forward model, and then favors solutions that are of minimum energy by using

source covariance as a prior. To use this method, a noise covariance matrix was estimated for

each subject from the recordings using the pre-stimulus interval (-1,000 to -2ms before the

presentation of the cue), while constrained source covariance model was used to model one

dipole, oriented normally to the surface. Further processing conducted on the sources per

participant consisted of z-score transformation of the signal with reference to the baseline

(from -1,000 to -2ms). A spatial smoothing kernel (FWHM = 3mm) was then applied on the

sources, that were then re-interpolated (projected) on a common template (default anatomy).

MRI recordings. High-resolution T1-weighted structural MRI images (MPRAGE sequence,

flip-angle, 9; Repetition Time, 2300ms; Echo Time, 4.18ms; voxel size: $1 \times 1 \times 1$ mm) were

acquired for each participant using a 3-T Siemens, TRIO whole-body MRI scanner (Siemens

Medical Solutions, Erlangen, Germany) located at the CENIR MRI center (Salpetriere

Hospital, Paris, France). After acquisition, images were then segmented using the FreeSurfer

"recon-all" pipeline (Fischl, 2012), and imported in Brainstorm (Tadel et al., 2011) for co-

registration purposes. MEG sensors and structural MRI images were first manually aligned

using the nasion/left ear/right ear (NAS/LPA/RPA) fiducial points recorded in the MEG file

and in the MRI MNI coordinates. Co-registration was then further refined using the "refine

using head points" option on Brainstorm, which uses an iterative closest point algorithm to fit

the head shape and the digitized scalp points. Additional details about the MRI-MEG co-

registration steps as done in Brainstorm can be found here (Tadel et al., 2019).

Behavioral Data Analysis. Response times (RTs) below 150ms (anticipated responses) and

above 1,800ms (delayed responses) were excluded from subsequent analyses (below 2% of

total number of trials). Independent sample *t*-tests were used to assess potential differences in

the thresholds sampled during the calibration sessions separately for invalid and valid cue

trials. Analysis of Variance (ANOVA) and Signal Detection Theory (SDT) analyses were

conducted to examine whether and how nonpredictive and predictive cues modulated visual

conscious perception. The Kolmogorov-Smirnov and the Shapiro Wilk test for normality were

conducted to examine whether the data on accuracy was normally distributed. For both RT

and Accuracy (in percentage), a 2 (seen, unseen) $\times$ 2 (valid, invalid) repeated measures

ANOVA was conducted on correctly discriminated responses. These analyses were conducted

to examine behavioral differences between trials reported as seen or unseen under the valid

and invalid cue conditions. SDT was conducted to estimate changes in the signal to noise ratio

as a function of cue validity condition (*d'*), and to investigate whether the presence of a cue

could bias the observer towards a more liberal reporting threshold (*criterion*). For the *d'*,

participants' performance was estimated using the standardized difference between target-

present (Hits: $z(H)$) and target-absent (false alarms: $z(F)$) trials, by diving the percentage of

correctly detected trials by the number of false alarms (FA: trials in which participants

reported having seen the stimulus); zero false alarm rates (n = 12) were corrected using the

following formula (FA = (FA + 0,5) / (FA + CR + 1,0)) (Snodgrass and Corwin, 1988). The

greater the value of *d'*, the higher the quantity between the noise and the signal distribution

(i.e., the signal/noise ratio). The *criterion (C)* summarizes the distance of the threshold

relative to the noise distribution *B* from the threshold of an ideal observer (*-0.5 * ([z(H) +*

z(F)])). A negative value of *C* represents a more liberal threshold, while a positive value of *C*

represents a more conservative threshold. Both *d'* and *C* were estimated separately for valid

and invalid trials, and paired sample *t*-test was used to assess differences in *d'* and *criterion*

between these two conditions.

MEG data analysis. In parallel with the behavioral analyses, 2 (seen, unseen) $\times$ 2 (valid,

invalid) $\times$ 2 (left, right) repeated-measures ANOVAs were conducted on individuals'

estimated activity (z-scores) of reconstructed source images to examine how neuronal activity

induced by nonpredictive cues can enhance conscious perception. Spatiotemporal cluster-

based permutation tests (Maris and Oostenveld, 2007) were used for comparing main effects

and interactions between brain activations in the time window between 0 and 800ms (locked

to cue onset), with the number of permutations set to 1000 and the alpha threshold level set to

0.05 for all tests.

Connectivity Analysis. To examine the modulation in connectivity between brain areas as a

result of the experimental conditions, we estimated leakage-controlled (orthogonalized) high-

gamma (60 to 90 Hz) amplitude envelope correlation (AEC) (Brookes et al., 2012; Colclough

et al., 2015) of eighteen regions of interest (ROIs). Analyses were conducted solely on the

high-gamma frequency band because of the extensive knowledge on the role of the oscillatory

synchronization in this frequency band in spatial attention and its modulation of visual areas

(Tallon-Baudry et al., 2005; Magazzini and Singh, 2018; Fiebelkorn and Kastner, 2019).

ROIs were selected on the basis of previous results (Corbetta and Shulman, 2002; Fan et al.,

2005; Tallon-Baudry et al., 2005; Bressler et al., 2008; Chica et al., 2013c; Chica et al.,

2016b; Xuan et al., 2016), showing their involvement in attention-related processes. In

addition, areas of the ventral visual stream were also added to the ROI analyses to examine

potential frontoparietal modulation of activity in early and late visual areas. The ROIs were

created using Brainstorm (Tadel et al., 2011), seed vertices and area (in cm^2) of the ROIs are

listed in **Table 1** (in MNI coordinates). Paired sample *t*-tests ($p < .05$, FDR corrected) were

used to compare the AEC estimates across experimental conditions.

Results

Does the validity of nonpredictive cues affect participants' response to visual Gabor

targets? To answer this question, we examined behavioral responses to seen and unseen

targets separately for valid and invalid nonpredictive cues. Valid cues, compared to invalid

cues, lowered the response criterion (mean $\pm$ SD valid: 0.85 ± 0.28; invalid: 1.03 ± 0.27; $t(17)$

$= -2.82$; $p < .05$), and increased the percentage of detected targets (mean $\pm$ SD valid: $0.57 \pm$

0.09; invalid: 0.52 ± 0.11; $t(17) = -3.08$; $p < .01$). However, sensitivity (*d'*) was similar for

valid and invalid trials (valid: 2.30 ± 0.57; invalid: 2.29 ± 0.47; $t(17) < 1$) (see **Fig 2**). These

results did not depend on different perceptual thresholds resulting from the calibration

session, because these thresholds were similar for valid trials (43.98%) and for invalid trials

(43.99%; $t < 1$). Left- and right-sided targets evoked a similar number of correct responses

(sum of the total number of correctly discriminated targets presented to the left visual field:

3,489; mean $\pm$ SD: 194.83 $\pm$ 37.05; sum of the total number of correctly discriminated targets

presented to the right visual field: 3,507; mean $\pm$ SD: 194.83 $\pm$ 33.6; Wilcoxon signed-rank

test = 76; $p = .70$; Bayesian Wilcoxon signed-rank test $BF_{10} = 0.56$, with median posterior $\delta =$

-0.079, 95% CI [-1.07, 0.91]).

As expected, participants were more accurate in discriminating the orientation of *Seen*

targets (79.2 $\pm$ 14.0%) than that of *Unseen* targets (50.2 $\pm$ 7.0%) [ANOVA with

Consciousness (seen, unseen), *Visual Field* (left, right), and *Validity* (valid, invalid) as factors,

$F_{(1,17)} = 85.60$; $p < .0001$; $\eta^2 = .84$] (**Fig 2a**). No other factors or interactions reached

statistical significance.

A similar ANOVA conducted on RTs (see **Fig 2b**) revealed a main effect of

Consciousness ($F_{(1,17)} = 102.65$; $p < .0001$; $\eta^2 = .86$), because participants were slower for

Seen targets (1034 $\pm$ 115ms) than for *Unseen* targets (601 $\pm$ 190ms). No other factors or

interactions reached significance.

Figure 2. Results of the behavioral analyses conducted on performance with nonpredictive cues (a.-e.) and with predictive cues (f.-j.). **a.** and **b.** report result of the ANOVA conducted on Accuracy (in percentage) and RTs (in ms), respectively, while **c.**, **d.**, and **e** report result of the Signal Detection Theory analysis for d', criterion, and detection (in percentage). **f.** and **g.** report result of the ANOVA conducted on Accuracy (in percentage) and RT (in ms), respectively, while **h.**, **i.**, and **j.** report results of the SDT analysis for d prime, criterion, and detection (in percentage).

Does the validity of nonpredictive cues affect neural responses associated with the

conscious report of visual Gabor targets? We addressed this question by examining brain

responses to seen and unseen targets separately for valid and invalid nonpredictive cues.

Source analysis of the MEG signal revealed that conscious perception was associated with

right-lateralized frontoparietal feed-forward and feedback sweeps. Two clusters exceeding the

threshold of randomization distribution under H0 emerged for the seen *vs* unseen comparison

(both *p*s < 0.001) in the time window of 400 - 800ms after cue onset. The first cluster was in

the right hemisphere, the second in the left hemisphere. Both clusters started in the occipital

cortex and afterwards extended to the frontoparietal network and temporal regions, bilaterally

(see **Fig 3**). The differences for the main effects of *Validity* (valid, invalid), for the main effect

of *Visual Field* (left, right), and for the interactions did not reach statistical significance.

Control analyses showed that there was no significant difference between the number of MEG

trials for left- and right-sided targets (left visual field, mean $\pm$ SD: 235.39 $\pm$ 32.49; right

visual field, 235.5 $\pm$ 30.02; Wilcoxon signed-rank test = 85; *p* = 1; Bayesian Wilcoxon

signed-rank test BF_{10} = 0.57, with median posterior δ = -0.05, 95% CI [-1.09, 0.96]).

Figure 3. a. When preceded by nonpredictive cues, seen targets evoked two clusters of brain

activity compared to unseen targets. Cluster 1 occurred in the 400 - 800ms time window after

cue onset, was lateralized to the right hemisphere, and encompassed a frontoparietal *feed-*

forward and *feedback sweeps* (around 456ms), with subsequent diffusion to widespread

bilateral activation. Cluster 2 occurred also in the 400 - 800ms time window after cue onset,

but was lateralized to the left hemisphere, and encompassed a widespread brain activation. **b.**

Average signal changes in the ROIs separately for the Seen (solid line) and Unseen (dashed

line) condition. The area in green highlights the time interval in which cluster-corrected

analysis showed a significant difference between the two signals.

Does the validity of nonpredictive cues modulate the connectivity between ROIs associated

with the conscious report of visual Gabor targets? To answer this question, we examined

functional connectivity associated to seen and unseen targets separately for valid and invalid

nonpredictive cues. Whole-trial (0 – 800ms) high-gamma band (60 - 90Hz) amplitude

envelope correlation analyses conducted among attention-related ROIs (see **Table 1**) revealed

a strong connectivity within visual ventral areas and between these areas and the anterior

insular cortex (AIC), bilaterally for *Seen* compared to *Unseen* trials ($p < .001$, FDR corrected;

Fig 4a). There was no evidence for a difference in connectivity between *Valid* and *Invalid*

trials. Differences ($ps < .005$, FDR corrected; **Fig 4b** and **c**) emerged in the connectivity

between *Seen Valid* and *Unseen Valid* trials across areas within the visual ventral stream as

well as with the right inferior parietal lobule (IPL) and with the right AIC. There was also

increased connectivity of the right AIC with areas in the visual ventral cortical stream and in

Seen Invalid trials than *Unseen Invalid* trials.

Figure 4. Pairwise functional connectivity (estimated using amplitude envelope correlation, AEC, of the whole duration of the trials – 0 to 800ms) between ROIs, with weighted edges denoting significant *t*-score values (overall *ps* < .005, FDR corrected) obtained from the contrast between the conditions of interest. **a.** difference in AEC between *Seen* and *Unseen* trials in the nonpredictive cue (*NonPredCue*) experiment; **b.** difference in the AEC between *Seen Valid* and *Unseen Valid* in the *NonPredCue* experiment; **c.** difference in the AEC *Seen Invalid* and *Unseen Invalid* in the *NonPredCue* experiment; **d.** difference in the AEC *Seen Valid* and *Seen Invalid* in the predictive cue (*PredCue*) experiment.

Thus, the validity of nonpredictive cues modulated participants' conscious reports of visual Gabors for both behavioral performance and MEG connectivity measures. Differences also emerged between seen and unseen trials that were not confined to the occipital lobe, but were quickly followed by frontoparietal activity (approximately 150ms after the target and 50ms after the occipital activation), and subsequently by a widespread activity that lasted to the end of the trial, and resembled the pattern described as the Global Neuronal Workspace (Mashour et al., 2020).

Does the validity of predictive cues affect participants' response to visual Gabor targets? To

address this issue, we examined behavioral responses to seen and unseen targets separately for

valid and invalid predictive cues. As did nonpredictive cues, also valid predictive cues,

compared to invalid predictive cues, lowered the response criterion (mean ± SD valid: (0.85 ±

0.28; 1.26 ± 0.15; (t(13) = -6.37; p <.001), because participants adopted a more liberal

response threshold in the valid cue condition, and increased the percentage of detected targets

(mean ± SD valid: 54.11 % ± 7.13%; invalid: 43.33 ± 5.14%; t(13) = 8.06; p <.001).

However, *sensitivity* (*d'*) was greater for invalid trials (2.20 ± 0.33) than for valid trials (1.91

± 0.54) (t(13) = -2.29; p < .05) (see **Fig 2**). These results did not depend on possible

differences between the thresholds established separately for valid and invalid trials cue trials

during the calibration session (valid trials: 59.21; invalid trials: 60.79; t<1). Unexpectedly,

right-sided targets evoked more correct responses that left-sided ones (right visual field, mean

± SD: 220.21 ± 35.42; left visual field, 194.72 ± 21.89; Wilcoxon signed-rank test = 6.00; p <

.01; Bayesian Wilcoxon signed-rank test BF_{10} = 4.06, with median posterior δ = -1.19, 95%

CI [-2.51, 0.01]).

Results of the ANOVA conducted on Accuracy percentages to the discrimination task

(after ensuring that these percentages followed a normal distribution by using the

Kolmogorov-Smirnov and the Shapiro Wilk tests for normality) are shown in **Fig 2f**. There

was a main effect of the factor *Consciousness* ($F_{(1,13)}$ = 31.62; p < .0001; η^2 = .71), because

participants were more accurate for *Seen* trials (72.87 ± 15.2%) than for *Unseen* (49.71 ±

5.9%) trials. No other effect or interaction reached significance, except for the interactions

between *Consciousness* and *Validity* ($F_{(1,13)}$ = 11.9; p < .01; η^2 = .48) and between *Visual*

Field and *Validity* ($F_{(1,13)}$ = 8.89; p < .05; η^2 = .41). Pairwise comparisons showed that the

difference between valid and invalid trials was significant for the seen trials (valid: 74.88 ±

14.41%; invalid: 70.86 ± 16.14%; p < .01), but not for the unseen trials (valid: 48.91 ± 5.13%;

invalid 50.51 ± 6.52%; $p = .14$). Participants detected more validly cued targets than invalidly

cued targets in the right visual field (valid: 64.24 ± 9.92%, invalid: 60.08 ± 12.26%, $p < .01$),

but not in the left visual field (valid: 59.56 ± 9.62%; invalid: 61.29 ± 10.04% $p = .21$).

432   Results of the ANOVA conducted on the RTs (see **Fig 2g**) showed that participants

responded slower to *Seen* targets (1048 ± 171ms) than to *Unseen* ones (640 ± 259ms; $F_{(1,13)} =$

28.91; $p < .0001$; $\eta^2 = .69$). The main effects of *Visual Field* and *Validity* did not reach

significance (both Fs < 1). *Consciousness* interacted with *Visual Field* ($F_{(1,13)} = 5.08$; $p < .05$;

$\eta^2 = .28$), because the seen/unseen difference was greater for left-sided targets (seen: 1,055 ±

168ms; unseen: 624 ± 249ms; $p < .001$) than for right-sided targets (seen: 1,040 ± 177;

unseen: 656 ± 272; $p < .001$).

Does the validity of predictive cues affect neural responses associated with the conscious

report of visual Gabor targets? We addressed this question by examining brain responses to

seen and unseen targets separately for valid and invalid predictive cues. Similar to the

experiment with nonpredictive cues, two clusters exceeding the threshold of randomization

distribution under H0 emerged for the seen *vs* unseen comparison ($ps < 0.001$), both in the

time window of 460 - 760ms after cue onset. The first cluster was lateralized to the right

hemisphere, the second was lateralized to the left hemisphere. Both started in the occipital

cortex and then spread to the parieto-temporal regions, without any evidence of involvement

of prefrontal regions. *Consciousness* interacted with *Validity*. Four clusters exceeding the

threshold of randomization distribution under H0 were found for the interaction term ((seen

valid *minus* unseen valid) *minus* (seen invalid *minus* unseen invalid)). The first cluster ($p <$

0.001) occurred in the 100 - 150ms time window after cue onset, and was lateralized to the

left visual cortex; the second cluster ($p < 0.05$), was also lateralized to the left visual cortex

and left IPL, in the 300 - 380ms time window after cue onset; the third cluster ($p < 0.05$)

occurred in the inferior temporal lobe of the right hemisphere and spanned across the time

window of 550 and 600ms after cue onset; the forth cluster ($p < 0.05$) was localized in the left

precuneus, extending into the left posterior cingulate cortex and spanned across the time

window of 580 and 600ms after cue onset. **Fig 5** displays the follow-up analysis conducted in

the valid *vs* invalid comparison for the seen condition only (seen valid *minus* seen invalid).

Three clusters exceeded the threshold of randomization distribution under H0. A first cluster

($p < 0.001$) was lateralized to the left hemisphere and occurred in the 50 - 400ms time

window after cue onset, spanning across visual cortex, to the parietal cortex (around 100ms

post-cue), prefrontal cortex (around 150ms post-cue), and temporoparietal cortex. A second

cluster ($p < 0.05$), occurred later, as a right-hemisphere occipito-parietal activation around

200ms post-cue, and then substantially resembled the activity found in the first cluster,

spanning across the frontoparietal cortex (around 240ms post-cue), and temporoparietal

cortex (around 280ms post-cue). The third cluster ($p < 0.05$) was lateralized to the left visual

cortex and included middle- and infero-temporal areas, the inferior frontal gyrus (around

470ms post-cue), the left frontal eye fields (524ms post-cue) and then temporoparietal,

inferior parietal, and inferior temporal areas.

Control analyses showed that there was no significant difference between the number

of MEG trials for left- and right-sided targets (left visual field mean ± SD: 265.43 ± 21.16;

right visual field; mean ± SD: 266.79 ± 26.91; Wilcoxon signed-rank test = 48.5; $p = 0.83$;

Bayesian Wilcoxon signed-rank test $BF_{10} = 0.58$, with median posterior $\delta = -0.073$, 95% CI [-

1.11, 0.94]).

Figure 5. When preceded by valid predictive cues, seen targets evoked three clusters of brain

activity, compared to invalidly cued targets. **a**) Cluster 1 occurred in the 50-400ms time

window after cue onset and was lateralized to the left hemisphere ($p < 0.001$); Cluster 2

occurred in 200ms time-window after cue onset and was lateralized to the right hemisphere (p

< 0.05); Cluster 3: occurred 470ms after cue onset and lateralized to the left hemisphere. The

same valid-invalid comparison for unseen targets did not reveal any significant cluster. **b)**

average signal changes in the ROIs separately for the Seen Valid (solid line) and Seen Invalid

(dashed line) condition. The area in green highlights the time interval in which cluster-

corrected analysis showed a significant difference between the two signals.

Does the validity of predictive cues modulate the connectivity between ROIs associated with

the conscious report of visual Gabor targets? We examined functional connectivity

associated to seen and unseen targets separately for valid and invalid predictive cues. High-

gamma band (60 - 90Hz) amplitude envelope correlation analyses conducted on the ROIs

revealed increased connectivity between the right AIC and the right temporo-occipital area for

Seen Valid vs Seen Invalid trials (see Fig 4d). The comparisons between connectivity patterns

in the *Seen Valid vs Unseen Valid* and the comparison between connectivity patterns in the

Seen Invalid vs Unseen Invalid did not yield statistically significant results.

Thus, similar to nonpredictive cues, also the validity of predictive cues modulated

participants' conscious reports of visual Gabors for both behavioral performance and neural

activity measures. Differences between seen and unseen trials with predictive cues emerged in

the post-target time period, but these were delayed in time compared to the same contrast with

nonpredictive cues. Bilateral activity started in the visual cortex, around 160ms after the

target, and was followed by mainly temporo-parietal activity, with prefrontal activity arising

only later in time (almost 300ms after the target). The interaction between attention and

consciousness showed four clusters of activation located in occipito-temporal regions that

spanned across both the cue-target and the post-target periods, possibly showing the *site* of

attentional modulation (Posner and Driver, 1992) rather than the *source* of it (Bressler et al.,

2008; Liu et al., 2016). The modulation produced by predictive cues on visual awareness was

also evident in the contrast between the functional connectivity patterns in valid and invalid

trials for seen targets, which again showed correlated activity in AIC and areas in the ventral

visual stream, bilaterally.

Cross-experiment comparison: different effects of nonpredictive and predictive cues on

visual conscious perception.

Did nonpredictive and predictive cues produce different effects on brain activity and

connectivity? Seen targets preceded by a nonpredictive cue evoked a significant cluster (Fig

6), occurring in the 210 - 300ms time window after cue onset, lateralized to the right

hemisphere, and encompassing an early activation in the middle and inferior frontal gyri, the

superior frontal gyrus and the temporoparietal junction (mainly located in the angular gyrus),

with subsequent diffusion to occipitotemporal regions (cluster-corrected Welch's t test

conducted in the cue-target period (0 – 300 ms), $p < .02$).

Figure 6. MEG activity evoked by nonpredictive vs. predictive cues for Seen vs. Unseen

targets in the cue-target period (0 – 300 ms). **a.** One right-hemispheric cluster emerged in the

210-300ms time window, and included the temporoparietal junction and the prefrontal cortex.

b. Average signal changes in the ROIs separately for the experiments with nonpredictive cue

(solid line) and with predictive cues (dashed line). The area in green highlights the time

interval in which cluster-corrected analysis showed a significant difference between the two

signals.

Whole-trial (0 – 800ms) high-gamma band (60-90Hz) pairwise functional connectivity

analyses conducted among attention-related ROIs (see Table 1) revealed a stronger

connectivity between the left TPJ and right fusiform gyrus ($p < .01$), the left anterior insular

cortex (AIC) and the left fusiform gyrus ($p < .05$), and between the left superior parietal lobe

(SPL) and the right IPL ($p < .05$) for the seen minus unseen contrast with nonpredictive cues,

as compared to same contrast with predictive cues.

Discussion

Our results establish the existence of interaction patterns between spatial attention and

consciousness that vary based on the predictivity of the cue, and specify how attention

modulates conscious perception. Altogether, the contributions of nonpredictive and predictive

cues in improving target detection indicate that distinct spatiotemporal dynamics support the

flexible allocation of attentional resources for near-threshold visual information to reach a

conscious level of processing.

Valid spatial cues influence conscious reports. Both predictive and nonpredictive valid

cues increased the proportion of correctly detected targets. These results align with behavioral

evidence showing the modulation of orienting cues in increasing visual awareness (Ling and

Carrasco, 2006; Liu et al., 2009; Chica et al., 2011b; Sergent et al., 2013; Botta et al., 2017;

Vernet et al., 2019). However, only predictive cues improved performance on the

discrimination task, while Chica et al. (2011b) found effects of both nonpredictive and

predictive cues in a similar task. One important methodological difference between our study

and the Chica et al. (2011b) study is that in the current experiments two supra-threshold

Gabor patches were presented 500ms after the target to collect the discrimination response.

This screen could have increased the perceptual difficulty of the task and therefore reduced or

abolished the effects of nonpredictive cues (Kerzel et al., 2009). Unexpectedly, valid

predictive cues decreased perceptual sensitivity. This result is at odds with previous

observations of increased perceptual sensitivity (Chica et al., 2011b) and contrast sensitivity

(Carrasco, 2018) at attended locations with predictive cues. Possible masking effects caused

by the post-target display might explain our finding. Validly cued targets might have suffered

interference from the preceding cue and the subsequent response Gabors, while invalid targets

were only masked by the response Gabors. Facilitatory effects induced by peripheral cues are

typically much larger for discrimination than for detection tasks (Lupiáñez et al., 1997;

Lupiáñez et al., 2001; Chica et al., 2006). Consequently, the possible increased difficulty of

detecting a near-threshold target, might have reduced the behavioral effects of spatial

attention. Peripheral cues (whether nonpredictive or predictive) produced a shift in the

response criterion towards a more liberal response in valid compared to invalid trials,

indicating that orienting cues modulated decision-related processes in our setting.

Distinct neural processes elicited by nonpredictive and predictive cues. MEG results

indicated distinct spatiotemporal dynamics across tasks, both at the source level and in the

functional connectivity patterns. Nonpredictive cues elicited a greater activation of the right

TPJ and right prefrontal cortex in the seen *minus* unseen contrast compared to predictive cues.

This network is connected by the ventral branch of the superior longitudinal fasciculus (SLF

III) (Thiebaut De Schotten et al., 2011), and involved in exogenous attention (Corbetta and

Shulman, 2002; Chica et al., 2011a). Interestingly, comparison of brain activity elicited by

nonpredictive *versus* predictive cues (irrespective of participants' responses), showed a right-

hemispheric TPJ-prefrontal cluster. The reduced TPJ activity for predictive cues compared to

nonpredictive cues is consistent with evidence of TPJ deactivation during endogenous

orienting (Shulman et al., 2007; Doricchi et al., 2010). Moreover, the combination of spatial

and temporal resolution of MEG results demonstrated that this reduced TPJ activation for

predictive as compared to nonpredictive cue was preceded by deactivation of the prefrontal

nodes of the SLF III network in the middle and inferior frontal gyri. The statistically higher

frequency of valid trials in the predictive cue experiment than in the nonpredictive cue

experiment might explain the occurrence of different preparatory states after cue presentation

between the two experiments. This possibility may account for the absence of such effects in

the task with nonpredictive cues, which did not induce any probabilistic expectations.

Together with the behavioral validity effect occurring with predictive cues, this result

suggests that cue predictivity enhances attentional modulation. Interestingly, while the

occipital activation was bilateral, the IPL-TPJ activations lasted longer (by around 100ms)

only in the right hemisphere, in line with abundant evidence on anatomical and functional

asymmetries in frontoparietal attention networks favoring the right hemisphere (reviewed in

Bartolomeo and Malkinson, 2019; see also Spagna et al., 2020b), and specifically with the

role of the right TPJ in attentional orienting (Doricchi et al., 2010; Chica et al., 2014a), and in

the processing of behaviorally relevant stimuli (Macaluso and Doricchi, 2013).

Frontoparietal networks and visual awareness. Dysfunction of SLF II and III

networks in the right hemisphere is a typical finding in brain-damaged patients with left visual

neglect, who have impaired awareness of left-sided events (Bartolomeo et al., 2012). The

frontoparietal activity found with nonpredictive cues seems consistent with the *forward* and

backward sweeps associated with the awareness of a visual target (Lamme, 2006), and

suggests a role for exogenous attention in this process. The relationship of these

forward/backward sweeps with behavioral changes in criterion, but not in sensitivity, might

be consistent with the known role of frontoparietal networks in the activation of general

processes, such as the alerting system (Périn et al., 2010; Chica et al., 2016a; Baria et al.,

2017; Petersen et al., 2017; Podvalny et al., 2019). An interaction of nonpredictive cues with

alerting is also supported by the functional connectivity analysis, which showed a role of the

right AIC in differentiating seen *vs.* unseen trials separately for valid and invalid trials (see

Fig 6). This result is consistent with evidence showing the critical role of the AIC in the

production and maintenance of alertness, both in terms of BOLD response and of functional

connectivity (Cai et al., 2014; Sadaghiani and D'Esposito, 2015; Chica et al., 2016a; Coste

and Kleinschmidt, 2016; Han et al., 2019; Haupt et al., 2019). As a node of the SLF III

network, the AIC may have an important role in signaling behaviorally relevant stimuli

(Uddin, 2015).

In conclusion, four results of this study are broadly consistent with the gateway

hypothesis (Posner, 1994), and demonstrate how the interaction between attention and visual

conscious processing heavily depends on the specific attentional component manipulated.

First, both predictive and nonpredictive visual cues modulated behavioral performance in the

form of a shift towards a more liberal response criterion. Second, peripheral predictive cues

were able to modulate visual conscious processing, as shown by the interaction with response

accuracy, in line with previous evidence (Chica et al., 2013a). Third, predictive cues also

interacted with consciousness at the neural level, and induced increased activity in visual

areas, possibly related to the site of attentional modulation. Last, high-gamma band functional

connectivity patterns involving the AIC and the visual areas for seen trials differed across task

conditions, with valid cues inducing greater modulation of the visual areas compared to

invalid cues.

It remains possible that attention is required for conscious processing only when there

is some competition between the stimuli to be resolved (Tsuchiya and Koch, 2014; Davidson

et al., 2018), which was the case in our setting with two possible target locations.

Alternatively, even an isolated stimulus might need some attentional capture to be

consciously processed. Evidence from visual mental imagery studies, showing the implication

of frontoparietal attention networks within the conscious imagination of an object in its

absence, and thus without any competition (Spagna et al., 2020a), might support this

possibility, which needs to be empirically assessed. Altogether, the comparison of the

spatiotemporal dynamics underlying the interaction between nonpredictive and predictive

attention with consciousness shown here confirmed that these distinct contributions span

across behavioral, neural, and connectivity measures, and underlines the role of the right AIC

in visual awareness.

Author Contributions

D.B., A.B.C., and P.B. designed the experiments; A.S., D.B., and Z.R. analyzed the data. All

authors discussed the results and contributed to writing up the report. Behavioral data and the

code to reproduce the figures (built using Python on Spyder) can be found on the GitHub page

of A.S. We thank Fabrizio Doricchi, Isabella Elaine Rosario, and Catherine Tallon-Baudry for

providing extensive comments on this manuscript.

Conflict of Interest Statement

The authors report no conflict of interest.
638

639   **Funding**
640
641   Funded by an ICM fellowship to A.S., by project eraNET-NEURON 2009 "BEYONDVIS" to

642   D.J.B and P.B.. P.B. is also supported by the *Agence Nationale de la Recherche* through

643   ANR-16-CE37-0005 and ANR-10-IAIHU-06, and by the *Fondation pour la Recherche sur*

644   *les AVC* through FR-AVC-017.

645
646
647

648 **References**
649
Baillet S (2017) Magnetoencephalography for brain electrophysiology and imaging. Nature
neuroscience 20:327.
Baria AT, Maniscalco B, He BJ (2017) Initial-state-dependent, robust, transient neural
dynamics encode conscious visual perception. PLoS Comput Biol 13:e1005806.
Bartolomeo P, Malkinson TS (2019) Hemispheric lateralization of attention processes in the
human brain. Current opinion in psychology 29:90-96.
Bartolomeo P, Decaix C, Siéroff E (2007) The phenomenology of endogenous orienting.
Consciousness and Cognition 16:144-161.
Bartolomeo P, Thiebaut De Schotten M, Chica AB (2012) Brain networks of visuospatial
attention and their disruption in visual neglect. Frontiers in human neuroscience
6:110.
Botta F, Ródenas E, Chica AB (2017) Target bottom-up strength determines the extent of
attentional modulations on conscious perception. Experimental brain research
235:2109-2124.
Bressler SL, Tang W, Sylvester CM, Shulman GL, Corbetta M (2008) Top-down control of
human visual cortex by frontal and parietal cortex in anticipatory visual spatial
attention. Journal of Neuroscience 28:10056-10061.
Brookes MJ, Woolrich MW, Barnes GR (2012) Measuring functional connectivity in MEG: a
multivariate approach insensitive to linear source leakage. Neuroimage 63:910-920.
Cai W, Ryali S, Chen T, Li C-SR, Menon V (2014) Dissociable roles of right inferior frontal
cortex and anterior insula in inhibitory control: evidence from intrinsic and task-
related functional parcellation, connectivity, and response profile analyses across
multiple datasets. Journal of Neuroscience 34:14652-14667.
Carrasco M (2018) How visual spatial attention alters perception. Cognitive processing
19:77-88.
Chica AB, Lupiánez J, Bartolomeo P (2006) Dissociating inhibition of return from endogenous
orienting of spatial attention: Evidence from detection and discrimination tasks.
Cognitive neuropsychology 23:1015-1034.
Chica AB, Bartolomeo P, Valero-Cabré A (2011a) Dorsal and ventral parietal contributions to
spatial orienting in the human brain. Journal of Neuroscience 31:8143-8149.
Chica AB, Bartolomeo P, Lupianez J (2013a) Two cognitive and neural systems for
endogenous and exogenous spatial attention. Behav Brain Res 237:107-123.
Chica AB, Bartolomeo P, Lupiáñez J (2013b) Two cognitive and neural systems for
endogenous and exogenous spatial attention. Behavioural brain research 237:107-
123.
Chica AB, Bourgeois A, Bartolomeo P (2014a) On the role of the ventral attention system in
spatial orienting. Frontiers in Human Neuroscience 8:235.
Chica AB, Paz-Alonso PM, Valero-Cabre A, Bartolomeo P (2013c) Neural bases of the
interactions between spatial attention and conscious perception. Cerebral Cortex
23:1269-1279.
Chica AB, Martín-Arévalo E, Botta F, Lupiánez J (2014b) The Spatial Orienting paradigm: How
to design and interpret spatial attention experiments. Neuroscience & Biobehavioral
Reviews 40:35-51.
Chica AB, Bayle DJ, Botta F, Bartolomeo P, Paz-Alonso PM (2016a) Interactions between
phasic alerting and consciousness in the fronto-striatal network. Sci Rep 6.

Chica AB, Bayle DJ, Botta F, Bartolomeo P, Paz-Alonso PM (2016b) Interactions between
phasic alerting and consciousness in the fronto-striatal network. Scientific reports
6:31868.
Chica AB, Lasaponara S, Chanes L, Valero-Cabré A, Doricchi F, Lupiáñez J, Bartolomeo P
(2011b) Spatial attention and conscious perception: the role of endogenous and
exogenous orienting. Attention, Perception, & Psychophysics 73:1065-1081.
Colclough GL, Brookes MJ, Smith SM, Woolrich MW (2015) A symmetric multivariate leakage
correction for MEG connectomes. Neuroimage 117:439-448.
Corbetta M, Shulman GL (2002) Control of goal-directed and stimulus-driven attention in the
brain. Nature reviews neuroscience 3:201-215.
Coste CP, Kleinschmidt A (2016) Cingulo-opercular network activity maintains alertness.
Neuroimage 128:264-272.
Davidson MJ, Alais D, van Boxtel JJ, Tsuchiya N (2018) Attention periodically samples
competing stimuli during binocular rivalry. Elife 7:e40868.
Dehaene S, Changeux J-P, Naccache L, Sackur J, Sergent C (2006) Conscious, preconscious,
and subliminal processing: a testable taxonomy. Trends Cogn Sci 10:204-211.
Doricchi F, Macci E, Silvetti M, Macaluso E (2010) Neural correlates of the spatial and
expectancy components of endogenous and stimulus-driven orienting of attention in
the Posner task. Cerebral Cortex 20:1574-1585.
Fan J, McCandliss BD, Sommer T, Raz A, Posner MI (2002) Testing the efficiency and
independence of attentional networks. Journal of cognitive neuroscience 14:340-347.
Fan J, McCandliss BD, Fossella J, Flombaum JI, Posner MI (2005) The activation of attentional
networks. Neuroimage 26:471-479.
Fiebelkorn IC, Kastner S (2019) A rhythmic theory of attention. Trends in cognitive sciences
23:87-101.
Fischl B (2012) FreeSurfer. Neuroimage 62:774-781.
Funes MJ, Lupiáñez J, Milliken B (2007) Separate mechanisms recruited by exogenous and
endogenous spatial cues: Evidence from a spatial Stroop paradigm. Journal of
Experimental Psychology: Human Perception and Performance 33:348.
Han SW, Eaton HP, Marois R (2019) Functional fractionation of the Cingulo-opercular
network: alerting insula and updating cingulate. Cerebral Cortex 29:2624-2638.
Haupt M, Ruiz-Rizzo AL, Sorg C, Finke K (2019) Phasic alerting effects on visual processing
speed are associated with intrinsic functional connectivity in the cingulo-opercular
network. NeuroImage 196:216-226.
Kentridge RW, Nijboer TC, Heywood CA (2008) Attended but unseen: Visual attention is not
sufficient for visual awareness. Neuropsychologia 46:864-869.
Kerzel D, Zarian L, Souto D (2009) Involuntary cueing effects on accuracy measures: Stimulus
and task dependence. Journal of Vision 9:16-16.
Koch C, Tsuchiya N (2007) Attention and consciousness: two distinct brain processes. Trends
in cognitive sciences 11:16-22.
Lamme VA (2006) Towards a true neural stance on consciousness. Trends in cognitive
sciences 10:494-501.
Ling S, Carrasco M (2006) Sustained and transient covert attention enhance the signal via
different contrast response functions. Vision research 46:1210-1220.
Liu T, Abrams J, Carrasco M (2009) Voluntary attention enhances contrast appearance.
Psychological science 20:354-362.

Liu Y, Bengson J, Huang H, Mangun GR, Ding M (2016) Top-down modulation of neural
742   activity in anticipatory visual attention: control mechanisms revealed by
743   simultaneous EEG-fMRI. Cerebral Cortex 26:517-529.
744   Lupiáñez J, Milán EG, Tornay FJ, Madrid E, Tudela P (1997) Does IOR occur in discrimination
745   tasks? Yes, it does, but later. Perception & psychophysics 59:1241-1254.
746   Lupiáñez J, Milliken B, Solano C, Weaver B, Tipper SP (2001) On the strategic modulation of
747   the time course of facilitation and inhibition of return. The Quarterly Journal of
748   Experimental Psychology: Section A 54:753-773.
749   Macaluso E, Doricchi F (2013) Attention and predictions: control of spatial attention beyond
750   the endogenous-exogenous dichotomy. Frontiers in human neuroscience 7:685.
751   Magazzini L, Singh KD (2018) Spatial attention modulates visual gamma oscillations across
752   the human ventral stream. Neuroimage 166:219-229.
753   Maris E, Oostenveld R (2007) Nonparametric statistical testing of EEG-and MEG-data. Journal
754   of neuroscience methods 164:177-190.
755   Mashour GA, Roelfsema P, Changeux J-P, Dehaene S (2020) Conscious processing and the
756   global neuronal workspace hypothesis. Neuron 105:776-798.
757   Oostenveld R, Fries P, Maris E, Schoffelen J-M (2011) FieldTrip: open source software for
758   advanced analysis of MEG, EEG, and invasive electrophysiological data. Comput Intell
759   Neurosci 2011:1.
760   Périn B, Godefroy O, Fall S, de Marco G (2010) Alertness in young healthy subjects: an fMRI
761   study of brain region interactivity enhanced by a warning signal. Brain and cognition
762   72:271-281.
763   Petersen A, Petersen AH, Bundesen C, Vangkilde S, Habekost T (2017) The effect of phasic
764   auditory alerting on visual perception. Cognition 165:73-81.
765   Podvalny E, Flounders MW, King LE, Holroyd T, He BJ (2019) A dual role of prestimulus
766   spontaneous neural activity in visual object recognition. Nature communications
767   10:1-13.
768   Portin K, Vanni S, Virsu V, Hari R (1999) Stronger occipital cortical activation to lower than
769   upper visual field stimuli. Neuromagnetic recordings. Exp Brain Res 124:287-294.
770   Posner MI (1980) Orienting of attention. Quarterly journal of experimental psychology 32:3-
771   25.
772   Posner MI (1994) Attention: the mechanisms of consciousness. Proceedings of the National
773   Academy of Sciences 91:7398-7403.
774   Posner MI, Driver J (1992) The neurobiology of selective attention. Current opinion in
775   neurobiology 2:165-169.
776   Sadaghiani S, D'Esposito M (2015) Functional characterization of the cingulo-opercular
777   network in the maintenance of tonic alertness. Cerebral Cortex 25:2763-2773.
778   Sergent C, Wyart V, Babo-Rebelo M, Cohen L, Naccache L, Tallon-Baudry C (2013) Cueing
779   attention after the stimulus is gone can retrospectively trigger conscious perception.
780   Current biology 23:150-155.
781   Shulman GL, Astafiev SV, McAvoy MP, d'Avossa G, Corbetta M (2007) Right TPJ deactivation
782   during visual search: functional significance and support for a filter hypothesis.
783   Cerebral Cortex 17:2625-2633.
784   Snodgrass JG, Corwin J (1988) Pragmatics of measuring recognition memory: applications to
785   dementia and amnesia. J Exp Psychol Gen 117:34.
786   Spagna A, Mackie M-A, Fan J (2015) Supramodal executive control of attention. Frontiers in
787   psychology 6:65.

Spagna A, Hajhajate D, Liu J, Bartolomeo P (2020a) Visual mental imagery engages the left
fusiform gyrus, but not the early visual cortex: a meta-analysis of neuroimaging
evidence. BioRxiv.
Spagna A, Kim TH, Wu T, Fan J (2020b) Right hemisphere superiority for executive control of
attention. Cortex 122:263-276.
Tadel F, Baillet S, Mosher JC, Pantazis D, Leahy RM (2011) Brainstorm: a user-friendly
application for MEG/EEG analysis. Comput Intell Neurosci 2011:879716.
Tadel F, Bock EA, Niso G, Mosher JC, Cousineau M, Pantazis D, Leahy RM, Baillet S (2019)
MEG/EEG group analysis with Brainstorm. Front Neurosci 13:76.
Tallon-Baudry C (2012) On the neural mechanisms subserving consciousness and attention.
Front Psychol 2:397.
Tallon-Baudry C, Bertrand O, Hénaff M-A, Isnard J, Fischer C (2005) Attention modulates
gamma-band oscillations differently in the human lateral occipital cortex and
fusiform gyrus. Cerebral cortex 15:654-662.
Taulu S, Simola J (2006) Spatiotemporal signal space separation method for rejecting nearby
interference in MEG measurements. Phys Med Biol 51:1759.
Thiebaut De Schotten M, Dell'Acqua F, Forkel S, Simmons A, Vergani F, Murphy DG, Catani M
(2011) A lateralized brain network for visuo-spatial attention. Nature Precedings:1-1.
Tsuchiya N, Koch C (2014) On the relationship between consciousness and attention: MIT
Press.
Tsuchiya N, Koch C (2016) The relationship between consciousness and top-down attention.
In: The neurology of conciousness, pp 71-91: Elsevier.
Uddin LQ (2015) Salience processing and insular cortical function and dysfunction. Nature
reviews neuroscience 16:55-61.
Van Boxtel JJ, Tsuchiya N, Koch C (2010) Opposing effects of attention and consciousness on
afterimages. Proceedings of the National Academy of Sciences 107:8883-8888.
Vernet M, Japee S, Lokey S, Ahmed S, Zachariou V, Ungerleider LG (2019) Endogenous
visuospatial attention increases visual awareness independent of visual
discrimination sensitivity. Neuropsychologia 128:297-304.
Wyart V, Tallon-Baudry C (2008) Neural dissociation between visual awareness and spatial
attention. J Neurosci 28:2667-2679.
Xuan B, Mackie M-A, Spagna A, Wu T, Tian Y, Hof PR, Fan J (2016) The activation of
interactive attentional networks. NeuroImage 129:308-319.
821
822

823 **Tables**

Table 1. MNI Coordinates and size of the area (in cm^2) used in the connectivity analyses.

ROI	x	y	z	area
Left V1	-26	-99	7	12.63
Right V1	25	99	9	17.03
Left OT	-45	-88	-10	12.69
Right OT	45	-80	-17	12.67
Left hV4	-25	-95	-20	10.78
Right hV4	16	-95	-16	7.62
Left Fusiform	-34	-50	-21	9.82
Right Fusiform	34	-50	-21	9.59
Left SPL	-28	-60	57	16.64
Right SPL	21	-39	62	16.17
Left IPL	-27	-82	45	16.65
Right IPL	31	-82	43	16.96
Left TPJ	-62	-55	17	42.86
Right TPJ	67	-38	20	34.75
Left Ant Insula	-35	21	3	13.5
Right Ant Insula	35	21	3	13.82
Left FEF	-47	14	52	14.17
Right FEF	47	14	53	12.28

Note: ROIs were selected based on results from previous studies (Corbetta and Shulman,

2002; Fan et al., 2005; Tallon-Baudry et al., 2005; Bressler et al., 2008; Chica et al., 2013c;

Chica et al., 2016b; Xuan et al., 2016) showing the involvement of these areas in attention-

related processes. In addition, areas of the ventral visual stream were also added to the ROI

analyses to examine the potential modulation of the activity in attention network regions on

early and late visual areas. Seed vertices of the ROIs are reported in MNI coordinates, and the

area of the ROI is reported in cm^2.

a.

| Fixation
1000 to 1500 ms | Cue
50 ms | Fixation
250 ms | Target
16 ms | Fixation
484 ms | Discrimination Task | Detection Task |

Time

b.

4,9°

3,6°

2,5°

4°

1°

6°

c. *NonPredCue*

Invalid
Cue (50%)

Fixation

Target

d. *PredCue*

Valid
Cue (67%)

Fixation

Target

a. *NonPredCue* AEC: Seen *vs* Unseen ($p_{fdr} <.001$)

b. *NonPredCue* AEC: Seen Valid *vs* Unseen Valid ($p_{fdr} <.005$)

c. *NonPredCue* AEC: Seen Invalid *vs* Unseen Invalid ($p_{fdr} <.005$)

d. *PredCue* AEC: Seen Valid *vs* Seen Invalid ($p_{fdr} <.005$)