The cellular architecture of memory modules in Drosophila supports stochastic input integration

Omar A. Hafez (ORCID:0000-0002-7846-9226)[1,5†], Benjamin Escribano (ORCID:0000-0002-1432-5952)[2†], Rouven L. Ziegler (ORCID:0000-0002-3050-7692)[2], Jan J. Hirtz (ORCID:0000-0002-4486-3057)[3], Ernst Niebur (ORCID:0000-0002-2815-9262)[1,4*] and Jan Pielage (ORCID:0000-0002-5115-5884)[2*]

[1]Zanvyl Krieger Mind/Brain Institute, Johns Hopkins University, Baltimore, MD 21218, USA.
[2]Division of Neurobiology and Zoology, Department of Biology, University of Kaiserslautern, D-67663 Kaiserslautern, Germany.
[3]Physiology of Neuronal Networks Group, Department of Biology, University of Kaiserslautern, D-67663 Kaiserslautern, Germany.
[4]Solomon Snyder Department of Neuroscience, Johns Hopkins University School of Medicine, Baltimore MD 21218, USA.
[5]Current address: Yale MD-PhD Program, Yale School of Medicine, New Haven, CT 06510, USA.

*Corresponding author(s). E-mail(s): niebur@jhu.edu; pielage@bio.uni-kl.de;
Contributing authors: ohafez1@alumni.jh.edu; benjamin.escribano@bio.uni-kl.de; roziegle@rhrk.uni-kl.de; hirtz@bio.uni-kl.de;
[†]These authors contributed equally to this work.

Keywords: Mushroom Body Output Neuron, Dendritic Signal Processing, Neuronal Output Tuning, Decision making, Long term memory formation

Abstract The ability to associate neutral stimuli with valence information and to store these associations as memories forms the basis for decision making. To determine the underlying computational principles, we build a realistic computational model of a central decision module within the *Drosophila* mushroom body (MB), the fly's center for learning and memory. Our model combines the electron microscopy-based architecture of one MB output neuron (MBON-α3), the synaptic connectivity of its 948 presynaptic Kenyon cells (KCs), and its *in vivo* membrane properties obtained from patch-clamp recordings. We show that this neuron is electrotonically compact and that synaptic input corresponding to simulated odor input robustly drives its spiking behavior. Therefore, sparse innervation by KCs can efficiently control and modulate MBON activity in response to learning with minimal requirements on the specificity of synaptic localization. This architecture allows efficient storage of large numbers of memories using the flexible stochastic connectivity of the circuit.

1 Introduction

Operating successfully within a complex environment requires organisms to discriminate between a vast number of potential rewarding and stressful interactions. Learning to avoid potentially harmful interactions will promote survival of individual animals and requires long-term memory mechanisms. In recent years, significant progress has been made toward our understanding of the cellular and circuit mechanisms underlying learning and memory. Despite these advances, at the computational level it remains largely unknown how the cellular and circuit architecture contributes to the efficient formation and storage of multiple memories.

Here, we use the *Drosophila melanogaster* mushroom body (MB) as a simple model system to investigate the computational principles underlying learning and memory in the context of decision making. A key advantage of the MB, the learning and memory center of the fly, is its relative simplicity in terms of the number and types of neurons compared to the mammalian brain. It has been demonstrated that the MB is required for the acquisition and recall of associative olfactory memories (de Belle and Heisenberg, 1994), and a wide variety of genetic tools are available to access and manipulate key cell types within the memory circuitry (Owald et al, 2015b; Lai and Lee, 2006)

Odor information is transmitted from the antennal lobe via projection neurons (PNs) to the Kenyon cells (KCs) of the MB. PNs are connected in a largely stochastic manner to the approximately 2000 KCs per brain hemisphere, resulting in a unique KC odor representation in individual flies (Litwin-Kumar et al, 2017; Aso et al, 2014a; Caron et al, 2013; Gruntman and Turner, 2013; Lin et al, 2007). The formation of olfactory memory within the MB circuit is assayed through experimental manipulations in which normally neutral odors are associated with either positive (reward learning) or negative (avoidance learning) valences (Livingstone et al, 1984).

Electrophysiological and calcium imaging studies demonstrated that any given odor activates approximately 5% of KCs (Campbell et al, 2013; Honegger et al, 2011; Turner et al, 2008; Siegenthaler et al, 2019). KCs then transmit this odor information to mushroom body output neurons (MBONs) that subdivide the MB lobes into distinct structural and functional modules by virtue of their dendritic arborizations (Owald et al, 2015a; Aso et al, 2014b; Hige et al, 2015a; Cognigni et al, 2018). During associative olfactory conditioning, the behavioral response to any given odor can be altered to encode positive (approach) or negative (avoidance) behavior. The valence of the conditioned response is encoded by the activation of dopaminergic neurons (DANs) that selectively innervate the structural MB modules defined by the MBON dendrites (Aso et al, 2014a).

DANs of the PPL1 cluster preferentially provide input to the vertical lobe of the MB and are activated during negative reinforcement, while DANs of the PAM cluster are activated during positive reinforcement and modulate activity in the horizontal lobe of the MB (Handler et al, 2019; Aso and Rubin, 2016; Tomchik and Davis, 2009; Séjourné et al, 2011; Cohn et al, 2015; Burke

et al, 2012). The majority of MBONs of the vertical lobe utilize acetylcholine as a neurotransmitter, and artificial activation of these MBONs often mediates approach behavior (Aso et al, 2014b). In contrast, the majority of MBONs of the horizontal lobe release glutamate as a neurotransmitter, and these neurons mediate avoidance behavior (Aso et al, 2014b).

In vivo Calcium imaging and electrophysiological experiments demonstrated that the activity of individual MBONs can be altered by conditioning paradigms (Hige et al, 2015a; Plaçais et al, 2013; Owald et al, 2015a; Jacob and Waddell, 2020; Perisse et al, 2016; Felsenberg et al, 2018). This modulation of MBON activity depends on the pairing of DAN and KC activity and results in either an enhancement or depression of the strength of the KC>MBON connection (Handler et al, 2019; Aso and Rubin, 2016; Owald et al, 2015a; Cohn et al, 2015; Burke et al, 2012). These changes in KC>MBON connection strength occur in either the horizontal or the vertical lobe depending on the administered stimulus-reward contingencies. As a consequence, the resulting net activity of all MBONs is shifted to either approach or avoidance behavior and the behavioral response towards the conditioned stimulus during memory recall is changed (Handler et al, 2019; Aso and Rubin, 2016; Cohn et al, 2015; Burke et al, 2012). Evidence for this MBON balance model (Cognigni et al, 2018) has been provided for short-term memory (Aso et al, 2014b; Hige et al, 2015a; Perisse et al, 2016; Cohn et al, 2015) but to what extent, if any, such a model applies to long term memory remains unclear. Recently, genetic tools have been created that enable genetic tagging of long-term memory engram cells in the MB (Miyashita et al, 2018; Siegenthaler et al, 2019). These studies indicate that modulation of a relatively small number of KCs is sufficient to significantly alter memory recall.

Here, we generate a realistic computational model of the core KC>MBON module that is essential for the incorporation of memories to determine the cellular basis for decision making. Prior studies (Gouwens and Wilson, 2009) used "synthetic" (randomly generated) models of dendritic trees of Drosophila projection neurons to gain first insights into the computational processes of central neurons. Using linear cable theory, they demonstrated that voltage differences are much smaller within the dendritic trees than between the dendrites and the soma. Using a similar approach, Scheffer and coworkers simulated voltage distributions in the dendritic tuft of another Drosophila neuron (EPG) but using its measured morphology, rather than random geometries (Scheffer et al, 2020). Here, we go beyond these studies by combining precise structural data including the full synaptic connectivity from the electron microscopy-based synaptic connectome (Takemura et al, 2017) of the Drosophila MB with functional (electrophysiological) data of an individual MBON. This approach allows us to determine the impact of populations of KCs on MBON activity and to resolve the potential computational mechanisms underlying memory encoding. We focus on MBON-α3 (MBON-14 in the terminology of (Aso et al, 2014a)) an MBON at the tip of the α lobe that is relevant for long term memory (Aso et al, 2014b; Plaçais et al, 2013). We first determine the physiological

parameters of MBON-α3 by patch clamp electrophysiology. We then use these
parameters to generate an anatomically and physiologically realistic *in silico*
model of MBON-α3 and its KC inputs. Using a variety of computational sim-
ulations, we provide evidence that MBON-α3 is electrotonically compact and
perfectly suited to incorporate a substantial number of memories based on
largely stochastic KC inputs.

2 Results

2.1 Electrophysiological characterization of MBON-α3

MBON-α3 is particularly suitable for computational modeling due to
the availability of its complete dendritic reconstruction through electron
microscopy (Takemura et al, 2017). In addition, we can access MBON-α3
genetically by using a specific Gal4 line (MB082C-Gal4) to label MBON-α3
by the expression of membrane-bound 10XUAS-IVS-mCD8::GFP (Figure 1A).
In immunohistochemical co-labellings with the active zone marker Bruchpi-
lot (Brp), we were able to visualize the dendritic input sites, the axonal
projections and the cell soma of MBON-α3 (Figure 1A, arrow). We used
this labelling to perform fluorescence-guided electrophysiological patch-clamp
whole-cell recordings of this neuron *ex vivo* (Figure 1C,D). We carried out
standard patch-clamp protocols in five cells of independent preparations and
recorded all essential physiological parameters (Table 1). First, we performed
a continuous current-clamp protocol over $60s$ at a sampling rate of $50kHz$
which allowed us to determine the average resting membrane potential as
$-56.7 \pm 2.0mV$. This value for the resting membrane potential is in line with
prior measurements of MBONs *in vivo* (Hige et al, 2015a). In these record-
ings, we observed spontaneous firing activity of MBON-α3 with an average
frequency of $12.1Hz$.

Next, we performed a current-clamp de- and hyper-polarization protocol
where we recorded $400ms$ trials at a sampling rate of $50kHz$. Each trial started
with $0pA$ for $10ms$, followed by a current injection of $-26pA$ to $32pA$, with $2pA$
increments for $400ms$. These protocols allowed us to evoke action potentials
and to determine the resulting changes in membrane potential. An example dis-
playing selected traces of a representative recording is shown in Figure 1F. Two
additional examples are illustrated in Figure 1 Supplement 1. From this data
we determined the average membrane time constant as τ_m of $16.06 \pm 2.3ms$.
We then performed a voltage-clamp step protocol without compensating for
the series resistance. In this experiment, we recorded at $20kHz$ while apply-
ing a $5mV$ voltage step for $100ms$. This allowed us to calculate the average
membrane resistance as $R_m = 926 \pm 55M\Omega$, a rather high membrane resis-
tance when compared to similarly sized neurons (Gouwens and Wilson, 2009).
MBON-α3 is thus highly excitable, with small input currents being sufficient
to significantly alter its membrane potential (Gouwens and Wilson, 2009; Sak-
mann and Neher, 1995). The amplitude of action potentials was rather small
but similar to previously reported values for different MBONs (Hige et al,

2015a). The small amplitudes are likely a result of the unipolar morphology
of the neuron with a long neurite connecting the dendritic input region to
the soma (see below). As a result, signal propagation may partially bypass
the soma (Gouwens and Wilson, 2009), Figure 1F. We measured the neuron's
membrane capacitance as $C_m = 16.76 \pm 1.90 pF$, which classifies MBON-α3 as
a mid-sized neuron (Gouwens and Wilson, 2009).

Table 1 Summary of passive membrane properties of MBON-α3, calculated from electrophysiological measurements from five neurons. From left to right: resting membrane potential (V_m) in mV, membrane time constant (τ_m) in ms, capacitance of the membrane (C_m) in pF, specific capacitance (C_{spec}) in $\mu F/cm^2$ and passive membrane conductance (C_{pass}) in S/cm^2.

Sample	$Vm[mV]$	$\tau_m[ms]$	$C_m[pF]$	$C_{spec}[\mu F/cm^2]$	$C_{pass}[S/cm^2]$
Cell 1	-59.2	13.54	12.35	0.200	1.87×10^{-5}
Cell 2	-52.3	14.46	20.79	0.337	2.12×10^{-5}
Cell 3	-56.2	24.58	21.60	0.350	1.48×10^{-5}
Cell 4	-52.8	15.50	15.83	0.257	1.66×10^{-5}
Cell 5	-63.1	12.20	13.21	0.214	1.76×10^{-5}
Mean	-56.7	16.06	16.76	0.272	1.78×10^{-5}
SD	4.5	4.92	4.26	0.069	0.24×10^{-5}
SEM	2.0	2.20	1.90	0.031	0.11×10^{-5}

To enable the calibration of the *in silico* model, we then recorded 35 trials
per cell in a current-clamp injection protocol with injections of $10pA$ current
pulses for $100ms$ at a sampling rate of $20kHz$. We averaged 105 traces from 3
selected example cells to a standard curve, describing the membrane kinetics
of MBON-α3 in response to current injections (Figure 1G). To to further char-
acterize MBON-α3, we analyzed the resulting data from the current-clamp de-
and hyper-polarization protocols in the three example cells (Figure 1F and
Supplement 1). We recorded the absolute deflection, the action potential firing
frequency, and the amplitude of action potentials for all individual cells. In all
three cells, we observed that the absolute deflection of the membrane poten-
tial was proportional to the increasing current injections (Figure 1H). The
action potential firing frequency increased gradually with increasing membrane
potential deflections (Figure 1I), with no significant change in action poten-
tial amplitudes (Figure 1J). The same effect was observed with an increasing
current injection (Figure 1 Supplement 1). This is consistent with the general
idea that action potentials are "all or nothing" events, and that MBON-α3
is a spike-frequency adapting neuron. These values and the tunability of the
MBON-α3 are in good agreement with prior recordings of a different class of
MBONs and of other central neurons in Drosophila (Gouwens and Wilson,
2009; Hige et al, 2015a). We next used these data to generate an *in silico* model
based on realistic passive membrane properties.

Figure 1 Electrophysiological characterization of the MBON-α3 neuron. (A) Immunohistochemical analysis of the MBON-α3 (green, marked by GFP expression) reveals the unique morphology of the MBON with its dendritic tree at the tip of the α-lobe and the cell soma (arrowhead) at a ventro-medial position below the antennal lobes. The brain architecture is revealed by a co-staining with the synaptic active zone marker Bruchpilot (Brp, magenta). (B) Artificial superimposition of a partial rendering of MBON-α3 (green) and of $\alpha\beta\gamma$-KCs (magenta) to illustrate KC>MBON connectivity. A simplified schematic $\alpha\beta$-KC projection is illustrated in white. (C) Transmitted light image of an MBON-α3 cell body attached to a patch pipette. (D) Visualisation of the GFP-expression of MBON-α3 that was used to identify the cell via fluorescence microscopy. The patch pipette tip is attached to the soma (right side). The scale bar in C applies to C and D. (E) Recording of spontaneous activity of an MBON-α3 neuron in current clamp without current injection. (F) Recording of evoked neuronal activity of an MBON-α3 neuron with step-wise increasing injection of $400ms$ current pulses. Pulses start at $-26pA$ (bottom) with increasing $2pA$ steps and end at $+32pA$ (top). (G) Mean trace of the induced membrane polarization resulting from a $200ms$ long current injection of $-10pA$. We combined 105 trials from three different cells (35 each) to determine the membrane time constant τ_m. Gray shading indicates the SD. (H) Relative change of membrane potential after current injection. Vm is equal to the maximum depolarization of the membrane. (I) Correlation between the frequency of action potential firing and changes in membrane potential. (J) Relative change of spike amplitude after current injections. Spike amplitude represents the peak of the largest action potential minus baseline. **Figure 1 Supplement 1** See appendix.

2.2 Constructing an *in silico* model of MBON-α3

For our *in silico* model of MBON-α3, we used the recently published electron microscopy-based connectome of the Drosophila mushroom body (Takemura et al, 2017; Zheng et al, 2018). This dataset includes the precise morphological parameters of the entire dendritic tree, the axon terminals and parts of the neurite connection to the cell soma. This dataset also includes precise location data for all 12,770 synaptic connections from the 948 innervating KCs. To build a complete morphological reconstruction of MBON-α3, we determined the length of the soma to dendrite neurite via confocal light microscopy and we incorporated this information together with all other values into the NEURON simulation environment (Hines and Carnevale, 1997) (Figure 2 A & C and Figure 2 Supplement 1).

Table 2 Electrophysiological values applied for the *in silico* model of MBON-α3-A. Parameter values were obtained from literature or based on the fitting to our electrophysiological data. The first column shows the parameters, with the names used in the NEURON environment in parentheses. The maximal conductance was determined to achieve the target MBON depolarization from monosynaptic KC innervation.

Electrophysiological Properties			
Variable	Description	Fitted	Literature
R_a (Ra)	Cytoplasm resistivity [$\Omega \times$cm]	85.41	30-400 (Borst and Haag, 1996; Gouwens and Wilson, 2009)
C_m (cm)	Specific membrane capacitance [μF/cm^2]	0.6961	0.6-2.6 (Borst and Haag, 1996)
g_{pas} (pas.g)	Passive membrane conductance [S/cm^2]	9.399×10^{-6}	$3.8 \times 10^{-5} - 1.2 \times 10^{-4}$ (Cassenaer and Laurent, 2007)
e_{pas} (pas.e)	Leak reversal potential [mV]	-55.64	-60 (Berger and Crook, 2015)
Synaptic Parameters			
τ_s (tau)	time to max. conductance [ms]	N.A.	0.44 (Su and O'Dowd, 2003)
E	synaptic current reversal potential [mV]	N.A.	8.9 (Su and O'Dowd, 2003)
G_{max} (gmax)	maximal conductance [μmho]	N.A.	1.5627×10^{-5} (Hige et al, 2015b)

Based on the published vector data (Takemura et al, 2017; Zheng et al, 2018) we subdivided the dendritic tree into 4,336 linear sections, of which 3,121 are postsynaptic to one or more of the 12,770 synaptic contacts from 948 KCs (Figure 2B). We used linear cable theory (Niebur, 2008) to generate the *in silico* model of MBON-α3. Linear cable theory is the baseline model for neurites and the most rational approach in the absence of detailed information about potential nonlinear currents.

Figure 2 Construction of a computational model of MBON-α3. (A) Electron microscopy based reconstruction of MBON-α3. Data was obtained from NeuPrint (Takemura et al, 2017) and visualized with NEURON (Hines and Carnevale, 1997). Data set MBON14 (ID 54977) was used for the dendritic architecture (A-C) and MBON14(a3)R (ID 300972942) was used to model the axon (light gray) and synaptic terminal (dark gray, A, C). The proximal neurite (bright green) was defined as the proximal axonal region next to the dendritic tree. This is the presumed site of action potential generation. Connectivity to the soma (blue) is included for illustration of the overall morphology and is not drawn to scale. (B) A magnified side view of the dendritic tree of MBON-α3. Within the dendritic tree a total of 4,336 individual membrane sections were defined (light gray area). The 3,121 membrane sections that contain one or more of the 12,770 synaptic input sites from the 948 innervating KCs are highlighted in magenta. (C) Simplified *in silico* model of MBON-α3 highlighting the size of the individual neuronal segments. We simulated recordings at the proximal neurite or at the soma. (D) Normalized trace of a simulated membrane polarization after injection of a $200ms$ long square-pulse current of $-10pA$ at the soma. (E) Normalized and averaged experimental traces (blue) with standard deviation (light blue) of our measured depolarization. (F) Comparison of a normalized induced depolarization from the model (red) and from the experimental approach (blue). The model was fitted within the NEURON environment to the measured normalized mean traces with a mean squared error between model and measured data of $0.048mV^2$. Two tailed non-parametric Spearman correlation results in r=0.9976 and p=<0.0001 between the two data sets. **Figure 2 Supplement 1** See appendix.

In previous simulations of *Drosophila* neurons, the kinetics of membrane
polarisation as a response to current injections were adjusted to *in vivo* mea-
surements through a fitting procedure (Gouwens and Wilson, 2009). We first

defined the boundaries of the passive membrane parameters within NEU-
RON's PRAXIS (principle axis) optimization algorithm and incorporated the
experimentally defined passive membrane properties (Table 1).

We then determined the biophysical parameters of the *in silico* neuron by
current step injections of $-10pA$ for $200ms$ at the soma and by recording
the neuronal voltage traces. This enabled us to fit the membrane parameters
of our model neuron to the experimental data of the three example cells,
Figure 2E. The biophysical parameters that resulted from the fitting procedure
are provided in Table 2.

We obtained membrane kinetics that closely resembles *in vivo* current injec-
tions of $-10pA$, with a mean squared error between model and measured data
of $0.0481mV^2$. For comparison, we normalized both the modelled current injec-
tion (Figure 2D) and the traces from the *in vivo* recordings (Figure 2E) and
overlayed the resulting traces in Figure 2F. These traces were highly correlated
with a Spearman's correlation coefficient of $r = 0.9976$. These results indicate
that we were able to generate a realistic *in silico* model of MBON-α3.

241 2.3 Effect of KC>MBON synaptic inputs

As KCs are cholinergic (Barnstedt et al, 2016) we utilized previously deter-
mined parameters of cholinergic synapses to simulate synaptic input to
MBON-α3. We used a reversal potential of 8.9mV and a synaptic time constant
$\tau_s = 0.44ms$ as target values (Su and O'Dowd, 2003). Individual synaptic con-
tacts were simulated as alpha functions $(g_{max}(t - t_i) \times \exp(-(t - t_i - \tau_s)/\tau_s))$
where t_i is the time of the incoming spike and $g_{max} = 1.56 \times 10^{-11}S$, chosen to
obtain response levels in agreement with the target values. The average num-
ber of synaptic contacts from a KC to this MBON is 13.47 (Takemura et al,
2017), Fig. 4B. To determine the computational constraints of MBON-α3, we
first stimulated each of the 3,121 dendritic sections that receive KC synaptic
innervation using the described alpha-function conductance change. We then
"recorded" (from our simulation) the resulting voltage excursions at the den-
dritic input site (Fig 3A, red), at the proximal neurite (PN, green) and at the
cell soma (blue). The comparative analysis of the responses at these three loca-
tions revealed two notable features: (1) voltage excursions directly within the
dendrite are faster and substantially larger than in the proximal neurite and
soma, and (2), the resulting voltages in the latter two compartments are much
less variable than in the dendrite and distributed within a very small voltage
range (Figure 3D,E). These results indicate that the morphological architec-
ture and biophysical parameters of MBON-α3 promote a "compactification,"
or "democratization," of synaptic inputs, with all inputs resulting in similar
voltage excursions at the PN and soma, regardless of the strength of the initial
dendritic voltage or the position on the dendritic tree.

To understand the morphological basis for the input computation, we ana-
lyzed the relation between synaptic position and local dendritic volume and
the resulting voltages in the dendrite, PN and soma. Representing the distance
between dendritic inputs and soma (Figure 3G) by a color code, we observed

the largest voltages at distal dendritic sites with the smallest section volume
(Fig. 3H,I).

Figure 3 Computational characterization of MBON-α3. (A) Simulated stimulation of individual synaptic input sites. Voltage deflections of synaptic currents were analyzed locally at the dendritic segment (red, next to electrode tip), the proximal neurite (green) and the soma (blue). Color code also applies to B-F. (B) Resulting mean voltages (lines) and standard deviations (shading) from synaptic activations in 3,121 individual stimulus locations (dendrite), and at the proximal neurite and soma. (C-E) Violin plots of the simulated maximal amplitudes at the dendritic section (C), the proximal neurite (D) and the soma (E) in response to synaptic activations. Dotted lines represents the quartiles and dashed lines the mean. Proportions of the violin body represent the distribution of individual data points. Note the difference in scale and compactification of the amplitudes in the proximal neurite and soma. (F) Distribution of the percentage of voltage attenuation recorded at the soma. (G) Distribution of distance between all individual dendritic sections and the soma. This color code is applied in H and I. (H) The elicited local dendritic depolarization is plotted as a function of the local dendritic section volume. (I) The soma amplitude is plotted as a function of the amplitude at individual dendritic segments. Note different scales between abscissa and ordinate; the blue line represents identity. **Figure 3 Supplement 1** See appendix.

This architecture with the smallest dendritic sections at the most dis-
tant sites may contribute to the compactness of the dendritic tree, ensuring
that even the most distant synaptic inputs result in somatic voltage deflec-
tions comparable to the most proximal ones. It is important to note again the
close range of somatic responses in comparison to the large range of voltages
recorded locally at the dendritic input sites (Fig. 3B). We observed a strong
mean attenuation of 75.99% of the dendritic inputs at the PN and the soma
which correlated with the distance from the recording site (Fig. 3F,H,I and
Supplement 1).

Figure 4 MBON-α3 is electrotonically compact. (A) Simulated activation of all synapses
from all 948 individual KC innervating MBON-α3. A representative example of a KC (KC12)
with 13 input synapses to MBONα3 is shown. (B) Histogram of the distribution of the
number of synapses per KC. (C) Membrane potential traces from simulated activation of all
948 KCs. (D) Distribution of the elicited amplitudes evoked by the individual activations
of all KCs. (E) Correlation between the somatic amplitude and the number of activated
synapses in the trials. (F) Distribution of the somatic amplitude evoked by the activation
of all KCs with exactly 13 input synapses (each dot represents one KC).

We next analyzed the impact of individual KCs on MBON-α3. We sim-
ulated the activation of all synapses of each individual KC that innervates
MBON-α3, using the synaptic parameters described above, and we recorded
the simulated voltage excursions in the soma, Figure 4A. The number of
synapses varied between KCs, ranging from 1 to 27 (plus one outlier with 38
synaptic contacts), with a mean of 13.47 synapses per KC (Figure 4B).
In these simulations, we observed a wide range of voltage excursions (note
the outlier in Figure 4B,C,D,E). The neuron is firmly in a small-signal oper-
ation mode, as we observed a highly linear relation between the voltage
excursion and the number of synapses per KC, Fig 4E. Activating a single
KC leads to a voltage excursion at the soma with a mean of 0.37mV, Fig 4D;

this value is in good agreement with *in vivo* data for MBONs (Hige et al,
2015b,a). To further analyze this linear relationship between synapse number
and somatic voltage excursion, we analyzed all KCs with exactly 13 synaptic
inputs to MBON-α3. This analysis revealed a highly stereotypical depolariza-
tion of the soma for all these KCs, despite wide variations in the position of
the individual synaptic contacts (Figure 4E,F). These results provide further
support for the compactness of MBON-α3.

2.4 Physiological and tuned activation of MBON-α3

Given that $\approx 5\%$ of the 984 KCs innervating MBON-α3 are typically activated
by an odor (Honegger et al, 2011; Campbell et al, 2013; Siegenthaler et al,
2019)d, we next simulated voltage excursions in the soma resulting from the
simultaneous activation of sets of 50 randomly selected KCs. In 1,000 indepen-
dent activation trials (Figure 5A), we observed highly stereotypical activation
patterns, with a mean somatic depolarization of 15.24mV and only small varia-
tions in its time course (Figure 5B,C (blue)). These results are consistent with
an activation of 50 KCs being sufficient to induce action potential generation
in MBON-α3 (Fig 1F). The small range over which the MBON depolarization
varies after input from different (random) populations of KCs, and the fact
that the depolarization was strong enough to robustly fire the cell for each of
the simulated odors, provides support for the current model that odor encoding
in KCs is, at least to a large extent, random (stochastic) (Caron et al, 2013).
Based on *in vivo* data, two different modes of plasticity may act at the
level of the KC-MBON module to alter MBON output to conditioned odors.
In vivo imaging of short-term memory formation demonstrated alterations of
KC input strength (Owald and Waddell, 2015; Hige et al, 2015a; Perisse et al,
2016; Plaçais et al, 2013; Pai et al, 2013; Séjourné et al, 2011; Jacob and
Waddell, 2020) while recent observations of long-term memory demonstrated
a specific change (increase) in the number of KCs representing the conditioned
odor (Delestro et al, 2020; Shyu et al, 2019).
To directly address the relative impact of these two modulations on MBON
depolarization, we repeated the same simulation but now modified input
strength by either altering activated KC numbers or by modulating the synap-
tic gain. To enable a direct comparison, we modulated each factor by 25%.
First, we increased the number of activated KCs per set by 25%, from 50 to
63. This lead to a mean somatic depolarization of 18mV, a significant increase
by 19% (Figure 5B,D; dark blue). Decreasing the number of activated KCs by
25%, from 50 to 38, lead to a mean somatic depolarization of 12mV, a sig-
nificant decrease by 20% (Figure 5B,D; light blue). Next, we either increased
or decreased synaptic strength within sets by 25%. Increase lead to a mean
somatic depolarization of 18mV, a significant increase by 19% (Figure 5C,E;
purple) while decrease lead to a mean somatic depolarization of 12mV, a sig-
nificant decrease by 21% (Figure 5C,E; red). Averaged over all conditions, a
change of input of 25% lead to a change of $20\pm1\%$ in membrane depolarization.

Figure 5 Effect of KC recruitment and synaptic plasticity on MBON-α3 responses. (A) Schematic of the simulated recording paradigm. We modeled the activation of $\approx$ 5% of KCs of random combinations of identified KCs, *i.e.* we activated the anatomically correct synapse locations of 50 randomly selected KCs to mimic activation by an odor. The panel shows the distribution of activated synapses for one of these trials. We then varied the number of activated KCs by $\pm$25% around its mean of 50, *i.e.* 38, 50 and 63 KCs, and we varied synaptic strength $\pm$25% around its original value, *i.e.* 75%, 100% and 125%. (B) Mean somatic depolarizations after activating 1000 different sets of either 38 (light blue), 50 (blue), or 63 (dark blue) KCs, all at standard synaptic strength (100%). Shades represent the standard deviations. (C) Mean somatic depolarizations after activating 1000 different sets of 50 KCs at either 75% (red), 100% (blue) or 125% (purple) synaptic strength. Shades represent the standard deviations. (D, E) Violin plots of the relative amplitudes evoked by the different KC sets from (B, C). (F-G) Violin plots for the comparison of the amplitudes between the different plasticity paradigms. (F) 63 KCs at 100% *vs.* 50 KCs at 125% synaptic strength. (G) 38 KCs at 100% *vs.* 50 KCs at 75% synaptic strength. Plots in D-G use the color code from (B) and (C). (H) 63 KCs at 75% (dark red) *vs.* 50 KCs at 100% (blue) and 38 KCs at 125% (light purple) synaptic strength. See Fig. 3 for symbols in violin plots. (I) Relation between somatic depolarization and the number and strength of activated synapses. Illustrated are all tested conditions for synaptic plasticity and KC recruitment. Stimulation paradigms are color coded as shown. (J) Relation between the slope of the condition-specific linear regressions from (I) and the synaptic strength of the activated KC sets. (K) Relation between the slope of the condition-specific linear regressions from (I) and the number of recruited KCs. (L) Slopes of the condition-specific linear regressions from (I). Color codes of (J-L) as in (I). Statistical significance was tested in multiple comparisons with a parametric one-way ANOVA test (D, E and H) resulting in p=<0.0001 for all conditions, or in an unpaired parametric t-test (F and G) resulting in p=<0.0001 for all conditions. Numerical results are summarized in Table 3.

Even though the integrated synaptic conductance was changed by the same
amount by our manipulation of either synaptic strength or number of acti-
vated KCs, we observed small but significant differences in the resultant mean
somatic amplitudes (Figure 5F,G). Finally, we combined both manipulations
by compensating an increase in KC number with a corresponding decrease in
synaptic strength and vice versa. An overview of the performed simulations
and numerical results are provided in Table 3.

Table 3 Summary of data from the plasticity simulations of Figure 5. Simulations vary
the number of active KCs and/or the strength of the KC-MBON synapses by scaling the
maximal conductance parameter of the synaptic alpha-conductance function. Results
include the mean number of activated synaptic sites as well as the mean amplitude of the
somatic voltage depolarization across the 1000 trial repetitions of each simulation. Ranges
are reported as parentheticals. The slope of the linear regression describing the relation
between the number of active synapses and somatic voltage amplitude is reported
(Fig. 5I-L).

Simulation Parameters			Results	
# KCs	Synaptic Strength	Mean # Activated Synapses	Mean Somatic Depolarization Amplitude [mV]	Slope
63	125%	847.88 (747 - 939)	21.55 (19.66 - 23.21)	0.0184
50	125%	672.55 (573 - 778)	18.14 (15.99 - 20.29)	0.0209
38	125%	511.77 (425 - 615)	14.60 (12.48 - 16.96)	0.0235
63	100%	847.08 (725 - 977)	18.26 (16.15 - 20.34)	0.0166
50	100%	673.76 (574 - 805)	15.24 (13.34 - 17.58)	0.0185
38	100%	512.48 (429 - 599)	12.14 (10.43 - 13.85)	0.0203
63	75%	849.05 (749 - 965)	14.56 (13.11 - 16.17)	0.0141
50	75%	672.86 (570 - 767)	11.98 (10.37 - 13.41)	0.0153
38	75%	511.97 (420 - 606)	9.44 (7.91 - 10.96)	0.0164

Both manipulations resulted, on average, in a significant decrease of MBON
membrane depolarization (Figure 5H). Our simulations revealed a nearly lin-
ear relationship between the activated number of synapses and the measured
amplitude in the soma regardless of synaptic strength (Figure 5I). Interest-
ingly, we observed some differences in the slope of the responses between the
different tuning modalities (Figure 5J,K,L), indicating that these modulations
are not completely equivalent.

348 3 Discussion

The ability of an animal to adapt its behavior to a large spectrum of sensory
information requires specialized neuronal circuit motifs (Scheffer et al, 2020).
For our understanding of the computational principles underlying learning and
memory, it is essential to determine the intrinsic contributions of neuronal
circuit architecture. Associative olfactory memory formation in Drosophila
provides an excellent model system to investigate such circuit motifs, as a large
number of different odors can be associated with either approach or rejection

behavior through the formation of long-term memory within the MB circuitry. In contrast to most axon guidance processes in Drosophila that are essentially identical in all wild-type individuals, processing of odor information in the MB largely depends on the stochastic connectivity of projection neurons to KCs that relay odor information from the olfactory glomeruli to the MBONs (Caron et al, 2013; Gruntman and Turner, 2013; Litwin-Kumar et al, 2017; Warth Pérez Arias et al, 2020). Individual flies display idiosyncratic, apparently random connectivity patterns that transmit information of specific odors to the output circuit of the MB. However, the role of MBONs within the circuit is largely fixed between animals and they can be mostly classified as either approach or avoidance neurons (Aso et al, 2014b,a; Litwin-Kumar et al, 2017; Li et al, 2020; Takemura et al, 2017; Scheffer et al, 2020). Furthermore, individual KCs are not biased in their MBON connectivity but innervate both kinds of output modules, for both approach and avoidance. Learning and memory, the modulation of odor response behavior through pairing of an individual odor with either positive or negative valence, is incorporated *via* the local activity of valence-encoding DANs that can either depress or potentiate KC>MBON synaptic connections (Aso and Rubin, 2016; Berry et al, 2018; Handler et al, 2019; Yamagata et al, 2015; Cohn et al, 2015; Burke et al, 2012; Hige et al, 2015a; Owald et al, 2015a; Jacob and Waddell, 2020; Pai et al, 2013; Plaçais et al, 2013; Séjourné et al, 2011; Bouzaiane et al, 2015). As KC odor specificity and connectivity differ significantly between flies with respect to the number and position of KC>MBON synapses, this circuit module must be based on architectural features supporting robust formation of multiple memories regardless of specific individual KC connectivity.

Prior computational work addressing properties of central nervous system neurons in Drosophila relied on synthetic (randomly generated) data (Gouwens and Wilson, 2009) or partial neuronal reconstructions (Scheffer et al, 2020). We go beyond these previous studies by combining precise structural data from the electron-microscopy based synaptic connectome (Takemura et al, 2017) with functional (electrophysiological) data. The structural data consists of the precise neuroanatomical structure of MBON-α3, including the 12,770 synaptic inputs of all its 948 innervating KCs. The neuron's functional properties were determined from *ex vivo* patch clamp recordings. Near-perfect agreement between experimentally observed and simulated voltage distributions in the dendritic tree shows that linear cable theory is an excellent model for information integration in this system. Together, we obtain a realistic *in silico* model of a central computational module of memory-modulated animal behavior, *i.e.* a mushroom body output neuron.

Our data show that the dendritic tree of the MBON is electrotonically compact, agreeing with a prior electrophysiological characterization of an MBON in locust (Cassenaer and Laurent, 2007). The location of an individual synaptic input within the dendritic tree has therefore only a minor effect on the amplitude of the neuron's output, despite large variations of local dendritic

potentials. This effect was particularly striking when we restricted the analy-
sis to a population of KCs with identical numbers of synapses that all elicited
highly stereotypical responses. The compactification of the neuron is likely
related to the architectural structure of its dendritic tree. Together with the
relatively small size of many central neurons in *Drosophila*, this indicates that
in contrast to large vertebrate neurons, local active amplification or other
compensatory mechanisms (Rumsey and Abbott, 2006; Sterratt et al, 2012;
Froemke et al, 2005; Gidon and Segev, 2009) may not be necessary to support
input normalization in the dendritic tree. In contrast, for axons it has been
recently reported that voltage-gated Na$^+$ channels are localized in putative
spike initiation zones in a subset of central neurons of *Drosophila* (Ravenscroft
et al, 2020). Once *in vivo* electrophysiological data quantitatively describing
local active currents become available, they can be incorporated into our model
to further increase the agreement between model and biological system.

Encoding of odor information and incorporation of memory traces is not
performed by individual KCs but by ensembles of KCs. Calcium imaging *in
vivo* demonstrated that individual odors evoke activity in approximately 3-
9% of KCs (Campbell et al, 2013; Honegger et al, 2011). Our simulations of
1,000 independent trials with random sets of 50 KCs, each representing one
distinct odor that activates approximately 5% of the KCs innervating the tar-
get MBON, demonstrate that such activation patterns robustly elicit MBON
activity to a level that matched our *in vivo* observations of odor-induced activ-
ity, as well as that found in previous studies (Siegenthaler et al, 2019; Hige
et al, 2015a; Warth Pérez Arias et al, 2020). The low variability of depolariza-
tions observed in these simulations indicates that information coding by such
activation patterns is highly robust. As a consequence, labeled line representa-
tions of odor identity are likely not necessary at the level of KCs since relaying
information *via* **any** set of ≈50 KCs is of approximately equal efficiency.
Such a model is supported by a recent computational study demonstrating
that variability in parameters controlling neuronal excitability of individual
KCs negatively affects associative memory performance. The authors provide
evidence that compensatory variation mechanisms exist that ensure similar
activity levels between all odor encoding KC sets to maintain efficient memory
performance (Abdelrahman et al, 2021).

Optical recordings of *in vivo* activity of MBONs revealed selective reduc-
tions in MBON activity in response to aversive odor training (Jacob and
Waddell, 2020; Pai et al, 2013; Owald and Waddell, 2015; Perisse et al, 2016;
Séjourné et al, 2011) or to optogenetic activation of selective DANs (Hige et al,
2015a). More generally, both depression and potentiation of MBON activity
have been previously observed in different MBON modules *in vivo* (Owald
and Waddell, 2015; Hige et al, 2015a; Perisse et al, 2016; Plaçais et al, 2013;
Pai et al, 2013; Séjourné et al, 2011; Jacob and Waddell, 2020). In addi-
tion, recent studies have observed changes in KC stimulus representations
after conditioning (Delestro et al, 2020; Shyu et al, 2019) that may be due to
learning-dependent modulations of synaptic PN input to KCs (Baltruschat

et al, 2021). Our computational model allowed us to implement and com-
pare these two mechanisms changing MBON output: One is a change in the
strength of the KC>MBON synapses (over a range of $\pm 25\%$); the other is a
change in the number of activated KCs (over the same range). Interestingly, we
found almost linear relationships between the number of active KCs and the
resulting depolarizations, and the same between the strength of synapses and
MBON depolarization. Decreasing either of these variables by 25% significantly
decreased the level of MBON activation.

Our simulation data thus shows that the KC>MBON architecture rep-
resents a biophysical module that is well-suited to simultaneously process
changes based on either synaptic and/or network modulation. Together with
the electrotonic nature of the MBONs the interplay between KCs and MBONs
thus ensures reliable information processing and memory storage despite the
stochastic connectivity of the memory circuitry. While our study focuses on
the detailed activity patterns within a single neuron, the availability of large
parts of the fly connectome at the synaptic level, in combination with realistic
models for synaptic dynamics, should make it possible to extend this work to
circuit models to gain a network understanding of the computational basis of
decision making.

464 4 Materials and Methods

465 Fly Stocks

466 Flies were reared on standard fly food at 25°C and 65% humidity. Following
467 stocks were used in this study: 10XUAS-IVS-mCD8::GFP (Pfeiffer et al, 2010)
468 (BDSC 32186), MB082C-Gal4 (Aso et al, 2014b) (BDSC 68286).

469 Immunohistochemistry

Male flies expressing GFP in MBON-α3 were fixed for 3.5h at 4°C in 4%
PFA containing PBST (0.2% Triton-X100). Flies were then washed for at
least 3 x 30 min at RT in PBST before dissection. Dissected brains were
then labelled with primary antibodies (rabbit anti-GFP (A6455, Life technolo-
gies, Carlsbad, USA 1:2000) and mouse anti-Brp nc82, Developmental Studies
Hybridoma Bank, Iowa, USA, 1:200) for two nights at 4°C. After incuba-
tion, brains were washed at least 6 x 30 min in PBST. Secondary antibodies
(Alexa488 (goatαrabbit) and Alexa568 (goatαmouse) coupled antibodies, Life
technologies, Carlsbad, USA, 1:1000) were applied for two nights at 4°C. After
a repeated washing period, brains were submerged in Vectashield (Vector Lab-
oratories, Burlingame, USA), mounted onto microscope slides, and stored at
4°C.

Images were acquired using a confocal scanning microscope (LSM 710, Carl
Zeiss GmbH, Jena, Germany) and a 25x (Plan-NEOFLUAR, NA 0.8 Korr DIC,
Carl Zeiss GmbH, Jena, Germany) oil objective. Raw images were projected

with Fiji (Schindelin et al, 2012) and cropped in Photoshop (Adobe, San José,
USA). Uniform adjustments of brightness and contrast were performed.

Electrophysiology

All patch clamp recordings were performed at room temperature similar to
prior descriptions (Hige et al, 2015a,b). Flies were flipped 1d after hatching
to obtain 2-3 d old flies. Due to the ventral location of the MBON-α3 somata
within the brain, we performed the recordings *ex vivo*. For preparation, flies
expressing GFP in MBON-α3 were briefly anesthetized on ice before removing
the entire brain out of the head capsule. The preparation was performed in
oxygenated (95% O_2 and 5% CO_2) high glucose ($260mM$) extracellular saline.
Analogous to a prior study, the brains were incubated for $30s$ in $0.5mg/ml$
protease (from Streptomyces griseus, CAS# 90036-06-0, Sigma-Aldrich, St.
Louis, USA) containing extracellular saline (Wilson et al, 2004). The brain was
then transferred to standard extracellular saline (Wilson et al, 2004) (in mM:
103 $NaCl$, 3 KCl, 5 TES, 10 trehalose, 260 glucose, 7 sucrose, 26 $NaHCO_3$,
1 NaH_2PO_4, 1.5 $CaCl_2$, 4 $MgCl_2$, pH 7.3, $280 - 290mOsmol/kg$ adjusted
with sucrose). For physiological recordings, the brain was transferred to a glass
bottom chamber with continuously perfused ($2ml/min$) oxygenated standard
extracellular saline and held in place by a custom made platinum frame.
We used glass capillaries (GB150 (F-)-10P, Science Products, Hofheim,
Germany) and a horizontal puller (Flaming Brown Micropipette Puller P-
1000, Sutter instruments, Novato, USA) to obtain pipettes with a resistance
of $7 - 10M\Omega$. For recordings, patch pipettes were filled with internal solu-
tion (Mauss et al, 2014) (in mM: 140 K-aspartate, 10 HEPES, 4 Mg-ATP, 0.5
Na-GTP, 1 EGTA, 1 KCl, pH 7.29 adjusted with KOH, $265mOsmol/kg$).
Whole-cell recordings were made using the EPC10 amplifier (HEKA, Reutlin-
gen, Germany) and the PatchMaster software (HEKA, Reutlingen, Germany).
Signals were low-pass filtered at $3kHz$, and digitized at $10kHz$ via a digital-to-
analog converter. The liquid junction potential of $13mV$ was corrected online.
Patch pipettes were guided under visual control with an upright microscope
(BX51WI; Olympus, Tokio, Japan) equipped with a 60x water immersion
objective (LUMPlanFl/IR; Olympus, Tokio, Japan) and a CCD camera (Orca
05G, Hamamatsu, Japan). GFP signal of target cells was visualized through
fluorescence excitation (100W fluorescence lamp, Carl Zeiss GmbH, Jena, Ger-
many), and a dual-band emission filter set (512/630 HC Dualband Filter
GFP/DsRed, AHF Analysentechnik, Germany). We used GFP expression in
soma and neurite to identify MBON-α3 neurons. During recordings, the fluo-
rescence excitation was shut off to minimize phototoxic effects. The complete
setup was mounted on an air-damped table while being shielded by a Faraday
cage. Series resistance was maintained below $90M\Omega$ and compensated for up
to 35% through the amplifier's compensation circuitry. In current-clamp mode,
cells were held at a baseline of $-60mV$ to measure the relevant parameters for
the model. Signals were recorded with a sample rate of $20kHz$ or $50kHz$ and
low-pass filtered at $5kHz$.

To determine passive electrical membrane properties and qualitative patch
characteristics, a series of standard protocols were performed:
VCstep without compensation. In voltage-clamp mode, a voltage-step
$(100ms, 5mV)$ is applied without compensating the series resistance. The cor-
responding current trace was then used to determine R_{series}, R_{input} and R_m
with Igor Pro (WaveMetrics, Portland, USA).
CCdehyper. In current-clamp mode, a step-protocol was performed and
recorded at $50kHz$. Each sweep starts with $0pA$ for $10ms$, which is followed by
a current injection ($-26pA$ to $32pA$, with $2pA$ increments) and a step duration
of $400ms$. The resulting changes in membrane potential and induced action
potentials were then analyzed and plotted with Matlab (MathWorks, Natick,
USA). The maximum depolarization was calculated as a mean of the last 70
sample points of the stimulation sweep in each trace. The resting membrane
potential was calculated as the mean of the first 500 sample points before the
stimulation in each trace. The absolute action potential was measured as the
maximum of each trace. The relative action potential amplitude resulted from
subtracting the baseline from the absolute value. The baseline resulted from
the mean of a 400 sample points window, starting 700 sample points upstream
of the maxima in each trace. The resulting trace of the $-10pA$ injection was
used to calculate the membrane time constant τ_m of each cell.
CCcontinuous. In current-clamp mode, no current was injected throughout
a $60s$ recording at $50kHz$. The resting membrane potential was determined
as the mean of the baseline in the recording with Igor Pro and spontaneous
activity was plotted with Matlab.
CCinject. In current-clamp mode, a short current pulse ($-10pA$, $200ms$)
was injected and recorded at $20kHz$. We performed 35 iterations per cell. The
resulting data of three example cells was used for fitting the membrane kinetics
of the model as described previously (Gouwens and Wilson, 2009). Averaging,
normalisation (V_m-V_{m0}/V_{m0}) and plotting was performed with Matlab. This
data was used to perform the model fitting. The baseline (V_{m0}) resulted from
the average of the sample points before stimulation.
Cells displaying a resting membrane potential higher than $-45mV$ and/or
a R_m smaller than $600M\Omega$ were excluded from the analysis. A list of the five
included cells and the corresponding calculated values is provided in Table 1.

Computational Model

Morphology data for the Drosophila MBON-α3-A was originally characterized
using scanning electron microscopy (Takemura et al, 2017) and for the present
study obtained from the "mushroombody" dataset hosted on the database
neuPrint (https://neuprint-examples.janelia.org/; cell ID 54977). These data
describe the structure of its dendritic arborization throughout the MBON
α3 compartment (Fig. 2B, and Supplement 1), with coordinates specified
at 8 nm pixel resolution and converted to μm for model implementation.
The reconstruction is limited to the portion of the neuron in the mushroom

body, not including its soma and axon. The geometry of the axon was determined by investigating a separate MBON, MBON14(a3)R (Fig. 2A), obtained from neuPrint's "hemibrain:v1.1" dataset (https://neuprint.janelia.org/; cell ID 300972942). The axon and synaptic terminal of this neuron were approximated as five separate sections, distinguished by the major branch points identified in the electron microscopic reconstruction (Fig. 2 Supplement 1). Each section was characterized by its total length and average diameter and divided into segments to maintain the average section length of that region in the original reconstruction. They were then appended to the MBON-α3-A proximal neurite as individual linear segments (Fig. 2 Supplement 1). The dimensions of the soma were characterized by confocal microscopy of MBON-α3-A (Fig. 1A). Synaptic locations of the KC innervation of MBON-α3-A were also obtained from neuPrint's "mushroombody" dataset. MBON-α3-A is innervated by 948 distinct KCs, forming a total of 12,770 synapses with an average of 13.47 synapses per KC (Takemura et al, 2017).

Table 4 Morphological parameters of the model

Parameter	Value
number of sections	4342
average section length	$1.24\mu m$
total length	$5377.39\mu m$
average diameter	$0.29\mu m$
average segment surface area	$1.26\mu m^2$
total surface area	$6168.32\mu m^2$
soma diameter	$6.45\mu m$

Electrophysiological properties of the model neuron (Table 2) were determined by fitting the model response to the recorded data, an approach consistent with prior work modeling *Drosophila* central neurons (Gouwens and Wilson, 2009). Somatic voltage excursions in response to 200ms current injections of -10pA were recorded *in vivo* (Fig. 1 G) and compared with the simulated membrane potential (Fig. 2D). This current injection was replicated *in silico* and the membrane potential change at the soma was fit to our recorded data by varying the cell's cytoplasm resistivity, specific membrane capacitance, and passive membrane conductance. The fitting was performed using NEURON's principle axis optimization algorithm, where parameters were tuned until the computed voltage excursion at the soma matched our electrophysiological recordings. The model optimization searched parameter space to minimize an error function defined as the squared differences between the averaged experimental responses to current pulses (Fig. 1 G) and the computed model responses (Fig. 2D-F). The fitting was then verified by normalizing both the model and experimental voltage excursions and performing two-tailed nonparametric Spearman correlation. The model was first allowed

to equilibrate for 30ms simulated time, after which the 200 ms current pulse
was initiated. The region of data used for the fitting ranged from 32.025ms
to 228.03ms, excluding approximately the first and last 2ms of current injec-
tion. The domain of the parameter space was restricted to physiologically
plausible values: $0.5\mu F/cm^2$ to $1.5\mu F/cm^2$ for specific membrane capacitance,
1E-7S/cm^2 to 1E-4S/cm^2 for passive membrane conductance, and $30\Omega cm$ to
$400\Omega cm$ for cytoplasm resistivity. The final values for each parameter are
well within these margins. Resting membrane potential and the leak current
reversal potential were set at -55.64mV based on the initial conditions of the
recorded data to which the model was fit.
Cholinergic KC to MBON synapses were modeled as localized conductance
changes described by alpha functions (Table 2). KC innervation of MBON-α3-
A was determined to be cholinergic (Takemura et al, 2017). Currents recorded
in *Drosophila* KCs were demonstrated to have a reversal potential of 8.9mV and
a rise time to a maximal conductance of 0.44ms (Su and O'Dowd, 2003). The
maximal conductance for the synapses was set at $1.5627 * 10^{-5}\mu S$, the value
determined to achieve the target MBON depolarization from monosynaptic
KC innervation (Hige et al, 2015b).
All simulations were performed using the NEURON 7.7.2 simulation envi-
ronment (Hines and Carnevale, 1997) running through a Python 3.8.5 interface
in Ubuntu 20.04.1. The morphology data divided the MBON-α3-A dendritic
tree into 4336 sections. We added 6 additional sections describing the axon,
synaptic terminals, and soma to a total of 4342 sections with an average of
$1.24\mu m$ in length and $0.29\mu m$ in diameter (Table 4). Passive leak channels
were inserted along the membrane of the dendrite sections as well as through-
out the axon and soma. Each dendritic section was automatically assigned an
index number by NEURON and modeled as a piece of cylindrical cable, with
connectivity specified by the neuron's morphology. The coordinate location of
each KC synapse was mapped to the nearest section of the MBON dendrite,
and the conductance-based synapse model was implemented in the center of
that section. Simulations were performed in which each synapse-containing
section was individually activated, followed by additional simulations activat-
ing groups of synapses corresponding to single or multiple active KCs. A list of
all performed simulations related to Figure 5 is provided in Table 3. Membrane
potential data was analyzed and plotted with MATLAB R2020b (MathWorks,
Natick, USA) and Prism 9 (GraphPad, San Diego, USA).

Statistical Analysis

Statistical analysis and visualization of data was performed in Prism 9 (Graph-
Pad, San Diego, USA) and Matlab (MathWorks, Natick, USA). Before a
statistical comparison was performed, individual groups were tested in Prism
for normality and lognormality with an Anderson-Darling test. Statistical sig-
nificance was tested with an unpaired parametric t-test or in case of multiple
comparisons, with an ordinary corrected parametric one-way ANOVA test.
Further information is provided in figure legends.

Acknowledgments. We would like to thank Glenn Turner and Toshihide
Hige for insightful comments on the patch clamp analysis.

650 Declarations

651 Funding

652 This work was supported by NIH R01DC020123, NSF 1835202, NIH
653 R01DA040990, NIH R01EY027544, NIH Medical Scientist Training Pro-
654 gram Training Grant T32GM136651 (all EN), and a BMBF grant (FKZ
655 01GQ2105)(JP).

656 Competing interests

657 The authors declare no competing interests.

658 Availability of data, materials, and code

659 All code and data files necessary to replicate the simulations are available at:
660 https://doi.org/10.7281/T1/HRK27V.

661 Authors' contributions

662 O.A.H., B.E, E.N. and J.P. designed research; O.A.H., B.E. and R.L.Z.
663 performed research; O.A.H., B.E., R.L.Z., J.J.H., E.N., J.P. analyzed data;
664 O.A.H., B.E., E.N. and J.P. wrote the paper

665 References

Abdelrahman NY, Vasilaki E, Lin AC (2021) Compensatory variability in
network parameters enhances memory performance in the Drosophila mush-
room body. Proceedings of the National Academy of Sciences of the
United States of America 118(49):e2102158,118. https://doi.org/10.1073/
pnas.2102158118

Aso Y, Rubin GM (2016) Dopaminergic neurons write and update memories
with cell-type-specific rules. eLife 5:e16,135. https://doi.org/10.7554/eLife.
16135

Aso Y, Hattori D, Yu Y, et al (2014a) The neuronal architecture of the
mushroom body provides a logic for associative learning. eLife 3:e04,577.
https://doi.org/10.7554/eLife.04577

Aso Y, Sitaraman D, Ichinose T, et al (2014b) Mushroom body output neurons
encode valence and guide memory-based action selection in Drosophila. eLife
3:e04,580. https://doi.org/10.7554/eLife.04580

Baltruschat L, Prisco L, Ranft P, et al (2021) Circuit reorganization in the
Drosophila mushroom body calyx accompanies memory consolidation. Cell
Reports 34(11):108,871. https://doi.org/10.1016/j.celrep.2021.108871

Barnstedt O, Owald D, Felsenberg J, et al (2016) Memory-Relevant Mushroom
Body Output Synapses Are Cholinergic. Neuron 89(6):1237–1247. https://
doi.org/10.1016/j.neuron.2016.02.015

Berger SD, Crook SM (2015) Modeling the Influence of Ion Channels on Neu-
ron Dynamics in Drosophila. Frontiers in Computational Neuroscience 9:139.
https://doi.org/10.3389/fncom.2015.00139

Berry JA, Phan A, Davis RL (2018) Dopamine Neurons Mediate Learning
and Forgetting through Bidirectional Modulation of a Memory Trace. Cell
Reports 25(3):651–662.e5. https://doi.org/10.1016/j.celrep.2018.09.051

Borst A, Haag J (1996) The intrinsic electrophysiological characteristics of
fly lobula plate tangential cells: I. Passive membrane properties. Jour-
nal of Computational Neuroscience 3(4):313–336. https://doi.org/10.1007/
BF00161091

Bouzaiane E, Trannoy S, Scheunemann L, et al (2015) Two Independent
Mushroom Body Output Circuits Retrieve the Six Discrete Components
of Drosophila Aversive Memory. Cell Reports 11(8):1280–1292. https://doi.
org/10.1016/j.celrep.2015.04.044

Burke CJ, Huetteroth W, Owald D, et al (2012) Layered reward signalling
through octopamine and dopamine in Drosophila. Nature 492(7429):433–
437. https://doi.org/10.1038/nature11614

Campbell RAA, Honegger KS, Qin H, et al (2013) Imaging a population
code for odor identity in the Drosophila mushroom body. The Jour-
nal of Neuroscience: The Official Journal of the Society for Neuroscience
33(25):10,568–10,581. https://doi.org/10.1523/JNEUROSCI.0682-12.2013

Caron SJC, Ruta V, Abbott LF, et al (2013) Random convergence of olfactory
inputs in the Drosophila mushroom body. Nature 497(7447):113–117. https:
//doi.org/10.1038/nature12063

Cassenaer S, Laurent G (2007) Hebbian STDP in mushroom bodies facil-
itates the synchronous flow of olfactory information in locusts. Nature
448(7154):709–713. https://doi.org/10.1038/nature05973

Cognigni P, Felsenberg J, Waddell S (2018) Do the right thing: Neural network
mechanisms of memory formation, expression and update in Drosophila.
Current Opinion in Neurobiology 49:51–58. https://doi.org/10.1016/j.conb.
2017.12.002

Cohn R, Morantte I, Ruta V (2015) Coordinated and Compartmentalized Neu-
romodulation Shapes Sensory Processing in Drosophila. Cell 163(7):1742–
1755. https://doi.org/10.1016/j.cell.2015.11.019

de Belle J, Heisenberg M (1994) Associative odor learning in Drosophila abol-
ished by chemical ablation of mushroom bodies. Science 263(5147):692–695.
https://doi.org/10.1126/science.8303280

Delestro F, Scheunemann L, Pedrazzani M, et al (2020) In vivo large-scale
analysis of Drosophila neuronal calcium traces by automated tracking
of single somata. Scientific Reports 10(1):7153. https://doi.org/10.1038/
s41598-020-64060-x

Felsenberg J, Jacob PF, Walker T, et al (2018) Integration of Parallel Opposing
Memories Underlies Memory Extinction. Cell 175(3):709–722.e15. https://
doi.org/10.1016/j.cell.2018.08.021

Froemke RC, Poo Mm, Dan Y (2005) Spike-timing-dependent synaptic plas-
ticity depends on dendritic location. Nature 434(7030):221–225. https://doi.
org/10.1038/nature03366

Gidon A, Segev I (2009) Spike-timing-dependent synaptic plasticity and synap-
tic democracy in dendrites. Journal of Neurophysiology 101(6):3226–3234.
https://doi.org/10.1152/jn.91349.2008

Gouwens NW, Wilson RI (2009) Signal Propagation in Drosophila Cen-
tral Neurons. Journal of Neuroscience 29(19):6239–6249. https://doi.org/10.
1523/JNEUROSCI.0764-09.2009

Gruntman E, Turner GC (2013) Integration of the olfactory code across
dendritic claws of single mushroom body neurons. Nature Neuroscience
16(12):1821–1829. https://doi.org/10.1038/nn.3547

Handler A, Graham TG, Cohn R, et al (2019) Distinct Dopamine Receptor
Pathways Underlie the Temporal Sensitivity of Associative Learning. Cell
178(1):60–75.e19. https://doi.org/10.1016/j.cell.2019.05.040

Hige T, Aso Y, Modi MN, et al (2015a) Heterosynaptic Plasticity Underlies
Aversive Olfactory Learning in Drosophila. Neuron 88(5):985–998. https:
//doi.org/10.1016/j.neuron.2015.11.003

Hige T, Aso Y, Rubin GM, et al (2015b) Plasticity-driven individualization of
olfactory coding in mushroom body output neurons. Nature 526(7572):258–
262. https://doi.org/10.1038/nature15396

Hines ML, Carnevale NT (1997) The NEURON Simulation Environment.
Neural Computation 9(6):1179–1209. https://doi.org/10.1162/neco.1997.9.

6.1179

Honegger KS, Campbell RAA, Turner GC (2011) Cellular-Resolution Popu-
lation Imaging Reveals Robust Sparse Coding in the Drosophila Mushroom
Body. Journal of Neuroscience 31(33):11,772–11,785. https://doi.org/10.
1523/JNEUROSCI.1099-11.2011

Jacob PF, Waddell S (2020) Spaced Training Forms Complementary Long-
Term Memories of Opposite Valence in Drosophila. Neuron 106(6):977–
991.e4. https://doi.org/10.1016/j.neuron.2020.03.013

Lai SL, Lee T (2006) Genetic mosaic with dual binary transcriptional systems
in Drosophila. Nature Neuroscience 9(5):703–709. https://doi.org/10.1038/
nn1681

Li F, Lindsey JW, Marin EC, et al (2020) The connectome of the adult
Drosophila mushroom body provides insights into function. eLife 9:e62,576.
https://doi.org/10.7554/eLife.62576

Lin HH, Lai JSY, Chin AL, et al (2007) A Map of Olfactory Representation
in the Drosophila Mushroom Body. Cell 128(6):1205–1217. https://doi.org/
10.1016/j.cell.2007.03.006

Litwin-Kumar A, Harris KD, Axel R, et al (2017) Optimal Degrees of Synap-
tic Connectivity. Neuron 93(5):1153–1164.e7. https://doi.org/10.1016/j.
neuron.2017.01.030

Livingstone MS, Sziber PP, Quinn WG (1984) Loss of calcium/calmodulin
responsiveness in adenylate cyclase of rutabaga, a Drosophila learning
mutant. Cell 37(1):205–215. https://doi.org/10.1016/0092-8674(84)90316-7

Mauss AS, Meier M, Serbe E, et al (2014) Optogenetic and Pharmacologic
Dissection of Feedforward Inhibition in Drosophila Motion Vision. Journal
of Neuroscience 34(6):2254–2263. https://doi.org/10.1523/JNEUROSCI.
3938-13.2014

Miyashita T, Kikuchi E, Horiuchi J, et al (2018) Long-Term Memory Engram
Cells Are Established by c-Fos/CREB Transcriptional Cycling. Cell Reports
25(10):2716–2728.e3. https://doi.org/10.1016/j.celrep.2018.11.022

Niebur E (2008) Neuronal cable theory. Scholarpedia 3(5):2674. https://doi.
org/10.4249/scholarpedia.2674

Owald D, Waddell S (2015) Olfactory learning skews mushroom body out-
put pathways to steer behavioral choice in Drosophila. Current Opinion in
Neurobiology 35:178–184. https://doi.org/10.1016/j.conb.2015.10.002

Owald D, Felsenberg J, Talbot CB, et al (2015a) Activity of Defined Mushroom
Body Output Neurons Underlies Learned Olfactory Behavior in Drosophila.
Neuron 86(2):417–427. https://doi.org/10.1016/j.neuron.2015.03.025

Owald D, Lin S, Waddell S (2015b) Light, heat, action: Neural control of fruit
fly behaviour. Philosophical Transactions of the Royal Society B: Biological
Sciences 370(1677):20140,211. https://doi.org/10.1098/rstb.2014.0211

Pai TP, Chen CC, Lin HH, et al (2013) Drosophila ORB protein in two
mushroom body output neurons is necessary for long-term memory forma-
tion. Proceedings of the National Academy of Sciences 110(19):7898–7903.
https://doi.org/10.1073/pnas.1216336110

Perisse E, Owald D, Barnstedt O, et al (2016) Aversive Learning and Appet-
itive Motivation Toggle Feed-Forward Inhibition in the Drosophila Mush-
room Body. Neuron 90(5):1086–1099. https://doi.org/10.1016/j.neuron.
2016.04.034

Pfeiffer BD, Ngo TTB, Hibbard KL, et al (2010) Refinement of Tools for
Targeted Gene Expression in Drosophila. Genetics 186(2):735–755. https:
//doi.org/10.1534/genetics.110.119917

Plaçais PY, Trannoy S, Friedrich AB, et al (2013) Two Pairs of Mushroom
Body Efferent Neurons Are Required for Appetitive Long-Term Mem-
ory Retrieval in Drosophila. Cell Reports 5(3):769–780. https://doi.org/10.
1016/j.celrep.2013.09.032

Ravenscroft TA, Janssens J, Lee PT, et al (2020) Drosophila Voltage-Gated
Sodium Channels Are Only Expressed in Active Neurons and Are Localized
to Distal Axonal Initial Segment-like Domains. The Journal of Neuroscience:
The Official Journal of the Society for Neuroscience 40(42):7999–8024. https:
//doi.org/10.1523/JNEUROSCI.0142-20.2020

Rumsey CC, Abbott LF (2006) Synaptic democracy in active dendrites. Jour-
nal of Neurophysiology 96(5):2307–2318. https://doi.org/10.1152/jn.00149.
2006

Sakmann B, Neher E (eds) (1995) Single-Channel Recording. Springer US,
Boston, MA, https://doi.org/10.1007/978-1-4419-1229-9

Scheffer LK, Xu CS, Januszewski M, et al (2020) A connectome and analysis
of the adult Drosophila central brain. eLife 9:e57,443. https://doi.org/10.
7554/eLife.57443

Schindelin J, Arganda-Carreras I, Frise E, et al (2012) Fiji: An open-source
platform for biological-image analysis. Nature Methods 9(7):676–682. https:
//doi.org/10.1038/nmeth.2019

Séjourné J, Plaçais PY, Aso Y, et al (2011) Mushroom body efferent neurons
responsible for aversive olfactory memory retrieval in Drosophila. Nature
Neuroscience 14(7):903–910. https://doi.org/10.1038/nn.2846

Shyu WH, Lee WP, Chiang MH, et al (2019) Electrical synapses between
mushroom body neurons are critical for consolidated memory retrieval
in Drosophila. PLOS Genetics 15(5):e1008,153. https://doi.org/10.1371/
journal.pgen.1008153

Siegenthaler D, Escribano B, Bräuler V, et al (2019) Selective suppression and
recall of long-term memories in Drosophila. PLoS biology 17(8):e3000,400.
https://doi.org/10.1371/journal.pbio.3000400

Sterratt DC, Groen MR, Meredith RM, et al (2012) Spine Calcium Transients
Induced by Synaptically-Evoked Action Potentials Can Predict Synapse
Location and Establish Synaptic Democracy. PLOS Computational Biology
8(6):e1002,545. https://doi.org/10.1371/journal.pcbi.1002545

Su H, O'Dowd DK (2003) Fast Synaptic Currents in *Drosophila* Mush-
room Body Kenyon Cells Are Mediated by α-Bungarotoxin-Sensitive Nico-
tinic Acetylcholine Receptors and Picrotoxin-Sensitive GABA Receptors.
The Journal of Neuroscience 23(27):9246–9253. https://doi.org/10.1523/
JNEUROSCI.23-27-09246.2003

Takemura Sy, Aso Y, Hige T, et al (2017) A connectome of a learning and
memory center in the adult Drosophila brain. eLife 6:e26,975. https://doi.
org/10.7554/eLife.26975

Tomchik SM, Davis RL (2009) Dynamics of Learning-Related cAMP Signal-
ing and Stimulus Integration in the Drosophila Olfactory Pathway. Neuron
64(4):510–521. https://doi.org/10.1016/j.neuron.2009.09.029

Turner GC, Bazhenov M, Laurent G (2008) Olfactory Representations
by Drosophila Mushroom Body Neurons. Journal of Neurophysiology
99(2):734–746. https://doi.org/10.1152/jn.01283.2007

Warth Pérez Arias CC, Frosch P, Fiala A, et al (2020) Stochastic and Arbi-
trarily Generated Input Patterns to the Mushroom Bodies Can Serve as
Conditioned Stimuli in Drosophila. Frontiers in Physiology 11:53. https:
//doi.org/10.3389/fphys.2020.00053

Wilson RI, Turner GC, Laurent G (2004) Transformation of Olfactory Rep-
resentations in the Drosophila Antennal Lobe. Science 303(5656):366–370.
https://doi.org/10.1126/science.1090782

Yamagata N, Ichinose T, Aso Y, et al (2015) Distinct dopamine neurons
mediate reward signals for short- and long-term memories. Proceedings of

the National Academy of Sciences 112(2):578–583. https://doi.org/10.1073/
pnas.1421930112

Zheng Z, Lauritzen JS, Perlman E, et al (2018) A Complete Electron
Microscopy Volume of the Brain of Adult Drosophila melanogaster. Cell
174(3):730–743.e22. https://doi.org/10.1016/j.cell.2018.06.019

Figure 1 Supplement 1 Additional information about the electrophysiology of MBON-α3. Second (A) and third (B) example cells for the evoked neuronal activity of MBON-α3 through a step-wise increasing injection of $400ms$ current pulses. Pulses start at $-26pA$ (bottom) with increasing $2pA$ steps and end at $+32pA$ (top). (C) Correlation between frequency of action potential firing and injected current. The frequency describes action potentials per second. (D) Relative change of spike amplitude after current injections. Spike amplitude corresponds to the peak of the highest action potential minus baseline. (E) Change of membrane potential relative to the resting membrane potential after current injection. ΔVm is equal to the maximum depolarisation minus the resting membrane potential.

Section	Length [μm]	Diameter [μm]	Number of Segments
Neurite 1 (red)	75.89	1.14	27
Neurite 2 (orange)	151.58	0.63	39
Branch 1 (blue)	65.98	0.69	17
Branch 2 (violet)	480.32	0.80	363
Branch 3 (brown)	129.02	0.79	97
Soma (black)	6.45	6.45	1

Figure 2 Supplement 1 Overview of the axonal sections and soma in the *in silico* model. The pink and brown sections represent the presynaptic release sites of MBON-α3. Color-coded electron-microscopy based reconstruction (A) and *in silico* model (B). The reconstruction of the hemibrain did not include the complete axonal area (yellow) and the soma. (B) illustrates the appended sections and the corresponding diameter and lengths. The soma diameter (black) and axon length (yellow) were estimated from confocal microscopy images and literature. The table shows each section of the model (colors as in (A), (B)) including length, diameter and number of segments, where the latter corresponds to NEU-RON's 'nseg' variable for the respective section.

Figure 3 Supplement 1 Additional information about voltage attenuation at the proximal neurite (PN) and soma. (A) Distance to the PN for each stimulated dendritic segment with the color code for the following graphs. (B) Scatter plot of the elicited amplitude relative to the resting membrane potential in each stimulated dendritic section as function of its surface area. (C) Scatter plot of the voltage in the PN as a function of the voltage in the dendrite, both relative to their baseline values. The blue line represents the line of identity. (D) Correlation of the measured distance to the PN with the percent of voltage attenuation in the PN for each synaptic section. (E) Distribution of the percentage of voltage attenuation in the soma per synapse. Vertical dotted line represents the mean attenuation in percent. (F) Correlation of the measured distance to the soma with the percent of voltage attenuation in the soma for each synaptic section. (G) Correlation of the attenuation in percent at the soma (blue) or at the proximal neurite (green) relative to the dendritic voltage.