**Transient *grb10a* Knockdown Permanently Alters Growth, Cardiometabolic Phenotype**

**and the Transcriptome in *Danio rerio***

Bridget L Evans[1], Terence Garner[1], Chiara De Leonibus[1], Oliver H Wearing[2], Holly A Shiels[2],

Adam F L Hurlstone[3], Peter E Clayton[1], Adam Stevens[1*]

**Affiliations:**

[1] Division of Developmental Biology and Medicine, School of Medical Sciences, Faculty of

Biology, Medicine, and Health, University of Manchester, Manchester, UK.

[2] Division of Cardiovascular Sciences, School of Medical Sciences, Faculty of Biology

Medicine and Health, University of Manchester, Manchester, UK.

[3] Division of Infection, Immunity, and Respiratory Medicine, School of Biological Sciences,

Faculty of Biology, Medicine, and Health, University of Manchester, Manchester, UK.

BLE - bridget.evans@postgrad.manchester.ac.uk  ORCID 0000-0001-7481-606X

TG - terence.garner@manchester.ac.uk  ORCID 0000-0003-3962-9730

CDL - c.deleonibus@tigem.it  ORCID 0000-0003-4440-3402

OHW- wearingo@mcmaster.ca ORCID 0000-0002-1866-0416

HAS - holly.shiels@manchester.ac.uk  ORCID 0000-0001-5223-5205.

AFLH - adam.hurlstone@manchester.ac.uk  ORCID 0000-0001-5260-9457.

PEC - peter.clayton@manchester.ac.uk  ORCID 0000-0003-1225-4537.

AS* - adam.stevens@manchester.ac.uk  ORCID 0000-0002-1950-7325.

*Corresponding author.

**Abstract**

Embryonic growth trajectory is a risk factor for chronic metabolic and cardiovascular disorder. Grb10 is a negative regulator of the main pathways driving embryonic growth. This study investigates the long-term cardiometabolic consequences and transcriptomic profiles of transient disruption of grb10a expression in Danio rerio. Knockdown was associated with increased embryonic growth (+7%) and metabolic rate (+25%), and decreased heart rate (-50%) in early life. Juvenile growth and respiratory rate were also elevated (+30% and 7-fold increase respectively). The transcriptome was permanently remodelled by this transient disruption, with dysregulation of multiple growth, cardiac, and metabolic pathways. Phenotypic alteration persisted into adulthood, resulting in a leaner body with elevated skeletal and cardiac muscle content and aerobic scope (43%). This study not only confirms for the first time that transient disruption of a single gene can result in permanent transcriptomic remodelling but correlates this remodelling with persistent alterations to the adult cardiometabolic phenotype.

## Introduction

The Foetal Origins of Adult Disease (FOAD) hypothesis proposes that diseases occurring in adulthood have their origins during development. While this hypothesis was first proposed in relation to coronary heart disease and foetal undernutrition in humans[1], it is now accepted that a wide range of diseases have early developmental origins. In humans, small and large for gestational age (SGA, LGA) status are early risk factors for chronic metabolic and cardiovascular disorders, including type II diabetes (TIID), obesity, cardiac dysfunction, and hypertension[1-3]. Therefore, understanding the impact of altered embryonic growth trajectory on later life disease is important to identify at-risk individuals.

During embryonic development, temperature, circulating glucose levels, and oxygen availability serve as indicators of the mature, external environment[2–4]. Small changes to the phenotype occur in response to external cues to promote immediate survival in a process termed "developmental plasticity". This is important for healthy development, though immediate survival can come at the expense of elevated disease risk in later life, particularly when there is mismatch between the developing and mature environments[3,5]. While these correlations have been observed across a wide number of species, targeted, longitudinal in-vivo studies to elucidate the developmental origins of health and disease, and the pathways involved, are lacking.

Zebrafish are an ideal model organism for both longitudinal and developmental study owing to a rapid generation time, large clutch sizes, and ease of access to embryos. As in mammals, embryonic growth in zebrafish is primarily driven by the insulin/insulin like growth factor (Ins/IGF) signaling pathway[6]. Circulating insulin and IGF levels give an indication of the caloric and nutrient condition of the mature environment, allowing

modulation of developmental rate to match the prevailing conditions and improve survival

prospects.

Growth factor receptor bound protein 10 (GRB10) is a negative regulator of the Insulin/IGF

signaling pathway. GRB10 downregulates the growth response, promoting a switch from

glucose to fat metabolism and halting cell cycle progression[7–9]. GRB10 expression limits

placental growth and efficiency[10] and correlates with small body size [11] in mammals.  In

humans, GWAS show *GRB10* is associated with TIID[12], and *GRB10* copy number variation is

associated with Silver Russell Syndrome[13], a rare growth disorder typically characterised by

intrauterine growth restriction, SGA status, hypoglycemia, poor muscle development, and

increased fat deposition. The role of GRB10 in the regulation of human growth is notably

associated with response to recombinant human growth hormone in children with growth

hormone deficiency, where lower GRB10 expression correlates with a greater response[14,15].

Variability in Grb10 expression is also linked to the dramatic range of body sizes observed

between cetaceans[16]. Average daily mass gain is elevated in beef cattle with a *grb10*

associated deletion[17], and *grb10* SNPs impact muscle and lipid mass, angularity, and body

conditioning score[18]. Global *grb10* knockout in mice correlates with a "leaner" phenotype,

including elevated muscle and reduced lipid mass, and insulin sensitivity[19,20]. Therefore,

understanding the impact transient disruption to early-life growth trajectory has on mature

organism size, average daily gain, and lipid to muscle ratio may provide the groundwork for

boosting meat yield and quality, necessary to match the growing global demand for protein.

In this study, *grb10a* expression was transiently supressed in wild-type zebrafish (*Danio*

*rerio*) embryos by antisense oligonucleotide directed blocking of mRNA splicing, resulting in

increased insulin/IGF signalling during development. The importance of *grb10a* as a

coordinator of growth, metabolism, and cardiac health was assessed over the first 5 days

post fertilisation (dpf). The impact of early-life growth disruption on the transcriptomic

landscape was also investigated, together with the lasting impact on later-life body

morphology, metabolism, and cardiac phenotype, linking together the distinct pathways

commonly associated with the FOAD hypothesis.

**Materials and Methods**

**Zebrafish Husbandry**

AB zebrafish were maintained under standard conditions (≈28 °C; 14/10 h light/dark cycle; <

5 fish per litre) within the Biological Services Unit of The University of Manchester.

Regulated procedures received ethical approval and were performed under a Home Office

Licence (PPL P005EFE9F9). To generate embryos, breeding pairs of similar ages were

selected at a ratio of 1 male to 1 female and fasted overnight in breeding tanks. Dividers

were removed at the start of the following light cycle and embryos were collected after 20

minutes of free breeding. Embryos were kept at a stocking density of < 50 per petri dish and

raised in embryo water (Instant Ocean salt 60 µg/mL) up to 5 dpf and transferred to the

main aquarium.

**Transient Knockdown of *grb10a* Expression**

Morpholino-modified antisense oligonucleotide knockdown (KD) of grb10a was validated in

accordance with current guidelines for morpholino use in zebrafish[21]. Morpholinos targeting

exon three (e3i3) and four (e4i4) were designed by and obtained from Gene Tools, LLC

(Philomath, OR, USA) along with a standard control (SC) oligonucleotide targeting human β-

globin, used to control for microinjection (sequences - *Table 1,* microinjection solutions -

*Table 2*[21]). Phenol red and nCerulean (nuclear-targeting blue fluorescent protein) mRNA

were included to ensure successful injection. Embryos received a single injection into the

yolk directly below the cell mass at the single-cell stage, as per established methods[21].

Embryos were screened for fluorescence at 48 hours post-fertilisation (hpf) to ensure

constitutive and even uptake of the injection material. Non-uniformly or weakly-fluorescent

embryos were removed.

**Validation of *grb10a* Knockdown**

Primer sequences, outlined in Table 1, were designed using SnapGene® (GSL Biotech, San

Diego, CA, USA) and synthesised by Thermo Fisher Scientific (Waltham, MA, USA). Specificity

was confirmed using Primer BLAST[22]. To confirm antisense oligonucleotide activity, RNA was

extracted from pooled zebrafish embryos at 24, 48, 72, 96, and 120 hpf (n=3, 5 embryos per

pool). Extraction was performed using QIAGEN RNeasy lipid extraction kit according to the

manufacturer's instructions, and cDNA was generated by reverse transcription using the

ProtoScript® II First Strand cDNA Synthesis Kit (NEB). cDNA was amplified with primers

flanking each splice site by Taq polymerase (NEB, Hitchin, UK) (thermocycling parameters

outlined in Table 3). β-actin (*actb1*) was used as a positive control for cDNA integrity.

**Analysing Downstream Signalling by Western Blot**

96 hpf embryos were deyolked in Ringer's Buffer to limit background interference (n=15,

performed in triplicate). Embryos were resuspended in 100 µl RIPA buffer (150 mM NaCl,

1% Nonident P-40, 0.5% Sodium deoxycholate, 0.1% SDS, 25 mM Tris pH 7.4) containing

protease and phosphatase inhibitors, and homogenised. Samples were incubated on ice for

30 minutes, clarified by centrifugation at 4 °C, and denatured at 98 °C for 5 minutes in

Laemmli buffer (2% SDS, 10% glycerol, 60 mM Tris-Cl, 0.01% bromophenol blue, 0.1% β-

Mercaptoethanol). Proteins were separated by 10% SDS acrylamide gel electrophoresis and

transferred using established methods. The transfer membrane was incubated in blocking

buffer (3% BSA in TBS-T) for one hour, primary antibody (Table 4) at 4 °C under constant

agitation overnight, and secondary antibody (Table 4) for one hour at room temperature.

The membrane was covered with ECL Western Blotting Substrate (Promega, Southampton,

UK) and imaged immediately. Protein expression was quantified from band intensity in

ImageJ[23].

**_grb10a_ mRNA for Overexpression and Rescue**

Total RNA was extracted, and cDNA generated from a pool of 96 hpf embryos, as previously

described. _Grb10a_ was amplified from cDNA by high specificity PCR (NEB Q5 Hot Start) with

primers flanking open reading frame (Table 1). The product was purified using QIAGEN

Quick Gel Extraction, blunt ligated into pCR-Blunt II-TOPO (Thermo Fisher Scientific) and

transformed into _E. coli_ following an established protocol. The purified plasmid was Sanger

sequenced to confirm successful cloning. The insert was liberated by digestion with _Cla I_ and

_Xba I_ restriction enzymes (NEB) and subcloned into pCS2+[24]. Capped RNA was generated

using the mMESSAGE mMACHINE® SP6 Transcription Kit (Thermo Fisher Scientific) according

to the manufacturer's instructions. RNA was purified by MEGAclear™ Transcription Clean-

Up. 1 µl of the purified RNA was analysed by gel electrophoresis to confirm amplification

and structural integrity.

**Embryonic Physiological and Metabolic Measurements**

Whole body length, the longest straight-line distance between the snout and tip of the

notochord, and yolk area measurements were taken at 24-hour intervals from 48 to 120

hpf. Embryos were dechorionated and acclimatised for ≥ one hour before imaging. Images

were imported into ImageJ, which was calibrated with an image of a graticule of known size.

Data were imported into GraphPad Prism version 7.00 for Windows (GraphPad Software, La

Jolla, CA, USA, www.graphpad.com).

To investigate embryonic cardiac phenotype, embryos were sedated using an anaesthetic

concentration known to have no impact on cardiac function (0.04% MS-222 solution[25]).

Heart beats were counted over a 20 second period and converted to beats per minute

(bpm).

A Glucose Uptake-Glo TM Assay (Promega) was performed on 96 hpf zebrafish to detect

differences in metabolic rate. Individual embryos (n = 5 per treatment) were injected with 1

mM 2-deoxyglucose-6-phosphate directly into the yolk and allowed to recover for 30

minutes. Embryos were processed according to an established protocol[26]. Luminescence

was measured by plate reader with 8 readings per well. Readings were adjusted for

background luminescence.

**Quantitative PCR**

QPCR primers amplifying *grb10a* and markers of cardiac dysfunction were designed (Table

1) and their efficiency validated[27]. RNA was extracted from pooled (n=10) embryonic

zebrafish and pooled (n=3) adult (> 1 year) heart samples by QIAGEN RNEasy Lipid Tissue

Extraction kit according to the manufacturer's instructions. Samples were repeated in

triplicate and tested for gene expression by qPCR (Applied Biosystems Power SyBr Green)

(Table 3). Relative fold change in gene expression was calculated using the ΔΔCt method

according to the following equation:

$$\Delta\Delta Ct = \Delta Ct_{e3i3} - \Delta Ct_{SC}$$

where

$$\Delta Ct = Ct_{gene\ of\ interest} - Ct_{housekeeping\ gene}$$

173    and

$$Fold\ change = 2^{-\Delta\Delta Ct}$$

Data were imported into GraphPad and unpaired t-tests were performed on ΔCt values to

assess the statistical difference between samples.

**Transcriptomic Analysis**

Age-associated changes in gene expression were assessed by computational analysis of

transcriptomic data. Pooled samples (n=5) were generated from each treatment group (SC,

KD) at 5, 10, 15, 20, and 30 dpf, with three repeats per sample. Zebrafish were culled under

terminal anaesthesia, and RNA was extracted from tissue anterior to the gills.

Transcriptomic data were generated using Affymetrix Zebgene 1.0st arrays. Data for all

75212 gene probes were imported into Qlucore Omics Explorer 2.2 (Lund, Sweden) as .cel

files and normalised using the robust multi-array average (RMA) approach with a gene level

summary. Zebrafish gene identities were assigned using Affymetrix gene definitions. Human

orthologues (GRCh 38) were mapped using the *biomaRt* R-package[28]. A workflow pipeline of

the transcriptomic analyses is outlined in *Supplementary Figure 1*.

Unsupervised analysis of gene expression by age group was conducted by generating

hierarchically clustered heat maps. Standard deviation filtering (standard deviation of

specific gene expression divided by maximum gene standard deviation $[s/s_{max}]$) was

performed on the dataset to remove genes with low variance, as these were unlikely to be

informative. Projection scores[29] were used to determine the threshold for filtering by

calculating the maximum separation in principal component analysis (PCA).

**Hypernetwork Modelling**

Hypernetwork analysis was performed to investigate higher order interactions between target

genes, a general model of which is outlined in *Supplementary Figure 1*[30]. All analyses were

performed in R (version 3.4.2). Pearson's correlation coefficients ($r$) were calculated

between age-associated genes identified by unsupervised analysis ($g$, KD = 119, SC = 297)

and the rest of the transcriptome ($g^c$, KD = 75093, SC = 74915). R-values were binarized to

generate the incidence matrix of the hypernetwork ($M$). Positive and negative correlations

greater than $\pm 1$ standard deviation (sd) from the mean of the R-values were assigned as '1' (i.e.,

present) and values closer to zero were assigned '0'. In this way, each element ($\in$) of $g$ can be

described as:

204
$$\in \text{ of } g \text{ in } M = \begin{cases} 1, & x > \pm 1sd|r| \\ 0, & x \leq \pm 1sd|r| \end{cases}$$

The resulting binary incidence matrix ($M$) was multiplied by its transpose ($M^t$) to generate

the adjacency matrix of the hypernetwork ($M.M^t$) which quantifies the shared correlations

between any pair of genes ($g$). This hypernetwork adjacency matrix represents the higher

order interactions between pairs of genes in a manner not captured by traditional

transcriptomic approaches. As a measurement of co-ordination, hypernetworks have been

suggested to model functional relationships[30]. Coordination between age-related genes and

the rest of transcriptome was investigated by interrogating the incidence matrix of the

hypernetwork and identifying a subset of edges and nodes which could form a complete

subgraph. This represents the subset of the transcriptome ($\subset g^c$) showing correlated

expression with all the transcripts from the hypernetwork central cluster ($\subset g$).

**Quantification of Network Topology**

All subsequent analyses focused on genes defined as the central cluster of the

hypernetwork (86 genes in SC fish, 67 in KD) of 20-30 dpf zebrafish. Correlation networks

model functional relationships within gene networks[32] and allow identification of clusters of

highly connected genes[33]. Quantification of hypernetwork properties (connectivity and

entropy) was performed on the hypernetwork ($M.M^t$), where connectivity is the sum of

connections shared by each element of the network ($g$) and entropy is the degree of

disorder within the distribution of shared correlations. Entropy is positively correlated with

the cellular differentiation potential[34], where a high entropy indicates an earlier cell lineage

and multiple potential signalling pathways, and low entropy indicates a more specific

function. Entropy has also been used as an index of regularity and patterning[35]. Entropy was

measured using R package BioQC[36].

To identify the impact of age-association on the transcriptome and compare the effect

between the SC and KD data sets, connectivity and entropy were calculated and averaged

for 1000 hypernetworks of genes randomly selected from each dataset (having no age-

association). The difference in connectivity and entropy that age-association conveyed on

the transcriptome could then be determined by comparing the age-associated datasets with

their corresponding random datasets. As entropy scales proportionally with the size of a set,

entropy was normalised for each dataset, generating a proportional measure of the

maximum entropy possible for a given set size (ranging from 0 - 1), calculated by dividing

the measured entropy by log(n) where n is the set size.

## Gene Ontology

Gene set enrichment analysis (GSEA) was performed[37] to associate gene expression with biological processes. Genes were mapped to human orthologues using Qlucore and GSEA was performed to rank genes by age group associated ANOVA p-values. Additional GSEA was carried out through Webgestalt[38] using genes ranked by R-value, derived from a rank regression analysis of gene expression against age. Over-representation analysis (ORA) was used to identify gene ontology associated with unranked gene sets (Webgestalt). All gene ontology analysis used the GO Biological Process Ontology gene list[37,39].

## Hypernetwork Modelling of Ontology

Pathways identified by GSEA were modelled using a hypernetwork approach to investigate the association between each pathway and the two treatment groups. Hypernetworks were generated separately for SC and KD fish using human genes associated with each pathway[40]. Pathways with fewer than 15 associated genes were removed, and all remaining gene sets were converted to zebrafish homologues using the Ensembl database (release 104)[41], queried using BiomaRt for R[28,42].

Hypernetworks were generated on 10 genes from each pathway, iterated 1000 times. Hypernetwork entropy was assessed on each iteration. A Bayesian approach was used to model the entropy distributions for each pathway to identify differences between SC and KD. This was performed using Bayesian generalized linear modelling via the r package rstanarm[43,44]. Differences between SC and KD entropy distributions were calculated as a $\beta$ value and significance was assigned to pathways for which the 89% credible interval of the beta values did not include 0, as per established methods[45].

**Stop-Flow Respirometry**

Individual 30 dpf zebrafish were placed into one of four stop-flow respirometry chambers

(volume 2 ml) and allowed to acclimate at 28 °C for > one hour. Optical oxygen sensors

paired with oxygen sensor spots (Pyroscience, Aachen, Germany) were used to measure

oxygen saturation within the chambers and recorded using a FireStingO2 Fiber-optic oxygen

and temperature meter, simultaneously recording and maintaining temperature at 28 ± 0.3

°C. Probes were calibrated according to the manufacturer's instructions. Chambers were

randomised per trial, with one chamber left empty to correct for background bacterial

respiration. Chambers were refreshed immediately prior to the experiment until oxygen

saturation measured 100%. Oxygen consumption curves were recorded in triplicate with

five trials per individual. Chambers were manually refreshed when oxygen saturation

reached 80%. To calculate the rate of oxygen consumption, linear regression in Microsoft

Excel was used to calculate the change in oxygen saturation during each trial. This was

normalised against the dry mass of the subject, length of the trial, and volume of the

respirometry chamber according to the following equation:

$$Oxygen\ consumption\ (\mu gO_2 h^{-1}g^{-1}) = \frac{\Delta O2\% \times \frac{7900\mu gL^{-1}}{100} \times chamber\ volume\ (L)}{time\ (h)\ \times\ mass\ (g)}$$

Where 7900 $\mu gL^{-1}$ is equivalent to 100% oxygen saturation at 28 °C.

A closed-circuit stop-flow respirometer was used to investigate aerobic scope in adult

zebrafish (18 months). Sealed chambers (70 ml) were combined with recirculation loops

containing an oxygen flow-through cell (Pyroscience, Aachen, Germany), paired with an

optical oxygen sensor (Pyroscience, Aachen, Germany), calibrated according to the

manufacturer's instructions. A stop-flow pump, controlled by a Cleware USB-Switch

(Cleware GmbH, Germany) programmable switch and AquaResp v.3 software (AquaResp, v3,

Python 3.6[46]) was incorporated into the circuit to automatically refresh the water in the

chambers after each trial (60 s flush, 30 s wait, 300 s measure). To maintain a constant

temperature of 28 °C ± 0.3 °C, the respirometry system was immersed in a recirculation

chamber under constant aeration. Oxygen saturation and water temperature were recorded

as previously described. Regression curves were automatically generated by AquaResp, and

oxygen consumption was extracted and normalised against the length of the trial and

volume of the respirometry chamber. Zebrafish were manually stressed for 2 minutes

immediately prior to the start of the first trial. Maximum oxygen consumption was

identified as the trial with the greatest difference in oxygen saturation. Standard metabolic

rate was calculated as the mean of the lowest 10% of the trials[47]. Aerobic scope was

calculated as the difference between the maximum oxygen consumption and standard

metabolic rate. Data are presented as individual data points alongside the mean and SEM.

**Glucose Tolerance and Insulin Sensitivity**

Adult zebrafish (18 months) were fasted overnight and allocated to either glucose tolerance

testing or insulin sensitivity testing. All blood samples were acquired following the protocol

outlined by Zhang et al.[48]. Mass, body length, and fasting blood glucose were measured

immediately prior to the start of the protocol.

Fish were anaesthetised in 0.02% MS-222, placed on their side and patted dry. A single IP

injection of glucose (0.5 mg glucose/g) or glucose and insulin (0.5 mg glucose/g and 0.0075

U insulin/g) was performed before recovery in 28 °C system water[49,50]. Blood samples were

taken at 30- and 120-minutes post-injection (glucose tolerance) or 30- and 60-minutes post

injection (insulin sensitivity). Blood samples were immediately tested for blood glucose

concentration (Sinocare Safe AQ blood glucose monitor). Individuals were culled in MS-222

during the final blood draw. Data are presented as the mean and the SEM (glucose

tolerance n = 10, insulin sensitivity n = 8-12).

**Adult Physiological Measurements**

To assess the end-stage body morphology induced by *grb10a* KD, dry mass and body length

were measured in adult (18 month) zebrafish and Fulton's condition factor was calculated.

Body length was measured as the greatest straight-line distance between the snout and the

end of the tail. The caudal fin was not included as fin length can be influenced by factors

such as damage or variation between strains. Fulton's condition factor was calculated

according to the formula[51]:

$$K = \frac{M}{L^3} \times 100$$

Where M = mass in grams and L = length in centimetres. Higher K values indicate thicker,

more rounded bodies. Data are presented as the mean ± SEM (n = 21-34).

**Histology**

Whole adult zebrafish (18 months) were embedded longitudinally in paraffin wax. 5 µm

sagittal sections were taken from each tissue at a consistent depth and stained with

Masson's Trichrome (IHC World Masson's Trichrome Staining Protocol for Collagen Fibres,

Woodstock, MD, USA) to differentiate skeletal muscle (red), connective tissue (blue), and

nuclei (black). Slides were scanned and visualised at 20x magnification (3D Histech

CaseViewer v2.4.0.119028, Budapest, Hungary).

Skeletal muscle measurements were performed on a site lateral to the dorsal fin. The

perpendicular width of individual muscle fibres was recorded and are presented as the

mean of five measurements of each muscle fibre (ten fibres from five individuals, 50 fibres

total).

Red blood cells were digitally removed from images of the ventricle before importing into

ImageJ. The ratio of compacta to spongiosa was calculated by measuring the area of each

tissue type. Data are presented as the ratio of compacta to spongiosa as a percentage.

Cardiac tissue density was calculated by restricting the region of interest to the boundary of

the ventricle and calculating the total number of pixels in the image. Threshold_Colour was

used to threshold the images, which were converted to 8-bit black and white images.

Voxel_Counter.class was used to calculate the number of black pixels in the image.

**Statistical Tests**

For transcriptomic analyses, rank regression (least squares method) was used to generate

the most appropriate linear model for each probe (the smallest degree of variance over the

sample). Multi group analysis of variance (ANOVA) was used to associate each gene ID with

time dependent gene expression. Wilcoxon rank sum test (ggpubR package for R[52]) was

used to test for differences in network topology. False discovery rate (FDR) adjustment was

made using the Benjamini-Hochberg method and applied to the gene ontology analysis[53].

All data were ROUT tested[54] for outliers and subject to D'Agostino and Pearson normality

tests. All comparisons between SC and KDs were performed using unpaired t-tests.

Comparisons of multiple groups were performed using one-way ANOVAs. Post-hoc power

calculations were performed to confirm sample sizes were sufficient, where $\alpha = 0.05$.

**Results**

**Knockdown of *grb10a* Expression by Splice-Blocking Antisense Oligonucleotide**

As *grb10* has been linked to embryonic growth trajectory, expression was examined over

the first 120 hpf. QPCR analysis of *grb10a* expression at 24-hour intervals revealed a strong

upregulation at 48 hpf (*Figure 1a*). Expression of the *grb10* paralogue, *grb10b*, was not

detectable at any time point.

To knock down *grb10a* expression, zygotes were microinjected with splice-blocking

antisense oligonucleotides e3i3 and e4i4. Exon 3 and 4 donor splice sites (*Figure 1b*) were

targeted in order to confirm the specificity of the phenotype, in accordance with current

guidelines[21]. Multiplexed RT-PCR amplification using primers flanking the splice sites (*Figure*

*1b*) showed a single product of the anticipated size for e3i3 and e4i4 embryos, and no

product was detected for SC embryos (*Figure 1c*), consistent with successful incorporation

of the corresponding intron.

To confirm grb10a KD induced a quantifiable impact on the downstream insulin signalling

pathway, phosphorylation of key proteins was analysed by Western Blot. As shown in *Figure*

*1d*, phosphorylated (active) versus total protein ratios of AKT and S6 were significantly

elevated in *grb10a* KD zebrafish at 96 hpf compared to SC (p = 0.0007 and 0.0413

respectively, n=3), consistent with the expected impact of grb10a KD.

**Growth Trajectory and Early Life Cardiometabolic Phenotype is Significantly Impacted by**

**Transient *grb10a* Perturbation**

To determine the effect of *grb10a* KD on growth, total body length was measured at 24-

hour intervals over the first 5 dpf. As shown in *Figure 2a,* total body length was initially

comparable between KD and SC zebrafish (2.857 ± 0.0549 mm vs 2.826 ± 0.0962 mm,

p=0.7896, n=9 and n=10 respectively). Subsequently, KD zebrafish began to diverge from the

SCs at 48 hpf, corresponding to the peak in *grb10a* expression observed in WT zebrafish

(*Figure 1a*). KD zebrafish were longer on average than SCs (3.411 ± 0.0165 mm vs 3.177 ±

0.0231 mm at 72 hpf, p<0.0001, n=46 and n=41 respectively). This phenotype was reversed

in zebrafish overexpressing *grb10a,* which were significantly shorter than SC counterparts

(3.361 ± 0.0239 mm vs 3.505 ± 0.0339, p=0.001, n=24 and n=25 respectively), as shown in

*Figure 2b*. Co-injection of e3i3 and *grb10a* RNA returned body length to SC levels (3.505 ±

0.0339 mm vs 3.564 ± 0.0265 mm, p=0.3792, n=25), confirming the validity of e3i3 induced

*grb10a* KD. Moreover, *grb10a* KD induced by e4i4 (*Figure 2c*) also resulted in increased body

length, and the ability of *grb10a* overexpression to suppress growth was shown to be dose

dependent (*Figure 2c*). Intriguingly, by 120 hpf, body length converged, indicating activation

of compensation to regulate growth and return to an "ideal" length post-hatch.

To investigate the impact of *grb10a* KD on the developing cardiac system, heart rate was

measured over the first 5 dpf. As shown in *Figure 2d*, average heart rate began to diverge

between the treatment groups at 48 hpf, again correlating with the WT peak in *grb10a*

expression. While heart rate increased slightly over time in SC zebrafish, in-line with

increasing body size, average KD heart rate fell. By 120 hpf, average heart rate was almost

50% lower in the KD compared to SC (61.5 ± 6.97 bpm vs 118.4 ± 2.83 bpm, p<0.0001, n=14

and n=19 respectively).

To determine whether *grb10a* KD had an impact on metabolic rate, yolk absorption was

measured to indicate energy demand. As shown in *Figure 2e*, there was initially no

difference in yolk area between the groups (p=0.8185, n=10). As with body length and heart

rate, SC and KD yolk consumption began to diverge at 48 hpf (0.298 ± 0.0055 mm$^2$ vs 0.390

± 0.0104 mm$^2$, p<0.0001, n=18 and n=19 respectively). The elevation in yolk consumption

observed in the KD fish suggests an elevated metabolic rate. To support this conclusion, a

Glucose Uptake-Glo™ Assay was performed to compare the rate of glucose uptake. As

shown in *Figure 2f*, 2D6P accumulation was significantly higher in KD zebrafish compared to

SC, an increase of almost 30% (p=0.0002, n=5), indicating glucose uptake was elevated.

These findings are consistent with the role of *grb10a* as a negative regulator of the insulin

signalling pathway[55] and a coordinator of growth and metabolism.

**Transient *grb10a* Knockdown Persistently Dysregulates Age-Associated Gene Expression**

To understand whether the observed changes were coupled with lasting changes in gene

expression, the transcriptomic landscape of SC and KD zebrafish was investigated over the

first 30 dpf. Unsupervised hierarchical clustering, standard deviation filtering, and

maximised projection scores (MPS) were used to define a set of genes with strong age-

association in the SC zebrafish (297 genes, MPS = 0.43). These genes fell into four distinct

clusters, associating with 5, 10, 15, and 20-30 dpf (163, 15, 32, and 87 genes respectively)

(*Figure 3a*).

When performing the same analysis in KD zebrafish, the 5 dpf cluster was significantly

disrupted (*Figure 3b*). While the clustering was largely conserved in the latter three clusters

(10 dpf – 14/15, 15 dpf - 29/32, and 20-30 dpf – 74/87 genes), genes strongly associated

with 5 dpf in the SCs were generally expressed at different time points in the KDs (dotted

white lines, *Figure 3a* and *Figure 3b*), with a mapping of only 38/163 genes (23%). Notably,

five genes associated with 5 dpf in SC zebrafish mapped to 20-30 dpf in the KD. This

suggests the expression of these genes is usually associated with early larval development,

but instead, in the KDs, is associated with the late-juvenile stage. Human orthologues of

these dysregulated genes include *DGAT2* (fatty acid metabolism), *GAMT* (energy storage,

muscle contraction, and fatty acid oxidation), and *PDIA2* (thiol-disulphide interchange,

particularly in the pancreas).

As the SC gene clusters were disrupted in the KD dataset, unsupervised analysis of the KD

data was performed to identify the subset of age-associated genes in the KD zebrafish. 119

genes were identified (MPS = 0.37) which segregated into 5-15, 15-30, and 20-30 dpf (37,

16, and 66 genes) (*Figure 3c*).

To assign functionality to these age-related genes, Gene Set Enrichment Analysis was

performed for genes identified in both the SC and KD datasets. Functionality conserved

between the SC and KD is described in *Supplemental Table 1*. Dissimilar pathways (*Figure*

*3e*) included several actin and collagen related pathways, and extracellular structure and

RNA processing, which were age-related in KD but not SC zebrafish. Conversely, several

metabolic pathways were age-associated in SC zebrafish but not in KD animals. This clear

dysregulation of age-related gene expression in the KD zebrafish implies transient *grb10a*

KD induces persistent remodelling of the transcriptome.

**Transient *grb10a* Knockdown Disrupts Transition Between Larval Gene Clusters**

To investigate the co-ordination of the whole transcriptome with age-associated gene

expression, hypernetwork models were constructed for the SC and KD datasets based on

the gene sets identified in the cluster analysis (297 and 119 genes respectively). This

approach has been used to model functional relationships[31]. *Figure 4* describes the results,

where colour intensity represents the number of shared interactions between each gene

pair. In SCs, shared interactions segregated into three groups, correlating to 5, 10-15, and

20-30 dpf (161/297, 50/297, and 86/297 genes) (*Figure 4a*). In KD animals, however, genes

segregated into only two groups, either 5-15 dpf or 20-30 dpf (31/119 and 65/119 genes)

(*Figure 4b*). Notably, the large group of co-ordinated interactions at 5 dpf was absent in the KD dataset, with a combined group of genes correlating with 5-15 dpf identified instead.

## *Grb10a* Knockdown is Associated with Increased Connectivity and Crosstalk Within the Transcriptome

To quantify these differences in the co-ordination of the transcriptome, two network topology parameters were used: connectivity and entropy. Hypernetwork connectivity quantifies the number of higher order interactions within the transcriptome and is related to function[31,56], while entropy is a measure of information content and serves as an indicator of "disorder". I.e. lower entropy (more order) describes a network with little crosstalk and a more discrete function[34,57], while a network with greater entropy has increased crosstalk and pleiotropic functions.

Changes in connectivity (*Figure 4c*) and entropy (*Figure 4d*) were calculated for genes identified as active at 20-30 dpf by the previously described analyses. The impact of KD on the co-ordination of age-associated genes in the network was identified by comparing the connectivity and entropy of SC and KD datasets. To provide a comparison between age-associated genes and genes associated with other functions, connectivity and entropy were also calculated for randomly selected gene sets (*Supplemental Figure 2*).

Genes associated with 20-30 dpf were more highly connected and less entropic (more ordered) than random genes in both the KD and SC datasets ($p < 2.2 \times 10^{-16}$). Comparison of age associated genes in KD and SC revealed a higher entropy (1.05-fold, $p < 2.2 \times 10^{-16}$) and higher connectivity in the KD (1.20-fold, $p < 2.2 \times 10^{-16}$). This suggests age-associated genes in the KD zebrafish transcriptome have more interactions and are less ordered (have more crosstalk) than in the SC.

**Pathways Associated with Transcriptome-Wide Remodelling Support an Alteration in Cardiometabolic Phenotype**

Having identified a core set of age-related genes, a broader set of genes which coordinate with these genes was defined as the complete subgraph between the central age-related genes of the hypernetwork and their correlates. 12775 (KD) and 459 (SC) genes were defined in this set from the 20 to 30 dpf hypernetwork clusters described previously, all significantly associated with age (rank-regression: KD q < 1.50 x10$^{-6}$, SC q < 4.44 x10$^{-2}$; *Supplementary Table 2*). 28 times more genes were implicated in the KD. Both datasets showed a skew towards a positive association with age (68% SC vs 59% KD) (*Figure 5a, Figure 5b*). The overlap in gene expression between the two datasets was 244 (77.9% of SC) and 60 (41.1% of SC) in the positively and negatively correlated sets, respectively. Thus, genes in the wider transcriptome with co-ordinated expression at 20-30 dpf in SC zebrafish demonstrated a similar pattern of expression in KD (*Figure 5c*). However, the inverse is not true, as genes with co-ordinated expression at 20-30 dpf in the KD zebrafish were dysregulated in the SC (*Figure 5d*).

To assign functionality, GSEA was performed as previously described (ranked by R-value, top 100 pathways with weighted set cover). Results of this ontology analysis are outlined in *Figure 5e* and *Figure 5f*. The two datasets featured distinctly different regulated pathways. RNA processing, a variety of metabolic pathways, and cardiovascular development featured in the KD dataset, whereas growth and immune signalling were featured in the SC dataset. Functional analysis of these determined that the greatest differences in activity between the KD and SC were in *Positive Regulation of Lipid Localization*, *Amino Acid Transport* and *Cellular Modified Amino Acid Biosynthetic Process* (*Figure 6*).

A distinct pattern of gene expression was identified in the group of genes dysregulated at

20-30 dpf in the KD dataset (white box, *Figure 5d*). These genes were upregulated at 15 dpf,

downregulated at 20 dpf, and re-upregulated at 30 dpf. This subgroup of 3460 genes

(*Supplemental Table 3*) was associated with a range of gene ontologies related to

metabolism and development (*Figure 5g*) and was synchronous with the spike in growth

identified in *Figure 7a*. Specific pathways following this pattern of expression included

cardiovascular system development, muscle structure development, and developmental

maturation.

**The Impact of Transient *grb10a* KD Persists into Adulthood**

As *grb10a* KD significantly impacted early-life growth, metabolism, and cardiovascular

development and was associated with persistent remodelling of the transcriptome,

investigation was conducted into the phenotypical differences in older zebrafish.

Total body length measurements up to 30 dpf are shown in *Figure 7a*. Growth was

significantly elevated in the KDs compared to SCs between 15 and 20 dpf (30% increase,

7.493 ± 0.2726 mm vs 4.320 ± 0.1594 mm, p < 0.0001, n = 10). This corresponded to the

cluster of dysregulated genes identified by hypernetwork modelling (*Figure 5g and white*

*box in Figure 5d*). The growth profile of the KDs was shifted to an earlier age compared to

the SC, suggesting a faster rate of maturation. The GSEA highlighted a reduction in activity in

developmental maturation pathways (normalised enrichment score < 1) in SCs vs. KDs,

which was reflected in the growth rate at approximately 20 dpf.

Late-life body length and mass measurements were recorded in 18-month zebrafish to

investigate the lasting impact of *grb10a* KD on the phenotype. Final body length (*Figure 7b.*)

was higher, with KDs 1.7 mm longer than SCs (3.20 ± 0.029 cm vs 3.03 ± 0.021 cm, p <

0.0001). These fish were also approximately 10% heavier (0.54 ± 0.012 mg vs 0.49 ± 0.011

mg, p = 0.005). On the other hand, the Fulton's condition factor (an indicator of body

condition) was closer to 1 in the KDs (1.64 ± 0.026 vs 1.75 ± 0.034, p=0.0215) despite the

increase in mass. To equate this to differences in body composition, skeletal muscle fibres,

isolated from the base of the dorsal fin, were embedded and stained with Masson's

Trichrome (*Figure 7c.*). The average muscle fibre diameter was approximately 20% greater in

the KDs (27.01 µm ± 0.728 vs 22.71 µm ± 0.698, p < 0.0001), likely contributing to the

increase in mass despite a decrease in roundness.

The impact on cardiac health was assessed by qPCR and histology. As shown in *Figure 8a*,

*myl7* expression (an index of muscle mass and hypertrophy) in the heart was over 20%

greater in KDs (p < 0.0001, n = 3), while *nppa* expression (activated in response to

ventricular stress during hypertrophy and heart failure[58-60]) was reduced by approximately

40% (p = 0.0012, n = 3). There was no difference in *pcna* expression (proliferating cell

nuclear antigen), suggesting there was no difference in proliferation in the cardiac tissue. To

support these findings, ventricular morphology was also assessed. As shown in *Figure 8b*,

the ratio of compact myocardial layer to trabeculated was significantly greater in the KD

zebrafish (0.369 +/- 0.07 vs 0.185 +/- 0.03, p-value = 0.0288). This was coupled with an

overall increase in tissue density (76.3% vs 64.4%, p-value = 0.0289).

To determine whether metabolic rate was persistently altered, stop-flow respirometry was

conducted on juvenile (30 dpf). As shown in *Figure 8b*, KDs consumed 7 times more oxygen

than SCs (134.4 ± 15.60 $\mu gO_2h^{-1}g^{-1}$ vs 19.0 ± 3.28 $\mu gO_2h^{-1}g^{-1}$, p < 0.0001, n = 5), suggesting

metabolic rate remained significantly elevated in juvenile fish.

To determine whether this was a permanent change, adult (18-month) zebrafish were also

investigated. Peak and basal oxygen consumption were recorded to further investigate the

"lean" phenotype. Maximum oxygen consumption, achieved following exhaustive activity,

was elevated by ~25% in KDs, as shown in *Figure 8c* (1645 ± 90.11 mgO$_2$kg$^{-1}$h$^{-1}$ vs 1309 ±

26.48 mgO$_2$kg$^{-1}$h$^{-1}$, vs p < 0.0001). There was no significant difference in basal metabolic rate

(486.5 ± 46.75 mgO$_2$kg$^{-1}$h$^{-1}$ vs 474.1 ± 27.78 mgO$_2$kg$^{-1}$h$^{-1}$, p = 0.8132). Consequently, aerobic

scope was greater in the KDs (1158 ± 100.7 vs 810.1 ± 42.95, p = 0.0007), supporting the

conclusion that the fish conform to a "leaner" phenotype.

As grb10a is involved in regulating the insulin signalling pathway, glucose tolerance and

insulin sensitivity tests were performed on adult zebrafish (18 months) to determine

whether there was a lasting biological impact. Fasting blood glucose (*Figure 8d*) was

significantly higher in KDs (7.96 ± 0.867 mmolL-1 vs 6.04 ± 0.493 mmolL-1, p = 0.0463).

Glucose tolerance testing showed both groups produced a similar response to glucose

challenge and were similarly able to respond to introduction of insulin.

**Discussion**

The first key finding from this study is that grb10a regulates embryonic growth in zebrafish,

consistent with its role in mammalian embryogenesis[47-49], and transient knockdown is

sufficient to have an impact on downstream pathways. This is consistent with its role as a

negative regulator of insulin signalling[50-51]. Western Blotting showed a clear upregulation in

insulin signalling pathway activation downstream of grb10a, which was coupled with

significant changes to the early phenotype, including elevated growth rate, increased

metabolism, and distinct cardiac alterations.

The observed changes in phenotype coincided with the peak in grb10a expression,

confirming the role of *grb10a* as a coordinator of embryonic growth and development. The

elevated growth coincided with higher metabolic rate, consistent with an increase in energy

demand due to a larger population of highly proliferating cells. Cardiac changes were also

present, supporting the conclusion that grb10a has a role in coordinating growth,

metabolism, and cardiac development. This is the first study, to our knowledge, to provide

*in vivo* evidence for the coordination of these distinct pathways.

This study also provides clear evidence for a compensatory growth mechanism activated

during early larval development, which regulates adherence to an "ideal" body length. Both

elevated embryonic growth and growth suppression were compensated for by 5 dpf, with

body length returning to comparable values in all treatment groups. This suggests there may

be a benefit to entering life at this controlled size. The altered phenotype established by day

5 served as a basis for life-long changes to the organism. Body length, metabolism, and

cardiac physiology were all significantly altered in 1.5-year-old zebrafish, despite no further

manipulation to the organism. This provides essential, novel, *in vivo* evidence for the Fetal

Origins of Adult Health hypothesis, which proposes that events during development can

have life-long impacts on an organism.

Furthermore, this study has shown for the first time that a transient disruption in the

expression of a single gene can result in permanent remodelling of the transcriptome. This

highlights the importance of regulated control during embryogenesis and the significant

impact small changes, such as increase in growth, can have on the developed organism.

Age-associated genes in both transcriptomes showed increased connectivity and decreased

entropy compared to non-age associated genes. This demonstrates that the genes identified

as age-associated are better connected to one another and more ordered in their

interactions than randomly selected genes. Entropy and connectivity were both higher in KD

than SC which demonstrates that age-associated genes share more interactions in KD but

with less discrete organization than in SC. These results support the conclusion that the KD

transcriptome is dysregulated, compared to the SC transcriptome.

Notably, gene expression at 5 dpf was significantly dysregulated in the KD zebrafish

compared to the SCs. An early transition in gene expression identified in the SCs, occurring

between 5 and 10 dpf, likely corresponding to a shift away from early developmental

pathways, was lost in KD zebrafish. This loss was reflected both in the age-related genes and

the coordination of the wider gene set. This fundamental difference in the transcriptome

during larval development was particularly notable in a subset of genes with fluctuating

gene expression in the KD zebrafish. Genes associated with cardiovascular development,

muscle development, and developmental maturation showed a distinct pattern of

upregulation at 15 dpf, downregulation at 20 dpf, and re-upregulation at 30 dpf, coinciding

with the spike in growth in the larval zebrafish. This pattern may contribute to the left-shift

in growth rate observed in the KD zebrafish, which is reflected in human cohorts

experiencing early puberty[61]. Furthermore, functional differences in cardiac and skeletal

muscle and metabolic phenotype were confirmed to persist into the adult zebrafish,

suggesting these transient changes in gene expression have long-lasting implications for the

organism.

This model of embryonic growth perturbation may also yield significant insights into the

propensity for early growth disruption to correlate with increased risk of cardiovascular and

metabolic disease in later life[62–65]. It is widely accepted that many disorders are likely to

have their origins during embryonic development[66–69], but the mechanisms involved are not

fully understood, and targeted *in vivo* research is lacking. Mammalian models have been

used to investigate the immediate impact of embryonic growth disruption[69–71], findings

which are replicated in this study. However, longitudinal studies are absent from the

literature, and little research has been conducted into the whole-life significance of early

growth disruption. Multiple distinct growth trajectories can yield similar birth weights,

including catch-up and catch-down growth[62] (both of which were achievable by modulation

of grb10a expression). Catch-up and catch-down growth have been reported to correlate

with increased risk of chronic health disorders[65,72,73], and as *grb10a* modulation is sufficient

to alter embryonic growth trajectory, metabolic rate, and cardiac function, the model

generated in this study may prove key in understanding the mechanisms involved in the

developmental origins of health and disease and identifying novel avenues for prevention

and treatment.

As identified in mammalian studies[19,20], *grb10a* KD was associated with a "leaner"

phenotype. KD zebrafish displayed an elevation in skeletal muscle fibre thickness and an

increase in body mass, likely as a result of altered body composition in favour of skeletal

muscle. Body condition also conformed to a leaner morphology in the KDs, where Fulton's

condition factor, used by fisheries and in research to assess the condition of fish stocks[74],

was closer to 1. Healthy fish in good condition have condition scores close to 1, while higher

or lower scores indicate overly fat or "skinny" fish respectively. Together, these findings

support the existence of a "lean" phenotype associated with suppressed grb10a activity and

greater insulin signalling activation. This is expected, as circulating insulin levels positively

correlate with growth, and serve as an indicator of resource availability in the mature

environment.

This study further extended the "lean" phenotype to include cardiovascular and metabolic

changes, the first *in vivo* work to combine these aspects. The establishment of an altered

cardiac phenotype during embryogenesis was followed by a notable change in ventricle

morphology. To maintain optimal cardiac output, a reduction in the rate of contraction must

be paired with increased cardiac efficiency and an increase in force or volume of

contraction. The increased demand on the cardiac tissue resulting from the reduction in

heart rate resulted in compacta thickening and an increase in ventricular density. This is

consistent with increased stroke volume and greater cardiac efficiency due to an increase in

contractility. Gene expression analysis of cardiac tissue, coupled with histological evidence,

supported this conclusion, and indicated the increase in musculature was due to an increase

in cell size rather than cell number. The greater degree of *myl7* expression in the absence of

an increase in *pcna* expression suggests an increase in hypertrophy (muscle size) rather than

hyperplasia (cell number) in the cardiac tissue.

Aerobic scope was also elevated as a result of grb10a KD, consistent with a "lean"

phenotype. This greater aerobic scope suggests the potential energy for non-essential

activities is elevated in KDs. As a result, these fish may have the capacity to outperform SC

counterparts in other energy-heavy tasks, such as swimming and courtship. The difference

in peak oxygen consumption between the groups suggests there is either increased oxygen

demand or an increased ability to extract oxygen from the environment in KD zebrafish.

Glucose homeostasis control was also altered in KD zebrafish. Glucose uptake was elevated

during embryogenesis and energy stores were depleted more rapidly, which was reflected

in elevation of fasting blood glucose in later life. Taken together with the elevation in

oxygen consumption observed in juvenile and adult zebrafish, this indicates greater

metabolic rate. This suggests KD zebrafish, while having a comparable basal energy demand,

have a greater demand for energy when performing non-essential activities. This may be

due to higher contribution of skeletal muscle to body composition, as heavily respiring

tissues require more energy when active than other tissues. KD zebrafish showed a greater

ability to maintain glucose homeostasis compared to SCs, responding more rapidly to

glucose challenge. As metabolic rate was greater in the KD zebrafish, this increase in the

rate of glucose clearance may be a direct consequence of heavily respiring tissue removing

glucose from the bloodstream at a higher rate.

Findings from this study may also provide important for the agri- and aquaculture industries.

The promotion of embryonic growth induced by *grb10a* KD may be a key alternative to

growth hormone treatment for improving meat yield and production efficiency, and larger

juvenile fish with greater aerobic scope are likely to be more capable of overwintering[75,76],

and thus improve fish stocks. This is particularly key, as the impact of climate change on

fisheries is an increasing global concern[77–79]. The increase in skeletal muscle contribution to

body composition also has relevance for industry, as lean products have greater value and

marketability, particularly as current health trends are increasing demand for low-fat

products.

**Conclusion**

This study proves for the first time that transient knockdown of *grb10a* expression is

sufficient to permanently alter growth trajectory, metabolic rate, and cardiac physiology.

We have presented significant evidence to suggest grb10a plays a previously unidentified,

fundamental role in the coordination of these distinct physiological pathways and could

represent a promising target for enhancement of meat yield and quality in agri- and aqua-

cultural.  Furthermore, this study shows that altering the expression of a single gene during

embryogenesis can remodel the entire transcriptome. We have shown that remodelling

established during embryogenesis provides a basis on which the adult phenotype is formed,

clearly demonstrating the lasting impact of early-life events on the whole life course of an

organism. This provides the first longitudinal *in vivo* support for the Foetal Origins of Health

hypothesis in *Danio rerio*. We have also expanded on the "lean" phenotype associated with

grb10 KO to include a distinctly altered metabolic phenotype and improved ventricular

efficiency. These life-long alterations may have ongoing implications for survival, and the

model generated in this study, featuring easily measurable phenotypic characteristics, a

short developmental window, and rapid generation time, will be indispensable for future

research into the mechanisms underpinning the foetal origins of health.

**References**

1.      Barker, D. J. P. Fetal origins of coronary heart disease. *BMJ* **311**, 171 (1995).

2.      Huss, M., Lindmark, M., Jacobson, P., van Dorst, R. M. & Gårdmark, A. Experimental

evidence of gradual size-dependent shifts in body size and growth of fish in response

to warming. *Glob. Chang. Biol.* **25**, 2285–2295 (2019).

3.      Hanson, M. A. & Gluckman, P. D. Early developmental conditioning of later health and

disease: physiology or pathophysiology? *Physiological reviews* **94**, 1027–1076 (2014).

4.      Ducsay, C. A. *et al.* Gestational hypoxia and developmental plasticity. *Physiol. Rev.* **98**,

1241–1334 (2018).

5.      Gluckman, P. D., Hanson, M. A. & Low, F. M. Evolutionary and developmental

mismatches are consequences of adaptive developmental plasticity in humans and

have implications for later disease risk. *Philosophical Transactions of the Royal Society*

*B: Biological Sciences* **374**, (2019).

6.    Harding, J. The nutritional basis of the fetal origins of adult disease. *Int. J. Epidemiol.*

**30**, 15–23 (2001).

7.    Murdaca, J. *et al.* Grb10 prevents Nedd4-mediated vascular endothelial growth factor

receptor-2 degradation. *J. Biol. Chem.* **279**, 26754–26761 (2004).

8.    Tezuka, N., Brown, A. M. C. & Yanagawa, S. ichi. GRB10 binds to LRP6, the Wnt co-

receptor and inhibits canonical Wnt signaling pathway. *Biochem. Biophys. Res.*

*Commun.* **356**, 648–654 (2007).

9.    Wick, K. R. *et al.* Grb10 inhibits insulin-stimulated insulin receptor substrate (IRS)-

phosphatidylinositol 3-kinase/Akt signaling pathway by disrupting the association of

IRS-1/IRS-2 with the insulin receptor. *J. Biol. Chem.* **278**, 8460–8467 (2003).

10.   Charalambous, M. *et al.* Maternally-inherited Grb10 reduces placental size and

efficiency. *Dev. Biol.* **337**, 1–8 (2010).

11.   Everson, T. M. *et al.* Placental expression of imprinted genes, overall and in sex-

specific patterns, associated with placental cadmium concentrations and birth size.

*Environ. Health Perspect.* **127**, 57005 (2019).

12.   Depetris, R. S. *et al.* Structural basis for inhibition of the insulin receptor by the

adaptor protein Grb14. *Mol. Cell* **20**, 325–333 (2005).

13.   Crippa, M. *et al.* Molecular Etiology Disclosed by Array CGH in Patients With Silver-

Russell Syndrome or Similar Phenotypes. *Front. Genet.* **10**, 955 (2019).

14.    Clayton, P. *et al.* A pharmacogenomic approach to the treatment of children with GH

711           deficiency or Turner syndrome. *Eur. J. Endocrinol.* **169**, 277–289 (2013).

712    15.    De Leonibus, C. *et al.* The in vitro functional analysis of single-nucleotide

713           polymorphisms associated with growth hormone (GH) response in children with GH

714           deficiency. *Pharmacogenomics J.* **19**, 200–210 (2019).

715    16.    Sun, Y. *et al.* Insights into body size variation in cetaceans from the evolution of body-

716           size-related genes. *BMC Evol. Biol.* **19**, 157 (2019).

717    17.    Xu, L. *et al.* Probe-based association analysis identifies several deletions associated

718           with average daily gain in beef cattle. *BMC Genomics* **20**, 31 (2019).

719    18.    Magee, D. A. *et al.* DNA sequence polymorphisms in a panel of eight candidate bovine

720           imprinted genes and their association with performance traits in Irish Holstein-

721           Friesian cattle. *BMC Genet.* **11**, 93 (2010).

722    19.    Smith, F. M., Garfield, A. S. & Ward, A. Regulation of growth and metabolism by

723           imprinted genes. *Cytogenet. Genome Res.* **113**, 279–291 (2006).

724    20.    Charalambous, M. *et al.* Disruption of the imprinted Grb10 gene leads to

725           disproportionate overgrowth by an Igf2-independent mechanism. *Proc. Natl. Acad.*

726           *Sci. U. S. A.* **100**, 8292–8297 (2003).

727    21.    Stainier, D. Y. R. *et al.* Guidelines for morpholino use in zebrafish. *PLoS Genet.* **13**,

728           (2017).

729    22.    Ye, J. *et al.* Primer-BLAST: a tool to design target-specific primers for polymerase

730           chain reaction. *BMC Bioinformatics* **13**, 134 (2012).

731     23.    Schindelin, J. *et al.* Fiji: An open-source platform for biological-image analysis. *Nat.*

732            *Methods* **9**, 676–682 (2012).

733     24.    Fink, M., Flekna, G., Ludwig, A., Heimbucher, T. & Czerny, T. Improved translation

734            efficiency of injected mRNA during early embryonic development. *Dev. Dyn. an Off.*

735            *Publ. Am. Assoc. Anat.* **235**, 3370–8 (2006).

736     25.    Avey, S. R., Ojehomon, M., Dawson, J. F. & Gillis, T. E. How the expression of green

737            fluorescent protein and human cardiac actin in the heart influences cardiac function

738            and aerobic performance in zebrafish danio rerio. *J. Fish Biol.* **92**, 177–189 (2018).

739     26.    Evans, B. L., Hurlstone, A. F. L., Clayton, P. E., Stevens, A. & Shiels, H. A. Glucose

740            Uptake as an Alternative to Stop-Flow Respirometry for Measuring Metabolic Rate in

741            Danio rerio Larvae. *bioRxiv* 2021.04.30.442098 (2021).

742            doi:10.1101/2021.04.30.442098

743     27.    SigmaAldrich. Assay Optimization and Validation A Technical Guide to PCR

744            Technologies Table of Contents. *MERCK* 1–11 (2015). Available at:

745            https://www.sigmaaldrich.com/technical-documents/articles/biology/assay-

746            optimization-and-validation.html#validating. (Accessed: 1st March 2020)

747     28.    Durinck, S., Spellman, P. T., Birney, E. & Huber, W. Mapping identifiers for the

748            integration of genomic datasets with the R/ Bioconductor package biomaRt. *Nat.*

749            *Protoc.* **4**, 1184–1191 (2009).

750     29.    Fontes, M. & Soneson, C. The projection score - an evaluation criterion for variable

751            subset selection in PCA visualization. *BMC Bioinformatics* **12**, 307 (2011).

30.    Chantzichristos, D. *et al.* Identification of human glucocorticoid response markers

using integrated multi-omic analysis from a randomized crossover trial. *Elife* **10**,

(2021).

31.   Battiston, F. *et al.* Networks beyond pairwise interactions: Structure and dynamics.

*Physics Reports* (2020). doi:10.1016/j.physrep.2020.05.004

32.   Zhang, B. & Horvath, S. A general framework for weighted gene co-expression

network analysis. *Stat. Appl. Genet. Mol. Biol.* **4**, (2005).

33.   Luo, F. *et al.* Modular organization of protein interaction networks. *Bioinformatics* **23**,

207–214 (2007).

34.   Banerji, C. R. S. *et al.* Cellular network entropy as the energy potential in

Waddington's differentiation landscape. *Sci. Rep.* **3**, 1–7 (2013).

35.   Gill, M. S., Tillmann, V., Veldhuis, J. D. & Clayton, P. E. Patterns of GH Output and

Their Synchrony with Short-Term Height Increments Influence Stature and Growth

Performance in Normal Children. *J. Clin. Endocrinol. Metab.* **86**, 5860–5863 (2001).

36.   Zhang, J. D. *et al.* Detect tissue heterogeneity in gene expression data with BioQC.

*BMC Genomics* **18**, 277 (2017).

37.   Subramanian, A. *et al.* Gene set enrichment analysis: A knowledge-based approach

for interpreting genome-wide expression profiles. *Proc. Natl. Acad. Sci. U. S. A.* **102**,

15545–15550 (2005).

38.   Liao, Y., Wang, J., Jaehnig, E. J., Shi, Z. & Zhang, B. WebGestalt 2019: gene set analysis

toolkit with revamped UIs and APIs. *Nucleic Acids Res.* **47**, W199–W205 (2019).

39.   Liberzon, A. *et al.* The Molecular Signatures Database Hallmark Gene Set Collection.

*Cell Syst.* **1**, 417–425 (2015).

40.    M, A. *et al.* Gene ontology: tool for the unification of biology. The Gene Ontology

776           Consortium. *Nat. Genet.* **25**, 25–29 (2000).

777    41.    KL, H. *et al.* Ensembl 2021. *Nucleic Acids Res.* **49**, D884–D891 (2021).

778    42.    S, D. *et al.* BioMart and Bioconductor: a powerful link between biological databases

779           and microarray data analysis. *Bioinformatics* **21**, 3439–3440 (2005).

780    43.    Goodrich, B., Gabry, J., Ali, I. & Brilleman, S. rstanarm: Bayesian applied regression

781           modeling via Stan. (2020).

782    44.    Brilleman, S. L., Crowther, M. J., Moreno-Betancur, M., Buros Novik, J. & Wolfe, R.

783           Joint longitudinal and time-to-event models via Stan. in *StanCon 2018*

784    45.    McElreath, R. Statistical Rethinking : A Bayesian Course with Examples in R and Stan.

785           *Stat. Rethink. A Bayesian Course with Examples R Stan* 1–469 (2018).

786           doi:10.1201/9781315372495

787    46.    Svendsen, M. B. S., Bushnell, P. G. & Steffensen, J. F. AquaResp. (2019).

788           doi:http://doi.org/10.5281/zenodo.2584015

789    47.    Clark, T. D., Sandblom, E. & Jutfelt, F. Aerobic scope measurements of fishes in an era

790           of climate change: Respirometry, relevance and recommendations. *Journal of*

791           *Experimental Biology* **216**, 2771–2782 (2013).

792    48.    Zang, L., Shimada, Y., Nishimura, Y., Tanaka, T. & Nishimura, N. Repeated blood

793           collection for blood tests in adult zebrafish. *J. Vis. Exp.* **2015**, 1–10 (2015).

794    49.    Zang, L., Shimada, Y. & Nishimura, N. Development of a Novel Zebrafish Model for

795           Type 2 Diabetes Mellitus. *Sci. Rep.* **7**, 1–11 (2017).

50.   Eames, S. C., Philipson, L. H., Prince, V. E. & Kinkel, M. D. Blood sugar measurement in

zebrafish reveals dynamics of glucose homeostasis. *Zebrafish* **7**, 205–213 (2010).

51.   Froese, R. Cube law, condition factor and weight-length relationships: History, meta-

analysis and recommendations. *Journal of Applied Ichthyology* **22**, 241–253 (2006).

52.   *Package 'ggpubr' Type Package Title 'ggplot2' Based Publication Ready Plots*. (2020).

53.   Benjamini, Y. & Hochberg, Y. *Controlling the False Discovery Rate: A Practical and*

*Powerful Approach to Multiple Testing. Source: Journal of the Royal Statistical Society.*

*Series B (Methodological)* **57**, (1995).

54.   Motulsky, H. & Brown, R. Detecting outliers when fitting data with nonlinear

regression - a new method based on robust nonlinear regression and the false

discovery rate. *BMC Bioinformatics* **7**, (2006).

55.   Desbuquois, B., Carré, N., Burnol, A.-F. F., Carre, N. & Burnol, A.-F. F. Regulation of

insulin and type 1 insulin-like growth factor signaling and action  by the Grb10/14 and

SH2B1/B2 adaptor proteins. *FEBS J.* **280**, 794–816 (2013).

56.   Johnson, J. Hypernetworks in the Science of Complex Systems. **3**, (2014).

57.   Banerji, C. R. S., Severini, S. & Teschendorff, A. E. Network transfer entropy and

metric space for causality inference. *Phys. Rev. E - Stat. Nonlinear, Soft Matter Phys.*

**87**, 052814 (2013).

58.   Cheng, C. *et al.* Mutation in NPPA causes atrial fibrillation by activating inflammation

and cardiac fibrosis in a knock-in rat model. *FASEB J.* **33**, 8878–8891 (2019).

59. Houweling, A. C., van Borren, M. M., Moorman, A. F. M. & Christoffels, V. M.

Expression and regulation of the atrial natriuretic factor encoding gene Nppa during

development and disease. *Cardiovasc. Res.* **67**, 583–593 (2005).

60. T, H. *et al.* Distinct regulation of developmental and heart disease-induced atrial

natriuretic factor expression by two separate distal sequences. *Circ. Res.* **102**, 849–

859 (2008).

61. Tirumuru, S. S., Arya, P., Latthe, P. & Kirk, J. Understanding precocious puberty in

girls. *Obstet. Gynaecol.* **14**, 121–129 (2012).

62. Godfrey, K. M., Inskip, H. M. & Hanson, M. A. The long-term effects of prenatal

development on growth and metabolism. *Semin. Reprod. Med.* **29**, 257–265 (2011).

63. Devaskar, S. U. & Chu, A. Intrauterine growth restriction: Hungry for an answer.

*Physiology* **31**, 131–146 (2016).

64. Jaquet, D., Gaboriau, A., Czernichow, P. & Levy-Marchal, C. Insulin Resistance Early in

Adulthood in Subjects Born with Intrauterine Growth Retardation [1]. *J. Clin. Endocrinol.*

*Metab.* **85**, 1401–1406 (2000).

65. Lei, X. *et al.* Childhood Health Outcomes in Term, Large-for-Gestational-Age Babies

with Different Postnatal Growth Patterns. *Am. J. Epidemiol.* **187**, 507–514 (2018).

66. Barker, D. J. P., Osmond, C., Kajantie, E. & Eriksson, J. G. Growth and chronic disease:

Findings in the Helsinki Birth Cohort. *Ann. Hum. Biol.* **36**, 444–458 (2009).

67. Savanur, M. S. & Ghugre, P. S. BMI, body fat and waist-to-height ratio of stunted v.

non-stunted Indian children: A case-control study. *Public Health Nutr.* **19**, 1389–1396

(2016).

68.    Painter, R. C. *et al.* Transgenerational effects of prenatal exposure to the Dutch

839            famine on neonatal adiposity and health in later life. *BJOG An Int. J. Obstet. Gynaecol.*

840            **115**, 1243–1249 (2008).

841    69.    Holemans, K., Aerts, L. & Van Assche, F. A. Fetal growth restriction and consequences

842            for the offspring in animal models. *Journal of the Society for Gynecologic Investigation*

843            **10**, 392–399 (2003).

844    70.    Wadley, G. D. *et al.* Growth restriction in the rat alters expression of metabolic genes

845            during postnatal cardiac development in a sex-specific manner. *Physiol. Genomics* **45**,

846            99–105 (2013).

847    71.    Robinson, J. S., Moore, V. M., Owens, J. A. & McMillen, I. C. Origins of fetal growth

848            restriction. *Eur. J. Obstet. Gynecol. Reprod. Biol.* **92**, 13–19 (2000).

849    72.    Finkielstain, G. P., Lui, J. C. & Baron, J. Catch-up growth: Cellular and molecular

850            mechanisms. *World Rev. Nutr. Diet.* **106**, 100–104 (2013).

851    73.    Singhal, A. Long-Term Adverse Effects of Early Growth Acceleration or Catch-Up

852            Growth. *Annals of Nutrition and Metabolism* **70**, 236–240 (2017).

853    74.    S, J., X, Y., H, Z. & W, F. Weight-length relationships and Fulton's condition factors of

854            skipjack tuna (Katsuwonus pelamis) in the western and central Pacific Ocean. *PeerJ* **3**,

855            (2015).

856    75.    Francová, K. & Ondračková, M. Overwinter Body Condition, Mortality and Parasite

857            Infection in Two Size Classes of 0+ Year Juvenile European Bitterling Rhodeus Amarus.

858            *J Fish Biol* **82**, 555–68 (2013).

76.    Shoup, D. & Wahl, D. Body Size, Food, and Temperature Affect Overwinter Survival of

Age-0 Bluegills. *Trans. Am. Fish. Soc.* **140**, 1298–1304 (2011).

77.    Myers, S. S. *et al.* Climate Change and Global Food Systems: Potential Impacts on

Food Security and Undernutrition. *Annual Review of Public Health* **38**, 259–277

(2017).

78.    Plagányi, É. Climate change impacts on fisheries: Assessment of past fisheries

productivity helps to predict and manage future changes. *Science (80-. ).* **363**, 930–

931 (2019).

79.    Free, C. M. *et al.* Impacts of historical warming on marine fisheries production.

*Science (80-. ).* **363**, 979–983 (2019).

**Acknowledgements**

This work was funded by the Biotechnology and Biological Sciences Research Council at the

University of Manchester, Manchester, UK, in combination with an unrestricted Merck

research grant, Darmstadt, Germany. Preliminary and the transcriptomic work was partially

funded by a University of Manchester Research Institute Pump Priming Fund entitled "The

quantitative comparison of the Zebrafish as a model of human development in relation to

paediatric medicine", RMS 104699.

The authors would also like to acknowledge the following people: Dr. Andrew Badrock for

his training, guidance, and donation of reagents, and Jack Broadbent and Joseph Whitehead

for their preliminary work in quantifying growth and metabolism in this model.

**Competing Interests**

The authors declare the research was conducted in the absence of any conflicts of interest.

**Data and Code Availability**

Transcriptomic data is available from the Gene Expression Omnibus (GSE162474). R code is

available online at: https://github.com/terencegarner/GRB10_KD_ZF.

**Figure Legends**

*Figure 1:* **Grb10a** *is successfully knocked down in zebrafish injected with splice-blocking*

*antisense oligonucleotides. 1a*. *Grb10a* qPCR of WT embryos (24-120 hpf, triplicated, n = 5

embryos per well). Data is shown as gene expression relative to β-actin and shows a

significant peak at 48 hpf. *1b.* Schematic of the first five exons of the zebrafish *grb10a* gene.

5' splice sites are highlighted with the forward and reverse primer triad indicated. *1c.*

Multiplexed PCR amplification of the e3i3 and e4i4 splice site in embryos treated with either

Standard Control morpholino, e3i3, or e4i4. β-actin was used as a positive control. *1d.*

Western blot results of phosphorylated vs total protein ratios for two major signalling

molecules of the insulin signalling pathway: AKT and S6. Quantitation using densitometry

depicts mean + SEM. Activation of both proteins was found to be significantly elevated in KD

zebrafish compared to SC (n = 3, unpaired t-test *** p = 0.0007, * p = 0.0413).

***Figure 2: Growth and cardiometabolic phenotype are significantly impacted by* grb10a**

***perturbation. 2a.*** Mean total body length ± SEM of SC and KD zebrafish from 24 to 120 hpf.

T-test significance *** = 0.0006 **** < 0.0001 *** = 0.0001. ***2b.*** Mean total body length ±

SEM and individual data points of 96 hpf zebrafish embryos (n = 25). *Grb10a* KD phenotype

was reversed in *grb10a* overexpression zebrafish. Coinjection resulted in phenotype rescue.

One-way ANOVA revealed KD zebrafish were significantly longer, while *grb10a*

overexpression zebrafish were significantly smaller than SC (*** = 0.0001). Rescue zebrafish

were of similar length to SC (ns = 0.3792). ***2c.*** Mean body length measurements relative to

SC ± SEM. E3i3 and e4i4 both exhibit a propensity to elevate body length, while *grb10a* RNA

shows a dose-dependent ability to inhibit body length. ***2d.*** Mean heart rate ± SEM in beats

per minute of SC and KD zebrafish. Heart rate was significantly lower in KD embryos

compared with SC after 48 hpf (*** = 0.0006, **** < 0.0001). ***2e.*** Mean yolk area ± SEM of

SC and KD embryos over the embryonic life stage. Following the initial 24 hours, KD zebrafish

had significantly smaller yolks compared to SC, indicating yolk content was metabolised at a

much higher rate (** = 0.0084 **** < 0.0001). ***2f.*** Glucose Uptake-Glo$^{TM}$ Assay of 96 hpf KD

and SC zebrafish, where higher luminescence indicates a greater accumulation of

intracellular 2D6P. Luminescence was approximately 30% greater in KD zebrafish compared

to SC (*** = 0.0002). All SC vs KD comparisons by unpaired t-test.

***Figure 3. Transcriptomic analysis of Standard Control and grb10a Knockdown gene***

***expression over the first 30 dpf.*** Hierarchically clustered heat maps of gene expression

generated from an Affymetrix GeneChip™ Zebrafish Genome Array of SC (***3a***) and KD (***3b*** and

***3c***) zebrafish RNA, taken at 5, 10, 15, 20, and 30 dpf. Expression segregates into three

clusters in SC zebrafish (***3a***). Clustering of the same genes identified in ***3a*** are disrupted in

the KD dataset (***3b***). Analysing the KD dataset independently shows age-related hierarchical

gene expression in the falls into two clusters (***3c***). ***3d***. Venn diagram of age associated genes.

***3e***. Gene set enrichment analysis, using the GO Biological Process Ontology gene list, of the

age-related genes in the SC and KD datasets. The top 20 most enriched pathways with

differential expression are included here, with associated normalised enrichment scores and

q-values.

***Figure 4: Coordination within the transcriptome is altered by grb10a KD, as assessed by***

***hypernetwork analysis 4a.*** Hypernetwork analysis defined three clusters of highly

connected genes in Standard Control zebrafish, corresponding to 5 dpf, 10-15 dpf, and 20-

30 dpf respectively. ***4b***. Two clusters were defined in the KD data, corresponding to 5-15 dpf

and 20-30 dpf. ***4c*** *and* ***4d***. Violin plots of connectivity (left) and entropy (right) in the 20-30

dpf cluster. The KD transcriptome was more connected with greater entropy compared to

Standard Controls, suggesting more diverse functionality.

***Figure 5: Analysis of the set of genes in the wider transcriptome shows a 27.8-fold increase***

***in the KD ZF. 5a-5b***. Venn diagrams of genes positively (***a***) and negatively (***b***) correlating

with age. ***5c-5d***. Hierarchically clustered heat maps of gene expression of the genes

identified in the wider transcriptome. Gene expression in the standard control (***5c***) cluster

into two age related groups, whereas expression in the knockdown (***5d***) show significant

dysregulation. **5e-5f** Gene set enrichment analysis (GSEA) ranked by R-value of rank age

regression in the standard control (**5e**) and knockdown (**5f**). **5g** GSEA ranked by R-value of

rank age regression of the cluster of genes in the white box in **5d**.

**_Figure 6: Hypernetwork modelling of ontology highlights pathways with functional_**

**_difference between SC and KD._** Hypernetworks were iteratively generated from genes

attributed to each pathway and entropy was modelled across the two treatment groups. β

values represent the difference in entropy between SC and KD, with a β value of 0 indicating

no difference between groups. Ontology classes are considered to be significantly different

if the distribution of β values (89% CI) does not include 0. _Actin filament based movement_

and _negative regulation of extrinsic apoptotic signalling pathway_ were the only two

pathways with no functional difference identified between the groups.

**_Figure 7: Length and cardiac function are permanently altered. 7a._** Mean total body length

of SC and KD zebrafish up to 30 dpf. Following the embryonic growth spurt, KD zebrafish

experienced an additional period of rapid growth between 15 and 20 dpf (**** < 0.0001).

**7b.** Individual total body length (left), mass (middle), and condition factor scores (right) for

18-month KD and SC zebrafish (n = 21-24). Length and mass were significantly higher in the

KD (**** $p < 0.0001$, ** $p = 0.005$), while condition factor was significantly lower (* $p =$

$0.0215$), indicating KD zebrafish have leaner bodies. **7c.** Thickness of 10 individual skeletal

muscle fibres stained with Masson's Trichrome from 5 individuals (left) and means grouped

by individual (right) taken from a position posterior to the dorsal fin. Fibres were consistently

thicker in the KD zebrafish (**** $p < 0.0001$). All comparisons by unpaired t-test. Data are

presented as mean or individual values ± SEM.

*Figure 8: Changes established during embryonic development perpetuate into variations in*

*adult function. 8a.* qPCR results of three genes associated with cardiac performance in adult

cardiac tissue, relative to β-actin. *Myl7* expression was significantly elevated in KD zebrafish

(**** $p < 0.0001$) while *nppa* expression was significantly down regulated (*** $p = 0.0012$)

compared to SC zebrafish. There was no significant difference between the expression of

*pcna* ($p = 0.3041$). 8b. Ventricular morphometrics obtained by Masson's Trichrome histology

comparing KD and SC compacta thickness and tissue density. The compacta layer was

significantly thicker (* $p = 0.0288$) and overall density was higher (* $p = 0.0289$) in the KD

zebrafish, suggesting greater cardiac efficiency (n = 5-6). *8c*. Maximum (MMR) and Basal

(BMR) oxygen consumption of adult (18 month) zebrafish, adjusted for body mass. BMR was

comparable between the two groups, while MMR was greater in the KD zebrafish (**** $p <$

$0.0001$), resulting in a greater aerobic scope (dotted line, *** $p = 0.0007$). *8d.* Fasting blood

glucose concentrations of adult zebrafish from each treatment group (n = 14-21). KD

zebrafish had significantly higher circulating blood glucose levels (* $p = 0.0463$). *8d.* Glucose

tolerance (top) (n = 10) and insulin sensitivity (bottom) (n = 8–12) trials in adult (18 month)

KD and SC zebrafish. The treatment groups were equally able to tolerate glucose challenge

and responded similarly to insulin. Treatment started immediately after the first

measurement. Data are presented as either mean or individual values ± SEM, and all

comparisons are by unpaired t-test.

**Tables**

**Table 1. Oligonucleotide sequences**

| A) Morpholino-modified antisense oligonucleotides | | | |
|---|---|---|---|
| **Name** | **Accession no.** | **Sequence** | |
| e3i3 | | CATACAGTATGCATTACCTGACAGC | |

| e4i4 | BX571825 | GTGTCTGTTTTCAGCTCTTACATGT |
| Standard Control | | CCTCTTACCTCAGTTACAATTTATA |

**B) Primers for cloning *grb10a* cDNA and for validating splicing interruption**

| Name | Accession no. | Forward | Reverse |
|---|---|---|---|
| *grb10a* cDNA | NM_0010042 | gacagaatcgatGACTGAGTATGGC | gacagatctagaTCATAAGGCC |
| *grb10a* exon3- | BX571825 | TGGTGAATGACATGGCCTCT | CTGCATGGTCAAGACACAC |
| *grb10a* exon3- | | TGGTGAATGACATGGCCTCT | CTGACGGATAATAGCTACA |

**B) Primers for qPCR**

| Gene | Accession no. | Forward | Reverse |
|---|---|---|---|
| *grb10a* | NM_0010042 | TGGATGACTGCTTTTAGACTGC | CGACCAGTCCTTCCAGAAA |
| *actb1* | NM_131031. | CTTCCAGCAGATGTGGATCA | GCCATTTAAGGTGGCAACA |
| *myl7* | NM_131329. | TCACTGTCTTCCTCACCCTC | CACGTCTATTGGAGCCACT |
| *pcna* | NM_131404. | AGGCAACATCAAGCTCTCAC | ATTTGACGTGTCCCATGTCT |
| *nppa* | NM_198800. | AAGCAAAAGCTTGTCTGG | ACTGTATCCGCGTATTGCA |

981    **Table 2. Microinjection Solutions**

| Injection | Oligonucleotide | nCerulean | Phenol Red |
|---|---|---|---|
| Standard Control (5 | 0.5 mM human beta globin | 50 ng/µl | 1 µl |
| *grb10a* KD (5 µl) | 0.5 mM e3i3 or e4i4 morpholino | 50 ng/µl | 1 µl |
| *grb10a* OE (5 µl) | 500 ng/µl *grb10a* RNA | 50 ng/µl | 1 µl |

982    **Table 3. Thermocycling Parameters**

| Procedure | Temperature (℃) | Time (S) | Cycles |
|---|---|---|---|
| NEB Q5 Hot Start | 95 | 300 | |
| | 95 | 30 | 35 |
| | 55 | 30 | |
| | 72 | 120 | |
| | 72 | 600 | |
| NEB Taq | 95 | 30 | |
| | 95 | 15 | 30 |
| | 55 | 15 | |
| | 68 | 30 | |
| | 68 | 300 | |
| SyBr Green qPCR | 95 | 30 | |
| | 95 | 5 | 40 |
| | 51 | 15 | |
| | 72 | 10 | |

983    **Table 4. Western Blot Antibodies**

| Target | Species | Supplier | Dilution |
|---|---|---|---|
| AKT | Mouse | Cell Signalling #2920 | 1:1000 |

| p-AKT | Rabbit | Cell Signalling - #5831 | 1:1000 |
|---|---|---|---|
| S6 | Mouse | Cell Signalling - #2317 | 1:1000 |
| p-S6 | Rabbit | Cell Signalling - #2215 | 1:1000 |
| Anti-Mouse HRP linked IgG | Horse | Cell Signalling - #7076 | 1:5000 |
| Anti-Rabbit HRP linked IgG | Goat | Cell Signalling - #7074 | 1:5000 |

**Supplementary Figures**

***Supplementary Figure 1: Pipeline of transcriptomic analysis and hypernetworks. 1a.***

Analysis pipeline of transcriptomic data. **(i)** Unsupervised hierarchical clustering was

performed to identify age-associated gene clusters. Genes were filtered by variance, using a

projection score to maximise the informativeness of the genes selected. Clusters of age-

associated genes were identified for SC and KD animals. **(ii)** Hypernetworks were generated

using age associated genes for each group. Hypernetwork structure was quantified using

connectivity and entropy. Clusters of highly connected genes were identified, and a wider

set of transcripts were implicated as important by identifying the complete subgraph

between cluster nodes and edges in the hypernetwork incidence matrix. **(iii)** GSEA was used

to investigate biological functions associated with genes clustered by the hypernetwork or

implicated by the complete subgraph in the hypernetwork incidence matrix. **(iv)** Biological

processes identified by GSEA were assessed for functional activity. Hypernetworks were

iterated, using subsets of genes associated with each process, and calculating hypernetwork

entropy. A Bayesian modelling approach was used to detect differences in entropy

distributions between processes. **1b.** A general model of a hypernetwork, shown as a three-

dimensional representation of genes (coloured tetrahedra) correlating with the expression

of other genes (black spheres). Shared correlations are represented by matching vertices,

edges, and faces of the tetrahedra. The dimensionality of the connection between genes is

defined by the number of shared correlations between those genes. *1c.* A hypernetwork

representation of the "higher order" interactions within the transcriptome. This summary of

the genes with shared correlations can be considered as the incidence matrix of a multi-

dimensional network.

***Supplementary Figure 2. Connectivity (A) and entropy (B) of 20-30 dpf associated genes in***

***the SC and KD zebrafish compared with randomly selected gene sets. 2A.*** Connectivity in

the experimental data was significantly greater than in the random iterative data, and

greater in the KD data than SC ($p < 0.0001$). ***2B.*** Entropy was significantly lower in the

experimental data than the random iterative data. The reduction in entropy between the

experimental data and interated data was greater in the SC than the KD.

**Supplementary Tables**

***Supplementary Table 1.*** Gene set enrichment analysis (GSEA) on age related gene

expression in morpahlino and control animals. 3733 orthologous human genes from 75212

probsets. Gene probe sets collapsed to gene summary by average. Group ANOVA by age

groups. q value is the false discovery rate (fdr) modified p-value of the GSEA, ES = edge

score, NES= normalised edge score, abs(ES)= absolute edge score. GSEA performed in

Qlucore Omics Explorer (v3.6) using Gene Ontology Biological Process. Significance level of

the fdr coloured from blue (highly significant) to red (not significant).

***Supplementary Table 2.*** Gene sets from the wider connected transcriptome of the 20-30dpf

Control and Morpholino Zebrafish (460 and 12775 respectively). Age rank regression

analysis of gene expression presented with R-statistic and p-value. Zebrafish gene symbol

and corresponding human orthologue shown.

*Supplementary Table 3.* A Gene probe sets (3460) from the wider connected transcriptome

of 20-30dpf Morpholino Zebrafish dysregulated in controls. Age rank regression analysis of

gene expression presented with p-value and q-value (false discovery rate modified p-value).

ZF gene symbol and corresponding human orthologue shown.

*Supplementary Table 4.* Dysregulated expression of cardiac phenotype marker genes in the

KD zebrafish. False discovery rate modified p-value (q-value) shown for age group ANOVA.

*Figure 1:* **Grb10a** *is successfully knocked down in zebrafish injected with splice-blocking antisense oligonucleotides*

*Figure 2: Growth and cardiometabolic phenotype are significantly impacted by grb10a perturbation.*

Age-associated genes SC data

Age-associated genes KD data

| Name | KD NES | KD q | SC NES | SC q |
|---|---|---|---|---|
| ACTIN FILAMENT BASED MOVEMENT | 2.088 | 0.012 | 2.176 | 1.000 |
| PLATELET ACTIVATION | 2.068 | 0.013 | 2.236 | 1.000 |
| EXTRACELLULAR STRUCTURE ORGANIZATION | 2.011 | 0.013 | 2.246 | 1.000 |
| COLLAGEN FIBRIL ORGANIZATION | 2.026 | 0.014 | 2.257 | 1.000 |
| REGULATION OF RNA SPLICING | 1.608 | 0.039 | 0.726 | 0.855 |
| GLUTATHIONE METABOLIC PROCESS | 1.660 | 0.031 | 0.735 | 0.848 |
| RIBOSOME ASSEMBLY | 1.700 | 0.026 | 0.743 | 0.840 |
| REGULATION OF MRNA PROCESSING | 1.609 | 0.039 | 0.810 | 0.772 |
| PLATELET DEGRANULATION | 1.753 | 0.022 | 2.155 | 0.757 |
| METANEPHROS DEVELOPMENT | 1.709 | 0.026 | 0.841 | 0.733 |
| DICARBOXYLIC ACID TRANSPORT | 0.690 | 0.895 | 1.693 | 0.048 |
| POSITIVE REGULATION OF WOUND HEALING | 0.876 | 0.703 | 1.672 | 0.048 |
| FAT SOLUBLE VITAMIN METABOLIC PROCESS | 0.950 | 0.597 | 1.655 | 0.049 |
| POSITIVE REGULATION OF SMOOTH MUSCLE CELL PROLIFERATION | 0.986 | 0.543 | 1.708 | 0.048 |
| REGULATION OF STEROL TRANSPORT | 1.012 | 0.503 | 1.704 | 0.049 |
| POSITIVE REGULATION OF PROTEIN BINDING | 1.033 | 0.471 | 1.663 | 0.049 |
| AMINO ACID TRANSPORT | 1.037 | 0.466 | 1.703 | 0.048 |
| POSITIVE REGULATION OF LIPID LOCALIZATION | 1.059 | 0.436 | 1.757 | 0.050 |
| CELLULAR MODIFIED AMINO ACID BIOSYNTHETIC PROCESS | 1.068 | 0.423 | 1.664 | 0.049 |
| NEGATIVE REGULATION OF EXTRINSIC APOPTOTIC SIGNALING PATHWAY | 1.097 | 0.383 | 1.674 | 0.049 |

*Figure 3. Transcriptomic analysis of Standard Control and grb10a Knockdown gene expression over the first 30 dpf.*

*Figure 4: Coordination within the transcriptome is altered by grb10a KD, as assessed by hypernetwork analysis.*

*Figure 5: Analysis of the set of genes in the wider transcriptome shows a 27.8-fold increase in the KD ZF.*

*Figure 6: Hypernetwork modelling of ontology highlights pathways with functional difference between SC and KD.*

***Figure 7: Length and cardiac function are permanently altered.***

**Figure 8: Changes established during embryonic development perpetuate into variations in adult function.**

**A.** **Analysis Pipeline**

(i)

**B.**

*Supplementary Figure 1: Pipeline of transcriptomic analysis and hypernetworks.*

*Supplementary Figure 2. Connectivity (A) and entropy (B) of 20-30 dpf associated genes in the SC and KD zebrafish compared with randomly selected gene sets.*