The defensive behavior of Indiana mite biting honeybees against *Varroa destructor* and the
structure of mandibles

Jada Smith[1], Xaryn L. Cleare[1], Krispn Given[2], Hongmei Li-Byarlay[1, 3] *

1. Department of Agricultural and Life Sciences, Central State University, Wilberforce, OH
45384, USA;
2. Department of Entomology, Purdue University, West Lafayette, IN 47906, USA;
3. Agricultural Research and Development Program, Central State University, Wilberforce,
OH 45384, USA;

* Corresponding author: Hongmei Li-Byarlay, hli-byarlay@centralstate.edu

Keywords: honeybee, *Apis mellifera*, defensive behavior, Varroa resistance, bite, grooming,
mandible, microCT

Abstract:

The honeybees (*Apis mellifera*) are the most important managed pollinator for sustainable
agriculture and our ecosystem. However, the managed honeybee colonies in the United States
experience 30-40% of losses annually. Among all the biotic stressors, the parasitic mite *Varroa*
destructor is considered as one of the main pests for colony losses. The mite biting behavior as a
Varroa tolerant or resistant trait has been selected in the State of Indiana for over a decade (13
years). A survey of damaged mites from the bottom of a colony can be used as an extended
phenotype of the mite biting behavior to evaluate a colony. On average 37% of mites sampled
were damaged or mutilated from the breeding stocks of 59 colonies of mite biters of Indiana,
which is significantly higher than 19% of damaged mites in commercial colonies from the
Southern United States. No significant difference was detected between the damage of missing
legs between breeding stocks and commercial colonies. In addition, the morphology of pollen
forager worker mandibles was compared between two populations via X-ray micro-computed
tomography using six parameters, and a difference was detected in one parameter. Our results
showed the novel scientific evidence to explain the potential defensive mechanism against
Varroa mites via potential structural changes in worker mandibles.

Introduction

The infestation of ectoparasitic mites *Varroa destructor* is the primary concern for honeybee (*Apis mellifera*) colony losses since these mites were introduced to the United States since 1987 (Guzmán-Novoa et al., 2010; Le Conte et al., 2010; Nazzi and Le Conte, 2016). Mature *Varroa* females are 1.1x1.5 mm, and the males are 0.8 x 0.7 mm in sizes. The majority of a mite's life cycle happens in the brood cells including developing stages of egg, six-legged larva, protonymph, deutonymph, and adult (Bailey, 1968; Genersch, 2010). *Varroa* mites can weaken, deform, and decrease the lifespan of the infested honeybee colony because mites feed on the honey bee pupae and adults, transmit viral pathogens, cause morphological and behavioral defects on bee development and behavior (Le Conte et al., 2010). Recent scientific evidence showed that *Varroa* mite feed on the fat body of adult bees (Ramsey et al., 2019). Furthermore, *Varroa* mite is an effiective vector to transmit honey bee viruses within the honeybee colony (Di Prisco et al., 2011; Wilfert et al., 2016).

The European Honeybee (*A. mellifera*) has developed defensive behavior to keep the mite population low by grooming, biting, and hygienic behaviors against *Varroa* mites (Arechavaleta-Velasco and Guzmán-Novoa, 2001; Guzman-Novoa et al., 2012; Ruttner and Hanel, 1992; Spivak, 1996; Tsuruda et al., 2012; Villa et al., 2017). The biting behavior, also considered a type of grooming behavior, enables worker bees to bite and remove adult mites from their bodies (Peng et al., 1987; Pritchard, 2016; Ruttner and Hanel, 1992). Colonies selected by instrumental insemination or open mating with feral colonies on mite biting behavior potentially have greater fitness over subsequent generations. Field reports showed that Indiana mite biters from Purdue University are being selected in the past decade, and dramatic increases in the average proportion of damaged mites in their breeding population (Andino and Hunt, 2011; Hunt et al., 2016; Morfin et al., 2019). However, to date, there is no report to compare the Indiana mite biter honeybee stocks to the commercial population in the United States.

The breeding of mite resistant bees is critical to maintaining sustainable apiculture for local pollination, food, and crop productivity (Oddie et al., 2017). Modern beekeeping and breeding techniques of instrumental insemination enable artificial selection effects on honeybee

queens and colonies (Meixner et al., 2010). Breeding efforts have been made in different

continents and countries to select mite tolerance or resistant traits (Büchler et al., 2010; Rinderer

et al., 2010; Spivak, 1996; Villa et al., 2017). Mite resistant bees will assist beekeepers in

managing the growing chemical miticide-resistance problems and play a critical role in

promoting sustainable agriculture practices (Hamiduzzaman et al., 2017; Kanga et al., 2016). In

the past, research efforts for breeding are focused on different bee breeding stocks to improve

colony health, including Russian bees, *Varroa* Sensitive Hygienic bees, Minnesota Hygienic

bees (Buechler et al., 2010; Danka et al., 2011; Guarna et al., 2017; Guarna et al., 2015; Rinderer

et al., 2010; Spivak, 1996; Villa et al., 2017).

Honeybee mandibles are considered to be the main mouthpart that worker bees use to bite

or chew the parasites, including mites and wax moths in the colony (Ruttner and Hanel, 1992).

Chemical analysis revealed 2-Heptanone was secreted from the mandibles, acting as an

anaesthetic on wax moth larva and *Varroa* mites (Papachristoforou et al., 2012). Micro-X-ray-

computed tomography (MicroCT) is a technology with a fast 3-D scanning in satisfactory spatial

resolution without complicated and lengthy sample preparation procedures. In the past, this

technique has been used to study the brain anatomy and evolution of bees, ants, and other insects

(Coty et al., 2014; Larabee et al., 2017; Li et al., 2011; Ribi et al., 2008). However, there is no

report to use MicroCT to demonstrate the shape of honeybee mandible in fine details.

In this study, we hypothesized that the Indiana mite biter breeding stocks have a higher

level of mite biting behavior than commercial bee colonies. Our goal is to characterize the

behavioral and morphological capacity for the mite biting behavior. Total number of mites, the

percentage of damaged mites as the parameter for mite biting behavior, and the number of mite

legs missing per colony were reported. In addition, we investigate the mechanism of mite biting

behavior by examining the 3-dimensional shapes of mandibles between Indiana mite biter

colonies and commercial colonies from the Southern United States (mainly the state of Georgia).

Material and Methods

Honeybee colonies

Fifteen colonies from five commercial source were sampled from beekeepers who bought their package colonies in 2018 originally from the State of Georgia (from five different commercial providers). Mites were collected in September of 2018 from different areas in the State of Ohio (Figure 1a – 1 colony from Defiance (Defianace1), OH; and b – 14 colonies from Bellbrook (NA1, NA2, NA3, NA5, NA6), Beavercreek (PBJohn1, PBJohn2), Cedarville (Dan4), and Wilberforce (AB1, CSU23, CSU24, CSU32, CSU51, CSU52) OH). Total of 59 colonies from Indiana mite biter honeybees were sampled at Lafayette, Indiana (Figure 1c, colony numbers are in the supplemental table 1). Mite samples were collected over a 5-day period as fresh mites from each colony at Purdue University's main apiary in the fall of 2018. Some colonies sampled twice during the fall season. Seven colonies of Wright Patterson Air Force Base (WPAFB) Huffman prairie site housed seven virgin queens from Purdue stock colonies and then open-mated with drones from feral colonies near WPAFB (within two miles).

For all of the worker bee samples of the mandible scanning, pollen foragers returning to the hive (either commercial or Indiana mite biter colony) entrance were collected using an insect vacuum (Cat: 2820GA, BioQuip). Then all bees were frozen on dry ice, transported back to the lab and kept in -80 freezer. At least 10 foragers were collected per colony per site. Indiana mite biter bee samples were collected in June 2018. Commercial bee samples were collected in July 2018.

Varroa Mites

Three groups of mites were compared: 1) commercial (comm), 2) mite biters (from Indiana), and 3) mite biter open mated (at WPAFB). Figure 2 showed one example of a worker bee biting a *Varroa* mite on the top of a hive. Mite samples of Indiana mite biters were collected following the previously described method (Andino and Hunt, 2011). For commercial bees and mite biter open mated in Ohio, mite samples were collected as previously reported (Andino and Hunt, 2011). A small paint brush was used to remove mites from the bottom boards to a plastic disposable cup with a lid (volume 1oz). Mite samples stayed in a -20 freezer overnight. Each mite was carefully glued onto a glass microscope slide (25 x 75 x 1mm, Globe Scientific Inc.) by a small paint brush. Slides were examined under a light microscope (Zeiss STEMI 580) with a magnify degree of 50x. Colonies with 15 or more mites sampled within a 5-day period were included in the data analysis. Mites collected in Ohio were examined for any missing legs (range

from 1 to 8) as visible damage in the view (Figure 3). The total number of mites sampled per colony and the number of damaged mites were compared among all three groups (comm, IN mite biters, and mite biter open mated). The number of missing legs were compared between commcial colonies and mite biter open mated colonies. Immature mites and empty shells of mite body were excluded.

MicroCT Scanning

From commercial colonies, five bees (five pairs of mandibles) were scanned from two colonies (three bees from colony PBJohn2 and two bees from colony Dan4). For the Indiana mite biters, nine bees from three colonies (three bees from colony 5, four bees from colony 15, and two bees from colony 41) were scanned. The scanning process was performed at the Center for Electron Microscopy and Analyses (CEMAS) of Ohio State University (Columbus, OH) using a HeliScan (FEI, Thermal Fisher, USA) for 3-dimensional imaging. A pair of mandibles was fixed to thin wooden posts (r = 1 mm and h = 148mm) using super glue to fit in the HeliScan. The scanning parameters were as follows: isotropic voxel size = 2.564 m per pixel; voltage =60 or 80 KV, current = 80 or 46 micro A; helical scan using space filling trajectory reconstruction (Kingston et al., 2018) without any filter, and 1440 X-ray raw images. The software AVIZO for FEI systems (version 9.4) was used to quantify the measurements (Height, Length, Width, Small Edge, Long Edge, and the Span of spike area) for each scanned sample.

Data Analysis

The average number of mite samples per colony was compared among three groups, commercial (comm) bees, Indiana mite biters, and the mite biters open mated at WPAFB. If a colony was sampled twice in the fall season, the average number of mites collected was used for statistical analysis. The percentage of damaged mites over total sampled mites was transformed using positive square root for normal distribution. One-way ANOVA (ANalysis Of VAriance) with post-hoc Tukey HSD (Honestly Significant Difference) Test calculated for comparing multiple treatments test was used for the difference among means of different populations as to the total mite numbers, mite biting rate, damage of legs, and parameters of mandibles. The online tool Interactive Dotplot (Weissgerber et al., 2017) was used to generate all boxplots.

Results

The behavior of mite biting or grooming refered to here describes how a worker bee uses

its two forlegs and two mandibles of its mouthpart to attack a *Varroa* mite in a honeybee colony.

To characterize the mite damage, we categorized the mites observed as four different types: (A)

mature adult female mites in dark brown colors with no damage and all eight legs (Left 1, 2, 3, 4,

Right 1, 2, 3, 4) are present, (B) damaged adult female mites with legs (in this case, four legs

Left 1, 2, 3, 4) missing, (C) young adult female mites which are pale in color and are not counted

as a damaged adult mite, (D) mites with partial body missing (Figure 2). Mite samples similar to

C type are excluded because no damage was detected by worker bees' grooming or biting

behavior.

To evaluate the total mite population, we collected all the *Varroa* mites that appeared on

the bottom board of each colony from each commercial colony (15 colonies, N_{mite} = 886), mite

biter colony (59 colonies, N_{mite} = 3,390), and mite biting open mated colony (7 colonies, N_{mite} =

569). There was no significant difference or significant interaction detected among the

commercial colonies, mite biter colonies, and mite biter open mated colonies (one-way ANOVA

of three independent treatments, $F(2, 80)$ = 1.1992, p = 0.3069, Figure 3).

To assess the mite biting behavior or grooming behavior, we surveyed all the damaged

mites collected from the bottom board of each colony from all three groups (commercial

colonies, N_{mite} = 172, mite biter colonies, N_{mite} =1,201, mite biter open mated colonies N_{mite} =

199). The mean of the percentage of damaged mites per colony among three treatments

(commercial, mite biter, and mite biter open mated) after transformation are 39.80%, 60.22%,

and 60.14%. One-way ANOVA analysis suggests that one or more groups are significantly

different ($F(2, 80)$ = 31.3785, p = 1.0031e-10). The Tukey HSD test identified that the levels of

mite biting behavior or grooming behavior in the mite biter are significantly higher than the

commercial colonies (Q = 11.0866, p < 0.01, Figure 4). Levels of mite biting behavior or

grooming behavior in the mite biter open mated colonies are significantly higher than the

commercial colonies (Q = 6.9771, p < 0.01, Figure 4). Levels of mite biting behavior or

grooming behavior between mite biter and mite biter open mated colonies are not significantly

different (Q = 0.0304, p > 0.05, Figure 4). Commercial bees have the lowest mite biting behavior

among all three groups.

To evaluate the damage of mites and potential difference between commercial colonies
and mite biter open mated colonies, we counted the number of legs missing from each damaged
mite. Our result showed no difference regarding the average number of legs missing per mite
between these two groups ($Q = 2.4278$, p > 0.05, Figure 5).

To compare the morphology of mandibles, six different parameters (Figure 6) including
the length (L), the width (W), the height (H), long edge (LE), short edge (SE), and span of the
spine area (S) are measured from the microCT data between commercial colonies and mite biter
colonies. Statistical analysis (ANOVA) between these two groups showed that the long ege (LE)
of mite biter colonies are significantly short than the commercial colonies (F = 5.7762, p =
0.0256, Figure 7). There is no significant difference of the other five parameters between two
groups (Figure 7), but in the length, height, short edge and span of the spine area, we noticed a
consistent trend of a smaller value in mite biter colonies.

Discussion

We have investigated the *Varroa* mite population and the mite biting behavior among
mite biter colonies, mite biter open mated colonies, and commercial colonies in the U.S. Also,
we evaluated the difference in the shape of bee mandibles between mite biter honeybee stock and
commercial colonies. The mite biter colonies displayed a higher level of mite biting behavior
than the commercial colonies and the difference in the long edge of their mandibles may provide
a potential explaination of a physical mechanism to mutilated mites using mandibles.

Previous research on grooming behavior and damaged mites in *Apis* showed that
grooming behavior is a selected trait in naturally mite-resistant colonies (Arechavaleta-Velasco
and Guzmán-Novoa, 2001; Boecking and Ritter, 1993; Fries et al., 1996; Peng et al., 1987;
Russo et al., 2020). As one of the defensive behaviors of honey bees for parasitic *Vaoora* mites,
the mite biting or grooming behavior can be used as a parameter to selection honey bee stocks
for *Varroa* mite resistance (Hunt et al., 2016; Morfin et al., 2019; Pritchard, 2016; Rinderer et
al., 2010; Spivak, 1996).

Our present comparisons between commercial and mite biter bees support the value of
selection of mite biting behavior in *A. mellifera*. With our evidence of freshly damaged mites, we

provided a strong argument that workers of *A. mellifera* are able to amputate the legs of *Varroa*

mites as described before (Ruttner and Hanel, 1992). Damaged mites collected from the bottom

board of each colony within a five-day time ensured the observed damage on mites was fresh. It

is possible for beekeepers to record the proportion of damaged mites from the bottom board of

colonies and create a collaborative regional effort for a selection in a region (Bienefeld et al.,

1999; Hunt et al., 2016).

As a defensive response, honeybees use biting or grooming behavior to decrease the

infestation of a mite population in the colony. It is known that worker bees can use their

mandibles to multilate or damage *Varroa* mites from reports published before (Ruttner and

Hanel, 1992). Further studies showed that certain chemical such as 2-Heptanone can be released

from worker's mandibles during a bite to anaestheze parasites including *Varroa* mites and wax

moths in honeybee colonies (Papachristoforou et al., 2012). Although there are reports on the

mite biting behavior, there is no report on the underlying mechanism for this behavior. Our

results provided empirial evidence on the changes of the structure of mandibles such as long

edge which can be an explanation for the mechanism of biting behavior. In addition, our data

showed that mite biters are under selelction which may lead to the structure changes for mite

resistance. Eastern honeybees (*A. cerana*) are original hosts of *Varroa* mites, and have evolved

to be *Varroa* mite resistant now in Asia. Their body size are slightly smaller than *A. mellifera*

(Peng et al., 1987; Yue et al., 2018).

Our data indicate the clear trend that mite biters are evolving a behavioral adaptation to

the parasitic mite *V. desctructor*. The sensitivity of workers to the mites in colonies selected for

mite-biting behavior is higher, and they perform higher level of mite biting behavior compared to

commercial colonies given similar numbers of mite population in all colonies. However, this

study cannot compare these stocks ability to reduce mite populations because they were in

different geographic locations and had different management histories.

The worker bee's ability to detect mites may be based on the olfactory mechanisms

considering the mite biting behavior happens in the dark hive most time. The stocks of mite

biters may show a better capability to detect and recognize mites as pest than non-selected

commercial colonies. There may be gradual changes in the relationship between *A. mellifera* and

Varroa mites. There is a potential change in the biting mechanism of mite biters, which Asian

honeybees displayed in previous research(Peng et al., 1987). Compared to the low frequency of

removing mites or low success in clearning the mites from commercial-source colonies, we

observed an improvement of these abilities in mite biter colonies.

Even though there is evidence for variation in the mite biting or grooming behavior in

different genotypes (Guzman-Novoa et al., 2012), more knowledge is needed on the genetic

architecture and the pattern of inheritance of this behavior for honeybee breeding and selections.

With rapid development of new sequencing technologies and genome editing tools, genome-

wide marker-assistant selelction may be applied in the future for honeybee breeding.

Acknowledgments

We thank Greg Hunt who guided sample collections at Purdue University and provided

comments for the manuscript, Dwight Wells and Danielle Trevino for assistance on sample

collections at the WPAFB, Nathan Arndt, Andy Bern, John Gazzerro, Danielle Kroh, and Jamie

Walters for the access to commercial colonies, Carley Goodwin at Ohio State University for the

assistance on the microCT process, and Kaila Young at Central State University for assistant to

mount and scan samples.

Author contributions

H.L.-B. conceived and designed experiments; J.S., X.L.C., K.G. & H.L.-B. collected samples &

data. J.S. and H.L.-B. performed data analysis and wrote the paper. All authors approved the

final manuscript.

Funding information

Financial support was granted to H.L.-B. by the USDA NIFA Evans-Allen fund

(NI171445XXXXG004), USDA SARE North Central Region Grant (ONC19-062), and USDA

AFRI Seed Grant (2020-67014-31557) and Levin Foundation. J.S. was supported by the NSF

IPSRG scholarship and the USDA NIFA Evans-Allen fund.

 Figure and Table legend

Figure 1. The map of locations for sample collection from the State of Ohio and Indiana. (a site)
Defiance, OH; (b site) Green county including cities of Bellbrook, Beavercreek, Cedarville, and
Wilberforce, OH. (c site) West Lafayette, IN for Indiana mite biter colonies.

Figure 2. Images of damaged *Varroa* mites from our sampled colonies. A: matured mite with no
damage, Labeled numbers of 1-8 indicate eight legs of the *Varroa* mite; B: a damaged mite with
legs missing; C: young mature mites; D: a matured mite with a missing body part. Arrows
indicated the damaged legs or the body part.

Figure 3. The box plot of the average number of mites collected per colony among three groups,
commercial colonies (Comm, N $_{colony}$ = 15), Indiana mite biters (mite biter, N $_{colony}$ = 59), and
mite biter open-mated colonies (mite biter open mated, N = 7). The open circle is the outlier of
colonie. $F(2, 80) = 1.1992$, $p = 0.3069$. The p-value corresponding to the F-statistic of one-way
ANOVA suggests that all three treatments are no significantly different.

Figure 4. The violin plot of the percentage of damaged mites identified versus total mites
collected per colony among three groups, commercial colonies (Comm, N $_{colony}$ = 15, N $_{mite}$ =
172), Indiana mite biters (mite biter, N $_{colony}$ = 59, N $_{mite}$ =1,201), and mite biter open mated
colonies (mite biter open mated, N $_{colony}$ = 7, N $_{mite}$ = 199). The mean of the percentage of
damaged mites per colony among three treatments (commercial, mite biter, and mite biter open
mated) after transformation are 39.80%, 60.22%, and 60.14%. One-way ANOVA analysis
suggests that one or more groups are significantly different (F(2, 80) = 31.3785, p = 1.0031e-10).
The Tukey HSD test identified that the levels of mite biting behavior or grooming behavior in
the mite biter are significantly higher than the commerican colonies (Q = 11.0866, p < 0.01).
Levels of mite biting behavior or grooming behavior in the mite biter open mated colonies are
significantly higher than the commercial colonies (Q = 6.9771, p < 0.01). Levels of mite biting
behavior or grooming behavior between mite biter and mite biter open mated colonies are not
significantly different (Q = 0.0304, p > 0.05).

Figure 5. The box plot of the average number of legs missing per mite per colony among two
groups, commercial colonies (Comm, N $_{colony}$ = 15) and mite biter open-mated colonies (mite
biter open mated, N $_{colony}$ = 7). The open circle was outliers. Our result showed no difference
regarding to the average number of legs missing per mite between two groups (Q = 2.4278, p >
0.05).

Figure 6. Six parameters of the scanned mandible are listed from the microCT image. L- length,
W –width, H-height, SE-short edge, LE – long edge, S-span of spine area.

Figure 7. The box plot of the six parameters of mandible morphology between two groups,
commercial colonies (Comm, N $_{mandible}$ = 10) and mite biters (mite biter, N $_{mandible}$ = 15). Each
filled circle is each data point. The p-value corresponding to the F-statistic of one-way ANOVA
suggests that two treatments are not significantly different in length, width, height, short edge,
the span of spine area, but significantly different in long edge (F (1, 21) = 5.7762, p < 0.05).
There is no significant difference of other five parameters between two groups, but in the length,

height, short edge and span of the spine area, we noticed a consistant trend of smaller value in
mite biter colonies.

Figure 1

Figure 2

Figure 3

Figure 4

Figure 5

Figure 6

Figure 7

Reference

Andino GK, Hunt GJ, 2011. A scientific note on a new assay to measure honeybee mite-grooming
behavior. Apidologie 42:481-484.
Arechavaleta-Velasco ME, Guzmán-Novoa E, 2001. Relative effect of four characteristics that restrain
the population growth of the mite Varroa destructor in honey bee (Apis mellifera) colonies.
Apidologie 32:157-174.
Bailey L, 1968. HONEY BEE PATHOLOGY. Annual Review of Entomology 13:191-+. doi:
10.1146/annurev.en.13.010168.001203.
Bienefeld K, Zautke F, Pronin D, Mazeed A, 1999. Recording the proportion of damaged Varroa jacobsoni
Oud. in the debris of honey bee colonies (Apis mellifera). Apidologie 30:249-256.
Boecking O, Ritter W, 1993. Grooming and removal behaviour of Apis mellifera intermissa in Tunisia
against Varroa jacobsoni. Journal of Apicultural Research 32:127-134.
Buechler R, Berg S, Le Conte Y, 2010. Breeding for resistance to Varroa destructor in Europe. Apidologie
41:393-408. doi: 10.1051/apido/2010011.
Büchler R, Berg S, Le Conte Y, 2010. Breeding for resistance to Varroa destructor in Europe. Apidologie
41:393-408.
Coty D, Aria C, Garrouste R, Wils P, Legendre F, Nel A, 2014. The first ant-termite syninclusion in amber
with CT-scan analysis of taphonomy. PloS one 9:e104410.
Danka RG, Harris JW, Villa JD, 2011. Expression of Varroa sensitive hygiene (VSH) in commercial VSH
honey bees (Hymenoptera: Apidae). Journal of economic entomology 104:745-749.
Di Prisco G, Pennacchio F, Caprio E, Boncristiani Jr HF, Evans JD, Chen Y, 2011. Varroa destructor is an
effective vector of Israeli acute paralysis virus in the honeybee, Apis mellifera. Journal of
General Virology 92:151-155.
Fries I, Wei HZ, Shi W, Chen SJ, 1996. Grooming behavior and damaged mites (Varroa jacobsoni) in Apis
cerana cerana and Apis mellifera ligustica. Apidologie 27:3-11. doi: 10.1051/apido:19960101.
Generisch E, 2010. Honey bee pathology: current threats to honey bees and beekeeping. Applied
Microbiology and Biotechnology 87:87-97. doi: 10.1007/s00253-010-2573-8.
Guarna MM, Hoover SE, Huxter E, Higo H, Moon K-M, Domanski D, Bixby MEF, Melathopoulos AP,
Ibrahim A, Peirson M, Desai S, Micholson D, White R, Borchers CH, Currie RW, Pernal SF, Foster
LJ, 2017. Peptide biomarkers used for the selective breeding of a complex polygenic trait in
honey bees. Scientific Reports 7. doi: 10.1038/s41598-017-08464-2.
Guarna MM, Melathopoulos AP, Huxter E, Iovinella I, Parker R, Stoynov N, Tam A, Moon K-M, Chan
QWT, Pelosi P, White R, Pernal SF, Foster LJ, 2015. A search for protein biomarkers links
olfactory signal transduction to social immunity. Bmc Genomics 16. doi: 10.1186/s12864-014-
1193-6.
Guzman-Novoa E, Emsen B, Unger P, Espinosa-Montano LG, Petukhova T, 2012. Genotypic variability
and relationships between mite infestation levels, mite damage, grooming intensity, and
removal of Varroa destructor mites in selected strains of worker honey bees (Apis mellifera L.).
Journal of Invertebrate Pathology 110:314-320. doi: 10.1016/j.jip.2012.03.020.
Guzmán-Novoa E, Eccles L, Calvete Y, Mcgowan J, Kelly PG, Correa-Benítez A, 2010. Varroa destructor is
the main culprit for the death and reduced populations of overwintered honey bee (Apis
mellifera) colonies in Ontario, Canada. Apidologie 41:443-450.

Hamiduzzaman MM, Emsen B, Hunt GJ, Subramanyam S, Williams CE, Tsuruda JM, Guzman-Novoa E, 2017. Differential Gene Expression Associated with Honey Bee Grooming Behavior in Response to Varroa Mites. Behavior Genetics 47:335-344. doi: 10.1007/s10519-017-9834-6.

Hunt G, Given JK, Tsuruda JM, Andino GK, 2016. Breeding mite-biting bees to control Varroa. Bee Culture 8:41-47.

Kanga LHB, Marshall K, Legaspi JC, 2016. Mechanisms of insecticide resistance in field populations of the varroa mite (Acari: Mesostigmata: Varroidae) in Florida. Florida Entomologist 99:324-326. doi: 10.1653/024.099.0231.

Kingston AM, Myers GR, Latham SJ, Recur B, Li H, Sheppard AP, 2018. Space-filling X-ray source trajectories for efficient scanning in large-angle cone-beam computed tomography. IEEE Transactions on Computational Imaging 4:447-458.

Larabee FJ, Gronenberg W, Suarez AV, 2017. Performance, morphology and control of power-amplified mandibles in the trap-jaw ant Myrmoteras (Hymenoptera: Formicidae). Journal of Experimental Biology 220:3062-3071.

Le Conte Y, Ellis M, Ritter W, 2010. Varroa mites and honey bee health: can Varroa explain part of the colony losses? Apidologie 41:353-363.

Li D, Zhang K, Zhu P, Wu Z, Zhou H, 2011. 3D configuration of mandibles and controlling muscles in rove beetles based on micro-CT technique. Analytical and bioanalytical chemistry 401:817-825.

Meixner MD, Costa C, Kryger P, Hatjina F, Bouga M, Ivanova E, Buechler R, 2010. Conserving diversity and vitality for honey bee breeding. Journal of Apicultural Research 49:85-92. doi: 10.3896/ibra.1.49.1.12.

Mondragón L, Spivak M, Vandame R, 2005. A multifactorial study of the resistance of honeybees Apis mellifera to the mite Varroa destructor over one year in Mexico. Apidologie 36:345-358.

Morfin N, Given K, Evans M, Guzman-Novoa E, Hunt GJ, 2019. Grooming behavior and gene expression of the Indiana "mite-biter" honey bee stock. Apidologie:1-9.

Nazzi F, Le Conte Y, 2016. Ecology of Varroa destructor, the Major Ectoparasite of the Western Honey Bee, Apis mellifera. Annual Review of Entomology, Vol 61 61:417-432. doi: 10.1146/annurev-ento-010715-023731.

Oddie MAY, Dahle B, Neumann P, 2017. Norwegian honey bees surviving Varroa destructor mite infestations by means of natural selection. Peerj 5. doi: 10.7717/peerj.3956.

Papachristoforou A, Kagiava A, Papaefthimiou C, Termentzi A, Fokialakis N, Skaltsounis A-L, Watkins M, Arnold G, Theophilidis G, 2012. The Bite of the Honeybee: 2-Heptanone Secreted from Honeybee Mandibles during a Bite Acts as a Local Anaesthetic in Insects and Mammals. Plos One 7. doi: 10.1371/journal.pone.0047432.

Peng YS, Fang YZ, Xu SY, Ge LS, 1987. THE RESISTANCE MECHANISM OF THE ASIAN HONEY-BEE, APIS-CERANA FABR, TO AN ECTOPARASITIC MITE, VARROA-JACOBSONI OUDEMANS. Journal of Invertebrate Pathology 49:54-60. doi: 10.1016/0022-2011(87)90125-x.

Pritchard DJ, 2016. Grooming by honey bees as a component of varroa resistant behavior. Journal of Apicultural Research 55:38-48.

Ramsey SD, Ochoa R, Bauchan G, Gulbronson C, Mowery JD, Cohen A, Lim D, Joklik J, Cicero JM, Ellis JD, 2019. Varroa destructor feeds primarily on honey bee fat body tissue and not hemolymph. Proceedings of the National Academy of Sciences 116:1792-1801.

Ribi W, Senden TJ, Sakellariou A, Limaye A, Zhang S, 2008. Imaging honey bee brain anatomy with micro-X-ray-computed tomography. Journal of neuroscience methods 171:93-97.

Rinderer TE, Harris JW, Hunt GJ, De Guzman LI, 2010. Breeding for resistance to Varroa destructor in North America. Apidologie 41:409-424.

Russo RM, Liendo MC, Landi L, Pietronave H, Merke J, Fain H, Muntaabski I, Palacio MA, Rodríguez GA,
Lanzavecchia SB, 2020. Grooming Behavior in Naturally Varroa-Resistant Apis mellifera Colonies
From North-Central Argentina. Frontiers in Ecology and Evolution 8:361.
Ruttner F, Hanel H, 1992. ACTIVE DEFENSE AGAINST VARROA MITES IN A CARNIOLAN STRAIN OF
HONEYBEE (APIS-MELLIFERA-CARNICA POLLMANN). Apidologie 23:173-187. doi:
10.1051/apido:19920210.
Spivak M, 1996. Honey bee hygienic behavior and defense against Varroa jacobsoni. Apidologie 27:245-
260. doi: 10.1051/apido:19960407.
Tsuruda JM, Harris JW, Bourgeois L, Danka RG, Hunt GJ, 2012. High-Resolution Linkage Analyses to
Identify Genes That Influence Varroa Sensitive Hygiene Behavior in Honey Bees. Plos One 7. doi:
10.1371/journal.pone.0048276.
Villa JD, Danka RG, Harris JW, 2017. Repeatability of measurements of removal of mite-infested brood to
assess Varroa Sensitive Hygiene. Journal of Apicultural Research 56:631-634.
Weissgerber TL, Savic M, Winham SJ, Stanisavljevic D, Garovic VD, Milic NM, 2017. Data visualization,
bar naked: A free tool for creating interactive graphics. Journal of Biological Chemistry
292:20592-20598.
Wilfert L, Long G, Leggett H, Schmid-Hempel P, Butlin R, Martin S, Boots M, 2016. Deformed wing virus is
a recent global epidemic in honeybees driven by Varroa mites. Science 351:594-597.
Yue M, Luo S, Liu J, Wu J, 2018. Apis cerana is less sensitive to most neonicotinoids, despite of their
smaller body mass. Journal of economic entomology 111:39-42.