Heterogeneity of monocyte subsets and susceptibility to influenza virus contribute to inter-population variability of protective immunity

Mary O'Neill[1], Hélène Quach[2], Julien Pothlichet[3], Yann Aquino[1], Aurélie Bisiaux[1], Nora Zidane[4], Matthieu Deschamps[1], Valentina Libri[5], Milena Hasan[5], Shen-Ying Zhang[6,7,8], Qian Zhang[6], Daniela Matuozzo[7,8], Aurélie Cobat[7,8], Laurent Abel[6,7,8], Jean-Laurent Casanova[6,7,8,9], Nadia Naffakh[10], Maxime Rotival[1,12], Lluis Quintana-Murci[1,11,12]

[1]Human Evolutionary Genetics Unit, Institut Pasteur, UMR 2000, CNRS, Paris 75015, France
[2]Muséum National d'Histoire Naturelle, UMR7206, CNRS, Université de Paris, Paris 75016, France
[3]DIACCURATE, Paris, 75014, France
[4]Biodiversity and Epidemiology of Bacterial Pathogens Unit, Institut Pasteur, Paris 75015, France
[5]Cytometry and Biomarkers UTechS, Institut Pasteur, Paris 75015, France
[6]St. Giles Laboratory of Human Genetics of Infectious Diseases, Rockefeller Branch, The Rockefeller University, New York, NY, 10065, USA
[7]Laboratory of Human Genetics of Infectious Diseases, Necker Branch, Necker Hospital for Sick Children, INSERM UMR 1163, Necker Hospital for Sick Children, Paris 75015, France
[8]Paris University, Imagine Institute, Paris 75015, France
[9] Howard Hughes Medical Institute, New York, NY, USA
[10]RNA Biology of Influenza Virus Unit, Institut Pasteur, Paris, 75015, France
[11]Chair of Human Genomics and Evolution, Collège de France, Paris, 75005, France

[12]M.R. and L.Q.M. contributed equally to this work.
Correspondence and requests for materials should be addressed to M.R or L.Q.-M. (email: maxime.rotival@pasteur.fr, quintana@pasteur.fr).

Authors contributions: M.B.O, M.R. and L.Q.M. conceived and designed the study. M.B.O., H.Q., J.P., Y.A., A.B., N.Z., and M.D. conducted the experiments at the Human Evolutionary Genetics Unit. M.B.O., M.R., Y.A. and D.M. developed computational methods and performed bioinformatic analyses. J.P., V.L., M.H., S.Z.Y., Q.Z., A.C. L.A., J.L.C. and N.N. provided resources, expertise and feedback. M.R. and L.Q.M supervised the study. L.Q.M secured funding. M.B.O., M.R. and L.Q.M. wrote the manuscript with input from all authors.

Competing Interest Statement: The authors declare no competing interests.

Keywords: Influenza, monocytes, single-cell, transcriptomics, ancestry

This PDF file includes:
    Main Text
    Figures 1 to 5

Abstract

There is considerable inter-individual and inter-population variability in response to viruses. The

potential of monocytes to elicit type-I interferon responses has attracted attention to their role in

viral infections. Here, we use an *ex vivo* model to characterize the role of cellular heterogeneity

in human variation of monocyte responses to influenza A virus (IAV) exposure. Using single-cell

RNA-sequencing, we show widespread inter-individual variability in the percentage of IAV-

infected monocytes. We show that cells escaping viral infection display increased mRNA

expression of type-I interferon stimulated genes and decreased expression of ribosomal genes,

relative to both infected cells and those never exposed to IAV. While this host defense strategy

is shared between $CD16^+/CD16^-$ monocytes, we also uncover $CD16^+$-specific mRNA expression

of *IL6* and *TNF* in response to IAV, and a stronger resistance of $CD16^+$ monocytes to IAV

infection. Notably, individuals with high cellular susceptibility to IAV are characterized by a lower

activation at basal state of an IRF/STAT-induced transcriptional network, which includes antiviral

genes such as *IFITM3*, *MX1*, and *OAS3*. Finally, using flow cytometry and bulk RNA-sequencing

across 200 individuals of African and European ancestry, we observe a higher number of $CD16^+$

monocytes and lower susceptibility to IAV infection among monocytes from individuals of

African-descent. Collectively, our results reveal the effects of IAV infection on the transcriptional

landscape of human monocytes and highlight previously unappreciated differences in cellular

susceptibility to IAV infection between individuals of African and European ancestry, which may

account for the greater susceptibility of Africans to severe influenza.

Significance Statement

Monocytes may play a critical role during severe viral infections. Our study tackles how

heterogeneity in monocyte subsets and activation contributes to shape individual differences in

the transcriptional response to viral infections. Using single-cell RNA-sequencing, we reveal

heterogeneity in monocyte susceptibility to IAV infection, both between $CD16^+/CD16^-$ monocytes

and across individuals, driven by differences in basal activation of an IRF/STAT-induced antiviral

program. Furthermore, we show a decreased ability of IAV to infect and replicate in monocytes

from African-ancestry individuals, with possible implications for antigen presentation and

lymphocyte activation. These results highlight the importance of early cellular activation in

determining an individuals' innate immune response to viral infection.

Main Text

Introduction

Respiratory viruses with pandemic potential pose enormous health and economic impacts on the

human population. In the last century, we have witnessed outbreaks of several coronaviruses,

including SARS-CoV-2, SARS-CoV-1 and MERS, and a number of avian and swine influenza A

viruses (IAV). A particularly harrowing and shared feature of these pandemics are the sudden

deaths of otherwise healthy individuals (1). A hyperinflammatory state characterized by high

levels of inflammatory cytokines, often referred to as a 'cytokine storm' (2, 3), has emerged as a

hallmark of these severe viral infections. While still controversial, there is increasing evidence to

suggest that the mononuclear phagocyte system is an important immunological determinant of

this phenotype (4-6). Upon viral infection, sentinel cells such as lung-resident macrophages

trigger complex signaling cascades that recruit leukocytes to the site of infection, among them

monocytes. These infiltrating monocytes differentiate into monocyte-derived dendritic cells or

macrophages, enabling viral clearance through the induction of the adaptive response, and help

replenish the pool of tissue-resident alveolar macrophages (4, 7).

In humans, circulating monocytes are divided into classical (~80%), intermediate (~15%),

and nonclassical (~5%) subsets, based on surface receptor expression of the cluster-

determinant antigens CD14 and CD16 (8). While nonclassical monocytes ($CD14^+CD16^{++}$) are

long-lived and 'patrol' healthy tissues through long-range crawling on the endothelium, classical

($CD14^{++}CD16^-$) and intermediate ($CD14^{++}CD16^+$) monocytes are recruited to the lung in

response to viral infection, where they secrete inflammatory cytokines and chemokines, as well

as type I interferons (IFNs) (7, 9-11). In most individuals, recruited cells help clear infection

despite being susceptible to infection themselves (12, 13); yet, in some individuals, a

dysfunctional immune response occurs resulting in widespread lung inflammation. Whether

monocyte subsets behave differently upon viral exposure, and how direct viral sensing and

exposure to secreted cytokines shape monocyte activation and differentiation are not well

understood.

Variation in blood composition and cellular proportions have been shown to be one of the

main factors underlying transcriptional variation in immune genes across individuals (14), with

these proportions being influenced by both genetic and non-heritable factors (15-17). Recently,

we characterized the genetic architecture of transcriptional responses of primary monocytes

from 200 individuals of African and European ancestry to *ex vivo* challenge with viral stimuli (18).

In this model, where we were able to control for viral determinants of disease (i.e. dose and

strain), we reported marked inter- and intra-population differences in transcriptional responses to

IAV. While our analyses revealed numerous *cis*-expression quantitative trait loci (18), genetic

variants could only account for a small fraction of expression variation, in line with other studies

(14, 19).

Here, we implemented single-cell RNA-sequencing (scRNA-seq) on human primary

monocytes exposed to IAV to investigate (i) the effects of direct viral infection versus activation

by exposure to secreted cytokines, (ii) the subset-specific responses of monocytes to viral

challenge, and (iii) the extent of interindividual and between-population variation in the

proportions of monocyte subsets and the degree of monocyte susceptibility to IAV infection.

Results

Using single-cell RNA-sequencing to investigate cellular heterogeneity. To investigate the

role of cellular heterogeneity in driving immune variability across individuals, we performed a

time-course experiment where we monitored the $CD14^+$ fraction of peripheral blood

mononuclear cells (PBMCs) from eight donors, both in the presence and absence of viral

challenge. To maximize inter-individual variability, we chose individuals from two distinct

ancestries whose cells demonstrated extreme responses to viral stimuli in a previous bulk RNA-

seq experiment (18). Droplet-based scRNA-seq was performed on monocytes from all eight

donors immediately before infection initiation (T_0), as well as at 2 (T_2), 4 (T_4), 6 (T_6), and 8 (T_8)

hours post challenge with A/USSR/90/1977(H1N1) at a multiplicity of infection (MOI) equal to 1

(IAV-challenged) and mock infection (non-infected). To mitigate batch effects, we pooled IAV-

challenged and non-infected cells from distinct donors in each library, assigning cells to their

condition in silico via genetic barcoding (20). After stringent quality control, our final dataset

contained 88,559 high-quality cells, among which we predicted >99% monocyte purity at T_0 (**Fig.**

1*A*; *SI Appendix*, Figs. S1 and S2). At later time points, a substantial fraction of non-infected

cells (up to 70% at T_8) were predicted to be macrophage-like, indicating monocyte differentiation

over the course of the experiment. For clarity, we refer to cells as monocytes at T_0 and as

monocyte-derived cells from T_2-T_8.

Stable *FCGR3A* expression distinguishes monocyte subsets over time. We next sought to

characterize each cell by its mRNA expression of the canonical monocyte markers, CD14 and

CD16, given that much of the structure in our data was associated with *FCGR3A* (aka *CD16*)

mRNA expression. In droplet-based scRNA-seq, encapsulation of ambient mRNAs emanating

from dying cells can occur during library preparation leading to spurious mRNA detection (21).

We thus used a statistical framework to test whether *CD14* and *CD16* were expressed at a level

significantly higher than expected when accounting for potential contamination from the ambient

pool (***Methods***). Despite having been positively selected for the CD14 antigen, only 32.4% of

monocytes significantly expressed *CD14* at T_0; this percentage further decreased at later time

points and remained <15% across all time points and conditions (average 6.4% s.d.: 5.0%, ***SI***

Appendix,* Fig. S3*A-C). On the other hand, 12.1% of monocytes significantly expressed

FCGR3A (*CD16*) (referred to as *CD16*$^+$) at T_0, this marker proving much more stable across

conditions and time points (9.3% of *CD16*$^+$ cells on average, s.d.: 1.8%, **Fig. 1*B* and *SI***

Appendix,* Fig. S3*D-F). While we deciphered classical, intermediate, and nonclassical

monocytes subsets at T_0 (***SI Appendix*, Note 1 and Fig. S4; Dataset S1**), we focus on the

simpler distinction of *CD16*$^-$ and *CD16*$^+$ subsets given that positive-selection for monocytes does

not capture the entire nonclassical population and that we were unable to distinguish the

intermediate and noncanonical subsets after T_0.

Functional features of monocyte subsets are conserved upon manipulation. To assess

how transcriptional profiles of *CD16*$^-$ and *CD16*$^+$ monocytes and their derived-cells differ, we

focused on the 5,681 genes expressed with a normalized log count > 0.1 in at least one

condition, time point, and subset (**Dataset S2*A***). We found that the $\log_2$ fold change in gene

expression between *CD16*$^{+/-}$ subsets remained relatively stable over the course of the

experiment (Pearson r between time points >0.42, and >0.52 for the non-infected and IAV-

challenged conditions respectively, *p*-values < 2.2×10^{-16}; ***SI Appendix,* Fig. S5*A***), and

differentially expressed genes between *CD16*$^{+/-}$ subsets were largely the same across conditions

(Pearson r = 0.92, *p*-value < 2.2×10^{-16}; ***SI Appendix,* Fig. S5*B***). We thus searched for genes

that were consistently differentially expressed between *CD16*$^+$ and *CD16*$^-$ cells across all time

points (including T_0), conditions, and donors. We identified 266 genes over-expressed

($\log_2$FC>0.2, FDR<1%) in *CD16*$^+$ cells relative to *CD16*$^-$ cells, and 389 genes that showed the

opposite pattern, and performed a GO-term enrichment analysis on these genes (**Dataset S2*B***).

Consistent with previous reports (22-24), *CD16*$^-$ subsets were characterized by high expression

of several proinflammatory *S100 Calcium Binding Proteins* (*S100A12*, *S100A9*, and *S100A8*),

contributing to a sizable GO-term enrichment in the defense response to fungus pathway

(GO:0050832: OR=41.3, FDR=4.9×10^{-4}), while *CD16*$^+$ subsets were characterized by high

expression of Fc-gamma receptor signaling pathway genes (GO:0038096: OR=8.7,

FDR=6.2×10^{-6}). Notably, *CD16*$^+$ subsets over-expressed several type I IFN stimulated genes

(ISGs) relative to *CD16*$^-$ subsets (e.g. GO:000071357: OR=5.3, FDR=2.6×10^{-3}), including the

well-known viral restriction factors *IFITM3* and *OAS1*. Collectively, these results demonstrate

CD16 is a reliable marker at the mRNA level and that *CD16*$^{+/-}$ monocyte subsets maintain

functional differences upon manipulation.

scRNA-seq highlights heterogeneity in monocyte susceptibility and viral transcription.

Using the presence of IAV transcripts as a proxy for infection (**Fig. 1*B***), we next sought to

distinguish cells that were successfully infected from those that were not. Among monocyte-

derived cells that were exposed to IAV, we found that 50.3% expressed IAV transcripts above

ambient levels when allowing up to 10% of mRNAs to come from the ambient pool. In contrast,

less than 1% of non-infected cells showed evidence of viral transcription, supporting the validity

of the threshold used to detect IAV expressing cells (**Fig. 1*C***). We deemed cells with statistical

evidence for expression of IAV transcripts from the IAV-challenged condition as 'infected', while

the remaining cells from this condition were considered as 'bystanders', as these either did not

come into contact with the virus or were able to fully repress viral mRNA transcription. When

comparing the percentage of infected cells between subsets, we noticed that *CD16*$^+$ cells were

slightly less likely to be infected than *CD16*$^-$ cells (42.3% sd: 4.0% for *CD16*$^+$ relative to 49.4%

sd: 5.4% for *CD16*$^-$, generalized linear model with CD16$^{+/-}$ status, donor, and time point as

covariates, *p*-value=0.006). This suggests a higher resistance of *CD16*$^+$ to IAV infection,

possibly related to the higher expression of ISGs observed in this subset (**Dataset S2 *A* and *B***).

We observed that the proportions of viral mRNAs among infected cells were bimodally

distributed and largely varied between the clusters identified in our unsupervised analysis (**Fig.**

1*D*). We used a Gaussian mixture model to locate the two modes of the distribution and further

sub-classify infected cells into those with lower IAV mRNA levels (<1-6%) and those with higher

IAV mRNA levels (6-83%); while viral mRNA levels are dictated by both the rate of transcription

and degradation, for simplicity we refer to these infected cell states as 'low IAV-transcribers' and

'high IAV-transcribers', respectively. The proportions of infected cells among individuals

remained largely unchanged over the course of the experiment; however, high-IAV transcribers

were virtually absent at 2h (<2% of infected cells), peaked to ~36% of IAV-infected cells at 4h,

and decreased to 8.5% by 8h, suggesting that high-IAV transcribers represent a transient state

of IAV-infection preceding IAV-induced apoptosis (**Fig. 1*E***). These results reveal profound

heterogeneity in monocyte susceptibility and subsequent viral transcription upon IAV-challenge.

Interplay of cytokine and ribosome networks drive cell states upon infection. To

characterize host transcriptional responses over time, we next subsampled each subset (*CD16*$^-$

/*CD16*$^+$), cell state (unexposed, bystander, infected), and time point in our scRNA-seq data to

100 cells, while ensuring a balanced representation of all donors. We then focused on the 6,669

host genes with average $\log_2$ normalized count >0.1 in at least one subgroup (**Dataset S3*A***).

Overall, *CD16*⁻ and *CD16*⁺ subsets behaved similarly upon stimulation with changes in gene

expression between cell states being strongly correlated among subsets (Pearson r = 0.83-0.95,

p-values < 2.2×10^{-16}; ***SI Appendix,* Fig. S6)**. GO term enrichment analyses of shared responses

(FDR<1% & $\log_2$FC >0.2 in same direction in both subsets) uncovered several functional

categories interacting to shape the activation state of cells (**Fig. 2*A***; **Dataset S3*B***). Both

bystander and infected cells showed increased mRNA expression of genes involved in antigen

processing and presentation via class I MHC (GO:0019885, OR=53.7, FDR < 2.0×10^{-6}) and

response to type I IFN (GO:0034340, OR >14.8, FDR < 3.3×10^{-20}). Yet, bystander cells showed

increased mRNA expression of type I IFN response and defense response to virus pathways

relative to infected cells (GO:0034340: OR=13.4, FDR=4.4×10^{-7}; GO:0051607: OR=9.0,

FDR=2.1×10^{-7}), while infected cells displayed higher mRNA expression of mitochondrial

(GO:0005743, OR=4.7, FDR=3.3×10^{-3}) and ribosomal genes (GO: 0005840, OR=117,

FDR=1.0×10^{-78}).

Among infected cells, ribosomal genes showed higher activity among high IAV-transcribing

cells relative to low IAV-transcribing cells (**Fig. 2*B***, comparison only made at T_4 due to sample

size constraints, e.g. GO:0019083: OR=137, FDR=6.1×10^{-65}). This observation is consistent with

the notion that the expression of viral proteins is dependent on cellular ribosomes, with recent

data suggesting that IAVs do not induce a global shut-off of cellular translation but rather a

reshaping of the translation landscape (25-27). Likewise, among bystander cells, numerous

ribosomal genes were downregulated at later time points relative to unexposed cells (**Fig. 2*A***

and *C*; GO:0019083, OR=5.3, FDR=4.2×10^{-6}), suggesting that repression of ribosomal subunits

plays an active role in limiting viral replication. Collectively, these results suggest that ISGs and

ribosomal expression interact to shape cell states upon IAV-challenge.

Increased IRF and STAT activity drives stronger antiviral response. Despite qualitatively

similar responses to infection between CD16⁻/CD16⁺ subsets (***SI Appendix,* Fig. S6)*,* we

hypothesized that subtle differences in the intensity of such responses might contribute to the

increased resistance of *CD16*⁺ cells to infection. We thus performed an interaction test on the

subsampled scRNA-seq data, and searched for genes for which transcriptional response upon

IAV-challenge differed between *CD16*⁻ and *CD16*⁺ subsets in either infected and/or bystander

cells (***SI Appendix,* Fig. S6 *A* and *B***; **Dataset S3*A***). At FDR≤1%, we identified a total of 335

such genes, of which 98 differed between subsets only in bystander cells, 144 only in infected

cells, and 93 in both. Hierarchical clustering highlighted eight major patterns of transcriptional

responses (modules) among the 335 genes, several of which were associated with specific

biological functions (**Fig. 3*A*; Dataset S3 *C* and *D***). Notably, module 1 (green) was enriched for

genes in the antiviral response pathway (GO:0051607, OR=23.2, FDR<5.43×10^{-7}) and displayed

a stronger response in infected *CD16$^+$* cells relative to *CD16$^-$* infected cells. Of additional interest

was the transient *CD16$^+$* specific transcription of the inflammatory cytokine genes *IL6* and *TNF*,

following viral challenge (**Fig. 3*B***). We also found that several genes involved in the regulation

and production of IL-6 and TNFα are over-expressed in *CD16$^+$* subsets at all time-points and

conditions (**Dataset S2*B***), but only see active transcription of the cytokines upon viral exposure.

These results reveal the strong antiviral and inflammatory potential of *CD16$^+$* relative to *CD16$^-$*

monocytes in response to viral infection (28).

We next sought to characterize the regulatory architecture underlying the 335 genes whose

transcriptional response to IAV-challenge differed between monocyte subsets. Using SCENIC

(29), we identified 113 high-confidence gene regulatory networks, or 'regulons', which are active

in non-infected and/or IAV-challenged cells, each composed of a transcription factor (TF) and a

set of predicted targets (genes). We used these 113 regulons to search for an

enrichment/depletion of TF targets among the eight modules of genes displaying subset-specific

response to infection (**Dataset S3*E***). Among modules associated with an increased expression

in cells exposed to IAV (modules 1-5), we observed a widespread over-representation of targets

of IFN regulatory factors (IRFs) and signal transducing and activators of transcription (STATs)

(**Fig. 3*C***), reinforcing the central role of the IFN response upon IAV challenge. Interestingly,

several of these factors displayed subset-specific activity themselves in response to IAV

(*IRF1/2/7* and *STAT1/2/3*, FDR<1%), mirroring the expression patterns of module 1 (Pearson

r>0.92). These results collectively highlight a *CD16$^+$*-specific inflammatory response upon IAV-

challenge and suggest stronger activation of IRF and STAT transcription factors as driver of the

increased antiviral response observed in *CD16$^+$* cells upon IAV infection.

Basal activation differences correlate with monocyte susceptibility. To explore the degree

of inter-individual variation upon viral challenge, we next quantified IAV transcripts in the

monocyte-derived cells of each individual, and created pseudo-bulk estimates by averaging the

percent of viral mRNAs per-cell across all cells from each donor at each time point (**Fig. 4*A***).

While viral mRNAs peaked at the same time for all individuals, we observed extensive variation

in the levels of viral mRNAs and percentages of infected cells across individuals (**Fig. 4*B***). To

identify specific genes that might underlie infection potential, we focused on the 4,589 genes

that were expressed at > 0.1 $\log_2$ normalized counts in at least one canonical monocyte subset

at T_0. We identified a total of 3,131 genes that differed among our eight donors in either

classical, intermediate, and/or nonclassical monocyte subsets (Kruskal-Wallis Rank Test, FDR

1%; **Dataset S4*A***). Within each subset, focusing on genes that significantly differ between

donors, we searched for those for which mean expression at basal state was correlated with the

percentage of infected cells at T_4 among our eight donors. Despite our limited sample size, we

found that cellular susceptibility was strongly correlated with basal expression of the well-known

host viral restriction factor *IFITM3.* Although it reached significance only in nonclassical

monocytes (FDR~1%), the association remained strong in other subsets ($p < 4.1 \times 10^{-4}$; **Fig. 4*C***).

We next relaxed our search to all genes for which basal expression showed nominal

correlation ($p < 0.01$) with the percentage of infected cells at T_4. Depending on the monocyte

subset, between 3.6 to 8.3% of genes matched these criteria, resulting in a set of 118 genes

displaying correlation with monocyte susceptibility in at least one subset. These 118 genes were

collectively enriched for several related biological processes such as defense response to virus

(GO:0051607, OR=15.3, FDR=9.2×10^{-19}) and response to type I IFN (GO:0034340, OR=19.6

FDR=8.4×10^{-15}) (**Dataset S4*B***). Among genes contributing to this enrichment, we found

additional antiviral genes such as *OAS3*, and *MX1*, as well as the critical TF, *IRF7*, involved in

the severity of IAV-infection both in mice and humans (30-32). Finally, overlap with the TF

targets identified by SCENIC revealed strong enrichments of several IRFs and STATs among

the 118 genes, including IRF7, as well as STAT1, STAT2 and IRF9 that form the tripartite IFN-

stimulated gene factor 3 (ISGF3) (**Fig. 4*D***; **Dataset S4*C***). Together, our results provide

evidence that the basal mRNA expression of genes related to IFN-induced and antiviral

responses are indicative of the proportion of cells that will become infected in the first cycle of

IAV infection.

African-ancestry monocytes are more resistant to infection. Lastly, we wondered how our

findings of inter-individual variation might extrapolate to the population level. In a previous study

(18), we challenged the primary monocytes from 200 Belgian individuals of African (AFB) and

European (EUB) ancestry with the same IAV strain and MOI used in the present study, and

performed bulk RNA-seq at 6 hours post infection (hpi). While basal (T_0) expression profiles

were not collected, flow cytometry labelling of CD14 and CD16 was performed on the CD14$^+$-

selected monocytes for the majority of donors. Interestingly, AFB individuals had higher

proportions of *CD16*$^+$ cells than EUB individuals (**Fig. 5*A***; ***SI Appendix,* Fig. S7**). In light of our

findings that *CD16*[+] cells are more resistant to IAV infection, we hypothesized that this might translate to lower infection rates among AFB monocytes relative to EUB monocytes.

To test this hypothesis, we mapped the bulk RNA-seq profiles collected 6hpi challenge with IAV for the 200 individuals to a combined human-IAV reference. Excluding 1 sample with low quality RNAs, we found that 0.02-13.5% of RNA-seq reads from each sample were of viral origin (**Fig. 5B**). Reassuringly, these percentages correlated with IAV mRNA levels estimated from the single cell experiment across all time points for the eight donors used in the present study (Pearson r > 0.84, *p*-value < 8.9×10^{-3}), with the strongest correlation being observed at the peak of viral transcription (T_4) (Pearson r = 0.97, *p*-value=5.1×10^{-5}). These observations indicate that *ex vivo* cellular susceptibility is highly reproducible among individuals, even across different experimental protocols and technologies. Among the 199 bulk profiles, AFB and EUB samples presented overlapping but significantly shifted distributions of total IAV-mapping reads (**Fig. 5B**, 4.9% vs. 6.8% of reads, respectively, Wilcoxon *p*-value = 5.3×10^{-8}), and of each of the 10 primary viral transcripts (**Fig. 5C**, Wilcoxon *p*-value < 5.5×10^{-4}).

Using the transcriptional profiles obtained from the scRNA-seq data at T_6, we estimated the proportion of reads coming from each inferred cell state in these bulk RNA-seq profiles (**Fig. 5D and 5E**; *SI Appendix*, **Note S2 and Fig. S8A**). We found that, on average, AFB monocytes were more resistant to IAV infection than EUB monocytes (39.2% vs. 48.9% infected, respectively, Wilcoxon *p*-value = 5.3×10^{-10}). Differences in the estimated percentage of infected cells alone explained 63% of the inter-individual variability in viral mRNA levels (**Fig. 5F**), and was sufficient to account for the observed difference in viral mRNA levels between AFB and EUB individuals (*p*-value=0.16 after adjusting on infected cells, compared to *p*-value=5.3×10^{-8} without adjustment). Nonetheless, variation in the percentage of high/low transcribers among infected cells accounted for an additional 19% of variance in viral mRNA expression (*SI Appendix*, **Note S2 and Fig. S8B**). Finally, the ratio of *CD16*[+]/*CD16*[-] cells negatively correlated with the percentage of infected cells, albeit weakly (-0.27, *p*-value = 0.0165 adjusted on population). Altogether, these results show that population differences in viral mRNA levels are primarily driven by the overall proportion of cells that will ultimately become infected, with only a fraction of the differences being attributable to the different proportions of *CD16*[+/-] subsets observed in individuals of African and European ancestry.

Discussion

We performed scRNA-seq on primary monocytes, before and after *ex vivo* IAV-challenge, to assess transcriptional differences between monocytes infected by IAV (i.e. infected) versus

those activated only by exposure to secreted cytokines (i.e. bystanders), and to identify subset-

specific responses of monocytes to viral challenge. We found that bystander cells display

increased mRNA expression of ISGs relative to infected cells; yet, we additionally observed both

an induction of ribosomal gene mRNA expression in IAV-transcribing cells and a down

regulation of these genes in bystander cells at later time points. While the former is likely

induced by the virus to enhance mRNA translation (33), the repression of ribosomal expression

observed in bystander cells may reflect a host mechanism to contain infection by shutting down

the translational machinery of neighboring cells. Interestingly, the interplay of ribosomal and ISG

expression also distinguished infected cells into two distinct states (high and low IAV-

transcribers), providing an explanation for the high cell-to-cell variation in IAV replication

observed among circulating monocytes, which has also been documented in other cell types

and during natural infection (34-41).

While these patterns are generally shared across $CD16^-$ and $CD16^+$ subsets, we found

$CD16^+$ cells to be slightly more resistant to infection. This is likely attributable to their higher

absolute expression of some ISGs relative to $CD16^-$ cells (independent of viral exposure), as

well as their more robust upregulation of antiviral genes upon IAV-challenge, which we found to

be driven by stronger activity of IRF transcription factors. Interestingly, $CD16^+$ cells displayed

transient mRNA expression of $IL6$ and TNF upon viral exposure (both infected and bystander

cells), two cytokines that have been widely implicated in cytokine storms (5). Collectively, these

findings highlight the opposing roles of ISG and ribosomal gene mRNA expression on viral

transcription, and reveal the stronger antiviral and pro-inflammatory potential of $CD16^+$ monocyte

subsets.

At the population level, we found that the ratio of $CD16^+/CD16^-$ at basal state was predictive

of the percentage of monocytes that were susceptible to IAV infection, and observed that

African-ancestry individuals harbored more $CD16^+$ monocytes on average than European-

ancestry individuals residing in the same city (Ghent, Belgium), consistent with previous

observations (42). Independently of monocyte subset proportions, we identified that individuals

presenting lower monocyte susceptibility to IAV had a higher basal activation of an IRF/STAT-

driven antiviral program. These findings suggest that the fate of a monocyte hinges upon its

basal activation state, and that the infection potential differs both within an individuals' monocyte

population, in part based on the differentiation status of the cell (i.e. $CD16$-positivity), but also

between individuals, where a $CD16^-$ cell from one individual may have a higher antiviral state

than a $CD16^+$ cell from another individual. This latter phenomenon likely reflects the influence of

both genetic and non-heritable factors on transcriptional variation in immune genes (14, 15).

These inter- and intra-population differences are noteworthy in and of themselves, and raise questions about how such differences in susceptibility of monocytic cells to infection - and differences in proportions of monocyte subsets - may relate to viral disease.

African Americans are more often hospitalized than other self-defined ethnic groups by both influenza and COVID-19, even when adjusting for age and various social factors such as poverty and vaccination status (43, 44). Thus, should monocyte infection play a role in the severity of viral infections, our results would imply a paradox where higher infection and replication of IAV in circulating monocytes is associated with an advantage to fight viral infection *in vivo*. Such a paradox could possibly be explained by an enhanced antigen presenting capacity of infected cells, with higher infection of European-monocytes facilitating the activation of T cell antiviral responses (13, 45). In support of this hypothesis, we observe that infected and bystander cells display comparable induction of antigen-presentation genes relative to non-exposed cells. This suggests that infected cells maintain their ability to present viral antigens to the adaptive immune system, while actively producing these antigens. Independently, but not mutually exclusively, proportions of monocyte subsets in circulation may play a role in viral disease; of note, patients with severe influenza and COVID-19 harbor higher proportions of intermediate monocytes in peripheral blood than patients with mild disease (46, 47). Given our finding that *CD16*[+] subsets are the main drivers of inflammatory cytokine gene expression such as *IL6* and *TNF*, and that African-ancestry individuals harbor a larger fraction of these monocyte subsets, it is tangible to conceive that monocyte subset composition prior to infection may influence disease outcome, and potentially serve as a biomarker.

Materials and methods

Experimental model and subjects. All individuals from this study were part of the

EVOIMMUNOPOP cohort, which has been previously described (18). Briefly, 200 healthy male

donors living in Belgium of self-reported African descent (AFB) or European descent (EUB) were

recruited. Inclusion was restricted to nominally healthy individuals between 19 and 50 years of

age at the time of sample collection. Serological testing was performed for all donors to exclude

those with serological signs of past or ongoing infection with human immunodeficiency virus

(HIV), hepatitis B virus (HBV) or hepatitis C virus (HCV).

Single-cell analyses and RNA sequencing. For eight selected donors (4 individuals from each

ancestry, selected from extremes of the first principal component of gene expression in our

previous study of monocyte response to IAV-challenge (18)), 100×10^6 PBMCs were thawed,

washed twice and resuspended in complete medium: pre-warmed RPMI-1640 Glutamax

medium, supplemented with 10% FCS and 1% penicillin/streptomycin (Cat# 15140-122, Life

Technologies). Monocytes were then positively selected with magnetic CD14 microbeads,

according to the manufacturer's instructions (Cat#130-050-201, Miltenyi Biotec). The number of

monocytes was determined with the Countless2 automated cell counter system (Cat#

AMQAX1000, ThermoFisher Scientific) in the presence of trypan blue. For each donor,

monocytes were seeded at 0.5×10^6 monocytes per well on 24-well NUNC plates in 500 µL of

complete media and allowed to rest for one hour at 37°C under 5% CO_2. Five-hundred

microliters of complete media (non-infected) or A/USSR/90/1977(H1N1) at a concentration of

1×10^6 pfu/mL in complete media (IAV-challenged, MOI=1) were added to each sample.

Following one hour of staging at 4°C, plates were centrifuged at 1300 rpm for 10 minutes at 4°C,

media was removed by pipette, and each well was washed with 1mL complete media. The spin

was repeated, media removed by pipette, and samples were resuspended in 1mL pre-warmed

complete media before being transferred to an incubator at 37°C under 5% CO_2 to initiate

infection (T_0).

At each time point (T_0, T_2, T_4, T_6, and T_8), samples were mixed by pipetting and transferred

to Eppendorf tubes. Wells were washed with 300uL of PBS + 0.04% BSA and transferred to the

same tubes. Collection tubes were centrifuged at 1300 rpm for 10 minutes, media was removed

and replaced with 1mL PBS + 0.04% BSA and an aliquot of 10µL was taken to count each

sample on a Countless2 automated cell counter system, before repeating the centrifugations.

Individual samples were adjusted to 2×10^6 live cells/mL.

Samples were multiplexed for running on the 10X Chromium (Cat# 120223 & 1000074, 10X

Genomics) by mixing equal proportions from 6-8 samples in a manner that balanced conditions

and allowed us to assess for batch effects across lanes (*SI Appendix,* **Table S1**). Multiplexed

samples were counted with the Countless2 automated cell counter system and adjusted to

target recovery of 10,000 cells per reaction of the Chromium Single Cell 3' Reagent Kits v3

(Cat# 1000092 & 1000078, 10X Genomics) assuming a recovery rate of 50%. GEM Generation

& Barcoding, Post GEM-RT Cleanup & cDNA Amplification, and 3' Gene Expression Library

Construction were performed as per manufacturer's instructions (48). All 13 libraries were mixed

prior to sequencing across 13 different lanes from an Illumina HiSeq X (28bp barcode + 91bp

insert – target 400 M reads pairs per lane), leading to a total of 5.3 billon reads.

Sample Genotyping. Genotyping data [accession EGAS00001001895] were obtained for all

200 individuals from the EvoImmunoPop cohort based on both Illumina HumanOmni5-Quad

BeadChips and whole-exome sequencing with the Nextera Rapid Capture Expanded Exome kit

(18). The 3,782,260 SNPs obtained after stringent quality control were then used for imputation,

based on the 1,000 Genomes Project imputation reference panel (Phase 1 v3.2010/11/23) (49),

leading to a final set of 19,619,457 high-quality SNPs, of which 7,766,248 SNPs had a MAF $\geq$5%

in our cohort.

Processing of scRNA-seq data. Basic pre-processing of the sequencing data was performed

with CellRanger v3.0.2 (50), including the *mkfastq*, *count*, and *aggr* commands. Default

parameters and our combined human-IAV reference were used, and batch correction was

disabled in the *aggr* command. Cell-containing droplets (*n*=132,130) were traced back to

individual donors using two independent methods, Demuxlet and SoupOrCell, which capitalize

on genetic variation in the sequencing reads (20, 51). Barcodes with ambiguous and/or non-

concordant calls between the two programs were used to establish suitable QC metrics. We

found that barcodes deemed as *doublets* (i.e. the droplet contained two or more cells originating

from different donors) were more likely to be nearest-neighbors in a *knn*-graph with other

doublets than assigned *singlets*. We used this feature to identify droplets presumed to contain

two or more cells originating from the same donor; barcodes with > 5 *doublets* as nearest-

neighbors were excluded from further analysis (*SI Appendix,* **Fig. S1** *A and B*). Additionally,

droplets containing low-quality cells (i.e. damaged, dying) were excluded using the following

thresholds: < 1500 total counts, < 500 genes, or > 50% mitochondrial gene content (*SI*

Appendix, **Fig. S1***C*). This QC resulted in a final data set of 96,386 single cells.

Transcriptomes (i.e. counts) were adjusted for the presence of ambient RNA with SoupX,

(https://github.com/constantAmateur/SoupX/, accessed November 28, 2019) (21), using

estimated contamination fractions (per 10X library) from SoupOrCell (51). SoupX-adjusted

counts were normalized using pool-based size factors followed by deconvolution as

implemented in the scran R package (52). Feature selection was performed by (i) constructing a

mean-variance trend in the log-counts and retaining genes found to exhibit more variation than

expected assuming Poisson-distributed technical noise, as implemented in the makeTechTrend

and TrendVar functions from package scran (52), and (ii) selecting genes expressed in at least

25 cells (n=22,603). The first 10 PCs of the data were retained for data visualization and

clustering analyses. Graph-based clustering was performed by building the shared nearest-

neighbor graph with the buildSNNGraph function from scran (52) using a series of k values, and

cell clusters were defined with the igraph Walktrap algorithm (53). Similar clustering results were

obtained based on the knn-graphs generated using k=25, 50, 75, and 100, and k=25 was used

for all downstream analyses (**SI Appendix, Fig. S2A**). Cell types were predicted using SingleR

and the built-in BluePrintEncodeData reference (54). Based on the clustering and cell-type

predictions, we removed cells belonging to clusters associated with lymphoid cell types or low

QC metrics from downstream analyses (**SI Appendix, Fig. S2 B and C**).

Accounting for ambient RNA contamination in scRNA-Seq Data and assigning cell states.

Droplet-based scRNA-seq methods capture ambient mRNAs present in the cell suspension in

addition to cell specific mRNAs. To estimate which cells in our experiment were genuinely

expressing mRNAs for *CD14*, *FCGR3A (CD16)*, and those originating from the virus, we

implemented a two-step strategy utilizing the estimateNonExpressionCells function of the SoupX

package (21). This function estimates whether each cell contains significantly more counts of a

provided gene-set than would be expected under a Poisson model, given the estimated ambient

RNA from its library of origin and the maximum contamination fraction. First, we used the viral

genes to estimate the true maximum contamination fraction, based on the assumption that cells

from the non-infected state should only contain viral reads from ambient mRNA captured in their

droplets. To do so, we modified the estimateNonExpressionCells function to return p-values, and

performed the test on each of our 13 libraries with a range of maximum contamination values

from 1-50% (step of 1%) using the viral genes. We then computed FDR adjusted p-values for

each maximum contamination value on the 88,559 high-quality, single monocytes. The number

of non-simulated cells deemed to significantly express IAV transcripts (FDR<0.01) was used as

a proxy for false positives. In examining the relationship between this number and the number of

IAV-challenged cells found to significantly express viral transcripts at FDR<0.01 (**Fig. *1C***), we

found that a maximum contamination fraction of 10% resulted in a 1% false positive rate

(defined as the percentage of non-infected cells from T_2-T_8 that were deemed to significantly

express IAV transcripts). This parameter value was then used to correct for contamination from

ambient for all genes considered (*CD14*, *FCGR3A* and IAV transcripts).

Assigning cell states and investigating sources of variability in IAV levels. We used a

maximum contamination fraction of 10% to test for significant expression of IAV transcripts in

each cell (**Fig. 1 *C-E***). IAV-challenged cells that contained a significant amount of IAV transcripts

were considered as infected, while the others were deemed bystanders. To distinguish low from

high IAV-transcribing cells, a Gaussian mixture model was fitted to the total percentage of viral

mRNAs per cell across all infected cells, using the normalmixEM function from mixtools R

package with k=2 (55). Each cell was assigned to the cluster with the highest posterior

probability, and the cluster of cells with higher IAV content was annotated as high IAV-

transcribing.

Characterizing monocyte subsets and transcriptional profiles from scRNA-seq data.

Principal components analysis of 6,601 cells at T_0 was used to order monocytes along a

differentiation axis separating *CD14*[+] cells from *CD16*[+] cells. We then computed the average

percentage of classical and nonclassical monocytes obtained by FACS across the eight donors,

weighting each individual by the number of high-quality cells in the scRNA-seq data at T_0. Based

on these percentages (87.1% for classical and 7.6% for nonclassical), we annotated the

monocytes on each side of the differentiation axis as classical and nonclassical, respectively,

with the remaining 5.3% of monocytes being annotated as intermediates. Validity of our

approach was confirmed by correlating the proportion of monocytes assigned to each subset

across the eight donors, with the percentage of classical, intermediate and nonclassical

monocytes estimated by FACS.

Differential expression between subsets was assessed for the 4,859 genes expressed at an

average logCount > 0.1 in any of the 3 subsets. Specifically, Wilcoxon rank tests were

implemented in the scran package (52), using the findMarkers function and blocking on donor.

We considered genes to be differentially expressed (DE) between monocyte subsets when gene

expression was significant at an FDR≤1% and $\log_2$FC>0.2. The 848 genes that differed between

classical (CL) and nonclassical (NC) monocyte subsets were classified according to their

behavior in intermediate monocytes (INT). They were either deemed 'similar to classical' (DE

between INT and NC, but not between INT and CL), 'similar to nonclassical' (DE between INT and CL, but not between INT and NC), or 'intermediate' (all other cases).

At later time points, comparisons between $CD16^+$ and $CD16^-$ monocytes subsets were done based on 5,681 genes expressed with a normalized log count > 0.1 on average in either subset, in at least one condition and time point. For each subset, $\log_2$ fold change in gene expression relative to T_0 were correlated across times points. Differential expression between $CD16^+$ and $CD16^-$ cells was assessed with findMarkers (52), based on Wilcoxon rank tests and blocking on donors, time points and condition. Again, an FDR≤1% and $\log_2$FC>0.2 were required to define differentially expressed genes. To assess how $CD16^{+/-}$ status alters the infection of monocytes by IAV, we used logistic regression to model bystander/infected status as a function of $CD16^{+/-}$ status, while adjusting on donor, and time point (as factors).

Characterizing subset-specific responses to IAV challenge. To allow comparison between responses of $CD16^+$ and $CD16^-$ monocytes, 100 cells were subsampled from each subset and cell state, and at each time point. When subsampling, we ensured balanced representation of all donors across each monocyte subset and cell state, by using sampling weights that were inversely proportional to each donor representation in the original dataset. After sampling, a total of 6,669 genes with normalized log counts >0.1 on average in at least one group (cell-state x subset x time point) was selected for further analyses. For each monocyte subset, differences in expression between cell states (unexposed, bystander, infected) as well as between high- and low-IAV transcribing infected cells were performed using the findMarkers function from the scran package (52) and blocked on time-point. For each comparison, genes were considered to be differentially expressed between cell states when gene expression was significant at an FDR=1% (Wilcoxon rank tests) and the $\log_2$ fold change was > 0.2. In addition, for each comparison between cell states, we tested for differences in response between subsets using a linear model of the form:

$$(10)\ Expr_i \sim State_i + subset_i + State_i . subset_i$$

where $Expr_i$ is the expression of the gene being tested in cell i, $State_i$ is an indicator variable that distinguishes the two cell states being compared (e.g. unexposed and bystander), and $subset_i$ is an indicator variable that reflects the $CD16^{+/-}$ status of cell i. The 335 genes with significant interactions at a 1% FDR (for unexposed-bystander and unexposed-infected

comparisons) were clustered using the hclust R function with method 'Ward.D2'.

DynamicTreeCut algorithm (56) was used to identify eight major patterns of response to IAV.

Transcription factor enrichment analyses. To estimate Transcription Factor (TF) activity and

define TF-targets relationships, we ran the R SCENIC pipeline (29) on the expression matrix

(pre-normalization) on a random subsample of 4800 cells (100 cells from each donor at each

time point and each condition, pre-exclusion of dying and contaminant cells) with default

parameters. For each gene, motif-enrichment was considered for either *cis*-regulatory regions

located <10kb from the TSS (distal regulatory elements), or between 500 bp upstream and 100

bp downstream of the promoter (proximal regulatory elements). To do so, motif-enrichment

scores for all human genes (hg38 build, refseq_r80), were retrieved from

https://resources.aertslab.org/cistarget and used as input for the Rcistarget package (29).

Sets of high-confidence targets for the 113 TFs whose activity could be quantified by

SCENIC were then extracted and used for enrichment analysis. For each gene module, TF

enrichment was assessed using a Fisher's exact test with the 6,669 expressed genes as

background (**Dataset 3**). Resulting *p*-values were adjusted using a global Benjamini-Hochberg

correction for all eight modules and 113 TFs.

For each TF, with its targets enriched among one of the eight modules, TF activity inferred

by SCENIC was used to test for subset-specific activity using a linear model of the form:

$(11)\ TF_i \sim State_i +\ subset_i + State_i\ .\ subset_i$

where TF_i is the activity of the TF being tested in cell i, $State_i$ is an indicator variable that

distinguishes the two cell states being compared (e.g. unexposed and bystander), and $subset_i$ is

an indicator variable that reflects the *CD16*$^{+/-}$ status of cell i. Average TF activity was then

computed for each cell state, subset and time point, and correlated with gene expression of the

associated module, to assess the link between TF activation and the TF-target enriched

modules.

Association of the outcome of IAV infection using basal gene expression. For each of the

three monocyte subsets detected at basal state, Kruskall-wallis test was used to search for

genes whose expression levels significantly differ across donors. Within each monocyte subset,

we then computed the average expression of each gene for all eight donors and correlated it

with the percentage of infected cells at 4hpi. Genes that differed in expression between donors

(FDR≤1%), and passed a nominal *p*-value threshold of 0.01 for association with IAV levels in

any of the 3 subsets, were selected for downstream enrichment analyses. For genes nominally

correlated with viral mRNA levels, TF enrichment was assessed as previously using a Fisher's

exact test with all 4,859 genes expressed at T_0 as background (**Dataset S4**), and Benjamini-

Hochberg correction for all 113 TFs was applied.

4
5   **Gene Ontology enrichment analyses.** All Gene Ontology (GO) enrichment analyses were

6   performed with GOSeq package using default settings (57). Background gene sets consisted of

7   all genes that had average log-normalized expression value > 0.1 in at least one of the

8   groupings being examined, and are described in the text. Only enrichments significant at

9   FDR≤5% are reported.

10
11 **Pseudo-bulk estimates from scRNA-seq data.** Pseudo bulk estimates of IAV mRNA levels

12 were computed by measuring, for each donor and time point, the mean percentage of reads of

13 viral origin across all cells from the sample. At each time point, we then used a Pearson's

14 correlation test to compare pseudo-bulk estimates for the 8 donors with IAV mRNA levels

15 obtained in bulk data at 6hpi.

16
Monocyte subset characterization of EVOIMMUNOPOP samples via FACS. For 174 of the

200 EVOIMMUNOPOP donors, proportions of classical, intermediate and nonclassical

monocytes were determined based on a fraction of 10^5 CD14$^+$ positively-selected monocytes,

stained according to the manufacturer's instructions, with fluorescent APC-conjugated anti-CD14

and PE-conjugated anti-CD16 antibodies (Cat#130-091-243 and Cat #130-091-245,

respectively, Miltenyi Biotec). Samples were then analyzed on a MACSQuant Analyzer 10

benchtop flow cytometer (Miltenyi Biotec).

24
Quantification of canonical monocyte subsets in EVOIMMUNOPOP samples. FlowJo

v10.6.1 software (58) was used with the gating strategy depicted in ***SI Appendix*, Fig. S1** to

quantify monocyte subsets for 174 EVOIMMUNOPOP donors. Population-level differences in

proportion of canonical monocyte subsets were assessed using Wilcoxon Rank tests.

Correlation of the ratio of CD16$^+$ to CD16$^-$ cells with IAV mRNA levels was assessed using a

linear model of the form

$(1)\ IAV \sim ratio + Pop,$

where '*IAV*' are IAV mRNA levels, '$ratio$' is the percentage of CD16$^+$ monocytes (non

classical+intermediates) divided by the percentage of CD16$^-$ monocytes (classical), and '*Pop*' is

and indicator variable separating AFB from EUB individuals.

Analysis of bulk RNA-seq profiles from the EVOIMMUNOPOP cohort. A combined human-

IAV reference was generated by concatenation of the primary human genome assembly

(GRCh38) with the 8 segments of the human influenza A virus (IAV) A/USSR/90/1977(H1N1)

genome (accession numbers CY010372-CY010379). Comprehensive human gene annotation

was obtained from GENCODE (release 27) and merged with the 12 known transcripts of

A/USSR/90/1977(H1N1). RNA-seq reads (FASTQs) for all 970 samples that passed quality

control in our previous study (18) [accession EGAS00001001895] were mapped to the

combined reference with the STAR aligner (v.2.5.0a) (59) and assessed for quality with

QualiMap 'bamqc' and 'rnaseq' (60, 61). Expression of viral mRNAs was measured as the

percentage of uniquely-mapped reads aligning to the IAV genome. Reassuringly, the mean

percentage of RNA-seq reads among samples from the IAV-challenged condition was 5.86%

versus <0.01% in the other four conditions. Comparison of the percentage of IAV reads between

populations was done using a Wilcoxon rank test. StringTie (v.1.3.3) (62) was used to quantify

expression levels in transcripts per million mapped reads (TPM) for each annotated transcript.

Gene expression data were filtered to remove genes with little evidence of activation (mean

zTPM score < -3) (63) in any of the 5 conditions, and their quality was checked by principal

component analysis (PCA). As GC content, $5'/3'$ bias, date of the experiment and library batch

were previously determined to be the strongest confounding factors on transcript expression

(18), we corrected the data for these factors. First, we adjusted the data for GC content and $5'/3'$

bias using linear models. Then, we imputed missing values by k-nearest neighbor imputation

and adjusted for experiment date and library batch by sequentially running ComBat (64) for each

batch effect, with condition and population as covariates. After batch effect correction, only IAV-

stimulated samples were kept for downstream analyses.

Cell states deconvolution from bulk RNA sequencing. To assess the percentage of total

transcripts that originate from each cell state across the 199 IAV-challenged samples, we pooled

cells from T_6 into 3 groups, based on their assigned cell-state (bystander, infected: high and low

IAV-transcribing) and to which we added a 4^{th} group containing all singlets that were either (i)

assigned to cluster numbers 3, 8, 10, and 11 (dying cells) or (ii) discarded based on their high

mitochondrial content or low read counts (dead cells). We then estimated pseudo-bulk profiles

for each group by summing UMIs across all cells and computing the number of UMIs associated

to each gene per million of sequenced UMIs. TPM profiles obtained from bulk data were then

normalized to improve comparison with pseudo-bulk. Specifically, we first computed a global

pseudo-bulk profile of the entire single cell dataset as the average of the pseudo bulk profiles from the 4 cell states (bystander, infected: high and low IAV-transcribing, or dying/dead), weighted by the percentage of UMIs they contribute to the overall pool of cells. To account for the difference in how gene expression is quantified between the two methods (3' end counts for scRNA-seq and full-length gene coverage for bulk RNA-seq), we computed for each gene i a normalization factor s_i given by

(5) $s_i = \log(\overline{TPM_i}) - \log(PB_i)$

where PB_i is the number of UMI per million for gene *i* in the global pseudo-bulk profile, and $\overline{TPM_i}$ is the average expression of the gene *i* in the 199 IAV-stimulated samples from the bulk RNA-seq data. For each gene, s_i was then subtracted from the log transformed TPM to yield a normalized TPM profiles. We next applied DeconRNAseq (65) to the normalized log TPM profiles from all individuals, using the log-transformed pseudo bulk profiles from the 4 cell states as a basis for deconvolution. Quality of the deconvolution was assessed using leave-one-out cross validation, based on the eight individuals for whom we had scRNA-seq data. Specifically, for each of these eight individuals, bulk mRNAs were decomposed using pseudo-bulk profiles recomputed based on the seven other individuals. The resulting proportions were then compared with the percentage of UMIs that originate in each cell-state in the scRNA-seq to assess the quality of the deconvolution. Comparisons between populations were performed using Wilcoxon rank tests.

The effect of the percentage of infected cells and percentage of high IAV-transcribing cells among infected cells on the total IAV mRNA levels were assessed by modeling

(6) *IAV ~ INF + POP*

and

(7) *IAV ~ HI + POP*

where *IAV* are the IAV mRNA levels across the 199 bulk mRNA samples, *INF* and *HI* are respectively the percentage of infected cells and the percentage of high IAV-transcribing cells among infected cells that we estimated from the deconvolution, and *POP* is a factor variable reflection the population (EUB or AFB). The fraction η of population differences attributable to difference in rate of infection was estimating by comparing model (6) with model (8) below

(8) *IAV ~ POP*

and computing $\eta = 100 \times \left(1 - \frac{\beta_6}{\beta_8}\right)$, with β_i the effect of population on IAV levels in model (i). To assess how the contribution of the percentage of high IAV transcribing cells to total IAV mRNA levels differed between populations, we used a linear model of the form

(9) *IAV ~ HI + POP + HI:POP*

1 and tested for significant effect of the interaction term *HI:POP* on *IAV* mRNA levels.

Data availability

The bulk RNA-seq data used in this study are available at the European Genome Phenome archive under accession number [EGAS00001001895]. The single cell RNA-seq data generated during this study are available at the European Genome Phenome archive under accession number [EGAS00001005000]. Code generated as part of this study is available on Github (https://github.com/h-e-g/PopDiff_MonocyteIAV).

Acknowledgments

This work was supported by the *Institut Pasteur*, the *Collège de France*, the French Government's Investissement d'Avenir program, *Laboratoires d'Excellence* "Integrative Biology of Emerging Infectious Diseases" (ANR-10- LABX-62-IBEID) and "Milieu Intérieur" (ANR-10- LABX-69-01), the *Fondation de France* (n°00106080), and the *Fondation pour la Recherche Médicale* (Equipe FRM DEQ20180339214). M.B.O was supported by a *European Molecular Biology Organization* long-term fellowship (ALTF 229-2017). We also wish to thank Tzachi Hagai and John Marioni for input and feedback, and Sylvie van Der Werf and Vincent Enouf for providing resources and protocols relating to influenza viruses.

References

1. V. A. Ryabkova, L. P. Churilov, Y. Shoenfeld, Influenza infection, SARS, MERS and COVID-19: Cytokine storm - The common denominator and the lessons to be learned. *Clinical immunology (Orlando, Fla.)* **223**, 108652 (2021).
2. F. Krammer *et al.*, Influenza. *Nat Rev Dis Primers* **4**, 3 (2018).
3. Q. Zhang *et al.*, Life-Threatening COVID-19: Defective Interferons Unleash Excessive Inflammation. *Med (N Y)* **1**, 14-20 (2020).
4. A. A. Stegelmeier *et al.*, Myeloid Cells during Viral Infections and Inflammation. *Viruses* **11** (2019).
5. D. C. Fajgenbaum, C. H. June, Cytokine Storm. *N Engl J Med* **383**, 2255-2273 (2020).
6. C. Guo *et al.*, Single-cell analysis of two severe COVID-19 patients reveals a monocyte-associated and tocilizumab-responding cytokine storm. *Nature communications* **11**, 3924 (2020).
7. R. Alon *et al.*, Leukocyte trafficking to the lungs and beyond: lessons from influenza for COVID-19. *Nat Rev Immunol* **21**, 49-64 (2021).
8. L. Ziegler-Heitbrock *et al.*, Nomenclature of monocytes and dendritic cells in blood. *Blood* **116**, e74-80 (2010).
9. F. Geissmann, S. Jung, D. R. Littman, Blood monocytes consist of two principal subsets with distinct migratory properties. *Immunity* **19**, 71-82 (2003).
10. C. Auffray *et al.*, Monitoring of blood vessels and tissues by a population of monocytes with patrolling behavior. *Science* **317**, 666-670 (2007).
11. L. Ziegler-Heitbrock, The CD14+ CD16+ blood monocytes: their role in infection and inflammation. *J Leukoc Biol* **81**, 584-592 (2007).
12. M. A. Hoeve, A. A. Nash, D. Jackson, R. E. Randall, I. Dransfield, Influenza virus A infection of human monocyte and macrophage subpopulations reveals increased susceptibility associated with cell differentiation. *PLOS ONE* **7**, e29443 (2012).
13. W. Hou *et al.*, Viral infection triggers rapid differentiation of human blood monocytes into dendritic cells. *Blood* **119**, 3128-3131 (2012).
14. B. Piasecka *et al.*, Distinctive roles of age, sex, and genetics in shaping transcriptional variation of human immune responses to microbial challenges. *Proceedings of the National Academy of Sciences of the United States of America* **115**, E488-E497 (2018).
15. P. Brodin *et al.*, Variation in the human immune system is largely driven by non-heritable influences. *Cell* **160**, 37-47 (2015).
16. W. J. Astle *et al.*, The Allelic Landscape of Human Blood Cell Trait Variation and Links to Common Complex Disease. *Cell* **167**, 1415-1429 e1419 (2016).
17. E. Patin *et al.*, Natural variation in the parameters of innate immune cells is preferentially driven by genetic factors. *Nat Immunol* **19**, 302-314 (2018).
18. H. Quach *et al.*, Genetic Adaptation and Neandertal Admixture Shaped the Immune System of Human Populations. *Cell* **167**, 643-656 e617 (2016).
19. K. G. Ouwens *et al.*, A characterization of cis- and trans-heritability of RNA-Seq-based gene expression. *Eur J Hum Genet* **28**, 253-263 (2020).
20. H. M. Kang *et al.*, Multiplexed droplet single-cell RNA-sequencing using natural genetic variation. *Nature biotechnology* **36**, 89-94 (2018).
21. M. D. Young, S. Behjati, SoupX removes ambient RNA contamination from droplet-based single-cell RNA sequencing data. *Gigascience* **9** (2020).
22. K. L. Wong *et al.*, Gene expression profiling reveals the defining features of the classical, intermediate, and nonclassical human monocyte subsets. *Blood* **118**, e16-31 (2011).
23. V. Segura *et al.*, In-Depth Proteomic Characterization of Classical and Non-Classical Monocyte Subsets. *Proteomes* **6** (2018).

24. C. Schmidl *et al.*, Transcription and enhancer profiling in human monocyte subsets. *Blood* **123**, e90-99 (2014).

25. A. Bercovich-Kinori *et al.*, A systematic view on influenza induced host shutoff. *eLife* **5** (2016).

26. H. M. Machkovech, J. D. Bloom, A. R. Subramaniam, Comprehensive profiling of translation initiation in influenza virus infected cells. *PLoS pathogens* **15**, e1007518 (2019).

27. S. Li, Regulation of Ribosomal Proteins on Viral Infection. *Cells* **8** (2019).

28. J. Cros *et al.*, Human CD14dim monocytes patrol and sense nucleic acids and viruses via TLR7 and TLR8 receptors. *Immunity* **33**, 375-386 (2010).

29. S. Aibar *et al.*, SCENIC: single-cell regulatory network inference and clustering. *Nature methods* **14**, 1083-1086 (2017).

30. E. K. Allen *et al.*, SNP-mediated disruption of CTCF binding at the *IFITM3* promoter is associated with risk of severe influenza in humans. *Nature Medicine* **23**, 975 (2017).

31. Q. Zhang, Human genetics of life-threatening influenza pneumonitis. *Human genetics* **139**, 941-948 (2020).

32. M. J. Ciancanelli *et al.*, Infectious disease. Life-threatening influenza and impaired interferon amplification in human IRF7 deficiency. *Science* **348**, 448-453 (2015).

33. B. Panthu *et al.*, The NS1 Protein from Influenza Virus Stimulates Translation Initiation by Enhancing Ribosome Recruitment to mRNAs. *J Mol Biol* **429**, 3334-3352 (2017).

34. C. Wang *et al.*, Cell-to-Cell Variation in Defective Virus Expression and Effects on Host Responses during Influenza Virus Infection. *mBio* **11** (2020).

35. Y. Steuerman *et al.*, Dissection of Influenza Infection In Vivo by Single-Cell RNA Sequencing. *Cell systems* **6**, 679-691.e674 (2018).

36. A. B. Russell, C. Trapnell, J. D. Bloom, Extreme heterogeneity of influenza virus infection in single cells. *eLife* **7** (2018).

37. J. Sun *et al.*, Single cell heterogeneity in influenza A virus gene expression shapes the innate antiviral response to infection. *PLoS pathogens* **16**, e1008671 (2020).

38. I. Ramos *et al.*, Innate Immune Response to Influenza Virus at Single-Cell Resolution in Human Epithelial Cells Revealed Paracrine Induction of Interferon Lambda 1. *Journal of virology* **93** (2019).

39. A. B. Russell, E. Elshina, J. R. Kowalsky, A. J. W. Te Velthuis, J. D. Bloom, Single-Cell Virus Sequencing of Influenza Infections That Trigger Innate Immunity. *Journal of virology* **93** (2019).

40. E. Kudo *et al.*, Low ambient humidity impairs barrier function and innate resistance against influenza infection. *Proceedings of the National Academy of Sciences of the United States of America* **116**, 10905-10910 (2019).

41. Y. Cao *et al.*, Single-cell analysis of upper airway cells reveals host-viral dynamics in influenza infected adults. *bioRxiv* 10.1101/2020.04.15.042978, 2020.2004.2015.042978 (2020).

42. L. J. Appleby *et al.*, Sources of heterogeneity in human monocyte subsets. *Immunol Lett* **152**, 32-41 (2013).

43. R. Chandrasekhar *et al.*, Social determinants of influenza hospitalization in the United States. *Influenza Other Respir Viruses* **11**, 479-488 (2017).

44. J. L. Hadler *et al.*, Influenza-Related Hospitalizations and Poverty Levels - United States, 2010-2012. *MMWR Morb Mortal Wkly Rep* **65**, 101-105 (2016).

45. N. J. Roberts, Jr., Diverse and Unexpected Roles of Human Monocytes/Macrophages in the Immune Response to Influenza Virus. *Viruses* **12** (2020).

46. S. L. Cole *et al.*, M1-like monocytes are a major immunological determinant of severity in previously healthy adults with life-threatening influenza. *JCI Insight* **2**, e91868 (2017).

47. Y. H. Zhou *et al.*, Pathogenic T cells and inflammatory monocytes incite inflammatory storm in severe COVID-19 patients. *Natl Sci Rev* 10.1093/nsr/nwaa041, nwaa041 (2020).

48. 10xGenomics, Chromium Single Cell 3' Reagent Kits User Guide (v3 Chemistry). (2020).

49. C. Genomes Project *et al.*, An integrated map of genetic variation from 1,092 human genomes. *Nature* **491**, 56-65 (2012).

50. G. X. Zheng *et al.*, Massively parallel digital transcriptional profiling of single cells. *Nature communications* **8**, 14049 (2017).

51. H. Heaton *et al.*, Souporcell: robust clustering of single-cell RNA-seq data by genotype without reference genotypes. *Nature methods* **17**, 615-620 (2020).

52. A. T. Lun, D. J. McCarthy, J. C. Marioni, A step-by-step workflow for low-level analysis of single-cell RNA-seq data with Bioconductor. *F1000Res* **5**, 2122 (2016).

53. P. Pons, M. Latapy (2005) Computing Communities in Large Networks Using Random Walks. in *Computer and Information Sciences - ISCIS 2005*, eds p. Yolum, T. Güngör, F. Gürgen, C. Özturan (Springer Berlin Heidelberg, Berlin, Heidelberg), pp 284-293.

54. D. Aran *et al.*, Reference-based analysis of lung single-cell sequencing reveals a transitional profibrotic macrophage. *Nat Immunol* **20**, 163-172 (2019).

55. T. Benaglia, D. Chauveau, D. R. Hunter, D. S. Young, mixtools: An R Package for Analyzing Mixture Models. *Journal of Statistical Software; Vol 1, Issue 6 (2010)* (2009).

56. P. Langfelder, B. Zhang, S. Horvath, Defining clusters from a hierarchical cluster tree: the Dynamic Tree Cut package for R. *Bioinformatics* **24**, 719-720 (2008).

57. M. D. Young, M. J. Wakefield, G. K. Smyth, A. Oshlack, Gene ontology analysis for RNA-seq: accounting for selection bias. *Genome Biol* **11**, R14 (2010).

58. D. a. C. Becton (2019) FlowJo™ Software (Ashland).

59. A. Dobin *et al.*, STAR: ultrafast universal RNA-seq aligner. *Bioinformatics* **29**, 15-21 (2013).

60. F. García-Alcalde *et al.*, Qualimap: evaluating next-generation sequencing alignment data. *Bioinformatics* **28**, 2678-2679 (2012).

61. K. Okonechnikov, A. Conesa, F. García-Alcalde, Qualimap 2: advanced multi-sample quality control for high-throughput sequencing data. *Bioinformatics* **32**, 292-294 (2016).

62. M. Pertea *et al.*, StringTie enables improved reconstruction of a transcriptome from RNA-seq reads. *Nature biotechnology* **33**, 290-295 (2015).

63. T. Hart, H. K. Komori, S. LaMere, K. Podshivalova, D. R. Salomon, Finding the active genes in deep RNA-seq gene expression studies. *BMC genomics* **14**, 778 (2013).

64. W. E. Johnson, C. Li, A. Rabinovic, Adjusting batch effects in microarray expression data using empirical Bayes methods. *Biostatistics (Oxford, England)* **8**, 118-127 (2007).

65. T. Gong, J. D. Szustakowski, DeconRNASeq: a statistical framework for deconvolution of heterogeneous tissue samples based on mRNA-Seq data. *Bioinformatics* **29**, 1083-1085 (2013).

Fig. 1. Single-Cell RNA-Sequencing of 88,559 Monocytes and Their Derived Cells. (*A*) Post-QC tSNE colored by unsupervised graph-based clusters. (*B*) Post-QC tSNE colored by *FCGR3A* (*CD16*) $\log_2$ normalized counts (top), or percentage of viral mRNAs (bottom). (*C*) Determination of the maximum contamination fraction by ambient RNA. The number of non-infected cells deemed to significantly express IAV transcripts (presumed false positives) versus the number of IAV-challenged cells deemed to significantly express IAV transcripts across a range of maximum contamination fractions from 1-50% (color bar). Dotted grey line is drawn at 1% on the x-axis. A maximum contamination fraction of 10% results in 1% of non-infected cells being classified as infected (false positive proxy), and half of IAV-challenged cells showing evidence of viral transcription. (*D*) Distribution of counts of viral origin across all donors, from T_2 to T_8. Cells are shown separately for non-infected (top) and IAV-challenged (bottom) conditions. Fill color reflects the cell state assignments. Note that the threshold used to define infected cells is dependent on the number of viral mRNAs in the ambient pool, and varies across libraries. (*E*) Post-QC tSNE stratified by time point. For each time point, cells are colored according to their $CD16^{+/-}$ status (see key) and their assigned cell state (same as depicted in *D*). For each condition and time point, stacked bar charts below the tSNE represent the relative proportions of the various cell states and subsets.

1
2
Fig. 2. Gradient of mRNA Expression from Ribosomal and IFN-Stimulated Genes
Separates Bystander and Infected Cells. (***A***) Transcriptional responses of cells upon IAV-
challenge (T_2-T_8) highlight the interplay between IFN response (GO:0034340), ribosomal
(GO:0005840), and mitochondrial (GO:0005743) genes. The log_2FC change in gene expression
between unexposed and bystander cells is plotted on the x-axis, while the log_2FC change in
gene expression between unexposed and infected cells is plotted on the y-axis. Values are
plotted based on a meta-analysis across time points and subsets, based on a subsampled data
set with balanced representation of all donors. (***B***) The interplay between IFN response
(GO:0034340), ribosomal (GO:0005840), and mitochondrial (GO:0005743) genes among cells
exposed to IAV. The log_2FC change in gene expression between low IAV-transcribing infected
and bystander cells is plotted on the x-axis, while the log_2FC change in gene expression
between low IAV-transcribing infected and high IAV-transcribing infected cells is plotted on the y-
axis. Values are plotted based on a meta-analysis across monocyte subsets at T_4. (***C***) mRNA
expression levels of representative IFN-stimulated (*MX1*) and ribosomal (*RPL34*) genes across
the subsampled dataset. Colors reflect the cell state and subset assignment depicted in **Fig. 1 *D***
and ***E***.
19
20

Fig. 3. IRFs and STATs Have a Central Role in the Subset-Specific Responses to IAV Infection. (**A**) Heatmap of scaled gene expression from 335 genes displaying a subset-specific response to infection challenge. Genes are grouped into 8 modules based on hierarchical clustering of their expression patterns. Representative genes from each module are labelled. (**B**) Mean expression over time of *IL6* and *TNF*, across the different monocyte subsets and cell states. (**C**) Network of transcription factors (round nodes) associated with each gene expression module (square nodes). Transcription factor nodes are colored according to the number of modules they are associated with. Black lines represent enrichments of the module in TF targets, while red lines represent depletions.

1
2
3
Fig. 4. Basal IRF/STAT-Induced Transcriptional Network Underlies Inter-Individual
Differences in Monocyte Susceptibility and IAV levels. (***A***) Pseudo-bulk estimates of the
percentage of counts of viral origin in IAV-challenged condition (T_2-T_8). Donors are colored
based on the rank of these pseudo-bulk estimates at the peak of viral transcription, T_4, from that
with highest observed viral mRNA level (D1) to that of the lowest (D8). (***B***) Proportions of cell
states from the IAV-challenged condition at T_2, T_4, T_6, and T_8, in the eight donors. X-axis is
ordered by decreasing viral mRNA levels found at T_4 (D1-D8). (***C***) Log normalized expression
values of *IFITM3* across all cells, stratified by canonical monocyte subsets, and separated by
donor. Colors reflect the different donors depicted in ***A***. For each donor and monocyte subset,
the violin plots show the full distribution of *IFITM3* expression across individual cells and
boxplots highlight the median and interquartile range. (***D***) Enrichment of SCENIC-predicted
targets among the 118 genes whose basal expression at T_0 correlates with the percentage of
infected cells at later time points (odds ratio and 95% confidence interval). Red line designates
an odds ratio equal to 1. Only TFs significantly enriched among the 118 candidate genes are
shown (FDR < 0.05). Abbreviations: classical (CL), intermediate (INT), and nonclassical (NC).
19
20

1
2
Fig. 5. African-Ancestry Individuals Display Increased Number of *CD16*+ Cells and Lower
Susceptibility to IAV Infection. (*A*) Variation in the number of classical (CD14++CD16-),
intermediate (CD14++CD16+), and nonclassical (CD14+CD16++) monocytes across African- and
European- ancestry individuals following CD14+ selection from PBMCs (n_{AFB}=89, n_{EUB}=85).
Colors reflect population (AFB in red and EUB in blue). All three subsets are significantly
different between populations (*p*-value<0.01). (*B*) Inter- and intra-population variation in the
percentage of RNA-seq reads mapping to the IAV genome (n_{AFB}=100, n_{EUB}=99). Colors reflect
population (AFB in red and EUB in blue). The percentage of RNA-seq reads mapping to the IAV
genome is significantly higher in European-ancestry individuals relative to African-ancestry
individuals (*p*-value=5.3×10⁻⁸). (*C*) Inter- and intra-population variation in viral mRNA expression
at 6hpi (n_{AFB}=100, n_{EUB}=99). Expression levels for each of the 10 primary transcripts of IAV are
plotted. Colors reflect population (AFB in red and EUB in blue). All IAV transcripts are
significantly higher in European-ancestry individuals on average (*p*-value<0.001). (*D*) Estimated
distribution of the percentage of cells from each cell state in the bulk RNA-seq data (n_{AFB} =100,
n_{EUB}=99). Fill colors reflect cell state assignments, while outlines of boxplots reflect population
(AFB in red and EUB in blue). (*E*) Distribution of the percentage of high IAV-transcribers among
infected cells, stratified by population. One individual with no infected cell was excluded
(n_{AFB}=99, n_{EUB}=99). (*F*) Percentage of RNA-seq reads of viral origin as a function of the
estimated proportion of infected cells (n_{AFB}=100, n_{EUB}=99), colored by population (AFB in red and
EUB in blue). *A and C:* Outlier points are not displayed. Abbreviations: African-ancestry
individuals from Belgium (AFB), European-ancestry individuals from Belgium (EUB), transcripts
per million (TPM), influenza A virus (IAV), mean fluorescent intensity (MFI), classical (CL),
intermediate (INT), and nonclassical (NC). * *p*-value < 0.01; ** *p*-value < 0.001; *** *p*-value <
0.0001