Joint species distributions reveal the combined effects of host

plants, abiotic factors and species competition as drivers of

species abundances in fruit flies

Benoit Facon[1,*], Abir Hafsi[2,*], Maud Charlery de la Masselière[2], Stéphane Robin[3], François

Massol[4], Maxime Dubart[5], Julien Chiquet[3], Enric Frago[6], Frédéric Chiroleu[2], Pierre-François

Duyck[2,7] & Virginie Ravigné[2]

[1] INRAE, UMR PVBMT, F-97410 Saint Pierre, France

[2] CIRAD, UMR PVBMT, F-97410 Saint Pierre, France

[3] Laboratoire MMIP – UMR INRA 518/AgroParisTech, Paris, France

[4] Univ. Lille, CNRS, Inserm, CHU Lille, Institut Pasteur de Lille, U1019 - UMR 9017 - CIIL -

Center for Infection and Immunity of Lille, F-59000 Lille, France

[5] Univ. Lille, CNRS, UMR 8198 – Evo-Eco-Paleo, F-59000 Lille, France

[6] CIRAD, UMR CBGP, France

[7] IAC, Equipe ARBOREAL, BP 98857, Nouméa, Nouvelle-Calédonie

Short title: Joint distributions of interacting fruit flies

Keywords:

Type of article: Letters

* Co-first authors

Corresponding author: benoit.facon@inrae.fr

Abstract

The relative importance of ecological factors and species interactions for phytophagous insect

species distributions has long been a controversial issue. Using field abundances of eight sympatric

Tephritid fruit flies on 21 host plants, we inferred flies' realized niches using joint species distribution

modelling and network inference, on the community as a whole and separately on three groups of

host plants. These inferences were then confronted to flies' fundamental niches estimated through

laboratory-measured fitnesses on host plants. Species abundances were mainly determined by host

plants followed by climatic factors, with a minor role for competition between species sharing host

plants. The relative importance of these factors mildly changed when we focused on particular host

plant groups. Despite overlapping fundamental niches, specialists and generalists had almost distinct

realized niches, with possible competitive exclusion of generalists by specialists on Cucurbitaceae

and different assembly rules: specialists were mainly influenced by their adaptation to host plants

while generalist abundances varied regardless of their fundamental host use.

Introduction

The search for fundamental processes underlying species distributions is among the oldest challenges in ecology (Diamond 1975, Gotelli and Graves 1996). Understanding assembly processes could also be crucial for coping with global changes and habitat loss currently affecting both abiotic conditions and species distributions (Adler and HilleRisLambers 2008). Species distributions can be determined by several factors such as environmental filtering, interspecific interactions, regional species pool and dispersal (Müller et al. 2011, D'Amen et al. 2018, Nakadai et al. 2018, Jabot et al. 2020). Despite decades of research, estimating the relative importance of these processes on species distributions has proven particularly complex (Pollock et al. 2014, Zurell et al. 2018). Most of these processes imprint species distributions in a scale-dependent manner (Meynard et al. 2013). For instance, abiotic factors are generally thought to determine large-scale species ranges, whereas interspecific interactions would influence species distributions at smaller spatial scales (Heikkinen et al. 2007, Thuiller et al. 2015, but see Gotelli et al 2010 and Araújo and Rozenfeld 2014).

Phytophagous insects are among the most diverse and abundant groups of terrestrial animals and a major component of ecosystems due to their tight interaction with primary producers, and their sometimes important economic impacts and invasive potential (Roy et al. 2015). Knowledge of the main determinants of insect occurrence on particular plant species and their potential to colonize and persist in a given area is however still limited. In particular, the importance of interspecific competition in structuring phytophagous insect communities has been a controversial issue (Kaplan and Denno 2007). Many experimental studies conclude that interspecific competition plays a primary role (Denno et al. 1995, Kaplan & Denno 2007), but the consistent absence of negative co-occurrence patterns in natural phytophagous insect communities suggests otherwise (Tack et al. 2009, Brazeau & Schamp 2019). This apparent discrepancy could result from the regulation of phytophagous insect populations below competitive levels through shared predators or parasites (Hairston et al. 1960). Phytophagous insects would also rarely overexploit their hosts, leaving sufficient plant material for competition with other species to be mild (Kaplan & Denno 2007). Ecological differences between

species could lower the intensity of competition (Stewart et al. 2015). Lastly, some phytophagous arthropods could benefit from previous attack of the host plant by other species (Godinho et al. 2016). The importance of competition relative to environmental characteristics in shaping phytophagous insect distributions thus remains an open question and demands appropriate testing (Augustyn et al. 2016, Nakadai et al. 2018).

A way to identify whether biotic interactions affect species distributions consists in analyzing patterns of co-occurrence. Species interactions are expected to affect species occurrence, e.g., competition should cause checkerboard patterns of occurrence. However, species co-occurrences also result from common or diverging species dependence on confounding environmental factors. Species that share the same abiotic niche will frequently co-occur without necessarily interacting (Wisz et al. 2013, Blanchet et al. 2020). Conversely, negative co-occurrence patterns may simply result from diverging ecological requirements. As a consequence, estimating the effect of species interactions on species distributions first requires properly characterizing species abundance's responses to environmental variables (Pollock et al. 2014), which is the object of species distribution modeling (SDM) approaches (Elith & Leathwick 2009). A particular class of SDM approaches, joint species distribution models (JSDM) attempts to infer the relationships between species abundances and environmental variables, explicitly accounting for the interdependence of species distributions using multivariate regression methods (Pollock et al. 2014). In addition to estimates of the effect of habitat filtering on species distributions, these approaches provide residual covariances between species abundances, i.e., covariances not explained by environmental factors. Residual covariances result from species interactions and a diversity of other factors, as e.g., missing covariates, so that there is no simple relationship between species interactions and residual covariances (Zurell et al. 2018).

To further track species interactions, a growing body of literature pleads for using independent knowledge of species traits in species distribution modelling (Lavorel et al. 1997, Kraft et al. 2008, Poisot et al. 2015), which is still seldom done. Explicitly comparing estimates of fundamental niche,

i.e., measures of fitness in controlled conditions and in absence of species interactions, to estimates

of realized niche, i.e., inferred species abundances' responses to environmental factors, could shed

new light on the gap between fundamental and realized niches and the importance of species

interactions in shaping species distributions.

In the case of phytophagous insects, an obvious feature of the environment to account for is host

plant identity. Host plants can be treated as any environmental cofactor, and their effects on species

abundances can be inferred directly from adequate abundance data (Ferrier and Guisan 2006). Host

plants impose a specific challenge because modelling a phytophagous community as a whole relies

on the assumption that species abundances' interdependence does not depend on host plants. Since

intraguild interactions mostly occur in/on plant organs, they may be modulated by plant species

identity, with possible consequences for species occurrence patterns (Ulrich et al. 2017). Analyzing

competition patterns on different plants or groups of plants could therefore allow detecting the role

of host plants in shaping species co-occurrences.

Here we aimed at disentangling the roles of host plant species, abiotic factors, and interspecific

interactions on the distributions of eight fruit fly species (Tephritidae) occurring in sympatry on a

diversity of host plants and in highly variable abiotic conditions. The study system, which comprises

four generalist species, three specialists of *Cucurbitaceae*, and one specialist of *Solanaceae*, presents

key advantages to tackle community assembly questions. First, these eight species occupy a small

island in South-western Indian Ocean (Réunion, 2512 km^2) where they are considered the main

actors in the guild of fruit-eating phytophagous arthropods (Quilici and Jeuffrault, 2001). Second, the

local environment is characterized by important variability in elevation (from 0 to 3000m), climatic

conditions, land use, and plant distributions (Duyck et al. 2006a). Observational and experimental

studies have suggested that climatic factors could influence local Tephritid distributions (Duyck et al.

2004). Climatic factors were even found more influential than host plant diversity in allowing

coexistence in an analysis of the distributions of the four generalist species on four host plants

(Duyck et al. 2008). Lastly, competition between the eight species has repeatedly been advocated to

shape this community. First, host-use strategies largely overlap, opening possibilities of competition (Quilici & Jeuffrault 2001, Duyck et al. 2008). Second, the arrival of one generalist species on the island has constrained the host ranges of some resident species, without complete exclusion (Charlery de la Masselière et al. 2017a). Third, larval competition experiments in a subset of plant species and abiotic conditions have evidenced hierarchical competition interactions among the four generalist species (Duyck et al. 2006b).

Here we confronted a long-term field dataset describing abundances of the eight fly species on 21 host plants with laboratory measures of fundamental host use obtained for seven of the fly species on the same plants. We first modelled joint species distributions using Poisson-LogNormal (PLN) modelling (Chiquet et al. 2019) and conducted model selection among various combinations of host plant species and ecological covariates (representing temperature, rainfall, elevation, land use and date). Residual correlations estimated under the selected model were further dissected to identify significant traces of unexplained co-occurences. Second, we assessed whether knowledge on fundamental host use was sufficient to explain field species abundances by accounting for host plant species either directly or through estimates of female preference and larval performance in laboratory conditions. Finally, we tested for a potential dependence of community structuring factors on host plants by replicating the analyses on three subsets of plants: *Cucurbitaceae*, *Solanaceae* and the other plants.

Materials and Methods

Species abundance table

Field campaigns were conducted over a period of 18 years, between 1991 and 2009, to identify potential host plants for Tephritidae on Réunion on the whole island including orchards, gardens, and natural areas. These surveys were assembled in a previous study (Charlery de la Masselière et al. 2017a), and used here as our species abundance table. Each observation corresponds to the number of individual flies of the eight species recorded from a set of fruits sampled in one location at a

specific date. For each sample, the collected fruits were counted and weighted, before being stored until adult fly emergence. To avoid keeping samples that could have suffered from a transportation or storage issue, only samples with at least one individual fly were kept. Among these we further selected samples with GPS coordinates, and belonging to one of the 21 host plants characterized in the laboratory (see below). Of the 12872 initial samples, we therefore kept 4918 samples and a total of 97351 individual flies. Samples covered 104 field sessions all year round over the study period (Tables S1 and S2) and originated from 380 sites well distributed over the whole island (Figure 1A). Additional details on sample collection can be found in Appendix S1.

Ecological covariables

The GPS coordinates of each sample were used to retrieve ecological and climatic characteristics from GIS information available on the CIRAD Agricultural Web Atlas for Research (AWARE, https://aware.cirad.fr). Each sample was associated with a month and a year, a land use category, an elevation, three pluviometry descriptors (minimal rainfall in the 20% most humid years, minimal rainfall in the 20% driest years, and median annual rainfall between 1986 and 2016) and three temperature variables (minimal, mean and maximal annual temperature over the 1987-2017 period) (Figure 1A, Appendix S1). To account for the correlations between some of the variables, a FAMD (factorial analysis of mixed data) was conducted on all 10 variables using *FactoMineR* (Lê et al. 2008). Ten uncorrelated dimensions were obtained and subsequently used as ecological covariables in the following analyses (see Figure S2 for details on the FAMD).

Species traits

For all species but *D. ciliatus*, fundamental host use, i.e., fly fitness on host plants in optimal abiotic conditions and in absence of antagonists, was characterized using four traits describing larval performances and female preferences for 21 plant species. Female preferences were the numbers of eggs laid by females during 24h on each of the 21 fruit species in the 'no-choice' experiment of Charlery de la Masselière et al. (2017b). Larval performances (survival probability until maturity *s*,

development time *T*, and pupal weight *w*) were obtained from Hafsi et al. (2016) for 17 plant species,

and in the current study for *Coffea arabica*, *Solanum mauritianum*, *Syzygium jambos*, and *S.*

samarangense, using the same methods. The three larval performance traits were combined into a

single performance trait using the formula:

$$Perf = s \times w/T.$$

Both preference and performance traits were log-transformed before being included as covariates in

statistical models.

8
9   Figure 1: Characteristics of the species abundance dataset. A) Sampling sites in Réunion. Dot
    colors refer to the family of the sampled plant (Solanaceae n=259, Cucurbitaceae n=2347, other
    families n=2285 samples). Elevation, represented by grey isoclines, ranges from 0 along the coast,
10 to >3000 m, in the center of the island, and strongly correlates with annual temperature.
    Pluviometry (here median rainfall over 1986-2016) is represented by green isohyets. The island is
11 separated into two contrasted rainfall regimes: very humid all year round in the east, and drier,
12 especially during winter, in the west. B) Fly species abundances on the 21 studied plant species.

Statistical analysis

Datasets

To account for the possibility that determinants of species distributions depend on host plant identity, we replicated all analyses on the full 21-plants dataset, and on the following three sub-datasets: (i) Cucurbitaceae only (3 fly species x 6 plant species, 2347 samples), (ii) Solanaceae only (3 fly species x 3 plant species, 259 samples), (iii) other plant families (4 fly species x 12 plant species, 2285 samples).

Statistical modelling

Joint variations in fly species abundances were modelled using Poisson-LogNormal (PLN) models with the *PLN*() function in the *PLNmodels* R package (Chiquet et al. 2018, 2019). A PLN model is a multivariate mixed generalized linear model, where each species count is assumed to arise from a Poisson distribution with a parameter resulting from fixed effects of covariates and a random log Normal effect. Random effects associated to all species observed in a sample are jointly sampled from a multivariate Log Normal distribution. The variance-covariance of the multivariate distribution reveals species abundance covariations still unexplained after controlling for confounding environmental covariates and differences in sampling efforts. Samples differed in fruit number and weight, inducing uncontrolled variation in sampling effort among samples potentially leading to spurious associations between fly species. Consequently, the (log-transformed) total fruit weight of each sample was added as an offset to every tested model. Using fruit number in each sample as an offset was also tested and yielded identical conclusions (results not shown). Model diagnostics were conducted using the R package DHARMa (Hartig 2020). Covariations between species abundances unexplained by covariates were further investigated using the *PLNnetwork*() function, which adjusts the considered model under a sparsity constraint on the inverse of the variance-covariance matrix, i.e., constraining the number of edges in the resulting estimated network. The stability of the

resulting species associations was further estimated as their selection frequency in bootstrap

subsamples of the StARS model selection procedure (range 0-1; Liu et al. 2010; Appendix S3).

Model selection design

The importance of plant species identity, ecological covariates and species interactions to explain

species abundances was evaluated by model selection using the extended BIC criteria (Chen & Chen

2008). First we focused on species response to environmental variables and compared models (listed

in Table 1) including either no covariate (Model 1-0), plant species identity as a cofactor (Model 1-1),

ecological variables (all 10 FAMD dimensions described above, Model 1-3) or both plant species and

ecological variables (Model 1-5). Second, we evaluated the importance of species interactions.

PLNmodels enables fitting models where the residual variance-covariance matrix is constrained to be

diagonal. Such models assume no possible interaction between fly species. We therefore compared

all models with their diagonal counterpart (Models 1-2, 1-4 and 1-6) as a way to identify unexplained

species abundance covariations possibly resulting from species interactions. Third we estimated how

well knowledge of the fundamental niche, here laboratory-measured preferences and performances,

explained field abundances. In its present form, *PLNmodels* does not allow accounting for species

traits (covariates describe samples but not species within samples). To cope with this limitation, we

considered that assuming that species distribute according to their fundamental niche implies that

species interactions are negligible, and species distribute independently from one another. Using this

assumption we fitted models separately for each fly species, obtained their likelihoods and numbers

of parameters, and computed the BIC of the seven-species dataset as:

$BIC = kln(n) - 2ln(L),$

where *k* and *L* are the sums of the numbers of parameters and likelihoods over the seven single-

species datasets and *n* the number of samples. Following this principle, we built models where the

host plant cofactor is replaced by either preference (Model 2-7 without ecological covariates and 2-8

with ecological covariates), or performance (Models 2-9 and 2-10), or both preference and

performance (Models 2-11 and 2-12). In addition, all previous models were reevaluated on the

datasets excluding the species lacking fundamental host use estimates (*D. ciliatus*; Models 2-0 to 2-

6).

Results

All plants

The variance-covariance structure of the complete dataset including 21 plant species was first

inferred by fitting a PLN model without any covariate (Model 1-0). The obtained residual variance-

covariance matrix (Figure 2A) revealed a sharp distinction between three groups of flies: (i) the four

generalist fly species (*B. zonata*, *C. capitata*, *C. catoirii*, and *C. quilicii*), (ii) the three specialists of

Cucurbitaceae (*D. ciliatus*, *D. demmerezi* and *Z. cucurbitae*), and the specialist of Solanaceae (*N.*

cyanescens). While *N. cyanescens* abundances showed very low covariances with those of other

species, the two other groups showed positive within-group covariances and negative among-group

covariances. This variance-covariance structure suggested strong separation of the realized niches of

the three groups, likely mediated by host plants.

Figure 2: Residual variance-covariance matrices obtained after PLN model fitting on species abundances. A) Without any covariate (Model 1-0). B) With ecological covariates (Model 1-3). C) With plant species as a cofactor (Model 1-1). D) With both plant species and ecological covariates (Model 1-5).

Considering model selection between models without species traits, i.e., models including

combinations of plant species and ecological covariates (Table 1), model ranking was equivalent on

the eight-species (Models 1-0 to 1-6) and seven-species (Models 2-0 to 2-6) datasets, except that

diagonal models tended to be slightly better than their full-matrix counterparts on the latter. This

result was further shown robust to sample bootstrapping (Appendix S3). For both datasets, the

selected model included both ecological variables and host plant species as covariates (Models 1-5

and 2-6), which strongly improved the BIC (Δ_{BIC} = 17087.4 and 18935.7 respectively) as compared to

the basic model (no covariate, Models 1-0 and 2-0). Models with host plant species ranked close to

the selected model with a moderately inflated BIC relative to the selected model (Model 1-2: Δ_{BIC} =

971.1, Model 2-2: Δ_{BIC} = 933.2) and an important reduction in residual covariances relative to the

basic model (compare Figures 2A and 2C). In comparison, including ecological variables alone

deteriorated model fit with a greatly increased BIC (Model 1-3: Δ_{BIC} = 15088.8, Model 2-4: Δ_{BIC} =

11387.9) and a mild reduction of residual covariances (compare Figures 2A and 2B).

The selected model showed a good fit, with a tendency to overestimate low abundance values

(Figure S3). As expected, model diagnostics revealed an excess of zeros and some over dispersion

(Appendix S3) which PLN modeling is robust to (Chiquet et al. 2019). No temporal or spatial

autocorrelation was detected in model residuals. The coefficients relative to host plant species,

which represent the response of species abundances to host plants, had a strongly bimodal

distribution (Figure 3A, right panel). For each fly, some plant species had very low coefficients (<-80),

meaning a negligible effect of the corresponding host plant on fly abundances (approx. exp(-80) ≈ 1E-

35) and we considered them as non-host plants. The other plants had much stronger coefficients (>-

14) and were interpreted as host plants. Overall, this inferred realized host range was narrower than

the laboratory-measured fundamental host range (Figure 3A, left panel), particularly for generalists.

Only 10 fly-plant associations (out of 147) showed the reverse pattern (zero laboratory-measured

fitness and yet strong inferred response to plant), suggesting marginal difficulties with measuring

fitness in laboratory conditions. Among plants inferred as hosts from species abundance patterns

(i.e., high coefficient values), coefficients correlated positively with fly laboratory-measured fitness

for specialists but not for generalists (Figure 3B).

Models	Covariates	Residual matrix	K	L	BIC	Δ_{BIC}
A) Model set 1 (21 plants x 8 flies)						
Model 1-5	Plant + Eco	Full	284	-27664.3	57742.8	0.0
Model 1-6	Plant + Eco	Diagonal	256	-27997.8	58171.7	428.9
Model 1-2	Plant	Diagonal	176	-28608.9	58713.9	971.1
Model 1-1	Plant	Full	204	-28784.0	59302.1	1559.2
Model 1-3	Eco	Full	124	-35888.8	72831.6	15088.8
Model 1-4	Eco	Diagonal	96	-36598.4	74012.9	16270.1
Model 1-0	None	Full	44	-37228.1	74830.3	17087.4
B) Model set 2 (21 plants x 7 flies)						
Model 2-6	Plant + Eco	Diagonal	224	-20753.9	43374.1	0.0
Model 2-5	Plant + Eco	Full	245	-20723.1	43487.5	113.4
Model 2-2	Plant	Diagonal	154	-21563.1	44409.4	1035.3
Model 2-1	Plant	Full	175	-21504.7	44467.5	1093.4
Model 2-12	Preference + Performance + Eco	Diagonal	98	-23924.4	48665.3	5291.2
Model 2-10	Performance + Eco	Diagonal	91	-24387.4	49533.0	6158.9
Model 2-11	Preference + Performance	Diagonal	28	-25689.9	51613.0	8238.9
Model 2-8	Preference + Eco	Diagonal	91	-25516.9	51792.1	8418.0
Model 2-9	Performance	Diagonal	21	-26443.3	53061.6	9687.5
Model 2-4	Eco	Diagonal	84	-27082.1	54864.0	11489.9
Model 2-7	Preference	Diagonal	21	-27628.9	55432.8	12058.7
Model 2-3	Eco	Full	105	-27321.5	55517.8	12143.7
Model 2-0	None	Full	35	-31060.1	62411.8	19037.7

Table 1: Model selection on the 21-plant dataset (n = 4918). Models are ranked by increasing BIC (from best to worst). k is the number of parameters. L is the log-likelihood. Δ_{BIC} is the BIC difference between any focal model and the best one.

Figure 3: Comparison of fly species' fundamental and realized host use. A) Fundamental host use as measured in the laboratory (left panel – fitness is the logarithm of the product preference and performance) and realized host use as inferred from regression coefficients relative to host plants in the fullest model (right panel – response is obtained from Model 2-5 on the seven-species dataset). B) Relationship between inferred responses to host plants and laboratory-measured fitness for specialists in orange and generalists in green on hosts detected as such in the field. Lines represent linear regressions with slope 95% confidence intervals as shadowed areas.

The coefficients relative to the first two axes of the FAMD on ecological covariates could be interpreted as responses of fly abundances to rainfall, temperature, and elevation (Figure 4). Rainfall had a low effect on the abundances of *B. zonata*, *C. capitata, C. quilicii*, and *D. demmerezi*. *Ceratitis catoirii, N. cyanenscens,* and *Z. cucurbitae* showed a propensity towards warm high-rainfall areas, while *D. ciliatus* seemed to prefer colder drier climates. *Ceratitis quilicii* and *C. catoirii* were not much affected by temperature. *Bactrocera zonata*, *C. capitata*, *D. ciliatus,* and *Z. cucurbitae* should thrive in

low-elevation warm climates. *Dacus demmerezi* and to a lesser extent *N. cyanescens* seemed to

prefer colder higher-elevation climates.

Figure 4: Species abundance responses to ecological variables. A) Correlation circle on the first and

second axes of the FAMD (factorial analysis of mixed data) on ecological variables (Dim1 and Dim2, respectively). The first axis contrasts warm low-altitude sites and colder high-altitude sites. The second axis is a gradient of rainfall and maximal temperature (see Supplementary figure S3 for details on axes contributions). B) Regression slopes relative to Dim1 and Dim2, inferred under the fullest model on the

21 plant x 8 fly species dataset (Model 1-5). Error bars represent approximate confidence intervals (1.96 x standard errors). For the first axis (Dim1), negative slopes (e.g., *B. zonata* and *C. capitata*) can be interpreted as a positive effect of temperatures and a negative effect of elevation on species

abundances. For the second axis (FAMD 2), positive slopes (e.g., *C. catoirii*) can be interpreted as a positive effect of rainfall on species abundances.

The selected model had full residual variance-covariance matrix on the eight-species dataset and

diagonal residual variance-covariance matrix on the seven-species dataset. On the eight-species

dataset its diagonal version ranked second with Δ_{BIC} = 428.9. Congruently, the residual variance-

covariance matrix inferred under the selected model (Figure 2D) had very low covariance values for

all pairs of fly species, all covariances being negative. This suggested possible, though weak,

antagonist interactions. The largest residual correlations were observed between the three

specialists *D. demmerezi, D. ciliatus* and *Z. cucurbitae* (residual correlations ranging from -0.031 to -

0.112) and between the generalists *B. zonata*, *C. quilicii* and *C. capitata* (residual correlations from -0.025 to -0.090). Network inference, applied to the selected model, converged to one significant interaction between *D. ciliatus* and *Z. cucurbitae*, with a stability of 0.99 (i.e., detected in 99 out of 100 network inferences on bootstrapped data). Overall these results suggested that at the scale of the whole community competitive interactions between fly species only weakly affected their joint distributions and that fly species abundances were mainly explained by environmental covariates.

Considering models accounting for species fundamental niche (Table 1B), the best model with species traits included female preference, larval performance and ecological variables (Model 2-12, Δ_{BIC} = 5189.2). It ranked intermediate between the model with ecological variables alone and the best model. Such good performance of models with species traits suggests that the fundamental host range of fly species is an important determinant of fly species joint distributions. Whether in combination with ecological covariates or not, models with larval performance were slightly better than models with female preference.

Cucurbitaceae

Focusing on Cucurbitaceae and the three fly species they hosted, *D. ciliatus*, *D. demmerezi* and *Z. cucurbitae*, the selected model was again the one including both host plant species and ecological covariates (Model 1-5, Table S5A). Its BIC was lower than the one of the basic model (Model 1-0, Δ_{BIC} = 1208.0). It assumed a full residual variance-covariance matrix and performed better than its diagonal version (Δ_{BIC} = 190.6). Network inference yielded two significant negative interactions between *D. ciliatus* and *Z. cucurbitae* (stability = 1.0) and between *D. demmerezi* and *Z. cucurbitae* (stability = 0.56). For all other models, versions with full residual matrix performed better than their diagonal counterparts. Besides, models including plant species alone or ecological covariates alone performed equivalently (Δ_{BIC} = 515.3 vs. Δ_{BIC} = 653.2). This is congruent with the idea that plant species are more similarly affecting fly distributions within the Cucurbitaceae family than at the scale

of the 21-plant dataset, so that the influence of plant identity is less important for species distributions among Cucurbitaceae than on a wider set of plants.

For *D. demmerezi* and *Z. cucurbitae,* for which traits were available, models with species traits alone had poor performance, and displayed higher BIC than the basic model (Table S5B). However, the model with both species traits and ecological covariates ranked third just after the diagonal version of the selected model (Model 2-10, Δ_{BIC} = 124.9).

Solanaceae

For Solanaceae and their associated fly species, *N. cyanescens, C. capitata,* and *C. quilicii*, the selected model included host plant species, ecological covariates and a diagonal residual variance-covariance matrix (Model 1-6, Table S6). It improved the BIC of the basic model by 201.8. It performed better than its full residual matrix version (Δ_{BIC} = 15.8) and congruently, applying network inference to the latter yielded no significant interaction between fly species. On this dataset all models with a diagonal matrix performed better than their full residual matrix counterpart, further confirming the absence of detected interactions. Models with ecological covariates alone performed poorly (Model 1-3 Δ_{BIC} = 176.4 and Model 1-4 Δ_{BIC} = 162.0, respectively). Models with species traits were almost as good as their equivalent with host plant as a cofactor. For instance, the model with species traits and ecological covariates (Model 2-10) ranked second with Δ_{BIC} = 9.5. Female preference and larval performance performed equally well (Δ_{BIC} = 44.6 and Δ_{BIC} = 38.5 respectively) and almost as good as both traits together (Δ_{BIC} = 33.4), suggesting important correlation between the traits.

Other plant families

The last dataset considers all families other than Cucurbitaceae or Solanaceae with *B. zonata*, *C. capitata*, *C. catoirii*, and *C. quilicii* (Table S7). The selected model included host plant species and

ecological covariates (Model 1-5, Δ_{BIC} = 2639.7 with the basic model 1-0). It assumed a full residual

variance-covariance matrix and performed slightly better than its diagonal version (Δ_{BIC} = 55.8). All

models with a full residual matrix performed better than their diagonal counterpart. Network

inference yielded one significant interaction between *B. zonata* and *C. quilicii* (stability = 1.0).

The model with only ecological covariates performed well (Model 1-3, Δ_{BIC} = 967.3), and almost as

good as the model with host plant alone (Model 1-1, Δ_{BIC} = 684.1), suggesting redundancy between

ecological information and plant identity. Models that included species traits without ecological

covariates performed badly (Δ_{BIC} > 2220), and the model with species traits and ecological covariates

(Model 2-10, Δ_{BIC} = 814.3) performed only slightly better than the model with ecological covariates

only (Model 1-3, Δ_{BIC} = 967.3). Of the two traits, only female preference really improved model fit

(Model 2-8 Δ_{BIC} = 888.9 vs. Model 2-12 Δ_{BIC} = 1050.0).

Discussion

The determinants underlying the structure of a community of eight Tephritid fly species were

deciphered. Modelling joint species abundances without accounting for any covariate (only

intercepts and an offset) confirmed a major role of host use strategy on fly species abundances.

Species abundances co-varied positively among generalists and among specialists and negatively

between species from these two groups. Common responses to environmental factors may cause

positive residual correlations in species abundances, while divergent responses will imply negative

correlations, potentially leading to incorrect interpretations of species interactions (Ovaskainen et al.

2016, Dormann et al. 2018). Accounting for environmental covariates strongly improved model fit

and made all residual covariances almost completely vanish, particularly among groups, suggesting

that no important environmental factor structuring the community has been missed. Obtaining long-

term abundance data on all *a priori* relevant species of a given community is rarely possible. Here

over the study period, one other fruit fly species (*Carpomya vesuviana*) was mentioned locally. It was

considered very rare and only present on Indian jujube (*Ziziphus mauritiana*) in dry low-elevation

areas (Quilici & Jeuffrault 2001, Franck et al. 2017). It was not detected in the 204 jujube samples of

the full dataset. Second, not all 108 plant species considered as potential hosts were included. To

evaluate the effect of omitted plants, we conducted the analysis on all samples with GPS coordinates

and at least 10 samples with emerging flies (6434 samples, 36 plant species including the 21 studied

in the laboratory) and obtained strikingly similar results (model ranking, and estimates of regression

coefficients, Appendix S3), comforting the idea that most relevant factors have been accounted for.

Detection of competitive interactions

Some residual covariances remained non-negligible after accounting for fly species' response to host

plants between some generalists and between some specialists of Cucurbitaceae. They were all

negative, suggesting a possible minor role of antagonistic interactions within specialists and within

generalists. Only one of these residual covariations resisted the network inference process on the

whole dataset (*D. ciliatus - Z. cucurbitae*, the two most abundant specialists of Cucurbitaceae). Two

more significant covariations were detected when focusing on Cucurbitaceae (*Z. cucurbitae - D.*

demmerezi) and on other plant families (*B. zonata - C. quilicii*) suggesting possible dependence of

species interactions on host plants. On Cucurbitaceae, although qualitatively congruent with other

independent empirical measures of host range (Vayssieres et al. 2008) and climatic niche (Vayssières

& Carel 1999), host plant species and abiotic factors only moderately improved model fit. All three

specialist flies found on Cucurbitaceae are able to thrive on any plant of this family (Charlery de la

Masselière et al. 2017b) and competitive interactions between these fly species are highly plausible

(Vayssieres et al. 2008). On other plant families, both host plants and abiotic factors clearly improved

model fit, congruently with former interpretations of the system (Duyck et al. 2006a, Duyck et al.

2008). Responses of generalist species to abiotic factors were strikingly congruent with former

independent laboratory experiments (Duyck & Quilici 2002, Duyck et al. 2004, Duyck et al. 2006a).

There was redundancy between host plants and abiotic factors. Many of these plants are exploited

but not planted (e.g., Myrtaceae). Their distributions are therefore more dependent on ecological

factors than those of Solanaceae and Cucurbitaceae, which are mainly cultivated throughout the year in Réunion. On Solanaceae, no residual covariation was detected. Plant identity was the main determinant of species abundances, congruently with the idea that Solanaceae impose adaptive challenges on their fruit consumers through a variety of toxic compounds (Brévault et al. 2008), rendering host adaptation the main factor driving species abundances.

A current competition ghost

Among-group covariation between specialists and generalists were mainly attributable to fly species' adaptation to host use with a minor contribution of abiotic factors. Previous studies have highlighted differences in host adaptation between these fly species (Hafsi et al. 2016). Contrary to the specialists, which are mainly able to use their preferred hosts, the four generalists can thrive in numerous plant species, and have weak female preferences (Charlery de la Masselière et al. 2017b). Accordingly, specialists were seldom found in plants other than Cucurbitcaeae or Solanaceae. However, these results do not explain why generalists were so rarely found on Cucurbitaceae and Solanaceae. Competitive exclusion with specialists would be a natural hypothesis to explain this absence (Nakadai et al. 2018). Here, no competition among groups was detected. It is possible, however, that competition has operated and that competitive exclusion has been so strong that generalists cannot be found on Cucurbitaceae. In PLN modelling, such absence could be interpreted as negative response of generalist abundances to Cucurbitaceae and be encapsulated in plant cofactor slope rather than in species interactions. It is precisely when competition is intense enough to cause niche partitioning that it can no longer be detected. This result evokes a well-known paradox in ecology formerly termed ''the ghost of competition past'' (Connell 1980) according to which the observed differentiation in niches is the result of past interspecific competition.

To escape the paradox, knowledge about the fundamental niches of species through eco-evolutionary approaches could help settle whether species interactions are an important driver of species assemblages (Augustyn et al. 2016, Dormann et al. 2018). Laboratory measurements of larval

performance and female preference on all host plants were used in replacement of plant identity. Congruently with the community being essentially driven by host use, preference and performance clearly improved model fit. Interestingly, performance was more informative than preference, which was expected from previous knowledge that generalists' preferences are uncorrelated to their performances (Charlery de la Masselière et al. 2017b). If competition truly shapes species abundances and has not been detected, it is to be found in the difference between models with plant identity as a cofactor and models with species traits instead. Here we found a difference suggesting that competitive exclusion is at work. In terms of importance, from model rankings, host use patterns were the most important factor shaping species abundances, followed by abiotic factors and possibly a dose of competition.

This predominance of host plants as a structuring factor of phytophagous insect communities has been much debated, but congruent studies exist. In analyses of insect communities along road verges, Schaffers et al. (2008) found that the composition of plant communities was a much better predictor of insect and spider assemblages than environmental variables. Similarly Nakadai et al. (2018) found that sharing of host species was predominant among butterflies of the Japanese archipelago, suggesting that interspecific resource competition may not effectively determine community assembly patterns at regional scales. In an earlier review on the importance of competition in insects, Denno et al. (1995) pointed that only weak to moderate effect of competition should certainly be expected in phytophagous insects such as Tephritids due to their high mobility and weak aggregation behaviors. Experimental manipulations of competitive interactions in the field could offer a promising way to test the validity of the present inferences. These experiments would also be useful to unveil the role of other biotic interactions (e.g., natural enemies), as forces capable of modulating interspecific competition between fruit flies.

Generalists vs. specialists

Overall specialists and generalists almost had very distinct realized host uses with different assembly rules. That specialists and generalists form separate interaction networks has already been highlighted, e.g., among soil microbial species (Barberan et al. 2012). It is well known that the predictability of assemblages differs between host-plant generalists and specialists (Müller et al. 2011). This has led to the hypothesis that specialists would have assemble according to the species-sorting paradigm of metacommunity ecology (Leibold et al. 2004), whereby species occurrences are mainly driven by habitat heterogeneity and local adaptation. Generalists' assembly rules, on the other hand, would rather follow a mass-effect paradigm (Shmida & Wilson 1985), according to which sink populations, where the species is maladapted, can persist through a migration influx from source populations (Müller et al. 2011). Our results confirm this hypothesis. For specialists, a good agreement between the inferred host plant effects on species abundances and the laboratory measures of host adaptation suggested that specialists were mainly filtered by host plant characteristics. In contrast, generalists displayed no relationship between inferred and laboratory-measured host plant effects, suggesting that generalists were found on some hosts where their fitness is low and at low density on good hosts. Besides generalists use fruits whose availability is highly variable over time. Contrary to specialists, most of their hosts are not available all year long (Figure S4). This temporally variable habitat may trigger a dynamics of local extinction and recolonization, in which the roles of migration and stochasticity become more important than that of host adaptation and in which coexistence is possible despite fundamental niche overlap (Connell 1980, Chesson 2000).

Acknowledgements

This work is dedicated to Serge Quilici who has been leading prolific research on the Tephritids of Réunion his whole career. Cirad technicians Jim Payet and Serge Glénac made this study possible through their invaluable expertise with fly rearing and ecology. We also thank them as well as

Antoine Franck, Christophe Simiand, and Patrick Turpin, for collecting field data over the years.

Thomas Brequigny contributed to measuring larval traits during his internship. This work used images

acquired within the framework of the CNES Kalideos device (Réunion site), which benefited from the

"Programme Investissements d'Avenir" EQUIPEX of the French "Agence Nationale de la Recherche"

on project GEOSUD bearing the reference ANR-10-EQPX-20. The images also required financial

support the French Ministry of Agriculture and field data transmitted by the "Syndicat du Sucre de la

Réunion" and the "SAFER de la Réunion". BF, FC, FM, JC, MD, SR, and VR received the financial

support of the French "Agence Nationale de la Recherche" project NGB (ANR-17-CE32-011). EF, FC,

PFD, and VR were funded by the European Union (European Regional Development Fund, ERDF

contract GURDT I2016-1731-0006632), the Conseil Régional de La Réunion, and the Centre de

Coopération Internationale en Recherche Agronomique pour le Développement (CIRAD). AH was

funded by the "Ministère de l'Enseignement supérieur et de la Recherche Scientifique de la Tunisie".

This study used the facilities provided by the Plant Protection Platform (3P, IBISA), Saint-Pierre,

Réunion, France.

Data and script accessibility

Data and script are available online: http://doi.org/10.5281/zenodo.4569532

Literature cited

Adler, P. B. & HilleRisLambers, J. (2008) The influence of climate and species composition on the population dynamics of ten prairie forbs. *Ecology*, 89, 3049-3050.

Araújo, M. B. & Rozenfeld, A. (2014) The geographic scaling of biotic interactions. *Ecography*, 37, 406-415.

Augustyn, W. J., Anderson, B. & Ellis, A. G. (2016) Experimental evidence for fundamental, and not realized, niche partitioning in a plant–herbivore community interaction network. *J. Anim. Ecol.*, 85, 994-1003.

Barberan, A., Bates, S. T., Casamayor, E. O. & Fierer, N. (2012) Using network analysis to explore co-occurrence patterns in soil microbial communities. *ISME J.*, 6: 343-351.

Blanchet, F. G., Cazelles, K. & Gravel D. (2020) Co-occurrence is not evidence of ecological interactions. *Ecol. Lett.*, 23, 1050-1063.

Brazeau, H. A. & Schamp, B. S. (2019) Examining the link between competition and negative co-occurrence patterns. *Oikos*, 128, 1358-1366.

Brévault, T., Duyck, P.F. & Quilici, S. (2008) Life-history strategy in an oligophagous tephritid: the tomato fruit fly, *Neoceratitis cyanescens*. *Ecol. Entomol.*, 33, 529-536

Charlery de la Masselière, M., Ravigné, V., Facon, B., Lefeuvre, P., Massol, F., Quilici, S. & Duyck, P. F. (2017a) Changes in phytophagous insect host ranges following the invasion of their community: Long-term data for fruit flies. *Ecol. Evol.*, 7, 5181-190.

Charlery de la Masselière, M., Facon, B., Hafsi, A. & Duyck, P. F. (2017b) Diet breadth modulates preference-performance relationships in a phytophagous insect community. Sci. Rep., 7, 16934.

Chen J. and Chen Z. (2008) Extended Bayesian information criteria for model selection with large model spaces. Biometrika, 95, 759-771.

Chesson, P. (2000) Mechanisms of maintenance of species diversity. *Ann. Rev. Ecol. Evol. Syst.*, 31, 343-366.

Chiquet, J., Mariadassou, M. & Robin, S. (2018) Variational inference for probabilistic poisson PCA. *Ann. Appl. Stat.*, 12, 2674–2698.

Chiquet, J., Mariadassou, M. & Robin, S. (2019) Variational Inference for Sparse Network Reconstruction from Count Data. In Proceedings of the 36th International Conference on Machine Learning, edited by Kamalika Chaudhuri and Ruslan Salakhutdinov, 97, 1162–71. Proceedings of Machine Learning Research. Long Beach, California, USA: PMLR.

Connell, J. H. (1980) Diversity and the coevolution of competitors, or the ghost of competition past. *Oikos*, 35, 131-138.

D'Amen, M., Mod, H. K., Gotelli, N. J. & Guisan, A. (2018) Disentangling biotic interactions, environmental filters, and dispersal limitation as drivers of species co-occurrence. *Ecography*, 41, 1233-1244.

Denno, R. F., McClure, M. S. & Ott, J. R. (1995) Interspecific interactions in phytophagous insects: competition reexamined and resurrected. *Annu. Rev. Entomol.*, 40, 297-331.

Diamond, J. M. (1975) Assembly of species communities. – In: Cody, M. L. and Diamond, J. M. (eds), Ecology and evolution of communities. Harvard Univ. Press, pp. 342–344.

Dormann, C. F., Bobrowski, M., Dehling, D. M., Harris, D. J., Hartig, F., Lischke, H., Moretti M. D., Pagel, J., Pinkert, S., Schleuning, M., Schmidt, S. I., Sheppard C. S., Steinbauer M. J., Zeuss D. & Kraan, C. (2018) Biotic interactions in species distribution modelling: 10 questions to guide interpretation and avoid false conclusions. *Glob. Ecol. Biogeogr.*, 27, 1004-1016.

Duyck, P. F. & Quilici, S. (2002) Survival and development of different life stages of three *Ceratitis* spp. (Diptera: Tephritidae) reared at five constant temperatures. *B. Entomol. Res.*, 92, 461-469.

Duyck, P. F., Sterlin, J. F. & Quilici, S. (2004) Survival and development of different life stages of *Bactrocera zonata* (Diptera: Tephritidae) reared at five constant temperatures compared to other fruit fly species. *B. Entomol. Res.*, 94, 89-93.

Duyck, P. F., David, P. & Quilici, S. (2006a) Climatic niches partitioning following successive invasions

by fruit flies in La Réunion. *J. Anim. Ecol.*, 75, 518-526.

Duyck, P. F., David, P., Junod, G., Brunel, C., Dupont, R. & Quilici, S. (2006b) Importance of

competition mechanisms in successive invasions by polyphagous Tephritids in La Réunion.

Ecology, 87, 1770-1780.

Duyck, P. F., David, P., Pavoine, S. & Quilici, S. (2008) Can host-range allow niche differentiation of

invasive polyphagous fruit flies (Diptera: Tephritidae) in La Réunion? *Ecol. Entomol.*, 33,

439-452.

Elith, J. & Leathwick, J. R. (2009) Species distribtuion models: ecological explanation and prediction

across space and time. *Ann. Rev. Ecol. Evol. Syst.*, 40, 677-697.

Ferrier, S. & Guisan, A. (2006) Spatial modelling of biodiversity at the community level. *J. Appl. Ecol.*,

43, 393-404.

Godinho, D. P., Janssen, A., Dias, T., Cruz, C. & Magalhães S. (2016) Down-regulation of plant defence

in a resident spider mite species and its effect upon con- and heterospecifics. *Oecologia*,

180, 161-167.

Gotelli, N. J. & Graves, G. R. (1996) Null models in Ecology. Smithsonian Institution Press, Washington

DC.

Gotelli, N. J., Graves, G. R. & Rahbek C. (2010) Macroecological signals of species interactions in the

Danish avifauna. *Proc. Natl. Acad. Sci. U.S.A.*, 107, 5030-5035.

Hafsi, A., Facon, B., Ravigné, V., Chiroleu, F., Quilici, S., Chermiti, B. & Duyck, P. F. (2016) Host plant

range of a fruit fly community (Diptera: Tephritidae): Does fruit composition influence larval

performance? *BMC Ecology*, 16, 40.

Hairston, N. G., Smith, F. E. & Slobodkin, L. W. (1960) Community Structure, Population Control, and

Competition. *Am. Nat.*, 94, 421-425.

Hartig, F. (2020) DHARMa: Residual Diagnostics for Hierarchical (Multi-Level / Mixed) Regression

Models. R package version 0.3.3.0. https://CRAN.R-project.org/package=DHARMa

Heikkinen, R. K., Luoto, M., Virkkala, R., Pearson, R. G. & Körber J. H. (2007) Biotic interactions

improve prediction of boreal bird distributions at macro-scales. *Global Ecol. Biogeogr.*, 16,

754-763.

Jabot, F., Laroche, F., Massol, F., Arthaud, F., Crabot, J., Dubart, M., Blanchet, S., Munoz, F., David, P.

& Datry, T. (2020) Assessing metacommunity processes through signatures in

spatiotemporal turnover of community composition. *Ecol. Lett.*, 23, 1330-1339.

Kaplan, I. & Denno, R. F. (2007) Interspecific interactions in phytophagous insects revisited: a

quantitative assessment of competition theory. *Ecol. Lett.*, 10, 977-994.

Kraft, N. J. B., Valencia, R. & Ackerly D. D. (2008) Functional traits and niche-based tree community

assembly in an Amazonian forest. *Science*, 322, 580-582.

Lavorel, S., McIntyre, S., Landsberg, J. & Forbes, T. D. A. (1997) Plant functional classifications: from

general groups to specific groups based on response to disturbance. *Trends Ecol. Evol.*, 12,

474-478.

Lê, S., Josse, J. & Husson, F. (2008). FactoMineR: An R Package for Multivariate Analysis. *J. Stat.*

Softw., 25, 1-18.

Leibold, M. A., Holyoak, M., Mouquet, N., Amarasekare, P., Chase, J. M., Hoopes, M. F., Holt, R. D.,

Shurin, J. B., Law, R., Tilman, D., Loreau, M. & Gonzalez, A. (2004) The metacommunity

concept: a framework for multi-scale community ecology. *Eco. Lett.*, 7, 601-613.

Liu, H., Roeder, K. & Wasserman, L. (2010) Stability Approach to Regularization Selection (Stars) for

High Dimensional Graphical Models. In Proceedings of the 23rd International Conference on

Neural Information Processing Systems - Volume 2, 1432–40. USA.

Meynard, C. N., Lavergne, S., Boulangeat, I. & Garraud, L. (2013). Disentangling the drivers of

metacommunity structure across spatial scales. *J. Biogeogr.*, 40, 1560–1571.

Müller, J., Stadler, J., Jarzabek-Müller, A., Hacker, H., ter Braak, C. & Brandl, R. (2011) The

predictability of phytophagous insect communities: host specialists as habitat specialists.

PLoS One, 6, e25986.

Nakadai, R., Hashimoto, K., Iwasaki, T. & Sato, Y. (2018) Geographical co−occurrence of butterfly

species: the importance of niche filtering by host plant species. *Oecologia*, 186, 995-1005.

Ovaskainen, O., Roy, D. B., Fox, R. & Anderson B. J. (2016) Uncovering hidden spatial structure in

species communities with spatially explicit joint species distribution models. *Methods Ecol.*

Evol., 7, 428-436.

Poisot, T., Stouffer, D. B. & Gravel, D. (2015) Beyond species: why ecological interaction networks

vary through space and time. *Oikos*, 124, 243-251.

Pollock, L. J., Tingley, R., Morris, W. K., Golding, N., O'Hara, R. B., Parris, K. M., Vesk, P. A. &

McCarthy, M. A. (2014) Understanding co-occurrence by modelling species simultaneously

with a Joint Species Distribution Model (JSDM). *Methods Ecol. Evol.*, 5, 397-406.

Quilici, S. & Jeuffrault, E. (2001) Plantes-hôtes des mouches des fruits: Maurice, Réunion, Seychelles.

PRMF/COI, Imp. Graphica, St André, La Réunion.

Roy, H. E., Rorke, S. L, Beckmann, B., Botham, M. S., Brown, P. M. J., Noble, D., Sewell, J. & Walker, K.

J. (2015) The contribution of volunteer recorders to our understanding of biological

invasions. *Biol. J. Linn. Soc.*, 115, 678-689.

Schaffers, A. P., Raemakers, I. P., Sykora, K. V. & ter Braak C. J. F. (2008) Arthropod assemblages are

best predicted by plant species composition. *Ecology*, 89, 782-794.

Shmida, A. & Wilson, M. V. (1985) Biological determinants of species diversity. *J. Biogeogr.*, 12, 1-20.

Stewart, A. J. A., Bantock, T. M., Beckmann, B. C., Botham, M. S., Hubble, D. & Roy, D. B. (2015) The

role of ecological interactions in determining species ranges and range changes. *Bio. J. Linn.*

Soc., 115, 647-663.

Tack, A. J. M., Ovaskainen, O., Harrison, P. J. & Roslin, T. (2009) Competition as a structuring force in

leaf miner communities. *Oikos*, 118, 809-818.

Thuiller, W., Pollock, L. J., Gueguen, M. & Münkemüller, T. (2015) From species distributions to meta-

2   communities. *Ecol. Lett.*, 18, 1321-1328.

3   Ulrich, W., Jabot, F. & Gotelli, N. J. (2017) Competitive interactions change the pattern of species co-

4   occurrences under neutral dispersal. *Oikos*, 126, 91-100.

5   Vayssières, J. F. & Carel, Y. (1999) Les Dacini (Diptera : Tephritidae) inféodés aux Cucurbitaceae à la

6   Réunion : gamme de plantes-hôtes et stades phénologiques préférentiels des fruits au

7   moment de la piqûre pour des espèces cultivées. *Ann. Soc. Entomol. Fr.*, 35, 197-202.

8   Vayssières, J. F., Carel, Y., Coubes, M. & Duyck P. F. (2008) Development of immature stages and

9   comparative demography of two cucurbit-attacking fruit flies in Reunion island: *Bactrocera*

cucurbitae and *Dacus ciliatus* (Diptera Tephritidae). *Env. Entomol.*, 37, 307-314.

Wisz, M. S., Pottier, J., Kissling, W. D., Pellissier, L., Lenoir, J., Damgaard, C. F., Dormann, C. F.,

Forchhammer, M. C., Grytnes, J.A., Guisan, A., Heikkinen, R. K., Høye, T. T., Kühn, I., Luoto,

M., Maiorano, L., Nilsson, M. C., Normand, S., Ockinger, E., Schmidt, N. M., Termansen, M.,

Timmermann, A., Wardle, D. A., Aastrup, P. & Svenning J. C. (2013) The role of biotic

interactions in shaping distributions and realised assemblages of species: implications for

species distribution modelling. *Biol. Rev.*, 88, 15-30.

Zurell, D., Pollock, L. J. & Thuiller, W. (2018) Do joint species distribution models reliably detect

interspecific interactions from co-occurrence data in homogenous environments?

Ecography, 41, 1812-1819.

Supplementary material

Appendix S1: Additional materials & methods

Details on species table

Among the 12872 initial samples, only samples that fulfilled to following conditions were kept: (i) GPS coordinates could be retrieved (n=9715 in 41 plant species), (ii) at least one Tephritid fly emerged (n=6455 in 41 plant species), (iii) from plant species which had been successfully sampled at least 10 times in the dataset (n=6434 in 36 plants), and (iv) from plant species that were also characterized in the laboratory (n=4918 in 21 plants). In the resulting dataset, 97351 individual flies were counted. Samples covered 104 field sessions all year round over the 1991-2009 period (Tables S1 and S2). The number of flies per sample varied from 1 to 2244. Samples differed in terms of fruit number and weight. The number of flies varied from 1 to 188 per individual fruit and from 1.1E-3 to 15.8 flies per gram of fruit.

Table S1: Number of samples (*n*) by year in the full 21-plant dataset

Year	1991	1992	1993	1994	1995	1996	1997	1998	2001	2002	2003	2004	2005	2008	2009
n	4	15	31	52	80	38	434	1143	83	523	634	19	205	191	1466

Table S2: Number of samples (*n*) by month in the full 21-plant dataset

Month	Jan	Feb	Mar	Apr	May	Jun	Jul	Aug	Sep	Oct	Nov	Dec
n	447	818	708	458	328	182	594	252	222	238	242	429

The number of samples per plant species (Table S3) varied from 12 (*Averrhoa carambola*) to 1105 (*Cucurbita maxima*). The most represented plant families were the Cucurbitaceae (six plant species and 2374 samples), the Myrtaceae (four plant species and 1102 samples) and the Combretaceae (one plant species with 853 samples).

Table S3: Number of samples (*n*) for each of the 21 plant species retained in the study

Family	Species	Common English name	*n*
Anacardiaceae	*Mangifera indica*	Mango	36
Annonaceae	*Annona reticulata*	Custard apple	27
Combretaceae	*Terminalia catappa*	Indian almond	853
Cucurbitaceae	*Citrullus lanatus*	Water melon	175
	Cucumis melo	Melon	30
	Cucumis sativus	Cucumber	440
	Cucurbita maxima	Pumpkin	1105
	Cucurbita pepo	Zucchini	504
	Sechium edule	Chayote	120
Myrtaceae	*Psidium cattleyanum*	Strawberry guava	422
	Psidium guajava	Guava	398
	Syzygium jambos	Rose apple	255
	Syzygium samarangense	Java apple	27
Oxalidaceae	*Averrhoa carambola*	Star fruit	12
Rosaceae	*Eriobotrya japonica*	Loquat	61
	Prunus persica	Peach	99
Rubiaceae	*Coffea arabica*	Coffee	81
Rutaceae	*Citrus reticulata*	Mandarin	14
Solanaceae	*Capsicum annuum*	Chilli	29
	Solanum lycopersicum	Tomato	68
	Solanum mauritianum	Bugweed	162

Plant species differed in terms of the patterns of occurrence of the eight fly species (Figure 2B). In particular, Cucurbitaceae were mostly used by the three species known as specialists of this plant family, *D. ciliatus* (14047 flies in 1264 samples), *D. demmerezi* (6335 flies in 276 samples) and *Z. cucurbitae* (21830 flies in 1463 samples). Other fly species were detected in fewer than 10 Cucurbitaceae samples. Solanaceae mainly hosted the specialist species *N. cyanescens* (1205 flies in 225 samples). They were also, to a lesser extent, hosts of *C. capitata* (116 flies in 32 samples), and *C. quilicii* (131 flies in 22 samples). Other fly species were detected in less than 5 Solanaceae samples. The other plant species (12 plant species) were mainly used by the four generalist species, i.e., *B. zonata* (11517 flies in 728 samples), *C. capitata* (8735 flies in 537 samples), *C. quilicii* (33149 flies in 1583 samples), *C. catoirii* (291 flies in 65 samples). Other fly species were detected in less than 5 samples.

1 Co-occurrences between fly species were frequent. 1148 samples out of 4918 (23.3%) contained

2 more than one species, and 759 out of 4335 samples with a single fruit (17%) contained more than

3 one fly species. Co-occurrences between fly species were therefore much structured by whether the

4 considered plant species belonged to Cucurbitaceae, Solanaceae or other families (Table S4).

6 Table S4: Species co-occurences in the full 21-plant dataset (n = 4918 samples)

	Bactrocera zonata	*Ceratitis quilicii*	*Ceratitis capitata*	*Ceratitis catoirii*	*Neoceratitis cyanescens*	*Dacus ciliatus*	*Dacus demmerezi*	*Zeugodacus cucurbitae*
Bactrocera zonata	728	198	120	1	0	0	0	0
Ceratitis quilicii	198	1605	309	57	12	1	0	1
Ceratitis capitata	120	309	570	31	15	1	0	2
Ceratitis catoirii	1	57	31	66	1	1	0	1
Neoceratitis cyanescens	0	12	15	1	237	3	1	5
Dacus ciliatus	0	1	1	1	3	1266	107	454
Dacus demmerezi	0	0	0	0	1	107	276	133
Zeugodacus cucurbitae	0	1	2	1	5	454	133	1467

Ecological covariables

Elevation (in meters) of each sample was obtained from the Digital Elevation Model Litto3D® co-

produced by the French IGN (National Geographic Institute) and the SHOM (Marine Oceanographic

Hydrographic Service). Pluviometry was obtained from layers produced by M. Mezino from 143

CIRAD and Meteo-France meteorological stations, containing isohyets of (i) the minimal rainfall

observed in the 20% most humid years of the 1986-2016 period (ii) the minimal rainfall observed in

the 20% driest years of the 1986-2016 period and (iii) the median annual rainfall over the 1986-2016

period. For each sample, the value of closest isohyet was retained for each of the three variables.

Temperature was characterized by the (i) minimal, (ii) mean and (iii) maximal annual temperature

over the 1987-2017 period as interpolated by M. Mezino from 73 CIRAD and Meteo-France

meteorological stations raw data. Land use around each sample location was obtained from a 12-

categories layer produced by supervised classification of Pléiades 2018 images (Dupuy and Gaetano,

2019 doi:10.18167/DVN1/WKAJZO, CIRAD Dataverse, V1).

Additional details on FAMD on ecological covariables

Figure S1: Interpretation of the axes of the FAMD on ecological covariables. A) Percentage of

variance explained by each of the 10 axes of the FAMD. The first and second FAMD axes explain

19.8 % of the variance. C-D) Contribution of individual ecological covariates to Axis 1, 2 and other

axes, respectively. Axis 1 is mainly contributed by temperature and elevation. Axis 2 is mainly

contributed by rainfall and maximal temperature. Other axes are contributed by qualitative variables

(year, month and land use).

1 *Appendix S2: Complementary results*

2 *Model selection on the three sub-datasets*

3 *Cucurbitaceae*

4 Table S5: Model selection on the Cucurbitaceae dataset ($n = 2347$). Models are ranked by increasing

5 BIC (from best to worst). k is the number of parameters. L is the log-likelihood. Δ_{BIC} is the BIC

6 difference between any focal model and the best one.

Model	Covariates	Residual matrix	k	L	BIC	Δ_{BIC}
A) Model set 1 (6 plants x 3 flies)						
Model 1-5	Plant + Eco	Full	54	-14168.3	28756.3	0.0
Model 1-6	Plant + Eco	Diagonal	51	-14275.3	28947.0	190.6
Model 1-1	Plant	Full	24	-14542.5	29271.6	515.3
Model 1-3	Eco	Full	39	-14553.2	29409.5	653.2
Model 1-2	Plant	Diagonal	21	-14661.3	29485.7	729.4
Model 1-4	Eco	Diagonal	36	-14649.7	29579.2	822.8
Model 1-0	None	Full	9	-14947.2	29964.3	1208.0
B) Model set 2 (6 plants x 2 flies)						
Model 2-5	Plant + Eco	Full	35	-7311.6	14881.6	0.0
Model 2-6	Plant + Eco	Diagonal	34	-7361.7	14974.4	92.8
Model 2-10	Preference + Performance + Eco	Diagonal	28	-7399.9	15006.5	124.9
Model 2-3	Eco	Full	25	-7419.8	15024.2	142.6
Model 2-8	Preference + Eco	Diagonal	26	-7421.5	15035.1	153.5
Model 2-12	Performance + Eco	Diagonal	26	-7423.9	15039.8	158.2
Model 2-4	Eco	Diagonal	24	-7461.8	15100.9	219.3
Model 2-1	Plant	Full	15	-7570.5	15251.8	370.2
Model 2-2	Plant	Diagonal	14	-7654.1	15411.5	529.9
Model 2-0	None	Full	5	-7722.3	15481.6	600.0
Model 2-7	Preference	Diagonal	6	-7739.0	15522.4	640.8
Model 2-9	Preference + Performance	Diagonal	8	-7745.6	15550.4	668.7

Model 2-11	Performance	Diagonal	6	-7801.2	15646.7	765.1

Solanaceae

Table S6: Model selection on the Solanaceae dataset (*n* = 259). Models are ranked by increasing BIC (from best to worst). *k* is the number of parameters. *L* is the log-likelihood. Δ_{BIC} is the BIC difference between any focal model and the best one.

Model	Covariates	Residual matrix	*k*	*L*	BIC	ΔBIC
Model 1-6	Plant + Eco	Diagonal	42	-824.2	1881.7	0.0
Model 2-10	Preference + Performance + Eco	Diagonal	42	-829.0	1891.2	9.5
Model 1-5	Plant + Eco	Full	45	-823.8	1897.5	15.8
Model 2-12	Performance + Eco	Diagonal	39	-842.6	1901.7	20.0
Model 2-8	Preference + Eco	Diagonal	39	-846.5	1909.5	27.8
Model 1-2	Plant	Diagonal	12	-922.3	1911.2	29.4
Model 2-9	Preference + Performance	Diagonal	12	-924.2	1915.1	33.4
Model 2-11	Performance	Diagonal	9	-935.1	1920.2	38.5
Model 1-1	Plant	Full	15	-921.1	1925.6	43.8
Model 2-7	Preference	Diagonal	9	-938.2	1926.3	44.6
Model 1-4	Eco	Diagonal	36	-921.9	2043.8	162.0
Model 1-3	Eco	Full	39	-920.8	2058.1	176.4
Model 1-0	None	Full	9	-1016.8	2083.5	201.8

Other plant families

Table S7: Model selection on the dataset with other plant families. Models are ranked by increasing BIC (from best to worst). *k* is the number of parameters. *L* is the log-likelihood. Δ_{BIC} is the BIC difference between any focal model and the best one.

Model	Covariates	Residual matrix	*k*	*L*	BIC	Δ_{BIC}
Model 1-5	Plant + Eco	Full	98	-12218.5	25195.0	0.0

Model 1-6	Plant + Eco	Diagonal	92	-12269.6	25250.8	55.8
Model 1-1	Plant	Full	58	-12715.3	25879.1	684.1
Model 1-2	Plant	Diagonal	52	-12796.6	25995.3	800.3
Model 2-10	Preference + Performance + Eco	Diagonal	56	-12788.1	26009.3	814.3
Model 2-8	Preference + Eco	Diagonal	52	-12840.9	26083.9	888.9
Model 1-3	Eco	Full	54	-12872.4	26162.4	967.3
Model 2-12	Performance + Eco	Diagonal	52	-12921.4	26245.1	1050.0
Model 1-4	Eco	Diagonal	48	-12968.4	26308.1	1113.1
Model 2-9	Preference + Performance	Diagonal	16	-13650.6	27424.9	2229.8
Model 2-7	Preference	Diagonal	12	-13855.9	27804.7	2609.7
Model 1-0	None	Full	14	-13863.2	27834.7	2639.7
Model 2-11	Performance	Diagonal	12	-13905.2	27903.1	2708.1

1 *Host plants availability*

2 Figure S2: Availability of host plants along the year for seven fly species from Charlery de la

3 Masselière et al. (2017a, Supplementary material Figure S1). Only major host plants are reported.

4 Specialist species tend to rely on hosts with year-long availability, while generalists also have host

5 with short fructification periods in their diet.

1 *Appendix S3: Stability and robustness of results*

2 *Model diagnostics*

3 Model diagnostics were conducted using the R package DHARMa (Hartig, 2020; Figure S3 below).

Figure S3

Even on the most complete
model, an excess of zeros was
detected likely in relation
with fly species host ranges:
for any given fly species not
all 21 studied plants are
compatible hosts, therefore
introducing many zeros in the
complete 8-fly dataset. No
particular tendency of
residuals on predictors was
identified (plant shown here,
FAMD axes not shown).

To further explore potential spatial or temporal trends in model residuals, we studied the variogram

of residuals, i.e., how the estimated variance of the difference between two sites of this quantity

changes as a function of spatial distance of distance in sampling date (equation below, Figure S4):

$$V(i, i') = \mathbb{E}\left(\|\tilde{\mathbf{Z}}_i - \tilde{\mathbf{Z}}_{i'}\|^2 \right) = \mathbb{E}\left(\|\tilde{\mathbf{Z}}_i\|^2 + \|\tilde{\mathbf{Z}}_{i'}\|^2 \right) \approx \left\| \mathbf{M}_i + \frac{1}{2}\mathbf{S}_i^2 \right\|^2 + \left\| \mathbf{M}_{i'} + \frac{1}{2}\mathbf{S}_{i'}^2 \right\|^2$$

Correlations between the variance of residuals and spatial distance, difference in sampling month or

difference in year of sampling were weak (0.0433, -0.0049, and 0.0211, respectively).

Figure S4: Variogram of the residuals of the fullest model on the full 21 plant x 8 fly species dataset

(Model 1-5) with respect to spatial distance, difference in the month of sampling or difference in year

of sampling.

Lastly, we explored the relationship between observed and fitted species abundances (log scale) in

the best models on the full 21 plant x 8 fly species dataset (Figure S5, Model 1-5 on the left panel and

model 1-6 on the right panel). Data were relatively well fitted except that both models tended to

overestimate low abundance values.

Figure S5

BIC robustness in PLN models comparisons

3　To estimate the magnitude of the uncertainty associated to model selection, we have proceeded to

Figure S6

simulations by bootstrapping samples 50

times and conducting model comparison on

the bootstrapped datasets for all models of

Table 1 (no covariate, plant, ecological factors,

both with either full or diagonal residual

matrix on the full 21 plant x 8 fly species

dataset).

Stability of sparse structure in residual matrix

12　　The function PLNnetwork() allows digging into the residual variance-covariance matrix and infer

13　which species pairs still have with significant unexplained covariation in abundance after accounting

14　for covariates. In this approach, the residual matrix is formalized as a network, which nonzero edges

15　point to species pairs with significantly unexplained covariations. The function builds a series of 40

16　models with varying penalties over the number of edges in the resulting network (i.e., nonzero

Figure S7

residual covariances), and allows comparing

them using model selection criteria (Figure

S7). The stability of network edges, can be

evaluated using StARS (Liu et al. 2010), which

performs resampling to evaluate the

robustness of the network along the path of

solutions. It allows selecting networks with

varying stability criteria (Figures S8 and S9).

Figure S8

2   Figure S9: Various networks inferred. A. Network with the best BIC criterion. B. Network with a

3   stability of 0.95. C. Network with a stability of 0.75. Whatever the model selection criterion,

4   significant residual associations are found between the generalists *B. zonata* and *C. quilicii*, and

5   between the specialists *Z. cucurbitae* and *D. ciliatius*.

1 *Inferred realized host use using 36 host plants*

2 The full initial dataset contained 12872 from 40 host plants. Keeping only samples with GPS

3 coordinates, at least one individual fly and at least 10 samples per host plant led to a filtered dataset

4 with 6434 samples and 36 host plants, including the 21 ones further studied in the laboratory.

5 Conducting PLN modelling on the 36-plant datatset led to the following results.

Figure S10: Residual variance-covariance matrices obtained after PLN model fitting on species abundances. A) Without any covariate (Model 1-0). B) With ecological covariates (Model 1-3). C) With plant species as a cofactor (Model 1-1). D) With both plant species and ecological covariates (Model 1-5).

16 Table S8: Model selection on the extended dataset (n = 6434). Models are ranked by increasing BIC

17 (from best to worst). k is the number of parameters. L is the log-likelihood. Δ_{BIC} is the BIC difference

18 between any focal model and the best one.

Model	Covariates	Residual Matrix	k	L	BIC	Δ_{BIC}
Model 1-6	Plant + Eco	Diagonal	376	-35018.92	73335.11	0.00
Model 1-5	Plant + Eco	Full	404	-34955.09	73452.99	117.87
Model 1-2	Plant	Diagonal	296	-35933.11	74461.95	1126.83
Model 1-1	Plant	Full	324	-35869.94	74581.15	1246.04
Model 1-4	Eco	Diagonal	96	-47094.34	95030.54	21695.42
Model 1-3	Eco	Full	124	-48796.13	98679.67	25344.55
Model 1-0	None	Full	44	-50013.67	100413.20	27078.08

Figure S11: Realized host use as inferred from regression coefficients relative to host plants in the fullest model (Model 1-5 on the eight-species dataset). The orange rectangle contains plant-fly associations studied in the main text (Figure 3).

15 Figure S12: Species abundance responses to ecological variables inferred through regression

16 coefficients relative to Dim1 and Dim2, inferred under the fullest model on the 36 plant x 8 fly

17 species dataset (Model 1-5). Error bars represent approximate confidence intervals (1.96 x standard

18 errors).

Literature cited

Hartig, Florian. 2020. DHARMa: Residual Diagnostics for Hierarchical (Multi-Level / Mixed) Regression Models. R package version 0.3.3.0. https://CRAN.R-project.org/package=DHARMa

Liu, Han, Kathryn Roeder, and Larry Wasserman. 2010. "Stability Approach to Regularization Selection (Stars) for High Dimensional Graphical Models." In *Proceedings of the 23rd International Conference on Neural Information Processing Systems - Volume 2*, 1432–40. USA.