Title (12 words; 93 characters with spaces)

Bone marrow adipocytes drive the development of tissue invasive Ly6C[high] monocytes during obesity

Running Title: Bone marrow fat linked to Ly6C[high] monocytes in obesity

Parastoo Boroumand[1,2], David Prescott[3,4], Tapas Mukherjee[3], Philip J. Bilan[1], Michael Wong[1], Jeff Shen[1],

Ivan Tattoli[4], Yuhuan Zhou[1], Angela Li[3], Tharini Sivasubramaniyam[3], Nancy Shi[1], Lucie Y Zhu[1], Zhi Liu[1],

Clinton Robbins[3], Dana J. Philpott[3], Stephen E. Girardin[3,4], Amira Klip[1,2*].

[1]Cell Biology Program, The Hospital for Sick Children, Toronto, Ontario, M5G 0A4; Canada;

[2]Department of Biochemistry, University of Toronto, Toronto, Ontario, M5S 1A8; Canada;

[3]Department of Immunology, University of Toronto, Toronto, Ontario, M5S 1A8; Canada;

[4]Department of Laboratory Medicine and Pathophysiology, University of Toronto, Toronto, Ontario, M5S 1A8;

Canada

*Corresponding Author:

Amira Klip, Ph.D.

Cell Biology Program

Hospital for Sick Children

Toronto, Ontario M5G 0A4, Canada

Phone: 416-813-6392

E-mail: amira@sickkids.ca

Summary (150 Words, 150):

During obesity and high fat-diet (HFD) feeding in mice, sustained low-grade inflammation includes not

only increased pro-inflammatory macrophages in the expanding adipose tissue, but also bone marrow (BM)

production of invasive Ly6C^{high} monocytes. As BM adiposity also accrues with HFD, we explored the

relationship between the gains in BM white adipocytes and invasive Ly6C^{high} monocytes *in vivo* and

through *ex vivo* paradigms. We find a temporal and causal link between BM adipocyte whitening and the

Ly6C^{high} monocyte surge, preceding the adipose tissue macrophage rise during HFD. Phenocopying this,

ex vivo treatment of BM cells with conditioned media from BM adipocytes or from *bona fide* white

adipocytes favoured Ly6C^{high} monocyte preponderance. Notably, the switch towards the Ly6C^{high} subset

was preceded by monocyte metabolic reprogramming towards glycolysis, reduced oxidative potential and

increased mitochondrial fission. In sum, short-term HFD changes BM cellularity, resulting in local

adipocyte whitening driving a gradual increase and activation of invasive Ly6C^{high} monocytes.

Keywords:

Obesity; Immunometabolism, Bone marrow monocytes, Bone marrow adipocytes, Tissue invasive

monocytes, Ly6C^{high} monocytes, Glycolytic shift.

Introduction

A continuum of chronic obesity leading to metabolic disorders such as insulin resistance and eventually type 2 diabetes is of paramount concern worldwide (Centers for Disease Control and Prevention, 2017; Hamann, 2017; WHO, 2016). An important contributor to the development and exacerbation of insulin resistance is a gain in macrophages in adipose tissue (AT) (Liu et al., 2019; Weisberg et al., 2003), muscle (Fink et al., 2014; Garvey et al., 2008) and liver (Lanthier et al., 2010) as well as their enhanced pro-inflammatory (Lumeng et al., 2007, 2008; De Souza et al., 2005) and glycolytic switch (Sharma et al., 2020). This low-grade inflammation is partly explained by circulating monocyte infiltration into tissues where they differentiate to pro-inflammatory macrophages (Kohyama et al., 2016).

Ly6C[high] monocytes in mice, and classical CD14[++] CD16[-] monocytes in humans, constitute the predominant tissue-infiltrating monocyte subset during chronic inflammation. These tissue-invasive monocytes are classified by high surface abundance of the Ly6C protein, the chemokine receptor CCR2 and the adhesion molecule L-selectin, all of which allow for monocyte extravasation from the BM and infiltration into the tissues (Auffray et al., 2007; Boniakowski et al., 2018; Fink et al., 2014; Geissmann et al., 2003; Weisberg et al., 2006). With obesity, the development of Ly6C[high] monocytes along with their stem and progenitor cells in the BM is boosted (Griffin et al., 2018; Manicone et al., 2016; Singer et al., 2014) leading to a rise in available circulating Ly6C[high] monocytes for tissue-infiltration (Friedrich et al., 2019; Singer et al., 2014).

Various obesity-driven factors amalgamate for greater Ly6C[high] monocyte infiltration into the AT including (a) increased AT-dependent chemoattraction (Kamei et al., 2006; Nagareddy et al., 2014; Nov et al., 2013; Renovato-Martins et al., 2017) and (b) enhanced BM myelopoiesis (Griffin et al., 2018; Manicone et al., 2016; Singer et al., 2014). However, it is not clear whether the obesity-induced BM myelopoiesis occurs prior to, concurrently, or after the elevation of AT macrophages. Since the BM is a highly-vascularized tissue, home to hematopoietic and mesenchymal lineages, and is dynamically reactive to environmental changes, we reasoned it might be an early responder to HFD-induced obesity, producing increased Ly6C[high] monocytes prior to their accumulation in the AT. Further, since BM-resident adipocytes

undergo hyperplasia and hypertrophy during obesity (Boroumand and Klip, 2020; Doucette et al., 2015;

Scheller et al., 2015; Tratwal et al., 2020) we contemplated that this process might be temporally and

potentially causally related to the HFD-driven BM myelopoiesis and Ly6C^{high} monocytosis. Finally, we

considered whether the response of the monocytic lineage to the BM environment during obesity includes

a metabolic shift leading to the increase in Ly6C^{high} monocytes. In fact, and surprisingly, little is known

about the metabolic orientation of the Ly6C^{high} monocytes in states of obesity, although there is mounting

evidence of a link between a macrophage-intrinsic metabolic shift and their pro-inflammatory orientation

in this condition (O'Neill et al., 2016; Sharma et al., 2020; Thapa and Lee, 2019; Verdeguer and Aouadi,

2017).

Hence, the overall hypothesis driving this study is that, with obesity, the increased production of

Ly6C^{high} monocytes in the BM is accompanied and possibly caused by a glycolytic shift in these cells,

driven by changes in bone marrow adipocytes (BMA). Accordingly, we investigated the temporality of

enhanced BM myelopoiesis, BM Ly6C^{high} monocyte accumulation, and AT macrophage accrual during

high fat diet (HFD) feeding in mice. Next, we explored the potential for a causal relationship between the

obesity-induced changes in BMA and increased Ly6C^{high} monocyte production. Using an *in vitro* platform,

we deconstructed the effects of obesity-induced BM cellular changes on Ly6C^{high} monocyte expansion. We

report that BM Ly6C^{high} monocytes accumulate in the BM prior to the gain in pro-inflammatory AT

macrophages. The rise in BM monocytes temporally correlated with the elevation in BMA. Moreover,

BMA whitening correlated with the timely increases in BM Ly6C^{high} monocytes, and remarkably, isolated

BMA from HFD-fed mice directly prompted Ly6C^{high} monocyte development *ex vivo*. This monocyte

developmental cue was also induced *ex vivo* by *bona fide* white adipocytes, while *bona fide* brown

adipocytes favoured Ly6C^{low} monocyte predominance. Noteworthy, the white adipocyte-induced swap in

proportion of Ly6C^{low} relative to Ly6C^{high} monocytes was preceded by a switch in monocyte metabolism

from oxidative to glycolytic. Mechanistically, brown adipocytes increased Ly6C^{low} monocyte proliferation

while white adipocytes appear to promote Ly6C^{low} monocyte conversion and potentially progenitor

expansion to increase Ly6C^{high} preponderance. We propose that HFD-induced changes in BMA disrupts

the normal balance of BM monocyte subsets leading to a higher output of tissue invasive Ly6C^{high}

monocytes.

Results

Increased fasting blood glucose, peripheral and BM adiposity by 3 weeks of HFD

To study the temporal relationship of HFD-induced myeloid changes in the BM relative to that in epididymal white AT (EWAT), C57BL/6 mice were fed HFD or control low fat diet (LFD) for either 3, 8 or 18 weeks. HFD-fed mice had higher body weight and impaired fasting blood glucose by 3 weeks, compared to controls (**Figure 1A,B**). Body weight and hyperglycemia continued to rise with prolonged feeding. HFD-fed mice also displayed higher EWAT weight (**Figure 1C**) and inter-scapular brown AT (BAT) weight at 3 weeks (not shown). Cross-sectional tibia and femur bone histology visualized via H & E and Oil Red O staining revealed an increase in BMA size by 3 weeks and an increase in BMA number by 8 weeks of HFD (**Figure 1D-G**). This indicated that 3 weeks of HFD sufficed to induce whole body metabolic disturbance and augmented adiposity in peripheral tissues and in the BM.

BM monocytes increase prior to accumulation of EWAT macrophages and monocytes with HFD

Although total EWAT weight increased by 3 weeks, myeloid cells in the stromal vascular fraction (SVF) of AT were not elevated until 8 weeks of HFD, as revealed by gene expression of the general myeloid marker *Itgam* (encoding CD11b) and monocyte markers $Gr1^- CD115^+ CD11b^+ Ly6C^+$ (determined by flow cytometry) (**Figures 2A,D**). Similarly, AT macrophages, assessed by gene expression of the AT macrophage marker *Adgre1* (encoding F4/80) and by flow cytometry counting of $CD11b^+ Ly6C^- F4/80^+$ macrophages in the SVF were not elevated at 3 weeks of HFD but were at 8 weeks (**Figure 2B,E**). Further, gene expression of the pro-inflammatory myeloid marker *Itgax* (encoding CD11c) and surface detection of the pro-inflammatory markers $CD11b^+ Ly6C^- F4/80^+ CD11c^+$ were also higher in the SVF by 8 weeks of HFD relative to control diet, but not at 3 weeks (**Figure 2C,F**). Of note, gene expression of the embryonically-conserved tissue-resident AT macrophage markers *Timd4* and *Marco* (Beattie et al., 2016; Gibbings et al., 2015) was not elevated at 3 or at 8 weeks and only rose at 18 weeks of HFD (**Figure 2G-H**). This indicates that the gain in AT macrophages at 8 weeks of HFD is unlikely due to proliferation of resident macrophages and instead likely results from monocyte recruitment from the circulation.

In the BM, total monocytes, gated via the $Gr1^-$ $CD115^+$ $CD11b^+$ $Ly6C^+$ markers, increased at 3

weeks and continued to rise by 8 and 18 weeks of HFD (**Figure 2I**). Using the surface abundance of the

Ly6C protein as well as the CX3CR1 and CCR2 markers (Auffray et al., 2007, 2009; Kratofil et al., 2017),

we subcategorized $Ly6C^{low}$ and $Ly6C^{high}$ monocyte subsets. The two populations were identified by FACS

analysis using the gating strategy illustrated in Figure S1. At 3 weeks into the feeding study, $Ly6C^{low}$ and

$Ly6C^{high}$ monocytes represented about 40% of the total BM monocyte population in the control diet group,

whereas HFD promoted a higher proportion of $Ly6C^{low}$ monocytes (**Figure 2J**). However, by 8 weeks of

HFD there was a marked shift, whereby the relative proportion of $Ly6C^{low}$ monocytes in the total monocyte

population was *lower* and that of $Ly6C^{high}$ monocytes was *higher* than under the control diet (**Figure 2J**).

These differences were more prominent by 18 weeks of HFD feeding (**Figure 2J**). Collectively, these

findings revealed that, in the BM, HFD favoured monocytosis at 3 weeks, with a subsequent and prominent

switch towards tissue invasive $Ly6C^{high}$ monocytes at 8 weeks that continued into 18 weeks of HFD. Singer

et al. also demonstrated an increase in BM monocytosis and elevated monocyte progenitor cells with 18

weeks of HFD, that persisted after BM transplantation into chow-fed recipients (Singer et al., 2014). In the

present study, while we clearly recorded an earlier and continuing rise in total monocytes in HFD-fed mice

(**Figure 2I**), we did not observe statistically significant increases in monocyte progenitor and stem cells at

any time point analyzed (**Figure S2A-C**).

Glycolytic metabolic reprograming in monocyte subsets with HFD

Recent discoveries have highlighted the importance of glycolytic metabolism in the activation of

AT macrophages with HFD (Lee and Murphy, 2019), but the metabolic status of monocytes in obesity has

not been reported, although some studies have investigated the metabolism of circulating human

$CD14^+CD16^-$ monocytes (homologous to murine tissue invasive $Ly6C^{high}$ monocytes) during aging (Lee

and Murphy, 2019; Pence and Yarbro, 2019). To assess whether HFD affects the metabolic status of BM

monocyte subsets in obesity we analyzed $Ly6C^{low}$ and $Ly6C^{high}$ from control and HFD mice at 3, 8 and 18

weeks of feeding. The mitochondrial activity-indicator dye MitoTracker Red CMXRos was incorporated

into the monocyte flow cytometry panel. MitoTracker Red CMXRos is recruited to the mitochondrial membrane proportionally to the degree of mitochondrial membrane potential and is thus an indicator of mitochondrial activity in oxidative phosphorylation. The results revealed that Ly6C^{low} monocytes have a higher mitochondrial membrane potential than Ly6C^{high} monocytes in the control mice (**Figure 3A-C**). Strikingly, 3 weeks of HFD reduced the mitochondrial membrane potential in both monocyte subsets indicated by the left-ward shift in the MitoTracker Red CMXRos peak (**Figure 3A**). The HFD-induced effects were more pronounced by 8 and 18 weeks of HFD (**Figure 3B,C**). To further understand the effects of HFD on the developing BM monocytes, Ly6C^{low} and Ly6C^{high} monocytes from mice fed control or HFD for 3, 8 or 18 weeks were isolated via fluorescent activated cell sorting (FACS) and plated for metabolic analysis with a Seahorse XF Analyzer. The results revealed a gradual lowering in the oxygen consumption rate (OCR) of Ly6C^{low} monocytes with HFD (**Figure 3D-I**). Comparison of the area under the curve (AUC) revealed that at 18 weeks, the OCR of the Ly6C^{low} monocytes was significantly lowered with HFD (**Figure 3I**). Interestingly, at 8 and 18 weeks, the AUC of the OCR of the Ly6C^{low} monocytes was significantly higher than the AUC of the OCR of Ly6C^{high} monocytes in the control group (**Figure 3H-I**). Moreover, this difference was lost in the HFD group as the AUC of the OCR of Ly6C^{low} monocytes diminished (**Figure 3H-I**).

Comparison of the various sections of the OCR graph revealed that the changes in OCR observed with 3 weeks of HFD were not manifest at basal respiration nor at the ATP-dependent respiration, rather they were evident in the maximal respiratory capacity portion of the metabolic stress test (**Figure 3D**). A drop in basal respiration of both monocyte subsets emerged with 8 weeks of HFD and became more pronounced with 18 weeks of HFD (**Figure 3E-F**). This showcased the sustained effects of HFD on monocyte metabolic status. Conversely, the extracellular acidification rate (ECAR) of Ly6C^{low} monocytes was higher than that of Ly6C^{high} monocytes, consistent with a more active metabolism in the former (**Figure 3J**). However, this difference was lost with HFD, as ECAR dropped in the Ly6C^{low} monocytes and rose in the Ly6C^{high} monocytes relative to the control diet group (**Figure 3J**). Overall, these results suggest that,

along with promoting the preponderance of Ly6C^{high} monocytes, HFD also lowers OCR in Ly6C^{low}

monocytes.

Endorsing these findings, an analysis of the BM revealed elevated lactate levels across all HFD

time points (**Figure 3K**). This increase in lactate may be due to the glycolytic shift in BM monocytes

described above, among other potential sources within the BM cellular composition. Further analysis of the

BM soluble fraction failed to evince any changes between control and 18-week HFD-fed mice in pro-

inflammatory cytokines TNFα, IL-1β, IL-6, CXCL1 or CCL2 (**Figure S3A-C**). Similarly, there were no

differences between diet groups in pro-inflammatory gene expression of *IL-6*, *IL-1β* or *Itgax* in FACS-

sorted Ly6C^{low} and Ly6C^{high} monocytes (Not Shown). However, *TNFα* gene expression was higher in

FACS-sorted Ly6C^{high} monocytes from 18-week HFD-fed mice compared to controls (**Figure S3D**).

Collectively, these findings reveal that HFD alters the BM metabolic micro-environment with little effect

on pro-inflammatory cytokine secretion (based on the panel tested).

Expression of mitochondrial fusion and fission genes in monocytes is altered by HFD

The efficiency of mitochondrial function is largely reliant on its structure. Mitochondria are highly

dynamic in structure, largely regulated by concentration of proteins that control fission and fusion (Gottlieb

and Bernstein, 2017). Dynamic transitions are closely aligned to metabolic performance. To further assess

the mitochondrial activity of monocytes, we enquired whether their expression of mitochondrial dynamic

genes is altered with HFD compared to the control group.

The analyzed genes involved in mitochondrial fusion were *Mfn1*, *Mfn2*, *Opa1*, *Tomm20* and

Tomm40, while those involved in mitochondrial fission were *Drp1*, *CypD* and *Fis1*. BM monocytes from

mice fed HFD for 3 weeks did not present with differences in gene expression in the tested mitochondrial

fusion genes compared to control diet counterparts (**Figure 4A**). BM monocytes isolated from mice fed

HFD for 8 weeks showed downregulated *Mfn1* and *Mfn2* compared to the controls (**Figure 4B**). BM

monocytes isolated from 18-week HFD-fed mice downregulated the mitochondrial fusion genes *Mfn2*,

Opa1, and *Tomm20* (**Figure 4C**). Furthermore, from the tested mitochondrial fission genes, *Drp1* was

upregulated in the BM monocytes of 3 weeks HFD-fed mice (**Figure 4D**). *Drp1* and *CypD* were also

upregulated in BM monocytes of 8 weeks HFD-fed mice (**Figure 4E**). At 18 weeks, all three tested

mitochondrial fission genes, *Drp1*, *CypD* and *Fis1* were upregulated (**Figure 4F**). This gene expression

pattern suggests that HFD changes the monocyte gene expression to favour downregulation of genes

involved in mitochondrial fusion and upregulation of genes involved in fission.

Given that Ly6C^{low} and Ly6C^{high} monocytes presented with differences in metabolic preference

(glycolytic and oxidative, respectively), and the known link of glycolytic metabolism with mitochondria

fragmentation and conversely oxidative metabolism to mitochondrial fusion, we assessed whether the

expression patterns of the genes governing mitochondrial fission and fusion would correlated distinctly

with the BM monocyte subset abundance in control and HFD-fed mice. Indeed, and notably, the abundance

of the Ly6C^{high} monocyte subset negatively correlated with *Mfn2* and positively correlated with *Drp1* levels

(**Figure 4G-H**). In contrast, the abundance of BM Ly6C^{low} monocytes did not correlate with the expression

patterns of either *Mfn2* or *Drp1* (**Figure 4G-H**). These data are consistent with our metabolic findings

suggesting Ly6C^{high} monocytes are more glycolytic and adopt a gene expression pattern that would promote

mitochondrial fragmentation.

A shift in BMA type at 8 weeks of HFD influences monocyte subset preference and mitochondrial

dynamic gene expression

Hematopoietic myeloid cells reside in specific niches of the BM that determine their cellular

interaction with other co-resident BM cells (Boroumand and Klip, 2020). BMA, while representing a unique

type of adipocytes, respond to obesity with an increase their white adipocyte-like fat storing properties and

a decrease in brown-like characteristics (Ambrosi et al., 2017; Doucette et al., 2015; Krings et al., 2012;

Scheller et al., 2016). To explore the temporal relation of BMA whitening in relation to the development

of monocyte subsets, we followed the gene expression of a suite of recently identified markers that are good

differentiators of adipocyte types (Nascimento et al., 2014; Ussar et al., 2014) in isolated BMA from femur

and tibia of control and HFD-fed mice. By 3 weeks into HFD, the white adipocyte marker Asc1 (encoding

the guanine nucleotide-binding protein subunit beta-like protein) was significantly higher and the brown

adipocyte marker P2rx5 (encoding purinergic receptor P2X) trended to be lower albeit without reaching

statistical significance, relative to control diet. White adipocyte markers *Lep* (leptin) and *AdipoQ*

(adiponectin) as well as the brown adipocyte markers *Ucp1* (UCP1) and *Pat2* (Integrin alpha pat-2) did not

change (**Figure 5A**). By 8 weeks of HFD, *Asc1* mRNA remained elevated and *Lep* and *AdipoQ* trended in

the same direction (**Figure 5B**). Brown adipocyte markers *P2rx5* and *Ucp1* in the isolated BMA of HFD

mice were markedly lower relative to those from control diet mice (**Figure 5B**). Thus, 8 weeks of HFD

sufficed to shift BMA towards white adiposity. By 18 weeks of HFD, all white adipocyte markers genes

showed were upregulated while brown adipocyte markers were downregulated compared to the control

group (**Figure 5C**). Analyses of adipocyte differentiation and progenitor cell markers were conducted in

the BM pelleted fraction. *Hoxc8*, a regulator of adipocyte progenitors (Nakagami, 2013), was significantly

upregulated at 8 weeks along with *Hoxc9* at 18 weeks of HFD (**Figure S4B,C**). The beige adipocyte

progenitor marker *Tmem26* was not significantly higher at any of the tested HFD durations (**Figure S4A-**

C). Expression of the brown adipocyte determinant genes *Tbx1* and *Lhx8* was undetectable in the isolated

BMA. Thus, during the course of HFD, while markers for white adipocyte progenitor cells were

upregulated, precursors of beige/brown adipocytes were not affected.

The rise in BM Ly6C^{high} monocytes and the shift in BMA whitening both manifested by 8 weeks

of HFD. Correlative analysis was performed for both Ly6C^{low} and Ly6C^{high} with the white adipocyte marker

Asc1 and the brown adipocyte marker *Ucp1*. The results revealed a significant positive correlation (P value

of 0.0123) for Ly6C^{high} monocytes with *Asc1* gene expression (**Figure 5D**). Conversely, the surge in

Ly6C^{high} monocytes presented a significant negative correlation (P value of 0.0415) relative to *Ucp1* gene

expression (**Figure 5E**). Changes in Ly6C^{low} monocytes did not significantly correlate with *Asc1* (P value

of 0.0768) or *Ucp1* (P value of 0.1299) (**Figure 5D-E**). Collectively, the above data suggest the possibility

that the mere shift in BMA towards white adiposity may directly favour induction of BM Ly6C^{high}

monocytes. To test this hypothesis, we generated conditioned media (CM) from BMA (BMA-CM) pooled

from three control and three HFD-fed mice, collected 3, 8 and 18 weeks into the diets. In parallel, BM of

control mice was used to generate monocytes through a 5-day *in vitro* differentiation protocol (as reported

by (Francke et al., 2011), see Methods). These monocytes were treated with BMA-CM from control and

HFD-fed mice for 18 h thereafter. Subsequently, the monocyte population was analyzed via flow cytometry.

BMA-CM from 3 weeks HFD-fed mice had no effect on the monocyte subset proportions compared to

BMA-CM from control diet-fed mice (**Figure 5F**). Strikingly, however, BMA-CM from 8-week HFD-fed

mice applied to BM-derived monocytes from control mice resulted a significant production of Ly6C^{high}

monocytes compared to the effect of BMA-CM from mice on the 8-week control diet (**Figure 5G**). This

skewing was further enhanced when exposing the control BM cells to BMA-CM from 18-week HFD-fed

mice (**Figure 5H**).

256   Lastly, we tested whether BMA-CM was capable to induce changes in gene expression of the

mitochondrial fusion and fission genes in monocytes, as observed with monocytes isolated from HFD-fed

mice. Accordingly, *in vitro*-generated monocytes were treated with BMA-CM from control and HFD-fed

mice. As before, the analyzed genes involved in mitochondrial fusion were *Mfn1*, *Mfn2*, *Opa1*, *Tomm20*

and *Tomm40*, while the tested genes involved in mitochondrial fission were *Drp1*, *CypD* and *Fis1*.

Monocytes treated with BMA-CM from HFD-fed mice for 18 weeks showed significant downregulated

mitochondrial fusion genes *Mfn2*, *Tomm20* and *Tomm40* (**Figure 5I**). Conversely, the mitochondrial fission

genes *Drp1* and *CypD* were significantly up-regulated in monocytes treated with HFD-derived BMA-CM

(**Figure 5J**).

265   These results not only show that HFD-derived BMA-CM can bias monocyte subset cellularity but

266   also can alter mitochondrial morphology via changes in mitochondrial dynamic gene expression. These

267   changes can have consequential effects on the metabolic properties on the monocytes.

268   **White adipocytes favour Ly6C^{high} monocytes and promote mitochondrial fragmentation**

269   To assess whether the effects of BMA on monocyte subset skewing and metabolic differences can

be attributed to BMA whitening, we designed a proof-of-concept experiment studying the influence of *bona*

fide white or brown adipocytes on monocyte subset determination and metabolic changes *ex vivo*. Isolated

adipocytes from EWAT and brown adipocyte tissue (BAT) of three mice were pooled to generate white AT

adipocytes (WATA) and brown AT adipocytes (BATA), respectively. Confirming the phenotype of the

isolated cells, classical white adipocytes markers were solely expressed in the isolated WATA, and classical

brown adipocyte markers were exclusively expressed in the BATA (**Figure S4D-E**). The isolated WATA

and BATA were cultured for 24 h and the corresponding CM were collected for subsequent treatment of

monocyte preparations. *In vitro*-generated BM-derived monocytes from chow-fed mice were treated with

WATA-CM, BATA-CM or left untreated. The proportion of Ly6C^{low} and Ly6C^{high} monocyte subsets was

determined by flow cytometry analysis (**Figure S1**). This *in vitro* paradigm allowed for direct mechanistic

analysis of the influence of white or brown adipocytes on monocytes. These results indicate that WATA-

CM favoured Ly6C^{high} monocyte predominance, while BATA-CM prompted Ly6C^{low} monocyte abundance

(**Figure 6A-B**), revealing a novel, differential influence of white and brown adipocytes on monocyte

orientation preference. Similarly, CCR2 surface receptors also increased in the monocytes incubated with

WATA-CM (Not Shown), validating the effects of white adipocytes in promoting an abundance of tissue

invasive Ly6C^{high} monocytes.

Since segregated Ly6C^{low} and Ly6C^{high} monocyte subsets had different metabolic orientations, we

investigated if the differential adipocyte influence on monocyte subset development corresponded to a

change in their metabolic status. To this end, *in vitro*-generated monocytes were treated with WATA-CM,

BATA-CM or left untreated for the last 18 h of the monocyte differentiation protocol. The mitochondrial

membrane potential was assessed using MitoTracker Red CMXRos and flow cytometry analysis.

Remarkably, Ly6C^{low} monocytes incubated *in vitro* with WATA-CM experienced a left-ward shift in

MitoTracker Red CMXRos peak revealing a reduction in mitochondrial membrane potential induced by

the WATA-CM (**Figure 6C**), reproducing what was observed with monocytes from HFD-fed mice (see

Figure 3A-C). Furthermore, the gene expression of mitochondrial fusion genes *Mfn1*, *Mfn2*, *Tomm20* and

Tomm40 were reduced in monocytes treated with WATA-CM (**Figure 6D**). The mitochondrial fission

genes *Drp1*, *CypD* and *Fis1* were up-regulated in monocytes treated with WATA-CM (**Figure 6E**).

Conversely, treating monocytes with BATA-CM downregulated the mitochondrial fission gene *CypD*

(**Figure 6E**). To explore if the changes in mitochondrial fusion and fission gene expression correlated

with the corresponding mitochondrial morphology, we utilized super-resolution microscopy to image the

mitochondrial protein TOM20 in *in vitro* monocytes adhered to coverslips.

WATA-CM induced a striking increase in the percentage of cells with fragmented mitochondria,

from 22% in control CM-treated cells to 64%. On the other hand, BATA-CM raised mitochondrial

elongation from 20% in CM-treated cells to 55% (**Figure 6F-H**). These observations suggest that white

adipocyte products alter the metabolic status of Ly6C^{low} monocytes by reducing their mitochondrial

membrane potential and altering mitochondrial dynamic gene expression and morphology, thereby

linking changes in BMA whitening to monocyte metabolic status. Moreover, the shift to mitochondrial

fragmentation coincided with the higher abundance Ly6C^{high} monocytes resulting from in vitro-generated

monocytes with WATA-CM (**Figure 6A,B**).

White and brown adipocytes influence monocytes through distinct mechanisms

Monocyte precursor cells in the BM in the tissues maintain cell numbers by cell proliferation

(Cassado et al., 2015). Although we had not detected HFD-induced changes in hematopoietic stem and

monocyte progenitor cells *in vivo*, we assessed the abundance of these progenitor cell populations upon

exposing *in vitro*-generated monocytes to WATA-CM and BATA-CM. We reasoned that the *ex vivo*

experimental setup lent itself well to gain understanding of the mechanisms whereby monocytes undergo

changes in subset proportion in response to *bona fide* white and brown adipocyte media. The abundance of

hematopoietic stem and monocyte progenitor cells was determined using appropriate surface markers and

flow cytometry analysis (**Table 1, Figure S1D**). A small gain in myeloid progenitor cell (CMP) was

observed in response to WATA-CM treatment (**Figure S5A-C**). However, no differences in the abundance

of CD117+, Sca1+, Sca1-, and GMP cells were detected (**Figure S5A-C**). GMP cells are downstream of

CMP during lineage progression of progenitor cells towards monocytes or erythroid cells (Yáñez et al.,

2017), and so it is noteworthy these cells did not become more abundant with WATA-CM treatment, even

though content of CMP, their upstream multipotential progenitors, was slightly elevated.

Next, we asked whether proliferation might contribute to the higher proportion of terminally

differentiated Ly6C^{high} or Ly6C^{low} monocyte subsets observed following treatment with WATA-CM (in

addition to the conversion of Ly6C^{low} into Ly6C^{high}) or BATA-CM, respectively. To this end, we treated

BM-derived *in vitro*-generated monocytes with WATA-CM, BATA-CM or left them untreated for 18 h,

with or without the presence of bromodeoxyuridine (BrdU), an indicator of S-phase. Using flow cytometry

analysis, DNA-incorporated BrdU was detected in the Ly6C^{low} or Ly6C^{high} monocyte gates of the CD11b

versus Ly6C plots, with BrdU as the third colorimetric axis. Although at the highest levels the CD1b signal

created a green background signal in the BrdU channel (as seen in samples not treated with BrdU), the

actual BrdU signal could be readily distinguished by the yellow to red signal corresponding to the illustrated

gradient scale shown. Proliferating cells were quantified as percentage of CD11b^{+} Ly6C^{+} cells that were

positive for BrdU incorporation (yellow to red signal). WATA-CM-treated monocytes rendered the

expected increase in Ly6C^{high} monocytes, yet these cells did not show any significant incorporation of BrdU,

implying that they are not actively replicating their DNA. In contrast, BATA-CM increased DNA-

incorporated BrdU in the sorted Ly6C^{low} monocyte subset, visualized as yellow to red hues in the CD11b^{+}

Ly6C^{low} cell gate (**Figure 7A**), and the number of proliferating cells was elevated by 25-fold (**Figure 7B**).

These findings revealed that brown adipocytes promote an abundance of Ly6C^{low} monocytes by inducing

their proliferation. Interestingly, this replicative behavior of BATA-CM-treated of Ly6C^{low} monocytes,

which we showed above are more oxidative than Ly6C^{high} monocytes, is consistent with recent findings

that proliferative cells have higher glucose consumption and oxidative phosphorylation relative to quiescent

cells (Yao et al., 2019).

To buttress the above results, we also assessed the colony forming potential of *in vitro*-generated

monocytes that received pre-treatment of WATA-CM or BATA-CM, to assess the potential for expansion

of progenitor cells present in the samples. Colonies were identified as BFU-E (giving rise to erythroid cells);

CFU-GEMM (giving rise to large mixed cultures of granulocyte, erythroid, macrophage, megakaryocyte;

also, known as CMP); CFU-G (giving rise to granulocytes) or CFU-M (giving rise to macrophages). BATA-

CM promoted growth of CMP/CFU-GEMM cultures over control Media (**$p<0.01$ Two-way ANOVA,

Tukey's multiple comparisons, **Figure 7C**) and biased granulocyte/macrophage progenitors (widely known

as GMP) towards macrophage over granulocyte differentiation, relative to control Media (**$p<0.01$) or

WATA-CM (*$p<0.05$ or ***$p<0.001$, **Figure 7C**). The total numbers of colonies that grew after 7-10 days

of culturing of each pre-treated cohort of monocytes were not different across the three treatments, although

trended upwards with BATA-CM (**Figure 7D**). These results revealed that while BATA-CM promoted

expansion of selected populations (CMP/CFU-GEMM and CFU-M), consistent with the increase in BrdU

incorporation shown above, WATA-CM was without effect relative to control Media.

The above findings suggest that while the proportional increase in Ly6C^{low} monocytes induced by

BATA-CM involves cell proliferation, the proportional increase in Ly6C^{high} monocytes induced by WATA-

CM does not. As a complementary approach, BM cells were analyzed by flow analysis for the presence of

monocyte progenitors within the Ly6C^{low} or Ly6C^{high} monocyte subsets. GMP progenitor cells were

essentially the only progenitors detected by this approach in the Ly6C^{low} monocyte subset, and they

represented 140 GMP per 1000 Ly6C^{low} cells (**Figure 7E**). During the incubation time of 18 h with

conditioned medium, we anticipate the progenitors could theoretically undergo only one doubling and

therefore unlikely to account for the overall changes in Ly6C^{low} cell numbers produced by WATA-CM.

The above findings indicate that WATA-CM treatment did not result in any appreciable expansion

of progenitor cells or colony formation. Accordingly, alternative mechanisms had to be explored that might

contribute to the WATA-CM induced shift towards Ly6C^{high} monocyte preponderance. It is known that

Ly6C^{high} monocytes give rise to Ly6C^{low} monocytes in mouse blood and BM, and this is paralleled by the

human homologous CD14 and CD16 monocyte subsets (Kratofil et al., 2017; Lessard et al., 2017; Patel et

al., 2017). However, while there has been speculation about the conversion of Ly6C^{low} to Ly6C^{high}

monocytes, there is little experimental evidence for this phenomenon. Hence, we contemplated the

possibility that BATA-CM may also cause Ly6C^{high} to Ly6C^{low} monocyte conversion, and especially

whether WATA-CM might cause $Ly6C^{low}$ to $Ly6C^{high}$ monocyte conversion, however unprecedented. To

this end, we collected each FACS-sorted, homogenous $Ly6C^{low}$ and $Ly6C^{high}$ monocyte populations and,

once isolated, treated them with WATA-CM, BATA-CM, or control media for 18 h. Following treatments,

we evaluated the presence of the alternative monocyte subtype relative to that in the respective starting

homogenous populations. Surprisingly, no conversion from $Ly6C^{high}$ (red) to $Ly6C^{low}$ (gray) monocytes

was observed in any condition (**Figure 8A-F**), but, remarkably, there was a switch of the FACS sorted

$Ly6C^{low}$ population, as $Ly6C^{high}$ monocytes became more prevalent following exposure to WATA-CM

(their proportion accounting to 49.7% of monocytes) (**Figure 8B,E**). This was in stark contrast to all other

isolated cell and CM treatment combinations tested, which gave rise to a maximum of 16.3% of the opposite

monocyte subset compared to the starting population. Taken together, the results from Figures 7 and 8

endorse the novel possibility of $Ly6C^{low}$ into $Ly6C^{high}$ monocyte conversion induced by secreted factors

from white adipocytes.

Discussion

In this study, we demonstrate a temporal correlation between HFD-induced whitening of BMA and a BM myeloid lineage bias towards Ly6C^{high} monocytes *in vivo*. The latter present a more glycolytic metabolism underpinned by a reduction in mitochondrial membrane potential, a rise in mitochondrial fission genes and a reduction in mitochondrial fusion genes. Causatively, we show that CM from BMA of HFD-fed mice promotes the generation of Ly6C^{high} monocytes *ex vivo.* Phenocopying the effect of BMA derived from HFD-fed mice, *bona fide* WATA favoured a Ly6C^{high} monocyte preponderance, while BATA promoted a dominance of Ly6C^{low} monocytes. Interestingly, these differential influences by the respective adipocytes are induced by distinct cellular pathways. Mechanistically, white adipocytes induced a shift in monocyte subset proportion, from Ly6C^{low} to Ly6C^{high} monocytes suggesting a novel conversion of one subset into the other, while brown adipocytes favoured Ly6C^{low} predominance by inducing their proliferation. These changes in monocytes were underpinned by alterations in monocyte metabolism, as WATA blunted the monocytes' mitochondrial membrane potential and OCR. Overall, our findings reveal a regulation of monocyte subsets directly influenced by adipocytes, whether from the BM or from AT. Along with the observation of dynamic whitening of BMA during HFD, our results contribute to the emerging evidence that BMA are not mere space fillers, rather they are important contributors to cellular metabolism and hematopoietic lineage decisions (Boroumand and Klip, 2020). That these changes occur early on in the course of diet-induced obesity suggests that the BM is an early responder to a change in feeding status and that adipocytes play a significant role in directing monocyte subset skewing. Moreover, they reveal novel mechanisms regulating the genesis of such skewing.

BM as an early sensor and an active contributor to HFD-induced AT macrophage accumulation

It is now widely recognized that many factors contribute to the accumulation of macrophages in AT that includes the contribution of monocytes. There is mounting evidence for both HFD-induced EWAT monocyte chemo-attraction and for enhanced BM myelopoiesis, that work together to increase the monocyte infiltration and pro-inflammatory AT macrophage accumulation (Singer et al., 2014; Trottier et

al., 2012 and Fig. 1D-H). Singer et al. demonstrated that transplantation of BM from 18-week HFD-fed

mice into control recipients increased the number of invasive circulating monocytes and pro-inflammatory

AT macrophages (Singer et al., 2014). These pioneering findings heralded the notion that HFD influences

hematopoietic myeloid bias in the BM upstream of monocyte tissue invasion, thereby contributing to

downstream systemic and peripheral tissue inflammation. Furthermore, HFD-driven chemoattracting

signals from the homeostatically-disrupted EWAT initiate tissue influx and depletion of invasive Ly6C^{high}

monocytes from the circulation, which in turn may acts as a signal for their increased production in the BM

(Kamei et al., 2006; Nagareddy et al., 2014; Nov et al., 2013; Renovato-Martins et al., 2017).

In the present study, by analyzing chronological changes in EWAT and BM myeloid composition,

we observed that total BM monocytes increase by 3 weeks of HFD, whereas the proportion of Ly6C^{high}

monocyte predominance was evident only at 8 weeks coinciding with a gain in AT macrophages that were

also pro-inflammatory oriented. We did not observe statistical significance in HFD-dependent changes in

myeloid progenitors, likely due to their scarcity since HSC make up only 0.01% of the entire BM nucleated

cells (Challen et al., 2009). The rise in total monocytes observed at 3 weeks of HFD and prior to elevation

of AT macrophages identifies the BM as an earlier-responding *locus* during HFD. Interestingly, the increase

was only apparent in Ly6C^{low} patrolling monocytes and not in the Ly6C^{high} tissue invasive monocyte subset

that is known for tissue infiltration in chronic obesity. This might be explained by an initial non-specific

upregulation of total myelopoiesis and an increased Ly6C^{high} monocyte BM extravasation. Therefore, it

appears that only Ly6C^{low} monocytes are elevated in the BM at 3 weeks of HFD. By 8 weeks of HFD, we

observed a shift in the monocyte subset proportion, favouring Ly6C^{high} monocytes. Thus, by 8 weeks, the

BM has adopted the 'typical' HFD-induced hematopoietic phenotype, in so far as it already shows the

increased Ly6C^{high} monocytes previously reported at the habitually studied 16-18 weeks of HFD.

Importantly, we detected a gain in pro-inflammatory AT macrophages by 8 weeks of HFD that is concurrent

with the predominance of invasive Ly6C^{high} monocytes in the BM. In contrast, the embryonic-derived tissue

resident AT macrophage markers, *Timd4* and *Marco* (Beattie et al., 2016; Gibbings et al., 2015), were not

elevated until 18 weeks of HFD. This brings to light the importance of the contribution of the BM-derived

myeloid cell pool to early EWAT pathology and AT macrophage accumulation.

Changes in BMA influence monocyte subset skewing

BMA are distinctive components of the BM cellular suite that is sensitive to HFD effects (Pham et

al., 2020). In lean mice, BMA present characteristics that are similar to those of beige or brown adipocytes

in peripheral AT (Cuminetti and Arranz, 2019). With HFD, BMA undergo hyperplasia and hypertrophy

triggered by PPARγ activation (Rosen and MacDougald, 2006). Furthermore, with obesity, adipocyte

precursor mesenchymal stem cells tilt their lineage commitment towards adipogenesis (Silva et al., 2016,

Bowers and Singer, 2021). While BMA are now known to be a distinct adipocyte population, they express

several characteristics of white and brown adipocytes (Boroumand and Klip, 2020; Krings et al., 2012;

Sulston and Cawthorn, 2016; Tratwal et al., 2020). Brown adipocyte markers of BMA are reduced in obese

mice and a parallel adoption of more white adipocyte markers has been observed (Adler et al., 2014b;

Doucette et al., 2015; Krings et al., 2012). The amalgamation of upregulated white adipocyte proteins and

elevated fat storage capacity is known as BM adipocyte whitening. Obesity-induced gains in whole body

adiposity was causally linked to elevated circulating CD11b$^+$ monocytes (Takahashi et al., 2003). However,

the influence of BMA from either lean or obese mice on the expansion of CD11b$^+$ monocytes had not been

previously investigated.

The limited spatial capacity of the BM interior as well as the proximity of the cellular lineages in

the various BM niches lead us to anticipate that the obesity-linked BMA whitening would influence the

monocytic lineage. Here, we found a temporal, positive link between BMA hypertrophy and monocytosis

at 3 weeks of HFD. By 8 weeks of HFD, BMA whitening also positively correlated with the rise in tissue

invasive Ly6C^{high} monocytes in the BM. We did not observe significant gains in progenitor cells (CMP,

GMP) at either 3, 8 or 18 weeks of HFD compared to control diet, whereas Singer et al. found these cells

elevated at 16 weeks of HF-feeding (Singer et al., 2014, Singer et al., 2015). However, in another study, at

4 weeks of HFD, they did not observe changes in progenitor populations (PMID: 25869128), when

monocyte skewing towards Ly6C^{high} phenotype was already observed, apparently dissociating the

progenitor proliferative potential from the change in proportion of this monocyte subset.

As proof-of-concept for a direct and distinctive influence of adipocytes on monocyte number and

subsets, we show that exposure to soluble factors from *bona fide* white adipocytes favours Ly6C^{high}

monocyte preponderance while exposure to soluble factors from brown adipocytes promotes the

preponderance of Ly6C^{low} monocytes. Mechanistically, we observed that white adipocytes induced a

change in the proportion of Ly6C^{low} towards Ly6C^{high} monocytes, an unprecedented process that is not well

understood at present. Although the Ly6C^{low} population contained some level of multipotential progenitor

cells, it is unlikely that these contribute in a major way to the switch in preponderance of Ly6C^{high}

monocytes in response to WATA-CM, since this population did not undergo changes in the proximal

lineage-committed precursors (CFU-M) when assayed for colony forming unit potential, and further did

not involve BrdU accrual (indicative of DNA replication and cell proliferation).

Brown adipocytes, on the other hand, induce their influence on Ly6C^{low} monocytes by increasing

their proliferation, evinced by promoting BrdU incorporation into monocytes and an inducing expansion of

monocyte lineage-committed precursor CFU-M colonies when tested for colony formation. Interestingly,

secretome studies have reported that BAT-derived VEGFA and VEGFB can stimulate proliferation in

adipocyte progenitors and endothelial cells (Bagchi et al., 2013; Leung et al., 1989). Adipocyte-specific

VEGFA transgenic mice were protected from the effects of HFD and presented altered macrophage

polarization (Elias et al., 2012). Therefore, it is conceivable that a reduction in brown adipocyte

characteristics of BMA also leads to a drop in the factors that would normally favour proliferation of

Ly6C^{low} monocytes. While the identification of the soluble unknown factor(s) responsible for the adipocyte-

specific myeloid influence is outside the focus of the present study, we have laid the foundation for a future,

comprehensive investigation of the molecular mediators involved.

How do adipocytes skew myeloid lineage subsets?

While the designation of macrophages as 'M1 and M2' (of pro- or anti-inflammatory

characteristics, respectively) represents the extremes of a wide spectrum of phenotypes, the 'M1' pro-

inflammatory orientation has been linked to glycolytic activation while the 'M2' anti-inflammatory

orientation has been linked to oxidative phosphorylation (Castoldi et al., 2016; Kelly and O'Neill, 2015).

A similar glycolytic metabolic switch with obesity has been reported for CD8 pro-inflammatory T cells

(Buck et al., 2017). Surprisingly, while pro-inflammatory pathways in monocytes are activated by

hyperglycemic conditions (Shanmugam et al., 2003), the metabolic preference of the different monocyte

subsets had not been explored. This is a compelling question, given that Ly6C^{high} monocytes penetrate

tissues and differentiate to pro-inflammatory 'M1' AT macrophages while Ly6C^{low} monocytes enter tissues

to promote healing and repair (Geissmann et al., 2003; Lin et al., 2009). Therefore, we hypothesized here

that, in the context of metabolic inflammation, there is a change in the metabolic phenotype of monocyte

subsets.

Our findings show that Ly6C^{high} and Ly6C^{low} monocytes had different metabolic preferences at

baseline, with Ly6C^{low} having higher mitochondrial membrane potential indicative of more active oxidative

phosphorylation. Interestingly, HFD consumption lowered the mitochondrial membrane potential of both

Ly6C^{low} and Ly6C^{high} monocytes, implicating a reduction in their mitochondrial oxidative capacity. This

change aligned with a HFD-induced reduction in oxygen consumption by Ly6C^{low} monocytes. Importantly,

this difference was observed as early as 3 weeks and exacerbated by 8 and 18 weeks of HFD. The elevated

lactate concentration measured in the BM at all HFD durations may be reflective of metabolic changes in

cells of the BM cavity. Since we did not observe changes in the lactate levels in WATA-CM or BATA-

CM, the higher lactate in BM of HFD-fed mice may not originate from changes in BMA.

Along with the change in monocyte subset proportions, we observed an increase in genes

responsible for mitochondrial fragmentation and a decrease in those involved in mitochondrial elongation

in monocytes from HFD-fed mice and treated with WATA-CM and BMA-CM. The physiological

consequence of these changes was revealed by the observation of a striking mitochondrial fragmentation in

monocytes treated with WATA-CM in comparison to control or BATA-CM treated monocytes. HFD is

known to induce mitochondrial fission in a number of cells that correlates with insulin resistance. A recent

report linked the two phenomena through *Drp1* action, as *Drp1* inhibition reversed HFD-induced insulin

resistance (Filippi et al., 2017). Furthermore, obese and diabetic humans have reduced *Mfn2* expression in

skeletal muscle. *Mfn2* positively correlates with insulin sensitivity (Bach et al., 2005), and *Mfn2* gene and

protein are significantly downregulated in liver cells of HFD-fed rats (Gan et al., 2013). In our analysis, the

expression of *Mfn2* negatively correlated with the abundance of Ly6C^{high} monocytes, while, the expression

of *Drp1* positively correlated with the BM abundance of Ly6C^{high} monocytes. Overall, our findings reveal

novel and distinct effects of adipocyte types on monocyte subset development that may occur through

changes in mitochondrial fitness.

Given that the proportion of Ly6C^{low} monocytes dropped in favour of a predominance of Ly6C^{high}

monocytes by 8 but not 3 weeks of HFD, it is possible that this subset change obeys a prior shift in their

metabolic orientation. Supporting these observations, exposure of monocytes to white adipocyte CM also

reduced their mitochondrial membrane potential. Furthermore, BMA-CM from 8-week HFD-fed mice also

lowered monocyte mitochondrial membrane potential compared to control BMA-CM and favoured

Ly6C^{high} predominance. Mechanistically, these data suggest that white adipocytes induce a monocyte

metabolic shift that may guide Ly6C^{high} monocyte prevalence.

In conclusion, we demonstrate an interconnectivity of cellular networks within the BM, a

vulnerability of the BM microenvironment to HFD, and a consequential myeloid lineage determination

likely due to BMA whitening. Given the vastly different metabolic and soluble factors between *bona fide*

white and brown adipocytes, it is easy to imagine that phenotypic changes in BMA can vastly alter the BM

microenvironment and the factors that surround hematopoietic cells. BMA whitening during HFD feeding

contributes to myelopoiesis and predominance of invasive Ly6C^{high} monocytes mediated by changes in

monocyte metabolism.

Materials and Methods

Animal Procedures

Mouse protocols were approved by the animal care committee (Protocol #20011850 to S.E.G. and D.J.P., University of Toronto and #1000047074 to A.K., The Hospital for Sick Children). Three groups of 9-week-old male C57BL/6 mice were fed either low fat diet (10% kcal from fat; D12450Ji, Research Diets) or HFD (60% kcal from fat; D12492i, Research Diets) for 3, 8, or 18 weeks. Each duration of feeding involved at least 3 mice within each dietary group, and the precise number of mice evaluated in each experiment is indicated in the figure legends. The feeding protocol was repeated with three independent cohorts of mice, initiated at different times. The graphs illustrated include data collected from the three cohorts. Logistically, not every mouse could be used for every type of experiment, and the number of mice involved is indicated in each figure. Mice were singly caged to avoid stress and weight inequalities induced by aggression of the dominant male to the subordinate male (Kappel et al., 2017). The cages were maintained at 22°C on a 12h light/dark cycle, the diets were replaced once per week and various cage enrichments were provided including switching the positioning of the cages throughout the study.

Body weight was measured twice per week. Upon completion of the respective feeding durations, mice were fasted for 4 h, blood was drawn via tail vein and fasting blood glucose was measured with a glucometer. Mice were euthanized by cervical dislocation. Tissues were dissected and flash frozen in liquid nitrogen for RNA isolation. EWAT, blood and BM were immediately processed for flow cytometry analysis.

EWAT-Derived Stromal Vascular Fraction (SVF) Isolation

EWAT-derived SVF was collected as previously described (Orr et al., 2013). Excised EWAT was rinsed in PBS, minced and digested with 0.47wv/mL Liberase solution (Millipore Sigma, Cat No 5401054001) shaking at 200 rpm for 20 min at room temperature. SVF was isolated using a 70 μm cell strainer, the enzymatic activity of Liberase was neutralized with DMEM containing HI-FBS for 5 min and

centrifuged at 500 ×g for 10 min. The supernatant containing adipocytes was removed, the pellet was re-

suspended in red blood cell lysis buffer for 5 min to remove erythrocytes and washed with PBS three times.

BM Histology

Extracted tibia and femur bones were fixed in 10% formalin for 48 h, decalcified in 14% EDTA

solution for 14 days, paraffin wax-embedded, and longitudinal sections were stained with H&E and Oil

Red O (Millipore Sigma; Cat No O0625) to visualize BM adiposity. Fifty imaging fields per mouse femur

were analyzed for the presence of adipocytes, assessed by blinded volunteers. Adipocyte area was

calculated using ImageJ software version 1.53.

BM cell fractionation

Following cervical dislocation, femurs and tibias were dissected from the mice fed control or the

HFD for 3, 8 or 18 weeks. Epiphyses were cut from one end of each bone, then bones were placed in 0.2

mL microtubes containing 18-gauge holes. Microtubes were placed inside 0.5 mL Eppendorf tubes

containing 100 μL media or PBS. This complex was centrifuged at 15,000 x g for 10 s to extract BM in the

pellet. BM supernatant was removed from the cellular pelleted fraction to be assessed by ELISA and used

for lactate quantification. BM cells were used for flow cytometry analysis, fluorescence-activated cell

sorting (FACS) to identify monocyte subsets and their gene expression, as well as for differentiation to

BM-derived monocytes.

Primary Adipocyte Isolation and Culture to generate Conditioned Media

Epididymal white and interscapular brown AT were dissected in sterile conditions, the tissues

underwent mechanical mincing and enzymatic digestion by collagenase type I (Worthington Biochemical

Corp, Cat No LS004197) at 4 mg/mL shaking for 1 hour. The digested adipocytes were then filtered through

a 200 μm cell strainer and centrifuged, washed twice and plated in adipocyte media: high glucose DMEM,

1% BSA, 0.5% FBS, and 1% Penicillin and Streptomycin for 24 h. BM adipocytes were isolated from the

floating fraction following BM isolation via centrifugation. The cells were counted and cultured in

adipocyte media overnight. Tissues from three mice were collected and the isolated cells were pooled in

each CM generating experiment.

***In vitro*-generated BM-Derived Monocytes**

Murine primary BM cells were plated at 500,000 cells/mL on Thermo Scientific™ Nunc™ Non-

Treated 6-well plates (Cat No 1256680) with complete RPMI (Cat No 350-000-EL, Wisent BioProducts,

Saint-Jean-Baptiste, QC, Canada) and supplemented with recombinant mouse M-CSF (Peprotech Inc. Cat

No 315-02) at 20ng/mL for 5 days (Francke et al., 2011). Following 5 days of BM cell incubation on the

ultra-low attachment surface plates, the suspended cells have the signature of mature monocytes and can

be used for experimentation. These BM-derived differentiated monocytes were then collected, washed with

RMPI, centrifuged and re-suspended with various adipocyte CM for 18 h. Typically when BM cells were

collected three mice were independently processed at the same time. This process yielded three cohorts of

differentiated monocytes that would each be treated with Media-, WATA- and BATA-CM.

Flow Cytometry

Following red blood cell lysis, SVF, blood and the BM were washed with FACS buffer (PBS, 1%

(v/v) heat-inactivated FBS, 0.2% (w/v) BSA), stained with Aqua Live/Dead Antibody (ThermoFisher,

L34957) for 15 min in the dark, washed with FACS buffer again, blocked with anti-mouse CD16/32

antibody (eBioscience, 14-0161-86) for 15 min in the dark following a 30 min incubation with

corresponding antibody for each antigen **(**see **Table 1** for list of antibodies**).** The samples were then fixed,

read on LSR Fortessa cytometer (BD Biosciences) and data gated based on appropriate fluorescence minus

one (FMO) controls, and analyzed on the FlowJo software version 10 (Becton, Dickinson & Company).

See Figure S1 for gating strategies.

Quantitative PCR (qPCR)

RNA was extracted from cells or tissues using TRIzol™ (Life Technologies, Cat No 15596018),

then cDNA synthesized using the SuperScript VILO cDNA kit (Life Technologies, Cat No 11754250)

according to manufacturer's instructions. qPCR reactions were performed with predesigned TaqMan probes

(Life Technologies) on a StepOne Plus Real-Time PCR System (Applied Biosystems-ThermoFisher).

Duplicates of each cDNA sample were used for gene expression analysis and values were normalized to

the average of a set of the following housekeeping genes: Abt1, Hprt, and Eef2 by the $\Delta\Delta C_T$ method.

BrdU incorporation

The abcam protocol was followed to detect BrdU through an anti-BrdU antibody

(https://www.abcam.com/protocols/brdu-staining-protocol#BrdU%20labeling). BM-derived *in vitro*-

generated monocytes were labelled with 10 μM BrdU (ThermoFisher, 000103) during the 18-hour

treatment at 37C. The cells were then washed with FACS buffer, surface markers were stained as described

above and incubated in Foxp3 Fixation/Permeabilization Buffer (eBioscience, 00-5523-00) overnight. They

were then treated with 30 μg of DNAse I.9 (Sigma, 260913) for 1 hour, stained with anti-BrdU antibody

(abcam, ab6326) for 1 hour and then washed with FACS buffer in preparation to run on LSR Fortessa

Cytometer (BD Biosciences).

Fluorescent Activated Cell Sorting of Monocytes

BM from two femurs and tibias of 8-10-week-old chow-fed mice was used in the isolation of

monocyte subsets via fluorescent activated cell sorting using the FACS Aria Ilu instrument. Gr1[-] CD115[+]

CD11b[+] and Ly6C[+] surface markers were used in identifying total monocytes and two subsets were

subsequently distinguished by the relative surface abundance (low and high) of Ly6C protein (Figure S1).

Isolated Ly6C[low] and Ly6C[high] monocytes were then validated for positive staining by antibodies CX3CR1

and CCR2 (Figure S1), respectively. Ly6C and Cd11b FMO controls shown in Figure S1B illustrated that

the gates drawn for Ly6C[low] and Ly6C[high] were confidently identifying cells positive for both these markers.

Lactate quantification

The BM content was fractioned into hematopoietic and supernatant portions by centrifugation at

1,600 rpm. The supernatant fraction was re-suspended in 500 μL of 1X PBS (supplemented with Protease

Cocktail Inhibitor, Sigma). One drop of the BM supernatant fraction was placed on the Blood Lactate Meter

Lactate Plus (Nova Biomedical) for the quantification of its lactate content. The values were normalized to

the undiluted BM isolated from the femur.

Cell Metabolism Analysis

Oxygen consumption (OCR) and extracellular acidification rates were measured using a Seahorse

XFe96 analyzer (Seahorse Bioscience). Seahorse 96-well cell culture plates were pretreated with 1 μg/cm^2

Corning™ Cell-Tak Cell and Tissue Adhesive (Fisher Scientific, Cat No CB40240) and FACS sorted

Ly6C^{low} or Ly6C^{high} monocytes were seeded at 100,000 cells/well, allowed to equilibrate for 1 h in XF base

medium supplemented with 11 mM glucose, 1 mM sodium pyruvate and 2 mM Glutamine. Following

baseline measurements, FCCP was injected at 1 mM and thereafter an injection of Antimycin A/ Rotenone.

Cells were counted following the seahorse assay to test for changes in cellular detachment between the

conditions.

Immunofluorescence Staining for Super-Resolution Microscopy

Monocytes were spun down onto coverslips and fixed with 4% paraformaldehyde for 15 min.

Thereafter, the cells were rinsed with PBS and permeabilized with 0.1% Triton X-100 for 10 min. The cells

were then blocked with 2% BSA in PBS at 4℃ overnight. The cells were then incubated with the primary

antibody, Tom20 (FL-145, Santa Cruz, sc-11415), in a 1:100 dilution in blocking buffer at 4℃ overnight.

Thereafter, the cells were incubated with Alexa Fluor 488 anti-rabbit secondary antibody (1:100) for 2 h at

room temperature. Cells were incubated with Alexa Fluor 568 Phalloidin (1:100) for 3 h at room

temperature. Coverslips were then mounted by ProLong Diamond (Thermo Fisher Scientific) and allowed

to dry for two days. The images were acquired via Zeiss LSM880 Airyscan super resolution microscopy

and the images were processed via Fiji software. The percentage of monocytes displaying each

mitochondrial morphology (fragmented, intermediate, or tubulated) was scored as described previously

(Rambold et al., 2011).

Colony-Forming Unit Assay

In vitro differentiated BM-derived monocytes (5×10^5 cell/ml) were treated as indicated, then

pelleted and suspended in Iscove's modified Dulbecco's medium to 2×10^5 cell/ml, diluted in MethoCult

GF-M3434 medium and plated at 20,000 cells/35 mm well in a 6-well plate of low-adherence culture-ware

(STEMCELL Technologies, Inc.), in triplicate. The colonies that grew had distinct cell lineages and

morphologies as defined in detail by the STEMCELL manual for MethoCult GF-M3434 methylcellulose

media. After 7-10 days, colonies were counted, using an inverted light microscope, and identified as: Burst-

forming unit-erythroid (BFU-E), Colony-forming unit-granulocyte, erythrocyte, macrophage,

megakaryocyte (CFU-GEMM), CFU-granulocyte (CFU-G) or CFU-macrophage (CFU-M), based on the

STEMCELL Technologies detailed instructions. Of note, CFU-GEMM progenitors are also known as

CMP, and GMP progenitors give rise to the CFU-G and CFU-M lineages (immediate precursors of

monoblasts and monocytes) that form colonies in this assay (https://www.stemcell.com/hematopoietic-

stem-and-progenitor-cells-lp.html).

ELISA Determinations

The BM content was fractioned into hematopoietic and supernatant portions by centrifugation at

1,600 rpm. The supernatant fraction was re-suspended in 500ul of 1X PBS (supplemented with Protease

Cocktail Inhibitor, Sigma). Total protein content was estimated by Pierce™ BCA Protein Assay Kit

(ThermoFisher Scientific). Subsequently, mouse TNF, IL-1b, IL-6, CXCL1 and CCL2 ELISAs were

carried out according to the manufacturers' instructions. All Mouse DuoSet ELISA kits were purchased

from R&D systems. The individual protein contents were finally normalised to total protein content.

Statistical Analysis

Data are expressed as means ± SEM. An unpaired Student's T test was used to detect differences

in data sets with one variable such as that in Figure S2. A one-way ANOVA with Tukey post-test was used

to detect differences in data sets of two or more variables. Statistical significance was set at $P < 0.05$. Data

analysis and statistics tests were done using GraphPad Prism version 6 (GraphPad Software), Flow Jo

Software version 10.4 and figures were prepared using Adobe Illustrator CS6 version 16.0.

Correlative analysis statistics were conducted using the linear regression analysis via GraphPad

Prism version 6 (GraphPad Software) testing whether slopes and intercepts are significantly different. The

r^2 represents the goodness of the fit while the P values greater than 0.05 represent whether the slope of the

line of best fit is significantly non-zero.

Acknowledgements

We thank Dr. Spencer Freeman for helpful discussion. This work was supported by Canadian Institutes of

Health Research (CIHR) Foundation Grant (FND-143203 to A.K. and FDN-14333 to D.J.P. Studentship

support for P.B. was from a Banting & Best Diabetes Centre-Novo Nordisk Graduate Studentship (2017-

2019) and a SickKids Research Training Centre Award (2019- 2020).

Author Contribution

P.B., A.K. and S.E.G. conceived the study design. P.B. coordinated and performed experiments

and analyzed the data. M.W., J.S., I.T., N.S., L.Z., Z.L., A.L., T.M., Y.Z. and T.S. participated in

experimental work and data collection. P.J.B. contributed with valuable advice, troubleshooting, and

performed and analyzed experiments. A.K., D.J.P., S.E.G. and C.R. provided access to resources and

facilities. P.B. and A.K. wrote the manuscript, and all authors read and approved it.

References:
Adler, B.J., Kaushansky, K., and Rubin, C.T. (2014). Obesity-driven disruption of
haematopoiesis and the bone marrow niche. Nat. Rev. Endocrinol. *10*, 737–748.

Ambrosi, T.H., Scialdone, A., Graja, A., Gohlke, S., Jank, A.M., Bocian, C., Woelk, L., Fan, H.,
Logan, D.W., Schürmann, A., et al. (2017). Adipocyte Accumulation in the Bone Marrow during
Obesity and Aging Impairs Stem Cell-Based Hematopoietic and Bone Regeneration. Cell Stem
Cell *20*, 771–784.e6.

Auffray, C., Fogg, D., Garfa, M., Gaelle, E., Join-Lambert, O., Kayal, S., Sarnacki, S., Ana, C.,
Lauvau, G., and Geissmann, F. (2007). Monitoring of Blood Vessels and Tissues by a Population
of Monocytes with Patrolling Behavior. *317*, 666–671.

Auffray, C., Fogg, D.K., Narni-Mancinelli, E., Senechal, B., Trouillet, C., Saederup, N.,
Leemput, J., Bigot, K., Campisi, L., Abitbol, M., et al. (2009). CX 3 CR1 + CD115 + CD135 +
common macrophage/DC precursors and the role of CX 3 CR1 in their response to inflammation
. J. Exp. Med. *206*, 595–606.

Bach, D., Naon, D., Pich, S., Soriano, F.X., Vega, N., Rieusset, J., Laville, M., Guillet, C.,
Boirie, Y., Wallberg-henriksson, H., et al. (2005). Expression of Mfn2, the Charcot-Marie-Tooth
Neuropathy Type 2A Gene, in Human Skeletal Muscle. Diabetes *54*, 2685–2693.

Bagchi, M., Kim, L.A., Boucher, J., Walshe, T.E., Kahn, C.R., and Amore, P.A.D. (2013).
Vascular endothelial growth factor is important for brown adipose tissue development and
maintenance. FASEB J. *27*, 3257–3271.

Beattie, L., Sawtell, A., Mann, J., Frame, T.C.M., Teal, B., de Labastida Rivera, F., Brown, N.,
Walwyn-Brown, K., Moore, J.W.J., MacDonald, S., et al. (2016). Bone marrow-derived and
resident liver macrophages display unique transcriptomic signatures but similar biological
functions. J. Hepatol. *65*, 758–768.

Boniakowski, A.E., Kimball, A.S., Joshi, A., Schaller, M., Davis, F.M., denDekker, A., Obi,
A.T., Moore, B.B., Kunkel, S.L., and Gallagher, K.A. (2018). Murine macrophage chemokine
receptor CCR2 plays a crucial role in macrophage recruitment and regulated inflammation in
wound healing. Eur. J. Immunol. *48*, 1445–1455.

Boroumand, P., and Klip, A. (2020). Bone marrow adipose cells – cellular interactions and
changes with obesity. J. Cell Sci. *1*, 1–8.

Bowers E., Singer K. (2021) Obesity-induced inflammation: The impact of the hematopoietic
stem cell niche. JCI Insight. *6*, e145295.

Buck, M.D., Sowell, R.T., Kaech, S.M., and Pearce, E.L. (2017). Metabolic Instruction of
Immunity. Cell *169*, 570–586.

Cassado, A.A., D'Império Lima, M.R., and Bortoluci, K.R. (2015). Revisiting mouse peritoneal
macrophages: Heterogeneity, development, and function. Front. Immunol. *6*, 1–9.

Castoldi, A., De Souza, C.N., Saraiva Câmara, N.O., and Moraes-Vieira, P.M. (2016). The
macrophage switch in obesity development. Front. Immunol. *6*, 1–11.

Centers for Disease Control and Prevention, C. (2017). National Diabetes Statistics Report:
Estimates of Diabetes and Its Burden in the United States. Atlanta, GA: Centers for Disease
Control and Prevention; 2017. US Dep. Heal. Hum. Serv.

Challen, G.A., Boles, N., Lin, K.K.Y., and Goodell, M.A. (2009). Mouse hematopoietic stem
cell identification and analysis. Cytom. Part A *75*, 14–24.

Cuminetti, V., and Arranz, L. (2019). Bone Marrow Adipocytes: The Enigmatic Components of
the Hematopoietic Stem Cell Niche. J. Clin. Med. *8*, 707.

Doucette, C.R., Horowitz, M.C., Berry, R., MacDougald, O.A., Anunciado-Koza, R., Koza,
R.A., and Rosen, C.J. (2015a). A High Fat Diet Increases Bone Marrow Adipose Tissue (MAT)
But Does Not Alter Trabecular or Cortical Bone Mass in C57BL/6J Mice. J Cell Physiol *230*,
2032–2037.

Doucette, C.R., Horowitz, M.C., Berry, R., Macdougald, O.A., Anunciado-Koza, R., Koza, R.A.,
and Rosen, C.J. (2015b). A High Fat Diet Increases Bone Marrow Adipose Tissue (MAT) But
Does Not Alter Trabecular or Cortical Bone Mass in C57BL/6J Mice. J. Cell. Physiol. *230*,
2032–2037.

Elias, I., Franckhauser, S., Ferré, T., Vilà, L., Tafuro, S., Muñoz, S., Roca, C., Ramos, D., Pujol,
A., Riu, E., et al. (2012). Adipose Tissue Overexpression of Vascular Endothelial Growth Factor
Protects Against Diet-Induced Obesity and Insulin Resistance. Diabetes *61*, 1801–1813.

Filippi, B.M., Abraham, M.A., Silva, P.N., Bauer, P. V, Rocheleau, J. V, Lam, T.K.T., Filippi,
B.M., Abraham, M.A., Silva, P.N., Rasti, M., et al. (2017). Mitochondrial Fission Changes in the
Dorsal Vagal Report Dynamin-Related Protein 1-Dependent Mitochondrial Fission Changes in
the Dorsal Vagal Complex Regulate Insulin Action. CellReports *18*, 2301–2309.

Fink, L.N., Costford, S.R., Lee, Y.S., Jensen, T.E., Bilan, P.J., Oberbach, A., Blüher, M.,
Olefsky, J.M., Sams, A., and Klip, A. (2014a). Pro-Inflammatory macrophages increase in
skeletal muscle of high fat-Fed mice and correlate with metabolic risk markers in humans.
Obesity *22*, 747–757.

Fink, L.N., Costford, S.R., Lee, Y.S., Jensen, T.E., Bilan, P.J., Oberbach, A., Blüher, M.,
Olefsky, J.M., Sams, A., and Klip, A. (2014b). Pro-Inflammatory macrophages increase in
skeletal muscle of high fat-Fed mice and correlate with metabolic risk markers in humans.
Obesity *22*, 747–757.

Francke, A., Herold, J., Weinert, S., Strasser, R.H., and Braun-Dullaeus, R.C. (2011). Generation
of mature murine monocytes from heterogeneous bone marrow and description of their
properties. J. Histochem. Cytochem. *59*, 813–825.

Friedrich, K., Sommer, M., Strobel, S., Thrum, S., and Blüher, M. (2019). Perturbation of the
Monocyte Compartment in Human Obesity. Front. Immunol. *10*, 1–10.

Gan, K., Wang, C., Chen, J., Zhu, C., and Song, G. (2013). Mitofusin-2 ameliorates high-fat diet-
induced insulin resistance in liver of rats. *19*, 1572–1581.

Garvey, W.T., Henry, R.R., Kern, P.A., Saghizadeh, M., and Ong, J.M. (2008). The expression

of TNF alpha by human muscle. Relationship to insulin resistance. J. Clin. Invest. *97*, 1111–
1116.

Geissmann, F., Jung, S., and Littman, D.R. (2003). Blood monocytes consist of two principal
subsets with distinct migratory properties. Immunity *19*, 71–82.

Gibbings, S.L., Goyal, R., Desch, A.N., Leach, S.M., Prabagar, M., Atif, S.M., Bratton, D.L.,
Janssen, W., and Jakubzick, C. V. (2015). Transcriptome analysis highlights the conserved
difference between embryonic and postnatal-derived alveolar macrophages. Blood *126*, 1357–
1366.

Gottlieb, R.A., and Bernstein, D. (2017). Mitochondrial Remodeling: Rearranging, Recycling,
and Reprogramming. Cell Calcium *60*, 88–101.

Griffin, C., Eter, L., Lanzetta, N., Abrishami, S., Varghese, M., McKernan, K., Muir, L., Lane,
J., Lumeng, C.N.-K., and Singer, K. (2018). TLR4, TRIF, and MyD88 are essential for
myelopoiesis and CD11c+ adipose tissue macrophage production in obese mice. J. Biol. Chem.
jbc.RA117.001526.

Hamann, A. (2017). Obesity Update 2017. Diabetologe.

Kamei, N., Tobe, K., Suzuki, R., Ohsugi, M., Watanabe, T., Kubota, N., Ohtsuka-kowatari, N.,
Kumagai, K., Sakamoto, K., Kobayashi, M., et al. (2006). Overexpression of Monocyte
Chemoattractant Protein-1 in Adipose Tissues Causes Macrophage Recruitment and Insulin
Resistance *. J. Biol. Chem. *281*, 26602–26614.

Kappel, S., Hawkins, P., and Mendl, M. (2017). To Group or Not to Grou? Good Practice for
Housing Male Laboratory Mice. Animals 1–25.

Kelly, B., and O'Neill, L.A. (2015). Metabolic reprogramming in macrophages and dendritic
cells in innate immunity. Cell Res. *25*, 771–784.

Kohyama, M., Matsuoka, S., Shida, K., Sugihara, F., Aoshi, T., Kishida, K., Ishii, K.J., and
Arase, H. (2016). Monocyte infiltration into obese and fibrilized tissues is regulated by PILRα.
Eur. J. Immunol. *46*, 1214–1223.

Kratofil, R.M., Kubes, P., and Deniset, J.F. (2017). Monocyte conversion during inflammation
and injury. Arterioscler. Thromb. Vasc. Biol. *37*, 35–42.

Krings, A., Rahman, S., Huang, S., Lu, Y., Czernik, P.J., and Lecka-Czernik, B. (2012). Bone
marrow fat has brown adipose tissue characteristics, which are attenuated with aging and
diabetes. Bone *50*, 546–552.

Lanthier, N., Molendi-Coste, O., Horsmans, Y., van Rooijen, N., Cani, P.D., and Leclercq, I.A.
(2010). Kupffer cell activation is a causal factor for hepatic insulin resistance. Am. J. Physiol.
Liver Physiol. *298*, G107–G116.

Lee, M.K.S., and Murphy, A.J. (2019). Glycolysis Is Required for LPS-Induced Activation and
Monocytes. *10*, 1–10.

Lessard, A.J., LeBel, M., Egarnes, B., Préfontaine, P., Thériault, P., Droit, A., Brunet, A., Rivest,

S., and Gosselin, J. (2017). Triggering of NOD2 Receptor Converts Inflammatory Ly6Chighinto
Ly6ClowMonocytes with Patrolling Properties. Cell Rep. *20*, 1830–1843.

Leung, D.W., Cachianes, G., Kuang, W., Goeddel, D. V, and Ferrara, N. (1989). Vascular
Endothelial Growl th Factor Is a Secreted Angiogenic Mitogen. Science (80-.). *246*, 1306–9.

Lin, S.L., Castano, A.P., Nowlin, B.T., Lupher, M.L., and Duffield, J.S. (2009). Bone Marrow
Ly6Chigh Monocytes Are Selectively Recruited to Injured Kidney and Differentiate into
Functionally Distinct Populations. J. Immunol. *183*, 6733–6743.

Liu, T., Sun, Y.-C., Cheng, P., and Shao, H.-G. (2019). Adipose tissue macrophage-derived
exosomal miR-29a regulates obesity-associated insulin resistance. Biochem. Biophys. Res.
Commun.

Lumeng, C.N., Deyoung, S.M., Bodzin, J.L., and Saltiel, A.R. (2007). Increased Inflammatory
Properties of Adipose Tissue Macrophages Recruited During Diet-Induced Obesity. Diabetes *56*,
16–23.

Lumeng, C.N., Delproposto, J.B., Westcott, D.J., and Saltiel, A.R. (2008). Phenotypic Switching
of Adipose Tissue Macrophages With Obesity Is Generated by Spatiotemporal Differences in
Macrophage Subtypes. Diabetes *57*, 3239–3246.

Manicone, A.M., Gong, K., Johnston, L.K., and Giannandrea, M. (2016). Diet-induced obesity
alters myeloid cell populations in naïve and injured lung. Respir. Res. *17*, 24.

Nagareddy, P.R., Kraakman, M., Masters, S.L., Stirzaker, R.A., Gorman, D.J., Grant, R.W.,
Dragoljevic, D., Hong, E.S., Abdel-Latif, A., Smyth, S.S., et al. (2014). Adipose tissue
macrophages promote myelopoiesis and monocytosis in obesity. Cell Metab. *19*, 821–835.

Nakagami, H. (2013). The Mechanism of White and Brown Adipocyte Differentiation. Diabetes
Metab. J. *37*, 85–90.

Nascimento, E.B.M., Boon, M.R., and van Marken Lichtenbelt, W.D. (2014). Fat cells gain new
identities. Sci. Transl. Med. *6*, 1–4.

Nov, O., Shapiro, H., Ovadia, H., Tarnovscki, T., Dvir, I., Shemesh, E., Kovsan, J., Shelef, I.,
Carmi, Y., Voronov, E., et al. (2013). Interleukin-1β Regulates Fat-Liver Crosstalk in Obesity by
Auto-Paracrine Modulation of Adipose Tissue Inflammation and Expandability. PLoS One *8*.

O'Neill, L., Kishton, R., and Rathmell, J. (2016). A guide to immunometabolism for
immunologists. Nat. Rev. Immunol. *16*, 553–565.

Orr, J.S., Kennedy, A.J., and Hasty, A.H. (2013). Isolation of Adipose Tissue Immune Cells. J.
Vis. Exp. 1–7.

Patel, A.A., Zhang, Y., Fullerton, J.N., Boelen, L., Rongvaux, A., Maini, A.A., Bigley, V.,
Flavell, R.A., Gilroy, D.W., Asquith, B., et al. (2017). The fate and lifespan of human monocyte
subsets in steady state and systemic inflammation. J. Exp. Med. *214*, 1913–1923.

Pence, B.D., and Yarbro, J.R. (2019). Classical monocytes maintain ex vivo glycolytic
metabolism and early but not later inflammatory responses in older adults. Immun. Ageing *16*,

1–8.

Pham, T.T., Ivaska, K.K., Hannukainen, J.C., Virtanen, K.A., Lidell, M.E., Parkkola, R., Piirola,
S., Oikonen, V., Nuutila, P., Biology, C., et al. (2020). Human bone marrow adipose tissue is a
metabolically active and insulin sensitive distinct fat depot. J. Clin. Endocrinol. Metab. *pii*,
dgaa216.

Rambold A.S., Kostelecky B., Elia N., Lippincott-Schwartz J. (2011) Tubular network formation
protects mitochondria from autophagosomal degradation during nutrient starvation. Proc Natl
Acad Sci U S A. *108*, 10190-10195.

Renovato-Martins, M., Matheus, M.E., de Andrade, I.R., Moraes, J.A., da Silva, S.V., Citelli dos
Reis, M., de Souza, A.A.P., da Silva, C.C., Bouskela, E., and Barja-Fidalgo, C. (2017).
Microparticles derived from obese adipose tissue elicit a pro-inflammatory phenotype of CD16 +
, CCR5 + and TLR8 + monocytes. Biochim. Biophys. Acta - Mol. Basis Dis. *1863*, 139–151.

Rosen, E.D., and MacDougald, O.A. (2006). Adipocyte differentiation from the inside out. Nat.
Rev. Mol. Cell Biol. *7*, 885–896.

Scheller, E.L., Doucette, C.R., Learman, B.S., Cawthorn, W.P., Khandaker, S., Schell, B., Wu,
B., Ding, S.Y., Bredella, M.A., Fazeli, P.K., et al. (2015). Region-specific variation in the
properties of skeletal adipocytes reveals regulated and constitutive marrow adipose tissues. Nat.
Commun. *6*, 1–13.

Scheller, E.L., Khoury, B., Moller, K.L., Wee, N.K.Y., Khandaker, S., Kozloff, K.M.,
Abrishami, S.H., Zamarron, B.F., and Singer, K. (2016). Changes in skeletal integrity and
marrow adiposity during high-fat diet and after weight loss. Front. Endocrinol. (Lausanne). *7*, 1–
13.

Shanmugam, N., Reddy, M.A., Guha, M., and Natarajan, R. (2003). High glucose-induced
expression of proinflammatory cytokine and chemokine genes in monocytic cells. Diabetes *52*,
1256–1264.

Sharma, M., Boytard, L., Hadi, T., Koelwyn, G., Simon, R., Ouimet, M., Seifert, L., Spiro, W.,
Yan, B., Hutchison, S., et al. (2020). Enhanced glycolysis and HIF-1 α activation in adipose
tissue macrophages sustains local and systemic interleukin-1 β production in obesity. Sci. Rep.
10, 1–12.

Silva, S. V., Renovato-martins, M., Ribeiro-pereira, C., Citelli, M., and Barja-fidalgo, C. (2016).
Obesity Modifies Bone Marrow Microenvironment and Directs Bone Marrow Mesenchymal
Cells to Adipogenesis. Obes. Biol. Integr. Physiol. *24*, 2522–2532.

Singer, K., DelProposto, J., Lee Morris, D., Zamarron, B., Mergian, T., Maley, N., Cho, K.W.,
Geletka, L., Subbaiah, P., Muir, L., et al. (2014). Diet-induced obesity promotes myelopoiesis in
hematopoietic stem cells. Mol. Metab. *3*, 664–675.

Singer K., Maley N., Mergian T., DelProposto J., Cho K.W., Zamarron B.F., Martinez-
Santibanez G., Geletka L., Muir L., Wachowiak P., et al. (2015). Differences in Hematopoietic
Stem Cells Contribute to Sexually Dimorphic Inflammatory Responses to High Fat Diet-induced
Obesity. J Biol Chem. *290*, 13250-13262.

De Souza, C.T., Araujo, E.P., Bordin, S., Ashimine, R., Zollner, R.L., Boschero, A.C., Saad,
M.J.A., and Velloso, L.A. (2005). Consumption of a fat-rich diet activates a proinflammatory
response and induces insulin resistance in the hypothalamus. Endocrinology *146*, 4192–4199.

Sulston, R.J., and Cawthorn, W.P. (2016). Bone marrow adipose tissue as an endocrine organ:
close to the bone? Horm. Mol. Biol. Clin. Investig. *28*, 21–38.

Takahashi, K., Mizuarai, S., Araki, H., Mashiko, S., Ishihara, A., Kanatani, A., Itadani, H., and
Kotani, H. (2003). Adiposity Elevates Plasma MCP-1 Levels Leading to the Increased CD11b-
positive Monocytes in Mice *. J. Biol. Chem. *278*, 46654–46660.

Thapa, B., and Lee, K. (2019). Metabolic influence on macrophage polarization and
pathogenesis. BMB Rep. *52*, 360–372.

Tratwal, J., Labella, R., Bravenboer, N., and Kerckhofs, G. (2020). Reporting Guidelines ,
Review of Methodological Standards , and Challenges Toward Harmonization in Bone Marrow
Adiposity Research . Report of the Methodologies Working Group of the International Bone
Marrow Adiposity Society. Front. Endocrinol. (Lausanne). *11*, 1–36.

Ussar, S., Lee, K., Dankel, S., Boucher, J., Haering, M., Kleinridders, A., Thomou, T., Xue, R.,
Macoteia, Y., Cypess, A., et al. (2014). Asc-1, PAT2 and P2RX5 are novel cell surface markers
for white, beige and brown adipocytes. Sci Transl Med *247*, 247ra103.
doi:10.1126/scitranslmed.3008490. Asc-1,.

Verdeguer, F., and Aouadi, M. (2017). Macrophage heterogeneity and energy metabolism. Exp.
Cell Res. *360*, 35–40.

Weisberg, S.P., McCann, D., Desai, M., Rosenbaum, M., Leibel, R.L., and Ferrante, A.W.
(2003). Obesity is associated with macrophage accumulation in adipose tissue. J. Clin. Invest.
112, 1796–1808.

Weisberg, S.P., Hunter, D., Huber, R., Lemieux, J., Slaymaker, S., Vaddi, K., Charo, I., Leibel,
R.L., and Ferrante Jr, A.W. (2006). CCR2 modulates inflammatory and metabolic effects of
high-fat feeding. J. Clin. Invest. *116*, 115–124.

WHO (2016). World Health Organization. Global Report on Diabetes. Geneva, Switzerland.

Yao C.H., Wang R., Wang Y., Kung C.P., Weber J.D., Patti G.J. (2019) Mitochondrial fusion
supports increased oxidative phosphorylation during cell proliferation. Elife. *8*, e41351.

Yáñez A., Coetzee S.G., Olsson A., Muench D.E., Berman B.P., Hazelett D.J., Salomonis N.,
Grimes H.L., Goodridge H.S. (2017) Granulocyte-Monocyte Progenitors and Monocyte-
Dendritic Cell Progenitors Independently Produce Functionally Distinct Monocytes. Immunity.
47, 890-902.

Figure 1

A Body Weight

B Fasting Blood Glucose

C EWAT Weight

D Bone Marrow Adipocyte Area

E Bone Marrow Adipocyte Count

F Control   HFD

H

Figure 1. Mice fed HFD increased body weight, fasting blood glucose, visceral and bone marrow

adipose tissue. [A] Body weight was measured twice a week throughout the course of feeding. (n=14 per

dietary group). [B] Mice were fasted for 4 h after which fasting blood glucose was measured. (n=9-14 per

dietary group). [C] EWAT was weighed the day of sacrifice. (n=4 per dietary group at each feeding

duration). [D] Area covered by BM adipocytes were measured following H&E to visualize adipocyte

ghosts using quantified using ImageJ software. (n=6 mice in each dietary group at each feeding duration).

[E] BM adipocyte numbers per field were blindly counted following H&E staining using ImageJ software

version 1.53 (n=4-14 mice in each dietary group per feeding duration). [F] Representative images of

femurs from mice fed control and HFD diet for 3 weeks, that were fixed in formalin for 2 days followed

by 14 days EDTA decalcification and subsequently stained with Oil Red O (ORO). [G] Representative

images of femurs from mice fed control and HFD diet for 8 weeks, that were fixed in formalin for 2 days

followed by 14 days EDTA decalcification and subsequently stained with H&E. Scale bar is 25μm.

Results are expressed as mean ± SEM, n=4-14 mice per group. One-way ANOVA was used for statistical

analysis at each dietary group and across feeding periods. *$P < 0.05$, **$P < 0.01$, ***$P < 0.001$, ****$P <$

0.0001 and ns = not significant $P > 0.05$.

Figure 2

Figure 2. Bone marrow monocytes increased prior to adipose tissue macrophages. Analysis of

monocytes and macrophages in the stromal vascular fraction (SVF) of adipose tissue via flow cytometry

analysis, identified first by size and granularity through forward and side scatter and then using the

following markers: CD11b CD115 F4/80 Ly6C GR1 CD11c. (n=7-10 mice in each dietary group at each

feeding duration): [A] monocytes, [B] total macrophages, [C] pro-inflammatory macrophages. [D-F] Whole

adipose tissue gene expression of the myeloid marker *Itgam,* macrophage marker *Adgre* and pro-

inflammatory myeloid marker *Itgax* by RTqPCR. Each symbol represents a biological replicate analyzed

in duplicate and normalized to average of a group of housekeeping genes. (n=5 mice in each dietary group

at each feeding duration). [G-H] Gene expression analysis of embryonic derived tissue resident macrophage

markers *Timd4* and *Marco* by RTqPCR. Each symbol represents a biological replicate analyzed in duplicate

and normalized to average of a group of housekeeping genes. (n=3-5 mice in each dietary group at each

feeding duration). [I] Flow cytometry analysis of total BM monocytes (n=7 mice in each dietary group at

each feeding duration). [J] Monocyte subsets in the bone marrow with HFD-feeding (n=7 mice in each

dietary group at each feeding duration). Results are expressed as mean ± SEM. One-way ANOVA was used

for statistical analysis at each dietary group and across feeding periods. *P < 0.05, **P < 0.01, ***P <

0.001, ****P < 0.0001 and ns = not significant P > 0.05.

Figure 3

Figure 3. HFD blunted the mitochondrial oxidation of monocyte subsets. [A-C] Bone marrow cells

from mice fed control or HFD for 3, 8 or 18 weeks (n=3 mice per dietary group) were incubated with

Mitotracker Red CMXRos for 17 min at 37C. This is a fixable red-fluorescent dye that stains mitochondria

dependent on their membrane potential. Cells were then fixed, stained with fluorescent antibodies for flow

cytometry, and subjected to fluorescent activated cell sorting (FACS) to identify Ly6C^{low} and Ly6C^{high}

monocyte populations. [D-F] Ly6C^{low} and Ly6C^{high} monocytes from mice fed control or HFD for 3, 8 or 18

weeks, n=3 mice per dietary group) were isolated via FACS and plated for metabolic analysis of their

Oxygen Consumption Rate (OCR) with a Seahorse XF Analyzer. [G-I] Area under the curve of OCR of

FACS-sorted Ly6Clow and Ly6Chigh monocytes from control and HFD-fed mice measured by the

Seahorse Assay (n=3 mice per dietary group). [J] Area under the curve of ECAR measurement from FACS-

sorted Ly6C^{low} and Ly6C^{high} monocytes of control and HFD-fed mice measured by the Seahorse Assay and

pooled from all feeding durations. [K] BM lactate measurement of the supernatant fraction in HFD and

control samples quantified using a lactate Plus Meter and normalized to the initial volume of BM isolated

from the femur. (n=7 mice per dietary group). Results are expressed as mean ± SEM. One-way ANOVA

was used for statistical analysis. $*P < 0.05$, $**P < 0.01$ and ns = not significant $P > 0.05$. Graph in panels C

was created using FlowJo software (Becton, Dickinson Company).

Figure 4

Figure 4. HFD downregulated expression of mitochondrial fusion genes and upregulated

mitochondrial fission genes. Gene expression of mitochondrial fusion genes *Mfn1, Mfn2, Opa1, Tomm20,*

and *Tomm40* in monocytes isolated from control and HFD-fed mice for [A] 3, [B] 8, [C] 18 weeks. Gene

expression of mitochondrial fission genes *Drp1, CypD* and *Fis1* in monocytes isolated from control and

HFD-fed mice for [D] 3, [E] 8, [F] 18 weeks. (n=3 mice per dietary group and samples were analyzed in

duplicates). [G] Correlative analysis of monocyte subset BM cellular abundance and BM-derived monocyte

Mfn2 gene expression. [H] Correlative analysis of the average of monocyte subset BM cellular abundance

(n=8 per dietary group) and BM-derived monocyte *Drp1* gene expression (n=3 per dietary group) per

dietary group at each time point. Correlative analysis statistics were conducted using the linear regression

analysis testing whether slopes and intercepts are significantly different via GraphPad Prism version 6

(GraphPad Software). The r^2 represents the goodness of the fit while the P values greater than 0.05 represent

whether the slope of the line of best fit is significantly non-zero. Results are expressed as mean ± SEM.

One-way ANOVA was used for statistical analysis. *P < 0.05 and ns = not significant P > 0.05.

Figure 5

Figure 5. Bone marrow adipocyte whitening detected at 8 and 18 weeks of HFD favours increased bone marrow Ly6C^{high} monocytes and induced changes in mitochondrial dynamic gene expression. Volcano plots showing the gene expression of white and brown adipocyte markers in red and blue respectively [A] at 3, [B] 8 and [C] 18 weeks of HFD (n=3-9 per dietary group). Correlative graphs of the [D] white adipocyte marker ASC1 (n=8-9 per dietary group) and [E] brown adipocyte marker UCP1 (n=6 per dietary group) with Ly6Clow or Ly6Chigh monocyte subset abundance in the BM (n=8). Correlative analysis statistics were conducted using the linear regression analysis testing whether slopes and intercepts are significantly different via GraphPad Prism version 6 (GraphPad Software). The r^2 represents the goodness of the fit while the P values greater than 0.05 represent whether the slope of the line of best fit is significantly non-zero. [F-H] Flow cytometry quantification of naïve Ly6C^{low} and Ly6C^{high} monocytes (n=3-4 per dietary group) from *in vitro*-generated monocytes pre-treated for 18 h with pooled BMA-CM collected from three mice fed the control and HFD for 3, 8 and 18 weeks. [I] Gene expression of mitochondrial fusion genes *Mfn1, Mfn2, Opa1, Tomm20,* and *Tomm40* in naïve monocytes treated with BMA-CM from control and HFD-fed mice (n=3 per dietary group and samples were run in duplicates). [J] Gene expression of mitochondrial fission genes *Drp1, CypD* and *Fis1* in naïve monocytes treated with BMA-CM from control and HFD-fed mice (n=3 per dietary group and samples were run in duplicates). Results are expressed as mean ± SEM. One-way ANOVA was used for statistical analysis at each dietary group and across feeding periods. *P < 0.05, ****P < 0.0001 and ns = not significant > 0.05.

Figure 6

Figure 6. *Bona fide* white adipocytes favoured Ly6C^{high} monocytes preponderance, blunted monocyte mitochondrial membrane potential, and induced mitochondrial fragmentation *in vitro*. Naïve BM-derived monocytes were treated with CM from isolated *bone fide* white and brown adipocytes from EWAT and inter-scapular BAT for 18 h. [A] Flow cytometry dot plots detecting the effects of 18 h of treatment with media alone, pooled WATA-CM and BATA-CM from three mice (n=3 technical replicates each time with pooled CM from 3 mice) on monocyte subset abundance. [B] quantification of the gates drawn for Ly6C^{low} and Ly6C^{high} monocytes. (n=9 biological replicates from naïve monocytes treated). [C] Representative plot of MitoTracker Red CMXRos detected by flow cytometry in Ly6C^{low} monocytes treated with media alone, pooled WATA-CM and BATA-CM for 18 h. [D] Gene expression of mitochondrial fusion genes *Mfn1*, *Mfn2*, *Opa1*, *Tomm20*, and *Tomm40* in naïve BM-derived monocytes treated with control Media, pooled WATA-CM and BATA-CM. (n=6 mice in each CM group and samples were analyzed in duplicates). [E] Gene expression of mitochondrial fission genes *Drp1*, *CypD* and *Fis1* in naïve BM-derived monocytes treated with control Media, WATA-CM and BATA-CM. (n=6 mice in each CM group and samples were analyzed in duplicates). Results in [B], [D] and [E] are expressed as mean ± SEM, with one-way ANOVA used for statistical analysis: *$P < 0.05$, **$P < 0.01$ and ***$P < 0.001$, ****$P < 0.0001$ and ns = not significant $P > 0.05$. Graphs in [A] and [C] were created using FlowJo software (Becton, Dickinson Company). [F] Super-resolution microscopy images of mitochondrial morphology via TOM20 staining in BM-derived monocytes treated with control Media, pooled WATA-CM and BATA-CM (representative of n=3 mice in each CM group). Scale bar = $5\mu m$. Images were analyzed in Fiji (version 2.1.0/1.53c). [G] Schematic illustration of mitochondrial morphologies as fragmented, intermediate, or tubulated. [H] Quantification of the percentage of monocytes with the indicated mitochondrial morphology, in each treatment condition. Values are from three independent experiments; n=3 mice in each CM group; about 40 cells counted for each condition, within each experiment. Results are expressed as mean ± SEM. One-way ANOVA was used for statistical analysis. *$P < 0.05$, **$P < 0.01$ and ***$P < 0.001$, ****$P < 0.0001$ and ns = not significant $P > 0.05$.

Figure 7

Figure 7. Brown adipocytes favour Ly6C^{low} monocytes through progenitor cell differentiation and

proliferation. [A] BM-derived, *in vitro*-generated monocytes were treated with media alone, WATA-CM

or BATA-CM in the presence of the pyrimidine analogue BrdU to detect cellular proliferation. [B]

Quantification of BrdU abundance in Ly6C^{low} and Ly6C^{high} monocyte subsets with the treatments in panel

A, was determined by flow analysis (yellow to red signal, see Methods). Results are expressed as mean ±

SEM, n=3-6 individual mice per group. One-way ANOVA test was used for statistical analysis, ****$p <$

0.0005. [C, D] Colony Forming Unit assays were performed as described in Methods using six independent

cultures of BM-derived, *in vitro*-generated monocytes, (n=6 individual mice), were treated with WATA-

CM, BATA-CM or control adipocyte media (Media) for 18 h prior to mixing with methylcellulose-based

media and plating at 20,000 cells/ml to 6-well plates, in triplicate. [C] BATA-CM promoted CFU-GEMM

cultures over Media alone (**$p<0.01$ Two-way ANOVA, Tukey's multiple comparisons) and tilting

granulocyte/macrophage progenitors towards macrophage (CFU-M) over granulocyte (CFU-G)

differentiation, relative to Media alone (**$p<0.01$) or WATA-CM (*$p<0.05$ or ***$p<0.001$). [D] The counts

of the different colonies for each treatment (Media, WATA-CM or BATA-CM) were totaled and plotted as

the mean ± SEM (n=6 individual mice). There was no statistical difference between the media treatments.

[E] Isolated BM cells were processed for flow analysis to determine the progenitor cell populations present

in the Ly6C^{low} or Ly6C^{high} monocyte subsets. The percent GMP ratio was calculated relative to the total

cells detected in either monocyte subset as the mean ± SEM (isolated from n=3 individual mice). Graphs

in panel A were created using FlowJo software (Becton, Dickinson Company).

1048

Figure 8

Figure 8. WATA-CM tilts BM-derived Ly6C^{low} monocytes towards Ly6C^{high} monocytes.

Representative plots of FACS-sorted Ly6C^{low} (gray) and Ly6C^{high} (red) monocytes from control mice were

independently treated with [A] Media alone, [B] WATA-CM, or [C] BATA-CM (n=6 mice per FACS-

sorted monocyte group) for 18 h after which the presence of the alternate monocyte subset was assessed

via flow cytometry. [D-F] Quantification of Ly6C^{low} and Ly6C^{high} gates in panels A-C. [G] BM-derived, *in*

vitro-generated monocytes were analyzed for gene expression of the Ly6C^{low} marker SPN and the Ly6C^{high}

marker Ly6C2 by qPCR as described in the Methods.

Graphs in panels A-C were created using FlowJo software (Becton, Dickinson Company).

1057

Figure S1

A

B

C

D

SUPPLEMENTAL MATERIAL

Figure S1: Flow cytometry gating strategy for identification of Ly6C^{low} and Ly6C^{high} monocytes and

their progenitor cells. Following red blood cell lysis, SVF, blood and the BM were washed with FACS

buffer (PBS, 1% (v/v) heat-inactivated FBS, 0.2% (w/v) BSA), stained with Aqua Live/Dead Antibody

(ThermoFisher, L34957) for 15 min in the dark, washed with FACS buffer again, blocked with anti-mouse

CD16/32 antibody (eBioscience, 14-0161-86) for 15 min in the dark following a 30 min incubation with

the indicated antibodies. [A] Using side and forward scatter to identify singlet cells, CD115+Gr1- cells

were identified that were further analyzed using CD11b and Ly6C. [B] Ly6C^{low} and Ly6C^{high} monocyte

gates were drawn using fluorescence minus one (FMO) controls for CD11b and Ly6C. [C] The subsets

were further confirmed via CX3CR1 and CCR2, respectively. [D] CD45+ cells were analyzed using a

lineage cocktail of antibodies (**Table 1**) and CD117+, further broken down into Sca1+ and – cells. Sca1+

cells were further classified using CD48 and CD150 while Sca1- cells were further characterized via

CD16/32 and CD34. Graphs were created using FlowJo software (Becton, Dickinson Company).

1071

Figure S2

1072   **Figure S2. Monocyte progenitors and stem cells were not significantly altered with HFD *in vivo*.** Flow

1073   cytometry analysis of HSC and monocyte progenitor cells in mice fed a Control or HFD for [A] 3, [B] 8,

1074   and [C] 18 weeks. (n = 6-7 per dietary group at each feeding duration) Flow cytometry antibodies used are

1075   listed in Table 1. Student's T test was used and no statistical significance was observed (ns = not significant

1076   > 0.05).

1077

Figure S3

A 3 weeks BM Supernatant

B 8 weeks BM Supernatant

C 18 weeks BM Supernatant

D Sorted Monocyte Subsets TNFa

Figure S3: BM pro-inflammatory factors did not change with HFD. No changes in pro-inflammatory

cytokines were detected in BM supernatant fraction using the ELISA assay in mice fed control or HFD for

[A] 3, [B] 8, or [C] 18 weeks. (n = 3-6 mice in each dietary group at feeding duration) [D] FACS-sorted

monocyte subsets were analyzed for changes in pro-inflammatory gene expression; shown in TNFα gene

expression increased with 18 weeks of HFD in Ly6C^{high} monocytes, but no other changes were detected. (n

= 3 mice in each dietary group at each feeding duration) Results are expressed as mean ± SEM. One way

ANOVA test was used for statistical analysis. *P < 0.05, **P < 0.01 and ns = not significant P > 0.05.

1085

Figure S4

A

3 weeks

B

8 weeks

C

18 weeks

D

WATA

E

BATA

Figure S4. BMA precursor cells with HFD. RT qPCR of isolated pelleted fraction were assessed for pre-

adipocyte markers of mice fed a Control or HFD for [A] 3, [B] 8, and [C] 18 weeks. (n=3 mice per dietary

group and samples were analyzed in duplicates). Results are expressed as mean ± SEM. One way ANOVA

test was used for statistical analysis. *P < 0.05, **P < 0.01 and ns = not significant P > 0.05. [D] classical

white and brown adipocyte markers were confirmed in isolated WATA and BATA. (n=2 mice per tissue

and samples were analyzed in duplicates) No statistical analysis was conducted on data presented in panel

D as the brown adipocyte genes were undetectable in WATA and vice versa.

1093

1094

Figure S5

1095 **Figure S5: HSC and their progenitors following the white and brown CM treatment for 18 h.** [A-C]

1096 Flow cytometry analysis was used to test for the presence and changes in HSC and monocyte progenitor

1097 cells in BM-derived, *in vitro*-generated BM-derived monocytes treated with media, WATA-CM, and

1098 BATA-CM. Flow cytometry antibodies used are listed in Table 1 (n=3 individual mice per CM group). One

1099 way ANOVA test was used for statistical analysis. *P < 0.05 and ns = not significant P > 0.05.

1100

Figure S6

Figure S6: Flow cytometry gating strategy for identification of progenitor cells within Ly6C^low and

Ly6C^high monocyte subsets. To prevent contamination of blood in BM cells, mice (n=3 individual mice)

were placed under deep anesthesia using isoflurane and underwent intra-cardiac perfusion with 10 mL of

cold PBS, before sacrifice. Femurs were surgically removed and crushed with a mortar and pestle and bone

marrow cells suspended in PBS were filtered through a 40-μm-nylon mesh. Cells were pelleted at $300 \times g$,

5 min and resuspended in 500 μl FACS Buffer. 100 μl was aliquoted and stained with the antibody cocktail

for 30 min. Antibodies for CD117, CD16/32 and CD34, were used at 1:50 dilution, all others at 1:100

dilution. Samples were fixed (BD, Cytofix) before being run on a BD LSR Fortessa cytometer. [A] Gating

strategy for Ly6C^low and Ly6C^high. [B] Gating strategy for progenitor cell identification in the Ly6C^high

monocytes. [C] Gating strategy for progenitor cell identification in the Ly6C^low monocytes.

1111

Table 1. List of Antibodies used for flow cytometry analysis.

Markers	Fluorophore	Manufacturer	Clone	Catalog Number
CD11b	APC-eFluor 780	eBioscience	M1/70	47-0112-82
Ly6C	PE-Cy7	Biolegend	HK1.4	128018
Gr-1	Alexa Fluor 700	eBioscience	RB6-8C5	56-5931-82
CX3CR1	Brilliant Violet 421	Biolegend	SA011F11	149023
CCR2	Alexa594	R&D Systems	475301	FAB5538T
CD115	APC	eBioscience	AFS98	17-1152-82
F4/80	PerCP-Cy5.5	BioLegend	BM8	123128
CD11c	FITC	eBioscience	N418	11-0114-81
CD34	FITC	BD Biosciences	RA1434	553733
CD16/32	APC-R700	BD Biosciences	2.4G2	565502
CD45.2	BV650	BD Biosciences	30-F11	563410
CD150	PEPCP710	invitrogen	mShad150	46-1502-82
CD117	APC/Fire 750	BioLegend	2B8	105838
SCA1	PE-Cy7	Invitrogen	D7	25-5981-82
CD48	BV605	BioLegend	MM48-1	103441
CD3e*	PE	BD Biosciences	17A2	555725
B220*	PE	BD Biosciences	RA3-6B2	553089
Ly6G*	PE	BD Biosciences	1A8	551461
CD11b*	PE	BD Biosciences	M1/70	553311
CD11c*	PE	BioLegend	N418	117308
Ter119*	PE	BD Biosciences	TER-119	553623
CD127*	PE	eBioscience	A7R34	12-1271-82
NK1.1*	PE	BD Biosciences	PK138	553165

*LINEAGE is a cocktail of antibodies containing CD3e, B220, Ly6G, CD11b, CD11c, Ter119, CD127, NK1.1 with the same fluorophore to exclude all terminally differentiated cells.