Title (12 words; 93 characters with spaces)

Bone marrow adipocytes drive the development of tissue invasive Ly6C[high] monocytes during obesity

Running Title: Bone marrow fat linked to Ly6C[high] monocytes in obesity

Parastoo Boroumand[1,2], David Prescott[3,4], Michael Wong[1], Jeff Shen[1], Ivan Tattoli[4], Yuhuan Zhou[1], Nancy

Shi[1], Lucie Y Zhu[1], Zhi Liu[1], Angela Li[3], Tapas Mukherjee[3], Philip J. Bilan[1], Dana J. Philpott[3], Stephen E.

Girardin[3,4], Amira Klip[1,2*].

[1]Cell Biology Program, The Hospital for Sick Children, Toronto, Ontario, M5G 0A4; Canada;

[2]Department of Biochemistry, University of Toronto, Toronto, Ontario, M5S 1A8; Canada;

[3]Department of Immunology, University of Toronto, Toronto, Ontario, M5S 1A8; Canada;

[4]Department of Laboratory Medicine and Pathophysiology, University of Toronto, Toronto, Ontario, M5S 1A8;

Canada

*Corresponding Author:

Amira Klip, Ph.D.

Cell Biology Program

Hospital for Sick Children

Toronto, Ontario M5G 0A4, Canada

Phone: 416-813-6392

E-mail: amira@sickkids.ca

Summary (150 Words, 150):

During obesity and high fat-diet (HFD) feeding in mice, sustained low-grade inflammation includes not

only increased pro-inflammatory macrophages in the expanding adipose tissue, but also bone marrow (BM)

production of invasive Ly6C^{high} monocytes. As BM adiposity also accrues with HFD, we explored the

relationship between the gains in BM white adipocytes and invasive Ly6C^{high} monocytes *in vivo* and

through *ex vivo* paradigms. We find a temporal and causal link between BM adipocyte whitening and the

Ly6C^{high} monocyte surge, preceding the adipose tissue macrophage rise during HFD. Phenocopying this,

ex vivo treatment of BM cells with conditioned media from BM adipocytes or from *bona fide* white

adipocytes favoured Ly6C^{high} monocyte preponderance. Notably, Ly6C^{high} skewing was preceded by

monocyte metabolic reprogramming towards glycolysis, reduced oxidative potential and increased

mitochondrial fission. In sum, short-term HFD changes BM cellularity, resulting in local adipocyte

whitening driving a gradual increase and activation of invasive Ly6C^{high} monocytes.

Keywords:

Introduction:

A continuum of chronic obesity leading to metabolic disorders such as insulin resistance and eventually type 2 diabetes is of paramount concern worldwide (Centers for Disease Control and Prevention, 2017; Hamann, 2017; WHO, 2016). An important contributor to the development and exacerbation of insulin resistance is a gain in macrophages in adipose tissue (AT) (Liu et al., 2019; Weisberg et al., 2003), muscle (Fink et al., 2014; Garvey et al., 2008) and liver (Lanthier et al., 2010) as well as their enhanced pro-inflammatory (Lumeng et al., 2007, 2008; De Souza et al., 2005) and glycolytic skewing (Sharma et al., 2020). This low-grade inflammation is partly explained by circulating monocyte infiltration into tissues where they differentiate to pro-inflammatory macrophages (Kohyama et al., 2016).

Ly6C^{high} monocytes in mice, and classical CD14^{++} CD16$^-$ monocytes in humans, constitute the predominant tissue-infiltrating monocyte subset during chronic inflammation. These tissue-invasive monocytes are classified by high surface abundance of the Ly6C protein, the chemokine receptor CCR2 and the adhesion molecule L-selectin, all of which allow for monocyte extravasation from the BM and infiltration into the tissues (Auffray et al., 2007; Boniakowski et al., 2018; Fink et al., 2014; Geissmann et al., 2003; Weisberg et al., 2006). With obesity, the development of Ly6C^{high} monocytes along with their precursor cells in the BM is boosted (Griffin et al., 2018; Manicone et al., 2016; Singer et al., 2014) leading to a rise in available circulating Ly6C^{high} monocytes for tissue-infiltration (Friedrich et al., 2019; Singer et al., 2014).

Various obesity-driven factors amalgamate for greater Ly6C^{high} monocyte infiltration into the AT including (a) increased AT-dependent chemoattraction (Kamei et al., 2006; Nagareddy et al., 2014; Nov et al., 2013; Renovato-Martins et al., 2017) and (b) enhanced BM myelopoiesis (Griffin et al., 2018; Manicone et al., 2016; Singer et al., 2014). However, it is not clear whether the obesity-induced BM myelopoiesis occurs prior to, concurrently, or after the elevation of AT macrophages. Since the BM is a highly-vascularized tissue, home to hematopoietic and mesenchymal lineages, and is dynamically reactive to environmental changes, we reasoned it might be an early responder to HFD-induced obesity, producing increased Ly6C^{high} monocytes prior to their accumulation in the AT. Further, since BM-resident adipocytes

undergo hyperplasia and hypertrophy during obesity (Boroumand and Klip, 2020; Doucette et al., 2015;

Scheller et al., 2015; Tratwal et al., 2020) we contemplated that this process might be temporally and

potentially causally related to the HFD-driven BM myelopoiesis and Ly6C^{high} monocytosis. Finally, we

considered whether the response of the monocytic lineage to the BM environment during obesity includes

a metabolic shift leading to the increase in Ly6C^{high} monocytes. In fact, and surprisingly, little is known

about the metabolic orientation of the Ly6C^{high} monocytes in states of obesity, although there is mounting

evidence of a link between a macrophage-intrinsic metabolic shift and their pro-inflammatory orientation

in this condition (O'Neill et al., 2016; Sharma et al., 2020; Thapa and Lee, 2019; Verdeguer and Aouadi,

2017).

Hence, the overall hypothesis driving this study is that, with obesity, the increased production of

Ly6C^{high} monocytes in the BM is accompanied and possibly caused by a glycolytic shift in these cells,

driven by changes in bone marrow adipocytes (BMA). Accordingly, we investigated the temporality of

enhanced BM myelopoiesis, BM Ly6C^{high} monocyte accumulation, and AT macrophage changes during

high fat diet (HFD) feeding in mice. Next, we explored the potential causal relationship between the obesity-

induced changes in BMA and increased Ly6C^{high} monocyte production. Using an *in vitro* platform, we

deconstructed the effects of obesity-induced BM cellular changes on Ly6C^{high} monocyte expansion. We

report that BM Ly6C^{high} monocytes accumulate in the BM prior to the gain in pro-inflammatory AT

macrophages. The rise in BM monocytes temporally correlated with the elevation in BMA. Moreover,

BMA whitening correlated with the timely increases in BM Ly6C^{high} monocytes, and most notably, isolated

BMA from HFD-fed mice directly favoured Ly6C^{high} monocyte development *ex vivo*. This monocyte

developmental change was also induced *ex vivo* by *bona fide* white adipocytes while *bona fide* brown

adipocytes favoured Ly6C^{low} monocyte predominance. Notably, the conversion of Ly6C^{low} to Ly6C^{high}

monocytes was preceded by a switch in their metabolism, from oxidative to glycolytic. Mechanistically,

white adipocytes induced conversion of Ly6C^{low} to Ly6C^{high} monocytes, whereas brown adipocytes

increased Ly6C^{low} monocyte proliferation. We propose that HFD-induced changes in BMA disrupts the

normal balance of BM monocyte subsets leading to a higher output of tissue invasive Ly6C^{high} monocytes.

Results

Increased fasting blood glucose, peripheral and BM adiposity by 3 weeks of HFD

To study the temporal relationship of HFD-induced myeloid changes in the BM relative to that in epididymal white AT (EWAT), C57BL/6 mice were fed HFD or control low fat diet (LFD) for either 3, 8 or 18 weeks. HFD-fed mice had higher body weight and impaired fasting blood glucose by 3 weeks, compared to controls (**Figure 1A,B**). Body weight and hyperglycemia continued to rise with prolonged feeding. HFD-fed mice also displayed higher EWAT weight (**Figure 1C**) and inter-scapular brown AT (BAT) weight at 3 weeks (not shown). Cross-sectional tibia and femur bone histology visualized via H & E and Oil Red O staining revealed an increase in BMA size by 3 weeks and an increase in BMA number by 8 weeks of HFD (**Figure 1D-G**). This indicated that 3 weeks of HFD sufficed to induce whole body metabolic disturbance and augmented adiposity in peripheral tissues and in the BM.

BM monocytes increase prior to accumulation of EWAT macrophages and monocytes with HFD

Although total EWAT weight increased by 3 weeks, myeloid cells in the stromal vascular fraction (SVF) of AT did were not elevated until 8 weeks of HFD, as revealed by gene expression of the general myeloid marker *Itgam* (encoding CD11b) and monocyte markers Gr1$^-$ CD115$^+$ CD11b$^+$ Ly6C$^+$ (determined by flow cytometry) (**Figures 2A,D**). Similarly, AT macrophages, assessed by gene expression of the AT macrophage marker *Adgre1* (encoding F4/80) and by flow cytometry counting of CD11b$^+$ Ly6C$^-$ F4/80$^+$ macrophages in the SVF were not elevated at 3 weeks of HFD but were at 8 weeks (**Figure 2B,E**). Further, gene expression of the pro-inflammatory myeloid marker *Itgax* (encoding CD11c) and surface detection of the pro-inflammatory markers CD11b$^+$ Ly6C$^-$ F4/80$^+$ CD11c$^+$ were also higher in the SVF by 8 weeks of HFD relative to control diet, but not at 3 weeks (**Figure 2C,F**). Of note, gene expression of the embryonically-conserved tissue-resident AT macrophage markers *Timd4* and *Marco* (Beattie et al., 2016; Gibbings et al., 2015) was not elevated at 3 or at 8 weeks and only rose at 18 weeks of HFD (**Figure 2G-H**). This indicates that the gain in AT macrophages at 8 weeks of HFD is unlikely due to proliferation of resident macrophages and instead likely results from monocyte recruitment from the circulation.

In the BM, total monocytes, gated via the Gr1$^-$ CD115$^+$ CD11b$^+$ Ly6C$^+$ markers, increased at 3

weeks and continued to rise by 8 and 18 weeks of HFD (**Figure 2I**). Using the surface abundance of the

Ly6C protein as well as the CX3CR1 and CCR2 markers (Auffray et al., 2007, 2009; Kratofil et al., 2017),

we subcategorized Ly6C^{low} and Ly6C^{high} monocyte subsets. At 3 weeks into the feeding study, Ly6C^{low} and

Ly6C^{high} monocytes represented about 40% of the total BM monocyte population in the control diet group,

whereas HFD promoted a higher proportion of Ly6C^{low} monocytes (**Figure 2J**). However, by 8 weeks of

HFD there was a marked switch, whereby the relative proportion of Ly6C^{low} monocytes in the total

monocyte population was *lower* and that of Ly6C^{high} monocytes was *higher* than under the control diet

(**Figure 2J**). These differences were more prominent by 18 weeks of HFD feeding (**Figure 2J**).

Collectively, these findings revealed that, in the BM, HFD favoured monocytosis at 3 weeks, with a

subsequent and prominent skewing towards tissue invasive Ly6C^{high} monocytes at 8 weeks that continued

into 18 weeks of HFD. Singer et al. also demonstrated an increase in BM monocytosis and elevated

monocyte precursor cells with 18 weeks of HFD, that persisted after BM transplantation into chow-fed

recipients (Singer et al., 2014). In the present study, while we clearly recorded an earlier and continuing

rise in total monocytes in HFD-fed mice (**Figure 2I**), we did not observe statistically significant increases

in monocyte precursor and stem cells at any time point analyzed (**Figure S1A-C**).

Glycolytic metabolic reprograming in monocyte subsets with HFD

Recent discoveries have highlighted the importance of glycolytic metabolism in the activation of

AT macrophages with HFD (Lee and Murphy, 2019), but the metabolic status of monocytes in obesity has

not been reported, although some studies have investigated the metabolism of circulating human

CD14$^+$CD16$^-$ monocytes (homologous to murine tissue invasive Ly6C^{high} monocytes) during aging (Lee

and Murphy, 2019; Pence and Yarbro, 2019). To assess whether HFD affects the metabolic status of BM

monocyte subsets in obesity, we analyzed Ly6C^{low} and Ly6C^{high} from control and HFD mice at 3, 8 and 18

weeks of feeding. The mitochondrial activity-indicator dye MitoTracker Red CMXRos was incorporated

into the monocyte flow cytometry panel (Table 1). MitoTracker Red CMXRos is recruited to the

mitochondrial membrane proportionally to the degree of mitochondrial membrane potential and is thus an

indicator of mitochondrial activity in oxidative phosphorylation. The results revealed that $Ly6C^{low}$

monocytes have a higher mitochondrial membrane potential than $Ly6C^{high}$ monocytes in the control mice

(**Figure 3A-C**). Notably, 3 weeks of HFD reduced the mitochondrial membrane potential in both monocyte

subsets indicated by the left-ward shift in the MitoTracker Red CMXRos peak (**Figure 3A**). The HFD-

induced effects were more pronounced by 8 and 18 weeks of HFD (**Figure 3B,C**). To further understand

the effects of HFD on the developing BM monocytes, $Ly6C^{low}$ and $Ly6C^{high}$ monocytes from mice fed

control or HFD for 3, 8 or 18 weeks were isolated via fluorescent activated cell sorting (FACS) and plated

for metabolic analysis with a Seahorse XF Analyzer. The results revealed a gradual lowering in the oxygen

consumption rate (OCR) of $Ly6C^{low}$ monocytes with HFD (**Figure 3D-I**). Comparison of the area under

the curve (AUC) revealed that at 18 weeks, the OCR of the $Ly6C^{low}$ monocytes was significantly lowered

with HFD (**Figure 3I**). Interestingly, at 8 and 18 weeks, the AUC of the OCR of the $Ly6C^{low}$ monocytes

was significantly higher than the AUC of the OCR of $Ly6C^{high}$ monocytes in the control group (**Figure 3H-**

I). Moreover, this difference was lost in the HFD group as the AUC of the OCR of $Ly6C^{low}$ monocytes

diminished (**Figure 3H-I**).

158   Comparison of the various sections of the OCR graph revealed that the changes in OCR observed

with 3 weeks of HFD were not manifest at basal respiration nor at the ATP-dependent respiration, rather

they were evident in the maximal respiratory capacity portion of the metabolic stress test (**Figure 3D**). A

drop in basal respiration of both monocyte subsets emerged with 8 weeks of HFD and became more

pronounced with 18 weeks of HFD (**Figure 3E-F**). This showcased the sustained effects of HFD on

monocyte metabolic status. Conversely, the extracellular acidification rate (ECAR) of $Ly6C^{low}$ monocytes

was higher than that of $Ly6C^{high}$ monocytes, consistent with a more active metabolism in the former (**Figure**

3J). However, this difference was lost with HFD, as ECAR dropped in the $Ly6C^{low}$ monocytes and rose in

the $Ly6C^{high}$ monocytes relative to the control diet group (**Figure 3J**). Overall, these results suggest that,

along with promoting the preponderance of $Ly6C^{high}$ monocytes, HFD also lowers OCR in $Ly6C^{low}$

monocytes.

Notably, an analysis of the BM revealed elevated lactate levels across all HFD time points (**Figure**

3K). This increase in lactate may be due to the glycolytic shift in BM monocytes described above, among

other potential sources within the BM cellular composition. Further analysis of the BM soluble fraction

failed to evince any changes between control and 18-week HFD-fed mice in pro-inflammatory cytokines

TNFα, IL-1β, IL-6, CXCL1 or CCL2 (**Figure S2A-C**). Similarly, there were no differences between diet

groups in pro-inflammatory gene expression of *IL-6*, *IL-1β* or *Itgax* in FACS-sorted Ly6C^{low} and Ly6C^{high}

monocytes (Not Shown). However, *TNFα* gene expression was higher in FACS-sorted Ly6C^{high} monocytes

from 18-week HFD-fed mice compared to controls (**Figure S2D**). Collectively, these findings reveal that

HFD alters the BM metabolic micro-environment with little effect on pro-inflammatory cytokine secretion

(based on the panel tested).

Expression of mitochondrial fusion and fission genes are altered monocytes with HFD

The efficiency of mitochondrial function is largely reliant on its structure. Mitochondria are highly

dynamic in structure, largely regulated by concentration of proteins that control fission and fusion (Gottlieb

and Bernstein, 2017). Dynamic transitions are closely aligned to metabolic performance. To further assess

the mitochondrial activity of monocytes, we enquired whether their expression of mitochondrial dynamic

genes is altered with HFD compared to the control group.

The analyzed genes involved in mitochondrial fusion were *Mfn1*, *Mfn2*, *Opa1*, *Tomm20* and

Tomm40, while those involved in mitochondrial fission were *Drp1*, *CypD* and *Fis1*. BM monocytes from

mice fed HFD for 3 weeks did not present with differences in gene expression in the tested mitochondrial

fusion genes compared to control diet counterparts (**Figure 4A**). BM monocytes isolated from mice fed

HFD for 8 weeks showed downregulated *Mfn1* and *Mfn2* compared to the controls (**Figure 4B**). BM

monocytes isolated from 18 week HFD-fed mice downregulated the mitochondrial fusion genes *Mfn2*,

Opa1, and *Tomm20* (**Figure 4C**). Furthermore, from the tested mitochondrial fission genes, *Drp1* was

upregulated in the BM monocytes of 3 weeks HFD-fed mice (**Figure 4D**). *Drp1* and *CypD* were also

upregulated in BM monocytes of 8 weeks HFD-fed mice (**Figure 4E**). At 18 weeks, all three tested

mitochondrial fission genes, *Drp1*, *CypD* and *Fis1* were upregulated (**Figure 4F**). This gene expression

pattern suggests that HFD skews the monocyte gene expression to favour downregulation of genes involved

in mitochondrial fusion and upregulation of genes involved in fission.

Given that Ly6C^{low} and Ly6C^{high} monocytes presented with differences in metabolic preference

(glycolytic and oxidative, respectively), and the known link of glycolytic metabolism with mitochondria

fragmentation and conversely oxidative metabolism to mitochondrial fusion, we assessed whether the

expression patterns of the genes governing mitochondrial fission and fusion would correlated distinctly

with the BM monocyte subset abundance in control and HFD-fed mice. Indeed, and notably, the abundance

of the Ly6C^{high} monocyte subset negatively correlated with *Mfn2* and positively correlated with *Drp1* levels

(**Figure 4G-H**). In contrast, the abundance of BM Ly6C^{low} monocytes did not correlate with the expression

patterns of either *Mfn2* or *Drp1* (**Figure 4G-H**). These data are consistent with our metabolic findings

suggesting Ly6C^{high} monocytes are more glycolytic and adopt a gene expression pattern that would promote

mitochondrial fragmentation.

A shift in BMA type at 8 weeks of HFD influences monocyte subset preference and mitochondrial

dynamic gene expression

Hematopoietic myeloid cells reside in specific niches of the BM that determine their cellular

interaction with other co-resident BM cells (Boroumand and Klip, 2020). BMA, while representing a unique

type of adipocytes, respond to obesity with an increase their white adipocyte-like fat storing properties and

a decrease in brown-like characteristics (Ambrosi et al., 2017; Doucette et al., 2015; Krings et al., 2012;

Scheller et al., 2016). To explore the temporal relation of BMA whitening in relation to the development

of monocyte subsets, we followed the gene expression of a suite of recently identified markers that are good

differentiators of adipocyte types (Nascimento et al., 2014; Ussar et al., 2014) in isolated BMA from femur

and tibia of control and HFD-fed mice. By 3 weeks into HFD, the white adipocyte marker Asc1 (encoding

the guanine nucleotide-binding protein subunit beta-like protein) was significantly higher and the brown

adipocyte marker P2rx5 (encoding purinergic receptor P2X) trended to be lower albeit without reaching

statistical significance, relative to control diet. White adipocyte markers *Lep* (leptin) and *AdipoQ*

(adiponectin) as well as the brown adipocyte markers *Ucp1* (UCP1) and *Pat2* (Integrin alpha pat-2) did not

change (**Figure 5A**). By 8 weeks of HFD, *Asc1* mRNA remained elevated and *Lep* and *AdipoQ* trended in

the same direction (**Figure 5B**). Brown adipocyte markers *P2rx5* and *Ucp1* in the isolated BMA of HFD

mice were markedly lower relative to those from control diet mice (**Figure 5B**). Thus, 8 weeks of HFD

sufficed to shift BMA towards white adiposity. By 18 weeks of HFD, all white adipocyte markers genes

showed were upregulated while brown adipocyte markers were downregulated compared to the control

group (**Figure 5C**). Analyses of adipocyte differentiation and progenitor cell markers were conducted in

the BM pelleted fraction. *Hoxc8*, a regulator of adipocyte progenitors (Nakagami, 2013), was significantly

upregulated at 8 weeks along with *Hoxc9* at 18 weeks of HFD (**Figure S3B,C**). The beige adipocyte

progenitor marker *Tmem26* was not significantly higher at any of the tested HFD durations (**Figure S3A-**

C). Expression of the brown adipocyte determinant genes *Tbx1* and *Lhx8* was undetectable in the isolated

BMA. Thus, during the course of HFD, while markers for white adipocyte progenitor cells were

upregulated, precursors of beige/brown adipocytes were not affected.

The rise in BM Ly6C^{high} monocytes and the shift in BMA whitening both manifested by 8 weeks

of HFD. Correlative analysis was performed for both Ly6C^{low} and Ly6C^{high} with the white adipocyte marker

Asc1 and the brown adipocyte marker *Ucp1*. The results revealed a significant positive correlation (P value

of 0.0123) for Ly6C^{high} monocytes with *Asc1* gene expression (**Figure 5D**). Conversely, the surge in

Ly6C^{high} monocytes presented a significant negative correlation (P value of 0.0415) relative to *Ucp1* gene

expression (**Figure 5E**). Changes in Ly6C^{low} monocytes did not significantly correlate with *Asc1* (P value

of 0.0768) or *Ucp1* (P value of 0.1299) (**Figure 5D-E**). Collectively, the above data suggest the possibility

that the mere shift in BMA towards white adiposity may directly favour induction of BM Ly6C^{high}

monocytes. To test this hypothesis, we generated conditioned media (CM) from BMA (BMA-CM) pooled

from three control and three HFD-fed mice, collected 3, 8 and 18 weeks into the diets. In parallel, BM of

control mice was used to generate monocytes through a 5-day *in vitro* differentiation protocol. These

monocytes were treated with BMA-CM from control and HFD-fed mice for 18 hours thereafter.

Subsequently, the monocyte population was analyzed via flow cytometry. BMA-CM from 3 weeks HFD-

fed mice had no effect on the monocyte subset proportions compared to BMA-CM from control diet-fed

mice (**Figure 5F**). Strikingly, however, BMA-CM from 8-week HFD-fed mice applied to BM-derived

monocytes from control mice resulted a significant production of Ly6C^{high} monocytes compared to the

effect of BMA-CM from mice on the 8-week control diet (**Figure 5G**). This skewing was further enhanced

when exposing the control BM cells to BMA-CM from 18-week HFD-fed mice (**Figure 5H**).

Lastly, we tested whether BMA-CM was capable to induce changes in gene expression of the

mitochondrial fusion and fission genes in monocytes, as observed with monocytes isolated from HFD-fed

mice. Accordingly, *in vitro-generated* monocytes were treated with BMA-CM from control and HFD-fed

mice. As before, the analyzed genes involved in mitochondrial fusion were *Mfn1*, *Mfn2*, *Opa1*, *Tomm20*

and *Tomm40*, while the tested genes involved in mitochondrial fission were *Drp1*, *CypD* and *Fis1*.

Monocytes treated with BMA-CM from HFD-fed mice for 18 weeks showed significant downregulated

mitochondrial fusion genes *Mfn2*, *Tomm20* and *Tomm40* (**Figure 5I**). Conversely, the mitochondrial fission

genes *Drp1* and *CypD* were significantly up-regulated in monocytes treated with HFD-derived BMA-CM

(**Figure 5J**).

These results not only show that HFD-derived BMA-CM can skew monocyte subset cellularity but

also can alter mitochondrial morphology via changes in mitochondrial dynamic gene expression. These

changes can have consequential effects on the metabolic properties on the monocytes.

White adipocytes favour Ly6C^{high} monocytes and promote mitochondrial fragmentation

To assess whether the effects of BMA on monocyte subset skewing and metabolic differences can

be attributed to BMA whitening, we designed a proof of concept experiment studying the influence of *bona*

fide white or brown adipocytes on monocyte subset determination and metabolic changes *ex vivo*. Isolated

adipocytes from EWAT and brown adipocyte tissue (BAT) of three mice were pooled to generate white AT

adipocytes (WATA) and brown AT adipocytes (BATA), respectively. Confirming the phenotype of the

isolated cells, classical white adipocytes markers were solely expressed in the isolated WATA and classical

brown adipocyte markers were exclusively expressed in the BATA (**Figure S3D-E**). The isolated WATA

and BATA were cultured for 24 hours and the corresponding CM were collected for subsequent monocyte

treatment. BM-derived monocytes from chow-fed mice were treated with WATA-CM, BATA-CM or left

untreated. The proportion of Ly6C^{low} and Ly6C^{high} monocyte subsets was determined by flow cytometry

analysis (**Figure S4A-C**). This *in vitro* paradigm allowed for direct mechanistic analysis of the influence

of white or brown adipocytes on monocytes. These results indicate that WATA-CM favoured Ly6C^{high}

monocyte predominance, while BATA-CM propitiated Ly6C^{low} monocyte abundance (**Figure 6A-B**),

revealing a novel, differential influence of white and brown adipocytes on monocyte orientation preference.

Moreover, CCR2 surface receptors also increased in the monocytes incubated with WATA-CM (Not

Shown), validating the effects of white adipocytes in promoting an abundance of tissue invasive Ly6C^{high}

monocytes.

Since segregated Ly6C^{low} and Ly6C^{high} monocyte subsets had different metabolic orientations, we

investigated if the differential adipocyte influence on monocyte subset development corresponded to a

change in their metabolic status. To this end, *in vitro*-generated monocytes were treated with WATA-CM,

BATA-CM or left untreated for the last 18 hours of the monocyte differentiation protocol. The

mitochondrial membrane potential was assessed using MitoTracker Red CMXRos and flow cytometry

analysis. Remarkably, Ly6C^{low} monocytes incubated *in vitro* with WATA-CM experienced a left-ward shift

in MitoTracker Red CMXRos peak revealing a reduction in mitochondrial membrane potential induced by

the WATA-CM (**Figure 6C**), reproducing what was observed with monocytes from HFD-fed mice (see

Figure 3A-C). Furthermore, the gene expression of mitochondrial fusion genes *Mfn1*, *Mfn2*, *Tomm20* and

Tomm40 were reduced in monocytes treated with WATA-CM (**Figure 6D**). The mitochondrial fission

genes *Drp1*, *CypD* and *Fis1* were up-regulated in monocytes treated with WATA-CM (**Figure 6E**).

Conversely, treating monocytes with BATA-CM downregulated the mitochondrial fission gene *CypD*

(**Figure 6E**). To explore if the changes in mitochondrial fusion and fission gene expression correlated with

the corresponding mitochondrial morphology, we utilized super-resolution microscopy to image the

mitochondrial protein TOM20 in *in vitro* monocytes adhered to coverslips. Compared to control-medium

treated monocytes, WATA-CM induced a striking mitochondrial fragmentation while BATA-CM

promoted a more complete mitochondrial elongation (**Figure 6F**). These observations suggest that white

adipocytes, which favour Ly6C^{high} monocyte abundance, can alter the metabolic status of Ly6C^{low}

monocytes by reducing their mitochondrial membrane potential and altering mitochondrial dynamic gene

expression and morphology, thereby linking changes in BMA whitening to monocyte metabolic status.

White and brown adipocytes influence monocytes through distinct mechanisms

Finally, we investigated the mechanisms whereby monocytes undergo the change in subsets in

response to *bona fide* white and brown adipocyte media. Although we had not detected HFD-induced

changes in hematopoietic stem and monocyte precursor cells *in vivo*, we re-assessed the abundance of these

populations upon exposing *in vitro*-generated monocytes to WATA-CM and BATA-CM. The abundance

of hematopoietic stem and monocyte precursor cells was determined using appropriate surface markers and

flow cytometry analysis (**Table 1, Figure S4D**). A gain in myeloid precursor cell (CMP) was observed in

response to WATA-CM treatment (**Figure S5A-C**). However, no differences in the abundance of CD117+,

Sca1+, Sca1-, GMP and MEP cells were detected (**Figure S5A-C**). Given these results we investigated

other mechanisms by which monocyte subsets may be altered in response to WATA-CM and BATA-CM.

We first explored whether the proportion of monocyte subsets was altered through conversion of

one subset to another. Ly6C^{high} monocytes have been shown to give rise to Ly6C^{low} monocytes in mouse

blood and BM, and this is paralleled by the human homologous CD14 and CD16 monocyte subsets (Kratofil

et al., 2017; Lessard et al., 2017; Patel et al., 2017). However, while there has been speculation about the

conversion of Ly6C^{low} to Ly6C^{high} monocytes, there is little experimental evidence for this phenomenon.

To explore this possibility, we first separately isolated Ly6C^{low} and Ly6C^{high} monocytes through FACS.

Each FACS-sorted homogenous monocyte population was treated with WATA-CM, BATA-CM or left

untreated for 18 hours. Following treatments, we evaluated the presence of the alternative monocyte

subtype relative to that in the respective starting homogenous populations. Surprisingly, no conversion from

Ly6C^{high} to Ly6C^{low} monocytes was observed in any condition (**Figure 7A-F**), but, remarkably, there was

conversion of the FACS sorted Ly6C^{low} into Ly6C^{high} monocytes following exposure to WATA-CM

(**Figure 7B,E**). While the FACS-sorted starting cells were all Ly6C^{low} monocytes, following the 18-hour

WATA-CM treatment the monocyte population consisted of 49.7% Ly6C^{high} monocytes (only Ly6C^{low}

monocytes 50.3%) (**Figure 7E**). This was in stark contrast to all other isolated cell and CM treatment

combinations tested, which gave rise to a maximum of 16.3% of the opposite monocyte subset compared

to the starting population. These results revealed the interesting phenomenon of Ly6C^{low} into Ly6C^{high}

monocyte conversion induced by white adipocytes.

Monocyte precursor cells in the BM and their derivative macrophages in the tissues maintain their

cell numbers by cell proliferation (Cassado et al., 2015). Thus, we asked whether proliferation might

contribute to the higher proportion of terminally differentiated Ly6C^{high} or Ly6C^{low} monocyte subsets

observed following treatment with WATA-CM (in addition to the conversion of Ly6C^{low} into Ly6C^{high}) or

BATA-CM, respectively. To this end, we treated BM-generated monocytes with WATA-CM, BATA-CM

or left them untreated for 18 hours, with or without the presence of bromodeoxyuridine (BrdU). Using flow

cytometry analysis, DNA-incorporated BrdU was detected in the Ly6C^{low} or Ly6C^{high} monocyte gates of

the CD11b versus Ly6C plots, with BrdU as the third colorimetric axis. BrdU incorporation is illustrated

by a scale of yellow to red hues compared to the corresponding cells not exposed to BrdU. Proliferating

cells were quantified as percentage of CD11b^{+} Ly6C^{+} cells that were highly positive for BrdU incorporation.

Interestingly, BATA-CM increased DNA-incorporated BrdU in the sorted Ly6C^{low} monocyte subset,

present as yellow to red hues in the CD11b^{+} Ly6C^{low} cell gate (**Figure 7G**). This treatment with BATA-

CM increased the number of proliferating cells by 25-fold (**Figure 7H**). In contrast, BrdU incorporation in

the sorted Ly6C^{high} monocytes did not change with any treatment conditions. These findings revealed that

brown adipocytes promote an abundance of Ly6C^{low} monocytes by inducing their proliferation. Moreover,

this replicative behavior is consistent with recent findings that proliferative cells have higher glucose

consumption, lactate excretion and oxidative phosphorylation relative to quiescent cells (Yao et al., 2019).

Discussion

In this study, we demonstrate a temporal correlation between HFD-induced whitening of BMA and a BM myeloid lineage bias towards Ly6C^{high} monocytes *in vivo*. The latter present a more glycolytic metabolism underpinned by a reduction in mitochondrial membrane potential, a rise in mitochondrial fission genes and a reduction in mitochondrial fusion genes. Causatively, we show that CM from BMA of HFD-fed mice promotes the generation of Ly6C^{high} monocytes *ex vivo*. Phenocopying the effect of BMA derived from HFD-fed mice, *bona fide* WATA favoured a Ly6C^{high} monocyte preponderance, while BATA promoted a dominance of Ly6C^{low} monocytes. Interestingly, these differential influences by the respective adipocytes are induced by distinct cellular pathways. Mechanistically, white adipocytes induced conversion of Ly6C^{low} to Ly6C^{high} monocytes, while brown adipocytes favoured Ly6C^{low} predominance by inducing their proliferation. These changes in monocytes were underpinned by alterations in the monocytes metabolism, as WATA blunted the monocyte's mitochondrial membrane potential and OCR. Overall, our findings reveal a regulation of monocyte subsets directly influenced by adipocytes, whether from the BM or from AT. Along with the observation of dynamic whitening of BMA during HFD, our results contribute to the emerging evidence that BMA are not mere space fillers, rather they are important contributors to cellular metabolism and hematopoietic lineage decisions (Boroumand and Klip, 2020). That these changes occur early on in the course of diet-induced obesity suggests that the BM is an early responder to a change in feeding status and that adipocytes play a significant role in directing monocyte subset skewing. Moreover, they reveal novel mechanisms regulating the genesis of such skewing.

BM as an early sensor and an active contributor to HFD-induced AT macrophage accumulation

It is now widely recognized that many factors contribute to the accumulation of macrophages in AT that includes the contribution of monocytes. There is mounting evidence for both HFD-induced EWAT monocyte chemo-attraction and for enhanced BM myelopoiesis, that work together to increase the monocyte infiltration and pro-inflammatory AT macrophage accumulation (Singer et al., 2014; Trottier et al., 2012 and Fig. 1D-H). Singer *et al.* demonstrated that transplantation of BM from 18-week HFD-fed

mice into control recipients increased the number of invasive circulating monocytes and pro-inflammatory

AT macrophages (Singer et al., 2014). These findings support the notion that HFD influences hematopoietic

myeloid skewing in the BM upstream of monocyte tissue invasion, thereby contributing to downstream

systemic and peripheral tissue inflammation. Furthermore, HFD-driven chemoattracting signals from the

homeostatically-disrupted EWAT initiate tissue influx and depletion of invasive Ly6C^{high} monocytes from

the circulation, which in turn may acts as a signal for their increased production in the BM (Kamei et al.,

2006; Nagareddy et al., 2014; Nov et al., 2013; Renovato-Martins et al., 2017).

In the present study, by analyzing chronological changes in EWAT and BM myeloid composition,

we observed that total BM monocytes increase by 3 weeks of HFD, whereas the proportion of Ly6C^{high}

monocyte predominance was evident only at 8 weeks coinciding with a gain in AT macrophages that were

also pro-inflammatory skewed. We did not observe statistical significance in HFD-dependent changes in

myeloid precursors, likely due to their scarcity since HSC make up only 0.01% of the entire BM nucleated

cells (Challen et al., 2009). The rise in total monocytes observed at 3 weeks of HFD and prior to elevation

of AT macrophages identifies the BM as an earlier-responding *locus* during HFD. Interestingly, the increase

was only apparent in Ly6C^{low} patrolling monocytes and not the Ly6C^{high} tissue invasive monocyte subset

that is known for tissue infiltration in chronic obesity. This might be explained by an initial non-specific

upregulation of total myelopoiesis and an increased Ly6C^{high} monocyte BM extravasation. Therefore, it

appears that only Ly6C^{low} monocytes are elevated in the BM at 3 weeks of HFD. By 8 weeks of HFD, we

observed a shift in the monocyte subset proportion, favouring Ly6C^{high} monocytes. Thus, by 8 weeks, the

BM has adopted the 'typical' HFD-induced hematopoietic phenotype, in so far as it already shows the

increased Ly6C^{high} monocytes previously reported at the habitually studied 16-18 weeks of HFD.

Importantly, we detected a gain in pro-inflammatory AT macrophages by 8 weeks of HFD that is concurrent

with the predominance of invasive Ly6C^{high} monocytes in the BM. In contrast, the embryonic-derived tissue

resident AT macrophage markers, *Timd4* and *Marco* (Beattie et al., 2016; Gibbings et al., 2015), were not

elevated until 18 weeks of HFD. This brings to light the importance of the contribution of the BM-derived

myeloid cell pool to early EWAT pathology and AT macrophage accumulation.

Changes in BMA influence monocyte subset skewing

BMA are a distinctive member of the BM cellular suite that is sensitive to HFD effects (Pham et al., 2020). In lean mice, BMA present characteristics that are similar to those of beige or brown adipocytes in peripheral AT (Cuminetti and Arranz, 2019). With HFD, BMA undergo hyperplasia and hypertrophy triggered by PPARγ activation (Rosen and MacDougald, 2006). Furthermore, with obesity, adipocyte precursor mesenchymal stem cells tilt their lineage commitment towards adipogenesis (Silva et al., 2016). While BMA are now known to be a distinct adipocyte population, they express several characteristics of white and brown adipocytes (Boroumand and Klip, 2020; Krings et al., 2012; Sulston and Cawthorn, 2016; Tratwal et al., 2020). Brown adipocyte markers of BMA are reduced in obese mice and a parallel adoption of more white adipocyte markers has been observed (Adler et al., 2014b; Doucette et al., 2015; Krings et al., 2012). The amalgamation of upregulated white adipocyte proteins and elevated fat storage capacity is known as BM adipocyte whitening. Obesity-induced gains in whole body adiposity was causally linked to elevated circulating CD11b$^+$ monocytes (Takahashi et al., 2003). However, the influence of BMA from either lean or obese mice on the expansion of CD11b$^+$ monocytes had not been previously investigated.

The limited spatial capacity of the BM interior as well as the proximity of the cellular lineages in the various BM niches lead us to anticipate that the obesity-linked BMA whitening would influence the monocytic lineage. Here, we found a temporal, positive link between BMA hypertrophy and monocytosis at 3 weeks of HFD. By 8 weeks of HFD, BMA whitening also positively correlated with the rise in tissue invasive Ly6C^{high} monocytes in the BM. Interestingly, these findings were not associated with changes in pro-inflammatory signals in either monocyte subset. As a proof of concept for a direct and distinctive influence of adipocytes on monocyte number and subsets, we show that exposure to soluble factors from *bona fide* white adipocytes favours Ly6C^{high} monocyte preponderance while exposure to soluble factors from brown adipocytes favours Ly6C^{low} monocytes. Mechanistically, we discovered that white adipocytes induced conversion of Ly6C^{low} to Ly6C^{high} monocytes, a process that is not well understood at present. Brown adipocytes however induce their influence on Ly6C^{low} monocytes by increasing their proliferation. Interestingly, secretome studies have reported that BAT-derived VEGFA and VEGFB can stimulate

proliferation in adipocyte progenitors and endothelial cells (Bagchi et al., 2013; Leung et al., 1989).

Adipocyte-specific VEGFA transgenic mice were protected from the effects of HFD and presented altered

macrophage polarization (Elias et al., 2012). Therefore, it is conceivable that a reduction in brown adipocyte

characteristics of BMA also leads to a drop in the factors that would normally favour proliferation of

Ly6C^{low} monocytes. While the identification of the soluble unknown factor(s) responsible for the adipocyte-

specific myeloid influence is outside the focus of the present study, we have laid the foundation for a future,

comprehensive investigation of the molecular mediators involved.

How do adipocytes skew myeloid lineage subsets?

While the designation of macrophages as 'M1 and M2' (of pro- or anti-inflammatory

characteristics, respectively) represents the extremes of a wide spectrum of phenotypes, the 'M1' pro-

inflammatory orientation has been linked to glycolytic activation while the 'M2' anti-inflammatory

orientation has been linked to oxidative phosphorylation (Castoldi et al., 2016; Kelly and O'Neill, 2015).

A similar glycolytic metabolic switch with obesity has been reported for CD8 pro-inflammatory T cells

(Buck et al., 2017). Surprisingly, while pro-inflammatory pathways in monocytes are activated by

hyperglycemic conditions (Shanmugam et al., 2003), the metabolic preference of the different monocyte

subsets had not been explored. This is a compelling question, given that Ly6C^{high} monocytes penetrate

tissues and differentiate to pro-inflammatory 'M1' AT macrophages while Ly6C^{low} monocytes enter tissues

to promote healing and repair (Geissmann et al., 2003; Lin et al., 2009). Therefore, we hypothesized here

that, in the context of metabolic inflammation, there is a change in the metabolic phenotype of monocyte

subsets.

Our findings show that Ly6C^{high} and Ly6C^{low} monocytes had different metabolic preferences at

baseline, with Ly6C^{low} having higher mitochondrial membrane potential indicative of more active oxidative

phosphorylation. Interestingly, HFD consumption lowered the mitochondrial membrane potential of both

Ly6C^{low} and Ly6C^{high} monocytes, implicating a reduction in their mitochondrial oxidative capacity. This

change aligned with a HFD-induced reduction in oxygen consumption by Ly6C^{low} monocytes. Importantly,

this difference was observed as early as 3 weeks and exacerbated by 8 and 18 weeks of HFD. The elevated

lactate concentration measured in the BM at all HFD durations may be reflective of metabolic changes in

cells of the BM cavity. Since we did not observe changes in the lactate levels in WATA-CM or BATA-

CM, the higher lactate in BM of HFD-fed mice may not originate from changes in BMA.

Along with the change in monocyte subset proportions, we observed an increase in genes

responsible for mitochondrial fragmentation and a decrease in those involved in mitochondrial elongation

in monocytes from HFD-fed mice and treated with WATA-CM and BMA-CM. The physiological

consequence of these changes was revealed by the observation of a striking mitochondrial fragmentation in

monocytes treated with WATA-CM in comparison to control or BATA-CM treated monocytes. HFD is

known to induce mitochondrial fission in a number of cells that correlates with insulin resistance. A recent

report linked the two phenomena through *Drp1* action, as *Drp1* inhibition reversed HFD-induced insulin

resistance (Filippi et al., 2017). Furthermore, obese and diabetic humans have reduced *Mfn2* expression in

skeletal muscle. *Mfn2* positively correlates with insulin sensitivity (Bach et al., 2005), and *Mfn2* gene and

protein are significantly downregulated in liver cells of HFD-fed rats (Gan et al., 2013). In our analysis, the

expression of *Mfn2* negatively correlated with the abundance of Ly6C^{high} monocytes, while, the expression

of *Drp1* positively correlated with the BM abundance of Ly6C^{high} monocytes. Overall, our findings reveal

novel and distinct effects of adipocyte types on monocyte subset development that may occur through

changes in mitochondrial fitness.

Given that the proportion of Ly6C^{low} monocytes dropped in favour of a predominance of Ly6C^{high}

monocytes by 8 but not 3 weeks of HFD, it is possible that this subset change obeys a prior shift in their

metabolic orientation. Supporting these observations, exposure of monocytes to white adipocyte CM also

reduced their mitochondrial membrane potential. Furthermore, BMA-CM from 8-week HFD-fed mice also

lowered monocyte mitochondrial membrane potential compared to control BMA-CM and favoured

Ly6C^{high} predominance. Mechanistically, these data suggest that white adipocytes induce a monocyte

metabolic shift that may guide Ly6C^{high} monocyte prevalence.

In conclusion, we demonstrate an interconnectivity of cellular networks within the BM, a

vulnerability of the BM microenvironment to HFD, and a consequential myeloid lineage determination

likely due to BMA whitening. Given the vastly different metabolic and soluble factors between *bona fide*

white and brown adipocytes, it is easy to imagine that phenotypic changes in BMA can vastly alter the BM

microenvironment and the factors that surround hematopoietic cells. BMA whitening during HFD feeding

contributes to myelopoiesis and predominance of invasive Ly6C^{high} monocytes mediated by changes in

monocyte metabolism.

Materials and Methods

Animal Procedures

Mouse protocols were approved by the animal care committee (Protocol #20011850 to S.E.G. and D.J.P., University of Toronto and #1000047074 to A.K., The Hospital for Sick Children). Three groups of 9-week-old male C57BL/6 mice were fed either low fat diet (10% kcal from fat; D12450Ji, Research Diets) or HFD (60% kcal from fat; D12492i, Research Diets) for 3, 8, or 18 weeks. Mice were singly caged to avoid stress and weight inequalities induced by aggression of the dominant male to the subordinate male (Kappel et al., 2017). The cages were maintained at 22°C on a 12h light/dark cycle, the diets were replaced once per week and various cage enrichments were provided including switching the positioning of the cages throughout the study.

Body weight was measured twice per week. Upon completion of the respective feeding durations, mice were fasted for 4 hours, blood was drawn via tail vein and fasting blood glucose was measured with a glucometer. Mice were euthanized by cervical dislocation. Tissues were dissected and flash frozen in liquid nitrogen for RNA isolation. EWAT, blood and BM were immediately processed for flow cytometry analysis.

EWAT-Derived Stromal Vascular Fraction (SVF) Isolation

EWAT-derived SVF was collected as previously described (Orr et al., 2013). Excised EWAT was rinsed in PBS, minced and digested with 0.47wv/mL Liberase solution (Millipore Sigma, Cat No 5401054001) shaking at 200 rpm for 20 mins at room temperature. SVF was isolated using a 70 μm cell strainer, the enzymatic activity of Liberase was neutralized with DMEM containing HI-FBS for 5 minutes and centrifuged at 500 ×g for 10 minutes. The supernatant containing adipocytes was removed, the pellet was re-suspended in red blood cell lysis buffer for 5 minutes to remove erythrocytes and washed with PBS three times.

BM Histology

Extracted tibia and femur bones were fixed in 10% formalin for 48 hours, decalcified in 14% EDTA solution for 14 days, paraffin wax-embedded, and longitudinal sections were stained with H&E and Oil Red O (Millipore Sigma; Cat No O0625) to visualize BM adiposity.

BM-Derived Monocyte Suspension Cell Culture and Differentiation

Murine primary BM cells were plated at 500,000 cells/mL on Thermo Scientific™ Nunc™ Non-Treated 6-well plates (Cat No 1256680) with complete RMPI (company) and supplemented with recombinant mouse M-CSF (Peprotech Inc. Cat No 315-02) at 20ng/mL for 5 days (Francke et al., 2011). Following 5 days of BM cell incubation on the ultra-low attachment surface plates, the suspended cells have the signature of mature monocytes and can be used for experimentation. These BM-derived differentiated monocytes were then collected, washed with RMPI, centrifuged and re-suspended with various adipocyte CM for 18 hours.

Primary Adipocyte Isolation and Culture

Epididymal white and interscapular brown AT were dissected in sterile conditions, the tissues underwent mechanical mincing and enzymatic digestion by collagenase type I (Worthington Biochemical Corp, Cat No LS004197) at 4mg/mL shaking for 1 hour. The digested adipocytes were then filtered through a 200 μm cell strainer and centrifuged, washed twice and plated in adipocyte media: high glucose DMEM, 1% BSA, 0.5% FBS, and 1% Penicillin and Streptomycin for 24 hours. BM adipocytes were isolated from the floating fraction following BM isolation via centrifugation. The cells were counted and cultured in adipocyte media overnight. Tissues from three mice were collected and the isolated cells were pooled in each CM generating experiment.

Flow Cytometry

Following red blood cell lysis, SVF, blood and the BM were washed with FACS buffer, stained with Aqua Live/Dead Antibody (ThermoFisher, L34957) for 15 minutes in the dark, washed with FACS

buffer again, blocked with anti-mouse CD16/32 antibody (eBioscience, 14-0161-86) for 15 minutes in the

dark following a 30 minutes' incubation with corresponding antibody for each tissue **(Table 1).** The

samples were then fixed, read on LSR Fortessa cytometer (BD Biosciences) and data gated based on

appropriate fluorescence minus one (FMO) controls and analyzed on the FlowJo software version 10

(Becton, Dickinson & Company).

Quantitative PCR (qPCR)

RNA was extracted from cells or tissues using TRIzol^{TM} (Life Technologies, Cat No 15596018),

then cDNA synthesized using the SuperScript VILO cDNA kit (Life Technologies, Cat No 11754250)

according to manufacturer's instructions. qPCR reactions were performed with predesigned TaqMan probes

(Life Technologies) on a StepOne Plus Real-Time PCR System (Life Technologies). Gene expression was

normalized to that of housekeeping genes, Abt1, Hprt, and Eef2 by the $\Delta\Delta C_T$ method.

BrdU incorporation

BM derived monocytes were labelled with 10 µM BrdU (abcam, ab6326) during the 18-hour

treatment at 37C. The cells were then washed with FACS buffer, surface markers were stained as described

above and incubated in Foxp3 Fixation/Permeabilization Buffer (eBioscience, 00-5523-00) overnight. They

were then treated with 30 µg of DNAse I.9 (Sigma, 260913). for 1 hour, stained with anti-BrdU antibody

for 1 hour and then washed with FACS buffer in preparation to run on LSR Fortessa Cytometer.

Fluorescent Activated Cell Sorting of Monocytes

BM from two femurs and tibias of 10-week-old chow-fed mice was used in the isolation of

monocyte subsets via fluorescent activated cell sorting using the FACS Aria Ilu instrument. Gr1$^-$ CD115$^+$

CD11b$^+$ Ly6C$^+$ surface markers were used in identifying total monocytes and the two subsets were

subsequently distinguished by the relative surface abundance of Ly6C protein. Isolated Ly6C^{low} and

Ly6C^{high} monocytes were then validated using antibodies to CX3CR1 and CCR2 (**Table 1**), respectively.

Cell Metabolism Analysis

Oxygen consumption (OCR) and extracellular acidification rates were measured using a Seahorse XFe96 analyzer (Seahorse Bioscience). Seahorse 96-well cell culture plates were pretreated with 1 µg/cm^2 Corning™ Cell-Tak Cell and Tissue Adhesive (Fisher Scientific, Cat No CB40240) and FACS sorted Ly6C^{low} or Ly6C^{high} monocytes were seeded at 100,000 cells/well, allowed to equilibrate for 1h in XF base medium supplemented with 11 mM glucose, 1 mM sodium pyruvate and 2mM Glutamine. Following baseline measurements, FCCP was injected at 1mM and thereafter an injection of Antimycin A/ Rotenone. Cells were counted following the seahorse assay to test for changes in cellular detachment between the conditions.

Immunofluorescence Staining for Super-Resolution Microscopy

Monocytes were spun down onto coverslips and fixed with 4% paraformaldehyde for 15 minutes. Thereafter, the cells were rinsed with PBS and permeabilized with 0.1% Triton X-100 for 10 minutes. The cells were then blocked with 2% BSA in PBS at 4°C overnight. The cells were then incubated with the primary antibody, Tom20 (FL-145, Santa Cruz, sc-11415), in a 1:100 dilution in blocking buffer at 4°C overnight. Thereafter, the cells were incubated with Alexa Fluor 488 anti-rabbit secondary antibody (1:100) for 2 hours at room temperature. Cells were incubated with Alexa Fluor 568 Phalloidin (1:100) for 3 hours at room temperature. Coverslips were then mounted by ProLong Diamond (Thermo Fisher Scientific) and allowed to dry for two days. The images were acquired via Zeiss LSM880 Airyscan super resolution microscopy and the images were processed via Fiji software.

ELISA

The BM content was fractioned into hematopoietic and supernatant portions by centrifugation at 1,600 rpm. The supernatant fraction was re-suspended in 500ul of 1X PBS (supplemented with Protease Cocktail Inhibitor, Sigma). Total protein content was estimated by Pierce™ BCA Protein Assay Kit (ThermoFisher Scientific). Subsequently, mouse TNF, IL-1b, IL-6, CXCL1 and CCL2 ELISAs were

carried out according to the manufacturers' instructions. All Mouse DuoSet ELISA kits were purchased

from R&D systems. The individual protein contents were finally normalised to total protein content.

Statistical Analysis

Data are expressed as means ± SEM. An unpaired Student's T test was used to detect differences

in data sets with one variable. A one-way ANOVA with Tukey post-test was used to detect differences in

data sets of two or more variables. Statistical significance was set at $P < 0.05$. Data analysis and statistics

tests were done using GraphPad Prism version 6 (GraphPad Software), Flow Jo Software version 10.4 and

figures were prepared using Adobe Illustrator CS6 version 16.0.

Acknowledgements

A.K. was supported by Canadian Institutes of Health Research (CIHR) Foundation Grant (FND-143203)

and D.J.P. was supported by CIHR Foundation Grant (FDN-14333). P.B. was supported by a Banting &

Best Diabetes Centre-Novo Nordisk Graduate Studentship from 2017 to 2019 and by a SickKids

Restracomp Award from 2019 to 2020. We thank Dr. Clinton Robbins for support with hematopoietic

lineage analysis.

Author Contribution

P.B., S.E.G. and A.K. conceived the study design. P.B. performed experiments and analyzed the

data. M.W., J.S., I.T., N.S., L.Z., Z.L., A.L., T.M. and Y.Z. participated in experimental work and data

collection. A.K., D.J.P., and S.E.G. provided access to resources and facilities. P.J.B. contributed with

advice and trouble-shooting. P.B. and A.K. wrote the manuscript, and all authors read and approved it.

References:

Adler, B.J., Kaushansky, K., and Rubin, C.T. (2014). Obesity-driven disruption of haematopoiesis and the bone marrow niche. Nat. Rev. Endocrinol. *10*, 737–748.

Ambrosi, T.H., Scialdone, A., Graja, A., Gohlke, S., Jank, A.M., Bocian, C., Woelk, L., Fan, H., Logan, D.W., Schürmann, A., et al. (2017). Adipocyte Accumulation in the Bone Marrow during Obesity and Aging Impairs Stem Cell-Based Hematopoietic and Bone Regeneration. Cell Stem Cell *20*, 771–784.e6.

Auffray, C., Fogg, D., Garfa, M., Gaelle, E., Join-Lambert, O., Kayal, S., Sarnacki, S., Ana, C., Lauvau, G., and Geissmann, F. (2007). Monitoring of Blood Vessels and Tissues by a Population of Monocytes with Patrolling Behavior. *317*, 666–671.

Auffray, C., Fogg, D.K., Narni-Mancinelli, E., Senechal, B., Trouillet, C., Saederup, N., Leemput, J., Bigot, K., Campisi, L., Abitbol, M., et al. (2009). CX 3 CR1 + CD115 + CD135 + common macrophage/DC precursors and the role of CX 3 CR1 in their response to inflammation . J. Exp. Med. *206*, 595–606.

Bach, D., Naon, D., Pich, S., Soriano, F.X., Vega, N., Rieusset, J., Laville, M., Guillet, C., Boirie, Y., Wallberg-henriksson, H., et al. (2005). Expression of Mfn2, the Charcot-Marie-Tooth Neuropathy Type 2A Gene, in Human Skeletal Muscle. Diabetes *54*, 2685–2693.

Bagchi, M., Kim, L.A., Boucher, J., Walshe, T.E., Kahn, C.R., and Amore, P.A.D. (2013). Vascular endothelial growth factor is important for brown adipose tissue development and maintenance. FASEB J. *27*, 3257–3271.

Beattie, L., Sawtell, A., Mann, J., Frame, T.C.M., Teal, B., de Labastida Rivera, F., Brown, N., Walwyn-Brown, K., Moore, J.W.J., MacDonald, S., et al. (2016). Bone marrow-derived and resident liver macrophages display unique transcriptomic signatures but similar biological functions. J. Hepatol. *65*, 758–768.

Boniakowski, A.E., Kimball, A.S., Joshi, A., Schaller, M., Davis, F.M., denDekker, A., Obi, A.T., Moore, B.B., Kunkel, S.L., and Gallagher, K.A. (2018). Murine macrophage chemokine receptor CCR2 plays a crucial role in macrophage recruitment and regulated inflammation in wound healing. Eur. J. Immunol. *48*, 1445–1455.

Boroumand, P., and Klip, A. (2020). Bone marrow adipose cells – cellular interactions and changes with obesity. J. Cell Sci. *1*, 1–8.

Buck, M.D., Sowell, R.T., Kaech, S.M., and Pearce, E.L. (2017). Metabolic Instruction of Immunity. Cell *169*, 570–586.

Cassado, A.A., D'Império Lima, M.R., and Bortoluci, K.R. (2015). Revisiting mouse peritoneal macrophages: Heterogeneity, development, and function. Front. Immunol. *6*, 1–9.

Castoldi, A., De Souza, C.N., Saraiva Câmara, N.O., and Moraes-Vieira, P.M. (2016). The macrophage switch in obesity development. Front. Immunol. *6*, 1–11.

Centers for Disease Control and Prevention, C. (2017). National Diabetes Statistics Report: Estimates of Diabetes and Its Burden in the United States. Atlanta, GA: Centers for Disease Control and Prevention; 2017. US Dep. Heal. Hum. Serv.

Challen, G.A., Boles, N., Lin, K.K.Y., and Goodell, M.A. (2009). Mouse hematopoietic stem cell identification and analysis. Cytom. Part A *75*, 14–24.

Cuminetti, V., and Arranz, L. (2019). Bone Marrow Adipocytes: The Enigmatic Components of the Hematopoietic Stem Cell Niche. J. Clin. Med. *8*, 707.

Doucette, C.R., Horowitz, M.C., Berry, R., MacDougald, O.A., Anunciado-Koza, R., Koza, R.A., and Rosen, C.J. (2015a). A High Fat Diet Increases Bone Marrow Adipose Tissue (MAT) But Does Not Alter Trabecular or Cortical Bone Mass in C57BL/6J Mice. J Cell Physiol *230*, 2032–2037.

Doucette, C.R., Horowitz, M.C., Berry, R., Macdougald, O.A., Anunciado-Koza, R., Koza, R.A., and Rosen, C.J. (2015b). A High Fat Diet Increases Bone Marrow Adipose Tissue (MAT) But Does Not Alter Trabecular or Cortical Bone Mass in C57BL/6J Mice. J. Cell. Physiol. *230*, 2032–2037.

Elias, I., Franckhauser, S., Ferré, T., Vilà, L., Tafuro, S., Muñoz, S., Roca, C., Ramos, D., Pujol, A., Riu, E., et al. (2012). Adipose Tissue Overexpression of Vascular Endothelial Growth Factor Protects Against Diet-Induced Obesity and Insulin Resistance. Diabetes *61*, 1801–1813.

Filippi, B.M., Abraham, M.A., Silva, P.N., Bauer, P. V, Rocheleau, J. V, Lam, T.K.T., Filippi, B.M., Abraham, M.A., Silva, P.N., Rasti, M., et al. (2017). Mitochondrial Fission Changes in the Dorsal Vagal Report Dynamin-Related Protein 1-Dependent Mitochondrial Fission Changes in the Dorsal Vagal Complex Regulate Insulin Action. CellReports *18*, 2301–2309.

Fink, L.N., Costford, S.R., Lee, Y.S., Jensen, T.E., Bilan, P.J., Oberbach, A., Blüher, M., Olefsky, J.M., Sams, A., and Klip, A. (2014a). Pro-Inflammatory macrophages increase in skeletal muscle of high fat-Fed mice and correlate with metabolic risk markers in humans. Obesity *22*, 747–757.

Fink, L.N., Costford, S.R., Lee, Y.S., Jensen, T.E., Bilan, P.J., Oberbach, A., Blüher, M., Olefsky, J.M., Sams, A., and Klip, A. (2014b). Pro-Inflammatory macrophages increase in skeletal muscle of high fat-Fed mice and correlate with metabolic risk markers in humans. Obesity *22*, 747–757.

Francke, A., Herold, J., Weinert, S., Strasser, R.H., and Braun-Dullaeus, R.C. (2011). Generation of mature murine monocytes from heterogeneous bone marrow and description of their properties. J. Histochem. Cytochem. *59*, 813–825.

Friedrich, K., Sommer, M., Strobel, S., Thrum, S., and Blüher, M. (2019). Perturbation of the Monocyte Compartment in Human Obesity. Front. Immunol. *10*, 1–10.

Gan, K., Wang, C., Chen, J., Zhu, C., and Song, G. (2013). Mitofusin-2 ameliorates high-fat diet-induced insulin resistance in liver of rats. *19*, 1572–1581.

Garvey, W.T., Henry, R.R., Kern, P.A., Saghizadeh, M., and Ong, J.M. (2008). The expression of TNF alpha by human muscle. Relationship to insulin resistance. J. Clin. Invest. *97*, 1111–1116.

Geissmann, F., Jung, S., and Littman, D.R. (2003). Blood monocytes consist of two principal subsets with distinct migratory properties. Immunity *19*, 71–82.

Gibbings, S.L., Goyal, R., Desch, A.N., Leach, S.M., Prabagar, M., Atif, S.M., Bratton, D.L., Janssen, W., and Jakubzick, C. V (2015). Transcriptome analysis highlights the conserved difference between embryonic and postnatal-derived alveolar macrophages. Blood *126*, 1357–1366.

Gottlieb, R.A., and Bernstein, D. (2017). Mitochondrial Remodeling: Rearranging, Recycling, and Reprogramming. Cell Calcium *60*, 88–101.

Griffin, C., Eter, L., Lanzetta, N., Abrishami, S., Varghese, M., McKernan, K., Muir, L., Lane, J., Lumeng, C.N.-K., and Singer, K. (2018). TLR4, TRIF, and MyD88 are essential for myelopoiesis and CD11c+ adipose tissue macrophage production in obese mice. J. Biol. Chem. jbc.RA117.001526.

Hamann, A. (2017). Obesity Update 2017. Diabetologe.

Kamei, N., Tobe, K., Suzuki, R., Ohsugi, M., Watanabe, T., Kubota, N., Ohtsuka-kowatari, N., Kumagai, K., Sakamoto, K., Kobayashi, M., et al. (2006). Overexpression of Monocyte Chemoattractant Protein-1 in Adipose Tissues Causes Macrophage Recruitment and Insulin Resistance *. J. Biol. Chem. *281*, 26602–26614.

Kappel, S., Hawkins, P., and Mendl, M. (2017). To Group or Not to Grou? Good Practice for
Housing Male Laboratory Mice. Animals 1–25.

Kelly, B., and O'Neill, L.A. (2015). Metabolic reprogramming in macrophages and dendritic
cells in innate immunity. Cell Res. *25*, 771–784.

Kohyama, M., Matsuoka, S., Shida, K., Sugihara, F., Aoshi, T., Kishida, K., Ishii, K.J., and
Arase, H. (2016). Monocyte infiltration into obese and fibrilized tissues is regulated by PILRα.
Eur. J. Immunol. *46*, 1214–1223.

Kratofil, R.M., Kubes, P., and Deniset, J.F. (2017). Monocyte conversion during inflammation
and injury. Arterioscler. Thromb. Vasc. Biol. *37*, 35–42.

Krings, A., Rahman, S., Huang, S., Lu, Y., Czernik, P.J., and Lecka-Czernik, B. (2012). Bone
marrow fat has brown adipose tissue characteristics, which are attenuated with aging and
diabetes. Bone *50*, 546–552.

Lanthier, N., Molendi-Coste, O., Horsmans, Y., van Rooijen, N., Cani, P.D., and Leclercq, I.A.
(2010). Kupffer cell activation is a causal factor for hepatic insulin resistance. Am. J. Physiol.
Liver Physiol. *298*, G107–G116.

Lee, M.K.S., and Murphy, A.J. (2019). Glycolysis Is Required for LPS-Induced Activation and
Monocytes. *10*, 1–10.

Lessard, A.J., LeBel, M., Egarnes, B., Préfontaine, P., Thériault, P., Droit, A., Brunet, A., Rivest,
S., and Gosselin, J. (2017). Triggering of NOD2 Receptor Converts Inflammatory Ly6Chighinto
Ly6ClowMonocytes with Patrolling Properties. Cell Rep. *20*, 1830–1843.

Leung, D.W., Cachianes, G., Kuang, W., Goeddel, D. V, and Ferrara, N. (1989). Vascular
Endothelial Growl th Factor Is a Secreted Angiogenic Mitogen. Science (80-.). *246*, 1306–9.

Lin, S.L., Castano, A.P., Nowlin, B.T., Lupher, M.L., and Duffield, J.S. (2009). Bone Marrow
Ly6Chigh Monocytes Are Selectively Recruited to Injured Kidney and Differentiate into
Functionally Distinct Populations. J. Immunol. *183*, 6733–6743.

Liu, T., Sun, Y.-C., Cheng, P., and Shao, H.-G. (2019). Adipose tissue macrophage-derived
exosomal miR-29a regulates obesity-associated insulin resistance. Biochem. Biophys. Res.
Commun.

Lumeng, C.N., Deyoung, S.M., Bodzin, J.L., and Saltiel, A.R. (2007). Increased Inflammatory
Properties of Adipose Tissue Macrophages Recruited During Diet-Induced Obesity. Diabetes *56*,
16–23.

Lumeng, C.N., Delproposto, J.B., Westcott, D.J., and Saltiel, A.R. (2008). Phenotypic Switching
of Adipose Tissue Macrophages With Obesity Is Generated by Spatiotemporal Differences in
Macrophage Subtypes. Diabetes *57*, 3239–3246.

Manicone, A.M., Gong, K., Johnston, L.K., and Giannandrea, M. (2016). Diet-induced obesity
alters myeloid cell populations in naïve and injured lung. Respir. Res. *17*, 24.

Nagareddy, P.R., Kraakman, M., Masters, S.L., Stirzaker, R.A., Gorman, D.J., Grant, R.W.,
Dragoljevic, D., Hong, E.S., Abdel-Latif, A., Smyth, S.S., et al. (2014). Adipose tissue
macrophages promote myelopoiesis and monocytosis in obesity. Cell Metab. *19*, 821–835.

Nakagami, H. (2013). The Mechanism of White and Brown Adipocyte Differentiation. Diabetes
Metab. J. *37*, 85–90.

Nascimento, E.B.M., Boon, M.R., and van Marken Lichtenbelt, W.D. (2014). Fat cells gain new
identities. Sci. Transl. Med. *6*, 1–4.

Nov, O., Shapiro, H., Ovadia, H., Tarnovscki, T., Dvir, I., Shemesh, E., Kovsan, J., Shelef, I.,
Carmi, Y., Voronov, E., et al. (2013). Interleukin-1β Regulates Fat-Liver Crosstalk in Obesity by
Auto-Paracrine Modulation of Adipose Tissue Inflammation and Expandability. PLoS One *8*.

O'Neill, L., Kishton, R., and Rathmell, J. (2016). A guide to immunometabolism for
immunologists. Nat. Rev. Immunol. *16*, 553–565.
Orr, J.S., Kennedy, A.J., and Hasty, A.H. (2013). Isolation of Adipose Tissue Immune Cells. J.
Vis. Exp. 1–7.
Patel, A.A., Zhang, Y., Fullerton, J.N., Boelen, L., Rongvaux, A., Maini, A.A., Bigley, V.,
Flavell, R.A., Gilroy, D.W., Asquith, B., et al. (2017). The fate and lifespan of human monocyte
subsets in steady state and systemic inflammation. J. Exp. Med. *214*, 1913–1923.
Pence, B.D., and Yarbro, J.R. (2019). Classical monocytes maintain ex vivo glycolytic
metabolism and early but not later inflammatory responses in older adults. Immun. Ageing *16*,
1–8.
Pham, T.T., Ivaska, K.K., Hannukainen, J.C., Virtanen, K.A., Lidell, M.E., Parkkola, R., Piirola,
S., Oikonen, V., Nuutila, P., Biology, C., et al. (2020). Human bone marrow adipose tissue is a
metabolically active and insulin sensitive distinct fat depot. J. Clin. Endocrinol. Metab. *pii*,
dgaa216.
Renovato-Martins, M., Matheus, M.E., de Andrade, I.R., Moraes, J.A., da Silva, S.V., Citelli dos
Reis, M., de Souza, A.A.P., da Silva, C.C., Bouskela, E., and Barja-Fidalgo, C. (2017).
Microparticles derived from obese adipose tissue elicit a pro-inflammatory phenotype of CD16 +
, CCR5 + and TLR8 + monocytes. Biochim. Biophys. Acta - Mol. Basis Dis. *1863*, 139–151.
Rosen, E.D., and MacDougald, O.A. (2006). Adipocyte differentiation from the inside out. Nat.
Rev. Mol. Cell Biol. *7*, 885–896.
Scheller, E.L., Doucette, C.R., Learman, B.S., Cawthorn, W.P., Khandaker, S., Schell, B., Wu,
B., Ding, S.Y., Bredella, M.A., Fazeli, P.K., et al. (2015). Region-specific variation in the
properties of skeletal adipocytes reveals regulated and constitutive marrow adipose tissues. Nat.
Commun. *6*, 1–13.
Scheller, E.L., Khoury, B., Moller, K.L., Wee, N.K.Y., Khandaker, S., Kozloff, K.M.,
Abrishami, S.H., Zamarron, B.F., and Singer, K. (2016). Changes in skeletal integrity and
marrow adiposity during high-fat diet and after weight loss. Front. Endocrinol. (Lausanne). *7*, 1–
13.
Shanmugam, N., Reddy, M.A., Guha, M., and Natarajan, R. (2003). High glucose-induced
expression of proinflammatory cytokine and chemokine genes in monocytic cells. Diabetes *52*,
1256–1264.
Sharma, M., Boytard, L., Hadi, T., Koelwyn, G., Simon, R., Ouimet, M., Seifert, L., Spiro, W.,
Yan, B., Hutchison, S., et al. (2020). Enhanced glycolysis and HIF-1 α activation in adipose
tissue macrophages sustains local and systemic interleukin-1 β production in obesity. Sci. Rep.
10, 1–12.
Silva, S. V, Renovato-martins, M., Ribeiro-pereira, C., Citelli, M., and Barja-fidalgo, C. (2016).
Obesity Modifies Bone Marrow Microenvironment and Directs Bone Marrow Mesenchymal
Cells to Adipogenesis. Obes. Biol. Integr. Physiol. *24*, 2522–2532.
Singer, K., DelProposto, J., Lee Morris, D., Zamarron, B., Mergian, T., Maley, N., Cho, K.W.,
Geletka, L., Subbaiah, P., Muir, L., et al. (2014). Diet-induced obesity promotes myelopoiesis in
hematopoietic stem cells. Mol. Metab. *3*, 664–675.
De Souza, C.T., Araujo, E.P., Bordin, S., Ashimine, R., Zollner, R.L., Boschero, A.C., Saad,
M.J.A., and Velloso, L.A. (2005). Consumption of a fat-rich diet activates a proinflammatory
response and induces insulin resistance in the hypothalamus. Endocrinology *146*, 4192–4199.
Sulston, R.J., and Cawthorn, W.P. (2016). Bone marrow adipose tissue as an endocrine organ:
close to the bone? Horm. Mol. Biol. Clin. Investig. *28*, 21–38.

Takahashi, K., Mizuarai, S., Araki, H., Mashiko, S., Ishihara, A., Kanatani, A., Itadani, H., and
Kotani, H. (2003). Adiposity Elevates Plasma MCP-1 Levels Leading to the Increased CD11b-
positive Monocytes in Mice *. J. Biol. Chem. *278*, 46654–46660.
Thapa, B., and Lee, K. (2019). Metabolic influence on macrophage polarization and
pathogenesis. BMB Rep. *52*, 360–372.
Tratwal, J., Labella, R., Bravenboer, N., and Kerckhofs, G. (2020). Reporting Guidelines ,
Review of Methodological Standards , and Challenges Toward Harmonization in Bone Marrow
Adiposity Research . Report of the Methodologies Working Group of the International Bone
Marrow Adiposity Society. Front. Endocrinol. (Lausanne). *11*, 1–36.
Ussar, S., Lee, K., Dankel, S., Boucher, J., Haering, M., Kleinridders, A., Thomou, T., Xue, R.,
Macoteia, Y., Cypess, A., et al. (2014). Asc-1, PAT2 and P2RX5 are novel cell surface markers
for white, beige and brown adipocytes. Sci Transl Med *247*, 247ra103.
doi:10.1126/scitranslmed.3008490. Asc-1,.
Verdeguer, F., and Aouadi, M. (2017). Macrophage heterogeneity and energy metabolism. Exp.
Cell Res. *360*, 35–40.
Weisberg, S.P., McCann, D., Desai, M., Rosenbaum, M., Leibel, R.L., and Ferrante, A.W.
(2003). Obesity is associated with macrophage accumulation in adipose tissue. J. Clin. Invest.
112, 1796–1808.
Weisberg, S.P., Hunter, D., Huber, R., Lemieux, J., Slaymaker, S., Vaddi, K., Charo, I., Leibel,
R.L., and Ferrante Jr, A.W. (2006). CCR2 modulates inflammatory and metabolic effects of
high-fat feeding. J. Clin. Invest. *116*, 115–124.
WHO (2016). World Health Organization. Global Report on Diabetes. Geneva, Switzerland.
802

Figure 1. Mice fed HFD increased body weight, fasting blood glucose, visceral and bone marrow adipose tissue. [A] Body weight was measured twice a week throughout the course of feeding. [B] Mice were fasted for 4 hours after which fasting blood glucose was measured. [C] Epididymis white adipose tissue was weighed the day of sacrifice. [D] Area covered by bone marrow adipocytes were measured

following hematoxylin and eosin (H&E) to visualize adipocyte ghosts using quantified using ImageJ. [E]

Bone marrow adipocyte numbers were blindly counted following H&E staining. [F] Femurs of control

and HFD-fed mice were fixed in formalin for 2 days followed by 14 days EDTA decalcification and

subsequently stained with ORO and [G] H&E. Scale bar is 25μm. Results are expressed as mean ± SEM,

n=4-14 mice per group. One-way ANOVA was used for statistical analysis at each dietary group and

across feeding periods. *$P < 0.05$, **$P < 0.01$, ***$P < 0.001$, ****$P < 0.0001$ and ns = not significant.

Figure 2

Figure 2. Bone marrow monocytes increased prior to adipose tissue macrophages. Quantification of

stromal vascular fraction (SVF) of [A] adipose tissue monocytes [B] total macrophages and [C] pro-

inflammatory macrophages via flow cytometry analysis identified first by size and granularity through

forward and side scatter and then using the following markers: CD11b CD115 F4/80 Ly6C GR1 CD11c.

[D-F] Whole adipose tissue gene expression of the myeloid marker *Itgam,* macrophage marker *Adgre* and

pro-inflammatory myeloid marker *Itgax* by RTqPCR. [G-H] Gene expression analysis of embryonic

derived tissue resident macrophage markers *Timd4* and *Marco* by RTqPCR. [I] Flow cytometry analysis of

total and [J] monocyte subsets in the bone marrow with HFD-feeding. Results are expressed as mean ±

SEM, n=5-10 mice per group. One-way ANOVA was used for statistical analysis at each dietary group and

across feeding periods. *P < 0.05, **P < 0.01, ***P < 0.001, ****P < 0.0001 and ns = not significant.

Figure 3

Figure 3. HFD blunted the mitochondrial oxidation of monocyte subsets. Flow cytometry detection of

[A-C] MitoTracker Red CMXRos in bone marrow monocyte subsets of control and HFD-fed mice. [D-F]

OCR of FACS-sorted Ly6C^{low} and Ly6C^{high} monocytes from control and HFD-fed mice measured by the

Seahorse Assay. [G-I] Area under the curve of OCR of FACS-sorted Ly6C^{low} and Ly6C^{high} monocytes from

control and HFD-fed mice measured by the Seahorse Assay. [J] Area under the curve of ECAR

measurement from FACS-sorted Ly6C^{low} and Ly6C^{high} monocytes of control and HFD-fed mice measured

by the Seahorse Assay and pooled from all feeding durations. [K] BM lactate measurement of the

supernatant fraction in HFD and control samples. Results are expressed as mean ± SEM, n=3 mice per

group where applicable. One-way ANOVA was used for statistical analysis. *P < 0.05 and **P < 0.01.

Graph in panels C was created using FlowJo software (Becton, Dickinson Company).

Figure 4

Figure 4. **HFD downregulated gene expression of mitochondrial fusion genes and upregulated mitochondrial fission genes.** Gene expression of mitochondrial fusion genes *Mfn1, Mfn2, Opa1, Tomm20,* and *Tomm40* in monocytes isolated from control and HFD-fed mice for [A] 3, [B] 8, [C] 18 weeks. Gene expression of mitochondrial fission genes *Drp1, CypD* and *Fis1* in monocytes isolated from control and HFD-fed mice for [D] 3, [E] 8, [F] 18 weeks. [G] Correlative analysis of monocyte subset BM cellular

842   abundance and BM-derived monocyte *Mfn2* gene expression. [H] Correlative analysis of monocyte subset

843   BM cellular abundance and BM-derived monocyte *Drp1* gene expression. Results are expressed as mean ±

844   SEM, n=3-9 mice per group. One-way ANOVA was used for statistical analysis. *P < 0.05.

Figure 5

Figure 5. Bone marrow adipocyte whitening detected at 8 and 18 weeks of HFD favours increased

bone marrow Ly6C^{high} monocytes and induced changes in mitochondrial dynamic gene expression.

Volcano plots showing the gene expression of white and brown adipocyte markers in red and blue

respectively [A] at 3, [B] 8 and [C] 18 weeks of HFD. Correlative graphs of the [D] white adipocyte marker

ASC1 and [E] brown adipocyte marker UCP1 with Ly6C^{low} and Ly6C^{high} monocyte subsets. Flow

cytometry quantification of naïve Ly6C^{low} and Ly6C^{high} monocytes upon 18 hours of treatment with BMA-

CM from control and HFD-fed mice for [F] 3, [G] 8 and [H]18 weeks. [I] Gene expression of mitochondrial

fusion genes *Mfn1, Mfn2, Opa1, Tomm20,* and *Tomm40* in naïve monocytes treated with BMA-CM from

control and HFD-fed mice. [J] Gene expression of mitochondrial fission genes *Drp1, CypD* and *Fis1* in

naïve monocytes treated with BMA-CM from control and HFD-fed mice. Results are expressed as mean ±

SEM, n=3-9 mice per group. One-way ANOVA was used for statistical analysis at each dietary group and

across feeding periods. *P < 0.05, ****P < 0.0001 and ns = not significant.

Figure 6

Figure 6. *Bona fide* white adipocytes favoured Ly6C^high monocytes preponderance, blunted

monocyte mitochondrial membrane potential, and induced mitochondrial fragmentation *in vitro*.

Naïve monocytes were treated with CM from isolated *bone fide* white and brown adipocytes from EWAT

and inter-scapular BAT for 18 hours. [A] Flow cytometry dot plots detecting the effects of 18 hours of

treatment with media alone, WATA-CM and BATA-CM on monocyte subset abundance with the [B]

quantification of the gates drawn for Ly6C^{low} and Ly6C^{high} monocytes. [C] MitoTracker Red CMXRos

detected by flow cytometry in Ly6C^{low} monocytes treated with media alone, WATA-CM and BATA-CM

for 18 hours. [D] Gene expression of mitochondrial fusion genes *Mfn1*, *Mfn2*, *Opa1*, *Tomm20*, and

Tomm40 in naïve monocytes treated with Media, WATA-CM and BATA-CM. [E] Gene expression of

mitochondrial fission genes *Drp1*, *CypD* and *Fis1* in naïve monocytes treated with Media, WATA-CM

and BATA-CM. [F] Super-resolution microscopy analysis of mitochondrial morphology via TOM20

staining in monocytes treated with Media, WATA-CM and BATA-CM. Scale bar = $5\mu m$. Images

analyzed in Fiji (version 2.1.0/1.53c). WATA-CM = white adipose tissue adipocyte-conditioned media;

BATA-CM = brown adipocyte tissue adipocyte-conditioned media. EWAT = epididymal white adipose

tissue; BAT = brown adipose tissue. Results are expressed as mean ± SEM, n=3-9 mice per group. One-

way ANOVA was used for statistical analysis. *$P < 0.05$, **$P < 0.01$ and ***$P < 0.001$, ****$P < 0.0001$

and ns = not significant. Graphs in panels A-C were created using FlowJo software (Becton, Dickinson

Company).

Figure 7

A Media

B WATA-CM

C BATA-CM

D Media

E WATA-CM

F BATA-CM

G Media   WATA-CM   BATA-CM

H

Figure 7. White and brown adipocytes favour Ly6C^{high} and Ly6C^{low} monocytes through different

mechanisms. FACS-sorted Ly6C^{low} and Ly6C^{high} monocytes from control mice were independently treated

with [A] media alone, [B] WATA-CM and [C] BATA-CM for 18 hours after which the presence of the

alternate monocyte subset was assessed via flow cytometry. [D-F] Quantification of Ly6C^{low} and Ly6C^{high}

gates in panels A-C. [G] *In vitro* generated monocytes were treated with media alone, WATA-CM or

BATA-CM in the presence of the pyrimidine analogue BrdU to detect cellular proliferation. [H]

Quantification of BrdU abundance in Ly6C^{low} and Ly6C^{high} with treatments in panel C. Results are

expressed as mean ± SEM, n=3-6 mice per group. One way ANOVA test was used for statistical analysis.

****$P < 0.0005$. Graphs in panels A-C and G were created using FlowJo software (Becton, Dickinson

Company).

Figure S1

Figure S1. Monocyte precursor and stem cells were not significantly altered with HFD *in vivo*. Flow

cytometry analysis of HSC and monocyte precursor cells in mice fed a Control or HFD for [A] 3, [B] 8,

and [C] 18 weeks. Flow cytometry antibodies used are listed in Table 1. Student's T test was used and no

statistical significance was observed.

Figure S2

Figure S2: BM pro-inflammatory factors did not change with HFD. No changes in pro-inflammatory cytokines were detected in BM supernatant fraction using the ELISA assay in mice fed control or HFD for [A] 3, [B] 8, or [C] 18 weeks. FACS-sorted monocyte subsets were analyzed for changes in pro-inflammatory gene expression; shown in [D] TNFα gene expression increased with 18 weeks of HFD in Ly6Chigh monocytes but no other changes were detected. Results are expressed as mean ± SEM, n=3-6 mice per group. One way ANOVA test was used for statistical analysis. *P < 0.05 and **P < 0.01.

Figure S3

902 **Figure S3. BMA precursor cells with HFD.** RT qPCR of isolated pelleted fraction were assessed for pre-

903 adipocyte markers of mice fed a Control or HFD for [A] 3, [B] 8, and [C] 18 weeks. [D] classical white and

904 brown adipocyte markers were confirmed in isolated WATA and BATA.

Figure S4

A

B

C

D

Figure S4: Flow cytometry gating strategy for identification of Ly6C^{low} and Ly6C^{high} monocytes and their precursor cells. [A] Using side and forward scatter to identify singlet cells, CD115+Gr1- cells were

identified that were further analyzed using CD11b and Ly6C. [B] Ly6C^{low} and Ly6C^{high} monocyte gates

were drawn using fluorescence minus one (FMO) controls for CD11b and Ly6C. [C] The subsets were

further confirmed via CX3CR1 and CCR2, respectively. [D] CD45+ cells were analyzed using a lineage

cocktail (table 1) and CD117+, further broken down into Sca+ and – cells. Sca+ cells were further classified

using CD48 and CD150 while Sca- cells were further characterized via CD16/32 and CD34. Graph in panels

C was created using FlowJo software (Becton, Dickinson Company).

Figure S5

Figure S5: HSC and their progenitors using the panel of markers listed in table 1 following the white

and brown CM treatment for 18 hours. [A-C] Flow cytometry analysis was used to test for the presence

and changes in HSC and monocyte precursor cells in cultured monocytes treated with media, WATA-CM,

and BATA-CM. Flow cytometry antibodies used are listed in Table 1. One way ANOVA test was used for

statistical analysis. $*P < 0.05$.

Table 1. List of Antibodies used for flow cytometry analysis.

Markers	Fluorophore	Manufacturer	Clone	Catalog Number
CD11b	APC-eFluor 780	eBioscience	M1/70	47-0112-82
Ly6C	PE-Cy7	Biolegend	HK1.4	128018
Gr-1	Alexa Fluor 700	eBioscience	RB6-8C5	56-5931-82
CX3CR1	Brilliant Violet 421	Biolegend	SA011F11	149023
CCR2	Alexa594	R&D Systems	475301	FAB5538T
CD115	APC	eBioscience	AFS98	17-1152-82
F4/80	PerCP-Cy5.5	BioLegend	BM8	123128
CD11c	FITC	eBioscience	N418	11-0114-81
CD34	FITC	BD Biosciences	RA1434	553733
CD16/32	APC-R700	BD Biosciences	2.4G2	565502
CD45.2	BV650	BD Biosciences	30-F11	563410
CD150	PEPCP710	invitrogen	mShad150	46-1502-82
CD117	APC/Fire 750	BioLegend	2B8	105838
SCA1	PE-Cy7	invitrogen	D7	25-5981-82
CD48	BV605	BioLegend	MM48-1	103441
CD3e*	PE	BD Biosciences	17A2	555725
B220*	PE	BD Biosciences	RA3-6B2	553089
Ly6G*	PE	BD Biosciences	1A8	551461
CD11b*	PE	BD Biosciences	M1/70	553311
CD11c*	PE	BioLegend	N418	117308
Ter119*	PE	BD Biosciences	TER-119	553623
CD127*	PE	eBioscience	A7R34	12-1271-82
NK1.1*	PE	BD Biosciences	PK138	553165

922 *LINEAGE is a cocktail of antibodies containing CD3e, B220, Ly6G, CD11b, CD11c, Ter119, CD127,
923 NK1.1 with the same fluorophore to exclude all terminally differentiated cells.