Applications of the indole-alkaloid gramine shape the prokaryotic microbiota thriving at the barley root-soil interface

Mauro Maver[1,2,3], Carmen Escudero-Martinez[2], James Abbott[4], Jenny Morris[5], Pete E. Hedley[5],

Tanja Mimmo[1,3*] and Davide Bulgarelli[2*]

[1] Faculty of Science and Technology, Free University of Bozen-Bolzano, Bolzano, Italy
[2] Plant Sciences, School of Life Sciences, University of Dundee, Dundee, United Kingdom
[3] Competence Centre for Plant Health, Free University of Bozen-Bolzano, Bolzano, Italy
[4] Data Analysis Group, School of Life Sciences, University of Dundee, Dundee, United Kingdom
[5] Cell and Molecular Sciences, The James Hutton Institute, Dundee, United Kingdom

*Corresponding Authors:

Tanja Mimmo
Faculty of Science and Technology, Free University of Bozen-Bolzano, Bolzano, Italy
Email address: tmimmo@unibz.it

Davide Bulgarelli
Plant Sciences, School of Life Sciences, University of Dundee, Dundee, United Kingdom
Email address: d.bulgarelli@dundee.ac.uk

Abstract

Microbial communities proliferating at the root-soil interface, collectively referred to as the rhizosphere microbiota, represent an untapped beneficial resource for plant growth, development and health. Integral to a rational manipulation of the microbiota for sustainable agriculture is the identification of the molecular determinants of these communities. In plants, biosynthesis of allelochemicals is centre stage in defining inter-organismal relationships in the environment. Intriguingly, this process has been moulded by domestication and breeding selection. The indole-alkaloid gramine, whose occurrence in barley (*Hordeum vulgare* L.) is widespread among wild genotypes but has been counter selected in several modern varieties, is a paradigmatic example of this phenomenon. This prompted us to investigate how exogenous applications of gramine impacted on the rhizosphere microbiota of two, gramine-free, elite barley varieties grown in a reference agricultural soil. High throughput 16S rRNA gene amplicon sequencing revealed that applications of gramine interfere with the proliferation of a subset of soil microbes with a relatively broad phylogenetic assignment. Strikingly, growth of these bacteria appeared to be rescued by barley plants in a genotype- and dosage-independent manner. In parallel, we discovered that host recruitment cues can interfere with the impact of gramine application in a host genotype-dependent manner. Interestingly, this latter effect displayed a bias for members of the phyla Proteobacteria. These initial observations indicate that gramine can act as a determinant of the prokaryotic communities inhabiting the root-soil interface.

Introduction

The interface between roots and soil host distinct microbial communities, collectively referred to as rhizosphere microbiota (Turner, James & Poole, 2013). These plant-microbial assemblages define a continuum of parasitic, commensal, and mutualistic interactions (Schlaeppi & Bulgarelli, 2015). For example, so-called plant growth-promoting rhizobacteria can enhance plant mineral uptake and protect their hosts from pathogens (Lugtenberg & Kamilova, 2009). Studies conducted with multiple plant species supports the notion that members of the rhizosphere microbiota are not passively assembled from the surrounding soil biota, rather a multi-step selection process differentiating plant-associated communities from those identified in unplanted soils (Bulgarelli et al., 2013; Edwards et al., 2015). This selection process operates at multiple taxonomic and functional ranks, with the enrichment of members of the phyla Actinobacteria, Bacteroidetes, Proteobacteria and Firmicutes representing the distinct signature of this selection process at the root-soil interface (Bulgarelli et al., 2013; Alegria Terrazas et al., 2016). The host genome represents one of the "checkpoints" in the selection process defining the microbiota thriving at the root-soil interface (Schlaeppi & Bulgarelli, 2015; Hacquard et al., 2015; Alegria Terrazas et al., 2016). Integral to host-mediated molecular mechanisms shaping the rhizosphere microbiota is the root exudation of metabolites capable of modulating the interactions among plants, microbes and the surrounding environment (Dakora & Phillips, 2002; Jones, Nguyen & Finlay, 2009; Pascale et al., 2020). Consistently, an increasing number of plant primary (Canarini et al., 2019), as well as secondary (Rolfe, Griffiths & Ton, 2019), metabolites have recently been implicated in shaping the plant microbiota.

Interestingly, the processes of crop domestication and breeding selection, which gave rise to

modern cultivated varieties, impacted on the plant's ability to assemble a rhizosphere microbiota

(Pérez-Jaramillo, Mendes & Raaijmakers, 2016). As wild ancestors of modern cultivated

varieties may hold the capacity to adapt to marginal soil conditions, there is a growing interest in

discerning the molecular mechanisms underpinning microbiota recruitment in crop wild relatives

and their contribution to plant's adaptation to the environment (Escudero-Martinez & Bulgarelli,

2019). This is particularly attractive for crops like barley (*Hordeum vulgare*), the fourth most

cultivated cereal worldwide, for which modern and wild genotypes are readily available for

experimentation (Bulgarelli et al., 2013; Alegria Terrazas et al., 2020).

The genus *Hordeum* has evolved two main indole alkaloids with allelopathic and defensive

functions, the benzoxazinoid DIBOA and gramine, whose biosynthesis appear mutually

exclusive within barley lineages (Grün, Frey & Gierl, 2005). In particular, gramine is the main

allelochemical of the lineage *H. vulgare* which has historically been implicated in defensive

responses against insects (Corcuera, 1993; Cai et al., 2009; Sun et al., 2013), as well as foliar

pathogens (Sepulveda & Corcuera, 1990; Matsuo et al., 2001) although the genetic basis of this

trait appears complex (Åhman, Tuvesson & Johansson, 2000; Macaulay, Ramsay & Åhman,

2020). Intriguingly, crop selection left a footprint on the biosynthesis of this secondary

metabolite: modern cultivated, so called 'elite', varieties (*H. vulgare* subp. *vulgare*) often fail to

accumulate gramine to levels identified in their wild relatives (*H. vulgare* subp. *spontaneum*)

(Matsuo et al., 2001; Maver et al., 2020). Of note, this apparent counter-selection for gramine

within domesticated material has been exerted on at least two distinct biosynthetic genes in

barley (Larsson et al., 2006).

To gain novel insights into the ecological significance of gramine for plant-environment

interactions, we hypothesized that the release of this secondary metabolite act as a recruitment

cue for the barley microbiota. To test this hypothesis, we exposed two 'elite', gramine-free,

barley genotypes, the cultivars Morex and Barke (Larsson et al., 2006), to exogenous

applications of gramine and we assessed the impact of these treatments on the taxonomic

composition of the prokaryotic microbiota thriving at the root-soil interface using a cultivation

independent approach.

Materials & Methods

Soil substrate

The experiments were carried out in "Quarryfield" soil, an unfertilized Scottish agricultural soil collected at the site of The James Hutton Institute, Invergowrie, Scotland (UK) (56°27'5"N 3°4'29"W). This soil was used previously to grow barley and left unplanted at least 4 years before being used for the experiments. Physical and chemical characterization: Silt: 39.35%; Clay: 11.08%; Sand: 49.58%; pH in water: 5.8; C.E.C.: 15.05 [meq/100g]; Org. Matter DUMAS: 5.95%.

Gramine adsorption

Five grams of Quarryfield soil were mixed with 10 mL of 10 mmol L^{-1} $CaCl_2$ solution containing gramine at the following concentrations: 0, 1.5, 5, 10, 15, 20, 30, 60, 100, 150, 200, 250, 300, 400, 600, 800, 1000 mg L^{-1}. Five replicates were prepared for each concentration. Soil suspensions were shaken for 24 h at room temperature. Then, suspensions were centrifuged for 5 min at 5000 g, the supernatant was collected, filtered (0.45 µm, Phenomenex) and analysed by liquid chromatography (HPLC) using the method reported in (Maver et al., 2020). Pure reagent grade gramine was used to prepare two stock solutions in water of 0.5 mol L^{-1} and 10 mmol L^{-1}. Adsorption solutions have been prepared by dilution from the stock solution with milliQ water. The adsorption rate of gramine in soil was obtained by difference between the initial and final concentration measured in the supernatant by HPLC.

Adsorption isotherms of gramine were fitted applying several nonlinear models: the two-parameter Langmuir and Freundlich isotherms and the three-parameter Sigmoidal Langmuir, Redlich-Peterson and Sips isotherms (Limousin et al., 2007; Foo & Hameed, 2010).

Plant material

Seeds of elite barley genotypes, two-row malting Barke and six-row malting Morex were

selected for this experiment. Both are well represented in barley studies: Barke as a parental

donor in the development of a nested associated mapping (NAM) population (Maurer et al.,

2015), and Morex for being the first sequenced barley genotype (International Barley Genome

Sequencing Consortium, 2012).

Growth conditions

Barley seeds were cleaned using deionized water and gently shaken for 1 min. After that water

was discarded and the whole process repeated 3 times. For seed germination, seeds were placed

on petri dishes containing semi-solid agar media at 0.5 % concentration. After a week, seedlings

displaying a comparable development were individually transferred into 12-cm pots containing

approximately 500 g of Quarryfield agricultural soil (Robertson-Albertyn et al., 2017; Alegria

Terrazas et al., 2020) previously sieved to remove stones and large debris. Plants were grown in

a randomized design in a glasshouse at 18/14°C (day/night) temperature regime with 16 h

daylight that was supplemented with artificial lighting to maintain a minimum light intensity of

200 μ mol quanta m^{-2} s^{-1}. The stock solution of gramine was prepared by adding pure reagent

grade gramine (Sigma-Aldrich, >99%), in water, sonicated for 20 min and then stored at 4 ^{o}C.

After 4 days of growing in soil, two different final gramine concentrations, 24 μmol L^{-1} and 46

μmol L^{-1}, were added directly on the soil of selected pots. Mock controls (i.e., gramine 0 μmol L^{-}

^{1}) were supplemented with sterilised water. Additional watering was performed every 2 days

with the application of 50 mL of deionized water to each pot. For each gramine treatment (i.e., 0,

24 and 46 μmol L^{-1}) we used five replicates (i.e., five individual pots) per barley genotype and

unplanted pots containing the same soil substrates used as 'bulk' soil controls (i.e., 45 pots in

total). Individual replicated pots were maintained in the glasshouse for 4 weeks post-

transplantation, when the tested genotypes reached early stem elongation, corresponding to

Zadoks stages 30–35 (Tottman, Makepeace & Broad, 1979).

Rhizosphere fractionation and sampling of soil-grown barley plants and bulk soil

Four-week-old barley plants were carefully removed from the soil, and the shoot and root

separated. The shoot was dried at 70°C for 48 h and the dry weight collected. The roots were

gently shaken to remove loosely bound soil particles, and the resulting root system and tightly

adhered soil, operationally defined as rhizosphere, was further sectioned to retain the uppermost

6 cm of the seminal root system of each sample. This root material was transferred in a sterile 50

mL falcon tube containing 15 mL of phosphate buffered saline solution (PBS). Samples were

then vortexed for 30 s, the soil sedimented for 2-3 mins, and transferred the roots in a new 50 mL

falcon tube with 15 mL PBS, in which the samples were vortexed again for 30 s to separate the

remaining rhizosphere soil from roots. The roots were discarded, the two falcon tubes were

combined in one single falcon tube, now containing the rhizosphere soil fraction, and then

centrifuged at 1,500 g for 20 mins. After centrifugation, the supernatant was discarded, and the

pellet immediately stored at -80°C. In the unplanted soil controls (i.e., the bulk soil pots), a

portion of soil corresponding to the area explored by roots was collected with a spatula and

processed as described for planted soils. Until DNA extraction, all the samples were stored at -

80°C.

Metagenomic DNA extraction from rhizosphere and bulk soil specimens

Total DNA was extracted from the rhizosphere and unplanted soil samples using FastDNA™

SPIN kit for soil (MP Biomedicals, Solon, USA) following the instructions by the manufacturer.

To assess the concentration and the quality (260/280 nm and 260/230 nm ratios) of the extracted

DNA, a Nanodrop 2000 spectrophotometer (Thermo Fisher Scientific, Waltham, USA) was used

and samples were stored at -20°C until further analysis. Aliquots at a final DNA concentration of

10 ng mL^{-1} were prepared for each sample using sterilized deionized water, and stored at -20°C.

16S rRNA gene amplicon library construction

The amplicon library was generated via a selective PCR amplification of the hypervariable V4

region of the 16S rRNA gene using the PCR primers 515F (5'-GTGCCAGCMGCCGCGGTAA-

3') and 806R (5'-GGACTACHVGGGTWCTAAT-3'). The PCR primer sequences were fused

with Illumina flow cell adapter sequences at their 5' termini and the 806R primers contained 12-

mer unique 'barcode' sequences to enable the multiplexed sequencing of several samples in a

single pool (Caporaso et al., 2012).

For each individual bulk and rhizosphere preparations, 50 ng of DNA was subjected to PCR

amplification using the Kapa HiFi HotStart PCR kit (Kapa Biosystems, Wilmington, USA). The

individual PCR reactions were performed in 20 µL final volume and contained:

• 4 µL of 5X Kapa HiFi Buffer

• 10 ng Bovine Serum Albumin (BSA) (Roche, Mannheim, Germany)

• 0.6 µL of a 10 mM Kapa dNTPs solution

• 0.6 µL of 10 µM solutions of the 515F and 806R PCR primers

• 0.25 µL of Kapa HiFi polymerase

Reactions were performed in a G-Storm GS1 thermal cycler (Gene Technologies, Somerton,

UK) using the following programme: 94°C (3 min), followed by 35 cycles of 98°C (30 s), 50°C

(30 s) 72°C (1 min) and a final step of 72°C (10 min). For each 515F-806R primer combination,

a no template control (NTC) was subjected to the same process. To minimize potential biases

originating during PCR amplifications, individual reactions were performed in triplicate and 2

independent sets of triplicate reactions per barcode were performed.

To check the amplification and/or any possible contamination, prior to purification, 6 µL aliquots

of individual replicates and the corresponding NTCs were inspected on 1.5% agarose gel. Only

samples that display the expected amplicon size and no detectable contamination in NTCs on gel

were retained for library preparation.

Individual PCR amplicons replicates were then pooled in a single plate, moving each sample to a

specific position according to their barcode. They were purified using Agencourt AMPure XP

Kit (Beckman Coulter, Brea, USA) with 0.7 µL AmPure XP beads per 1µL of sample. Following

purification, 6 µL of each sample was quantified using PicoGreen (Thermo Fisher Scientific,

Watham, USA). Once quantified, individual barcode samples were pooled to a new tube in an

equimolar ratio to generate amplicons libraries.

Illumina 16S rRNA gene amplicon sequencing

Amplicon libraries were supplemented with 15 % of a 4 pM phiX solution and run at 10 pM final

concentration on an Illumina MiSeq system with paired end 2x 150 bp reads (Caporaso et al.,

2012) as recommended, to generate FASTQ sequence files for processing and analysis.

Amplicon sequencing reads processing

Sequencing reads were processed using a customized bioinformatics pipeline as described before

(Alegria Terrazas et al., 2020). Briefly:

• Sequencing reads were subjected to quality assessment using FastQC (Andrews et al.,

2015).

• Amplicon Sequencing Variants (ASVs) were then generated using DADA2 version 1.10

(Callahan et al., 2016) and R 3.5.2 (Team R Development Core, 2018) following the

basic methodology outlined in the 'DADA2 Pipeline Tutorial (1.10)'.

• Read filtering was carried out using the DADA2 pairedFastqFilter method, trimming

10bp of sequence from the 5' of each reads using a truncQ parameter of 2 and maxEE of

2. The remainder of the reads were of high quality so no 3' trimming was deemed

necessary. The dada method was run to determine the error model with a

MAX_CONSIST parameter of 20, following which the error model converged after 9

and 12 rounds for the forward and reverse reads respectively.

• The DADA2 method was then run with the resulting error model to denoise the reads

using sample pooling, followed by read merging, then chimera removal using the

consensus method. Taxonomy assignment was carried out using the RDP Naive Bayesian

Classifier through the 'assign Taxonomy' method, with the SILVA database (Quast et al.,

2012) version 138, using a minimum bootstrap confidence of 50. The DADA2 outputs

were converted to a Phyloseq object (version 1.26.1) (McMurdie & Holmes, 2013).

• The resulting Phyloseq object was pruned in silico of ASVs assigned to chloroplast and

mitochondria, putatively representing host contamination, as well as of ASVs previously

identified as putative lab contaminant (Pietrangelo et al., 2018). Likewise, ASVs lacking

taxonomic classification at Phylum level (i.e., classified as "NAs") were removed from

the dataset.

• Next, we applied an abundance filtering for ASVs to be retained in the final dataset of 20

reads in at least 11% of the samples (i.e., representing the number of replicates for one

sample type/condition).

• Upon completion of this additional filtering step, individual ASVs were agglomerated at

Genus level.

• Finally, to control for sample-to-sample differences exceeding a factor of ~10X in

sequencing depth (Weiss et al., 2017), we downsized the Phyloseq object at 25,000 reads

per sample.

Statistical analysis

Data analysis was performed in R software v 3.5.2. The following R packages were used:

• Phyloseq v.1.26.1 (McMurdie & Holmes, 2013) for Alpha and Beta-diversity indexes;

• DESeq2 v1.22.2 (Love, Huber & Anders, 2014) for the differential analysis of microbial

enrichment;

• ggplot2 v.3.3.2 (Wickham, 2016) for data visualization;

• Vegan v.2.5-6 (Oksanen et al., 2019) for statistical analysis of beta-diversity;

• PMCMR v.4.3 (Pohlert, 2018) for non-parametric analysis of variance.

The analysis of the microbiota data was performed on filtered "Phyloseq object" described above

and linking into the analysis the mapping file (metadata information). For the Alpha-diversity

analysis, Chao1, Observed ASVs and Shannon indices were calculated using the function to

estimate richness included in the Phyloseq package.

For the Beta-diversity analysis, the rarefied ASV table was used as input to compute a Bray-

Curtis, dissimilarity matrices. This dissimilarity matrix was visualized using Principal

Coordinates Analysis (PCoA) and Canonical Analysis of Principal coordinates (CAP) (Anderson

& Willis, 2003). Beta-diversity dissimilarity matrices were tested by Permutational Multivariate

Analysis of Variance (Permanova) using Adonis function in Vegan package over 5,000

permutations, to calculate the statistical significance.

Microbial enrichment analysis

The analysis of the microbial enrichment was performed using the DESeq2 package (Love,

Huber & Anders, 2014), in order to identify the number of genera significantly enriched in pair-

wise comparisons with an adjusted p value (False Discovery Rate, FDR $p < 0.05$). The microbial

enrichment analysis was carried out between bulk soil and planted soil to evaluate the impact of

gramine on the 'rhizosphere effect'. The number of genera simultaneously enriched in the pair-

wise comparisons among the three different concentrations of gramine, in both Morex and

Barke, was plotted using the package UpSetR (Conway, Lex & Gehlenborg, 2017). In parallel

we performed a series of pair-wise comparisons between Morex and Barke using the genera

enriched in these latter specimens from unplanted soil controls. Differentially enriched genera

were visualized using ternary plots as previously described (Bulgarelli et al., 2012).

Results

Gramine application modulates microbiota composition in a microhabitat-dependent manner

Gramine availability and mobility in Quarryfield soil are two prerequisites to fulfill its allelopathic ability towards target plants and inter-organismal relationships. Thus, we assessed its adsorption applying several two- and three-parameter non-linearized isotherms models, with the best fit based on the residual sum of squares (RSSs; Fig. S1). This choice was dictated by the fact that non-linear forms permit greater accuracy in predicting parameters compared to linear forms (Foo & Hameed, 2010). The model with the best fit, the Sigmoidal Langmuir isotherm model, resulted in a sigmoidal curve, describing a cooperative adsorption phenomenon, which is common for non-polar compounds (Sparks, 2003; Limousin et al., 2007). A K_{foc}, defined as organic-carbon normalized Freundlich distribution coefficient, estimated at 1390 (OECD, 2001) classifies gramine as slightly mobile in this particular soil according to FAO Mobility Classification (FAO, 2000), yet remaining available in the soil solution. In addition, no significant differences were recorded by comparing plant dry weight at sampling time (Fig. S2, ANOVA followed by Tukey HSD test, p value > 0.05). These data indicate that Quarryfield soil is a suitable substrate to investigate the impact of the exogenous application of gramine and the root-soil interface and that this latter is not associated to obvious pleiotropic effect on barley growth.

Next, we generate a 16S rRNA gene amplicon library from 45 bulk soil and rhizosphere samples exposed to different concentration of gramine. Upon processing of the sequencing reads in silico (Material and Methods) with a protocol comparable to previous studies conducted in the same soil type (Robertson-Albertyn et al., 2017; Alegria Terrazas et al., 2020), we failed to identify a

significant effect of the treatment on the alpha-diversity parameters of the tested communities

(Fig. 1, Kruskal-Wallis test followed by Dunn's post-hoc test, individual p values > 0.05 upon

Bonferroni correction). Conversely, we observed a clear impact of the external application of

gramine on the bacterial communities thriving at the root soil interface regardless of the applied

concentration: both a Canonical Analysis of Principal Coordinates (CAP, Fig. 2) and a Principal

Coordinates Analysis (PCoA, Fig. S3) built on a Bray-Curtis distance matrix revealed a partition

of the microbiota associated to the applied treatment. Of note, the effect of gramine appeared

more pronounced on bulk soil samples than rhizosphere specimens. Congruently, a

permutational analysis of variance computed on both matrices indicated a significant effect of

the individual microhabitat, bulk soil, Morex and Barke rhizosphere, respectively (R2 ~38.6%,

Adonis test p value = 0.00020; 5 000 permutations; Table 1) and the interaction term between

gramine application and microhabitat (R2 ~6.6; Adonis test p value = 0.01540; 5 000

permutations; Table 1). Conversely, the significance of the impact of gramine *per se* failed

marginally to pass the significance threshold imposed (R2 ~4.9%; Adonis test p value = 0.05559;

5 000 permutations; Table 1). Interestingly, we obtained strikingly similar results when this

calculation was performed using individual ASVs (Fig. S4 and Fig. S5), suggesting that gramine

responsiveness in the bacteria proliferating at the root-soil interface is coded by taxonomically

conserved portions of their genomes.

Taken together, this indicates that the gramine application exerts a selective pressure on the

proliferation of a subset of members of the microbiota and this pressure appears stronger on

unplanted soil communities.

A bi-directional interference between gramine application and the barley recruitment cues of
the rhizosphere microbiota

The observation that rhizosphere profiles tend to converge on the computed ordinations (see Fig. 2 and Fig. S3), suggested that the two barley genotypes evolved the capacity of reverting, at least in part, the selective pressure of gramine on soil bacteria. To quantify this phenomenon, we implemented pair-wise comparisons between individual genera retrieved from unplanted soil and rhizosphere communities at the three levels of gramine tested. Congruently with the initial observation, we determined that the majority of genera enriched in the rhizosphere of either genotype does not respond to gramine treatment (Fig. 3, Wald test, individual p values <0.05, FDR corrected). Conversely, 18 genera whose cumulative relative abundance represented ~1.66% and ~1.12% of the Morex and Barke rhizosphere communities respectively, were identified as gramine-responsive in a genotype-independent manner (Fig. 4, Wald test, individual p values <0.05, FDR corrected). Interestingly, we noticed that the host genotype drives the enrichment of genera whose abundance in gramine-treated bulk soil samples was almost obliterated. When looking at the taxonomic affiliation of these rhizosphere- and gramine-responsive genera we observed a broad phylogenetic affiliation, encompassing members of the phyla Acidobacteria, Actinobacteria, Bacteroidetes and Proteobacteria. Next, we inspected how gramine application impacted on the genotype-driven diversification of the rhizosphere bacterial microbiota, using the number of genera a) enriched from soil and b) differentially enriched between the tested genotypes as a readout for this analysis. This analysis revealed that plants exposed to no gramine or the lowest dosage displayed a differential enrichment between cultivars of 31 and 42 genera, respectively (Fig. 5A and 5B, Wald test, individual p values <0.05, FDR corrected). Conversely, at the highest gramine dosage the host genotype effect was limited to 12 differentially enriched genera (Fig. 5C, Wald test, individual p values <0.05, FDR

corrected). When we inspected the taxonomic affiliation of these differentially enriched genera,

we made two observations. First, members of the phylum Proteobacteria accounted for the

majority of differentially recruited genera (14 out of 31 genera at Gramine 0 µmol L^{-1}, 24 out of

42 at Gramine 24 µmol L^{-1} and 6 out of 12 at Gramine 46 µmol L^{-1}, respectively). The second

observation was that despite the number of differentially regulated genera at the intermediate

dosage of gramine was comparable to mock-treated specimens, the magnitude of the host

selection was modulated by the host genotype itself, as Barke enriched for almost three times the

number of Proteobacteria upon gramine application.

Collectively our results suggest a two-pronged effect of gramine application on the barley-

microbiota. A subset of soil bacteria, whose proliferation in soil would be otherwise suppressed

by gramine application, appeared rescued by host-derived mechanisms and successfully colonise

the barley rhizosphere regardless of the treatment. At the same time, other or additional host-

derived mechanisms underpinning the genotype diversification in the rhizosphere, modulate the

effect gramine application at the barley root-soil interface.

Discussion

In this study we demonstrated that the exogenous application of the indole-alkaloid gramine produced reproducible perturbations of the microbiota thriving at the barley root-soil interface without triggering a discernable negative effect on plant growth performance on two barley elite varieties.

The observation that gramine applications impact on the soil microbiota is aligned with the findings of a recent survey performed by Schütz and co-workers to monitor the impact of the application of several plant secondary metabolites, including gramine, on bacterial thriving in a German agricultural soil (Schütz et al., 2021). Similar to our findings, this study failed to identify an effect of gramine on community richness (i.e., alpha diversity, Fig. 1), while observing a shift in community composition (i.e., beta diversity, Fig. 2). Of note, the impact of gramine applications on individual bacterial enrichments displayed an experimental effect: in our investigation this was manifested with an apparent suppression of bacterial proliferation in unplanted soil controls more pronounced than previously reported (Schütz et al., 2021). In addition to differences in applications *per se*, it is important to consider that those experiments were conducted using different soil types. For instance, soil pH, one of the main drivers of bacterial community composition in soil (Rousk et al., 2010), in the two studies differed of ~0.5 unit. This latter scenario not only contributes to explain apparent discrepancies between the studies but would also be congruent with observations that the exogenous application to grassland soils of low-molecular weight carbon compounds, mimicking plants primary metabolites, altered microbiota composition in a soil- and substrate-dependent manner (Eilers et al., 2010).

Crucially, our experimental design allowed us to discern for the first time the impact of gramine

application on plant-associated bacterial communities and compare this effect with the ones on

other secondary metabolites identified in other grasses such as maize, sorghum and oat.

For instance, we identified a limited, but significant, effect of gramine application on the

composition of the bacterial communities populating the root-soil interface (Fig. 3), indicating

that gramine *per se* (or lack thereof) does not disrupt the capacity of individual barley genotypes

of assembling a distinct rhizosphere microbiota. This is congruent with data gathered from study

conducted using maize lines impaired in the biosynthesis of benzoxazinoids grown in

agricultural soils: despite mutants were capable of recruiting a distinct microbiota, this latter was

compositionally different from the one associated with wild type lines (Hu et al., 2018;

Kudjordjie et al., 2019). We consider these observations particularly relevant as benzoxazinoids

are secondary metabolites produced by several grasses (Frey et al., 2009) with the important

exception of the *Hordeum vulgare* clade (Grün, Frey & Gierl, 2005). Similar to gramine,

benzoxazinoid display allelochemical, antimicrobial and insecticidal properties (Niemeyer &

Perez, 1994; Niemeyer, 2009). A prediction of this observation is that, within the Poaceae

family, different classes of secondary metabolites may have evolved to fine-tune microbiota

composition. Congruently, sorghum produces a species-specific allelopathic compound

designated sorgoleone (Czarnota et al., 2001; Dayan et al., 2010) capable of selectively

modulating bacterial microbiota composition as demonstrated by experiments conducted using

RNA-interference lines impaired in sorgoleone biosynthesis grown under soil conditions (Wang

et al., 2021). Likewise, oat plants impaired in the production of avenacin, a triterpenoid

defensive compound active against fungal pathogens (Papadopoulou et al., 1999), recruit a

taxonomically distinct rhizosphere microbiota compared to cognate wild type plants (Turner et

al., 2013). Interestingly, the effect of avenacin manifested predominantly at the level of the

eukaryotic component of the microbiota, particularly the protists Amoebozoa and Alveolata,

rather than the prokaryotic counterpart (Turner et al., 2013). Although differences in the

experimental and sequencing procedures existing among the aforementioned studies hinder the

capacity of establishing first principles, these observations suggest that, in cereals, species-

specific secondary metabolites act as a "gatekeepers" in a the multi-step selection process

proposed for the diversification of the plant microbiota from the surrounding soil communities

(Bulgarelli et al., 2013; Edwards et al., 2015).

The impact of gramine application on the taxonomic composition of the barley rhizosphere

microbiota revealed a relatively broad phylogenetic impact: members of 17 prokaryotic orders

responded to gramine application in a dosage- and host genotype-independent manner (Fig. 4).

Among those, of particular interest is the order Nitrosotaleales represented by the *Candidatus*

genus *Nitrosotalea*. Member of this lineage have previously been characterized as ammonia-

oxidizing archaea, i.e., responsible for the rate-limiting step in the process of nitrification

(Treusch et al., 2005), and, despite being autotrophic organisms, are capable of differential

physiological responses in the presence of organic substrates (Lehtovirta-Morley et al., 2014).

For instance, our data indicate that the application of gramine to unplanted soil communities

suppress the proliferation of this member of ammonia oxidizing archaea, pointing at a role of this

compound in the biological inhibition of nitrification, as observed for others plant-derived

compounds (Tesfamariam et al., 2014; Kaur-Bhambra et al., 2021). Yet, this scenario is difficult

to reconcile with the observation that *Candidatus* genus *Nitrosotalea* is enriched in the

rhizosphere of gramine-treated plants. A possible explanation could be derived by the niche

adaptation of ammonia oxidizing archaea: barley seedlings, similar to other grasses, have high a

rate of ammonia uptake and, in turn, this may create the optimum substrate conditions for the

proliferation of organisms like *Candidatus* genus *Nitrosotalea* despite the presence of putative

inhibitors (Thion et al., 2016). Congruently, previous experiments conducted with different

barley genotypes identified ammonia oxidizing archaea among members of the resident

rhizosphere microbiota in different soil types (Glaser et al., 2010). An alternative, but not

mutually exclusive, scenario is that the gramine inhibitory effect, is reduced by the activity of

other microorganisms in the rhizosphere compared to the unplanted soil control. As the

biological inhibition of nitrification has positive implications for sustainable crop production

(Coskun et al., 2017), it will be interesting to further investigate the relationships between

gramine biosynthesis and ammonia oxidation in prokaryotes.

Conversely, when we inspected the impact of applications in a genotype-dependent manner we

observed that the majority of microbes responding to gramine belong to the phylum

Proteobacteria (Fig. 5). Members of this phylum also respond to differential exudation of

benzoxazinoid (Jacoby, Koprivova & Kopriva, 2021), possibly representing another element of

selection of cereal secondary metabolites towards the soil biota. As Proteobacteria represent the

most abundant members of the plant microbiota across host species (Hacquard et al., 2015), the

observed gramine effect may simply mirror the dominance of this group at the root-soil interface.

Yet, a recent study conducted on maize demonstrated that Proteobacteria, specifically members

of the family Oxalobacteraceae, promote lateral root density and shoot dry weight (proxies for

plant growth) under nitrogen limiting conditions (Yu et al., 2021). As the soil tested in our

experiments is limited in the availability of nitrogen for barley growth (Terrazas et al., 2020) and

considering the enrichment of putative ammonia oxidizers archaea in the rhizosphere of treated

plants (see above), it can be speculated that the differential Proteobacterial enrichment triggered

by gramine application may be linked to nitrogen turnover in the rhizosphere. An additional

observation derived from these experiments is that gramine application triggers a differential

microbial recruitment in the two tested genotypes. A recent investigation conducted in tomato

revealed that a component of the root exudates can be induced by the exposure to a given

microbiota composition (Korenblum et al., 2020): it can therefore to be hypothesized that the

differential compositional differences observed upon gramine application may be the result of a,

genotype-dependent, fine-tuning of exudate profiles. An alternative, not mutually exclusive

scenario, is a differential rate of gramine degradation in the rhizosphere (Ghini, Burton & Gros,

1991) of the two genotypes as this would alter the bioavailability of this compound to the

resident members of the microbiota. Regardless of the scenario, it is interesting to note that

gramine applications failed to trigger a sustained enrichment of members of the Actinobacteria,

which can be considered as a hallmark of elite, gramine-free, barley genotypes grown in the

same soil type (Alegria Terrazas et al., 2020) and in other modern/ancestral plant pairs (Pérez-

Jaramillo et al., 2018).

Conclusions

Our results indicate that the application of the indole-alkaloid gramine modulates the proliferation of a subset of soil microbes with relatively broad phylogenetic assignments. This effect is two-pronged: a component of the barley microbiota responds to gramine application in a genotype- and dosage-independent manner while other or additional host-derived mechanisms, underpinning the genotype diversification in the rhizosphere, modulate the effect of gramine application with a bias for members of the phylum Proteobacteria. As gramine biosynthesis has previously been reported as stress induced (Velozo et al., 1999; Matsuo et al., 2001), we anticipate that exposure to different soil characteristics, including different microbiomes, is likely to amplify (or obliterate) the effect of this metabolite on edaphic microbes. We therefore propose to capitalise on these initial observations, and the expanding genomic resources for barley (Maurer et al., 2015; Jayakodi et al., 2020), to resolve the genetic basis of gramine biosynthesis and ultimately elucidate its adaptive value for plant-microbe interactions.

Data availability

The sequences generated in the 16S rRNA gene sequencing survey are deposited in the European Nucleotide Archive (ENA) under the accession number PRJEB39836. The version of the individual packages and scripts used to analyse the data and generate the figures of this study are available at https://github.com/Stramon1um/gramine_microbiome.

Figure legend

Figure 1. Boxplot of alpha diversity indexes. Alpha diversity Observed genera, Chao1 and Shannon indexes, of Bulk soil, Barke and Morex, at three different gramine concentrations. Individual dots depict individual biological replicates; no significant differences observed for

gramine treatment upon Kruskal-Wallis test followed by Dunn's post-hoc test, individual p

values > 0.05, Bonferroni corrected.

Figure 2. Canonical analysis of Principal Coordinates (CAP) constructed on a Bray-Curtis

dissimilarity matrix of Bulk soil, Barke and Morex, at three different gramine concentrations.

Individual shapes depict individual samples, color-coded according the gramine treatment

imposed on them. The ordination was constrained for genotype and gramine concentrations.

Figure 3. Gramine modulates bacterial abundances at the barley root-soil interface UpSetR

plots of A) Genera simultaneously enriched in pair-wise comparisons retrieved from unplanted

soil and Morex rhizosphere, B) and from unplanted soil and Barke rhizosphere, at the three

levels of gramine tested. Vertical bars denote the number of genera enriched shared or unique for

each comparison, while the horizontal bars the number of genera enriched in the indicated

gramine concentration. In A and B genera differentially enriched at individual p values <0.05,

Wald Test, FDR corrected.

Figure 4. Taxonomy and abundances of the gramine-responsive genera. Heatmap of the 18

bacteria, classified either at Order or Phylum level, significantly enriched in rhizosphere samples

in a genotype- and gramine dosage- (24 and 46 μM, respectively) independent manner.

Microbial enrichments defined at individual p values <0.05, Wald Test, FDR corrected.

Figure 5. Gramine application attenuates the genotype effect on the rhizosphere microbiota

Ternary plots depicting bacteria distribution across the indicated microhabitats in sample

exposed to A) no gramine, B) gramine 24 μM or C) gramine 46 μM. In each plot, individual dots

depict individual microbes whose size is proportional their sequencing abundances. Position of

the dots within the plots reflects the contribution of each microhabitats to microbial abundances.

Coloured dots denote genera differentially enriched between Morex and Barke (individual p

values <0.05, Wald Test, FDR corrected) color-coded according to their taxonomic affiliation at

phylum level.

Table 1. Permutational analysis of rhizosphere microbiota variance computed on Bray-Curtis

matrix, variance explained by the indicated variables and corresponding statistical significance.

Figure S1. Non-linearized isotherm adsorption models of the alkaloid gramine in Quarryfield

soil. Adsorption isotherms of gramine in Quarryfield soil were fitted applying several nonlinear

models: the two-parameter Langmuir and Freundlich isotherms and the three-parameter

Sigmoidal Langmuir, Redlich-Peterson and Sips isotherms, along with their corresponding

Residual Sum of Squares (RSSs) indicating the fitness of the model to the data. Ce = gramine

concentration at equilibrium, qe = gramine concentration adsorbed on the soil.

Figure S2. Above-ground performance of the tested genotypes. Stem dry weight of Barke and

Morex barley plants subjected to different gramine concentrations at the time of sampling. No

significant differences using ANOVA followed by Tukey HSD test, p value > 0.05

Figure S3. Principal Coordinates Analysis (PCoA) constructed on a Bray-Curtis dissimilarity

matrix of Bulk soil, Barke and Morex, at three different gramine concentrations. Individual

shapes depict individual samples, color-coded according the gramine treatment imposed on

them. The ordination was constrained for genotype and gramine concentrations.

Figure S4. Alpha diversity computed at Amplicon Sequencing Variants level. Alpha diversity

Observed ASVs, Chao1 and Shannon indexes, of Bulk soil, Barke and Morex, at three different

gramine concentrations. Individual dots depict individual biological replicates; no significant

differences observed.

Figure S5. CAP Bray-Curtis computed at Amplicon Sequencing Variants level. Individual

shapes depict individual samples, color-coded according the gramine treatment imposed on

them. The ordination was constrained for genotype and gramine concentrations.

Funding

The work presented in this manuscript was supported by a Royal Society of Edinburgh/Scottish Government Personal Research Fellowship co-funded by Marie Curie Actions and a UK Research and Innovation grant (BB/S002871/1) awarded to DB.

Acknowledgement

We thank Malcolm Macaulay, (The James Hutton Institute, UK) for the technical assistance in preparing the amplicon sequencing library and Rodrigo Alegria Terrazas (Mohammed VI Polytechnic University, Morocco) for the critical comments on the manuscript.

References

Åhman I, Tuvesson S, Johansson M. 2000. Does indole alkaloid gramine confer resistance in barley to aphid *Rhopalosiphum padi*? *Journal of chemical ecology* 26:233–255. DOI: 10.1023/A:1005405915031.

Alegria Terrazas R, Balbirnie-Cumming K, Morris J, Hedley PE, Russell J, Paterson E, Baggs EM, Fridman E, Bulgarelli D. 2020. A footprint of plant eco-geographic adaptation on the composition of the barley rhizosphere bacterial microbiota. *Scientific Reports* 10:1–14. DOI: 10.1038/s41598-020-69672-x.

Alegria Terrazas R, Giles C, Paterson E, Robertson-Albertyn S, Cesco S, Mimmo T, Pii Y, Bulgarelli D. 2016. *Plant-microbiota interactions as a driver of the mineral Turnover in the Rhizosphere*. Elsevier Ltd. DOI: 10.1016/bs.aambs.2016.03.001.

Anderson MJ, Willis TJ. 2003. Canonical analysis of principal coordinates: A useful method of constrained ordination for ecology. *Ecology* 84:511–525. DOI: 10.1890/0012-9658(2003)084[0511:CAOPCA]2.0.CO;2.

Andrews S, Krueger F, Seconds-Pichon A, Biggins F, Wingett S. 2015. FastQC. A quality control tool for high throughput sequence data. Babraham Bioinformatics. *Babraham Institute* 1:1.

Bulgarelli D, Rott M, Schlaeppi K, Ver Loren van Themaat E, Ahmadinejad N, Assenza F, Rauf P, Huettel B, Reinhardt R, Schmelzer E, Peplies J, Gloeckner FO, Amann R, Eickhorst T, Schulze-Lefert P. 2012. Revealing structure and assembly cues for Arabidopsis root-inhabiting bacterial microbiota. *Nature* 488:91–5. DOI: 10.1038/nature11336.

Bulgarelli D, Schlaeppi K, Spaepen S, Ver Loren van Themaat E, Schulze-Lefert P. 2013. Structure and functions of the bacterial microbiota of plants. *Annual review of plant biology* 64:807–38. DOI: 10.1146/annurev-arplant-050312-120106.

Cai QN, Han Y, Cao YZ, Hu Y, Zhao X, Bi JL. 2009. Detoxification of gramine by the cereal aphid sitobion avenae. *Journal of Chemical Ecology* 35:320–325. DOI: 10.1007/s10886-009-9603-y.

Callahan BJ, McMurdie PJ, Rosen MJ, Han AW, Johnson AJA, Holmes SP. 2016. DADA2: High-resolution sample inference from Illumina amplicon data. *Nature Methods* 13:581–583. DOI: 10.1038/nmeth.3869.

Canarini A, Kaiser C, Merchant A, Richter A, Wanek W. 2019. Root exudation of primary metabolites: Mechanisms and their roles in plant responses to environmental stimuli. *Frontiers in Plant Science* 10:157. DOI: 10.3389/fpls.2019.00157.

Caporaso JG, Lauber CL, Walters WA, Berg-Lyons D, Huntley J, Fierer N, Owens SM, Betley J, Fraser L, Bauer M, Gormley N, Gilbert JA, Smith G, Knight R. 2012. Ultra-high-throughput microbial community analysis on the Illumina HiSeq and MiSeq platforms. *ISME Journal* 6:1621–1624. DOI: 10.1038/ismej.2012.8.

Conway JR, Lex A, Gehlenborg N. 2017. UpSetR: An R package for the visualization of intersecting sets and their properties. *Bioinformatics* 33:2938–2940. DOI: 10.1093/bioinformatics/btx364.

Corcuera LJ. 1993. Biochemical basis for the resistance of barley to aphids. *Phytochemistry* 33:741–747. DOI: 10.1016/0031-9422(93)85267-U.

Coskun D, Britto DT, Shi W, Kronzucker HJ. 2017. Nitrogen transformations in modern agriculture and the role of biological nitrification inhibition. *Nature Plants* 3:1–10. DOI: 10.1038/nplants.2017.74.

Czarnota MA, Paul RN, Dayan FE, Nimbal CI, Weston LA. 2001. Mode of Action, Localization

of Production, Chemical Nature, and Activity of Sorgoleone: A Potent PSII Inhibitor in Sorghum spp. Root Exudates 1. *Weed Technology* 15:813–825. DOI: 10.1614/0890-037x(2001)015[0813:moalop]2.0.co;2.

Dakora FD, Phillips DA. 2002. Root exudates as mediators of mineral acquisition in low-nutrient environments. In: *Food Security in Nutrient-Stressed Environments: Exploiting Plants' Genetic Capabilities*. Springer Netherlands, 201–213. DOI: 10.1007/978-94-017-1570-6_23.

Dayan FE, Rimando AM, Pan Z, Baerson SR, Gimsing AL, Duke SO. 2010. Sorgoleone. *Phytochemistry* 71:1032–1039. DOI: 10.1016/j.phytochem.2010.03.011.

Edwards J, Johnson C, Santos-Medellín C, Lurie E, Podishetty NK, Bhatnagar S, Eisen JA, Sundaresan V, Jeffery LD. 2015. Structure, variation, and assembly of the root-associated microbiomes of rice. *Proceedings of the National Academy of Sciences of the United States of America* 112:E911–E920. DOI: 10.1073/pnas.1414592112.

Eilers KG, Lauber CL, Knight R, Fierer N. 2010. Shifts in bacterial community structure associated with inputs of low molecular weight carbon compounds to soil. *Soil Biology and Biochemistry* 42:896–903. DOI: 10.1016/j.soilbio.2010.02.003.

Escudero-Martinez C, Bulgarelli D. 2019. Tracing the evolutionary routes of plant–microbiota interactions. *Current Opinion in Microbiology* 49:34–40. DOI: 10.1016/j.mib.2019.09.013.

FAO. 2000. *FAO PESTICIDE DISPOSAL SERIES 8 Assessing soil contamination A reference manual*.

Foo KY, Hameed BH. 2010. Insights into the modeling of adsorption isotherm systems. *Chemical Engineering Journal* 156:2–10. DOI: 10.1016/J.CEJ.2009.09.013.

Frey M, Schullehner K, Dick R, Fiesselmann A, Gierl A. 2009. Benzoxazinoid biosynthesis, a model for evolution of secondary metabolic pathways in plants. *Phytochemistry* 70:1645–1651. DOI: 10.1016/j.phytochem.2009.05.012.

Ghini AA, Burton G, Gros EG. 1991. Biodegradation of the indolic system of gramine in Hordeum vulgare. *Phytochemistry* 30:779–784. DOI: 10.1016/0031-9422(91)85251-T.

Glaser K, Hackl E, Inselsbacher E, Strauss J, Wanek W, Zechmeister-Boltenstern S, Sessitsch A. 2010. Dynamics of ammonia-oxidizing communities in barley-planted bulk soil and rhizosphere following nitrate and ammonium fertilizer amendment. *FEMS Microbiology Ecology* 74:575–591. DOI: 10.1111/j.1574-6941.2010.00970.x.

Grün S, Frey M, Gierl A. 2005. Evolution of the indole alkaloid biosynthesis in the genus Hordeum: Distribution of gramine and DIBOA and isolation of the benzoxazinoid biosynthesis genes from Hordeum lechleri. *Phytochemistry* 66:1264–1272. DOI: 10.1016/j.phytochem.2005.01.024.

Hacquard S, Garrido-Oter R, González A, Spaepen S, Ackermann G, Lebeis S, McHardy AC, Dangl JL, Knight R, Ley R, Schulze-Lefert P. 2015. Microbiota and host nutrition across plant and animal kingdoms. *Cell Host and Microbe* 17:603–616. DOI: 10.1016/j.chom.2015.04.009.

Hu L, Robert CAM, Cadot S, Zhang X, Ye M, Li B, Manzo D, Chervet N, Steinger T, van der Heijden MGA, Schlaeppi K, Erb M. 2018. Root exudate metabolites drive plant-soil feedbacks on growth and defense by shaping the rhizosphere microbiota. *Nature Communications* 9:2738. DOI: 10.1038/s41467-018-05122-7.

Jacoby RP, Koprivova A, Kopriva S. 2021. Pinpointing secondary metabolites that shape the composition and function of the plant microbiome. *Journal of Experimental Botany* 72:57–69. DOI: 10.1093/jxb/eraa424.

Jayakodi M, Padmarasu S, Haberer G, Bonthala VS, Gundlach H, Monat C, Lux T, Kamal N,
Lang D, Himmelbach A, Ens J, Zhang X-Q, Angessa TT, Zhou G, Tan C, Hill C, Wang P,
Schreiber M, Fiebig A, Budak H, Xu D, Zhang J, Wang C, Guo G, Zhang G, Mochida K,
Hirayama T, Sato K, Chalmers KJ, Langridge P, Waugh R, Pozniak CJ, Scholz U, Mayer
KFX, Spannagl M, Li C, Mascher M, Stein N. 2020. The barley pan-genome reveals the
hidden legacy of mutation breeding. DOI: 10.1038/s41586-020-2947-8.
Jones DL, Nguyen C, Finlay RD. 2009. Carbon flow in the rhizosphere: Carbon trading at the
soil-root interface. *Plant and Soil* 321:5–33. DOI: 10.1007/s11104-009-9925-0.
Kaur-Bhambra J, Wardak DLR, Prosser JI, Gubry-Rangin C. 2021. Revisiting plant biological
nitrification inhibition efficiency using multiple archaeal and bacterial ammonia-oxidising
cultures. *Biology and Fertility of Soils*. DOI: 10.1007/s00374-020-01533-1.
Korenblum E, Dong Y, Szymanski J, Panda S, Jozwiak A, Massalha H, Meir S, Rogachev I,
Aharoni A. 2020. Rhizosphere microbiome mediates systemic root metabolite exudation by
root-to-root signaling. *Proceedings of the National Academy of Sciences of the United*
States of America 117:3874–3883. DOI: 10.1073/pnas.1912130117.
Kudjordjie EN, Sapkota R, Steffensen SK, Fomsgaard IS, Nicolaisen M. 2019. Maize
synthesized benzoxazinoids affect the host associated microbiome. *Microbiome* 7:1–17.
DOI: 10.1186/s40168-019-0677-7.
Larsson KAE, Zetterlund I, Delp G, Jonsson LM V. 2006. N-Methyltransferase involved in
gramine biosynthesis in barley: Cloning and characterization. *Phytochemistry* 67:2002–
2008. DOI: 10.1016/j.phytochem.2006.06.036.
Lehtovirta-Morley LE, Ge C, Ross J, Yao H, Nicol GW, Prosser JI. 2014. Characterisation of
terrestrial acidophilic archaeal ammonia oxidisers and their inhibition and stimulation by
organic compounds. *FEMS Microbiology Ecology* 89:542–552. DOI: 10.1111/1574-
6941.12353.
Limousin G, Gaudet JP, Charlet L, Szenknect S, Barthès V, Krimissa M. 2007. Sorption
isotherms: A review on physical bases, modeling and measurement. *Applied Geochemistry*
22:249–275. DOI: 10.1016/j.apgeochem.2006.09.010.
Love MI, Huber W, Anders S. 2014. Moderated estimation of fold change and dispersion for
RNA-seq data with DESeq2. *Genome Biology* 15:1–21. DOI: 10.1186/s13059-014-0550-8.
Lugtenberg B, Kamilova F. 2009. Plant-growth-promoting rhizobacteria. *Annual Review of*
Microbiology 63:541–556. DOI: 10.1146/annurev.micro.62.081307.162918.
Macaulay M, Ramsay L, Åhman I. 2020. Quantitative trait locus for resistance to the aphid
Rhopalosiphum padi L. in barley (Hordeum vulgare L.) is not linked with a genomic region
for gramine concentration. *Arthropod-Plant Interactions* 14:57–65. DOI: 10.1007/s11829-
019-09727-7.
Matsuo H, Taniguchi K, Hiramoto T, Yamada T, Ichinose Y, Toyoda K, Takeda K, Shiraishi T.
2001. Gramine increase associated with rapid and transient systemic resistance in barley
seedlings induced by mechanical and biological stresses. *Plant & cell physiology* 42:1103–
11.
Maurer A, Draba V, Jiang Y, Schnaithmann F, Sharma R, Schumann E, Kilian B, Reif JC, Pillen
K. 2015. Modelling the genetic architecture of flowering time control in barley through
nested association mapping. *BMC Genomics* 16:290. DOI: 10.1186/s12864-015-1459-7.
Maver M, Miras-Moreno B, Lucini L, Trevisan M, Pii Y, Cesco S, Mimmo T. 2020. New
insights in the allelopathic traits of different barley genotypes: Middle Eastern and Tibetan
wild-relative accessions vs. cultivated modern barley. *PLoS ONE* 15:1–18. DOI:

10.1371/journal.pone.0231976.

McMurdie PJ, Holmes S. 2013. Phyloseq: An R Package for Reproducible Interactive Analysis
and Graphics of Microbiome Census Data. *PLoS ONE* 8. DOI:
10.1371/journal.pone.0061217.

Niemeyer HM. 2009. Hydroxamic acids derived from 2-hydroxy-2h-1,4-benzoxazin-3(4h)-one:
Key defense chemicals of cereals. *Journal of Agricultural and Food Chemistry* 57:1677–
1695. DOI: 10.1021/jf8034034.

Niemeyer HM, Perez FJ. 1994. Potential of Hydroxamic Acids in the Control of Cereal Pests,
Diseases, and Weeds. In: 260–270. DOI: 10.1021/bk-1995-0582.ch019.

OECD. 2001. *OECD Guidline for testing chemicals Estimation of the Adsorption Coefficient (K*
oc) on Soil and on Sewage Sludge using High Performance Liquid Chromatography
(HPLC) 121.

Oksanen AJ, Blanchet FG, Friendly M, Kindt R, Legendre P, Mcglinn D, Minchin PR, Hara
RBO, Simpson GL, Solymos P, Stevens MHH, Szoecs E. 2019. Package ' vegan .'

Papadopoulou K, Melton RE, Leggett M, Daniels MJ, Osbourn AE. 1999. Compromised disease
resistance in saponin-deficient plants. *Proceedings of the National Academy of Sciences of*
the United States of America 96:12923–12928. DOI: 10.1073/pnas.96.22.12923.

Pascale A, Proietti S, Pantelides IS, Stringlis IA. 2020. Modulation of the Root Microbiome by
Plant Molecules: The Basis for Targeted Disease Suppression and Plant Growth Promotion.
Frontiers in Plant Science 10:1–23. DOI: 10.3389/fpls.2019.01741.

Pérez-Jaramillo JE, Carrión VJ, de Hollander M, Raaijmakers JM. 2018. The wild side of plant
microbiomes. *Microbiome* 6:143. DOI: 10.1186/s40168-018-0519-z.

Pérez-Jaramillo JE, Mendes R, Raaijmakers JM. 2016. Impact of plant domestication on
rhizosphere microbiome assembly and functions. *Plant Molecular Biology* 90:635–644.
DOI: 10.1007/s11103-015-0337-7.

Pietrangelo L, Bucci A, Maiuro L, Bulgarelli D, Naclerio G. 2018. Unraveling the composition
of the root-associated bacterial microbiota of Phragmites australis and Typha latifolia.
Frontiers in Microbiology 9:1–13. DOI: 10.3389/fmicb.2018.01650.

Pohlert T. 2018. Package 'PMCMRplus': Calculate Pairwise Multiple Comparisons of Mean
Rank Sums Extended. :1–8. DOI: 10.18637/jss.v080.i01>.

Quast C, Pruesse E, Yilmaz P, Gerken J, Schweer T, Yarza P, Peplies J, Glöckner FO. 2012. The
SILVA ribosomal RNA gene database project: improved data processing and web-based
tools. *Nucleic acids research* 41:D590--D596.

Robertson-Albertyn S, Alegria Terrazas R, Balbirnie K, Blank M, Janiak A, Szarejko I,
Chmielewska B, Karcz J, Morris J, Hedley PE, George TS, Bulgarelli D. 2017. Root Hair
Mutations Displace the Barley Rhizosphere Microbiota. *Frontiers in Plant Science* 8:1094.
DOI: 10.3389/fpls.2017.01094.

Rolfe SA, Griffiths J, Ton J. 2019. Crying out for help with root exudates: adaptive mechanisms
by which stressed plants assemble health-promoting soil microbiomes. *Current Opinion in*
Microbiology 49:73–82. DOI: 10.1016/j.mib.2019.10.003.

Rousk J, Bååth E, Brookes PC, Lauber CL, Lozupone C, Caporaso JG, Knight R, Fierer N. 2010.
Soil bacterial and fungal communities across a pH gradient in an arable soil. *ISME Journal*
4:1340–1351. DOI: 10.1038/ismej.2010.58.

Schlaeppi K, Bulgarelli D. 2015. The Plant Microbiome at Work. *Molecular Plant-Microbe*
Interactions 28:212–217. DOI: 10.1094/MPMI-10-14-0334-FI.

Schütz V, Frindte K, Cui J, Zhang P, Hacquard S, Schulze-Lefert P, Knief C, Schulz M,

Dörmann P. 2021. Differential Impact of Plant Secondary Metabolites on the Soil Microbiota. *Frontiers in Microbiology* 12:666010. DOI: 10.3389/fmicb.2021.666010.

Sepulveda BA, Corcuera LJ. 1990. Effect of gramine on the susceptibility of barley leaves to Pseudomonas syringae. *Phytochemistry* 29:465–467. DOI: 10.1016/0031-9422(90)85098-Z.

Sparks DL. 2003. Sorption Phenomena on Soils. *Environmental Soil Chemistry*:133–186. DOI: 10.1016/b978-012656446-4/50005-0.

Sun XQ, Zhang MX, Yu JY, Jin Y, Ling B, Du JP, Li GH, Qin QM, Cai QN. 2013. Glutathione S-Transferase of Brown Planthoppers (Nilaparvata lugens) Is Essential for Their Adaptation to Gramine-Containing Host Plants. *PLoS ONE* 8. DOI: 10.1371/journal.pone.0064026.

Team R Development Core. 2018. A Language and Environment for Statistical Computing. *R Foundation for Statistical Computing* 2:https://www.R-project.org.

Terrazas RA, Robertson-Albertyn S, Corral AM, Escudero-Martinez C, Balbirnie-Cumming K, Morris J, Hedley PE, Paterson E, Baggs EM, Abbott J, Bulgarelli D. 2020. Nitrogen availability modulates the host control of the barley rhizosphere microbiota. *bioRxiv*.

Tesfamariam T, Yoshinaga H, Deshpande SP, Srinivasa Rao P, Sahrawat KL, Ando Y, Nakahara K, Hash CT, Subbarao G V. 2014. Biological nitrification inhibition in sorghum: The role of sorgoleone production. *Plant and Soil* 379:325–335. DOI: 10.1007/s11104-014-2075-z.

Thion CE, Poirel JD, Cornulier T, De Vries FT, Bardgett RD, Prosser JI. 2016. Plant nitrogen-use strategy as a driver of rhizosphere archaeal and bacterial ammonia oxidiser abundance. *FEMS Microbiology Ecology* 92:91. DOI: 10.1093/femsec/fiw091.

Tottman DR, Makepeace RJ, Broad H. 1979. An explanation of the decimal code for the growth stages of cereals, with illustrations. *Annals of Applied Biology* 93:221–234. DOI: 10.1111/j.1744-7348.1979.tb06534.x.

Treusch AH, Leininger S, Kietzin A, Schuster SC, Klenk HP, Schleper C. 2005. Novel genes for nitrite reductase and Amo-related proteins indicate a role of uncultivated mesophilic crenarchaeota in nitrogen cycling. *Environmental Microbiology* 7:1985–1995. DOI: 10.1111/j.1462-2920.2005.00906.x.

Turner TR, James EK, Poole PS. 2013. The plant microbiome. *Genome Biology* 14. DOI: 10.1186/gb-2013-14-6-209.

Turner TR, Ramakrishnan K, Walshaw J, Heavens D, Alston M, Swarbreck D, Osbourn A, Grant A, Poole PS. 2013. Comparative metatranscriptomics reveals kingdom level changes in the rhizosphere microbiome of plants. *ISME Journal* 7:2248–2258. DOI: 10.1038/ismej.2013.119.

Velozo JA, Alvarez RI, Wächter GA, Timmermann BN, Corcuera LJ. 1999. Increase in gramine content in barley infested by the aphid Schizaphis graminum R. *Phytochemistry* 52:1059–1061. DOI: 10.1016/S0031-9422(99)00358-1.

Wang P, Chai YN, Roston R, Dayan FE, Schachtman DP. 2021. The Sorghum bicolor Root Exudate Sorgoleone Shapes Bacterial Communities and Delays Network Formation. *mSystems* 6. DOI: 10.1128/msystems.00749-20.

Weiss S, Xu ZZ, Peddada S, Amir A, Bittinger K, Gonzalez A, Lozupone C, Zaneveld JR, Vázquez-Baeza Y, Birmingham A, Hyde ER, Knight R. 2017. Normalization and microbial differential abundance strategies depend upon data characteristics. *Microbiome* 5:1–18. DOI: 10.1186/s40168-017-0237-y.

Wickham H. 2016. *Ggplot2: elegant graphics for data analysis*. Springer.

Yu P, He X, Baer M, Beirinckx S, Tian T, Moya YAT, Zhang X, Deichmann M, Frey FP, Bresgen V, Li C, Razavi BS, Schaaf G, von Wirén N, Su Z, Bucher M, Tsuda K,

803   Goormachtig S, Chen X, Hochholdinger F. 2021. Plant flavones enrich rhizosphere
804   Oxalobacteraceae to improve maize performance under nitrogen deprivation. *Nature Plants*
805   7:481–499. DOI: 10.1038/s41477-021-00897-y.
806

1 **Table 1:**
2
3 Permutational analysis of rhizosphere microbiota variance computed on Bray-Curtis matrix, variance
4 explained by the indicated variables and corresponding statistical significance.
5

Factor	R2	Pr (>F)
Bray-Curtis		
Microhabitat	0.38595	0.00020
Treatment	0.04911	0.05559
Microhabitat:Treatment	0.06623	0.01540

G0 effect

Bulk

0.8

0.6

0.4

0.2

Morex   Barke

G24 effect

Bulk

0.8

0.6

0.4

0.2

Morex   Barke

G46 effect

Bulk

0.8

0.6

0.4

0.2

Morex   Barke

Phylum

🟢 Proteobacteria

🔵 Bacteroidota

🔴 Actinobacteriota

⚫ Others