1 **A holobiont view of island biogeography: unraveling patterns driving the nascent**
2 **diversification of a Hawaiian spider and its microbial associates**
3

4 Ellie E. Armstrong*[,1], Benoît Perez-Lamarque*[,2,3], Ke Bi[4,5,6], Cerise Chen[7,8], Leontine E.
5 Becking[9,10], Jun Ying Lim[11], Tyler Linderoth[12], Henrik Krehenwinkel[7,13], Rosemary Gillespie[7]
6
[1] Department of Biology, Stanford University, Stanford, CA, USA
[2] Institut de Biologie de l'ENS (IBENS), Département de biologie, École normale supérieure,
CNRS, INSERM, Université PSL, Paris, France
[3] Institut de Systématique, Évolution, Biodiversité (ISYEB), Muséum national d'Histoire
naturelle, CNRS, Sorbonne Université, EPHE, UA, Paris, France
[4] Computational Genomics Resource Laboratory, California Institute for Quantitative
Biosciences, University of California, Berkeley, CA, USA 94720
[5] Museum of Vertebrate Zoology, University of California, Berkeley, CA, USA 94720
[6] Ancestry, 153 Townsend St., Ste. 800 San Francisco, CA, USA 94107
[7] Department of Environmental Science, Policy and Management, University of California,
Berkeley, CA, USA
[8] Long Marine Laboratory, University of California, Santa Cruz, CA, USA
[9] Marine Animal Ecology Group, Wageningen University & Research, Wageningen, The
Netherlands
[10] Wageningen Marine Research, Den Helder, The Netherlands
[11] School of Biological Sciences, Nanyang Technological University, 60 Nanyang Drive,
Singapore 637551
[12] Department of Genetics, University of Cambridge, UK
[13] Department of Biogeography, Trier University, Trier, Germany
26
27 * Contributed equally
28
Corresponding Author: elliea@stanford.edu, benoit.perez@ens.psl.eu

Abstract (250 words)

The diversification of a host organism can be influenced by both the external environment and its

assemblage of microbes. Here, we use a young lineage of spiders, distributed along a

chronologically arranged series of volcanic mountains, to determine the evolutionary history of a

host and its associated microbial communities, altogether forming the "holobiont". Using the stick

spider *Ariamnes waikula* (Araneae, Theridiidae) on the island of Hawai'i, and outgroup taxa on

older islands, we tested whether the host spiders and their microbial constituents have responded

in similar ways to the dynamic abiotic environment of the volcanic archipelago. The expectation

was that each component of the holobiont (the spider hosts, intracellular endosymbionts, and gut

microbiota) should show a similar pattern of sequential colonization from older to younger

volcanoes. In order to investigate this, we generated ddRAD data for the host spiders and 16S

rRNA gene amplicon data from their microbiota. Results showed that the host *A. waikula* is

strongly structured by isolation, suggesting sequential colonization from older to younger

volcanoes. Similarly, the endosymbiont communities were markedly different between *Ariamnes*

species on different islands, but more homogenized among *A. waikula* populations. In contrast,

the gut microbiota was largely conserved across all populations and species, and probably mostly

environmentally derived. Our results highlight the different evolutionary trajectories of the distinct

components of the holobiont, showing the necessity of understanding the interplay between

components in order to assess any role of the microbial communities in host diversification.

Keywords: Host-associated microbes, endosymbiont, speciation, population structure, adaptive

radiation, Ariamnes, Hawaiian Islands

Introduction

Patterns of biodiversity are influenced by both ecological and evolutionary processes operating within the dynamic context of a community (Weber et al. 2017). The external environment can serve to isolate populations for various periods, and select for traits that influence the evolutionary trajectory. At the same time, a given organism also represents a community by hosting a diverse array of microbial species, many of which perform essential functions for their host. Among arthropods, associated microbial communities are often highly diverse assemblages, accounting for an extensive range of interactions with their host (Engel & Moran 2013). Many arthropods host different microbial communities occupying various niches such as the gut microbiota or intracellular endosymbionts (Hansen & Moran 2014). The importance of microbial communities for promoting the isolation of their hosts (Sharon et al. 2010) and facilitating their adaptation to novel ecological niches (O'Connor et al. 2014) has been increasingly recognized. It is thus assumed that a species' response to the dynamic changes in the environment can be dictated by the "holobiont" of host and microbial associates (Margulis & Fester 1991). Therefore, understanding the nature and the interplay between different components of the holobiont – the host and the different communities of microbes - in response to external drivers, is essential for understanding potential drivers of evolution (McFall-Ngai et al. 2013).

First considering the gut microbiota, its composition is often determined by complex interactions of environment, diet, developmental stage, and host evolutionary history (Yun et al. 2014), contributing to various functions such as host nutrition or protection against pathogens (Engel & Moran 2013). However, for some arthropod taxa, recent work also suggests that a large proportion of the arthropod gut microbiota is purely environmentally derived, highly transient, and does not always have an apparent functional relevance (Hammer et al. 2017). For example, predators may have a microbiota derived from their prey items (Kennedy et al. 2020). In contrast, functional reliance of the host on its microbial communities could warrant more stable and predictable gut microbial communities, which may otherwise be less deterministic. In such a case

(*i.e.* host dependence), the observed microbial communities may even co-evolve with their host

(Engel & Moran 2013). That may lead to a co-diversification of microbial communities and host

taxa. On the other hand, host and microbial evolutionary histories may not be tightly coupled if the

environment of the host dictates microbial assemblage on short time scales.

In contrast to the gut microbiota, endosymbionts are mostly vertically-transmitted

intracellular bacteria. They can comprise tightly coevolved taxa, supplying their host with essential

nutrients, such as bacteria of the genus *Buchnera* in aphids (Koga et al. 2003). Many other

endosymbionts manipulate the reproduction of their host, such as species in the genera

Wolbachia, *Rickettsia*, *Rickettsiella,* and *Cardinium* (Duron et al. 2017; Hoy & Jeyaprakash 2005;

Vanthournout & Hendrickx 2015; White et al. 2020; Zhang et al. 2017). These taxa can promote

cytoplasmic incompatibilities between hosts and thus enhance genetic isolation (Shropshire &

Bordenstein 2016). Some endosymbionts can also affect dispersal ability (Goodacre et al. 2006;

Pekár & Šobotník 2007, 2008), which can further impact their host's diversification. Considering

their strong effect on the reproductive system, endosymbionts often evolve in concert with their

host. The dominant endosymbiont taxon in a lineage of arthropods is often stable, and the

endosymbiont's phylogeny commonly reflects that of their host, with major endosymbiont

switching events being infrequent (Bailly-Bechet et al. 2017). Recent evolutionary divergence in

the host may thus be mirrored by differentiation among associated endosymbionts.

In summary, various environmental and evolutionary factors can differentially influence a

microbial assemblage depending on the nature of the host/microbe relationship. Some microbes

may be purely environmentally sourced, while others may closely track their host's adaptation and

diversification. A key point of interest is then dissecting the extent, conditions, and mechanisms

under which hosts and their microbial communities influence one another's evolutionary

trajectories. We pursue this task by focusing on a lineage of spiders that shows recent divergence

between populations on the youngest island of Hawaiʻi (Gillespie et al. 2018).

The Hawaiian archipelago, a hotspot volcanic chain with the islands showing a geological

chronosequence of increasing age from southeast to northwest, provides an ideal system for

tracking the interplay between a host and the different components of its microbial community,

within a constrained setting (Shaw & Gillespie 2016). Even on the youngest island of Hawaiʻi, the

volcanoes are arranged chronologically, from approximately 430,000 years old (Kohala), to the

active flows of Kīlauea, with population structure of local flora and fauna generally shaped by

progressive colonization of the newly emerged volcanoes (e.g. Blankers et al. 2018; Eldon et al.

2019; Goodman et al. 2019).

Stick-spiders in the genus *Ariamnes* (Theridiidae) have diversified rapidly across the

landscapes in the Hawaiian archipelago (Gillespie & Rivera 2007) and exhibit repeated

diversification into ecomorphs adapted to specific microhabitats (Gillespie et al. 2018). However,

their diet is conserved and they are specialized consumers of other spiders (Kennedy et al. 2018).

While the activity of *Ariamnes* is exclusively nocturnal, the ecomorphs are defined by the

microhabitat with which they are associated during the day, with the "gold" ecomorph on the

underside of leaves, the "dark" ecomorph on dark vegetation and rocks, and the "matte white"

ecomorph on white lichen (Gillespie & Rivera 2007; Gillespie *et al.* 2018). The ecomorphs are

entirely cryptic on their daytime microhabitat, suggesting that the primary selective agent for morph

differences is diurnal predators, most likely birds (Gillespie et al. 2018).

The current study focuses on a single species of the Hawaiian *Ariamnes* (*A. waikula*),

endemic to the youngest island of Hawaiʻi, to understand how the early differentiation of the host

might be linked to the different components of its holobiont. We aim to determine whether this

highly specialized spider lineage and its microbial associates have responded in the same way to

recurrent colonization events across volcanoes within the island. We hypothesize that the

population structure of *A. waikula* reflects a stepping stone colonization from older to younger

volcanoes, and that populations from geologically older sites will show increased differentiation

and higher within population diversity compared to younger sites. If microbial associates are

largely conserved (because of transmissions or host-filtering (Moran & Sloan 2015)), we predict

that the microbiota of the *A. waikula* holobiont will closely mirror the population structure of the

host, including a diversity bottleneck in younger sites (Minard et al. 2015; Brooks et al. 2016). In

addition, we do not expect significant taxonomic changes in the endosymbionts across populations

given their typically vertical transmission.

To test these predictions, we examined the population genetic structure of *A. waikula* on

Hawai'i Island, along with several outgroup species from other islands, using genome-wide single

nucleotide polymorphism (SNP) data generated using double digest RAD sequencing (ddRAD).

We then investigated how different components of their microbiota have changed as the spiders

colonized new locations. To do so, we compared the genetic structure of microbial populations to

that of their host individual using 16S rRNA gene amplicon sequencing, capturing the diversity of

both the endosymbionts and the gut microbiota.

Material and Methods

Sampling

We sampled *Ariamnes* across Hawaiʻi Island, focusing on individuals of *A. waikula*, from 6

populations, while including 2 individuals of the related *A. hiwa* (brown ecomorph). We also

sampled individuals from two other species: *A. melekalikimaka* on West Maui and *A. n. sp.*

Molokaʻi (Gillespie et al. 2018). We included *A. hiwa*, *A. melekalikimaka*, and *A. n. sp.* Molokaʻi to

confirm monophyly of the clade on the Hawaiʻi Island (as outgroups) and to compare the diversity

of the microbial communities of other species between and within islands. Individuals were

collected by hand and immediately preserved in 90% EtOH. We collected a total of 133 individuals

for sequencing (Table 1; Supplementary Tables S1 & S2). Only adults were collected for this

study, to decrease the likelihood of capturing differences driven by age.

Table 1: Individual, population, and sampling locality information for *Ariamnes* spiders in this
study. Substrate age (in years) are approximate and based on geologic estimates of the youngest
lava flow making up each sampling locality. * Samples with sufficient coverage for microbiota
analysis. ¶ Approximate ages and estimates of time available for colonization (Carson & Clague
1995). For full details see Supplementary Tables S1 and S2.

159
160

Species	Island	Population	Volcano, age (mill. years) ¶	Substrate age (years)	Individuals (ddRAD)	Microbiota analysis *
A. waikula	Hawaiʻi	Kohala	Kohala, 0.43	300,000	18	6
A. waikula	Hawaiʻi	Saddle	Mauna Loa, Kea 0.01-0.38	4,000	6	3
A. waikula	Hawaiʻi	Alili	Mauna Loa, 0.01	20,000	18	13
A. waikula	Hawaiʻi	Puʻu Makaʻala	Mauna Loa, 0.01	11,000	18	13
A. waikula	Hawaiʻi	Olaʻa	Mauna Loa, 0.01	7,500	16	11
A. waikula	Hawaiʻi	Thurston	Kīlauea, 0.004	600	13	9
A. hiwa	Hawaiʻi	Puʻu Makaʻala	Mauna Loa, 0.01	11,000	1	N/A
A. hiwa	Hawaiʻi	Thurston	Kīlauea, 0.004	600	1	N/A
A. melekalikimaka	Maui	Puʻu Kukui	W. Maui, 1.3	1,500,000	16	9
A. n. sp	Molokaʻi	Kamakou	E. Molokaʻi, 1.8	1,400,000	16	7
					Total: 123	**Total: 71**

Ariamnes - ddRAD library preparation and sequencing

To examine the population structure of the spider host, we used ddRAD to obtain reduced

representation genome-wide SNP data. Genomic DNA was extracted from spider legs with several

modifications to the Qiagen DNeasy kit protocol. Legs were first removed from each specimen

using sterile tweezers so that the abdomen remained intact for the microbial DNA analysis. DNA

was then extracted by placing the tissue in Proteinase K and lysis buffer and grinding them with a

sterile pestle to break up the exoskeleton. We then added 4uL RNase A (100 mg/ml) and

incubated the extractions for two minutes at room temperature. Tubes with tissue and extraction

solution were then placed overnight in a heat block at 56°C. The remainder of the extraction

protocol was performed following the manufacturer's instructions. We built ddRAD libraries

following an adapted protocol of Peterson *et al.* (2012) (Saarman & Pogson, 2015; see Maas et

al. 2018 for protocol optimization steps). Briefly, we started the ddRAD protocol with a total of 100

nanograms of DNA per sample. The DNA was digested using SphI-HF (rare-cutting) and MlucI

(frequent-cutting) restriction enzymes. We assessed fragmentation with a Bioanalyzer High

Sensitivity chip (Agilent). We multiplexed 15-20 individuals per library for a total of eight ddRAD

libraries. We used a Sage Science Pippen Prep to size select 451-551bp (including internal

adapters) fragments, and confirmed the sizes using a Bioanalyzer. Ten indexing polymerase chain

reaction cycles (PCRs) were run on each library to enrich for double-digested fragments and to

incorporate a unique external index for each library pool. The eight libraries were sequenced using

100bp paired-end sequencing on one Illumina HiSeq 2500 lane at the Vincent J. Coates Genomic

Sequencing Facility at UC Berkeley.

Ariamnes - ddRAD data filtering and processing

We used a custom perl script invoking a variety of external programs to filter and process

the ddRAD data (RADTOOLKIT v0.13.10; https://github.com/CGRL-QB3-UCBerkeley/RAD). Briefly,

raw fastq reads were first de-multiplexed based on the sequence composition of internal barcodes

with tolerance of one mismatch. De-multiplexed reads were removed if the expected cutting site

was not found at the beginning of the 5'-end of the sequences. The reads were then filtered using

cutadapt (Martin 2011) and Trimmomatic using default parameters (Bolger et al. 2014) to trim off

Illumina adapter contaminations and low-quality reads. The resulting cleaned forward reads of

each individual were first clustered using cd-hit (Fu et al. 2012; Li & Godzik 2006), keeping only

those clusters with at least two reads. In each cluster, we pulled out the corresponding reverse

reads based on the identifiers. Both forward and reverse clusters at the same time were kept only

if the corresponding reverse reads also formed one cluster. If reverse reads were grouped into

more than one cluster, then only the forward read cluster was kept. For each paired cluster, the

representative sequences for each forward and reverse cluster determined by cd-hit were

retained. We then merged the forward and reverse sequences using FLASH (Magoč & Salzberg

2011). If they could not be merged then they were joined by placing "N"s between the two

sequences. The resulting loci were then masked for putative repetitive and low complexity

elements, and short repeats using RepeatMasker (Smit et al. 2004) with "spider" as a database.

After masking, we eliminated loci if more than 60% of the nucleotides were Ns. The resulting

ddRAD loci from each individual were combined and clustered for all individuals. Contigs that were

at least 40 nucleotides in length as well as shared by at least 60% of all the individuals served as

a reference. Cleaned sequence reads from each individual were then aligned to the reference

using Novoalign (http://www.novocraft.com) and reads that mapped uniquely to the reference

were kept. We used Picard (http://picard.sourceforge.net) to add read groups and GATK

(McKenna et al. 2010) to perform realignment around indels in BAM format generated by

SAMtools (Li et al. 2009). We then used SAMtools/BCFtools to generate data quality control

information in VCF format. These data were then further filtered using a custom perl script,

SNPcleaner (Bi et al. 2013). We filtered out any loci with more than two called alleles. We masked

sites within 10 bp upstream and downstream of an indel. We discarded sites with a total depth

outside of the genome-wide 1st and 99th percentile. To avoid excessive heterogeneity in sample

representation among sites, we also removed alignments if more than 40% of the samples had

less than 3X coverage. The resulting sites passing all of the above filters were analyzed using

ANGSD (Korneliussen *et al.* 2014).

Since most of the individuals were expected to have low coverage (between 5-15x) we

used ANGSD (Korneliussen et al. 2014) to calculate genotype likelihoods and genotypes for the

analyses. This tool was specifically developed for population and evolutionary genomic analyses

of low coverage data. We used genotype likelihoods whenever the downstream tools allowed us

to, since genotypes called from low coverage sites involve a non-negligible amount of uncertainty

arising from randomness in allele sampling as well as sequencing or mapping errors (Crawford &

Lazzaro 2012)).

Ariamnes - Phylogenetic Analyses

First, we investigated the phylogenetic relationships between populations of *A. waikula*

and the other *Ariamnes* species to better understand the colonization patterns. We used the

Stacks pipelines (Catchen et al. 2013) to group reads into homologous loci across all individuals

and extract phylogenetically informative sites (i.e. fixed within individuals but variable between

individuals). Next, we obtained an alignment composed of 58,899 sites and performed

phylogenetic reconstruction using IQtree (Nguyen et al. 2015) combining model selection with

ModelFinder Plus (Kalyaanamoorthy et al. 2017) and assessing branch supports with 1,000

ultrafast bootstrap (Hoang et al. 2018). Finally, we rooted the tree using the *A. hiwa* individual and

calibrated it with r8s (Sanderson 2003) without specifying any absolute dating (*i.e.* setting the root

age to 1). In addition, we also reconstructed the phylogeny using pairwise genetic distances

calculated from genotype likelihoods using the software ngsDist (Viera et al. 2015) and balanced

minimum evolution using FastME (Lefort et al. 2015). To do so, we first calculated genotype

likelihoods using ANGSD, which were used to construct a pairwise distance matrix between all

individuals using ngsDist. The phylogeny was then inferred from this matrix using FastME and

bootstrapping was carried out using RaxML (Stamatakis 2014, 100 replicates).

A. waikula - Population Genetic Analyses

We explored *A. waikula* population structure using Principal Components Analysis (PCA).

We used the previously generated genotype likelihoods to calculate genotype posterior

probabilities under an allele frequency prior (-doPost 1) in binary format (-doGeno 32) for use with

ngsCovar (part of the ngsTools package; (Fumagalli et al. 2014). We used ngsCovar to calculate

a genetic covariance matrix among individuals from the genotype probabilities at sites having a

minor allele frequency of at least 0.004 to avoid noise from very rare alleles (which could be due

to sequencing errors). Comparisons between the first three principal component axes were then

plotted using R. Pairwise F_{ST} values were calculated in ANGSD from allele frequency likelihoods

(-doSaf 1) using the respective pair's genome-wide, unfolded, joint SFS as a prior for jointly

observing any combination of allele frequencies between the two populations. We used unfolded

allele frequencies by supplying the reference sequences as a pseudo-ancestral sequence, as

ANGSD is only able to accurately estimate F_{ST} using unfolded data. In order to visualize F_{ST}

distances, we performed multidimensional scaling (to 2 dimensions) with R (R Core Team, 2020;

using the *cmdscale* function).

A crucial factor underlying population structure is gene flow between populations. In order

to investigate signatures of connectivity between populations of *A. waikula*, we used two different

approaches: ngsAdmix (Skotte *et al.* 2013) and EEMS (Petkova et al. 2016). ngsAdmix is a

genotype likelihood based-tool for estimating individual admixture proportions, while EEMS uses

genotype calls to infer effective migration surfaces. We inferred ancestry proportions indicative of

admixture for different values of K (number of ancestral populations; ranging from two to six) using

ngsAdmix and R for visualization. We then generated genotype calls for the EEMS analysis using

ANGSD with the following flags: '-doMaf2', '-doGeno 2', '-doPost 1', '-doSaf 1', '-fold 1', a SNP p-

value cut-off of 1e-6, and a postCutoff value of 0.75. We only considered sites with a minimum

genotype depth of 3. We then converted the output into the adegenet (R package) input format

using PopGenTools (https://github.com/CGRL-QB3-UCBerkeley/PopGenTools) to calculate

genetic distances between all individuals. We then divided Hawaiʻi Island into grids of 10km

(Supplementary Fig. S1). We then added the samples to the grid using one GPS coordinate per

population (Supplementary Table S1). Using a stepping stone model, we then calculated migration

rates between the demes. Convergence of the MCMC runs was assessed by plotting and

inspecting the traces by eye (Supplementary Fig. S2). We performed 10 independent runs for

each of the analyses.

Next, we calculated population genetic statistics such as nucleotide diversity (Pi),

Watterson's Theta, and Tajima's D. To do so, we generated a folded SFS for each population

using ANGSD and realSFS. We then ran ANGSD using –doTheta 1 and –pest (which provides

the genome-wide SFS prior to ANGSD) and the respective output formats were converted to bed

format using *thetaStat make_bed*. Subsequently, we calculated the per population statistics using

thetaStat do_stat. The average, min, and max Tajima's D were then calculated and we further

extracted the genome-wide average Watterson's theta and Pi values. To assess whether

populations show genetic signatures indicative of serial founder events and expansions, we used

linear models in R to test whether genetic diversity (Pi or Watterson's theta) was positively related

to the age of the youngest lava flow of each sampling site (referred to as the volcano age), as

more time would allow for genetic diversity to recover in a large population.

Microbial Communities

To characterize the microbial community within the *A. waikula* hosts, a subset of 71

individuals from the eight populations on Hawaiʻi and three additional islands (Table 1;

Supplementary Tables S1 & S2) were selected for analysis. We focused on the mid and hindgut,

both located in the spider's opisthosoma. The preservation in ethanol led to considerable

shrinkage of the opisthosoma and thus did not allow us to separately dissect out the gut. Instead,

we used the whole opisthosoma to extract DNA, as described in Kennedy et al. (2020). Specimens

which did not have the opisthosoma intact were not used. The digestive tract comprises the

majority of the opisthosoma's cavity. In addition, it contains silk glands, the heart, lungs, and

gonads. The opisthosoma was removed with a sterile razor blade and then washed in ethanol to

remove external bacteria (Hammer et al. 2015). We considered it to be representative of the "gut

microbiota", even if it technically consists of the "opisthosoma microbiota", but previous studies

have shown that the gut microbiota dominates in the opisthosoma (Sheffer et al. 2020, Kennedy

et al. 2020). The tissue was then transferred into lysis buffer and finely ground with a sterile pestle.

DNA was extracted using the Gentra Puregene Tissue Kit (Qiagen, Hilden, Germany) according

to the manufacturer's protocol. Spider abdominal tissue can contain PCR inhibitors (Schrader et

al. 2012), thus we cleaned the DNA extract with 0.9X AmPure Beads XP.

We next amplified a ~300 bp fragment of the V1-V2 region of the bacterial 16S rRNA using

the Qiagen Multiplex PCR kit according to the manufacturer's protocols and using the primer pair

MS-27F (AGAGTTTGATCCTGGCTCAG) and MS-338R (TGCTGCCTCCCGTAGGAGT) (Gibson

et al. 2014). PCRs were run with 20ng of template DNA and 30 cycles at an annealing temperature

of 55°C. PCR products were separated from leftover primer by 1X AmPure Beads XP. A six-cycle

indexing PCR was performed on the cleaned products, adding dual indexes to every sample using

the Qiagen Multiplex PCR kit. Indexing was performed according to (Lange et al. 2014). The dual

indexed libraries were isolated from leftover primer as described above, quantified using a Qubit

fluorometer, and pooled in equal amounts into a single tube. The library was sequenced on an

Illumina MiSeq using V3 chemistry and 300 bp paired reads. In order to discard contaminants from

our final dataset, we also performed blank extraction controls and negative PCR controls (without

DNA template), which were sequenced along the other samples.

We used the 16S profiling analysis pipeline for Illumina paired-end sequences of the

Brazilian Microbiome Project (Pylro et al. 2014), including QIIME 1.8.0 (Caporaso et al. 2010) and

USEARCH 11 (Edgar 2013). We modified some steps of these pipelines, using our own Bash and

R scripts (R Core Team, 2020; see Data Accessibility section). The QIIME script

join_paired_ends.py was used to merge paired reads. The fastq_filter command in USEARCH

was used for quality filtering the assemblies (below a base calling error probability of 0.5, using

an average Q-score for each read). We used the Stream EDitor in UNIX to remove PCR primers

from all assembled sequences. Sequences were de-replicated using USEARCH, removing all

singletons. OTUs were generated at a similarity cutoff of 3 % or 0% (0 radius OTUs, or "Z-OTU")

from the de-replicated sequences and chimera removed de novo using USEARCH. The following

analyses were thus independently applied on the two distinct sets of OTUs. We assigned

taxonomy to the resulting OTUs using the assign_taxonomy.py script based on the Greengenes

database (http://greengenes.secondgenome.com). We removed sequences corresponding to

OTUs found in high prevalence and abundance in the different negative controls from all samples,

as these could represent contaminants.

Spiders are known to carry various endosymbiotic bacteria (White et al. 2020). These can

be vastly overrepresented in microbial analyses and thus may completely dominate the microbial

community structure. Since we did not extract the gut from individuals, endosymbionts from

outside of the gut could be particularly prevalent in our analysis. We thus separated known

endosymbionts from remaining bacterial sequences (referred to as the "gut microbiota"), resulting

in two OTU sequence files. Both these datasets were used for the following analyses separately.

OTU tables were prepared by mapping sequences back to the filtered OTU sequence files

using USEARCH. The OTU tables were rarefied to an even coverage (from 400 to 8,000 reads

with 20 replications per rarefaction depth) using the multiple_rarefactions.py script in QIIME. Given

the rarefaction curves (Supplementary Fig. S3), we chose rarefied depths at 3,200, 110 and 3,000

reads for the whole microbiota, the endosymbionts, and the gut microbiota respectively at

replicated 20 times this rarefaction, and these rarefied OTU tables were used in all the following

analyses. We plotted the relative abundances of different microbial taxa at the genus and order levels for the different *Ariamnes* populations.

We then investigated whether microbial associates showed similar diversity patterns compared to their *Ariamnes* hosts along the chronosequence. Chao1 indexes of alpha diversity were calculated from the rarefied OTU tables using QIIME (alpha_diversity.py), and to evaluate the presence of a bottleneck in microbial diversity, linear mixed models were used to test for an effect of the volcano age and the genetic diversity of *Ariamnes* host populations (Pi and Watterson's theta) on the bacterial alpha diversity. Homoscedasticity and normality of the model residuals were verified.

To measure microbiota differentiation across *Ariamnes* species and populations, we computed beta diversity between microbial communities of each individual using QIIME (beta_diversity.py with Bray-Curtis dissimilarities). Beta diversity of microbial populations was also visualized with a Principal Coordinate Analysis (PCoA) and as dendrograms using a neighbor joining reconstruction with the R-package ape (Paradis et al., 2004). To test whether individuals from the same population tend to host similar bacterial communities (in all the populations, or within Hawai'i Island only), we performed a Permutational analysis of variance (PERMANOVA; *adonis* function, vegan R-package) on the beta diversity matrices with 10,000 permutations, after having verified the homogeneity of the variances (*betadispers* function). Finally, we analyzed whether the microbiota of the *Ariamnes* holobiont mirror the host's phylogeny by testing the correlations between microbiota differentiation and host genetic distances (ngsDist distances) or between microbiota differentiation and host phylogenetic distances, using Mantel tests with 10,000 permutations (vegan R-package). These analyses were performed between populations of *A. waikula* in Hawai'i Island, and were compared to the analyses performed between populations of different *Ariamnes* species (*i.e.* including *A. melekalikimaka* on West Maui and *A. n. sp.* Moloka'i).

During all analyses of diversity, we also controlled for any batch effects during the PCR steps by assessing the correlations between proximal samples in the PCR plates.

Results

Ariamnes - Phylogenetic Analyses

We obtained approximately 210 million paired-end reads from the Illumina sequencing.

After filtering for contaminants and low-quality reads, there were approximately 83 million

remaining across the 123 demultiplexed samples. From these reads, we retained a total of

2,957,301 sites passing filters out of a possible 7,378,384 that were used in downstream analyses

and resulting in 123 individuals for population genetic analysis. On average, samples had a

coverage of 12x (3-42x) across all loci.

All phylogenetic analyses (maximum likelihood and genetic distance based) confirm that

the *A. melekalikimaka* (West Maui), *A. n. sp.* (Moloka'i) and *A. waikula* (Hawai'i Island) are

monophyletic groups (Fig. 1). Among *A. waikula* individuals, the population from Kohala (the oldest

volcano on Hawai'i Island) is sister to the clade that contains all other individuals. Except for one

individual from Saddle and two from Alili, the populations from the Saddle (between Mauna Loa

and Mauna Kea) and Alili form monophyletic clades. The populations from Ola'a, Pu'u Maka'ala

and Thurston, all sites that are close together in the saddle between the volcanoes of Mauna Loa

and Kilauea (MLKS), form one mixed clade.

Figure 1: Host phylogenetic history partially recapitulates microbiota differentiation

Phylogenetic tree of the *Ariamnes waikula* individuals across the island of Hawai'i (A). Two

specimens of *A. hiwa* (brown ecomorph) were used as outgroup taxa. Microbiota dendrograms

reconstructed from the endosymbiont community (B) and the gut microbiota (C) for the Z-OTU.

(D) map of the sampled host populations and the corresponding age of the youngest lava on each

of the areas.

A. waikula - Population genetics

We analyzed population differentiation and structure for the full data set, and a subset

including only *A. waikula* from Hawaiʻi Island. We did not include the two *A. hiwa* individuals in the

population genetic analyses since they represent a different species used as outgroup for the

phylogenetic analyses.

We found the highest genetic differentiation (F_{ST}) between the different species on these

three islands with similar pairwise levels (0.54 between Maui and Hawaiʻi Island, 0.48 for Molokaʻi

and Hawaiʼi Island and 0.48 between Molokaʻi and Maui; Table S3). This closely matches the

results from COI data (Roderick et al. 2012). *A. waikula* on Hawaiʻi Island were primarily structured

according to locality with Kohala, Alili, and Saddle being distinct from the MLKS sites (pairwise F_{ST}

range 0.03-0.15; Puʻu Makaʻala, Thurston and Olaʻa; Fig. 1A).

Regarding potential admixture between populations, using ngsAdmix first, we found good

convergence between the 50 independent runs for higher values of K for the Hawaiʻi only sampling

(Supplementary Fig. S4). Kohala forms a separate group at K=2, next is Allili at K=3 and Saddle

at K=4. At K=6 we see a slightly closer grouping of Thurston and Olaʻa, then either of the two with

Puʻu Makaʻala, but overall the MLKS populations make up one group (Fig. 2B). Second, EEMS

analyses with *A. waikula* individuals indicate potential gene flow between the MLKS populations:

Puʻu Makaʻala, Thurston, and Olaʻa (Fig. 2C). These patterns were reiterated using PCA (Fig. 1A,

Supplementary Fig. S5). PC1 explained approximately 16% of the variation and separated Kohala

and the younger sites, placing Saddle in the middle (Fig. 1A). PC2 explained approximately 4% of

the variation and primarily separated Alili from the MLKS sites, while PC3 (3%) interestingly

clustered Kohala with Puʻu Makaʻala, Thurston, and Olaʻa, while separating the Alili and Saddle

populations (Supplementary Fig. S5).

 Figure 2: Population genetic analyses of *Ariamnes waikula*. A) ngsAdmix results for K=2 to

K=8 for all *A. waikula* specimens. B) ngsAdmix results for K=2 to K=6 (from top to bottom) for all

A. waikula from Hawai'i Island only. C) EEMS analysis of all *A. waikula* specimens. Brown color

indicates barriers for gene flow (the stronger the darker), and cyan indicates gene flow between

populations (the stronger the darker).

We found that Watterson's theta was fairly even across localities on the youngest island

of Hawai'i, with the populations on younger volcanoes possessing slightly higher thetas (e.g. Pu'u

Maka'ala: 0.0085, Ola'a: 0.0086; Table 2). The Maui population had the overall lowest theta

(0.0053). Pi estimates showed a similar pattern, with little differentiation between values on the

youngest volcanic sites and Maui (0.0051-0.0056). However, Moloka'i populations had the highest

value of Pi (0.0073). Despite low variance in Pi, the positive correlation between Pi genetic

diversities and the volcano age (linear regression: t=4.6, p-value=0.06) suggested that *Ariamnes*

likely experienced successive bottlenecks along the island chronosequence during the sequential

colonizations from older to younger volcanoes. However, this correlation was of lower significance

for *A. waikula* populations within Hawai'i (Figure 3) and no other correlations between theta and

volcano age or between *Ariamnes* genetic diversity (Pi or Theta) and *Ariamnes* microbial diversity

were found.

Table 2: Genetic diversity estimates for *Ariamnes* spiders across the different sampling sites

(excluding *A. hiwa*). The average volcano age of each site used for the statistical analyses is

indicated.

Sites	Volcano age (in Myr)	Average Pi	Average Watterson's theta
Puʻu Makaʻala	0.01	0.00518	0.00847
Alili	0.01	0.00535	0.00775
Kohala	0.43	0.00538	0.00749
Olaʻa	0.01	0.00535	0.00857
Thurston	0.004	0.00520	0.00712
Saddle	0.19	0.00526	0.00636
Molokaʻi	1.8	0.00727	0.00747
West Maui	1.3	0.00559	0.00532

438 We also investigated Tajima's D, which reflects population size changes as a skew in the

439 site frequency spectrum (Supplementary Fig. S6). This statistic compares the number of pairwise

440 differences to the number of segregating sites. We found that most sampling sites showed only

441 marginally negative values, but show no significant indications of population bottlenecks or

442 expansions (average: -1.0 to -0.06; Supplementary Table S4).

Figure 3: The bottleneck in diversity along the volcano age tends to affect the *Ariamnes* hosts, but not their microbial associates

Note that Maui and Molokaʻi correspond to different *Ariamnes* species (*A. melekalikimaka* and *A. n. sp respectively*) and are only represented as a reference here: the following linear model testing the effect of volcano age on host (A-B) or microbial (C-D) diversities were only performed on *A. waikula* populations in Hawai'i.

(A) *Ariamnes* genetic diversity (Pi) estimated per population as a function of the volcano age (in Myr). We found no significant relationship between *A. waikula* genetic diversity (Pi) and the volcano age (linear model: F=1.7, df=4, p-value=0.27), despite the observed trend, probably because of the small number of sampled populations in Hawai'i (6 only).

(B) *Ariamnes* genetic diversity (Watterson's theta) estimated per population as a function of the volcano ago (in Myr). We found no significant relationship between *A. waikula* genetic diversity (Watterson's theta) and the volcano age (linear model: F=0.023, df=4, p-value=0.87).

(C) Microbial alpha diversity (Chao1 index on Z-OTUs) estimated for the endosymbionts as a function of the volcano ago (in Myr). Each dot corresponds to the average of the 20 rarefactions performed for each individual. We found no significant relationship between endosymbionts alpha diversity and the volcano age (linear mixed model: t=1.10, p-value=0.24).

(D) Microbial alpha diversity (Chao1 index on Z-OTUs) estimated for the gut community as a function of the volcano ago (in Myr). Each dot corresponds to the average of the 20 rarefactions performed for each individual. We found no significant relationship between gut microbial alpha diversity and the volcano age (linear mixed model: t=-0.08, p-value=0.94).

In addition, we did not find any correlation between the *Ariamnes* genetic diversities and the microbial alpha diversities (not shown; p-value>0.05). Similar results were obtained with 97% OTUs.

Microbial Communities

We obtained a total of 4,932,236 bacterial reads. After the quality filtering steps, demultiplexing, and removal of contaminants based on the negative controls, we obtained a total of 1,315,469 sequences, which range from 5,000 sequences to 45,000 per individual *Ariamnes* sample. Due to the largely unexplored nature of the microbial communities of Hawaiian arthropods, we could not assign species level taxonomy to the majority of microbial OTUs (538 OTUs of 571 at 97%, 615 OTUs over 1,357 Z-OTUs), but a higher proportion at the genus level (330 at 97%, and 1,193 Z-OTUs).

We did not find any significant differences in the alpha diversities of the whole microbiota

among *Ariamnes* populations or species (Supplementary Table S5). We also did not find a

significant association of the microbial alpha diversity with volcano age or the genetic diversity of

the host population (Fig. 3), even if the alpha diversities of the gut microbiota seem to be slightly

higher for host populations present on old volcanos (Fig. 3D).

Gut microbiota: After removing endosymbionts, the remaining gut microbiota showed a

homogeneous taxonomic composition of microbial taxa at the genus level across different

Ariamnes populations or species (Fig. 4B & Supplementary Fig. S7). This stability of the

composition of the gut microbiota was also confirmed at the order level (Supplementary Fig. S8).

No clear differentiation according to the islands or the host populations could be detected for the

beta diversity of the gut microbiota, either from PCoA decomposition (Supplementary Fig. S9) or

hierarchical clustering (dendrogram; Fig. 1C & Supplementary Fig. S10). However, the gut

microbiota of individuals from the same population were significantly more similar than gut

microbiota from different populations or species, even within *A. waikula* populations from Hawai'i

Island (PERMANOVA; Supplementary Table S6). Such slight patterns of clustering by host

populations were also visually detectable for some individuals from the same populations on the

PCoA decomposition (Supplementary Fig. S9), or on the dendrogram of the gut microbiota (Fig.

1C). These results indicate that, although the composition of the gut microbiota seems

taxonomically similar, there is a slight trend toward microbiota differentiation according to the host

populations. However, using Mantel tests we found no significant correlations between the gut

microbial community beta diversities and the *Ariamnes* genetic distances nor the *Ariamnes*

phylogenetic distances (Table 3 & Supplementary Table S7), suggesting that there is likely no

pattern of sequential colonization from older to younger volcanoes (stepping stone colonization)

in these gut microbial communities in opposition to their spider hosts.

Figure 4: Relative abundances of endosymbiont genera per population (A) and gut symbionts per

spider population (B) defined from rarefied Z-OTU table. The relative abundance of each OTU is

delimited using the horizontal grey lines and are colored according to the OTU genera. Rare

genera (representing less than 1% of the abundance are merged together). Note that Maui and

Moloka'i correspond to different *Ariamnes* species (*A. melekalikimaka* and *A. n. sp respectively*)

whereas the other populations in Hawai'i correspond to *A. waikula*.

Endosymbiont community: In contrast to the gut microbiota, the endosymbiont community

showed considerable variation between the different species of *Ariamnes* (Fig. 1B). *Ariamnes*

melekalikimaka from West Maui mostly carry *Wolbachia*, while *Rickettsia* dominate the

endosymbiont community on *A. n. sp.* Moloka'i, and *Rickettsiella* the populations of *A. waikula* on

Hawai'i Island (Fig. 4A & Supplementary Fig. S7). The pronounced turnover of endosymbiont taxa

between host populations is also mirrored in differences of beta diversity. Individuals of the

different species from different islands form clearly separated clusters in our PCoA plots and

cluster dendrograms for the endosymbiont communities (Fig. 1B, Supplementary Figs. S9 & S10).

PCoA plots and cluster dendrograms exhibit a strong pattern of phylosymbiosis, where the

divergences between microbiota reflect the phylogenetic distances between the *Ariamnes* hosts.

Mantel tests showed a significant correlation between the endosymbiont community beta diversity

and the *Ariamnes* genetic distances (P<0.0001; Table 3 & Supplementary Table S7). Within

Hawaiʻi Island, endosymbiont communities of *A. waikula* from the same population tend to be more

similar than between populations (Supplementary Table S6; *e.g.* individuals from Olaʻa tend to

cluster together in Fig. 1B), suggesting a slight population differentiation. However, we found no

strong correlation between microbial beta diversity and *A. waikula* genetic distances using Mantel

tests (Table 3 & Supplementary Table S7).

Table 3: Mantel tests comparing the microbial beta diversity dis-similarities to the genetic

distances or the phylogenetic distances of their associated *Ariamnes*. Beta diversity dissimilarities

were computed on rarefied Z-OTU tables using the Bray-Curtis dissimilarities. Bold values

represent significant correlations. Mantel tests were either performed on all populations (between

Ariamnes species and *A. waikula* populations) or only on *A. waikula* populations.

Matrix 1	Matrix 2	All populations		*A. waikula* populations	
		Correlation	p-value	Correlation	p-value
Whole microbiota beta diversity distances	***Ariamnes* genetic distances**	**0.324**	**9.9e-05**	0.063	0.20
	***Ariamnes* phylo-genetic distances**	**0.334**	**9.9e-05**	0.012	0.40
Endosymbionts beta diversity distances	***Ariamnes* genetic distances**	**0.619**	**9.9e-05**	0.015	0.49
	***Ariamnes* phylo-genetic distances**	**0.642**	**9.9e-05**	0.047	0.32
Gut microbiota beta diversity distances	***Ariamnes* genetic distances**	-0.048	0.66	-0.052	0.10
	***Ariamnes* phylo-genetic distances**	-0.038	0.63	-0.139	0.93

Discussion

In this study, we examined the early stages of diversification of Hawaiian *Ariamnes* spiders in concert with the different components of their associated microbiota on the youngest island of the Hawaiian archipelago. Our results show a strong effect of isolation by distance in the host and contrasting phylogenetic patterns among both the endosymbiont community and the gut microbiota they harbor, suggesting eco-evolutionary distinctiveness of their host and each of the two microbial components.

Isolation by distance explains Ariamnes diversification

Our analyses suggest that the nascent diversification of *A. waikula* is mainly associated with isolation by distance, where island structure or isolation (on separate volcanoes) is the main barrier to gene flow. The different populations of *A. waikula* show a predictable pattern of colonization across Hawaiʻi Island from the oldest (Kohala) to the youngest (MLKS) sites (Fig. 1), with little genetic structure between the geographically close MLKS sites of Olaʻa, Puʻu Makaʻala, and Thurston (PCA, Supplementary Fig. S5; FST, Supplementary Table S3; Admixture, Fig. 2B). Our analyses suggest that gene flow is still occurring between these three populations, and that recent colonization events have not yet led to differentiation in this area (Roderick et al. 2012). Interestingly, the nucleotide diversity (Pi) tends to increase with volcano age, but their relationship is not significant for *A. waikula* populations. In addition, the pattern is less clear for Watterson's theta, likely due to loss of rare alleles during colonization and the short time frame for new mutations to arise subsequent to the colonization event. Notably, despite the popular hypothesis that sequential colonizations result in genetic bottlenecks, we show little detectable genetic evidence for such events here. Larger sample sizes and additional genetic data should be obtained to investigate these patterns further.

Patterns of genetic differentiation suggest a potentially complicated course of colonization

across Hawaiʻi Island for *A. waikula*. Although Kohala, which represents the oldest volcano, was

repeatedly characterized as distinct from other populations in PCA, phylogenetic, and admixture

analyses, both Saddle and Alili also presented as distinct. These patterns are potentially explained

by a shifting mosaic population structure (Carson et al 1990) with multiple colonization events,

historical connectivity, or ongoing admixture. Sampling of additional sites, individuals, and species

will be required to disentangle the historical sequence of events explaining the relationship

between these sites. Our results are consistent with other population genetic studies on Hawaiʻi

Island showing strong differentiation and potential speciation of lineages on the different volcanoes

within the island (planthoppers, Goodman et al. 2019; flies, Eldon et al. 2019; crickets, Blankers

et al. 2018).

Major endosymbiont changes after potential colonization event

We found major shifts among the dominant endosymbiont genera across the different

islands and *Ariamnes* species: Species of *Ariamnes* from the two geologically oldest sites

(Molokaʻi and Maui) each harbor a unique endosymbiont genus (*Rickettsia* and *Wolbachia*

respectively) while the different populations of *A. waikula* all have a mix of endosymbionts

dominated by the genus *Rickettsiella*. These shifts in endosymbionts likely contribute to the

congruence between the host phylogeny and the dendrogram of microbiota dissimilarity (a pattern

referred as phylosymbiosis (Brooks et al. 2016). These patterns are consistent with the maternal

inheritance and vertical transmission of the endosymbionts (Duron et al. 2008) and such pattern

have been similarly observed in other arthropods (e.g. a *Camponotus* ant system and its

associated endosymbiont *Candidatus Blochmania* (Degnan et al. 2004). Therefore, the overall

evolutionary patterns of the *Ariamnes* host are reflected in the endosymbiont component of the

microbiota. However, from our study, we cannot say whether endosymbionts played a role in

promoting the genetic isolation of *A. waikula* populations; this hypothesis would require

experimental manipulation of microbial communities within different host populations.

The observed turnover of dominant endosymbiont genera between islands and host

populations suggests that the endosymbiont vertical transmission happens unexpectedly fast. A

previous study reported turnover rates over time for the bacterial genus *Wolbachia* colonizing

arthropods from French Polynesia of 7 Myr on average and found that *Wolbachia* colonization did

not seem related to recent events of isolation mediated by island structure (Bailly-Bechet et al.

2017). Given that the islands and thus the spider populations studied here are separated by <2

Myr (Gillespie et al. 2018), this suggests that the turnover in *Ariamnes* endosymbiotic

communities, strongly linked to the Hawaiian archipelago structure, is much faster than the

extinction dynamics of *Wolbachia* in the French Polynesian arthropods. Predation on other

arthropods and cannibalism, which are particularly important for *Ariamnes* spiders (Whitehouse et

al. 2002), including *A. waikula* (Kennedy et al. 2018), are a potential avenue for increasing the

chance of endosymbiont horizontal transfer (Su et al. 2019). Such a mechanism may explain the

high endosymbiont turnover observed in this system. However, from our current analyses, it is

impossible to separate a scenario of recent horizontal shifts of the endosymbionts, where

endosymbionts are acquired from the environment, from a scenario of older shifts, where

endosymbionts are acquired primarily during colonization events. Testing this will require

modeling host-microbiota evolution, which is currently challenging on such short evolutionary

timescales (Groussin et al. 2017; Perez-Lamarque & Morlon 2019).

Several previous studies have suggested that endosymbionts such as *Rickettsia* enhance

spider dispersal (Goodacre et al. 2009; Pekár & Šobotník 2007, 2008). However, although

colonized by various endosymbionts including *Rickettsia*, most populations of *A. waikula* are highly

locally endemic which suggests that species within a given genus of endosymbiont are not

associated with the loss of the host's dispersal here.

Similarity of the gut microbiota

Across all spider populations, we observe a conserved gut microbiota: Over approximately

one million years, the composition seems to have remained stable and does not reflect the

Ariamnes genetic structure. The core lineages at the generic level reflect typical gut microbiota of

arthropods (Engel & Moran 2013). However, slight but significant differences in the gut microbiota

are detectable between the different spider species and between the *A. waikula* populations within

Hawai'i Island.

We would expect transmission of the gut microbes to generate a strong pattern of

phylosymbiosis and potentially evidence of bottlenecks in the microbial diversity of the youngest

populations (Minard et al. 2015). Instead, we find a strong similarity in terms of microbial diversity

across the different islands and populations, decoupled from the spider's phylogeny. This absence

of a correlation with the host's phylogenetic structure suggests that the gut microbiota is likely

acquired from the spider's environment, most probably through the spider's diet (Kennedy et al.

2020; Zhang et al. 2018). The pattern of slight differentiation according to the host populations

could either be due to gradual changes in the host-filtering process (Mazel et al. 2018), or only

reflect the local variations of the bacterial pool available in the host environment.

The overall similarity of the gut microbiota in terms of the identity and diversity of genera

represented, has been shown in other systems, such as recently diversified *Anolis* lizards (Ren et

al. 2016) and *Cephalotes* turtle ants (Sanders et al. 2014). Such similarity likely conserves the

functional properties of the gut microbiota (Muegge et al. 2011), which could be particularly

advantageous for arthropods which often rely on gut symbionts to complement imbalanced

nutrition (Engel & Moran 2013) or detoxify toxin-rich diets (Adams et al. 2013). This gut microbiota

conservatism of *A. waikula* may reflect its niche conservatism, despite changes in geographical

range.

Conclusion

Host organisms can harbor a large microbial diversity, and it is likely that the different components of the microbial community experience different ecological and evolutionary dynamics. In this work, we demonstrated a dichotomy between the endosymbiont communities and the gut microbiota: For the host Hawaiian *Ariamnes* spiders, abiotic factors, in particular isolation by distance along the volcano chronosequence, appear to have the strongest influence in explaining divergence patterns. The phylogeny of the host strongly impacts the evolution of the transmitted endosymbionts, which show clear shifts in their composition between species on different islands. However, we find little differentiation in endosymbionts across populations of *A. waikula* within Hawaʻi Island, despite the strong genetic structure of the host. In contrast to the differences in the endosymbionts between taxa, the composition of the gut microbiota is similar across both species and populations despite strong geographic isolation of their hosts. Our results stress the high heterogeneity within the different components of the microbiota in terms of host-associated evolution and calls for further investigations of the role on these heterogenous microbial symbionts on promoting host diversification. Specifically, the conservation of the microbiota across ecomorph and island in concert with broader sampling of both *Ariamnes* and other arthropods would significantly contribute to our understanding of the dynamics of co-evolution of microbe and host as it pertains to rapid speciation events.

Acknowledgements

We would like to acknowledge support and assistance from the following: The permit processing and access to different reserves and private land was possible thanks to Steve Bergfeld (DOFAW Big Island), Pat Bily (TNC Maui), Tabetha Block (HETF), Pomaikaʻi Kaniaupio-Crozier (Maui Land and Pineapple), Lance DaSilva (DOFAW Maui), Charmian Dang (NAR), Melissa Dean (HETF), Betsy Gagne (NAR), Lisa Hadway (DOFAW Big Island), Cynthia King (DLNR), Russell Kallstrom

(TNC Molokaʻi), Joey Mello (DOFAW Big Island), Ed Misaki (TNC Molokaʻi). For support and

advice in the lab, we are very grateful to Lydia Smith (Evolutionary Genetics Lab, Museum of

Vertebrate Zoology, UC Berkeley), Shana McDevitt (Vincent J. Coates Genomics Sequencing

Laboratory, QB3, UC Berkeley), and Anna Sellas (California Academy of Sciences, San

Francisco). Samples were provided by Susan Kennedy and Andrew Rominger. We thank

Benjamin Peter, Thorfinn S. Korneliussen, and Line Skotte for advice on the analyses. L.K.B was

supported by the Netherlands Organisation for Scientific Research VENI #863.14.020. B.P.L was

supported by a master fellowship from the École Normale Supérieure of Paris. H.K. was supported

by a postdoctoral fellowship by the German Research Foundation (DFG). Support for the project

was provided by the NSF DEB 1241253 to R.G.G.

Data Accessibility

The pipelines used for processing ddRAD seq data are available in https://github.com/CGRL-QB3-

UCBerkeley/RAD. All scripts used to analyze the Ariamnes microbiota are available in

https://github.com/BPerezLamarque/Scripts. The raw data can be found on dryad

(https://datadryad.org) DOI: https://doi.org/10.5061/dryad.nzs7h44qj

Author Contributions

R.G.G and H.K. conceived of the study. E.A. and B.P-L. performed DNA extractions. C.C.

and L.K.B. performed ddRAD lab work. E.A., K.B., and T.L. performed ddRAD genetic analyses.

B.P-L. and H.K. performed lab work and analyses for the microbial component of the study. E.A.,

B.P-L., H.K., and R.G.G wrote the manuscript. All authors edited and approved the manuscript

before final submission.

References

Adams AS, Aylward FO, Adams SM*, et al.* (2013) Mountain pine beetles colonizing historical and naive host trees are associated with a bacterial community highly enriched in genes contributing to terpene metabolism. *Appl. Environ. Microbiol.* **79**, 3468-3475.

Bailly-Bechet M, Martins-Simões P, Szöllősi GJ*, et al.* (2017) How long does Wolbachia remain on board? *Molecular Biology and Evolution* **34**, 1183-1193.

Bi K, Linderoth T, Vanderpool D*, et al.* (2013) Unlocking the vault: next–generation museum population genomics. *Molecular ecology* **22**, 6018-6032.

Blankers T, Oh KP, Bombarely A, Shaw KL (2018) The genomic architecture of a rapid island radiation: recombination rate variation, chromosome structure, and genome assembly of the Hawaiian cricket Laupala. *Genetics* **209**, 1329-1344.

Bolger AM, Lohse M, Usadel B (2014) Trimmomatic: a flexible trimmer for Illumina sequence data. *Bioinformatics* **30**, 2114-2120.

Brooks AW, Kohl KD, Brucker RM, van Opstal EJ, Bordenstein SR (2016) Phylosymbiosis: relationships and functional effects of microbial communities across host evolutionary history. *PLoS biology* **14**, e2000225.

Brucker RM, Bordenstein SR (2012) Speciation by symbiosis. *Trends in ecology & evolution* **27**, 443-451.

Caporaso JG, Kuczynski J, Stombaugh J*, et al.* (2010) QIIME allows analysis of high-throughput community sequencing data. *Nature methods* **7**, 335.

Carson, H.L., Lockwood, J.P. and Craddock, E.M., 1990. Extinction and recolonization of local populations on a growing shield volcano. *Proceedings of the National Academy of Sciences*, *87*(18), pp.7055-7057.

Carson H, Clague D (1995) Geology and biogeography of the Hawaiian Islands. In Wagner, WL & Funk, VA (eds). Hawaiian Biogeography, Evolution on a Hot Spot Archipelago, Smithsonian Institution Press, Washington: pp. 14–29.

Catchen J, Hohenlohe PA, Bassham S, Amores A, Cresko WA (2013) Stacks: an analysis tool set for population genomics. *Molecular ecology* **22**, 3124-3140.

Crawford JE, Lazzaro BP (2012) Assessing the accuracy and power of population genetic inference from low-pass next-generation sequencing data. *Frontiers in genetics* **3**.

Croucher PJ, Oxford GS, Lam A, Mody N, Gillespie RG (2012) Colonization history and population genetics of the color–polymorphic Hawaiian happy–face spider Theridion grallator (Araneae, Theridiidae). *Evolution: International Journal of Organic Evolution* **66**, 2815-2833.

Degnan PH, Lazarus AB, Brock CD, Wernegreen JJ (2004) Host–symbiont stability and fast evolutionary rates in an ant–bacterium association: cospeciation of Camponotus species and their endosymbionts, Candidatus Blochmannia. *Systematic biology* **53**, 95-110.

Duron O, Binetruy F, Noël V*, et al.* (2017) Evolutionary changes in symbiont community structure in ticks. *Molecular ecology* **26**, 2905-2921.

Duron O, Bouchon D, Boutin S*, et al.* (2008) The diversity of reproductive parasites among arthropods: Wolbachiado not walk alone. *BMC biology* **6**, 27.

Edgar RC (2013) UPARSE: highly accurate OTU sequences from microbial amplicon reads. *Nature methods* **10**, 996.

Eldon J, Bellinger MR, Price DK (2019) Hawaiian picture–winged Drosophila exhibit adaptive population divergence along a narrow climatic gradient on Hawai'i Island. *Ecology and evolution* **9**, 2436.

Engel P, Moran NA (2013) The gut microbiota of insects–diversity in structure and function. *FEMS microbiology reviews* **37**, 699-735.

Fu L, Niu B, Zhu Z, Wu S, Li W (2012) CD-HIT: accelerated for clustering the next-generation
sequencing data. *Bioinformatics* **28**, 3150-3152.
Fumagalli M, Vieira FG, Linderoth T, Nielsen R (2014) ngsTools: methods for population genetics
analyses from next-generation sequencing data. *Bioinformatics* **30**, 1486-1487.
Gibson J, Shokralla S, Porter TM*, et al.* (2014) Simultaneous assessment of the macrobiome and
microbiome in a bulk sample of tropical arthropods through DNA metasystematics.
Proceedings of the National Academy of Sciences **111**, 8007-8012.
Gillespie RG, Benjamin SP, Brewer MS, Rivera MAJ, Roderick GK (2018) Repeated
Diversification Of Ecomorphs In Hawaiian Stick Spiders. *Current Biology* **28**, 941-947.
Gillespie RG, Rivera MA (2007) Free-living spiders of the genus Argyrodes (Araneae: Theridiidae)
in Hawai'i. *Journal of Arachnology* **35**, 11-37.
Goodacre SL, Martin OY, Bonte D*, et al.* (2009) Microbial modification of host long-distance
dispersal capacity. *Bmc Biology* **7**, 32.
Goodacre SL, Martin OY, Thomas CG, Hewitt GM (2006) Wolbachia and other endosymbiont
infections in spiders. *Molecular Ecology* **15**, 517-527.
Goodman KR, Prost S, Bi K, Brewer MS, Gillespie RG (2019) Host and geography together drive
early adaptive radiation of Hawaiian planthoppers. *Molecular ecology* **28**, 4513-4528.
Groussin M, Mazel F, Sanders JG, Smillie CS, Lavergne S, Thuiller W, & Alm EJ (2017).
Unraveling the processes shaping mammalian gut microbiomes over evolutionary time.
Nature Communications, 8(1), 163–185. doi:10.1038/ncomms14319
Hammer TJ, Dickerson JC, Fierer N (2015). Evidence-based recommendations on storing and
handling specimens for analyses of insect microbiota. *PeerJ*, *3*, e1190.
doi:10.7717/peerj.1190
Hammer TJ, Janzen DH, Hallwachs W, Jaffe SP, Fierer N (2017) Caterpillars lack a resident gut
microbiome. *Proceedings of the National Academy of Sciences* **114**, 9641-9646.
Hansen AK, Moran NA (2014) The impact of microbial symbionts on host plant utilization by
herbivorous insects. *Molecular ecology* **23**, 1473-1496.
Hiller AE, Koo MS, Goodman KR*, et al.* (2019) Niche conservatism predominates in adaptive
radiation: comparing the diversification of Hawaiian arthropods using ecological niche
modelling. *Biological Journal of the Linnean Society* **127**, 479-492.
Hoang DT, Chernomor O, Von Haeseler A, Minh BQ, Vinh LS (2018) UFBoot2: improving the
ultrafast bootstrap approximation. *Molecular biology and evolution* **35**, 518-522.
Hoy MA, Jeyaprakash A (2005) Microbial diversity in the predatory mite Metaseiulus occidentalis
(Acari: Phytoseiidae) and its prey, Tetranychus urticae (Acari: Tetranychidae). *Biological
Control* **32**, 427-441.
Huson, D. H., & Bryant, D. (2006). Application of phylogenetic networks in evolutionary
studies. *Molecular biology and evolution*, *23*(2), 254-267.
Kalyaanamoorthy S, Minh BQ, Wong TK, von Haeseler A, Jermiin LS (2017) ModelFinder: fast
model selection for accurate phylogenetic estimates. *Nature methods* **14**, 587.
Kennedy SR, Dawson TE, Gillespie RG (2018) Stable isotopes of Hawaiian spiders reflect
substrate properties along a chronosequence. *PeerJ* **6**, e4527.
Kennedy SR, Tsau S, Gillespie R, Krehenwinkel H (2020) Are you what you eat? A highly transient
and prey–influenced gut microbiome in the grey house spider Badumna longinqua.
Molecular Ecology.
Koga R, Tsuchida T, Fukatsu T (2003) Changing partners in an obligate symbiosis: a facultative
endosymbiont can compensate for loss of the essential endosymbiont Buchnera in an
aphid. *Proceedings of the Royal Society of London. Series B: Biological Sciences* **270**,
2543-2550.
Korneliussen TS, Albrechtsen A, Nielsen R (2014) ANGSD: analysis of next generation
sequencing data. *BMC bioinformatics* **15**, 356.

Lange V, Böhme I, Hofmann J*, et al.* (2014) Cost-efficient high-throughput HLA typing by MiSeq
amplicon sequencing. *BMC genomics* **15**, 63.
Lefort, V., Desper, R., & Gascuel, O. (2015). FastME 2.0: a comprehensive, accurate, and fast
distance-based phylogeny inference program. *Molecular biology and evolution*, **32**(10),
2798-2800.
Li H, Handsaker B, Wysoker A*, et al.* (2009) The sequence alignment/map format and SAMtools.
Bioinformatics **25**, 2078-2079.
Li W, Godzik A (2006) Cd-hit: a fast program for clustering and comparing large sets of protein or
nucleotide sequences. *Bioinformatics* **22**, 1658-1659.
Maas, D. L., Prost, S., Bi, K., Smith, L. L., Armstrong, E. E., Aji, L. P., Toha, A.H.A., Gillespie,
R.G., & Becking, L. E. (2018). Rapid divergence of mussel populations despite incomplete
barriers to dispersal. *Molecular ecology*, **27**(7), 1556-1571.
Magoč T, Salzberg SL (2011) FLASH: fast length adjustment of short reads to improve genome
assemblies. *Bioinformatics* **27**, 2957-2963.
Margulis L, Fester R (1991) *Symbiosis as a source of evolutionary innovation: speciation and*
morphogenesis Mit Press.
Martin M (2011) Cutadapt removes adapter sequences from high-throughput sequencing reads.
EMBnet. journal **17**, pp. 10-12.
Mazel F, Davis KM, Loudon A*, et al.* (2018) Is host filtering the main driver of phylosymbiosis
across the tree of life? *mSystems* **3**, e00097-00018.
McFall-Ngai M, Hadfield MG, Bosch TC*, et al.* (2013) Animals in a bacterial world, a new
imperative for the life sciences. *Proceedings of the National Academy of Sciences* **110**,
3229-3236.
McKenna A, Hanna M, Banks E*, et al.* (2010) The Genome Analysis Toolkit: a MapReduce
framework for analyzing next-generation DNA sequencing data. *Genome research* **20**,
1297-1303.
Minard G, TRAN F-H, Goubert C*, et al.* (2015) French invasive Asian tiger mosquito populations
harbor reduced bacterial microbiota and genetic diversity compared to Vietnamese
autochthonous relatives. *Frontiers in Microbiology* **6**, 970.
Moeller AH, Caro-Quintero A, Mjungu D*, et al.* (2016) Cospeciation of gut microbiota with
hominids. *Science* **353**, 380-382.
Moran NA, Sloan DB (2015) The hologenome concept: helpful or hollow? *PLoS biology* **13**.
Muegge BD, Kuczynski J, Knights D*, et al.* (2011) Diet drives convergence in gut microbiome
functions across mammalian phylogeny and within humans. *Science* **332**, 970-974.
Nguyen L-T, Schmidt HA, Von Haeseler A, Minh BQ (2015) IQ-TREE: a fast and effective
stochastic algorithm for estimating maximum-likelihood phylogenies. *Molecular biology*
and evolution **32**, 268-274.
O'Connor TK, Humphrey PT, Lapoint RT, Whiteman NK, O'Grady PM (2014) Microbial
interactions and the ecology and evolution of Hawaiian Drosophilidae. *Frontiers in*
microbiology **5**, 616.
Oksanen J, Kindt R, Pierre L, O'Hara B, Simpson GL, Solymos P, Stevens MH. HH, Wagner H,
Blanchet FG, Kindt R, et al. 2016. vegan: Community Ecology Package, R package version
2.4-0. R package version 2.2-1.
Papadopulos AS, Kaye M, Devaux C*, et al.* (2014) Evaluation of genetic isolation within an island
flora reveals unusually widespread local adaptation and supports sympatric speciation.
Philosophical Transactions of the Royal Society B: Biological Sciences **369**, 20130342.
Paradis, E., Claude, J., & Strimmer, K. (2004). APE: Analyses of phylogenetics and evolution in
R language. *Bioinformatics*, *20*(2), 289–290. doi:10.1093/bioinformatics/btg412
826

Pekár S, Šobotník J (2007) Comparative study of the femoral organ in Zodarion spiders (Araneae:
Zodariidae). *Arthropod structure & development* **36**, 105-112.
Pekár S, Šobotník J (2008) Erratum to "Comparative study of the femoral organ in Zodarion
spiders (Araneae: Zodariidae)"[Arthropod Structure & Development 36 (2)(2007) 105–
112]. *Arthropod Structure & Development* **37**, 93-94.
Perez-Lamarque B, Morlon H (2019). Characterizing symbiont inheritance during host–microbiota
evolution: Application to the great apes gut microbiota. Molecular Ecology Resources, 1–
14. doi:10.1111/1755-0998.13063
Petkova, D., Novembre, J., & Stephens, M. (2016). Visualizing spatial population structure with
estimated effective migration surfaces. *Nature genetics*, *48*(1), 94-100.
Pylro VS, Roesch LFW, Morais DK*, et al.* (2014) Data analysis for 16S microbial profiling from
different benchtop sequencing platforms. *Journal of microbiological methods* **107**, 30-37.
Ren T, Kahrl AF, Wu M, Cox RM (2016) Does adaptive radiation of a host lineage promote
ecological diversity of its bacterial communities? A test using gut microbiota of Anolis
lizards. *Molecular ecology* **25**, 4793-4804.
Rennison DJ, Rudman SM, Schluter D (2019) Parallel changes in gut microbiome composition
and function during colonization, local adaptation and ecological speciation. *Proceedings*
of the Royal Society B **286**, 20191911.
Roderick GK, Croucher PJ, Vandergast AG, Gillespie RG (2012) Species differentiation on a
dynamic landscape: shifts in metapopulation genetic structure using the chronology of the
Hawaiian archipelago. *Evolutionary Biology* **39**, 192-206.
Sanders JG, Powell S, Kronauer DJ*, et al.* (2014) Stability and phylogenetic correlation in gut
microbiota: lessons from ants and apes. *Molecular Ecology* **23**, 1268-1283.
Sanderson MJ (2003) r8s: inferring absolute rates of molecular evolution and divergence times in
the absence of a molecular clock. *Bioinformatics* **19**, 301-302.
Schrader C, Schielke A, Ellerbroek L, Johne R (2012) PCR inhibitors–occurrence, properties and
removal. *Journal of applied microbiology* **113**, 1014-1026.
Sharon G, Segal D, Ringo JM*, et al.* (2010) Commensal bacteria play a role in mating preference
of Drosophila melanogaster. *Proceedings of the National Academy of Sciences* **107**,
20051-20056.
Shaw KL, Gillespie RG (2016) Comparative phylogeography of oceanic archipelagos: Hotspots
for inferences of evolutionary process. *Proceedings of the National Academy of Sciences*
113, 7986-7993.
Sheffer MM, Uhl G, Prost S, Lueders T, Urich T, Bengtsson MM (2019). Tissue- and Population-
Level Microbiome Analysis of the Wasp Spider Argiope bruennichi Identified a Novel Dominant
Bacterial Symbiont. Microorganisms, 8(1), 8. doi:10.3390/microorganisms8010008
Sherrod DR, Sinton JM, Watkins SE, Brunt KM (2007) Geologic map of the State of Hawai'i. *US*
geological survey open-file report **1089**.
Shropshire JD, Bordenstein SR (2016) Speciation by symbiosis: the microbiome and behavior.
MBio **7**, e01785-01715.
Skotte L, Korneliussen TS, Albrechtsen A (2013) Estimating individual admixture proportions from
next generation sequencing data. *Genetics* **195**, 693-702.
Smit A, Hubley R, Green P (2004) RepeatMasker Open-3.0. 1996–2004. *Institute for Systems*
Biology.
Stamatakis, A. (2014). RAxML version 8: a tool for phylogenetic analysis and post-analysis of
large phylogenies. *Bioinformatics*, *30*(9), 1312-1313.
Su Q, Hu G, Yun Y, Peng Y (2019) Horizontal transmission of Wolbachia in Hylyphantes
graminicola is more likely via intraspecies than interspecies transfer. *Symbiosis* **79**, 123-
128.

Team RC (2018) R Foundation for Statistical Computing; Vienna, Austria: 2015. *R: A language*
and environment for statistical computing, 2013.
Vanthournout B, Hendrickx F (2015) Endosymbiont dominated bacterial communities in a dwarf
spider. *PLoS One* **10**.
Vieira FG, Lassalle F, Korneliussen TS, Fumagalli M (2015) Improving the estimation of genetic
distances from Next-Generation Sequencing data. Biological Journal of the Linnean
Society **117**(1):139-149
Wang IJ (2013) Examining the full effects of landscape heterogeneity on spatial genetic variation:
a multiple matrix regression approach for quantifying geographic and ecological isolation.
Evolution **67**, 3403-3411.
Weber MG, Wagner CE, Best RJ, Harmon LJ, Matthews B (2017) Evolution in a community
context: on integrating ecological interactions and macroevolution. *Trends in ecology &*
evolution **32**, 291-304.
White JA, Styer A, Rosenwald LC*, et al.* (2020) Endosymbiotic bacteria are prevalent and diverse
in agricultural spiders. *Microbial ecology* **79**, 472-481.
Whitehouse M, Agnarsson I, Miyashita T*, et al.* (2002) *Argyrodes*: phylogeny, sociality and
interspecific interactions—a report on the *Argyrodes* symposium, Badplaas 2001. *Journal*
of Arachnology **30**, 238-245.
Yun J-H, Roh SW, Whon TW*, et al.* (2014) Insect gut bacterial diversity determined by
environmental habitat, diet, developmental stage, and phylogeny of host. *Appl. Environ.*
Microbiol. **80**, 5254-5264.
Zhang L, Yun Y, Hu G, Peng Y (2018) Insights into the bacterial symbiont diversity in spiders.
Ecology and evolution **8**, 4899-4906.
Zhang L, Zhang G, Yun Y, Peng Y (2017) Bacterial community of a spider, Marpiss magister
(Salticidae). *3 Biotech* **7**, 371.

Supplementary Figures

Figure S1: **Grid system used for the EEMS analyses on Hawaii Island (left) and**

corresponding results (right)

A: Alili, K: Kohala, O: Olaa, P: Puʻu Makaʻala, S: Saddle, T: Thurston

Figure S2. Traces of the MCMC runs of the 10 replicates to assess convergence.

Only individuals of the gold ecomorph from Hawaii Island were used here.

Figure S3: Rarefaction curve of the whole microbiota: number of observed OTU in each sample according to the number of rarefied reads per sample. Results of OTU at 97% are presented on the left, whereas Z-OTU are on the right. Rarefactions are performed 20 times independently and the mean value is plotted here. Note that one sample (from Thurston) present an unexpectedly high diversity compared to other samples.

Figure S4: NgsAdmix convergence likelihoods for 50 iterations K values 2-6 for *A. waikula* from

Hawai'i Island.

926 .

Figure S5: PCA analyses of ddRAD data of *A. waikula* from Hawai'i Island

933   **Figure S6**: Site frequency spectrum of all *Ariamnes* ddRAD data after filtering.

936   **Figure S7:** Relative abundances of endosymbiont genera per population (A) and gut symbionts

937   per spider population (B) defined from rarefied OTU table at 97%. Rare genera (representing less

938   than 1% of the abundance are merged together).

Figure S8: Relative abundances of orders: endosymbiont orders per population (A-B) and gut symbionts per spider population (C-D) defined from rarefied OTU table. Rare orders (representing less than 1% of the abundance are merged together). A and C (resp. B and D) are from 97% OTU (resp. Z-OTU).

949 **Figure S9:** PCoA decomposition of the microbial communities according to the type of OTU

950 clustering (OTU at 97% or Z-OTU) and the type of microbial communities (whole microbiota,

951 endosymbionts, or gut microbiota).

Figure S10: Microbiota dendrograms reconstructed from the gut microbiota (A) or the endosymbiont community (B) for the OTU at 97%.

953
954

Supplementary Tables

957   **Table S1:** Specimen collection detail for population genetic and microbial samples.

Species	Eco-morph	Island	Population	Volcano (age)	Substrate age (years)	Latitude	Longitude	Collector	Date	Individuals (ddRAD)	Microbiota
A. waikula	Gold	Hawai'i	Kohala	Kohala, 0.43	300,000	20.1048	-155.7242	Rominger	2014-01-12	18	6
A. waikula	Gold	Hawai'i	Saddle	Mauna Loa, Kea 0.01-0.38	4,000	19.6746	-155.3316	Rominger	2014-01-07	6	3
A. waikula	Gold	Hawai'i	Puu Makaala	Kilauea, 0.004	11,000	19.3748	-155.2202	Rominger	2014-01-05	18	13
A. waikula	Gold	Hawai'i	Thurston	Kilauea, 0.004	600	19.4078	-155.2388	Rominger	2014-01-08	13	9
A. waikula	Gold	Hawai'i	Ola'a	Kilauea, 0.004	7,500	19.4530	-155.2466	Rominger	2014-01-09	16	11
A. waikula	Gold	Hawai'i	Alili	Mauna Loa, 0.01	20,000	19.2309	-155.5134	Rominger	2014-01-04	18	13
A. hiwa	Brown	Hawai'i	Puu Makaala	Kilauea, 0.004	11,000	19.3748	-155.2202	Rominger	2014-01-05	1	N/A
A. hiwa	Brown	Hawai'i	Thurston	Kilauea, 0.004	600	19.4078	-155.2388	Rominger	2014-01-08	1	N/A
A. melekalikimaka	Gold	W. Maui	Puu Kukui	Puu Kukui, 1.3	1,500,000	20.9194	-156.5972	Gillespie	2012-06-29	16	9
A. n. sp	Gold	Moloka'i	Kamakou	Moloka'i, 1.8	1,400,000	21.1108	-156.8906	Gillespie	2012-06-23	16	7

960 **Table S2:** Specimen collection detail for population genetic and microbial samples.

Island	Population	Name ddRAD	Name microbiota	SeqID	Pool	Sample ID
Hawai'i	Pu'u Maka'ala	1	83	RGCC001A_S25_AACCA	A2	AJR923
Hawai'i	Pu'u Maka'ala	2	NA	RGCC001A_S25_AAGGA	A8	AJR913
Hawai'i	Pu'u Maka'ala	3	82	RGCC001A_S25_ACACA	A10	AJR948
Hawai'i	Pu'u Maka'ala	4	NA	RGCC001A_S25_ACTGG	A13	AJR956
Hawai'i	Pu'u Maka'ala	5	70	RGCC001A_S25_ACGGT	A12	AJR955
Hawai'i	Pu'u Maka'ala	6	65	RGCC001A_S25_ACTTC	A14	AJR964
Hawai'i	Pu'u Maka'ala	7	47	RGCC001A_S25_AGCTA	A9	AJR915
Hawai'i	Pu'u Maka'ala	8	NA	RGCC001A_S25_ATACG	A15	AJR965
Hawai'i	Pu'u Maka'ala	9	59	RGCC001A_S25_ATGAG	A16	AJR1017
Hawai'i	Pu'u Maka'ala	10	NA	RGCC001A_S25_CAACC	A6	AJR950
Hawai'i	Pu'u Maka'ala	11	85	RGCC001A_S25_CGATC	A3	AJR924
Hawai'i	Pu'u Maka'ala	12	80	RGCC001A_S25_CTGCG	A17	AJR1033
Hawai'i	Pu'u Maka'ala	13	60	RGCC001A_S25_CTGTC	A18	AJR1034
Hawai'i	Pu'u Maka'ala	14	52	RGCC001A_S25_CTTGG	A19	AJR1036
Hawai'i	Pu'u Maka'ala	15	74	RGCC001A_S25_GACAC	A20	AJR1037
Hawai'i	Pu'u Maka'ala	16	NA	RGCC001A_S25_GCATG	A1	AJR922
Hawai'i	Pu'u Maka'ala	17	NA	RGCC001A_S25_GGTTG	A7	AJR951
Hawai'i	Pu'u Maka'ala	18	NA	RGCC001A_S25_TCGAT	A4	AJR946
Hawai'i	Pu'u Maka'ala	19	61	RGCC001A_S25_TGCAT	A5	AJR947
Hawai'i	Alili	20	66	RGCC001B_S26_AACCA	B2	AJR839
Hawai'i	Alili	21	62	RGCC001B_S26_AAGGA	B8	AJR850
Hawai'i	Alili	22	88	RGCC001B_S26_AATTA	B11	AJR855
Hawai'i	Alili	23	100	RGCC001B_S26_ACACA	B10	AJR854
Hawai'i	Alili	24	45	RGCC001B_S26_ACGGT	B12	AJR857
Hawai'i	Alili	25	NA	RGCC001B_S26_ACTTC	B14	AJR859
Hawai'i	Alili	26	78	RGCC001B_S26_AGCTA	B9	AJR852
Hawai'i	Alili	27	118	RGCC001B_S26_ATACG	B15	AJR860
Hawai'i	Alili	28	104	RGCC001B_S26_ATGAG	B16	AJR861
Hawai'i	Alili	29	98	RGCC001B_S26_CAACC	B6	AJR848
Hawai'i	Alili	30	NA	RGCC001B_S26_CGATC	B3	AJR844
Hawai'i	Alili	31	107	RGCC001B_S26_CTGCG	B17	AJR891
Hawai'i	Alili	32	NA	RGCC001B_S26_CTGTC	B18	AJR893

Hawai'i	Alili	33	NA	RGCC001B_S26_GACAC	B20	AJR897
Hawai'i	Alili	34	NA	RGCC001B_S26_GCATG	B1	AJR838
Hawai'i	Alili	35	102	RGCC001B_S26_GGTTG	B7	AJR849
Hawai'i	Alili	36	NA	RGCC001B_S26_TCGAT	B4	AJR846
Hawai'i	Alili	37	NA	RGCC001B_S26_TGCAT	B5	AJR847
Hawai'i	Kohala	38	NA	RGCC001C_S27_AACCA	C2	AJR1413
Hawai'i	Kohala	39	55	RGCC001C_S27_AAGGA	C8	AJR1425
Hawai'i	Kohala	40	120	RGCC001C_S27_AATTA	C11	AJR1428
Hawai'i	Kohala	41	99	RGCC001C_S27_ACACA	C10	AJR1427
Hawai'i	Kohala	42	84	RGCC001C_S27_ACTGG	C13	AJR1430
Hawai'i	Kohala	43	NA	RGCC001C_S27_ACTTC	C14	AJR1476
Hawai'i	Kohala	44	NA	RGCC001C_S27_AGCTA	C9	AJR1426
Hawai'i	Kohala	45	NA	RGCC001C_S27_ATACG	C15	AJR1477
Hawai'i	Kohala	46	NA	RGCC001C_S27_ATGAG	C16	AJR1478
Hawai'i	Kohala	47	NA	RGCC001C_S27_CAACC	C6	AJR1417
Hawai'i	Kohala	48	NA	RGCC001C_S27_CGATC	C3	AJR1414
Hawai'i	Kohala	49	NA	RGCC001C_S27_CTGCG	C17	AJR1518
Hawai'i	Kohala	50	NA	RGCC001C_S27_CTGTC	C18	AJR1519
Hawai'i	Kohala	51	NA	RGCC001C_S27_CTTGG	C19	AJR1520
Hawai'i	Kohala	52	NA	RGCC001C_S27_GCATG	C1	AJR1412
Hawai'i	Kohala	53	NA	RGCC001C_S27_GGTTG	C7	AJR1418
Hawai'i	Kohala	54	117	RGCC001C_S27_TCGAT	C4	AJR1415
Hawai'i	Kohala	55	NA	RGCC001C_S27_TGCAT	C5	AJR1416
Hawai'i	Ola'a	56	NA	RGCC001D_S28_AACCA	D2	AJR1151
Hawai'i	Ola'a	57	NA	RGCC001D_S28_AAGGA	D8	AJR1158
Hawai'i	Ola'a	58	67	RGCC001D_S28_AATTA	D11	AJR1161
Hawai'i	Ola'a	59	64	RGCC001D_S28_ACACA	D10	AJR1160
Hawai'i	Ola'a	60	81	RGCC001D_S28_ACTGG	D13	AJR1187
Hawai'i	Ola'a	61	49	RGCC001D_S28_ACTTC	D14	AJR1188
Hawai'i	Ola'a	62	58	RGCC001D_S28_AGCTA	D9	AJR1159
Hawai'i	Ola'a	63	48	RGCC001D_S28_ATACG	D15	AJR1190
Hawai'i	Ola'a	64	72	RGCC001D_S28_CAACC	D6	AJR1156
Hawai'i	Ola'a	65	53	RGCC001D_S28_CGATC	D3	AJR1152
Hawai'i	Ola'a	66	119	RGCC001D_S28_CTGCG	D17	AJR1192

Hawai'i	Ola'a	67	NA	RGCC001D_S28_CTGTC	D18	AJR1232
Hawai'i	Ola'a	68	NA	RGCC001D_S28_CTTGG	D19	AJR1233
Hawai'i	Ola'a	69	NA	RGCC001D_S28_GCATG	D1	AJR1150
Hawai'i	Ola'a	70	NA	RGCC001D_S28_GGTTG	D7	AJR1157
Hawai'i	Ola'a	71	NA	RGCC001D_S28_TCGAT	D4	AJR1153
Hawai'i	Ola'a	72	109	RGCC001D_S28_TGCAT	D5	AJR1155
Hawai'i	Thurston	105	NA	RGCC001H_S31_AACCA	H2	AJR1061
Hawai'i	Thurston	106	44	RGCC001H_S31_AAGGA	H8	AJR1067
Hawai'i	Thurston	107	111	RGCC001H_S31_AATTA	H11	AJR1107
Hawai'i	Thurston	108	87	RGCC001H_S31_ACACA	H10	AJR1106
Hawai'i	Thurston	109	NA	RGCC001H_S31_ACGGT	H12	AJR1108
Hawai'i	Thurston	110	NA	RGCC001H_S31_ACTGG	H13	AJR1109
Hawai'i	Thurston	111	NA	RGCC001H_S31_ACTTC	H14	AJR1110
Hawai'i	Thurston	112	NA	RGCC001H_S31_AGCTA	H9	AJR1068
Hawai'i	Thurston	114	108	RGCC001H_S31_CGATC	H3	AJR1062
Hawai'i	Thurston	115	115	RGCC001H_S31_GCATG	H1	AJR1060
Hawai'i	Thurston	116	79	RGCC001H_S31_GGTTG	H7	AJR1066
Hawai'i	Thurston	117	97	RGCC001H_S31_TGCAT	H5	AJR1064
Hawai'i	Saddle	118	NA	RGCC001I_S32_AACCA	H2	AJRA2
Hawai'i	Saddle	119	NA	RGCC001I_S32_CAACC	H6	AJRA6
Hawai'i	Saddle	120	63	RGCC001I_S32_CGATC	H3	AJRA3
Hawai'i	Saddle	121	103	RGCC001I_S32_GCATG	H1	AJRA1
Hawai'i	Saddle	122	NA	RGCC001I_S32_TCGAT	H4	AJRA4
Hawai'i	Saddle	123	57	RGCC001I_S32_TGCAT	H5	AJRA5
Hawai'i	Thurston	NA	42	RGCC001H_S31_TCGAT	H4	AJR1063
Hawai'i	Kohala	NA	51	RGCC001C_S27_ACGGT	C12	AJR1429
Hawai'i	Thurston	NA	69	RGCC001H_S31_CAACC	H6	AJR1065
Hawai'i	Alili	NA	71	RGCC001B_S26_ACTGG	B13	AJR858
Hawai'i	Alili	NA	77	RGCC001B_S26_CTTGG	B19	AJR894
Hawai'i	Pu'u Maka'ala	NA	101	RGCC001A_S25_AATTA	A11	AJR949
Hawai'i	Ola'a	NA	113	RGCC001D_S28_ACGGT	D12	AJR1183
Hawai'i	Kohala	NA	NA	RGCC001C_S27_GACAC	C20	AJR1521
Hawai'i	Ola'a	NA	NA	RGCC001D_S28_ATGAG	D16	AJR1191
Moloka'i	Moloka'i	73	NA	RGCC001F_S29_AACCA	F2	RGGA10

Moloka'i	Moloka'i	74	NA	RGCC001F_S29_AAGGA	F8	RGG43
Moloka'i	Moloka'i	75	106	RGCC001F_S29_AATTA	F11	RGG46
Moloka'i	Moloka'i	76	86	RGCC001F_S29_ACACA	F10	RGG45
Moloka'i	Moloka'i	77	68	RGCC001F_S29_ACGGT	F12	RGG47
Moloka'i	Moloka'i	78	73	RGCC001F_S29_ACTGG	F13	RGG48
Moloka'i	Moloka'i	79	NA	RGCC001F_S29_ACTTC	F14	RGG5
Moloka'i	Moloka'i	80	114	RGCC001F_S29_AGCTA	F9	RGG44
Moloka'i	Moloka'i	81	NA	RGCC001F_S29_ATACG	F15	RGG6
Moloka'i	Moloka'i	82	NA	RGCC001F_S29_ATGAG	F16	RGG9
Moloka'i	Moloka'i	83	105	RGCC001F_S29_CAACC	F6	RGG41
Moloka'i	Moloka'i	84	NA	RGCC001F_S29_CGATC	F3	RGGA11
Moloka'i	Moloka'i	85	NA	RGCC001F_S29_GCATG	F1	RGGA9
Moloka'i	Moloka'i	86	112	RGCC001F_S29_GGTTG	F7	RGG42
Moloka'i	Moloka'i	87	NA	RGCC001F_S29_TCGAT	F4	RGG2
Moloka'i	Moloka'i	88	NA	RGCC001F_S29_TGCAT	F5	RGG3
Maui	W_Maui	89	56	RGCC001G_S30_AACCA	G2	RGG121
Maui	W_Maui	90	NA	RGCC001G_S30_AAGGA	G8	RGG135
Maui	W_Maui	91	43	RGCC001G_S30_AATTA	G11	RGG138
Maui	W_Maui	92	50	RGCC001G_S30_ACACA	G10	RGG137
Maui	W_Maui	93	116	RGCC001G_S30_ACGGT	G12	RGG151
Maui	W_Maui	94	NA	RGCC001G_S30_ACTGG	G13	RGG152
Maui	W_Maui	95	54	RGCC001G_S30_ACTTC	G14	RGG153
Maui	W_Maui	96	46	RGCC001G_S30_AGCTA	G9	RGG136
Maui	W_Maui	97	76	RGCC001G_S30_ATACG	G15	RGG154
Maui	W_Maui	98	41	RGCC001G_S30_ATGAG	G16	RGG155
Maui	W_Maui	99	NA	RGCC001G_S30_CAACC	G6	RGG133
Maui	W_Maui	100	110	RGCC001G_S30_CGATC	G3	RGG122
Maui	W_Maui	101	NA	RGCC001G_S30_GCATG	G1	RGGA2
Maui	W_Maui	102	NA	RGCC001G_S30_GGTTG	G7	RGG134
Maui	W_Maui	103	NA	RGCC001G_S30_TCGAT	G4	RGG131
Maui	W_Maui	104	NA	RGCC001G_S30_TGCAT	G5	RGG132
Hawai'i	Ola'a	NA	NA	RGCC001D_S28_GACAC	D20	AJR1186
Hawai'i	Thurston	113	NA	RGCC001H_S31_ATACG	H15	AJR1111

962   **Table S3**: F$_{ST}$ values for ddRAD data from *Ariamnes* spiders.

963
964
965

	Alili	Kohala	Olaʻa	Thurston	Saddle	Molokaʻi	Maui
Puʻu Makaʻala	0.0650	0.1466	0.0291	0.0461	0.1066	0.4870	0.5454
Alili		0.1394	0.0544	0.0629	0.0978	0.4829	0.5407
Kohala			0.1367	0.1496	0.1315	0.4742	0.5264
Olaʻa				0.0334	0.0937	0.4817	0.5404
Thurston					0.1120	0.4815	0.5468
Saddle						0.4629	0.5410
Molokaʻi							0.4781

966   **Table S4: Average Tajima's D values for each population of *A. waikula* and Maui (*A.***

967   ***melekelikimaka*) and Molokai (*A. n. spp.*).**

Population	Average	Min	Max
Alili	-0.77	-2.68	3.33
Ola'a	-0.95	-2.83	3.84
Pu'u Maka'ala	-1.00	-2.77	3.25
Saddle	-0.61	-2.97	3.79
Thurston	-0.74	-2.87	3.73
Kohala	-0.71	-2.82	3.52
Maui	-0.06	-2.63	3.94
Molokai	-0.23	-2.46	3.66

970 **Table S5:** Mean alpha diversity of the microbial communities per host populations. Alpha

971 diversities were computed on rarefied OTU tables defined at 97% or Z-OTUs using the Chaos 1

972 index.

Island	Population	Number individuals	OTU at 97%		Z-OTU	
			Alpha diversity (Chaos 1)	Standard deviation	Alpha diversity (Chaos 1)	Standard deviation
Hawai'i	Alili	13	42.4	7.2	95.4	4.0
Hawai'i	Kohala	6	55.8	12.3	114.8	7.8
Hawai'i	Ola'a	9	41.6	3.0	96.3	8.5
Hawai'i	Pu'u Maka'ala	13	49.5	3.2	108.0	3.4
Hawai'i	Saddle	3	44.8	1.6	103.1	5.2
Hawai'i	Thurston	9	40.8	4.6	103.2	5.7
Maui	Puu Kukui	9	46.0	5.7	106.6	4.4
Moloka'i	Kamakou	7	52.2	6.7	106.6	5.3

974
975
976

Table S6: PERMANOVA testing the of host populations on the dissimilarities between microbial communities according to the host populations. We indicated the adjusted R-squared and the associated p-value for each test. Significant relationships indicate a differentiation of the microbial communities according to the host populations.

Populations	OTU at 97%			Z-OTU		
	Whole microbiota	Gut microbiota	Endo-symbionts	Whole microbiota	Gut microbiota	Endo-symbionts
All populations	0.37905 (0.0001)	0.31109 (0.0001)	0.7146 (0.0001)	0.37709 (0.0001)	0.31739 (0.0001)	0.7115 (0.0001)
Populations within Hawai'i Island	0.21715 (0.0062)	0.29671 (0.0007)	0.28093 (0.0190)	0.21542 (0.0057)	0.29 (0.0003)	0.29072 (0.0120)

984 **Table S7:** Mantel test comparing the microbial beta diversity dis-similarities to the genetic

985 distances or the phylogenetic distances of their associated *Ariamnes*. Beta diversity dissimilarities

986 were computed on rarefied OTU tables defined at 97% using the Bray-Curtis dissimilarities. Bold

987 values represent significant correlations. Mantel tests were either performed on all populations

988 (between *Ariamnes* species and *A. waikula* populations) or only on *A. waikula* populations.

Matrix 1	Matrix 2	All populations		*A. waikula* populations	
		Correlation	p-value	Correlation	p-value
Whole microbiota beta diversity distances	*Ariamnes* genetic distances	**0.312**	**9.9e-05**	0.117	0.066
	Ariamnes phylogenetic distances	**0.323**	**9.9e-05**	0.026	0.321
Endosymbionts beta diversity distances	*Ariamnes* genetic distances	**0.620**	**9.9e-05**	0.011	0.500
	Ariamnes phylogenetic distances	**0.646**	**9.9e-05**	0.061	0.257
Gut microbiota beta diversity distances	*Ariamnes* genetic distances	-0.096	0.837	-0.043	0.668
	Ariamnes phylogenetic distances	-0.095	0.861	-0.130	0.912