Canonical Wnt signaling exerts bidirectional control on choroid plexus epithelial development

Arpan Parichha*[1], Varun Suresh*[1], Mallika Chatterjee[1$], Aditya Kshirsagar[2], Lihi Ben-Reuven[2], Tsviya Olender[2], M. Mark Taketo[3], Velena Radosevic[4,5], Mihaela Bobic-Rasonja[4,5], Sara Trnski[4], Nataša Jovanov Milošević[#4,5], Orly Reiner[#2], Shubha Tole[#1]

[1]Department of Biological Sciences, Tata Institute of Fundamental Research, Mumbai 400005, India; [2]Department of Molecular Genetics, Weizmann Institute of Science, Israel; [3]Division of Experimental Therapeutics Graduate School of Medicine, Kyoto University (Yoshida-Konoé-Cho, Sakyo), Kyoto 606-8501, Japan; [4]Croatian Institute for Brain Research, Department of Medical Biology, School of Medicine University of Zagreb, Šalata 12, Zagreb, Croatia, [5]University Hospital Centre Zagreb, Department of Gynecology and Department of Pathology and Cytology, Petrova 13, Zagreb, Croatia

*These authors contributed equally to this work
#Authors for correspondence: stole@tifr.res.in; orly.reiner@weizmann.ac.il; njovanov@gmail.com
$Current address: Amity Institute of Neuropsychology and Neurosciences, Amity University, Noida, India.

Abstract

The choroid plexus (CP) secretes cerebrospinal fluid and is critical for the development and function of the brain. In the telencephalon, the CP epithelium (CPe) arises from the *Wnt-* and *Bmp-* expressing cortical hem. We examined the role of canonical Wnt signaling in CPe development and report that the mouse and human embryonic CPe expresses molecules in this pathway. Either loss of function or constitutive activation of β-catenin, a key mediator of canonical Wnt signaling, causes a profound disruption of mouse CPe development. Loss of β-catenin results in a dysmorphic CPe, while constitutive activation of β-catenin causes a loss of CPe identity and a transformation of this tissue to a hippocampal-like identity. Aspects of this phenomenon are recapitulated in human embryonic stem cell (hESC)-derived organoids. Our results indicate that canonical Wnt signaling is required in a precisely regulated manner for normal CP development in the mammalian brain.

Introduction

The Choroid plexus (CP) produces cerebrospinal fluid (CSF) which plays an important role in the developing and adult brain. The CSF provides protective functions such as cushioning against mechanical damage; nutritive functions; and also signaling functions, providing molecular cues for the developing brain (Johansson, 2014; Lun, et al., 2015). The CP separates the blood from the CSF, creating a checkpoint for regulating the entry and exit of factors (Zappaterra and Lehtinen, 2012; Lun et al., 2015). The CP is also the gateway for immune cells entering or exiting the brain (Meeker et al., 2012).

The CP epithelium (CPe) arises from adjacent neuroepithelial tissue in the telencephalon, diencephalon, and the hindbrain. In the telencephalon, the non-neuronal CPe as well as a sub-population of Cajal-Retzius (CR) neurons arise from the cortical hem (Yoshida et al., 2006; Louvi et al., 2007; Gu *et al.*, 2011). This midline telencephalic signaling center expresses genes that belong to the BMP and WNT families in mouse (Grove et al., 1998) and human (Abu-Khalil et al., 2004). In the marsupial *Monodelphis Domestica,* BrdU positive cells arise from the hem and populate the growing telencephalic CPe in a "conveyor belt" like manner (Liddelow *et al.*, 2010). Consistent with this model, when the size of the cortical hem increases (Monuki et al., 2001), decreases (Konno *et al.*, 2012) or is missing altogether (Grove et al., 1998), there is a parallel change in the extent of the CP.

The mechanisms that regulate the specification of the CPe are important from a developmental, disease, and evolutionary perspective. Obvious candidates for the control of CPe development are the signaling molecules expressed in the neuroepithelial progenitor domains. The hem expresses several members of the Bmp and Wnt families (Furuta et al., 1997; Grove et al., 1998). Of particular interest are mechanisms that govern the decision between the neuronal (Cajal-Retzius cells) versus non-neuronal (CPe) lineages arising from progenitors in the hem.

Bmp signaling is critical for CPe specification: the CP is missing in the *BmpR1a* null mutant mouse (Hebert *et al.*, 2002; Watanabe *et al.*, 2012), and *Bmp4* is sufficient for CPe induction *in vitro* (Eiraku et al., 2008; Watanabe et al., 2012). The role of hem-derived Wnt ligands in CPe development appears to be equally important, though well explored. Non-canonical Wnt5a, expressed in the hem, appears to be required for maintaining the size and the cytoarchitecture of the CPe (Langford et al., 2020). A role for canonical Wnt signaling in CPe development is indicated by the observation that the addition of WNT3a augments the CPe-inducing ability of BMP4 in three-dimensional aggregation culture derived from embryonic stem cells (ESCs). How the CPe responds to canonical Wnt signaling, and whether this pathway

is required for CPe development, is an open question, and the focus of this study. We

demonstrate that molecules downstream of canonical Wnt signaling are expressed *in vivo* at

embryonic stages in mouse and human CPe. Loss of β-catenin, a key member of the canonical

Wnt pathway transcriptional complex, results in a dysmorphic CP in the mouse, whereas

constitutive activation of β-CATENIN, or treatment with a canonical Wnt agonist of human ESC

(hESC)- derived organoids, results in a loss of CPe identity. Together, our results indicate that a

precise balance of canonical Wnt signaling is critical for CPe specification and morphogenesis,

and that this mechanism for CPe regulation appears to be conserved in humans and rodents.

Results

Canonical Wnt signaling is necessary for normal CP development in mice

We examined whether the molecular machinery required for processing canonical Wnt signaling is present in the developing CPe. The hem and the CPe are identified by the expression of *Wnt3a* and the thyroid hormone T4 carrier transthyretin (*Ttr*), respectively. Several molecules involved in the canonical Wnt signaling pathway, including a receptor, *Frizzled 1* (*Fzd1*), transcriptional regulators β-CATENIN and LEF1, as well as a target gene *Axin2*, are each expressed in the CPe at E12.5 and E14.5 in mice (Figure 1A, B). These data suggest that the embryonic CPe has the capability of processing canonical Wnt signaling. *Lmx1a* is expressed in the hem and the CPe (Chizikov and Millen., 2004). To genetically manipulate the developing Cpe we used the Lmx1aCre line which faithfully labels the hem lineage comprising of Cajal-Retzius cells and CPe cells, when crossed to the Ai9 reporter line (Figure 1A).

Figure 1. Canonical Wnt signaling components are expressed in the developing mouse choroid plexus epithelium. (A) *In situ* hybridization (*Wnt3a, Ttr, Axin2, Fzd1*) and immunohistochemistry (β-CATENIN and LEF1) reveal the expression of canonical Wnt signaling components in E12.5 mouse brain sections. *Lmx1a* is expressed in the hem and the CPe. Lmx1a Cre drives the expression of the Ai9 reporter in these tissues and in the hem-derived Cajal-Retzius cells. (B) Immunohistochemistry reveals the expression of AXIN2, FZD1, β-CATENIN, and LEF1 in the E14.5 mouse CPe. All scale bars are 50 μm.

We tested whether the CPe requires β-CATENIN function by using a conditional mouse
line in which loxP sites flanking exon 2 and 6 of the *ß-catenin* gene (Brault et al., 2001). Cre-
mediated recombination results in the removal of a portion of the armadillo repeat encoding
segment, resulting in a non-functional protein (loss of function, LOF). We used Lmx1aCre to
specifically disrupt *ß-catenin* in the hem and CPe. In the Lmx1aCre:: β-catenin LOF
telencephalon, the CP appears dysmorphic and disorganized (Figure 2). From E14.5, some
regions of the CP lose their characteristic ribbon-like appearance and fail to express the
transcription factor *Otx2*, which is required for CP specification (Johansson et al. 2013), or
Aquaporin 1 (Aqp1), a channel critical for CSF production (Kim and Jung 2012; Castañeyra-Ruiz
et al. 2016). Though TTR expression appears relatively unaffected, the expression of E-
CADHERIN is perturbed, consistent with the abnormally folded appearance of the CP. This
disruption of the CP does not ameliorate with time, and at birth, TTR and E-CADHERIN
expression reveals an abnormally organized CP (Figure 2). In summary, loss of β-CATENIN
function causes profound molecular and cellular perturbations in the CP during its
development, demonstrating the requirement for the canonical Wnt pathway in early CP
development.

103
104
Figure 2. Loss of canonical Wnt signaling (β-catenin LOF) in the hem and CPe disrupts
choroid plexus development. (A) OTX2 and AQP1 are expressed throughout the control
mouse CP, but are missing in the distal tip of the Lmx1aCre:: β-catenin LOF CP at E14.5
(arrowheads). (B, C) TTR (E16.5 and P0) expression reveals a dysmorphic CP in Lmx1aCre:: β-
catenin LOF embryos compared with controls (arrowheads). (B, D) The expression of AQP1
(E16.5) and adherens junction marker E-CADHERIN is uniform in the control CP, but irregular in
the Lmx1aCre:: β-catenin LOF CP (arrowheads). All scale bars are 50 μm.

Constitutively active β-catenin disrupts CPe development

To examine the effects of constitutive and persistent activation of canonical Wnt signaling in the CPe, we used a mouse line in which exon 3 is flanked by loxP sites. Cre-mediated recombination results in a deletion of a domain that regulates β-CATENIN stability (Harada et al., 1999). The recombined β-catenin allele encodes a protein that remains constitutively active (gain of function, GOF). Analysis of sections from Lmx1aCre:: β-catenin GOF embryos reveal a progressive loss of *Ttr* expression from E12.5, and the CP loses its ribbon-like shape (Figure 3A). LEF1 expression is induced in the CP from E12.5, reflecting the enhanced canonical Wnt signaling as a result of β-catenin GOF. By E15.5, Aqp1 expression is not detectable, and the ventricles are collapsed, possibly due to a lack of fluid resulting from reduced AQP1 expression. E-CADHERIN staining in the Lmx1aCre:: β-catenin GOF CP appears disorganized, consistent with an overall disruption of CP differentiation and morphogenesis (Figure 3B).

Figure 3: Constitutively active β-catenin in the hem and CPe disrupts choroid plexus differentiation and morphogenesis (A) *In situ* hybridization for *Ttr* from E12.5-P0 reveals a patchy expression in Lmx1aCre:: β-catenin GOF brains compared with controls. (B) Immunostaining for LEF1 at E12.5 reveals increased expression at E12.5 consistent with activation of the canonical Wnt pathways in Lmx1aCre:: β-catenin GOF brains. At E15.5, AQP1 expression is lost and E-CADHERIN expression is disorganized in the Lmx1aCre:: β-catenin GOF choroid plexus compared with controls (arrowheads). All scale bars are 50 µm.

Gain of hippocampal-like identity and loss of CPe identity upon constitutive activation of β-catenin

The loss of CPe markers in Lmx1aCre:: β-catenin GOF brains prompted us to examine the identity of the TTR/AQP1-negative tissue. It is well established that canonical Wnt signaling from the cortical hem is necessary and sufficient for the induction of hippocampal fate in adjacent tissue (Lee et al., 200; Machon et al., 2007; Mangale et al., 2008). Therefore, we tested whether neuronal or in particular hippocampal markers were upregulated in the Lmx1aCre:: β-catenin GOF CPe. Indeed, regions devoid of *Ttr* expression displayed expression of the pro-neuronal transcription factor *Neurog2* at E13.5 and E15.5, and also of the dentate gyrus marker *Prox1* (Figure 4A, B). To examine this transformation more comprehensively, we microdissected the CP and hippocampus from E14.5 control and β-catenin GOF embryos and processed these samples for bulk RNA-seq (Figure 4C, D). 2439 genes were downregulated and 2163 genes upregulated in the CP upon constitutive activation of β-catenin, using a 1.5-fold cutoff (p.adj < 0.05; Supplementary Figure 1A). Principal Component Analysis (PCA) and Hierarchical clustering of top 500 genes expressed in the CP and the hippocampus revealed that the overall transcriptome of the Lmx1aCre:: β-catenin GOF CP appears to be more similar to the control hippocampus than the control CP (Figure 4E; Supplementary Figure 1B).

We curated a list of CP-enriched genes (Dani et al., 2019) and found that they were downregulated in the β-catenin GOF CP. In contrast, genes related to hippocampal development and dentate gyrus development (GO:0021766, GO:0021542), including bonafide markers of the developing hippocampus, *Zbtb20,* and *Lhx9*, were upregulated (Supplementary Figure 1. Gene Ontology (GO) analysis of the Differentially Expressed (DE) genes indicated that pathways involved in neurogenesis are upregulated in the β-catenin GOF CP (Supplementary Figure 1). In summary, transcriptomic analysis is consistent with the interpretation that upregulation of canonical Wnt signaling leads to a loss of CPe identity and a transformation to a neuronal/hippocampal-like fate.

Figure 4. Constitutive activation of β-catenin transforms the CPe to a neuronal and hippocampal-like molecular identity.

(A, B) The Lmx1aCre:: β-catenin GOF CP appears to have lost *Ttr* expression, which is normally seen throughout the control CP at E15.5 (A), E13.5, and E18.5 (B). In contrast, *Prox1*, *Axin2*, and *Neurog2,* which are not expressed in the control CP, are aberrantly expressed in the Lmx1aCre:: β-catenin GOF CP.

(C) A cartoon depicting the dissection of E14.5 mouse CP and hippocampus for RNA-seq analysis. (D) Principal Component Analysis and (E) Heat map of the top 2000 genes plotted as normalized read counts of the RNA-seq data from control and Lmx1aCre:: β-catenin GOF CP and hippocampus. All scale bars are 50 μm.

The human fetal CP displays components of Wnt signaling.
Given the critical role of CSF in brain development and the finding that both β-catenin
LOF and GOF disrupt CPe specification and morphogenesis, we sought to explore whether this
signaling pathway is important in the development of the human CP. The telencephalic CP
develops between gestational weeks 8 to 12 in the human. We collected human fetal samples
from GW11 and GW13 and performed immunohistochemical analysis for Wnt signaling
components. The human CP is identified by AQP1 and OTX2 expression as well as by its
elaborate fan-like morphology in the large ventricles seen in the human embryonic
telencephalon (Figure 5). The canonical Wnt receptor, FZD1, is expressed throughout the CPe,
as are β-CATENIN, LEF1, and AXIN2 (Figure 5). These findings indicate that canonical Wnt
signaling is likely to be active in the human CPe at early stages during CP development.

195
196
Figure 5: The fetal human CP displays canonical Wnt signaling components
(A) A scheme of the human fetal telencephalic CP (B-C) Immunostaining of human fetal
sections at GW11(B) and GW13 (C) for FDZ1, β-CATENIN, AQP1, LEF1, and AXIN 2 reveals
the presence of canonical Wnt signaling components and also of the CP marker OTX2.
Scale bars are 500 µm (low magnification panels) and 50 µm (high magnification panels) in B
and C.

CPe-like structures arise in hESC organoids upon a combination of BMP and canonical Wnt pathway activation

We sought to test whether constitutive activation of canonical Wnt signaling disrupts CPe identity in the human, as it does in the mouse. First, we used established protocols to achieve CPe differentiation in hESC-derived organoids (Sakaguchi et al., 2015). In this protocol, hESCs are aggregated and seeded in a 96 well low-adhesion plate, and over 18 days of incubation, they form spherical structures which grow in diameter (Figure 6A, B). At this stage, the morphogen cocktail containing BMP4 and CHIR, a GSK3 inhibitor, is added. GSK3 is responsible for phosphorylating β-CATENIN and marking it for degradation, so CHIR addition has the effect of creating a constitutively active β-CATENIN (Ying et al., 2008). However, unlike the β-catenin GOF mouse, CHIR can be added in graded amounts to control the extent of canonical Wnt pathway activation (Naujok et al 2014). A cocktail containing a low level of CHIR (3 μM) together with BMP4 (0.5nM) promotes CPe-like differentiation (Sakaguchi et al., 2015). The spherical organoids became slightly tapering and developed multiple protrusions within a few days of exposure to the cocktail (Figure 6B). Immunohistochemistry reveals CPe markers TTR, AQP1, and OTX2 to be expressed along the perimeter of the protrusions, with the cells displaying a cuboidal morphology (Figure 6C, D). To better characterize these organoids, we performed RNA-seq after 12 days of exposure to the cocktail. Even though the CPe-like differentiation was seen only along the perimeter of the organoid, the transcriptomic analysis of the sample revealed characteristic markers of the CPe such as *TTR, AQP1, KRT18, and CLIC6* (Figure 6E). Interestingly, genes associated with the blood-CSF barrier (BCSFB) such as *CLDN3, TJP1, TJP2, CDH1*, and *JAM2* were also upregulated, as were several transporters and ion channels encoded by the SLC gene family and genes associated with ciliogenesis, consistent with the ciliated epithelium nature of the CPe (Supplementary Figure 2). In summary, consistent with Sakaguchi et al. (2015), CPe-like structures develop in organoids upon exposure to a cocktail of BMP4 and low levels of CHIR.

Figure 6: Differentiation of hESC organoid cultures using a CPe differentiation-promoting

protocol (A) Schematic presentation describing the organoid culture protocol (B) Bright-field

images of organoids at different stages organoid (C, D) Immunostaining reveals the expression

of OTX2, TTR, and AQP1 in the perimeter of the protrusions of a day 40 organoid. The AQP1

immunostaining corresponds to that seen in the E14.5 mouse CP. (E) Heat map generated from

normalized reads obtained by MARS-Seq shows upregulation of CP-enriched genes by day 30

of the organoid culture. Scale bars are 200 μm in (B), 100 μm in (C), and 10 μm in (D).

Loss of CPe-like identity upon increased activation of canonical Wnt signaling in hESC-derived

organoids.

We sought to examine the effect of increased activation of canonical Wnt signaling in

the hESC-derived organoids. We tested a range of concentrations of CHIR (3 μM to 12 μM) in

the cocktail added to the organoids after 18 days *in vitro*, keeping the BMP4 concentration

constant. The organoids were examined for TTR, AQP1, and OTX2 expression by

immunohistochemistry, qPCR analysis, and RNA-seq (Figure 7). As early as 7 days post CHIR

treatment (25 days *in vitro*), 3 μM-treated organoids displayed multiple domains of TTR

decorating the perimeter, but 12 μM-treated organoids displayed no detectable expression of

TTR (Figure 7C; Supplementary Figure 3C). We tested the dose-response to CHIR treatment by

examining the expression of TTR, AQP1, and OTX2 in organoids using the different CHIR

regimens. Whereas all three proteins appeared along the perimeter of 3 μM CHIR-treated

organoids, their expression was undetectable at all higher concentrations (Figure 7D;

Supplementary Figure 3D). qPCR analysis confirmed this finding: upon treatment with 3 µM

CHIR, *TTR, AQP1,* and *OTX2* displayed a significant increase in fold-expression over the baseline

(untreated 18-day organoids). However, treatment with 12 µM CHIR appeared to suppress the

expression of these genes to levels below the baseline (Figure 7E). In contrast, *LEF1* expression

correctly reflected the increased activation of canonical Wnt signaling by displaying a graded

increase in fold expression above baseline upon 3 µM and 12 µM CHIR treatment (Figure 7F).

To obtain a comprehensive picture of the effect of increased canonical Wnt pathway

activation on hESC-derived organoids, we performed RNA-seq of 3 µM and 12 µM CHIR-treated

samples. Several major CPe markers detected in the 3 µM-treated condition are suppressed

upon 12 µM CHIR treatment (Figure 7G). These include markers of differentiated CPe such as

AQP1, TTR, OTX2; genes that encode proteins that participate in forming tight junctions that are

typical of the CPe, such as *CLDN1, 2, 5,* and *TJP1,3*; and those that encode transporters or ion

channels such as *SLC4A10, CLIC6, SLC12A7, and SLC12A8.*

In summary, multiple lines of analysis confirm that the CPe-like differentiation seen in

hESC-derived organoids upon low activation of the canonical Wnt pathway is suppressed upon

increased activation of this pathway.

Figure 7. High activation of the canonical Wnt pathway in hESC-derived organoids causes suppression of CPe markers.

(A) Schematic presentation of the protocol for treatment of organoids with canonical Wnt agonist CHIR. (B, C) The CP marker TTR is expressed in the periphery of a day 25 organoid exposed to 3 μM, but not when exposed to 12 μM CHIR, from day 18. (D) High magnification images (boxed region shown in B) of serial sections of a day 40 organoid exposed to 3 μM reveal the expression of TTR, AQP1, and OTX2 in the periphery. This expression is reduced or undetectable upon exposure to higher concentrations of CHIR (6/ 9/12 μM). Additional examples are in Supplementary Figure 3.

 (E) q-PCR analysis shows that the expression of the CP markers *TTR*, *AQP1*, and *OTX2* is upregulated in organoids at day 30 organoids exposed to 3 μM CHIR from day 18.   This

increased expression is lost upon treatment with 12 µM CHIR for the same period. (F) *LEF1*
expression displays a stepwise increase upon 3 µM and 18 µM CHIR treatment, consistent with
increased activation of the canonical Wnt pathway. Group means were compared using one-
way *ANOVA* followed by posthoc Tukey's multiple comparison test, * p-value<0.05, **
p-value<0.01, *** p-value<0.001(N=4). (G) Heatmap comparing normalized RNA-seq reads for
3 µM and 12 µM CHIR-treated organoids. Several CP-enriched genes are downregulated with
increased activation (12 µM CHIR) of canonical Wnt signaling. All scale bars are 100 µm.

Figure 8: Summary

(A) Schematic presentation showing that decreased Wnt signaling in the CPe leads to a
dysmorphic appearance of the choroid plexus while an increase in Wnt signaling transforms the
CPe into PROX1 positive DG granule cell-like identity. (B) hESC derived CP organoids lose
their choroid identity upon an increase in Wnt signaling following treatment with high CHIR
concentration.

Discussion

We report that β-CATENIN, which mediates canonical Wnt signaling, is required for the normal development of the telencephalic CP. Furthermore, hyper-activation of canonical Wnt signaling resulting from constitutive activation of β-CATENIN alters gene expression and disrupts CPe fate both in the mouse embryo and in hESC-derived CP organoids. Therefore, a finely regulated balance of canonical Wnt signaling appears to be critical for CPe differentiation in the mammalian brain.

As one of the major signaling pathways that regulate multiple aspects of cellular, tissue, and organ development and function, canonical Wnt signaling is tightly controlled, and perturbations of this pathway cause an array of defects: In human hair follicle morphogenesis, loss or constitutive activation of β-CATENIN levels leads to tumor formation or defective differentiation, respectively (Huelsken et al., 2001, Kajino et al 2001). In several cases, constitutive activation of β-CATENIN appears to transform the identity of the cells in which this perturbation is introduced: the mouse epiblast prematurely displays epithelial to mesenchymal transition (Kemler et al., 2004); endothelial cells of the midbrain transform to a "barrier cell" identity (Wang et al., 2019), and airway epithelial cells take on a neuroendocrine-like fate (Li et al., 2009). Our report of a transformation of CP epithelial cells to a neuronal fate upon constitutive activation of β-CATENIN has parallels with what has been seen in other systems.

An unexpected feature of the CPe-to-neuronal transformation, however, is that the transcriptomic profile of the β-catenin GOF CP is hippocampal-like, not CR cell-like. It has been well established that the hem is the progenitor domain for both CPe and CR cells (Louvi et al., 2007). This fate choice is regulated by transcription repressors of bHLH family members HES1, 2, and 3, which promote epithelial identity by suppressing neurogenic genes and inducing BMP signaling. Upon loss of all three *Hes* genes, the hem upregulates the pro-neuronal factor *Neurog2* and produces CR cells instead of CPe cells (Imayoshi *et al.*, 2008). How might hippocampal-like cells arise from this lineage? A study using Lmx1aCre-driven lineage analysis reported that in addition to CR and CPe cells, a small population of hippocampal cells also appears to originate in the hem, regulated by LMX1a itself (Chizhikov et al., 2010). Taking into consideration that canonical Wnt signaling is known to be both necessary and sufficient for hippocampal fate specification (Lee et al., 2000; Machon et al., 2007; Solberg et al., 2008), β-catenin GOF may drive the hem/CPe to this third "hippocampal" branch of the hem lineage. Together, the results suggest that the two neuronal derivatives of the hem may be regulated by distinct mechanisms, and this motivates a careful analysis of how different regulatory pathways may interact to drive differentiation along the non-neuronal and neuronal lineages arising from the hem.

Activation of the canonical Wnt pathway is often tumorigenic (Kim and Jeong 2019).
Human cases of CP carcinoma are associated with mutations in exon 3 of the *ß-CATENIN* gene
that lead to its constitutive activation (Kim & Jeong, 2019). Mouse models of CP papilloma that
are generated by constitutive activation of NOTCH signaling show concomitant upregulation of
the canonical Wnt pathway (L. Li et al., 2016). It is not clear what mechanisms control whether
canonical Wnt activation causes cell fate transformations or tumorigenic changes, but it is likely
that a crosstalk with other major signaling pathways is involved. The β-catenin GOF mouse CP
offers a useful and accessible model in which this question can be examined.
346
The hem also expresses the non-canonical WNT5A, which is required for normal CPe
cytoarchitecture, apicobasal polarity, and overall size and complexity of the CP (Langford et al.,
2000). Together with the established role for BMPs in the specification of the CP (Hebert *et al.*,
2002; Watanabe *et al.*, 2012), it is apparent that the known signaling molecule families
expressed in the hem, the BMPs, canonical and non-canonical WNTs, are each critical for CP
development. Our findings regarding the role of the canonical Wnt pathway motivate a careful
examination of how these pathways crosstalk with each other to execute the specification,
cytoarchitecture, morphogenesis, and expansion of the non-neuronal choroid plexus arising
from neuroepithelium.
356
Key features of β-catenin GOF in the CP include dysregulation of factors that are
essential for the CP to produce the CSF. These include several anion cation exchangers, ion
channels, and solute transporters each of which acts as regulators of CSF composition: a
sodium-potassium ATPase, *Atp1a1* (Ernst et al., 1986); an epithelial sodium channel, *Scnn1a*
(Amin et al., 2009); an inward rectifying potassium channel, *Kcnj13* (Cornejo et al., 2018); a
voltage-gated potassium channel, *Kcna1* (Speake et al., 2003); and a sodium-potassium-chloride
cotransporter 1, *Nkcc1*, that modulates water transport underlying CSF production (Steffensen
et al., 2018) and is also associated with Blood CSF barrier disruption (Kim & Jung., 2012).
Furthermore, the expression of several genes encoding transporters for glucose, chloride, and
amino acids, is also downregulated in the β-catenin GOF CP. Together, the data indicate that
canonical Wnt signaling is critical to enabling the CPe to produce normal CSF. Mutations in exon
3 of *ß-CATENIN* in humans result in stabilization of β-CATENIN (Bjorklund et al., 2008), similar to
that seen in the β-catenin GOF mouse model (Harada et al., 1999). Human studies have focused
on the association of these mutations with carcinomas (Gao and Zang., 2018; Kim & Jeong.,
2019), but no associated effects on the CSF have been explored. Our study identifies a major
developmental outcome of the β-catenin GOF perturbation in terms of its effect on the
composition of the CSF.
374
The formation of CPe in hESC-derived organoids requires low activation of the canonical

Wnt pathway together with BMP signaling. Loss of CPe fate in these organoids upon high

activation of the canonical Wnt pathway indicates conservation of the role of this pathway in

the development of the CPe. hESC-derived organoids are not only an excellent system to study

conserved developmental mechanisms but also offer the opportunity to test for subtle effects

of altered CP/CSF status on neuronal cells. Depending on the protocol used, the organoids

develop into fluid-filled cysts that contain several human CSF proteins (Pellegrini et al., 2020).

These organoids are permeable to small molecules and are excellent model systems to test

modalities of drug delivery to the CNS (Pellegrini et al., 2020). Thus, hESC-derived CP organoids

offer a suitable system to model both diseases as well as modalities for treatment.

In conclusion, we demonstrate that the developing choroid plexus expresses

components of the canonical Wnt signaling pathway in embryonic mouse and human brains.

Both loss and gain of function of β-CATENIN disrupt CPe development, indicating that this

pathway is likely to be under a finely tuned regulation *in vivo*. These findings motivate future

studies on how this regulation is achieved and may provide a mechanistic basis for subtle

deficits in CP/CSF function connected to the disruptions of the canonical Wnt pathway function

in this system.

Supplementary Figures

Supplementary Figure 1:   Analysis of differentially expressed gene sets in the control and β-
catenin GOF CP and Hippocampus.
(A-E) Analysis of RNA-seq data from the control (Ctrl) versus Lmx1aCre:: β-catenin GOF CP and
Hippocampus (HP). (A) Volcano plot showing Differentially Expressed Genes in the control and
β-catenin GOF CP (B) Sample correlation matrix revealing the similarity of the β-catenin GOF CP
to wild type HP rather than the wild type CP. (C) DEG reveals the loss of CP enriched genes and
the gain of HP genes in the GOF CP. (D) GO analysis of upregulated and downregulated genes
from (A). (E) Heatmap of normalized read counts of genes belonging to different classes of
transporters expressed in wild type and the β-catenin GOF CP.

404
405
Supplementary Figure 2: Characterization of CPe-like differentiation in hESC organoids
(A-E) A comparison of transcriptomics data from day 18 and day 30 organoids. (A) Sample
correlation matrix (B, C). Heat map of the normalized reads reveals that several BCSF genes are
upregulated (B) and ciliogenesis-related genes are differentially regulated (C). (D) GO analysis
(E) Heat map plotted from normalized reads shows an overall upregulation of transporter genes
(belonging to the Slc gene family) in day 30 organoids.

Supplementary Figure 3: High activation of the canonical Wnt pathway in hESC-derived

organoids causes suppression of CPe markers.

Additional examples of data shown in Figure 7 C and D. (A) Scheme showing the protocol for
treatment with canonical Wnt agonist CHIR (C) CP marker TTR is expressed in the periphery of
day 25 organoids (n=3) exposed to 3 μM, but not 12 μM CHIR, from day 18.   (D) High
magnification images (boxed region shown in B) of serial sections of four examples of day 40
organoids exposed to 3 μM reveal the expression of TTR, AQP1, and OTX2 in the

421 periphery. This expression progressively decreases and is undetectable upon increasing
422 exposure to higher concentrations of CHIR (6/ 9/ 12 μM). All scale bars are 100 μm.
423
424 **Material and methods**
425
Tissue collection:
Mice: All animal protocols were approved by the Institutional Animal Ethics Committee of the
Tata Institute of Fundamental Research, Mumbai, India. The mouse lines used in this study
were kind gifts from Kathy Millen (Center for Integrative Brain Research, Seattle Children's
Research Institute; Lmx1aCre line, Chizikov and Millen., 2004), Raj Awatramani (Department of
Neurology and Center for Genetic Medicine, Northwestern University Feinberg Medical School
Chicago, Illinois; β-catenin LOF mice, Joksimovic and Awatramani., 2014). The Ai9 mouse line
was obtained from JAX labs Stock No. 007909. Noon of the day of the vaginal plug was
designated as embryonic day 0.5 (E0.5). Control embryos were littermates that did not carry
the Lmx1aCre allele. Since the Lmx1aCre transgene is on the X-Chromosome and females
showed mosaic expression of the transgene due to random X inactivation, only male
embryos/pups with the Lmx1aCre were used for analysis. A total of 34 wild-type (Figures 1-4), 9
β-catenin LOF (Figure 2), and 15 β-catenin GOF (Figure 3, and 4) embryos/pups were used for in
situ hybridisation and immunohistochemistry analysis. Breeding colonies were maintained with
the female breeders carrying the Lmx1aCre transgene since this Cre line shows germline
recombination in the testis. The sex of the embryos was determined by examining the gonads
from E12.5 onwards.
Primers used were: Cre F: ATTTGCCTGCATTACCGGTC, Cre R: ATCAACGTTTTCTTTTCGG, Cre
positive DNA shows a band at 350bp. Other mouse lines were genotyped using references
mentioned before.
446
Human: Fetal brain specimens were obtained from either medically indicated or spontaneous
abortions. The Institutional Ethical Review Board of the School of Medicine, University of
Zagreb and University Hospital Center Zagreb has approved the procedure for the collection of
postmortem brain samples (UHC Zagreb EP02/21AG; UZSM: 641-01/19-02/01). The fetal age
was estimated based on crown-rump length (CRL, in mm), pregnancy records, and histological
findings, and expressed as GW. In addition, the correlation of maturational parameters (CRL,
body mass, pregnancy records, and sonographic examination) revealed no evidence of growth
retardation or malformations. Finally, specimens in which subsequent postmortem analysis
revealed macroscopically or microscopically identified pathological changes, or postmortem
autolytic changes, were excluded.
457
458

Immunohistochemistry

<u>Mouse sections and organoids:</u> Mouse brain sections /organoid sections were mounted on plus
slides (Catalogue number: EMS 71869-11) and dried for 2-3 hours. Slides were transferred to a
slide mailer (Catalogue number: EMS 71549-08) containing PBS+0.01% TritonX-100 for 10
minutes followed by a wash with PBS+0.03% TritonX-100 for 5 minutes. For antigen retrieval,
sections were boiled in a 10mM sodium citrate buffer (pH=6) at 90^0C for 10 minutes using a
water bath.   Slides were cooled to room temperature and washed with PBS+0.01% TritonX-
100 for 10 minutes. Blocking (5% horse serum /Lamb serum in PBS + 0.1% TritonX-100) for 1
hour followed by overnight primary antibody incubation at 4°C.   Secondary antibody
incubation was performed at room temperature for 2 hours followed by 3 washes with 1x PBS.
Slides were mounted using vector shield mounting media (Vector laboratories H-1000-10) and
imaged in an Olympus FluoView 1200 confocal microscope. The primary antibodies used were:
LEF1 (rabbit,1:200 CST catalogue # C12A5), β-CATENIN (mouse 1:200,BDbiosciences catalogue #
610153),RFP(rabbit,1:200 Abcam catalogue # ab62341)FZD1 (rabbit,1:100 catalogue #
LS-A4150),AXIN2 (rabbit ,1:200 Abcam catalogue # ab185821),TTR (rabbit 1:100 ,DAKO
catalogue # A0002), AQP1 (mouse 1:200 SCBT catalogue # sc25287),OTX2(mouse,1:200SCBT
catalogue # sc-514195),OTX2(1:500,invitrogen,PA5-39887),AQP1 (rabbit, 1:200,invitrogen
catalogue # MA5-32593),E-CADHERIN(mouse,1:200,SCBTcatalogue#sc21791).   Secondary
antibodies used were following: Goat Anti Rabbit Alexa fluor 488 (1:200, Invitrogen catalogue #
A11034), Goat Anti mouse Alexa fluor 594 (1:200, Invitrogen catalogue # R37121), Goat Anti
Rabbit Alexa fluor 568 (1:200, Invitrogen catalogue # A11011), Donkey Anti goat Alexa fluor 568
(1:200, Invitrogen catalogue # A32814).

<u>Human:</u> The entire brains were fixed by immersion in 4% paraformaldehyde in 0.1 M phosphate
buffer saline (pH=7.4). Subsequently, tissue blocks were embedded in paraffin wax. Serial
sections were cut (5 µm) and stained. The cresyl violet staining was used to delineate
cytoarchitectonic boundaries, while adjacent sections were labelled using immunofluorescence
with various antibodies. The following primary antibodies were used: (1) rabbit polyclonal anti-
Axin 2, (ab185821, Abcam, Cambridge, United Kingdom); (2) the rabbit polyclonal Frizzled 1
antibody, FZD1 (LS-A4150, LSBio, Seattle, Washington, SAD); (3) the monoclonal rabbit LEF1
antibody (C12A5, Cell Signaling, Leiden, Netherlands); (4) the mouse monoclonal aquaporin1
antibody, AQP1 (sc-25287, Santa Cruz, Dallas, Texas, SAD); (5) the rabbit non-phospho (Active)
βCatenin (D13A1, Cell Signaling, Leiden, Netherlands); (6) the monoclonal mouse OTX2
antibody (MA5-15854, Invitrogen, Carlsbad, California, SAD); (7) the mouse monoclonal Beta-
Catenin, (M3539, Dako Agilent, Santa Clara, California, SAD).

The dewaxing of sections was performed in xylol for 5 min, and a series of alcohol solutions
(100%, 96% 70%) for 2 min.   Slides were then washed in PBS with 0.03% triton (10min).

The antigen retrieval was performed in citrate buffer pH6.0 and microwave (800W) for 10
minutes. The slides were cooled at room temperature and immersed in a blocking solution (PBS
containing 3 % bovine serum albumin, BSA, and 0.01 % Triton X-100, all from Sigma, St. Louis,
MO) for 30 min. to prevent nonspecific background staining. Sections were then incubated with
primary antibodies for 48 h at 4˚C (AXIN2 1:100, FZD1 1:100, LEF1 1:100, AQP1 1:100, OTX2
1:50, β-Catenin cell signaling 1:400, β-Catenin Dako 1:200), washed and subsequently incubated
with secondary anti-mouse or anti-rabbit antibodies diluted in PBS (at 1:1000) for 2 h at the
room temperature. For visualization of the specific immune reactivity following secondary
antibodies were used according to the laboratory protocol (Culjat and Jovanov-Milosevic,
2019):   the donkey anti-mouse IgG Alexa fluor-plus 488, donkey anti-rabbit IgG Alexa fluor-
plus 647, donkey anti-rabbit IgG 488, donkey anti-mouse IgG Alexa-fluor 546 were used.
Sections were covered with a mounting medium containing DAPI. Negative controls were
included in all experiments by replacing the primary antibody with a blocking solution. No
immunolabeling was detected in control sections.

In situ hybridization

Mouse brains were sectioned using a freezing microtome (Leica SM2000R Sliding Microtome)
to a thickness of 20 µM. Sections were mounted on super frost plus slides (Catalogue #: EMS
71869-11), fixed with 4% (w/v) paraformaldehyde for 15 minutes, washed three times with PBS.
The sections were then treated with proteinase K dissolved in Tris-EDTA buffer (1 µg/ml).   Post
fixation was performed in 4% PFA for 15 minutes and the sections were washed three times in
PBS. Hybridization was performed for 16 h at 70°C in a buffer containing 50% (v/v) formamide,
2X SSC, and 1% (w/v) SDS. Digoxigenin (DIG)-labelled cRNA probes were used for hybridization
and were prepared from respective plasmids using in-vitro transcription. After hybridization,
three stringent washes (for 45 minutes each) were performed in solution X (50% formamide, 2X
SSC, and 1% SDS) followed by a wash with 2XSSC and then 0.2XSSC.   Blocking was performed
for 1 hr with 10% horse serum in TBST (Tris-buffered saline pH 7.5 with 0.1% Tween-20).
Sections were incubated in alkaline phosphatase-conjugated anti-DIG (Digoxygenin) antibody
Fab fragments (1:5000; Roche, catalogue #12486523) at 1:5000 in the blocking buffer and
incubated at 4°C overnight.   The color reaction was performed using NBT/BCIP substrate
(Roche, 4-nitroblue tetrazolium chloride, catalogue #70210625; 5-Bromo- 4-chloro-3-indolyl
phosphate, catalogue #70251721). Counterstaining was performed using Fast Red (Sigma-
Aldrich, catalogue # N3020). Slides were dried and coverslipped using DPX mounting reagent
(SDFCL, catalogue #46029).

Image acquisition and analysis

Bright field images were acquired using Zeiss Axioskop-2 plus microscope equipped with a Nikon DS-fi2 camera and associated software. Mouse sections were imaged in Olympus FluoView 1200 confocal microscope and the human brain sections were imaged in Olympus FluoView 3000 confocal microscope. All the organoid sections were imaged using the Andor Dragonfly spinning disk confocal microscope system. All the image analysis was done on Fiji-ImageJ and/or Adobe Photoshop CS6.

Generation of hESC derived organoids

The human ES cell line WIBR3, obtained from the Whitehead Institute for Biomedical Research, (https://hpscreg.eu/cell-line/WIBRe001-A), was used in compliance with all applicable statutes and regulations and compliance with applicable guidelines (including the NAS Guidelines or guidelines of the International Society for Stem Cell Research as relevant to the conduct of human embryonic stem cell research ["ISSCR Guidelines"]. The line was cultured on a MEF (mouse embryonic fibroblast) feeder layer. To maintain the pluripotent state, a customized human Naïve media with 40 μM Rock Inhibitor Y-27632 was used as described in Karzbrun et al 2018. Differentiation of hES aggregates into Choroid plexus like organoid was done using a protocol described in Sakaguchi et al 2015 (Sakaguchi et al., 2015). Briefly, hESC colonies were grown in a Matrigel plate till confluency, then the cells were dissociated with trypsin and rapidly reaggregated in an ultra-low attachment 96 well V bottom plate (9000 cells per well) containing a media termed as SA1 media. This media contains DMEM (Invitrogen cat # 11965092) supplemented with 20% (vol/vol) KSR (Invitrogen cat # 10828028), 0.1mM non-essential amino acids (Invitrogen cat # 11140050), 1mM pyruvate (Invitrogen cat # 11360070), 0.1mM 2-mercaptoethanol (Invitrogen cat # 21985023), 100U/ml penicillin and 100 mg/ml streptomycin. 0.3 mM IWR1e (tankyrase inhibitor, MedChemExpress cat# HY-12238) and 5 mM SB431542 (TGFβ inhibitor; MedChemExpress HY-10431) were added to culture from day 0 to day 18 (day of aggregation was defined as day 0). The culture media used after day 18 till day 40 was termed as SA2, which contains DMEM/F-12 (Invitrogen cat # 21331020), GlutaMAX(TM) (Gibco) supplemented with 1% N-2 supplement (Invitrogen cat # 17502048), 1% Chemically Defined Lipid Concentrate (Invitrogen cat # 11905031), fetal bovine serum (FBS; 10% vol/vol), 3 μM CHIR 99021 (GSK3 inhibitor; MedChemExpress cat # HY-10182) and 0.5 nM BMP4 (R&D cat # 314-BP-050) under normoxia condition. The medium was changed once every 2 days. From day 5 onwards the plates were kept on an orbital shaker placed inside the incubator.   On day 18 the organoids were transferred from 96 well plates to 10 cm ultra-low attachment plate (6 organoids per plate) and kept on an orbital shaker inside the incubator. For sample collection, the organoids were rinsed with 1x PBS for 2 times, then either flash-frozen for RNA-seq experiments or transferred to a small dish containing 4% PFA solution for

567 immunohistochemistry. A total of 12 control organoids (3 µM CHIR) were used for Figure 6. For
568 Figure 7, 14 organoids (3 µM CHIR), 9 organoids (6 µM CHIR), 5 organoids (9 µM CHIR treated),
569 and 14 organoids (12 µM CHIR treated), were used immunohistochemistry analysis.

570 **RNA sequencing, library preparation, and analysis of mouse tissues**
571

The mouse CP was dissected from E14.5 control and β-catenin GOF brains and stored in Trizol$^{®}$
reagent. Extracted RNA of 1 µg (RIN > 7.5, measured using the Agilent 2100 bio analyzer) from
CPe dissected from 4 embryos was pooled for each of two biological replicates. After library
preparation, sequencing was performed on the Illumina platform to achieve 100bp or 150bp
reads to generate 30 Million paired-end reads per sample. FastQ QC was performed as
described in (Andrews, S 2010), and reads > 30 Phred scores were aligned using HISAT2 (D.Kim
et al., 2019). Feature counts were used to quantify the number of reads per transcript.
Differential expression analysis was performed using DESeq2 (Love et al., 2014) on the R
platform (v3.6.3). Genes showing |log2 Fold change|≥1 were used for further analysis. Gene
ontology analysis was performed using geneontology.org or gprofiler.org.   Semantics were
summarized using "REVIGO" (Supek et al., 2011), and bar plots were created with GraphPad
Prism V7.0 for Windows, GraphPad Software, San Diego California USA, www.graphpad.com).
Heat maps and sample correlation plots were plotted using R studio. Gene-based heatmaps
were plotted using normalized reads on Morpheus (Morpheus,
https://software.broadinstitute.org/morpheus)
587

588 **RNA sequencing, library preparation, and analysis of organoids**
589

Organoids generated in figure 6 and supplementary figure 2, were pooled such that for one
biological replicate 3 organoids were used and three such biological replicates were performed
for day 18 and day 30 organoids. Bulk MARS-seq libraries were produced from 50ng of total
RNA as previously described (Meador et al.,2014). Libraries were then sequenced with 75bp
single-end read on Illumina Nextseq500 platform and FASTQ files were processed and analyzed
using UTAP (Kohen et al., 2019). Briefly, sequenced reads were trimmed using cutadapt
(parameters: -a adaptor -a "A {10}" –times 2 -u 3 -u -3 -q 20 -m 25) (M Martin., 2011) and were
mapped to hg38 indexed reference genome using STAR (Dobin et al., 2013) v2.4.2a with the
following parameters: –alignEndsType EndToEnd, –out Filter Mismatch Nover Lmax 0.05, –two
pass Mode Basic). UMIs were counted after marking duplicates using HTSeq-count (Anders et
al., 2015) in union mode. For the number of reads per gene, 1000bp of 3' end Gencode
annotated transcripts were counted. The total RNA-seq from 3 µM and 12 µM CHIR treated
organoids (Figure 7) were done using TruSeq stranded mRNA library kit and sequenced by DNA
Link Sequencing Lab, Korea, on NovaSeq 6000, 100PE. Data were analyzed with the UTAP

pipeline (Kohen et al., 2019), where adapter trimming and reference genome alignment was
performed as described above. Read count per gene was done with STAR. Normalization of the
counts and differential expression analysis was performed using DESeq2 (Love et al., 2014) with
the parameters: betaPrior=True, cooks Cutoff=FALSE, independentFiltering=FALSE. Raw P
values were adjusted for multiple testing using the procedure of Benjamini and Hochberg.
Genes with log2FC >1 or <-1, p adjust < 0.05 and baseMean > 10 were considered as
differentially expressed. GO Biological process overrepresentation test on differentially
expressed genes between different time points of organoid growth was performed with the
package clusterProfiler (Yu et al., 2012).
613
614 **qPCR Analysis of organoids**
615
Total RNA of Day 18 and Day 30 organoids was extracted using the RNeasy Mini kit (Qiagen,
Germany) following the manufacturer's protocol and followed by DNAseI treatment. RNA
concentration was measured using Nanodrop (Thermo Scientific, MA, USA). cDNA was
synthesized using M-MLV reverse transcriptase (M3682, Promega, Wisconsin, USA). Real-Time
reactions were performed in triplicates using KAPA SYBR FAST qPCR Kit (2X) on QuantStudio 5
Real-time PCR system (Bio-Rad, CA, USA) following the manufacturer's recommendation.
Expression levels were normalized against GAPDH using the $\Delta\Delta$ threshold cycle (Ct) method.
The fold changes were represented as mean±SEM. Primers: *AQP1* Fw: 5'-
TGGACACCTCCTGGCTATTG-3' Rw: 5'-GGGCCAGGATGAAGTCGTAG-3', *LEF1* Fw: 5'-
AGAACACCCCGATGACGGA-3' Rw: 5'-GAGGGTCCCTTGTTGTAGAGG-3', *TTR* Fw: 5'
ATCCAAGTGTCCTCTGATGGT 3' Rw: 5' GCCAAGTGCCTTCCAGTAAGA 3', *OTX2* Fw: 5'-
AGAGGACGACGTTCACTCG-3' Rw: 5'-TCGGGCAAGTTGATTTTCAGT-3'.
628
Data availability: The RNA-seq data described in the manuscript are deposited in the GEO
database (Accession number: GSE162784 and GSE162808). All other codes are available on
request with the corresponding authors.

Acknowledgments

We thank K. Millen (Seattle Children's hospital) for the kind gift of the Lmx1aCre mouse line;
Shital Suryavanshi and the animal house staff of the Tata Institute for Fundamental Research
(TIFR) for excellent support. This work was supported by a Wellcome Trust-Department of
Biotechnology India Alliance Early Career Fellowship (IA-E-12-1-500765; MC); by the Canada-
Israel Health Research Initiative, jointly funded by the Canadian Institutes of Health Research,
the Israel Science Foundation, the International Development Research Centre, Canada and the
Azrieli Foundation (ST); intramural funds from TIFR-DAE (12-R&D-TFR-5.10-0100RTI2001).

Research in the Reiner lab is supported by ISF-National Science Foundation of China (NSFC) joint

research program (Grant No. 2449/16), with the aid of a grant no. 2397/18 from the Canadian
Institutes of Health Research (CIHR), the International Development Research Center (IDRC),
the ISF, a grant from the Ministry of Science & Technology, Israel & The Ministry of Science and
Technology of the People's Republic of China., German-Israeli Foundation (GIF; Grant no. I-
1476-203.13/2018), and United States-Israel Binational Science Foundation (BSF; Grant No.
2017006). In addition, by the Helen and Martin Kimmel Institute for Stem Cell Research, Nella
and Leon Benoziyo Center for Neurological Diseases, David and Fela Shapell Family Center for
Genetic Disorders Research, Brenden-Mann Women's Innovation Impact Fund, Richard F.
Goodman Yale/Weizmann Exchange Program, The Irving B. Harris Fund for New Directions in
Brain Research, Irving Bieber, M.D. and Toby Bieber, M.D. Memorial Research Fund, Leff Family,
Barbara & Roberto Kaminitz, Sergio & Sônia Lozinsky, Debbie Koren. Jack and Lenore
Lowenthal, Dears Foundation. O. R. is incumbent of the Berstein-Mason professorial chair of
Neurochemistry and Head of M. Judith Ruth Institute for Preclinical Brain Research.

The study on the human brains was financed by Croatian Science Foundation projects IP-2019-
04-3182; DOK-2018-01-3771; DOK-2015-10-3939 to NJM and the Scientific Centre of Excellence
for Basic, Clinical, and Translational Neuroscience (GA KK01.1.1.01.0007 funded by the
European Union through the European Regional Development Fund). The collaboration was
initiated throughout COST Action 16118.

We thank the members of Tole lab, Reiner lab, and Jovanov lab for their valuable inputs and
discussion.

References

Abu-Khaklil, A., Fu, L., Grove, E. A., Zecevic, N. & Geschwind, D. H. Wnt genes define distinct boundaries in the developing human brain: Implications for human forebrain patterning. *J. Comp. Neurol.* **474**, 276–288 (2004).

Amin, M. S., Reza, E., Wang, H., & Leenen, F. H. H. (2009). Sodium transport in the choroid plexus and salt-sensitive hypertension. *Hypertension*, *54*(4), 860–867. https://doi.org/10.1161/HYPERTENSIONAHA.108.125807

Anders, S., Pyl, P. T., & Huber, W. (2015). HTSeq-A Python framework to work with high-throughput sequencing data. *Bioinformatics*, *31*(2), 166–169. https://doi.org/10.1093/bioinformatics/btu638

Andrews, S. (2010). **FastQC**: A Quality Control Tool for High Throughput Sequence Data http://www.bioinformatics.babraham.ac.uk/projects/fastqc

Björklund, P., Lindberg, D., Åkerström, G., & Westin, G. (2008). Stabilizing mutation of CTNNB1/beta-catenin and protein accumulation analyzed in a large series of parathyroid tumors of Swedish patients. *Molecular Cancer*, *7*, 1–8. https://doi.org/10.1186/1476-4598-7-53

Brault, V., Moore, R., Kutsch, S., Ishibashi, M., Rowitch, D. H., McMahon, A. P., Sommer, L., Boussadia, O., & Kemler, R. (2001). Inactivation of the β-catenin gene by Wnt1-Cre-mediated deletion results in dramatic brain malformation and failure of craniofacial development. *Development*, *128*(8), 1253–1264.

Castañeyra-Ruiz, L., González-Marrero, I., Hernández-Abad, L. G., Carmona-Calero, E. M., Meyer, G., & Castañeyra-Perdomo, A. (2016). A Distal to Proximal Gradient of Human Choroid Plexus Development, with Antagonistic Expression of Glut1 and AQP1 in Mature Cells vs. Calbindin and PCNA in Proliferative Cells. *Frontiers in Neuroanatomy*. https://doi.org/10.3389/fnana.2016.00087

Chip, H. B. B., Park, T., Mustafaoglu, N., Herland, A., Hasselkus, R., Mannix, R., Fitzgerald, E. A., Prantil-baun, R., Watters, A., Henry, O., Benz, M., Sanchez, H., Mccrea, H. J., Goumnerova, L. C., Song, H. W., Palecek, S. P., Shusta, E., & Ingber, D. E. (2019). Hypoxia-enhanced Blood-Brain Barrier Chip recapitulates human barrier function and shuttling of drugs and antibodies. *Nature Communications*, 1–12. https://doi.org/10.1038/s41467-019-10588-0

Chizhikov, V. V., Lindgren, A. G., Mishima, Y., Roberts, R. W., Aldinger, K. A., Miesegaes, G. R., Currle, D. S., Monuki, E. S., & Millen, K. J. (2010). Lmx1a regulates fates and location of cells

originating from the cerebellar rhombic lip and telencephalic cortical hem. *Proceedings of the*

National Academy of Sciences. https://doi.org/10.1073/pnas.0910786107

Chizhikov, Victor V., & Millen, K. J. (2004). Mechanisms of roof plate formation in the vertebrate

CNS. *Nature Reviews Neuroscience*, *5*(10), 808–812. https://doi.org/10.1038/nrn1520

Cornejo, I., Villanueva, S., Burgos, J., López-Cayuqueo, K. I., Chambrey, R., Julio-Kalajzic, F.,

Buelvas, N., Niemeyer, M. I., Figueiras-Fierro, D., Brown, P. D., Sepúlveda, F. V., & Cid, L. P.

(2018). Tissue distribution of Kir7.1 inwardly rectifying K+ channel probed in a knock-in mouse

expressing a haemagglutinin-tagged protein. *Frontiers in Physiology*, *9*(APR), 1–12.

https://doi.org/10.3389/fphys.2018.00428

Culjat, M., & Milošević, N. J. (2019). Callosal septa express guidance cues and are paramedian

guideposts for human corpus callosum development. *Journal of Anatomy*, *235*(3), 670–686.

https://doi.org/10.1111/joa.13011

Dani, N., Herbst, R. H., Habib, N., Head, J., Dionne, D., Nguyen, L., Mccabe, C., Cui, J., Shipley, F.

B., Jang, A., Rodman, C., Riesenfeld, S. J., Zhang, F., Rozenblatt-Rosen, O., Regev, A., & Lehtinen,

M. K. (n.d.). *A cellular and spatial map of the choroid plexus across brain ventricles and ages*.

https://doi.org/10.1101/627539

Dobin, A., Davis, C. A., Schlesinger, F., Drenkow, J., Zaleski, C., Jha, S., Batut, P., Chaisson, M., &

Gingeras, T. R. (2013). STAR: Ultrafast universal RNA-seq aligner. *Bioinformatics*, *29*(1), 15–21.

https://doi.org/10.1093/bioinformatics/bts635

Eiraku, M., Watanabe, K., Matsuo-Takasaki, M., Kawada, M., Yonemura, S., Matsumura, M.,

Wataya, T., Nishiyama, A., Muguruma, K., & Sasai, Y. (2008). Self-Organized Formation of

Polarized Cortical Tissues from ESCs and Its Active Manipulation by Extrinsic Signals. *Cell Stem*

Cell, *3*(5), 519–532. https://doi.org/10.1016/j.stem.2008.09.002

Ernst, S. A., Palacios, J. R., & Siegel, G. J. (1986). Immunocytochemical localization of Na+,K+-

ATPase catalytic polypeptide in mouse choroid plexus. *Journal of Histochemistry and*

Cytochemistry, *34*(2), 189–195. https://doi.org/10.1177/34.2.3003182

Furuta, Y., Piston, D. W. & Hogan, B. L. M. Bone morphogenetic proteins (BMPs) as regulators of

dorsal forebrain development. *Development* **124**, 2203–2212 (1997).

Gao, C., Wang, Y., Broaddus, R., Sun, L., Xue, F., & Zhang, W. (2018). Exon 3 mutations of

CTNNB1 drive tumorigenesis: A review. *Oncotarget*, *9*(4), 5492–5508.

https://doi.org/10.18632/oncotarget.23695

Grove, E. A., Tole, S., Limon, J., Yip, L. W., & Ragsdale, C. W. (1998). The hem of the embryonic
cerebral cortex is defined by the expression of multiple Wnt genes and is compromised in Gli3-
deficient mice. *Development*, *125*(12), 2315–2325.

Gu, X., Liu, B., Wu, X., Yan, Y., Zhang, Y., Wei, Y., Pleasure, S. J., & Zhao, C. (2011). Inducible
genetic lineage tracing of cortical hem derived Cajal-Retzius cells reveals novel properties. *PLoS*
ONE. https://doi.org/10.1371/journal.pone.0028653

Harada, N., Tamai, Y., Ishikawa, T. O., Sauer, B., Takaku, K., Oshima, M., & Taketo, M. M. (1999).
Intestinal polyposis in mice with a dominant stable mutation of the β-catenin gene. *EMBO*
Journal, *18*(21), 5931–5942. https://doi.org/10.1093/emboj/18.21.5931

Hébert, J. M., Mishina, Y., & McConnell, S. K. (2002). BMP signaling is required locally to pattern
the dorsal telencephalic midline. *Neuron*, *35*(6), 1029–1041. https://doi.org/10.1016/S0896-
6273(02)00900-5

Huelsken, J., Vogel, R., Erdmann, B., Cotsarelis, G., & Birchmeier, W. (2001). β-Catenin controls
hair follicle morphogenesis and stem cell differentiation in the skin. *Cell*, *105*(4), 533–545.
https://doi.org/10.1016/S0092-8674(01)00336-1

Imayoshi, I., Shimogori, T., Ohtsuka, T., & Kageyama, R. (2008). Hes genes and neurogenin
regulate non-neural versus neural fate specification in the dorsal telencephalic midline.
Development. https://doi.org/10.1242/dev.021535

Johansson, P. A., Irmler, M., Acampora, D., Beckers, J., Simeone, A., & Gotz, M. (2013). The
transcription factor Otx2 regulates choroid plexus development and function. *Development*,
140(5), 1055–1066. https://doi.org/10.1242/dev.090860

Johansson, Pia A. (2014). The choroid plexuses and their impact on developmental
neurogenesis. *Frontiers in Neuroscience*, *8*(OCT), 1–9. https://doi.org/10.3389/fnins.2014.00340

Kajino, Y., Yamaguchi, A., Hashimoto, N., Matsuura, A., Sato, N., & Kikuchi, K. (2001). β-Catenin
gene mutation in human hair follicle-related tumors. *Pathology International*, *51*(7), 543–548.
https://doi.org/10.1046/j.1440-1827.2001.01231.x

Karzbrun, E., Kshirsagar, A., Cohen, S. R., Hanna, J. H., & Reiner, O. (2018). Human brain
organoids on a chip reveal the physics of folding. *Nature Physics*, *14*(5), 515–522.
https://doi.org/10.1038/s41567-018-0046-7

Kemler, R., Hierholzer, A., Kanzler, B., Kuppig, S., Hansen, K., Taketo, M. M., de Vries, W. N.,
Knowles, B. B., & Solter, D. (2004). Stabilization of β-catenin in the mouse zygote leads to

premature epithelial-mesenchymal transition in the epiblast. *Development*, *131*(23), 5817–
5824. https://doi.org/10.1242/dev.01458

Kim, D., Paggi, J. M., Park, C., Bennett, C., & Salzberg, S. L. (2019). Graph-based genome
alignment and genotyping with HISAT2 and HISAT-genotype. *Nature Biotechnology*, *37*(8), 907–
915. https://doi.org/10.1038/s41587-019-0201-4

Kim, H., Wu, J., Ye, S., Tai, C. I., Zhou, X., Yan, H., Li, P., Pera, M., & Ying, Q. L. (2013). Modulation
of β-catenin function maintains mouse epiblast stem cell and human embryonic stem cell self-
renewal. *Nature Communications*, *4*(61). https://doi.org/10.1038/ncomms3403

Kim, J., & Jung, Y. (2012). Increased aquaporin-1 and Na+-K+-2Cl-cotransporter 1 expression in
choroid plexus leads to blood-cerebrospinal fluid barrier disruption and necrosis of
hippocampal CA1 cells in acute rat models of hyponatremia. *Journal of Neuroscience Research*,
90(7), 1437–1444. https://doi.org/10.1002/jnr.23017

Kim, S., & Jeong, S. (2019). Mutation hotspots in the β-catenin gene: Lessons from the human
cancer genome databases. *Molecules and Cells*, *42*(1), 8–16.
https://doi.org/10.14348/molcells.2018.0436

Konno, D., Iwashita, M., Satoh, Y., Momiyama, A., Abe, T., Kiyonari, H., & Matsuzaki, F. (2012).
The Mammalian DM Domain Transcription Factor Dmrta2 Is Required for Early Embryonic
Development of the Cerebral Cortex. *PLoS ONE*, *7*(10), 1–13.
https://doi.org/10.1371/journal.pone.0046577

Langford, M. B., O'Leary, C. J., Veeraval, L., White, A., Lanoue, V., & Cooper, H. M. (2020).
WNT5a Regulates Epithelial Morphogenesis in the Developing Choroid Plexus. *Cerebral Cortex*.
https://doi.org/10.1093/cercor/bhz330

Lee, S. M., Tole, S., Grove, E., & McMahon, a P. (2000). A local Wnt-3a signal is required for
development of the mammalian hippocampus. *Development*, *127*(3), 457–467.

Li, C., Li, A., Xing, Y., Li, M., Chan, B., Ouyang, R., Taketo, M. M., Kucherlapati, R., Borok, Z., &
Minoo, P. (2013). Apc deficiency alters pulmonary epithelial cell fate and inhibits Nkx2.1 via
triggering TGF-beta signaling. *Developmental Biology*, *378*(1), 13–24.
https://doi.org/10.1016/j.ydbio.2013.03.018

Li, L., Grausam, K. B., Wang, J., Lun, M. P., Ohli, J., Lidov, H. G. W., Calicchio, M. L., Zeng, E.,
Salisbury, J. L., Wechsler-Reya, R. J., Lehtinen, M. K., Schüller, U., & Zhao, H. (2016). Sonic
Hedgehog promotes proliferation of Notch-dependent monociliated choroid plexus tumour
cells. *Nature Cell Biology*, *18*(4), 418–430. https://doi.org/10.1038/ncb3327

Liddelow, S. A., Dziegielewska, K. M., VandeBerg, J. L., & Saunders, N. R. (2010). Development of
the lateral ventricular choroid plexus in a marsupial, Monodelphis domestica. *Cerebrospinal*
Fluid Research, *7*, 1–10. https://doi.org/10.1186/1743-8454-7-16

Louvi, A., Yoshida, M., & Grove, E. A. (2007). The derivatives of the Wnt3a lineage in the central
nervous system. *Journal of Comparative Neurology*, *504*(5), 550–569.
https://doi.org/10.1002/CNE.21461

Love, M. I., Huber, W., & Anders, S. (2014). Moderated estimation of fold change and dispersion
for RNA-seq data with DESeq2. *Genome Biology*, *15*(12), 1–21. https://doi.org/10.1186/s13059-
014-0550-8

Lun, M. P., Johnson, M. B., Broadbelt, K. G., Watanabe, M., Kang, Y. J., Chau, K. F., Springel, M.
W., Malesz, A., Sousa, A. M. M., Pletikos, M., Adelita, T., Calicchio, M. L., Zhang, Y., Holtzman,
M. J., Lidov, H. G. W., Sestan, N., Steen, H., Monuki, E. S., & Lehtinen, M. K. (2015). Spatially
heterogeneous choroid plexus transcriptomes encode positional identity and contribute to
regional CSF production. *Journal of Neuroscience*, *35*(12), 4903–4916.
https://doi.org/10.1523/JNEUROSCI.3081-14.2015

Machon, O., Backman, M., Machonova, O., Kozmik, Z., Vacik, T., Andersen, L., & Krauss, S.
(2007). A dynamic gradient of Wnt signaling controls initiation of neurogenesis in the
mammalian cortex and cellular specification in the hippocampus. *Developmental Biology*,
311(1), 223–237. https://doi.org/10.1016/j.ydbio.2007.08.038

Mangale, V. S., Hirokawa, K. E., Satyaki, P. R. V., Gokulchandran, N., Chikbire, S., Subramanian,
L., Shetty, A. S., Martynoga, B., Paul, J., Mai, M. V., Li, Y., Flanagan, L. A., Tole, S., & Monuki, E. S.
(2008). Lhx2 selector activity specifies cortical identity and suppresses hippocampal organizer
fate. *Science*, *319*(5861), 304–309. https://doi.org/10.1126/science.1151695

MARTIN, Marcel. Cutadapt removes adapter sequences from high-throughput sequencing
reads. EMBnet.journal, [S.l.], v. 17, n. 1, p. pp. 10-12, may 2011. ISSN 2226-6089. Available at:
<https://journal.embnet.org/index.php/embnetjournal/article/view/200/479>. Date accessed:
21 nov. 2020. doi:https://doi.org/10.14806/ej.17.1.200.

Meador, J. P., Lech, J. J., Rice, S. D., Hose, J. E., Short, J. W., Rice, S. D., Collier, T. K., Scholz, N. L.,
Collier, T. K., Scholz, N. L., Campana, S. E., Carlsson, L., Duker, G., Kass, R. S., Roden, D. M.,
Darbar, D., Cheng, H., Lederer, W. J., Freund, E. V, … White, E. (2014). Science Volume 343 issue
6172 2014 [doi 10.1126%2Fscience.1247651] Jaitin, D. A.; Kenigsberg, E.; Keren-Shaul, H.;
Elefant, N.; Pau -- Massively Parallel Single-Cell RNA-Seq for Marker-Free Decomposit.pdf.
February.

Meeker, R. B., Williams, K., Killebrew, D. A. & Hudson, L. C. Cell trafficking through the choroid
plexus. *Cell Adhes. Migr.* **6**, 390–396 (2012).

Monuki, E. S., Porter, F. D., & Walsh, C. A. (2001). *Monuki ES, 2001.pdf*. *32*, 591–604.

Pellegrini, L., Bonfio, C., Chadwick, J., Begum, F., Skehel, M., & Lancaster, M. A. (2020). Human
CNS barrier-forming organoids with cerebrospinal fluid production. *Science*, *369*(6500), 1–21.
https://doi.org/10.1126/science.aaz5626

Sakaguchi, H., Kadoshima, T., Soen, M., Narii, N., Ishida, Y., Ohgushi, M., Takahashi, J., Eiraku,
M., & Sasai, Y. (2015). Generation of functional hippocampal neurons from self-organizing
human embryonic stem cell-derived dorsomedial telencephalic tissue. *Nature Communications*.
https://doi.org/10.1038/ncomms9896

Saunders, N. R., Dziegielewska, K. M., Møllgård, K., Habgood, M. D., Wakefield, M. J., Lindsay,
H., Stratzielle, N., Ghersi-Egea, J. F., & Liddelow, S. A. (2015). Influx mechanisms in the
embryonic and adult rat choroid plexus: A transcriptome study. *Frontiers in Neuroscience*,
9(MAR). https://doi.org/10.3389/fnins.2015.00123

Solberg, N., Machon, O., & Krauss, S. (2008). Effect of canonical Wnt inhibition in the
neurogenic cortex, hippocampus, and premigratory dentate gyrus progenitor pool.
Developmental Dynamics, *237*(7), 1799–1811. https://doi.org/10.1002/dvdy.21586

Speake, T., Kibble, J. D., & Brown, P. D. (2004). Kv1.1 and Kv1.3 channels contribute to the
delayed-rectifying K + conductance in rat choroid plexus epithelial cells. *American Journal of*
Physiology - Cell Physiology, *286*(3 55-3), 611–620. https://doi.org/10.1152/ajpcell.00292.2003

Steffensen, A. B., Oernbo, E. K., Stoica, A., Gerkau, N. J., Barbuskaite, D., Tritsaris, K., Rose, C. R.,
& MacAulay, N. (2018). Cotransporter-mediated water transport underlying cerebrospinal fluid
formation. *Nature Communications*, *9*(1), 1–13. https://doi.org/10.1038/s41467-018-04677-9

Supek, F., Bošnjak, M., Škunca, N., & Šmuc, T. (2011). Revigo summarizes and visualizes long
lists of gene ontology terms. *PLoS ONE*, *6*(7). https://doi.org/10.1371/journal.pone.0021800

Wang, Y., Sabbagh, M. F., Gu, X., Rattner, A., Williams, J., & Nathans, J. (2019). Beta-catenin
signaling regulates barrier-specific gene expression in circumventricular organ and ocular
vasculatures. *ELife*, 1–36. https://doi.org/10.7554/eLife.43257

Watanabe, M., Kang, Y., Davies, L. M., Meghpara, S., Lau, K., Chung, C., Kathiriya, J.,
Hadjantonakis, A., & Monuki, E. S. (2012). *BMP4 Sufficiency to Induce Choroid Plexus Epithelial*
Fate from Embryonic Stem Cell-Derived Neuroepithelial Progenitors. *32*(45), 15934–15945.

https://doi.org/10.1523/JNEUROSCI.3227-12.2012

Ying, Q. L., Wray, J., Nichols, J., Batlle-Morera, L., Doble, B., Woodgett, J., Cohen, P., & Smith, A.
(2008). The ground state of embryonic stem cell self-renewal. *Nature*, *453*(7194), 519–523.
https://doi.org/10.1038/nature06968

Yoshida, M. (2006). Massive loss of Cajal-Retzius cells does not disrupt neocortical layer order.
Development, *133*(3), 537–545. https://doi.org/10.1242/dev.02209

Yu, G., Wang, L. G., Han, Y., & He, Q. Y. (2012). ClusterProfiler: An R package for comparing
biological themes among gene clusters. OMICS A Journal of Integrative Biology, 16(5), 284–287.
https://doi.org/10.1089/omi.2011.0118

Zappaterra, M. W., & Lehtinen, M. K. (2012). The cerebrospinal fluid: Regulator of
neurogenesis, behavior, and beyond. *Cellular and Molecular Life Sciences*, *69*(17), 2863–2878.
https://doi.org/10.1007/s00018-012-0957-x

Zhao Y, Sheng HZ, Amini R, Grinberg A, Lee E, Huang S, Taira M, Westphal H. Control of
hippocampal morphogenesis and neuronal differentiation by the LIM homeobox gene Lhx5.
Science. 1999; 284:1155–1158