Does arsenic contamination affect DNA methylation patterns in a wild bird population? An

experimental approach

Veronika N. Laine[1], Mark Verschuuren[1], Kees van Oers[1], Silvia Espín[2,3], Pablo Sánchez-

Virosta[2,3], Tapio Eeva[3], Suvi Ruuskanen[3]

[1]Department of Animal Ecology, Netherlands Institute of Ecology (NIOO-KNAW),

Wageningen, The Netherlands

[2]Area of Toxicology, Department of Socio-Sanitary Sciences, University of Murcia, Spain

[3]Department of Biology, University of Turku, Finland

Abstract

Pollutants, like toxic metals, negatively influence organismal health and performance, even

leading to population collapses. Studies in model organisms have shown that epigenetic

marks, such as DNA methylation, can be modulated by various environmental factors,

including pollutants, influencing gene expression and various organismal traits. Yet

experimental data on the effects of pollution on DNA methylation from wild animal

populations is largely lacking. We here investigated experimentally for the first time the

effects of early-life exposure to environmentally relevant levels of a key pollutant, arsenic

(As) on genome-wide DNA methylation status in a wild bird population. We experimentally

exposed nestlings of great tits (*Parus major*) to arsenic during their post-natal developmental

period (3 to 14 days post-hatching) and compared their DNA methylation levels to those of

respective controls. In contrast to predictions, we found no overall hypomethylation in the

arsenic group. We found evidence for loci to be differentially methylated between the

treatment groups, but for five CpG sites only. This may be explained by the heterogenous

environment, relatively low dose (compared to previous laboratory studies), or choice of

tissue. Further studies are needed to evaluate whether epigenetic dysregulation is a commonly

observed phenomena in polluted populations, and what are its consequences for organism

functioning and for populations.

Keywords: pollution, *Parus major*, environmental epigenetics, ecological epigenetics,

ecotoxicology

Running title: Arsenic pollution and DNA methylation in the wild

## Introduction

Environmental pollution can negatively affect organisms at multiple level of organization, from molecular and physiological level to performance, and even lead to population collapses (Dudgeon et al., 2006; McCrink-Goode, 2014; Saaristo et al., 2018; Vos et al., 2000). In wild populations, a largely unexplored mechanism mediating such pollution effects is the potential influence of the epigenome, such as DNA methylation. In human and animal models, it has been discovered that methylation patterns can be changed by various environmental factors, including metal and organic pollutants and other early-life stressors (reviewed by Baccarelli & Bollati, 2009; Cheng, Choudhuri, & Muldoon-Jacobs, 2012; Head, 2014; Head, Dolinoy, & Basu, 2012; Ray, Yosim, & Fry, 2014; Ruiz-Hernandez et al., 2015; Sharavanan et al.). DNA methylation is the addition of a methyl (-CH3) group to the 5' carbon site of cytosines catalyzed by DNA-methyltransferases, and is generally found to be negatively associated with gene expression (Goldberg, Allis, & Bernstein, 2007). Variation in DNA methylation is linked to variation in phenotypes and behavior, and associated with the prevalence of various diseases, including cancers in humans and model animals (Rosenfeld, 2010; Salameh, Bejaoui, & El Hajj, 2020; Skinner, Manikkam, & Guerrero-Bosagna, 2010). Epigenetic changes from early-life environment may persist and affect health throughout life-time and may even be transmitted to future generations (Horsthemke, 2018), which could potentially contribute to explaining delayed or persistent effects of pollutants (e.g. Head 2014). Yet the effects of pollutants on the epigenome have hardly been explored, and epigenetic research in wild animal populations is only emerging (Bossdorf, Richards, & Pigliucci, 2008; Head et al., 2012; Laine et al., 2016; Liebl, Schrey, Richards, & Martin, 2013; Riyahi, Sanchez-Delgado, Calafell, Monk, & Senar, 2015; Rubenstein et al., 2016; Sepers et al., 2019; Verhoeven, Vonholdt, & Sork, 2016; Wenzel & Piertney, 2014)

Arsenic (As) is a global, persistent pollutant, distributed in the environment due to

natural and anthropogenic sources such as mining, industrial activities or coal combustion

(Mandal & Suzuki, 2002) and the most highly ranked hazardous substances for animals and

plants (ATSDR, 2019). Across organisms, arsenic can have negative consequences for

basically all organ systems, often via causing oxidative stress, i.e. the imbalance between

harmful reactive oxygen species (ROS) and antioxidant defenses, and cancer (Jomova et al.,

2011; Sánchez-Virosta, Espín, García-Fernández, & Eeva, 2015).

Arsenic has been repeatedly observed to also modulate patterns of DNA methylation

*in vitro*, in laboratory animal models (with large doses) and in studies on human populations

(Alegria-Torres et al., 2016; Bjorklund, Aaseth, Chirumbolo, Urbina, & Uddin, 2018; Jensen,

Novak, Wnek, Gandolfi, & Futscher, 2009). The major pathway for arsenic to change the

DNA methylation of a cytosine, is via the depletion of the cellular availability of methyl

groups, since biotransformation of arsenic to less toxic forms includes the addition of methyl

group(s) (Minatel et al., 2018). The main methyl donor during the methylation of both arsenic

and DNA is s-adenosylmethionine (SAM). The high demand imposed on this molecule

during the biotransformation process can then lead to global DNA hypomethylation, as

shown in multiple biomedical studies (Baccarelli & Bollati, 2009; Hubaux et al., 2013).

Additionally, ROS created during arsenic biotransformation have been suggested to influence

DNA methylation by creating aberrant modifications (Ziech, Franco, Pappa, & Panayiotidis,

2011). Pre/postnatal exposure to arsenic is associated with epigenetic modifications related to

early onset of diseases, which could have long-term consequences (reviewed in Bantock,

Prys-Jones, & Lee, 2008; Gangopadhyay, Sharma, Chauhan, & Srivastava, 2019). For

example, prenatal arsenic exposure led to global hypomethylation of inflammatory and tumor

suppressor genes (Kile et al., 2014) and interfered with de novo methylation (Broberg et al.,

2014) in humans. Despite the extensive data on model animals and humans, the potential

effects of environmental arsenic on wild animals via epigenetic dysregulation has not been

studied up to date.

We here investigated the effects of experimental early-life (post-natal) exposure to

arsenic on genome-wide DNA methylation status in a wild population of great tits (*Parus*

*major*). To our knowledge, this is the first study on (i) environmentally relevant arsenic

exposure, and (ii) in a wild population, on epigenetic markers. We used a bird model, since

birds have been successfully used in biomonitoring of pollution and its effects (e.g. García-

Fernández, 2014). Arsenic exposure has been reported to negatively affect multiple fitness-

related traits (growth, physiology, behavior and even egg-laying) in several birds species

(reviewed in Sánchez-Virosta et al., 2015). For great tits specifically, we have previously

reported (using the same experimental protocol) that arsenic exposure increased nestling

mortality, reduced wing growth (Sánchez-Virosta et al., 2018) and decreased an intracellular

antioxidant, catalase, in nestlings (Sánchez-Virosta et al., 2020). More specifically, we here

experimentally exposed nestlings in non-polluted sites to environmentally relevant levels (1

µg/g body mass) of dietary arsenic during the entire post-hatching growth period, and

compared their DNA methylation levels to respective controls. We used reduced

representation bisulfite sequencing (RRBS) to assess genome-wide methylation and

characterized differential methylation across CpG sites between the experimental and the

control group. We predict that arsenic exposure will lead to genome-wide hypomethylation,

and potentially specifically on gene/hubs related to development.

**Methods**

*Arsenic treatment protocol and sampling*

The study was conducted in a nest box population of great tits (*Parus major),* a small

passerine bird and a popular model species in ecological and evolutionary research.

Importantly, as one of the only non-domesticated bird species, both the genome and

methylome are available (Derks et al., 2016; Laine et al., 2016; Verhulst et al., 2016).

The arsenic exposure, dosages and sampling are described in detail in Sánchez-

Virosta et al. (2020; 2018). In short, the experiment was conducted in a nest-box population

in western Finland, with known history of very low pollution levels (Eeva, Ryömä, &

Riihimaki, 2005). Breeding was monitored, and from day 3 after hatching until day 14 whole

broods were subjected to daily oral dosing with the following treatments: arsenic treatment (1

µg As/g bodymass, N = 16 broods) or control treatment (distilled water, N = 16 broods). The

dose aimed to represent environmentally relevant exposure levels occurring in polluted areas

in Europe: It was estimated combining data from several sources, such as (i) the lowest-

observed-adverse-effect level for different effects on mammals, (ii) fecal arsenic levels

reported for great tits at some metal polluted sites (reviewed in Sánchez-Virosta et al., 2015),

(iii) arsenic concentrations of food items (moth larvae, spiders and beetles) collected directly

from parent great tits feeding their nestlings in the polluted area (Sánchez-Virosta et al.,

2018, 2020), and (iv) a pilot experiment Sánchez-Virosta et al., 2018). Fecal matter was

sampled 8 days after hatching for metal analyses (see below). DNA methylation was

analyzed from red blood cells (RBCs) collected 14 days after hatching to avoid sacrificing the

individuals. Ten samples from the arsenic and ten from control treatment were randomly

selected for the DNA methylation analyses. These included five females and five males from

each treatment (molecularly sexed, following Bantock et al., 2008). Only one nestling per

nest was selected to avoid pseudoreplication. The experiment was conducted under licenses

from the Animal Experiment Committee of the State Provincial Office of Southern Finland

(license number ESAVI/11579/04.10.07/2014) and the Centre for Economic Development,

Transport and the Environment, ELY Centre Southwest Finland (license number

VARELY/593/2015).

*Metal analyses*

For detailed analyses, see (Sánchez-Virosta et al., 2018). Briefly, in both experimental

groups, two samples (one male and one female) from the same brood were combined to

assess brood level metal exposure (total N = 32 broods). We determined the concentrations of

As, but also other metals to confirm that the levels of other metals were low and similar

across the treatment groups (see Sánchez-Virosta et al., 2018). The determination of

pollutants was conducted with ICP-OES with detection limit of 1 ppt (ng/l) and below.

Calibration standards and certified reference materials were used for method validation. The

levels of other measured metals (Al, Pb, Ni, Zn, Mg, Mn, Fe, Cu) were low, and did not differ

among the treatment groups (all t <0.88, all p <0.38).

*DNA isolation*

DNA isolation was performed at the Center of Evolutionary Applications (University of

Turku, Finland). We used RBCs given that previous studies suggest that blood shows similar

methylation patterns as brain tissue in the study species (e.g. 80% similarity between brain

and blood methylation in CpGs Derks et al., 2016; Verhulst et al., 2016). DNA was extracted

from 10-20 µl RBCs using the salt extraction method modified from Aljnabi & Martinez

(1997). Extracted DNA was treated with RNase-I according to the manufacturer's protocol.

DNA concentration was measured fluorometrically with a Qubit High Sensitivity kit

(ThermoFisher Scientific) and we assessed DNA integrity by running each DNA sample on

an agarose gel.

***RRBS library preparation***

We used a reduced representation bisulfite sequencing (RRBS) approach, which enriches for

regions of the genome that have a high CpG content. It was previously shown in the study

species that a vast majority of methylated Cs (97%) were derived from CpG sites in blood

(Derks et al., 2016). Sequencing was conducted at the Finnish Microarray and Sequencing

Center in Turku, Finland. The library preparation was started from 200 ng of genomic DNA

and was carried out according to a protocol adapted from Boyle et al. (2012). The first step in

the workflow involved the fragmentation of genomic DNA with MspI where the cutting

pattern of the enzyme (C^CGG) was used to systematically digest DNA to enrich for CpG

dinucleotides. After a fragmentation step a single reaction was carried out to end repair and

A-tail (required for the adapter ligation) the MspI digested fragments using Klenow fragment

(3' => 5' exo) following the purification of A-tailed DNA with bead SPRI clean-up method

(AMPure magnetic beads). A unique Illumina TruSeq indexing adapter was then ligated to

each sample during adapter ligation step to be able to identify pooled samples of one flow

cell lane. To reduce the occurrence of adapter dimers, a lower concentration of adapters (1:10

dilution) was used than recommended by the manufacturer. These ligated DNA fragments

were purified with bead SPRI clean-up method before putting samples through bisulfite

conversion to achieve C-to-U conversion of unmethylated cytosines, whereas methylated

cytosines remain intact. Bisulfite conversion and sample purification were done according to

Invitrogen MethylCode Bisulfite Conversion Kit. Aliquots of converted DNA were amplified

by PCR (16 cycles) with Taq/Pfu Turbo Cx Polymerase, a proofreading PCR enzyme that

does not stall when it encounters uracil, the product of the bisulfite reaction, in the template.

PCR-amplified RRBS libraries were purified using two subsequent rounds of SPRI

beadclean-ups to minimize primer dimers in the final libraries. The high quality of the

libraries was confirmed with Advanced Analytical Fragment Analyzer and the concentrations

of the libraries were quantified with Qubit® Fluorometric Quantitation, Life Technologies.

We used an average fragment size of 250‑350 bp for sequencing.

***Sequencing***

The samples were normalized and pooled for the automated cluster preparation which was

carried out with Illumina cBot station. The 20 libraries were combined in two pools, 10

samples in each pool (treatments and sexes equally distributed between the pools) and

sequenced in two lanes. The samples were sequenced with an Illumina HiSeq 2500

instrument using TruSeq v3 sequencing chemistry. Paired‑end sequencing with 2 x 100 bp

read length was used with 6 bp index run.

***Sequence data processing and differential methylation expression analysis***

All the reads were checked for quality using FastQC (Babraham Bioinformatics) with

multiQC (Ewels, Magnusson, Lundin, & Kaller, 2016), and low-quality sequences were

trimmed with Trim Galore v. 0.4.4 (Brabraham Bioinformatics) by using --quality 20 --paired

--rrbs settings.

The trimmed reads were mapped to the *Parus major* reference genome build 1.1.

(https://www.ncbi.nlm.nih.gov/assembly/ GCF_001522545.2) using Bismark (Krueger &

Andrews, 2011) with default parameters. Methylation calling was conducted with Bismark,

first with default settings with paired-end mode and overlap removal (--p --no_overlap). After

this first calling round, we observed a methylation bias for the samples by plotting the

methylation proportion across each possible position in the read. Based on the plotting, the

three and two first bases of R1 and R2 respectively of the 5' prime end were omitted and the

first base in the R2 3' prime end was also omitted in the final methylation calling. Thereafter,

Methylkit (Akalin et al., 2012) implemented in R was used for filtering and differential

methylation analysis. We discarded bases that had coverage below 10x. To avoid a possible

PCR bias we also discarded bases that had more than 99.9th percentile of coverage in each

sample. Before differential methylation analysis we merged read counts from reads covering

both strands of a CpG dinucleotide and CpGs needed to be covered with at least 8 samples

per group (control and treatment).

Samples were thereafter clustered based on the similarity of their overall methylation

profile by (i) using the clustering method ward.D in Methylkit's clusterSamples -function and

(ii) using principal component analysis (PCA) with Methylkit's PCASamples -function. We

also checked for lane and sex effect by using Methylkit's assocComp -function where it

checks which principal components are statistically associated with the potential batch effects

such as the used lane and sex of the individuals. For the former, no missing data is allowed,

thus we created a separate data object where all the individuals needed to be covered.

For analyzing differential methylation of GpC sites between control and arsenic

treatment we used the beta-binomial model from DSS package (Feng, Conneely, & Wu,

2014) which is also included in Methylkit (calculateDiffMethDSS - function). DSS calculates

the differential methylation statistics using a beta-binomial model with parameter shrinkage.

Bonferroni correction was applied to account for multiple testing.

**Results**

*Arsenic exposure*

As reported in Sánchez -Virosta et al. (2020; 2018), dietary arsenic treatment successfully

increased arsenic load as fecal arsenic levels were on average 10 times higher in arsenic

exposure compared to control group (average±SD ppm: control 0.51±0.50, As exposure

4.92±4.57, $t_{15.4}$ = -3.83, p = 0.0015), and increased levels were also found in internal tissues.

*Sequencing and mapping*

The total number of read pairs was 341 million (Supplementary Table 1), varying from 14

million to 20 million per individual. After QC filtering the final number of read pairs was 337

million (Supplementary Table 1). The RRBS individual sequencing data have been deposited

in NCBI (Number will be added later). Mapping efficiency was on average 46.15% and on

average 3.1 million cytosines were covered before 10x coverage and percentile filtering.

After filtering, 1.3 million cytosines were identified in CpG context. When combining the Cs

from both strands and restricting our data to at least 8 individuals per group to be covered, we

ended up having 652 655 CpGs.

*Sample clustering and differential methylation*

Both the ward.D and PCA clustering methods showed that sample 14 was an outlier in its

methylation profile (Supplementary Figure 1). That particular sample also had a low number

of reads and showed lower duplication levels (Supplementary Table 1) and we therefore

decided to exclude this sample from further analysis. No lane effect was detected, but PC3

was associated with sex after Bonferroni correction (Supplementary Table 2, Supplementary

Figure 2), mostly driven by two samples, ctrl_3F and test_16F, since after removing these

two female samples from the data, PC3 was not significant anymore. Furthermore, when

removing the PC3 from the data, three CpG sites were significant in the differential

methylation analysis done with DSS: two of them were the same as when including all the

PCs (see below, Table 1). The three other significant sites found below were not covered by

all individuals as required in this PC-removal analysis.

In the differential methylation analysis when including all the PCs, five CpG sites showed a

significant difference in methylation level with a q-value below 0.05 and percent methylation

difference larger than 10% (Table 1, Figure 1, Supplementary Table 3). Lambda estimation

was close to 1 ($\lambda = 0.747$, SE 0.000136) (Supplementary Figure 3), suggesting no systematic

biases ($\lambda > 1$ indicates bias). Four of these sites were hypermethylated (higher methylation in

the arsenic treatment group) and one was hypomethylated (higher methylation in the control

group). Three of the sites were located in gene bodies, namely zinc finger and BTB domain

containing 47 (*ZBTB47*), HIVEP zinc finger 3 (*HIVEP3*) and insulin like growth factor 2

mRNA binding protein 1 (*IGF2BP1*) based on NCBI *P. major* annotation report 102.

**Table 1.** The differentially methylated CpG sites between arsenic exposed and control

individuals. Methylation diff% refers to the methylation difference, comparing arsenic

exposed to control group. Positive values therefore indicate hypermethylation in the arsenic

treatment group compared to the control group. PC3 indicates sites that were significant after

PC3 removal.

| Chr | Chr Genbank | Position | P-value | q-value | Methylation diff % | Gene | PC3 |
|---|---|---|---|---|---|---|---|
| 2 | NC_031769.1 | 2,448,788 | 1.00E-10 | 6.53E-05 | 36.40 | *ZBTB47* | |
| 12 | NC_031781.1 | 9,949,364 | 1.23E-08 | 8.05E-03 | 57.66 | - | |
| 23 | NC_031791.1 | 5,408,232 | 3.43E-08 | 2.24E-02 | 42.66 | *HIVEP3* | x |
| UN | NW_015379267.1 | 107,660 | 7.99E-10 | 5.21E-04 | 60.43 | *IGF2BP1* | |
| UN | NW_015379318.1 | 39,910 | 1.42E-08 | 9.27E-03 | -52.66 | - | x |
| **Only in PC3 removed** | | | | | | | |
| 14 | NC_031783.1 | 14,025,702 | 1.34E-07 | 0.042 | 27.09 | - | x |

**Figure 1.** Plots of significance of CpG sites from the differential methylation analysis

conducted with DSS implemented in Methylkit. (a) Manhattan plot with the significance of

differential methylation of the arsenic treatment against the control pool, against the great tit

reference genome version 1.1. The orange line depicts the genome-wide threshold based on a

Bonferroni correction: 7.11. (b) A volcano plot of the significance against the absolute

difference in methylation between the two pools, with delta methylation is arsenic treatment –

control. Orange points are the genome-wide significant sites after Bonferroni correction and

filtering for Delta methylation >10%.

**Discussion**

We studied for the first time whether early-life exposure to environmentally relevant levels of

experimental arsenic affects DNA methylation in a wild vertebrate population. The

experimental treatment increased arsenic levels significantly, but contrary to predictions, did

not lead to overall hypomethylation. We did find that treated individuals showed

hypermethylation in four and hypomethylation in one CpG site, indicating that increased

levels of arsenic exposure may affect methylation at specific parts of the genome only. Yet

also at these sites, the assumption of general hypomethylation was not met.

The lack of overall or site-specific hypomethylation may be explained by various

factors: first, contrary to our predictions, the methyl donor s-adenosylmethionine needed for

methylation may not have been limiting, potentially because oxidative status was not altered

dramatically in all individuals. Indeed, as reported from the exact same experiment and

samples by Sánchez-Virosta et al. (2020), most biomarkers of oxidative status and damage in

blood were only slightly (but not statistically significantly) elevated, and only antioxidant

enzyme catalase showed significant decrease. In the future, sampling before and after

exposure to e.g. pollutants may be advised to associate DNA methylation changes directly to

changes in oxidative status, for example in adult birds (in contrast to developing animals

where measurements are confounded by the changes in growth and associated changes in

physiology).

Second, the response is likely to depend on the tissue type studied. For example,

global hypomethylation in response to arsenic exposure is not consistently reported in blood:

in humans, where blood leucocytes have been used to characterize arsenic associated changes

no evidence for global hypo or hypermethylation was detected, yet arsenic was repeatedly

reported to induce hypermethylation in various genes (especially promoters) (Koestler,

Avissar-Whiting, Houseman, Karagas, & Marsit, 2013), whereas global hypomethylation was

detected in hepatic cells (Chen et al., 2001). Given that arsenic metabolism and SAM

production mostly takes part in liver, we may expect tissue-dependent hypomethylation

especially in liver, but not necessarily in other tissues. Unfortunately, we lack oxidative status

measurements from the liver in this experiment. Thus measuring methylation levels from red

blood cells might not be ideal in this case. Although studies have shown correlating

methylation values between different tissue (e.g. between absolute blood methylation levels

in blood and absolute methylation levels in liver, kidney and brain  Derks et al., 2016;

McKay et al., 2011), tissue-specific methylation patterns were more prominent for genes that

are expressed in a tissue-specific way (Derks et al., 2016).

Furthermore, contrary to many previous studies in laboratory animals, this experiment

was conducted with relatively low doses, mimicking exposure in polluted environments,

whereas effects via SAM may only be apparent when levels are higher. Finally as the

experiment was conducted in a wild population, in comparison to previous studies in

laboratory, the environmental variability and potential variability across sexes may have

masked some effects of the experimental treatments. The effects of (low) pollution may

further be evident only if there are other concurrent stressors.

We could annotate three of the five differentially methylated sites to genes. One of the

genes, *IGF2BP1* is especially interesting as it is associated with development and growth: it

has been showed that *IGF2BP1* plays important roles in various aspects of cell function, such

as cell proliferation, differentiation, migration, morphology and metabolism (Chen et al.,

2019; Hansen et al., 2004) but also embryogenesis and potentially even arsenic-related

carcinogenesis (Huang et al., 2018; Noubissi, Yedjou, Spiegelman, & Tchounwou, 2018).

*IGF2BP1* is abundantly expressed in fetal and neonatal tissues (Hansen et al., 2004).

Furthermore, two of the genes, *ZBTB47* and *HIVEP3* are both zinc-finger domains and are

associated with transcriptional regulation (Hicar, Liu, Allen, & Wu, 2001). Epigenetic

regulation of both *ZBTB47* and *HIVEP3* is known to be associated cancer (Lubecka et al.,

2018; Sarkar, Leung, Baguley, Finlay, & Askarian-Amiri, 2015)

All the three gene related differentially methylated CpG sites were found in

the gene body region, in both intron (*IGF2BP1*) and in exons (*ZBTB47* and *HIVEP3*).

Hypermethylation at CpG sites at promoter regions represses transcription of genes which is

a well-known mechanism operating in many scenarios. DNA methylation at intergenic

regions and gene bodies and its impact on gene expression is gaining more attention

especially in cancer studies (Jones, 2012). Interestingly, a recent study done with corals

showed that gene body methylation was altered by the environmental factors, which

facilitated acclimatization and adaptation to different habitats (Dixon, Liao, Bay, & Matz,

2018). However, in great tits the DNA methylation observed in gene body CpGs does not

seem to affect gene expression (Derks et al., 2016) thus future studies are needed to

determine the role of gene body methylation in gene expression control.

In conclusion, our study shows that early-life exposure to a toxic metal, arsenic, potentially

affects fitness via DNA methylation changes in specific pathways, but not via an overall

hypomethylation in the red blood cells. The effect might more profound in other tissues that

are more relevant to arsenic metabolism, such as liver. Thus, future studies should inspect

also other tissues as well. Other pathways of epigenetic alterations, known to be subject to

arsenic-related alternations in vitro, such as histone acetylation (Jensen (Jensen et al., 2009)

and micro-RNAs (Cao et al., 2011) could be further explored.

ASSOCIATED CONTENT

**Figure S1.** Clustering of samples based on ward.D (a) and principal component analysis

(PCA) (b) in Methylkit

**Table S1**. Number of reads before and after read trimming with mapping and methylation

calling success of Bismark. CpG site filtering was done with Methylkit.

**Table S2.** Results from the tests on the effects of sequencing lane and sex.

**Table S3**. A full description of all differentially methylated sites.

AUTHOR INFORMATION

Corresponding author: Suvi Ruuskanen (skruus@utu.fi)

**Conflict of interest**

We have no conflict of interest to declare.

**Data accessibility**

Data will be deposited in Dryad and in Genbank upon acceptance.

**Author contributions**

SR, SE, PSV, VNL, MV, KvO and TE designed the study. TE, SE and PSV collected the

data. SR and KvO designed the sequencing. VNL and MV conducted the bioinformati

analyses. SE and PSV conducted metal analyses. KvO and VNL provided the genome

resources. VNL, MV, KvO and SR interpreted the data. VNL and SR wrote the first draft. All

authors contributed to writing the manuscript.

**Funding sources**

TE, SE, PSV were funded by Academy of Finland (to TE), SR was also funded by Academy

of Finland and Turku University Foundation.

**Acknowledgements**

We thank Miia Rainio and Jorma Nurmi for their efforts in helping us with field work and Dr.

Fleur Gawehns-Bruning and William Sies for assistance with the methylation analysis. We

also thank the Center for Evolutionary Applications for molecular work and Finnish

Functional Genomics Centre for sequencing services. Our study was financed by KONE

foundation (SR) and Academy of Finland (TE: project 265859).

**References**

Akalin, A., Kormaksson, M., Li, S., Garrett-Bakelman, F. E., Figueroa, M. E., Melnick, A., & Mason, C. E.
(2012). methylKit: a comprehensive R package for the analysis of genome-wide DNA
methylation profiles. *Genome Biology, 13*(10), 9. doi:10.1186/gb-2012-13-10-R87
Alegria-Torres, J. A., Carrizales-Yanez, L., Diaz-Barriga, F., Rosso-Camacho, F., Motta, V., Tarantini, L.,
& Bollati, V. (2016). DNA Methylation Changes in Mexican Children Exposed to Arsenic From
Two Historic Mining Areas in San Luis Potosi. *Environmental and Molecular Mutagenesis,*
*57*(9), 717-723. doi:10.1002/em.22062
Aljanabi, S. M., & Martinez, I. (1997). Universal and rapid salt-extraction of high quality genomic DNA
for PCR-based techniques. *Nucleic Acids Research, 25*(22), 4692-4693.
doi:10.1093/nar/25.22.4692
ATSDR. (2019). The ATSDR 2019 Substance Priority List. In: Agency for Toxic Substances and Disease
Registry, U.S. Public Health Service, U.S.A.

Baccarelli, A., & Bollati, V. (2009). Epigenetics and environmental chemicals. *Current Opinion in*
*Pediatrics, 21*(2), 243-251. doi:10.1097/MOP.0b013e32832925cc

Bantock, T. M., Prys-Jones, R. P., & Lee, P. L. M. (2008). New and improved molecular sexing
methods for museum bird specimens. *Molecular Ecology Resources, 8*(3), 519-528.
doi:10.1111/j.1471-8286.2007.01999.x
Bjorklund, G., Aaseth, J., Chirumbolo, S., Urbina, M. A., & Uddin, R. (2018). Effects of arsenic toxicity
beyond epigenetic modifications. *Environmental Geochemistry and Health, 40*(3), 955-965.
doi:10.1007/s10653-017-9967-9
Bossdorf, O., Richards, C. L., & Pigliucci, M. (2008). Epigenetics for ecologists. *Ecology Letters, 11*(2),
106-115. doi:10.1111/j.1461-0248.2007.01130.x
Boyle, P., Clement, K., Gu, H. C., Smith, Z. D., Ziller, M., Fostel, J. L., . . . Meissner, A. (2012). Gel-free
multiplexed reduced representation bisulfite sequencing for large-scale DNA methylation
profiling. *Genome Biology, 13*(10), 10. doi:10.1186/gb-2012-13-10-R92
Broberg, K., Ahmed, S., Engstrom, K., Hossain, M. B., Mlakar, S. J., Bottai, M., . . . Vahter, M. (2014).
Arsenic exposure in early pregnancy alters genome-wide DNA methylation in cord blood,
particularly in boys. *Journal of Developmental Origins of Health and Disease, 5*(4), 288-298.
doi:10.1017/s2040174414000221
Cao, Y., Yu, S. L., Wang, Y., Guo, G. Y., Ding, Q. A., & An, R. H. (2011). MicroRNA-dependent
regulation of PTEN after arsenic trioxide treatment in bladder cancer cell line T24. *Tumor
Biology, 32*(1), 179-188. doi:10.1007/s13277-010-0111-z
Chen, H., Liu, J., Zhao, C. Q., Diwan, B. A., Merrick, B. A., & Waalkes, M. P. (2001). Association of c-
myc overexpression and hyperproliferation with arsenite-induced malignant transformation.
*Toxicology and Applied Pharmacology, 175*(3), 260-268. doi:10.1006/taap.2001.9253
Chen, J. H., Ren, X. Y., Li, L. M., Lu, S. Y., Chen, T., Tan, L. T., . . . Luo, W. (2019). Integrative Analyses
of mRNA Expression Profile Reveal the Involvement of IGF2BP1 in Chicken Adipogenesis.
*International Journal of Molecular Sciences, 20*(12), 15. doi:10.3390/ijms20122923
Cheng, T. F., Choudhuri, S., & Muldoon-Jacobs, K. (2012). Epigenetic targets of some toxicologically
relevant metals: a review of the literature. *Journal of Applied Toxicology, 32*(9), 643-653.
doi:10.1002/jat.2717
Derks, M. F. L., Schachtschneider, K. M., Madsen, O., Schijlen, E., Verhoeven, K. J. F., & van Oers, K.
(2016). Gene and transposable element methylation in great tit (Parus major) brain and
blood. *Bmc Genomics, 17*, 13. doi:10.1186/s12864-016-2653-y
Dixon, G., Liao, Y., Bay, L. K., & Matz, M. V. (2018). Role of gene body methylation in acclimatization
and adaptation in a basal metazoan. *Proceedings of the National Academy of Sciences of the
United States of America, 115*(52), 13342-13346. doi:10.1073/pnas.1813749115
Dudgeon, D., Arthington, A. H., Gessner, M. O., Kawabata, Z. I., Knowler, D. J., Leveque, C., . . .
Sullivan, C. A. (2006). Freshwater biodiversity: importance, threats, status and conservation
challenges. *Biological Reviews, 81*(2), 163-182. doi:10.1017/s1464793105006950
Eeva, T., Ryoma, M., & Riihimaki, J. (2005). Pollution-related changes in diets of two insectivorous
passerines. *Oecologia, 145*(4), 629-639. doi:10.1007/s00442-005-0145-x
Ewels, P., Magnusson, M., Lundin, S., & Kaller, M. (2016). MultiQC: summarize analysis results for
multiple tools and samples in a single report. *Bioinformatics, 32*(19), 3047-3048.
doi:10.1093/bioinformatics/btw354
Feng, H., Conneely, K. N., & Wu, H. (2014). A Bayesian hierarchical model to detect differentially
methylated loci from single nucleotide resolution sequencing data. *Nucleic Acids Research,
42*(8), 11. doi:10.1093/nar/gku154
Gangopadhyay, S., Sharma, V., Chauhan, A., & Srivastava, V. (2019). Potential facet for prenatal
arsenic exposure paradigm: linking endocrine disruption and epigenetics. *Nucleus-India,
62*(2), 127-142. doi:10.1007/s13237-019-00274-3
Goldberg, A. D., Allis, C. D., & Bernstein, E. (2007). Epigenetics: A landscape takes shape. *Cell, 128*(4),
635-638. doi:10.1016/j.cell.2007.02.006
Hansen, T. V. O., Hammer, N. A., Nielsen, J., Madsen, M., Dalbaeck, C., Wewer, U. M., . . . Nielsen, F.
C. (2004). Dwarfism and impaired gut development in insulin-like growth factor II mRNA-

binding protein 1-deficient mice. *Molecular and Cellular Biology, 24*(10), 4448-4464.
458             doi:10.1128/mcb.24.10.4448-4464.2004

459 Head, J. A. (2014). Patterns of DNA Methylation in Animals: An Ecotoxicological Perspective.
460             *Integrative and Comparative Biology, 54*(1), 77-86. doi:10.1093/icb/icu025

461 Head, J. A., Dolinoy, D. C., & Basu, N. (2012). Epigenetics for ecotoxicologists. *Environmental*
462             *toxicology and chemistry / SETAC, 31*(2), 221-227. doi:10.1002/etc.1707

463 Hicar, M. D., Liu, Y. L., Allen, C. E., & Wu, L. C. (2001). Structure of the human zinc finger protein
464             HIVEP3: Molecular cloning, expression, exon-intron structure, and comparison with
465             paralogous genes HIVEP1 and HIVEP2. *Genomics, 71*(1), 89-100.
466             doi:10.1006/geno.2000.6425

467 Horsthemke, B. (2018). A critical view on transgenerational epigenetic inheritance in humans. *Nature*
468             *Communications, 9*. doi:10.1038/s41467-018-05445-5

469 Huang, X. W., Zhang, H., Guo, X. R., Zhu, Z. X., Cai, H. B., & Kong, X. Y. (2018). Insulin-like growth
470             factor 2 mRNA-binding protein 1 (IGF2BP1) in cancer. *Journal of Hematology & Oncology, 11*,
471             15. doi:10.1186/s13045-018-0628-y

472 Hubaux, R., Becker-Santos, D. D., Enfield, K. S. S., Rowbotham, D., Lam, S., Lam, W. L., & Martinez, V.
473             D. (2013). Molecular features in arsenic-induced lung tumors. *Molecular Cancer, 12*, 11.
474             doi:10.1186/1476-4598-12-20

475 Jensen, T. J., Novak, P., Wnek, S. M., Gandolfi, A. J., & Futscher, B. W. (2009). Arsenicals produce
476             stable progressive changes in DNA methylation patterns that are linked to malignant
477             transformation of immortalized urothelial cells. *Toxicology and Applied Pharmacology,*
478             *241*(2), 221-229. doi:10.1016/j.taap.2009.08.019

479 Jomova, K., Jenisova, Z., Feszterova, M., Baros, S., Liska, J., Hudecova, D., . . . Valko, M. (2011).
480             Arsenic: toxicity, oxidative stress and human disease. *Journal of Applied Toxicology, 31*(2),
481             95-107. doi:10.1002/jat.1649

482 Jones, P. A. (2012). Functions of DNA methylation: islands, start sites, gene bodies and beyond.
483             *Nature Reviews Genetics, 13*(7), 484-492. doi:10.1038/nrg3230

484 Kile, M. L., Houseman, E. A., Baccarelli, A. A., Quamruzzaman, Q., Rahman, M., Mostofa, G., . . .
485             Christiani, D. C. (2014). Effect of prenatal arsenic exposure on DNA methylation and
486             leukocyte subpopulations in cord blood. *Epigenetics, 9*(5), 774-782. doi:10.4161/epi.28153

487 Koestler, D. C., Avissar-Whiting, M., Houseman, E. A., Karagas, M. R., & Marsit, C. J. (2013).
488             Differential DNA Methylation in Umbilical Cord Blood of Infants Exposed to Low Levels of
489             Arsenic in Utero. *Environmental Health Perspectives, 121*(8), 971-977.
490             doi:10.1289/ehp.1205925

491 Krueger, F., & Andrews, S. R. (2011). Bismark: a flexible aligner and methylation caller for Bisulfite-
492             Seq applications. *Bioinformatics, 27*(11), 1571-1572. doi:10.1093/bioinformatics/btr167

493 Laine, V. N., Gossmann, T. I., Schachtschneider, K. M., Garroway, C. J., Madsen, O., Verhoeven, K. J.
494             F., . . . Great Tit HapMap, C. (2016). Evolutionary signals of selection on cognition from the
495             great tit genome and methylome. *Nature Communications, 7*. doi:10.1038/ncomms10474

496 Liebl, A. L., Schrey, A. W., Richards, C. L., & Martin, L. B. (2013). Patterns of DNA Methylation
497             Throughout a Range Expansion of an Introduced Songbird. *Integrative and Comparative*
498             *Biology, 53*(2), 351-358. doi:10.1093/icb/ict007

499 Lubecka, K., Flower, K., Beetch, M., Qiu, J., Kurzava, L., Buvala, H., . . . Stefanska, B. (2018). Loci-
500             specific differences in blood DNA methylation in HBV-negative populations at risk for
501             hepatocellular carcinoma development. *Epigenetics, 13*(6), 605-626.
502             doi:10.1080/15592294.2018.1481706

503 Mandal, B. K., & Suzuki, K. T. (2002). Arsenic round the world: a review. *Talanta, 58*(1), 201-235.
504             doi:10.1016/s0039-9140(02)00268-0

505 McCrink-Goode, M. (2014). Pollution: A global threat. *Environment International, 68*, 162-170.
506             doi:10.1016/j.envint.2014.03.023

McKay, J. A., Xie, L., Harris, S., Wong, Y. K., Ford, D., & Mathers, J. C. (2011). Blood as a surrogate
marker for tissue-specific DNA methylation and changes due to folate depletion in post-
partum female mice. *Molecular Nutrition & Food Research, 55*(7), 1026-1035.
doi:10.1002/mnfr.201100008
Minatel, B. C., Sage, A. P., Anderson, C., Hubaux, R., Marshall, E. A., Lam, W. L., & Martinez, V. D.
(2018). Environmental arsenic exposure: From genetic susceptibility to pathogenesis.
*Environment International, 112*, 183-197. doi:10.1016/j.envint.2017.12.017
Noubissi, F. K., Yedjou, C. G., Spiegelman, V. S., & Tchounwou, P. B. (2018). Cross-Talk between Wnt
and Hh Signaling Pathways in the Pathology of Basal Cell Carcinoma. *International Journal of
Environmental Research and Public Health, 15*(7), 13. doi:10.3390/ijerph15071442
Ray, P. D., Yosim, A., & Fry, R. C. (2014). Incorporating epigenetic data into the risk assessment
process for the toxic metals arsenic, cadmium, chromium, lead, and mercury: strategies and
challenges. *Frontiers in Genetics, 5*, 201. doi:10.3389/fgene.2014.00201
Riyahi, S., Sánchez-Delgado, M., Calafell, F., Monk, D., & Senar, J. C. (2015). Combined epigenetic and
intraspecific variation of the DRD4 and SERT genes influence novelty seeking behavior in
great tit Parus major. *Epigenetics, 10*(6), 516-525. doi:10.1080/15592294.2015.1046027
Rosenfeld, C. S. (2010). Animal Models to Study Environmental Epigenetics. *Biology of Reproduction,
82*(3), 473-488. doi:10.1095/biolreprod.109.080952
Rubenstein, D. R., Skolnik, H., Berrio, A., Champagne, F. A., Phelps, S., & Solomon, J. (2016). Sex-
specific fitness effects of unpredictable early life conditions are associated with DNA
methylation in the avian glucocorticoid receptor. *Molecular Ecology, 25*(8), 1714-1728.
doi:10.1111/mec.13483
Ruiz-Hernandez, A., Kuo, C. C., Rentero-Garrido, P., Tang, W. Y., Redon, J., Ordovas, J. M., . . . Tellez-
Plaza, M. (2015). Environmental chemicals and DNA methylation in adults: a systematic
review of the epidemiologic evidence. *Clinical Epigenetics, 7*, 24. doi:10.1186/s13148-015-
0055-7
Saaristo, M., Brodin, T., Balshine, S., Bertram, M. G., Brooks, B. W., Ehlman, S. M., . . . Arnold, K. E.
(2018). Direct and indirect effects of chemical contaminants on the behaviour, ecology and
evolution of wildlife. *Proceedings of the Royal Society B-Biological Sciences, 285*(1885), 10.
doi:10.1098/rspb.2018.1297
Salameh, Y., Bejaoui, Y., & El Hajj, N. (2020). DNA Methylation Biomarkers in Aging and Age-Related
Diseases. *Frontiers in Genetics, 11*, 11. doi:10.3389/fgene.2020.00171
Sanchez-Virosta, P., Espin, S., Garcia-Fernandez, A. J., & Eeva, T. (2015). A review on exposure and
effects of arsenic in passerine birds. *Science of the Total Environment, 512*, 506-525.
doi:10.1016/j.scitotenv.2015.01.069
Sanchez-Virosta, P., Espin, S., Ruiz, S., Panda, B., Ilmonen, P., Schultz, S. L., . . . Eeva, T. (2020).
Arsenic-related oxidative stress in experimentally-dosed wild great tit nestlings.
*Environmental Pollution, 259*, 7. doi:10.1016/j.envpol.2019.113813
Sanchez-Virosta, P., Espin, S., Ruiz, S., Salminen, J. P., Garcia-Fernandez, A. J., & Eeva, T. (2018).
Experimental manipulation of dietary arsenic levels in great tit nestlings: Accumulation
pattern and effects on growth, survival and plasma biochemistry. *Environmental Pollution,
233*, 764-773. doi:10.1016/j.envpol.2017.10.113
Sarkar, D., Leung, E. Y., Baguley, B. C., Finlay, G. J., & Askarian-Amiri, M. E. (2015). Epigenetic
regulation in human melanoma: past and future. *Epigenetics, 10*(2), 103-121.
doi:10.1080/15592294.2014.1003746
Sepers, B., van den Heuvel, K., Lindner, M., Viitaniemi, H., Husby, A., & van Oers, K. (2019). Avian
ecological epigenetics: pitfalls and promises. *Journal of Ornithology, 160*(4), 1183-1203.
doi:10.1007/s10336-019-01684-5
Sharavanan, V. J., Sivaramakrishnan, M., Sivarajasekar, N., Senthilrani, N., Kothandan, R., Dhakal, N.,
. . . Naushad, M. Pollutants inducing epigenetic changes and diseases. *Environmental
Chemistry Letters*, 19. doi:10.1007/s10311-019-00944-3

Skinner, M. K., Manikkam, M., & Guerrero-Bosagna, C. (2010). Epigenetic transgenerational actions
of environmental factors in disease etiology. *Trends in Endocrinology and Metabolism, 21*(4),
214-222. doi:10.1016/j.tem.2009.12.007
Verhoeven, K. J. F., Vonholdt, B. M., & Sork, V. L. (2016). Epigenetics in ecology and evolution: what
we know and what we need to know INTRODUCTION. *Molecular Ecology, 25*(8), 1631-1638.
doi:10.1111/mec.13617
Verhulst, E. C., Mateman, A. C., Zwier, M. V., Caro, S. P., Verhoeven, K. J. F., & Van Oers, K. (2016).
Evidence from pyrosequencing indicates that natural variation in animal personality is
associated with DRD4 DNA methylation. *Molecular Ecology, 25*(8), 1801-1811.
doi:10.1111/mec.13519
Vos, J. G., Dybing, E., Greim, H. A., Ladefoged, O., Lambre, C., Tarazona, J. V., . . . Vethaak, A. D.
(2000). Health effects of endocrine-disrupting chemicals on wildlife, with special reference
to the European situation. *Critical Reviews in Toxicology, 30*(1), 71-133.
doi:10.1080/10408440091159176
Wenzel, M. A., & Piertney, S. B. (2014). Fine-scale population epigenetic structure in relation to
gastrointestinal parasite load in red grouse (Lagopus lagopus scotica). *Molecular Ecology,*
*23*(17), 4256-4273. doi:10.1111/mec.12833
Ziech, D., Franco, R., Pappa, A., & Panayiotidis, M. I. (2011). Reactive Oxygen Species (ROS)-Induced
genetic and epigenetic alterations in human carcinogenesis. *Mutation Research-*
*Fundamental and Molecular Mechanisms of Mutagenesis, 711*(1-2), 167-173.
doi:10.1016/j.mrfmmm.2011.02.015