Does arsenic contamination affect DNA methylation patterns in a wild bird population? An

experimental approach

Veronika N. Laine[1], Mark Verschuuren[1], Kees van Oers[1], Silvia Espín[2,3], Pablo Sánchez-

Virosta[2,3], Tapio Eeva[3], Suvi Ruuskanen[3, 4]

[1]Department of Animal Ecology, Netherlands Institute of Ecology (NIOO-KNAW),

Wageningen, The Netherlands

[2]Area of Toxicology, Department of Socio-Sanitary Sciences, University of Murcia, Spain

[3]Department of Biology, University of Turku, Finland

[4] Department of Biological and Environmental Science, University of Jyväskylä, Finland

Abstract

Pollutants, such as toxic metals, negatively influence organismal health and performance, even

leading to population collapses. Studies in model organisms have shown that epigenetic marks,

such as DNA methylation, can be modulated by various environmental factors, including

pollutants, influencing gene expression and various organismal traits. Yet experimental data

on the effects of pollution on DNA methylation from wild animal populations is largely

lacking. We here experimentally investigated for the first time the effects of early-life exposure

to environmentally relevant levels of a key pollutant, arsenic (As), on genome-wide DNA

methylation in a wild bird population. We experimentally exposed nestlings of great tits (*Parus*

major) to arsenic during their post-natal developmental period (3 to 14 days post-hatching) and

compared their erythrocyte DNA methylation levels to those of respective controls. In contrast

to predictions, we found no overall hypomethylation in the arsenic group. We found evidence

for loci to be differentially methylated between the treatment groups, but for five CpG sites

only. Three of the sites were located in gene bodies of zinc finger and BTB domain containing

47 (*ZBTB47*), HIVEP zinc finger 3 (*HIVEP3*) and insulin like growth factor 2 mRNA binding

protein 1 (*IGF2BP1*). Further studies are needed to evaluate whether epigenetic dysregulation

is a commonly observed phenomena in polluted populations, and what the consequences are

for organism functioning and for population dynamics.

Keywords: pollution, *Parus major*, environmental epigenetics, ecological epigenetics,

ecotoxicology

Running title: Arsenic pollution and DNA methylation in the wild

Introduction

Environmental pollution can negatively affect organisms at multiple level of organization, from

molecular and physiological level to performance, and even lead to population collapses (1-4).

In wild populations, a largely unexplored mechanism mediating such pollution effects is the

potential influence of the epigenome, such as DNA methylation. In human and animal models,

the effects of pollution on the epigenome is studied extensively, and it has been discovered that

methylation patterns can be changed by various environmental factors, including metal and

organic pollutants and other early-life stressors (reviewed by 5-11). DNA methylation is the

addition of a methyl (-CH3) group to the 5' carbon site of cytosines catalyzed by DNA-

methyltransferases, and is generally found to be negatively associated with gene expression

(12). Variation in DNA methylation is linked to variation in phenotypes and behavior, and

associated with the prevalence of various diseases, including cancers in humans and model

animals (13-16). Epigenetic changes from early-life environment may persist and affect health

throughout life-time and may even be transmitted to future generations (16), which could

potentially contribute to explaining delayed or persistent effects of pollutants (e.g. 7). Yet the

effects of pollutants on the epigenome have hardly been explored, and epigenetic research in

wild animal populations is only emerging (8, 17-24).

Arsenic (As) is a global, persistent pollutant, distributed in the environment due to

natural and anthropogenic sources such as mining, industrial activities or coal combustion (25)

and the most highly ranked hazardous substances for animals and plants (26). Across

organisms, arsenic can have negative consequences for basically all organ systems, often via

causing oxidative stress, i.e. the imbalance between harmful reactive oxygen species (ROS)

and antioxidant defenses, and cancer (27, 28).

Arsenic has been repeatedly observed to also modulate patterns of DNA methylation *in*

vitro (e.g. 29) in laboratory animal models (with levels exceeding environmental levels,

reviewed 30) and in studies on human populations (e.g. 31). Arsenic could influence DNA

methylation via multiple pathways: (i) arsenic can change the DNA methylation of a cytosine

via the depletion of the cellular availability of methyl groups. Biotransformation of arsenic to

less toxic forms includes the addition of methyl group(s) (32) with the main methyl-donor for

methylation of both arsenic and cytosines being s-adenosylmethionine (SAM). The high

demand imposed on this molecule during the biotransformation process can then lead to a

global DNA hypomethylation as shown in multiple (bio)medical studies in humans and mice

(reviewed e.g. 5, 33). (ii) Arsenic could influence epigenetic signaling by targeting the zinc

fingers of Tet proteins and perturbing the Tet-mediated oxidation of 5-mC (*in vitro*: 34-37).

(iii) Furthermore, ROS created during arsenic biotransformation have been suggested to

influence DNA methylation by creating aberrant modifications (humans: 38).

Pre/postnatal exposure to arsenic in humans is associated with epigenetic modifications

related to early onset of diseases, which could have long-term consequences (reviewed in 39).

For example, in humans, prenatal arsenic exposure led to global hypomethylation of

inflammatory and tumor suppressor genes (40) and interfered with de novo methylation (41)

in humans. Global hypomethylation can lead to chromosomal abnormalities, contributing to

overall genomic instability and malignant transformations (reviewed in 32). Studies have

demonstrated that widespread DNA hypomethylation induced by arsenic is also associated with

promoter activation and involved in carcinogenesis (reviewed in 32). Arsenic-related

hypomethylation of specific sets of genes has also be reported, and these include, for example,

genes related to neural development (e.g. 42), mitochondria biogenesis (e.g. 43) and

inflammation (e.g. 44). Despite the extensive data on model animals and humans, the potential

effects of environmental arsenic on wild animals via epigenetic dysregulation has not been

studied up to date.

We here investigated the effects of experimental early-life (post-natal) exposure to

arsenic on genome-wide DNA methylation status in a wild population of great tits (*Parus*

major). To our knowledge, this is the first study on the effect of arsenic on epigenetic marks in

a wild population. We used a bird model, since birds have been successfully used in

biomonitoring of pollution and its effects (e.g. 45). Arsenic exposure has been reported to

negatively affect multiple fitness-related traits (growth, physiology, behavior and even egg-

laying) in several bird species (reviewed in 28). For great tits specifically, we have previously

reported (results from the current experiment) that in nestlings, arsenic exposure increased

mortality, reduced wing growth (46) and decreased an intracellular antioxidant, catalase (47),

but did not largely influence body mass, plasma biochemistry (vitamins) or other biomarkers

of oxidative stress (46, 47). More specifically, we here experimentally exposed nestlings in

non-polluted sites to environmentally relevant levels (1 µg/g body mass) of dietary arsenic

during the entire post-hatching growth period, and compared their DNA methylation levels to

respective controls. We used reduced representation bisulfite sequencing (RRBS) to assess

genome-wide methylation and characterized differential methylation across CpG sites between

the experimental and the control group. We predict that arsenic exposure will lead to genome-

wide hypomethylation, and potentially specifically on gene/hubs related to development.

Methods

Arsenic treatment protocol and sampling

The study was conducted in the breeding season of 2015 (laying dates 4th May –10th June) in a

nest box population of great tits (*Parus major)* in western Finland. Great tit is a small passerine

bird and a popular model species in ecological and evolutionary research. Importantly, it is one

of the few non-domesticated bird species, for which the genome and methylome are available

(18, 48-49).

The arsenic exposure, dosages and sampling are described in detail in (46, 47). In short,

the experiment was conducted in a nest-box population with known history of relatively low

pollution levels (50). There are no air pollution samplers at the study sites but metal

biomonitoring studies have been done in this area, for example measuring forest floor moss

metal levels (a proxy for atmospheric fallout). In general, metal levels are relatively low in

moss samples (e.g. for arsenic <0.5 μg/g in 2014; 51) while this value is exceeded in large areas

in Central Europe (52). Mean topsoil arsenic concentration in the study site was 0.76 μg/g in

2014 (53).

Breeding was monitored, and from day 3 after hatching until day 13 whole broods were

subjected to daily oral dosing with the following treatments: arsenic treatment (1 μg arsenic/g

body mass in distilled water, N = 16 broods) or control treatment (distilled water, N = 16

broods). Dosing volumes were adjusted to estimated nestling mass based on average body

masses at different ages from large dataset on long-term averages from the study population

(54). Mass of individual nestlings was not measured every day to reduce handling time and

disturbance to the nest. The volumes dosed to the controls were exactly the same as for

treatments. We dosed the solution directly to the beak of the nestlings with a pipette. The range

of volumes was 50–170 μl and did not exceed the recommended volumes (20 ml/kg, e.g. 55).

The dose aimed to represent environmentally relevant exposure levels occurring in polluted

areas in Europe: It was estimated combining data from several sources, such as (i) the lowest-

observed-adverse-effect level for different effects on mammals (2-8μg/kg/day, 56), (ii) fecal

arsenic levels reported for great tits at some metal polluted sites (reviewed in 28): In previous

data, summarized in (28, Table 1), arsenic concentrations in feces of passerines are within the

range of 0.1–1.4 ppm in unpolluted sites and 5–16 ppm in polluted areas. The levels measured

in the samples from our experiment (ca 6.5 ppm, see results) overlap with these levels,

suggesting that the treatment levels were environmentally relevant, at the lower end of the

range. Yet, Sánchez-Virosta et al. (46) and Janssens et al. (57) report that great tit nestlings

from polluted areas in Harjavalta and Belgium have arsenic levels up to 48-52 ppm, thus levels

even this high are environmentally relevant. Other data sources were (iii) arsenic

concentrations of food items (moth larvae, spiders and beetles) collected directly from parent

great tits feeding their nestlings in the polluted area (46, 47), and (iv) a pilot experiment, to

ensure that the levels were environmentally relevant and were not causing excessive mortality

(46). Fecal matter was sampled 8 days after hatching for metal analyses (see below). DNA

methylation was analyzed from red blood cells (RBCs, 14 d after hatching) to avoid sacrificing

the individuals. Absolute methylation values between e.g. blood and liver or kidney and brain

are highly correlated (48, 58, 59), just like changes in methylation in red blood cells and liver

are correlated (59) and thus blood can be used as a proxy. Ten samples from the arsenic and

ten from control treatment were selected for the DNA methylation analyses. These included

five females and five males from each treatment (molecularly sexed, following 60). Only one

nestling per nest was selected to avoid pseudoreplication. We made use of the knowledge on

the fecal arsenic levels (see below), and selected individuals from 10 broods with highest

arsenic concentrations from the arsenic treatment and 10 lowest concentrations from the

control. All the dead nestlings found in the nests were collected and frozen at -20ºC until

necropsies could be performed in July 2015. Carcasses were necropsied to measure arsenic and

metal concentrations in liver and bone in order to compare arsenic accumulation among groups

and its distribution among tissues (46). The experiment was conducted under licenses from the

Animal Experiment Committee of the State Provincial Office of Southern Finland (license

number ESAVI/11579/04.10.07/2014) and the Centre for Economic Development, Transport

and the Environment, ELY Centre Southwest Finland (license number VARELY/593/2015).

Metal analyses

For detailed analyses, see Sánchez-Virosta et al. (46). Briefly, in both experimental groups,

several fecal samples (any sex) from the same brood were combined to assess brood level metal

exposure (total N = 32 broods). We determined the concentrations of arsenic, but also other

metals to confirm that the levels of other metals were low and similar across the treatment

groups (see 46). The determination of pollutants was conducted with inductively coupled

plasma optical emission spectrometry (ICP-OES) with detection limit of 1 ppt (ng/l) and below.

Calibration standards and certified reference materials were used for method validation. The

levels of other measured metals (aluminium, lead, nickel, zinc, manganese, iron, copper) were

low, and did not differ among the treatment groups (all t <0.88, all p <0.38).

DNA isolation

DNA isolation was performed at the Center of Evolutionary Applications (University of Turku,

Finland). We used RBCs given that previous studies suggest that blood shows similar

methylation patterns as brain tissue in the study species (e.g. 80% similarity between brain and

blood methylation in CpGs; 48, 49). DNA was extracted from 10-20 μl RBCs using the salt

extraction method modified from (61). Extracted DNA was treated with RNase-I according to

the manufacturer's protocol. DNA concentration was measured fluorometrically with a Qubit

High Sensitivity kit (ThermoFisher Scientific) and we assessed DNA integrity by running each

DNA sample on an agarose gel.

RRBS library preparation

We used a reduced representation bisulfite sequencing (RRBS) approach, which enriches for

regions of the genome that have a high CpG content. We chose the RRBS approach because

with the use of *MspI* as restriction enzyme, the method targets regions that are enriched for

CpG sites. These regions are typically situated in or near the promotor regions, which has the

advantage that CpGs in a relatively large proportion of the genes are covered (22, 62) making

this a cost-effective method for detecting sites that are likely functional (16). It was previously

shown in the study species that a vast majority of methylated Cs (97%) were derived from CpG

sites in blood (48). Sequencing was conducted at the Finnish Microarray and Sequencing

Center in Turku, Finland. The library preparation was started from 200 ng of genomic DNA

and was carried out according to a protocol adapted from (63). The first step in the workflow

involved the fragmentation of genomic DNA with MspI where the cutting pattern of the

enzyme (C^CGG) was used to systematically digest DNA to enrich for CpG dinucleotides.

After a fragmentation step a single reaction was carried out to end repair and A-tail (required

for the adapter ligation) the MspI digested fragments using Klenow fragment (3' => 5' exo)

following the purification of A-tailed DNA with bead SPRI clean-up method (AMPure

magnetic beads). A unique Illumina TruSeq indexing adapter was then ligated to each sample

during adapter ligation step to be able to identify pooled samples of one flow cell lane. To

reduce the occurrence of adapter dimers, a lower concentration of adapters (1:10 dilution) was

used than recommended by the manufacturer. These ligated DNA fragments were purified with

bead SPRI clean-up method before putting samples through bisulfite conversion to achieve C-

to-U conversion of unmethylated cytosines, whereas methylated cytosines remain intact.

Bisulfite conversion and sample purification were done according to Invitrogen MethylCode

Bisulfite Conversion Kit. Aliquots of converted DNA were amplified by PCR (16 cycles) with

Taq/Pfu Turbo Cx Polymerase, a proofreading PCR enzyme that does not stall when it

encounters uracil, the product of the bisulfite reaction, in the template. PCR-amplified RRBS

libraries were purified using two subsequent rounds of SPRI beadclean-ups to minimize primer

dimers in the final libraries. The high quality of the libraries was confirmed with Advanced

Analytical Fragment Analyzer and the concentrations of the libraries were quantified with

Qubit® Fluorometric Quantitation, Life Technologies. We used an average fragment size of

250-350 bp for sequencing.

Sequencing

The samples were normalized and pooled for the automated cluster preparation which was

carried out with Illumina cBot station. The 20 libraries were combined in two pools, 10 samples

in each pool (treatments and sexes equally distributed between the pools) and sequenced in two

lanes. The samples were sequenced with an Illumina HiSeq 2500 instrument using TruSeq v3

sequencing chemistry. Paired-end sequencing with 2 x 100 bp read length was used with 6 bp

index run.

Sequence data processing and differential methylation expression analysis

All the reads were checked for quality using FastQC (Babraham Bioinformatics) with multiQC

(64), and low-quality sequences were trimmed with Trim Galore v. 0.4.4 (Brabraham

Bioinformatics) by using --quality 20 --paired --rrbs settings.

The trimmed reads were mapped to the *Parus major* reference genome build 1.1.

(https://www.ncbi.nlm.nih.gov/assembly/ GCF_001522545.2) using Bismark (65) with default

parameters. Methylation calling was conducted with Bismark, first with default settings with

paired-end mode and overlap removal (--p --no_overlap). After this first calling round, we

observed a methylation bias for the samples by plotting the methylation proportion across each

possible position in the read. Based on the plotting, the three and two first bases of R1 and R2

respectively of the 5' prime end were omitted and the first base in the R2 3' prime end was also

omitted in the final methylation calling. Thereafter, Methylkit (66) implemented in R was used

for filtering and differential methylation analysis. We discarded bases that had coverage below

10x. To avoid a possible PCR bias we also discarded bases that had more than 99.9th percentile

of coverage in each sample. Before differential methylation analysis we merged read counts

from reads covering both strands of a CpG dinucleotide and CpGs needed to be covered with

at least 8 samples per group (control and treatment).

Samples were thereafter clustered based on the similarity of their overall methylation

profile by (i) using the clustering method ward.D in Methylkit's clusterSamples -function and

(ii) using principal component analysis (PCA) with Methylkit's PCASamples -function. We

also checked for lane and sex effect by using Methylkit's assocComp -function where it checks

which principal components are statistically associated with the potential batch effects such as

the used lane and sex of the individuals. For the former, no missing data is allowed, thus we

created a separate data object where all the individuals needed to be covered.

For analyzing differential methylation of CpG sites between control and arsenic

treatment we used the beta-binomial model from DSS package (67) which is also included in

Methylkit (calculateDiffMethDSS - function). DSS calculates the differential methylation

statistics using a beta-binomial model with parameter shrinkage. Bonferroni correction was

applied to account for multiple testing with q-value of 0.05. Furthermore, we also did the "tiling

window analysis" in Methylkit where methylation information is summarized over tiling

windows which are then used the in DSS analysis. We used the default values, win.size=1000,

step.size=1000, cov.bases = 10 for the tiling and ran DSS again for these regions.

Results

Arsenic exposure

As reported in Sánchez-Virosta et al. (46, 47), dietary arsenic treatment successfully increased

arsenic load as fecal arsenic levels were on average 10 times higher in arsenic exposure

compared to control group (average±SD ppm: control 0.51±0.50, arsenic exposure 4.92±4.57,

$t_{15.4}$ = -3.83, p = 0.0015). In the subsample of nests selected for RRBS, the values were

0.47±0.37 ppm for control nests and 6.50±5.10 ppm for arsenic treatment, respectively.

Furthermore, increased levels were also found in internal tissues: the mean (±SD) arsenic

concentrations in liver were 4.19 ± 5.92 µg/g, d.w. (N=21) for arsenic exposure and 0.058 ±

0.100 µg/g, d.w. (N= 16) for control group, and in the bone 3.37 ± 3.85 µg/g, d.w. and 0.074 ±

0.103 µg/g, d.w., respectively (see Table 2 in 46). The levels were statistically significantly

higher in arsenic exposure group compared to control group (p <0.001).

Sequencing and mapping

The total number of read pairs was 341 million (Supplementary Table 1), varying from 14

million to 20 million per individual. After QC filtering the final number of read pairs was 337

million (Supplementary Table 1). The RRBS individual sequencing data have been deposited

in NCBI (Number will be added later). Mapping efficiency was on average 46.15% and on

average 3.1 million cytosines were covered before 10x coverage and percentile filtering. After

filtering, 1.3 million cytosines were identified in CpG context. When combining the Cs from

both strands and restricting our data to at least 8 individuals per group to be covered, we ended

up having 652 655 CpGs.

Sample clustering and differential methylation

Both the ward.D and PCA clustering methods showed that sample 14 from the treatment group

was an outlier in its methylation profile (Supplementary Figure 1). That particular sample also

had a low number of reads and showed lower duplication levels (Supplementary Table 1) and

we therefore decided to exclude this sample from further analysis. No lane effect was detected,

but PC3 (explained 0.23% of the variance) was associated with sex after Bonferroni correction

(Supplementary Table 2, Supplementary Figure 2), mostly driven by two samples, ctrl_3F and

test_16F, since after removing these two female samples from the data, PC3 was not significant

anymore. Furthermore, when removing the PC3 from the data, three CpG sites were significant

in the differential methylation analysis done with DSS: two of them were the same as when

including all the PCs (see below, Table 1). The three other significant sites found below were

not covered by all individuals as required in this PC-removal analysis.

In the differential methylation analysis when including all the PCs, five CpG sites showed a

significant difference in methylation level with a q-value below 0.05 and percent methylation

difference larger than 10% (Table 1, Figure 1, Supplementary Table 3). Lambda estimation

was close to 1 ($\lambda = 0.747$, SE 0.000136) (Supplementary Figure 3), suggesting no systematic

biases ($\lambda > 1$ indicates bias). Four of these sites were hypermethylated (higher methylation in

the arsenic treatment group) and one was hypomethylated (higher methylation in the control

group). Three of the sites were located in gene bodies, namely zinc finger and BTB domain

containing 47 (*ZBTB47*), HIVEP zinc finger 3 (*HIVEP3*) and insulin like growth factor 2

mRNA binding protein 1 (*IGF2BP1*) based on NCBI *P. major* annotation report 102. None of

the regions from the tiling windows analysis were differentially methylated between control

and treatment samples.

Table 1. The differentially methylated CpG sites between arsenic exposed and control

individuals. Methylation diff% refers to the methylation difference, comparing arsenic exposed

to control group. Positive values therefore indicate hypermethylation in the arsenic treatment

group compared to the control group. PC3 indicates sites that were significant after PC3

removal.

Chr	Chr Genbank	Position	P-value	q-value	Methylation diff %	Gene	PC3
2	NC_031769.1	2,448,788	1.00E-10	6.53E-05	36.40	*ZBTB47*	
12	NC_031781.1	9,949,364	1.23E-08	8.05E-03	57.66	-	
23	NC_031791.1	5,408,232	3.43E-08	2.24E-02	42.66	*HIVEP3*	x
UN	NW_015379267.1	107,660	7.99E-10	5.21E-04	60.43	*IGF2BP1*	
UN	NW_015379318.1	39,910	1.42E-08	9.27E-03	-52.66	-	x
Only in PC3 removed							
14	NC_031783.1	14,025,702	1.34E-07	0.042	27.09	-	x

Figure 1. Plots of significance of CpG sites from the differential methylation analysis conducted with DSS implemented in Methylkit. (a) Manhattan plot with the significance of differential methylation of the arsenic treatment against the control pool, against the great tit reference genome version 1.1. The orange line depicts the genome-wide threshold based on a

317 Bonferroni correction: 7.11. (b) A volcano plot of the significance against the absolute

318 difference in methylation between the two pools, with delta methylation is arsenic treatment –

319 control. Orange points are the genome-wide significant sites after Bonferroni correction and

320 filtering for Delta methylation >10%.

Discussion

We investigated whether early-life exposure to environmentally relevant levels of experimental

arsenic affects DNA methylation in a wild vertebrate population. The experimental treatment

increased arsenic levels significantly, but contrary to predictions, did not lead to overall

hypomethylation. We found that treated individuals showed hypermethylation in four CpG

sites and hypomethylation in one CpG site, indicating that increased levels of arsenic exposure

appears to affect methylation at specific parts of the genome only. Yet also at these sites, the

assumption of general hypomethylation was not met.

The lack of overall or site-specific hypomethylation may be explained by various

factors: first, contrary to our predictions, the methyl donor s-adenosylmethionine needed for

methylation may not have been limiting, potentially because oxidative status was not altered

dramatically in all individuals. Indeed, as reported from the exact same experiment and samples

by Sánchez-Virosta et al. (47), most biomarkers of oxidative status and damage in blood were

only slightly (but not statistically significantly) elevated, and only the antioxidant enzyme

catalase showed a significant decrease. In the future, sampling before and after exposure to e.g.

pollutants may be advised to associate DNA methylation changes directly to changes in

oxidative status, for example in adult birds (in contrast to developing animals where

measurements are confounded by the changes in growth and associated changes in physiology).

Second, the response is likely to depend on the tissue type studied. For example, global

hypomethylation in response to arsenic exposure is not consistently reported in blood: in

humans, where blood leucocytes have been used to characterize arsenic associated changes no

evidence for global hypo or hypermethylation was detected, yet arsenic was repeatedly reported

to induce hypermethylation in various genes (especially promoters) (68), whereas global

hypomethylation was detected in hepatic cells (69). Given that arsenic metabolism and SAM

production mostly takes part in liver, we may expect tissue-dependent hypomethylation

especially in liver, but not necessarily in other tissues. Unfortunately, we lack oxidative status

measurements from the liver in this experiment. Studies have shown that absolute methylation

values between e.g. blood and liver or kidney and brain are highly correlated (48, 58), just like

changes in methylation in red blood cells and liver are correlated (59). Nevertheless tissue-

specific methylation differences were larger for genes that are expressed in a tissue-specific

way (48) and measuring methylation levels from red blood cells might therefore miss tissue-

dependent genes whose expression is expected to change (59).

Furthermore, contrary to many previous studies in laboratory animals, this experiment

was conducted with relatively low doses, mimicking exposure in polluted environments,

whereas effects via SAM may only be apparent when levels are higher. Also, we were only

interested in short-term, early-life effects while resident species inhabiting polluted

environments during their whole life-span may show marked effects due to cumulative arsenic

exposure. This is an interesting avenue for further research.

As the experiment was conducted in a wild population, in comparison to previous

studies in laboratory, the environmental or genetic variability and potential variability across

sexes may have masked some effects of the experimental treatments. Arsenic is known to have

sex-dependent effects in many model systems (though predominantly in adult animals;

reviewed e.g. 32). Furthermore, for example studies on mice report sex differences in DNA

methylation patters in response to arsenic (e.g. 70), and general methylation differences among

the sexes in young chickens (71). Yet, our initial models suggested that sex explained only a

very minor part of the variation in DNA methylation (and was therefore dropped from the final

model), which suggests that in our data sex-bias is unlikely to strongly mask the effects. DNA

methylation is known to be heavily influenced by the genetic background, for example in van

Oers et al. (62), the majority of the variation between individuals was explained by genetic

similarity. In the future, split-brood experimental designs may be used to distinguish genetic

effects from environmental. The arsenic exposure applied (as measured from the fecal samples)

was also at the lower range of variation if compared to polluted environments, which may

contribute to the findings of only limited differences − yet mortality was increased with these

levels, as reported in (46). We also report large variation in the fecal arsenic levels within the

arsenic exposure treatment. Several factors may affect those levels, such as the time elapsed

between last dosing and sampling, the times the nestling has been fed in that time and how

many droppings they have produced, among others. Feces dropped soon after arsenic

administration likely contain higher arsenic levels than later on.

We could annotate three of the five differentially methylated sites to genes. One of the

genes, *IGF2BP1* is especially interesting as it is associated with development and growth: it

has been showed that *IGF2BP1* plays important roles in various aspects of cell function, such

as cell proliferation, differentiation, migration, morphology and metabolism (72, 73) but also

embryogenesis and potentially even arsenic-related carcinogenesis (74, 75). *IGF2BP1* is

abundantly expressed in fetal and neonatal tissues (73). Furthermore, two of the genes, *ZBTB47*

and *HIVEP3* are both zinc-finger domains and are associated with transcriptional regulation

(76). Epigenetic regulation of both *ZBTB47* and *HIVEP3* is known to be associated cancer (77,

78). Because our sample size in combination with a stringent correction for repeated sampling

limits the power to detect subtle differences, we do expect to find a fraction of the number of

differentially methylated CpGs (79).

All the three gene-related differentially methylated CpG sites were found in the gene

body region, in both intron (*IGF2BP1*) and exons (*ZBTB47* and *HIVEP3*). Hypermethylation

at CpG sites at promoter regions represses transcription of genes which is a well-known

mechanism operating in many scenarios. DNA methylation at intergenic regions and gene

bodies and its impact on gene expression is gaining more attention especially in cancer studies

(80). Interestingly, a recent study on corals showed that gene body methylation was altered by

environmental factors, which facilitated acclimatization and adaptation to different habitats

(81). However, in great tits the DNA methylation observed in CpGs that are situated within

gene bodies do not seem to affect gene expression (48), thus future studies are needed to

determine the role of gene body methylation in gene expression control.

In conclusion, our study shows that early-life exposure to a toxic metal, arsenic, potentially

affects fitness via DNA methylation changes in specific pathways, but not via an overall

hypomethylation in the red blood cells. The effect might be more profound in other tissues that

are more relevant to arsenic metabolism, such as liver. Thus, future studies should inspect other

tissues as well. Other pathways of epigenetic alterations, known to be subject to arsenic-related

alternations in vitro, such as histone acetylation (29) and micro-RNAs (82) could be further

explored.

ASSOCIATED CONTENT

Figure S1. Clustering of samples based on ward.D (a) and principal component analysis (PCA)

(b) in Methylkit

Table S1. Number of reads before and after read trimming with mapping and methylation

calling success of Bismark. CpG site filtering was done with Methylkit.

Table S2. Results from the tests on the effects of sequencing lane and sex.

Table S3. A full description of all differentially methylated sites.

AUTHOR INFORMATION

Corresponding author: Suvi Ruuskanen (skruus@utu.fi)

Conflict of interest

We have no conflict of interest to declare.

Data accessibility

Data will be deposited in Dryad and in Genbank upon acceptance.

Author contributions

SR, SE, PSV, VNL, MV, KvO and TE designed the study. TE, SE and PSV collected the data.

SR and KvO designed the sequencing. VNL and MV conducted the bioinformatic analyses. SE

and PSV conducted metal analyses. KvO and VNL provided the genome resources. VNL, MV,

KvO and SR interpreted the data. VNL and SR wrote the first draft. All authors contributed to

writing the manuscript.

Funding sources

TE, SE, PSV were funded by Academy of Finland (to TE). SR was funded by Academy of

Finland and Turku University Foundation.

Acknowledgements

We thank Miia Rainio and Jorma Nurmi for their efforts in helping us with fieldwork and Fleur

Gawehns-Bruning and William Sies for assistance with the methylation analysis. We also thank

the Center for Evolutionary Applications for molecular work and Finnish Functional Genomics

Centre for sequencing services. Our study was financed by KONE foundation (SR) and

Academy of Finland (TE: project 265859). SE is currently funded by *Ministerio de Ciencia,*

Innovación y Universidades (IJCI-2017-34653 to SE).

References

1. Dudgeon, D., Arthington, A. H., Gessner, M. O., Kawabata, Z. I., Knowler, D. J.,
Leveque, C., . . . Sullivan, C. A. (2006). Freshwater biodiversity: importance, threats,
status and conservation challenges. *Biological Reviews, 81*(2), 163-182.
doi:10.1017/s1464793105006950
2. McCrink-Goode, M. (2014). Pollution: A global threat. *Environment International, 68*,
162-170. doi:10.1016/j.envint.2014.03.023
3. Saaristo, M., Brodin, T., Balshine, S., Bertram, M. G., Brooks, B. W., Ehlman, S. M., .
. . Arnold, K. E. (2018). Direct and indirect effects of chemical contaminants on the
behaviour, ecology and evolution of wildlife. *Proceedings of the Royal Society B-*
Biological Sciences, 285(1885), 10. doi:10.1098/rspb.2018.1297
4. Vos, J. G., Dybing, E., Greim, H. A., Ladefoged, O., Lambre, C., Tarazona, J. V., . . .
Vethaak, A. D. (2000). Health effects of endocrine-disrupting chemicals on wildlife,
with special reference to the European situation. *Critical Reviews in Toxicology, 30*(1),
71-133. doi:10.1080/10408440091159176
5. Baccarelli, A., & Bollati, V. (2009). Epigenetics and environmental chemicals. *Current*
Opinion in Pediatrics, 21(2), 243-251. doi:10.1097/MOP.0b013e32832925cc
464
6. Cheng, T. F., Choudhuri, S., & Muldoon-Jacobs, K. (2012). Epigenetic targets of some
toxicologically relevant metals: a review of the literature. *Journal of Applied*
Toxicology, 32(9), 643-653. doi:10.1002/jat.2717
7. Head, J. A. (2014). Patterns of DNA Methylation in Animals: An Ecotoxicological
Perspective. *Integrative and Comparative Biology, 54*(1), 77-86.
doi:10.1093/icb/icu025
8. Head, J. A., Dolinoy, D. C., & Basu, N. (2012). Epigenetics for ecotoxicologists.
Environmental toxicology and chemistry / SETAC, 31(2), 221-227.
doi:10.1002/etc.1707
9. Ray, P. D., Yosim, A., & Fry, R. C. (2014). Incorporating epigenetic data into the risk
assessment process for the toxic metals arsenic, cadmium, chromium, lead, and
mercury: strategies and challenges. *Frontiers in Genetics, 5*, 201.
doi:10.3389/fgene.2014.00201
10. Ruiz-Hernandez, A., Kuo, C. C., Rentero-Garrido, P., Tang, W. Y., Redon, J., Ordovas,
J. M., . . . Tellez-Plaza, M. (2015). Environmental chemicals and DNA methylation in
adults: a systematic review of the epidemiologic evidence. *Clinical Epigenetics, 7*, 24.
doi:10.1186/s13148-015-0055-7

11. Sharavanan, V. J., Sivaramakrishnan, M., Sivarajasekar, N., Senthilrani, N., Kothandan, R., Dhakal, N., . . . Naushad, M. Pollutants inducing epigenetic changes and diseases. *Environmental Chemistry Letters*, 19. doi:10.1007/s10311-019-00944-3

12. Goldberg, A. D., Allis, C. D., & Bernstein, E. (2007). Epigenetics: A landscape takes shape. *Cell, 128*(4), 635-638. doi:10.1016/j.cell.2007.02.006

13. Rosenfeld, C. S. (2010). Animal Models to Study Environmental Epigenetics. *Biology of Reproduction, 82*(3), 473-488. doi:10.1095/biolreprod.109.080952

14. Salameh, Y., Bejaoui, Y., & El Hajj, N. (2020). DNA Methylation Biomarkers in Aging and Age-Related Diseases. *Frontiers in Genetics, 11*, 11. doi:10.3389/fgene.2020.00171

15. Skinner, M. K., Manikkam, M., & Guerrero-Bosagna, C. (2010). Epigenetic transgenerational actions of environmental factors in disease etiology. *Trends in Endocrinology and Metabolism, 21*(4), 214-222. doi:10.1016/j.tem.2009.12.007

16. Horsthemke, B. (2018). A critical view on transgenerational epigenetic inheritance in humans. *Nature Communications, 9*. doi:10.1038/s41467-018-05445-5

17. Bossdorf, O., Richards, C. L., & Pigliucci, M. (2008). Epigenetics for ecologists. *Ecology Letters, 11*(2), 106-115. doi:10.1111/j.1461-0248.2007.01130.x

18. Laine, V. N., Gossmann, T. I., Schachtschneider, K. M., Garroway, C. J., Madsen, O., Verhoeven, K. J. F., . . . Great Tit HapMap, C. (2016). Evolutionary signals of selection on cognition from the great tit genome and methylome. *Nature Communications, 7*. doi:10.1038/ncomms10474

19. Liebl, A. L., Schrey, A. W., Richards, C. L., & Martin, L. B. (2013). Patterns of DNA Methylation Throughout a Range Expansion of an Introduced Songbird. *Integrative and Comparative Biology, 53*(2), 351-358. doi:10.1093/icb/ict007

20. Riyahi, S., Sánchez-Delgado, M., Calafell, F., Monk, D., & Senar, J. C. (2015). Combined epigenetic and intraspecific variation of the DRD4 and SERT genes influence novelty seeking behavior in great tit Parus major. *Epigenetics, 10*(6), 516-525. doi:10.1080/15592294.2015.1046027

21. Rubenstein, D. R., Skolnik, H., Berrio, A., Champagne, F. A., Phelps, S., & Solomon, J. (2016). Sex-specific fitness effects of unpredictable early life conditions are associated with DNA methylation in the avian glucocorticoid receptor. *Molecular Ecology, 25*(8), 1714-1728. doi:10.1111/mec.13483

22. Sepers, B., van den Heuvel, K., Lindner, M., Viitaniemi, H., Husby, A., & van Oers, K. (2019). Avian ecological epigenetics: pitfalls and promises. *Journal of Ornithology, 160*(4), 1183-1203. doi:10.1007/s10336-019-01684-5

23. Verhoeven, K. J. F., Vonholdt, B. M., & Sork, V. L. (2016). Epigenetics in ecology and evolution: what we know and what we need to know INTRODUCTION. *Molecular Ecology, 25*(8), 1631-1638. doi:10.1111/mec.13617

24. Wenzel, M. A., & Piertney, S. B. (2014). Fine-scale population epigenetic structure in relation to gastrointestinal parasite load in red grouse (Lagopus lagopus scotica). *Molecular Ecology, 23*(17), 4256-4273. doi:10.1111/mec.12833

25. Mandal, B. K., & Suzuki, K. T. (2002). Arsenic round the world: a review. *Talanta, 58*(1), 201-235. doi:10.1016/s0039-9140(02)00268-0

26. ATSDR (2019). The ATSDR 2019 Substance Priority List. In: Agency for Toxic Substances and Disease Registry, U.S. Public Health Service, U.S.A.

27. Jomova, K., Jenisova, Z., Feszterova, M., Baros, S., Liska, J., Hudecova, D., . . . Valko, M. (2011). Arsenic: toxicity, oxidative stress and human disease. *Journal of Applied Toxicology, 31*(2), 95-107. doi:10.1002/jat.1649

28. Sánchez-Virosta, P., Espín, S., García-Fernández, A. J., & Eeva, T. (2015). A review on exposure and effects of arsenic in passerine birds. *Science of the Total Environment, 512*, 506-525. doi:10.1016/j.scitotenv.2015.01.069

29. Jensen, T. J., Novak, P., Wnek, S. M., Gandolfi, A. J., & Futscher, B. W. (2009). Arsenicals produce stable progressive changes in DNA methylation patterns that are linked to malignant transformation of immortalized urothelial cells. *Toxicology and Applied Pharmacology, 241*(2), 221-229. doi:10.1016/j.taap.2009.08.019

30. Bjorklund, G., Aaseth, J., Chirumbolo, S., Urbina, M. A., & Uddin, R. (2018). Effects of arsenic toxicity beyond epigenetic modifications. *Environmental Geochemistry and Health, 40*(3), 955-965. doi:10.1007/s10653-017-9967-9

31. Alegria-Torres, J. A., Carrizales-Yanez, L., Diaz-Barriga, F., Rosso-Camacho, F., Motta, V., Tarantini, L., & Bollati, V. (2016). DNA Methylation Changes in Mexican Children Exposed to Arsenic From Two Historic Mining Areas in San Luis Potosi. *Environmental and Molecular Mutagenesis, 57*(9), 717-723. doi:10.1002/em.22062

32. Minatel, B. C., Sage, A. P., Anderson, C., Hubaux, R., Marshall, E. A., Lam, W. L., & Martinez, V. D. (2018). Environmental arsenic exposure: From genetic susceptibility to pathogenesis. *Environment International, 112*, 183-197. doi:10.1016/j.envint.2017.12.017

33. Hubaux, R., Becker-Santos, D. D., Enfield, K. S. S., Rowbotham, D., Lam, S., Lam, W. L., & Martinez, V. D. (2013). Molecular features in arsenic-induced lung tumors. *Molecular Cancer, 12*, 11. doi:10.1186/1476-4598-12-20

34. Liu, S., Jiang, J., Li, L., Amato. N.J., Wang, Z., Wang, Y. (2015) Arsenite Targets the Zinc Finger Domains of Tet Proteins and Inhibits Tet-Mediated Oxidation of 5-Methylcytosine. *Environmental Science & Technology* 49, 11923-11931.

35. Zhao, B., Yang, Y., Wang, X., Chong, Z., Yin, R., Song, S-H., … Wang, H. (2014). Redox-active quinones induces genome-wide DNA methylation changes by an iron-mediated and Tet-dependent mechanism. *Nucleic Acids Research*, 42, 1593–1605.

36. Yin, R., Mo, J., Dai, J., Wang, H. (2017). Nickel(II) Inhibits Tet-Mediated 5-Methylcytosine Oxidation by High Affinity Displacement of the Cofactor Iron(II). *ACS Chemical Biology* 12, 1494-1498.

37. Yin, R., Mo, J., Dai, J., Wang, H. (2018). Nickel(II) inhibits the oxidation of DNA 5-methylcytosine in mammalian somatic cells and embryonic stem cells. *Metallomics*, 10, 504–512.

38. Ziech, D., Franco, R., Pappa, A., & Panayiotidis, M. I. (2011). Reactive Oxygen Species (ROS)-Induced genetic and epigenetic alterations in human carcinogenesis. *Mutation Research-Fundamental and Molecular Mechanisms of Mutagenesis, 711*(1-2), 167-173. doi:10.1016/j.mrfmmm.2011.02.015

39. Gangopadhyay, S., Sharma, V., Chauhan, A., & Srivastava, V. (2019). Potential facet for prenatal arsenic exposure paradigm: linking endocrine disruption and epigenetics. *Nucleus-India, 62*(2), 127-142. doi:10.1007/s13237-019-00274-3

40. Kile, M. L., Houseman, E. A., Baccarelli, A. A., Quamruzzaman, Q., Rahman, M., Mostofa, G., . . . Christiani, D. C. (2014). Effect of prenatal arsenic exposure on DNA methylation and leukocyte subpopulations in cord blood. *Epigenetics, 9*(5), 774-782. doi:10.4161/epi.28153

41. Broberg, K., Ahmed, S., Engstrom, K., Hossain, M. B., Mlakar, S. J., Bottai, M., . . . Vahter, M. (2014). Arsenic exposure in early pregnancy alters genome-wide DNA methylation in cord blood, particularly in boys. *Journal of Developmental Origins of Health and Disease, 5*(4), 288-298. doi:10.1017/s2040174414000221

42. Zarazua, S., Rios, R., Delgado, J.M., Santoyo, M.E., Ortiz-Perez, D., Jimenez-Capdeville, M.E. (2010). Decreased argnine mehtylation and myelin alternations in arsenic exposed rats. Neurotoxicology 31, 94-100.

43. Sanval, T., Paul, M., Bhattacharjee, S., Bhattacharjee, P. 2020. Epigenetic alternation of mitochondrial biogenesis regulatory genes in arsenic exposure individuals (with and without skin lesions) and skin cancer tissues: A case control study. Chemosphere 258, 127305.

44. Phookphan, P., Navasumrit, P-, Waraprasit, S., Promvijit, J., Chaisatra, K., Ngaotepprutaram, T., Ruchirawat, M. 2017. Hypomethylation of inflammaratory genes (COX2, ERG1, and SOC23) and increased urinary 8-nitroguanine in arsenic-exposed newborns and children. Toxicology and Applied Pharmacology 316, 36-47.

45. García-Fernández, A.J., Espín, S., Martínez-López, E., (2013). Feathers as a Biomonitoring Tool of Polyhalogenated Compounds: A Review. Environmental Science & Technology 47, *3028-3043. DOI: 10.1021/es302758x*

46. Sánchez-Virosta, P., Espín, S., Ruiz, S., Panda, B., Ilmonen, P., Schultz, S. L., . . . Eeva, T. (2020). Arsenic-related oxidative stress in experimentally-dosed wild great tit nestlings. *Environmental Pollution, 259*, 7. doi:10.1016/j.envpol.2019.113813

47. Sánchez-Virosta, P., Espín, S., Ruiz, S., Salminen, J. P., García-Fernández, A. J., & Eeva, T. (2018). Experimental manipulation of dietary arsenic levels in great tit nestlings: Accumulation pattern and effects on growth, survival and plasma biochemistry. *Environmental Pollution, 233*, 764-773. doi:10.1016/j.envpol.2017.10.113

48. Derks, M. F. L., Schachtschneider, K. M., Madsen, O., Schijlen, E., Verhoeven, K. J. F., & van Oers, K. (2016). Gene and transposable element methylation in great tit (*Parus major*) brain and blood. *Bmc Genomics, 17*, 13. doi:10.1186/s12864-016-2653-y

49. Verhulst, E. C., Mateman, A. C., Zwier, M. V., Caro, S. P., Verhoeven, K. J. F., & Van Oers, K. (2016). Evidence from pyrosequencing indicates that natural variation in animal personality is associated with DRD4 DNA methylation. *Molecular Ecology, 25*(8), 1801-1811. doi:10.1111/mec.13519

50. Eeva, T., Ryoma, M., & Riihimaki, J. (2005). Pollution-related changes in diets of two insectivorous passerines. *Oecologia, 145*(4), 629-639. doi:10.1007/s00442-005-0145-x

51. Ruuth, J., Toivonen, H., Kuhmonen, I. and Kiljunen, A. (2016). Porin-Harjavallan alueen ilmanlaadun bioindikaattoritutkimus vuosina 2014–2015 - Nab Labs Oy. (in Finnish)

52. Harmens, H., Norris, D. A., Steinnes, E., et al. (2010). Mosses as biomonitors of atmospheric heavy metal deposition: Spatial patterns and temporal trends in Europe. - Environmental Pollution 158: 3144-3156.

53. Ruiz, S., Espín, S., Sánchez-Virosta, P., Salminen, J.-P., Lilley, T. M. and Eeva, T. (2017). Vitamin profiles in two free-living passerine birds under a metal pollution gradient - a calcium supplementation experiment. - Ecotoxicology and Environmental Safety 138: 242-252.

54. Eeva, T., Ahola, M. and Lehikoinen, E. (2009). Breeding performance of blue tits (*Cyanistes caeruleus*) and great tits (*Parus major*) in a heavy metal polluted area. Environmental Pollution 157: 3126–3131.

55. Gad, SC , Spainhour CB, Shoemake C, et. al. (2016). Tolerable Levels of Nonclinical Vehicles and Formulations Used in Studies by Multiple Routes in Multiple Species With Notes on Methods to Improve Utility, International Journal of Toxicology, 1-84.

56. ATSDR (2007). Toxicological profile for arsenic (Agency for Toxic Substances and Disease Registry, U.S. Public Health Service).

57. Janssens, E., Dauwe, T., Pinxten, R., Bervoets, L., Blust, R., Eens, M. (2003). Effects of heavy metal exposure on the condition and health of nestlings of the great tit (Parus major), a small songbird species. Environmental Pollution 126, 267–274

58. McKay, J. A., Xie, L., Harris, S., Wong, Y. K., Ford, D., & Mathers, J. C. (2011). Blood as a surrogate marker for tissue-specific DNA methylation and changes due to folate depletion in post-partum female mice. *Molecular Nutrition & Food Research, 55*(7), 1026-1035. doi:10.1002/mnfr.201100008

59. Lindner, M., Verhagen, I., Viitaniemi, H.M. *et al. 2021*. Temporal changes in DNA methylation and RNA expression in a small song bird: within- and between-tissue comparisons. *BMC Genomics* 22, 36. doi:10.1186/s12864-020-07329-9

60. Bantock, T. M., Prys-Jones, R. P., & Lee, P. L. M. (2008). New and improved molecular sexing methods for museum bird specimens. *Molecular Ecology Resources, 8*(3), 519-528. doi:10.1111/j.1471-8286.2007.01999.x

61. Aljanabi, S. M., & Martinez, I. (1997). Universal and rapid salt-extraction of high quality genomic DNA for PCR-based techniques. *Nucleic Acids Research, 25*(22), 4692-4693. doi:10.1093/nar/25.22.4692

62. Van Oers, K., Sepers, B., Sies, W., Gawehns, F., Verhoeven, K. J. F., and Laine, V. N. (2020). Epigenetics of animal personality: DNA methylation cannot explain the heritability of exploratory behavior in a songbird. *Integrative and Comparative Biology*. doi:https://doi.org/10.1093/icb/icaa138.

63. Boyle, P., Clement, K., Gu, H. C., Smith, Z. D., Ziller, M., Fostel, J. L., . . . Meissner, A. (2012). Gel-free multiplexed reduced representation bisulfite sequencing for large-scale DNA methylation profiling. *Genome Biology, 13*(10), 10. doi:10.1186/gb-2012-13-10-R92

64. Ewels, P., Magnusson, M., Lundin, S., & Kaller, M. (2016). MultiQC: summarize analysis results for multiple tools and samples in a single report. *Bioinformatics, 32*(19), 3047-3048. doi:10.1093/bioinformatics/btw354

65. Krueger, F., & Andrews, S. R. (2011). Bismark: a flexible aligner and methylation caller for Bisulfite-Seq applications. *Bioinformatics, 27*(11), 1571-1572. doi:10.1093/bioinformatics/btr167

66. Akalin, A., Kormaksson, M., Li, S., Garrett-Bakelman, F. E., Figueroa, M. E., Melnick, A., & Mason, C. E. (2012). methylKit: a comprehensive R package for the analysis of genome-wide DNA methylation profiles. *Genome Biology, 13*(10), 9. doi:10.1186/gb-2012-13-10-R87

67. Feng, H., Conneely, K. N., & Wu, H. (2014). A Bayesian hierarchical model to detect differentially methylated loci from single nucleotide resolution sequencing data. *Nucleic Acids Research, 42*(8), 11. doi:10.1093/nar/gku154

68. Koestler, D. C., Avissar-Whiting, M., Houseman, E. A., Karagas, M. R., & Marsit, C. J. (2013). Differential DNA Methylation in Umbilical Cord Blood of Infants Exposed to Low Levels of Arsenic in Utero. *Environmental Health Perspectives, 121*(8), 971-977. doi:10.1289/ehp.1205925

69. Chen, H., Liu, J., Zhao, C. Q., Diwan, B. A., Merrick, B. A., & Waalkes, M. P. (2001). Association of c-myc overexpression and hyperproliferation with arsenite-induced malignant transformation. *Toxicology and Applied Pharmacology, 175*(3), 260-268. doi:10.1006/taap.2001.9253

70. Solomon, E.R., Caldwell, K.K., Allan, A.M. (2020). Developmental arsenic exposure is associated with differences in the epigenetic regulation of stress geners in the adult mouse frontal cortex. *Tocicology and Applied Pharmacology*, 391, 114920.

71. Rancourt, R.C., Schellong, K:, Tzschentke, B., Henrich, W., Plagemann, A. 2018. DNA methylation and expression of pro-optiomelanocorting (POMC) gene in the hypothalamus of three-week-old chickens show sex-specific differences. *Febs OpenBio* 8, 932-939.

72. Chen, J. H., Ren, X. Y., Li, L. M., Lu, S. Y., Chen, T., Tan, L. T., . . . Luo, W. (2019). Integrative Analyses of mRNA Expression Profile Reveal the Involvement of IGF2BP1 in Chicken Adipogenesis. *International Journal of Molecular Sciences, 20*(12), 15. doi:10.3390/ijms20122923

73. Hansen, T. V. O., Hammer, N. A., Nielsen, J., Madsen, M., Dalbaeck, C., Wewer, U. M., . . . Nielsen, F. C. (2004). Dwarfism and impaired gut development in insulin-like growth factor II mRNA-binding protein 1-deficient mice. *Molecular and Cellular Biology, 24*(10), 4448-4464. doi:10.1128/mcb.24.10.4448-4464.2004

74. Huang, X. W., Zhang, H., Guo, X. R., Zhu, Z. X., Cai, H. B., & Kong, X. Y. (2018). Insulin-like growth factor 2 mRNA-binding protein 1 (IGF2BP1) in cancer. *Journal of Hematology & Oncology, 11*, 15. doi:10.1186/s13045-018-0628-y

75. Noubissi, F. K., Yedjou, C. G., Spiegelman, V. S., & Tchounwou, P. B. (2018). Cross-Talk between Wnt and Hh Signaling Pathways in the Pathology of Basal Cell Carcinoma. *International Journal of Environmental Research and Public Health, 15*(7), 13. doi:10.3390/ijerph15071442

76. Hicar, M. D., Liu, Y. L., Allen, C. E., & Wu, L. C. (2001). Structure of the human zinc finger protein HIVEP3: Molecular cloning, expression, exon-intron structure, and comparison with paralogous genes HIVEP1 and HIVEP2. *Genomics, 71*(1), 89-100. doi:10.1006/geno.2000.6425

77. Lubecka, K., Flower, K., Beetch, M., Qiu, J., Kurzava, L., Buvala, H., . . . Stefanska, B. (2018). Loci-specific differences in blood DNA methylation in HBV-negative populations at risk for hepatocellular carcinoma development. *Epigenetics, 13*(6), 605-626. doi:10.1080/15592294.2018.1481706

78. Sarkar, D., Leung, E. Y., Baguley, B. C., Finlay, G. J., & Askarian-Amiri, M. E. (2015). Epigenetic regulation in human melanoma: past and future. *Epigenetics, 10*(2), 103-121. doi:10.1080/15592294.2014.1003746

79. Lea, A.J., Vilgalys, T.P., Durst, P.A.P., Tung, J. 2017. Maximizing ecological and evolutionary insight in bisulfite sequencing data sets. Nat Ecol Evol. 1(8): 1074–1083

80. Jones, P. A. (2012). Functions of DNA methylation: islands, start sites, gene bodies and beyond. *Nature Reviews Genetics, 13*(7), 484-492. doi:10.1038/nrg3230

81. Dixon, G., Liao, Y., Bay, L. K., & Matz, M. V. (2018). Role of gene body methylation in acclimatization and adaptation in a basal metazoan. *Proceedings of the National Academy of Sciences of the United States of America, 115*(52), 13342-13346. doi:10.1073/pnas.1813749115

82. Cao, Y., Yu, S. L., Wang, Y., Guo, G. Y., Ding, Q. A., & An, R. H. (2011). MicroRNA-dependent regulation of PTEN after arsenic trioxide treatment in bladder cancer cell line T24. *Tumor Biology, 32*(1), 179-188. doi:10.1007/s13277-010-0111-z

.