# A Draft Genome for *Hirudo verbana*, the Medicinal Leech

**Authors and Affiliations**

Riley T. Paulsen[1,2], Diing D.M. Agany[3], Jason Petersen[4,5], Christel M. Davis[4], Erik A. Ehli[4,6],

Etienne Gnimpieba[2,3], and Brian D. Burrell[1,2*]

[1]Division of Basic Biomedical Sciences and Center for Brain and Behavior Research, Sanford

School of Medicine, University of South Dakota, Vermillion, SD, USA

[2]Center for Brain and Behavior Research, University of South Dakota, SD, USA

[3]Biomedical Engineering Program, University of South Dakota, Sioux Falls, SD, USA

[4]Avera Institute for Human Genetics, Sioux Falls, SD, USA

[5]Department of Internal Medicine, Sanford School of Medicine, University of South Dakota,

Sioux Falls, SD, USA

[6]Department of Psychiatry, Sanford School of Medicine, University of South Dakota, Sioux

Falls, SD, USA

**\* Corresponding author**

**E-mail:** bburrell@usd.edu (BDB)

# Abstract

The medicinal leech, *Hirudo verbana*, is a powerful model organism for investigating fundamental neurobehavioral processes. The well-documented arrangement and properties of *H. verbana*'s nervous system allows changes at the level of specific neurons or synapses to be linked to physiological and behavioral phenomena. Juxtaposed to the extensive knowledge of *H. verbana's* nervous system is a limited, but recently expanding, portfolio of molecular and multi-omics tools. Together, the advancement of genetic databases for *H. verbana* will complement existing pharmacological and electrophysiological data by affording targeted manipulation and analysis of gene expression in neural pathways of interest. Here, we present the first draft genome assembly for *H. verbana*, which is approximately 250 Mbp in size and consists of 61,282 contigs. Whole genome sequencing was conducted using an Illumina sequencing platform followed by genome assembly with CLC-Bio Genomics Workbench and subsequent functional annotation. Ultimately, the diversity of organisms for which we have genomic information should parallel the availability of next generation sequencing technologies to widen the comparative approach to understand the involvement and discovery of genes in evolutionarily conserved processes. Results of this work hope to facilitate comparative studies with *H. verbana* and provide the foundation for future, more complete, genome assemblies of the leech.

# Introduction

The medicinal leech (*Hirudo verbana*) has been repurposed from an ancient bloodletting instrument [1] to a widely utilized invertebrate model system in biomedical research [2]. The well-documented and accessible central nervous system of the leech allows for precise selection of neurons for electrophysiological studies based on their characteristic morphologies, positioning, and biophysical properties [3, 4]. Fundamental discoveries have been made using *Hirudo* in a variety of disciplines that include central pattern generators, behavioral choice, learning and memory, synaptic signaling, neuroethology, neuro-injury and repair, and neurodevelopment [5-9]. Extensive research has also been devoted to examining the proteins secreted during leech hematophagy, which has longstanding applications in inflammation and coagulation [10]. Genomic insights into the medicinal leech will facilitate a more comprehensive approach into the evolutionary conservation of genes involved in the mechanistic processes that the medicinal leech has been used to help elucidate.

Despite these well-documented advantages of the medicinal leech for addressing various research questions, the leech lacks the molecular and genetic tools in comparison to alternative model organisms [11, 12]. For example, *Caenorhabditis elegans* and *Drosophila melanogaster* have extensive resources for targeted genome engineering in addition to optogenetic tools for electrophysiology and behavior manipulation [13-16]. Improving the genomic resources of organisms like the medicinal leech will promote more inclusive comparative genomics approaches to identifying conserved structural and functional gene signatures involved in human health and disease. Moreover, for many years, the medicinal leech community had been inadvertently aggregating four species of medicinal leeches: *H. medicinalis*, *H. verbana*, *H.*

*orientalis*, and *H. troctina* [17-20]. This misunderstanding regarding the taxonomic classification

of these leech subspecies has led to some confusion surrounding appropriate cataloging of

preliminary leech omics databases [21, 22]. Finally, in spite of the advancements in sequencing

technology, most of the existing sequence repositories for the medicinal leeches have been

comprised of expressed sequence tag [23] and transcriptomic databases [24-26], with recent

leech genomics work centering around *H. medicinalis* despite the prominence of *H. verbana* in

neuroscience research [27, 28]. Moreover, other leech genomes are also becoming available,

including that for the Asian Buffalo leech, *Hirudinaria manillensis* [29]. This work presents the

first draft genome for *H. verbana,* which consists of 250 Mbp, 61,282 contigs, an N50 of 8,638

bp, and a GC content of 38%. This draft genome, in addition to the growing transcriptomic

resources for *H. verbana* [24, 26, 30] and the aforementioned genomic databases for the closely

related leech species, will help accelerate studies seeking to link the molecular basis of previous

and ongoing functional studies utilizing medicinal leeches.

# Materials and Methods

Tissue collection and DNA extraction: High molecular weight genomic DNA was isolated from
muscle of three specimens of *H. verbana* (obtained from Niagara Leeches, Cheyenne, WY) in
separate preparations using the QIAamp DNA Mini Kit Cat. No. 51304 (Qiagen; Hilden,
Germany). In accordance with the protocol, 25 mg of dissected muscle tissue from the body wall
was used as input for each of the DNA extractions. The DNA from the three different isolations
was pooled, concentrated with Ampure XP beads (Beckman Coulter; Brea, CA) and 500 ng (10
ng/µL) was utilized for sequencing library preparation.

Library preparation: Sequencing libraries were prepared using the TruSeq Synthetic Long-Read
DNA Library Prep Kit (Illumina, Inc; San Diego CA). Three sequencing libraries were prepared
by loading the sample into a Covaris g-TUBE (part no. 520079), and centrifuging twice at 4200
x g for 1 minute. Briefly, the DNA was fragmented to approximately 8-10 kb and ligated with
adapters, which mark the end of contigs during data analysis. Following a dilution to limit the
number of DNA molecules in each well of a 384-well plate, long-range PCR was performed to
enrich for DNA fragments with appropriate adapters. The DNA in each well was treated with
the Nextera transposome, which fragments and simultaneously adds adapters to DNA. Indexing-
PCR was used to barcode the DNA in each well of the 384-well plate. The resulting products
were pooled and bead size-selection was performed. The average size of the final libraries was
~725 bp as measured with a High Sensitivity DNA chip on a 2100 Bioanalyzer (Agilent; Santa
Clara, CA). The concentration of each library was determined by quantitative PCR (qPCR) via
the KAPA Library Quantification Kit for Next Generation Sequencing (KAPA Biosystems;
Woburn, MA).

Whole genome sequencing: Libraries were normalized to 2 nmol/L in 10 mM Tris-Cl, pH 8.5

with 0.1% Tween 20.  Prior to cluster amplification, the libraries were denatured with 0.05 N

NaOH and diluted to 20 pmol/L.  Paired-end cluster generation of denatured templates was

performed according to the manufacturer's instructions (Illumina, San Diego, CA) utilizing the

HiSeq Rapid PE Cluster Kit v2 chemistry and flow cells.  Libraries were optimally clustered at

11 pmol/L with a 1% PhiX spike-in.  Sequencing-by-synthesis was performed on a HiSeq 2500

utilizing v2 chemistry with paired-end 101 bp reads and an 8 bp index read.

Long read and genome assembly: A total of 1,862,297,140 bp of 2 x 101 bp reads were obtained

from three flow cells. Sequence read data were processed and converted to short-read FASTQ

format by Illumina BaseSpace analysis software (v2.0.13).  The short reads from each plate were

individually processed in three runs to construct primary contigs using the TruSPAdes assembly

software (v1.1.0) [31], which were merged into one assembly using CLC-Bio Genomics

Workbench De Novo Assembly (Qiagen, v11.0.1) with default parameters. Thorough quality

control was performed on the raw short read data using FastQC

(http://www.bioinformatics.babraham.ac.uk/projects/fastqc/) to assess the Phred score, presence

of repeat reads, non-nucleotide content, GC content, and duplicated read contents. The quality of

the primary contigs assembled by the TruSPAdes algorithm were assessed by Quast [32].

Repeat sequence and BUSCO annotation: Repeatmasker [33] was used to annotate repeating

sequences and transposable elements in the H. verbana genome assembly. Repeatmasker was

configured with the pooled databases RepBase-2017 [34] and Dfam-2017 consensus databases

[35], and Tandem Repeats Finder, ran with RMBlast (v2.2.27) search engine. Additionally, the

completeness and quality of the draft genome was evaluated using a BUSCO (Benchmarking

Universal Single-Copy Orthologs) assessment that matched our newly assembled sequences to

the metazoan OrthoDB v9.1 [36-38]. BUSCO.v3 was configured using AUGUSTUS gene

predictor [39], HMMER [40, 41], and NCBI-BLAST+ [42].

<u>Functional annotation and orthologous analysis</u>: In order to elucidate functional annotation and

gene ontology annotation, NCBI Blast+, UniprotKB [43], and the Blast2GO software suite [44]

integrated with InterProScan [45] were implemented. Furthermore, through locally constructed

databases in CLC-Bio Genomics Workbench, we utilized NCBI Blastn [46] on our genome

against closely related databases for the following closely related polychaete annelids: *H.*

*medicinalis* (GenBank: EY478949-EY505781) , *Helobdella robusta* (GCA_000326865.1), and

*Capitella teleta* (GCA_000328365.1).

<u>Gene prediction and macro-synteny analysis</u>: Gene predictions were performed using the

MAKER2 [47] genome annotation pipeline with SNAP [48] against the nematode

*Caenorhabditis elegans* and two other annelids: *C. teleta*, and *H. robusta*. The *ab-initio* gene

predictor, SNAP, was trained three times to improve performance. The *H. verbana* draft genome

assembly was aligned to the *C. elegans* genome and scaffolds of *H. robusta*. Circos [49] was

used to generate the circular genome alignment figures to analyze the anchoring of the top 600

*H. verbana* contigs onto the six chromosomes of *C. elegans* [50]. Additonally, the draft genome

top 600 contigs were mapped and anchored to the 1,991 scaffolds of the *H. robusta* genome.

Phylogenetic reconstruction: OrthoFinder2 [51] was utilized for comparative genomics between

our draft genome for *H. verbana*, and the protein sequence databases for six other organisms: *H.*

*medicinalis*, *H. robusta*, *C. teleta*, *C. elegans* (GCA_000002985.3), and the chordates *Mus*

*musculus* (GCA_000001635.8) and *Homo sapiens* (GCA_000001405.27). Orthofinder was

configured with the DIAMOND search engine [52], MCL clustering algorithm [53], and

FastTree [54] to construct the rooted phylogenetic tree which was visualized with Phylo.io [55].

Species Identification: The presence of mitochondrial contigs was assessed by comparing the

draft mitochondrial genomes for *H. medicinalis* and *H. verbana* [56] to our draft genome. NCBI

Blastn [46] was used on our draft genome against the mitochondrial DNA sequences

downloaded from NCBI GenBank. The phylogenetic relationships between the mitochondrial

contigs present in our draft genome, *H. medicinalis*, *H. verbana*, as well as other Hirudinea,

*Whitmania pigra* [57] and *Hirudo nipponia* [58] via their mitochondrial genomes was also

analyzed. The evolutionary analysis was performed in MEGA X (v11.0) [59, 60] using Multiple

Sequence Comparison by Log- Expectation (MUSCLE) [61], and the phylogenetic tree was

constructed using the Maximum Likelihood method with the Tamura-Nei model [62] and node

confidence was assessed with a 500 bootstrap trials.

## Results and Discussion

Next-generation sequencing leveraging an Illumina HiSeq-2500 platform was employed to construct the first draft genome for *H. verbana*, the medicinal leech. A total of 188 Gbp were generated that encompassed 1,862,297,140 bp of 101bp x 2 paired short reads (Table 1A). Quality control of the raw short reads performed by FastQC and MultiQC [63] revealed that the data had an average phred score >30 (S1 Appendix). The short reads were barcode assembled individually for each plate using TruSPAdes assembler software into TruSeq synthetic long reads. The long reads (Table 1B) were 190,514 bp, 198,741 bp, and 193,658 bp for each plate, respectively, and had an average N50 of 7623 bp. Together, these synthetic long reads consisted of 582,913 sequences, 3,429,493,670 bp, and had an estimated coverage of 6.9X. Quast assessment of both the synthetic long reads and final assembly maintained a phred score of 30. The draft genome was constructed from the TruSeq synthetic long reads using CLC-Bio Genomics Workbench to produce the draft genome assembly for *H. verbana*. Prior to arriving at the final assembly that used a combination of TruSPAdes synthetic long reads and CLC-Bio Genomics Workbench, we performed a thorough assessment of multiple assemblers using long reads formed both by Illumina BaseSpace analysis software and TruSPAdes in conjunction with SOAPdenovo2 [64], Megahit [65], Spades [66], Ray [67], and Velvet [68]. Ultimately, we are reporting the TruSPAdes and CLC-Bio assembly approach because it had the most coverage and performed best under downstream assessment described below.

176     **Table 1: Statistics of the *de novo* draft genome assembly for *Hirudo verbana***

| (A) Genome sequencing reads obtained from Illumina HiSeq via Basespace | | | |
|---|---|---|---|
| | Reads (Single, 101bp) | Reads (Pair, 101bp x 2) | Bases (Pair) |
| Plate 1 | 266,688,108 | 533,376,216 | 53,870,997,816 |
| Plate 2 | 328,593,254 | 657,186,508 | 66,375,837,308 |
| Plate 3 | 335,867,208 | 671,734,416 | 67,845,176,016 |
| Total reads | 931,148,570 | 1,862,297,140 | 188,092,011,140 |
| (B) TruSPAdes Truseq Synthetic Long-Read Assembly Statistics | | | |
| | Assembled reads | Total bases | N50 |
| Plate 1 | 190,514 | 1,117,298,449 | 7,612 |
| Plate 2 | 198,741 | 1,172,622,590 | 7,634 |
| Plate 3 | 193,658 | 1,139,572,631 | 7,624 |
| (C) Summary details of *Hirudo verbana* draft genome assembly | | | |
| Estimated genome size | | 250 Mb | |
| Total assembly length | | 250 Mb | |
| Total sequences | | 582,913 | |
| Short read coverage | | 627X | |
| Long read coverage | | 6.9X | |
| Contigs | | 61,282 | |
| N75 | | 4807 bp | |
| N50 | | 8638 bp | |
| N25 | | 14800 bp | |
| Minimum contig | | 200 bp | |
| Maximum contig | | 154993 bp | |
| Average | | 4084 bp | |
| Nucleotide frequency | | | |
| Adenine (A) | | 30.90% | |
| Cytosine (C) | | 19.10% | |
| Guanine (G) | | 19.00% | |
| Thymine (T) | | 31.00% | |

(A) Whole genome sequencing reads obtained from Illumina HiSeq for H. verbana de novo draft
genome assembly (B) Statistics for barcode-assembled synthetic long reads generated using
TruSPAdes (C) Summary of assembly statistics of the draft genome for H. verbana.

The final draft genome assembly presented here is 250,270,938 bp in size, which is

comparable to the genome assembly statistics for other annelids *C. teleta* (240 Mbp) and *H.*

*robusta* (310 Mbp). The genome assembly consists of 61,282 contigs that have a minimum

length of 200 bp, a maximum length of 154,993 bp, and an N50 of 8,638 bp (Table 1C). N50 is

important in an assembly assessment as it measures contigs lengths distribution and informs on

the internal parameter of the assembly. However, it does not inform or guarantee the

goodness/completeness of an assembly [69]. For example, filtering out of a shortest contigs or

the scaffolding that links contigs could increase the N50 without improving the assembly

completeness. Also, when analyzing the N50, a comparison should be made between draft

genomes of a similar length, or it will be misleading [70]. In this assembly, we picked the

assembly run with 200 bp parameter setting as the shortest contig cutoff. While assemblies were

tested with higher cutoffs, specifically 500 bp and 1000 bp, the chosen setting produced the best

outcome in in terms of balance of genome size coverage and N50 criteria. A condensed summary

comparing our draft genome to the 10 available annelid genomes is also provided in S1

Table.[27-29, 71-76] For preliminary validation of the quality of the draft genome assembly, the

61,282 contigs were mapped back to the raw short reads using the mapping module in CLC-Bio

Genomics Workbench. The result demonstrated that 86.72% of the assembly mapped back onto

the short reads, leaving 13.28% unmapped. Among the contigs that were reported to map back,

85.77% had identical base pair matching.

Next, the repeating segments and transposable elements of the draft genome were

annotated using RepeatMasker. An estimated 6.67% of the genome assembly (16,685,142 bp of

the total 250 Mbp) was determined to be repetitive or transposable elements, with a majority

consisting of simple repeats, interspersed repeats, and low complexity repeats (S2 Table).

Moreover, the completeness of the draft genome was assessed with a BUSCO analysis for the

presence of metazoan-specific orthologues. The metazoan BUSCO that we implemented

consisted of 978 genes, and our assembly returned 809 (82.70%) as complete, 533 (54.50%) as

complete and single-copy, 276 (28.20%) complete and duplicated, 70 (7.20%) fragmented, and

99 (10.10%) as missing (Table 2A). Overall, our genome has a completeness score of 89.9%

(82.70% complete + 7.20% fragmented).

**Table 2. Analysis of the completeness and orthology of the *H. verbana* genome.**

| (A) BUSCO statistics of genome based on 978 metazoan-conserved genes | | |
|---|---|---|
| BUSCO | Genes Present | Percentage (%) |
| Complete BUSCOs (C) | 809 | 82.70% |
| Complete and Single-copy BUSCOs (S) | 533 | 54.50% |
| Complete and duplicated BUSCOs (D) | 276 | 28.20% |
| Fragmented BUSCOs (F) | 70 | 7.20% |
| Missing BUSCOs (M) | 99 | 10.10% |
| (B) Genome functional annotation | | |
| | Number of contigs | Percentage (%) |
| Total | 61,282 | |
| Annotated | | |
| InterProScan | 5,000 | 8.16% |
| GO | 2,800 | 4.57% |
| UniProtKB | 44,000 | 71.80% |
| Unannotated | 9,482 | 15.47% |
| (C) Comparison of gene prediction and annotation to related organisms | | |
| Species | Size of genome assembly (Mbp) | Predicted number of genes |
| *Lottia gigantea* | 348 | 23,800 |
| *Capitella teleta* | 324 | 32,389 |
| *Helobdella robusta* | 228 | 23,400 |
| *Hirudo verbana* | 250 | 26,210 |

211    (A) BUSCO statistics assessing the completeness of the H. verbana draft genome based on 978
212    metazoan-conserved genes (B) Summary of structural functional annotation for H. verbana draft
213    genome (C) Comparison of H. verbana draft genome size and predicted number of genes to 3
214    closely related spiralian genomes (two annelids and one mollusc).

215  The sequence homology and similarity of the draft genome for *H. verbana* was assessed

216 by NCBI BLAST+ against the genomic and transcriptomic sequences available for *H.*

217 *medicinalis*, *H. robusta*, and *C. teleta*. Approximately 94% of the draft genome sequences had an

218 identity match within the queried databases, 5.5% exhibited at least 70% similarity, and the

219 remaining 0.5% was unidentified. Functional annotation was assessed using fast-BlastX [77]

220 against the Animalia NCBI Refseq database in Blast2GO [78]. From total assembly, 1,178

221 contigs returned significant blast hits at an e-value threshold of $10^{-10}$. The top 20 gene ontology

222 (GO) terms [79] for each classification – biological process (BP), molecular function (MF), and

223 cellular component (CC) – at GO level 5 are displayed in Fig 1. Moreover, the draft genome was

224 aligned to the genome of *C. elegans*. The draft genome contigs for *H. verbana* were mapped and

225 anchored to the 6 chromosomes of *C. elegans* (Fig 2A) and to the *H. robusta* scaffolds at

226 position 15000-17000 for contig *hv7* on scaffold *he49* (Fig 2B). A majority of the mapped

227 sequences achieved better fit onto chromosomes 1 and X of the *C. elegans* genome (S3 Table).

228  Using OrthoFinder, orthologues were generated from gene families of our draft genome

229 for *H. verbana*, *H. medicinalis*, *H. robusta*, C. *teleta*, *C. elegans*, *M. musculus*, and *H. sapiens*.

230 The phylogenetic tree was reconstructed using OrthoFinder after it identified the highest

231 similarity content between the draft genome and the reference organisms. The phylogenetic tree

232 (Fig 3A) appropriately placed *H. verbana* adjacent to its closest relative, *H. medicinalis*,

233 demonstrating their last known divergence in genus *Hirudo*. Predictive protein-coding gene

234 sequences were identified based on conserved protein signatures and domains with UniprotKB,

235 Blast2Go, and InterProScan. A total of 84.53% of the contigs were annotated for a protein-

236 coding function, of these, 8.16% were identified by InterProScan, 4.57% by Blast2GO, and

237 71.80% by UniProtKB (Table 2B). Lastly, two-pass annotation in MAKER predicted 26,210

protein-coding genes in the draft genome, which is similar to the first reports of draft genomes

for fellow lophotrochozoans *C. teleta*, *H. robusta*, and *L. gigantea,* a gastropod mollusc (Table

2C).

For further confidence that the leech used in this study were *H. verbana*, we searched our

draft genome for mitochondrial contigs using the draft mitochondrial genomes for *H. medicinalis*

and *H. verbana*. The results of the mitochondrial contig analysis suggest that the leeches used to

collect genomic DNA in our study were of the species *H. verbana*, which is emphasized by the

mitochondrial contigs found in our draft genome forming a clade with the *H. verbana* draft

mitochondrial genome in the phylogenetic tree (Fig 3B). The results of the mitochondrial

phylogenetic analysis, in addition to the characteristic coloration patterns observed on the dorsal

side of the leeches, indicate that the leeches used in this study were *H. verbana*.

This study is the first to publish an annotated draft genome sequence for the medicinal

leech, *H. verbana*. Overall, the genome assembly consists of 250 Mbp and 61,282 contigs,

84.53% of which have been predicted to contain a protein-coding function. The draft genome is

also predicted to contain 26,210 protein-coding genes and a repetitive content of 6.67%. The raw

short-read sequence data, synthetic long-reads, and assembled contigs for the present study have

been deposited into NCBI accession number JAIQDV000000000. The draft genome assembly

will assist in providing tools to understand the underlying molecular processes involved in

ongoing studies in neurophysiology, developmental biology, and neuroethology that utilize *H.*

*verbana* [80-83]. Whole-genome characterization for Hirudinae *H. medicinalis* [27, 28], *H.*

*manillensis* [29] and *H. verbana* is expanding and will help distinguish and clarify distinct

genetic undertones of these previously amalgamated species. Future efforts to better annotate and

complete a genome for *H. verbana* will enable insight into genetic mechanisms of processes

investigated with this model organism [83-85], and advance more robust cross-species validation

of comparative principles in next-generation biomedical research techniques.

## 263 Acknowledgements

This work was supported by a pilot grant from the University of South Dakota (USD) Center for

Brain and Behavior Research (CBBRe), the USD Neuroscience, Nanotechnology, and Networks

Program (DGE-1633213), and the National Science Foundation Graduate Research Fellowship

(DGE-1545679). Authors acknowledge the use of computational resources of the USD HPC

cluster and the Extreme Science and Engineering Discovery Environment (XSEDE), which is

supported by the National Science Foundation grant number ACI-1548562.

# References

1. Phillips, A.J. and M.E. Siddall, *Poly-paraphyly of Hirudinidae: many lineages of medicinal leeches.* BMC Evol Biol, 2009. **9**: p. 246.

2. Wagenaar, D.A., *A classic model animal in the 21st century: recent lessons from the leech nervous system.* J Exp Biol, 2015. **218**(Pt 21): p. 3353-9.

3. Nicholls, J.G. and D.A. Baylor, *Specific modalities and receptive fields of sensory neurons in CNS of the leech.* J Neurophysiol, 1968. **31**(5): p. 740-56.

4. Kristan, W.B., *Neuronal changes related to behavioral changes in chronically isolated segments of the medicinal leech.* Brain Research, 1979. **167**(1): p. 215-220.

5. Kristan, W.B., Jr., R.L. Calabrese, and W.O. Friesen, *Neuronal control of leech behavior.* Prog Neurobiol, 2005. **76**(5): p. 279-327.

6. Crisp, K.M. and B.D. Burrell, *Cellular and Behavioral Properties of Learning in Leech and Other Annelids* in *Annelids in Modern Biology*, D.H. Shain, Editor. 2009, John Wiley & Sons: New York.

7. Duan, Y., et al., *Repair and regeneration of functional synaptic connections: cellular and molecular interactions in the leech.* Cell Mol Neurobiol, 2005. **25**(2): p. 441-50.

8. Eisenhart, F.J., T.W. Cacciatore, and W.B. Kristan, Jr., *A central pattern generator underlies crawling in the medicinal leech.* J Comp Physiol A, 2000. **186**(7-8): p. 631-43.

9. Yuan, S. and B.D. Burrell, *Nonnociceptive afferent activity depresses nocifensive behavior and nociceptive synapses via an endocannabinoid-dependent mechanism.* J Neurophysiol, 2013. **110**(11): p. 2607-16.

292    10.     Lemke, S., et al., *May salivary gland secretory proteins from hematophagous leeches*

293             *(Hirudo verbana) reach pharmacologically relevant concentrations in the vertebrate*

294             *host?* PLoS One, 2013. **8**(9): p. e73809.

295     11.     Nobrega, M.A. and L.A. Pennacchio, *Comparative genomic analysis as a tool for*

296             *biological discovery.* J Physiol, 2004. **554**(Pt 1): p. 31-9.

297     12.     Clarke, S.L., et al., *Human developmental enhancers conserved between deuterostomes*

298             *and protostomes.* PLoS Genet, 2012. **8**(8): p. e1002852.

299     13.     Dickinson, D.J. and B. Goldstein, *CRISPR-Based Methods for Caenorhabditis elegans*

300             *Genome Engineering.* Genetics, 2016. **202**(3): p. 885-901.

301     14.     Gratz, S.J., et al., *CRISPR-Cas9 Genome Editing in Drosophila.* Curr Protoc Mol Biol,

302             2015. **111**: p. 31 2 1-31 2 20.

303     15.     Simpson, J.H. and L.L. Looger, *Functional Imaging and Optogenetics in Drosophila.*

304             Genetics, 2018. **208**(4): p. 1291-1309.

305     16.     Leifer, A.M., et al., *Optogenetic manipulation of neural activity in freely moving*

306             *Caenorhabditis elegans.* Nat Methods, 2011. **8**(2): p. 147-52.

307     17.     Kutschera, U. and J. Elliott, *The European medicinal leech Hirudo medicinalis L.:*

308             *Morphology and occurrence of an endangered species.* Zoosystematics and Evolution,

309             2014. **90**(2): p. 271-280.

310     18.     Siddall, M.E., et al., *Diverse molecular data demonstrate that commercially available*

311             *medicinal leeches are not Hirudo medicinalis.* Proc Biol Sci, 2007. **274**(1617): p. 1481-7.

312     19.     Petrauskiené, L., O. Utevska, and S. Utevsky, *Can different species of medicinal leeches*

313             *(Hirudospp.) interbreed?* Invertebrate Biology, 2009. **128**(4): p. 324-331.

20.    Trontelj, P. and S.Y. Utevsky, *Phylogeny and phylogeography of medicinal leeches*
*(genus Hirudo): fast dispersal and shallow genetic structure.* Mol Phylogenet Evol, 2012.
**63**(2): p. 475-85.

21.    Trontelj, P. and S.Y. Utevsky, *Celebrity with a neglected taxonomy: molecular*
*systematics of the medicinal leech (genus Hirudo).* Mol Phylogenet Evol, 2005. **34**(3): p.
616-24.

22.    Bely, A.E. and D.A. Weisblat, *Lessons from leeches: a call for DNA barcoding in the*
*lab.* Evol Dev, 2006. **8**(6): p. 491-501.

23.    Macagno, E.R., et al., *Construction of a medicinal leech transcriptome database and its*
*application to the identification of leech homologs of neural and innate immune genes.*
BMC Genomics, 2010. **11**: p. 407.

24.    Northcutt, A.J., et al., *An annotated CNS transcriptome of the medicinal leech, Hirudo*
*verbana: De novo sequencing to characterize genes associated with nervous system*
*activity.* PLoS One, 2018. **13**(7): p. e0201206.

25.    Hibsh, D., et al., *De novo transcriptome assembly databases for the central nervous*
*system of the medicinal leech.* Sci Data, 2015. **2**: p. 150015.

26.    Heath-Heckman, E., et al., *Transcriptional profiling of identified neurons in leech.* BMC
Genomics, 2021. **22**(1): p. 215.

27.    Babenko, V.V., et al., *Draft genome sequences of Hirudo medicinalis and salivary*
*transcriptome of three closely related medicinal leeches.* BMC Genomics, 2020. **21**(1): p.
331.

28.    Kvist, S., et al., *Draft genome of the European medicinal leech Hirudo medicinalis*

336           *(Annelida, Clitellata, Hirudiniformes) with emphasis on anticoagulants.* Sci Rep, 2020.

337           **10**(1): p. 9885.

338    29.    Guan, D.L., et al., *Draft Genome of the Asian Buffalo Leech Hirudinaria manillensis.*

339           Front Genet, 2019. **10**: p. 1321.

340    30.    Stowasser, A., et al., *Electrophysiology and transcriptomics reveal two photoreceptor*

341           *classes and complex visual integration in Hirudo verbana.* J Exp Biol, 2019. **222**(Pt 15).

342    31.    Bankevich, A. and P.A. Pevzner, *TruSPAdes: barcode assembly of TruSeq synthetic long*

343           *reads.* Nat Methods, 2016. **13**(3): p. 248-50.

344    32.    Gurevich, A., et al., *QUAST: quality assessment tool for genome assemblies.*

345           Bioinformatics, 2013. **29**(8): p. 1072-5.

346    33.    Tarailo-Graovac, M. and N. Chen, *Using RepeatMasker to identify repetitive elements in*

347           *genomic sequences.* Curr Protoc Bioinformatics, 2009. **Chapter 4**: p. Unit 4 10.

348    34.    Jurka, J., et al., *Repbase Update, a database of eukaryotic repetitive elements.* Cytogenet

349           Genome Res, 2005. **110**(1-4): p. 462-7.

350    35.    Hubley, R., et al., *The Dfam database of repetitive DNA families.* Nucleic Acids Res,

351           2016. **44**(D1): p. D81-9.

352    36.    Simao, F.A., et al., *BUSCO: assessing genome assembly and annotation completeness*

353           *with single-copy orthologs.* Bioinformatics, 2015. **31**(19): p. 3210-2.

354    37.    Waterhouse, R.M., et al., *BUSCO Applications from Quality Assessments to Gene*

355           *Prediction and Phylogenomics.* Mol Biol Evol, 2018. **35**(3): p. 543-548.

356    38.    Zdobnov, E.M., et al., *OrthoDB v9.1: cataloging evolutionary and functional annotations*

357           *for animal, fungal, plant, archaeal, bacterial and viral orthologs.* Nucleic Acids Res,

358           2017. **45**(D1): p. D744-D749.

359    39.    Stanke, M., A. Tzvetkova, and B. Morgenstern, *AUGUSTUS at EGASP: using EST,*

360           *protein and genomic alignments for improved gene prediction in the human genome.*

361           Genome Biol, 2006. **7 Suppl 1**: p. S11 1-8.

362    40.    Potter, S.C., et al., *HMMER web server: 2018 update.* Nucleic Acids Res, 2018. **46**(W1):

363           p. W200-W204.

364    41.    Finn, R.D., J. Clements, and S.R. Eddy, *HMMER web server: interactive sequence*

365           *similarity searching.* Nucleic Acids Res, 2011. **39**(Web Server issue): p. W29-37.

366    42.    Camacho, C., et al., *BLAST+: architecture and applications.* BMC Bioinformatics, 2009.

367           **10**: p. 421.

368    43.    UniProt, C., *UniProt: a worldwide hub of protein knowledge.* Nucleic Acids Res, 2019.

369           **47**(D1): p. D506-D515.

370    44.    Gotz, S., et al., *High-throughput functional annotation and data mining with the*

371           *Blast2GO suite.* Nucleic Acids Res, 2008. **36**(10): p. 3420-35.

372    45.    Jones, P., et al., *InterProScan 5: genome-scale protein function classification.*

373           Bioinformatics, 2014. **30**(9): p. 1236-40.

374    46.    Johnson, M., et al., *NCBI BLAST: a better web interface.* Nucleic Acids Res, 2008.

375           **36**(Web Server issue): p. W5-9.

376    47.    Holt, C. and M. Yandell, *MAKER2: an annotation pipeline and genome-database*

377           *management tool for second-generation genome projects.* BMC Bioinformatics, 2011.

378           **12**: p. 491.

48.  Korf, I., *Gene finding in novel genomes.* BMC Bioinformatics, 2004. **5**: p. 59.

49.  Krzywinski, M., et al., *Circos: an information aesthetic for comparative genomics.*

Genome Res, 2009. **19**(9): p. 1639-45.

50.  Hillier, L.W., et al., *Genomics in C. elegans: so many genes, such a little worm.* Genome

Res, 2005. **15**(12): p. 1651-60.

51.  Emms, D.M. and S. Kelly, *OrthoFinder: phylogenetic orthology inference for*

*comparative genomics.* Genome Biol, 2019. **20**(1): p. 238.

52.  Buchfink, B., C. Xie, and D.H. Huson, *Fast and sensitive protein alignment using*

*DIAMOND.* Nat Methods, 2015. **12**(1): p. 59-60.

53.  Enright, A.J., S. Van Dongen, and C.A. Ouzounis, *An efficient algorithm for large-scale*

*detection of protein families.* Nucleic Acids Res, 2002. **30**(7): p. 1575-84.

54.  Price, M.N., P.S. Dehal, and A.P. Arkin, *FastTree 2--approximately maximum-likelihood*

*trees for large alignments.* PLoS One, 2010. **5**(3): p. e9490.

55.  Robinson, O., D. Dylus, and C. Dessimoz, *Phylo.io: Interactive Viewing and Comparison*

*of Large Phylogenetic Trees on the Web.* Mol Biol Evol, 2016. **33**(8): p. 2163-6.

56.  Nikitina, A., et al., *Draft mitochondrial genomes of Hirudo medicinalis and Hirudo*

*verbana (Annelida, Hirudinea).* Mitochondrial DNA B Resour, 2016. **1**(1): p. 254-256.

57.  Shen, X., et al., *The complete mitochondrial genome sequence of Whitmania pigra*

*(Annelida, Hirudinea): the first representative from the class Hirudinea.* Comp Biochem

Physiol Part D Genomics Proteomics, 2011. **6**(2): p. 133-8.

58.  Xu, Y., et al., *Complete mitochondrial genome of Hirudo nipponia (Annelida,*

*Hirudinea).* Mitochondrial DNA A DNA Mapp Seq Anal, 2016. **27**(1): p. 257-8.

59.  Kumar, S., et al., *MEGA X: Molecular Evolutionary Genetics Analysis across Computing*

*Platforms.* Mol Biol Evol, 2018. **35**(6): p. 1547-1549.

60.  Stecher, G., K. Tamura, and S. Kumar, *Molecular Evolutionary Genetics Analysis*

*(MEGA) for macOS.* Mol Biol Evol, 2020. **37**(4): p. 1237-1239.

61.  Edgar, R.C., *MUSCLE: multiple sequence alignment with high accuracy and high*

*throughput.* Nucleic Acids Res, 2004. **32**(5): p. 1792-7.

62.  Tamura, K., *Estimation of the number of nucleotide substitutions when there are strong*

*transition-transversion and G+C-content biases.* Mol Biol Evol, 1992. **9**(4): p. 678-87.

63.  Ewels, P., et al., *MultiQC: summarize analysis results for multiple tools and samples in a*

*single report.* Bioinformatics, 2016. **32**(19): p. 3047-8.

64.  Luo, R., et al., *SOAPdenovo2: an empirically improved memory-efficient short-read de*

*novo assembler.* Gigascience, 2012. **1**(1): p. 18.

65.  Li, D., et al., *MEGAHIT: an ultra-fast single-node solution for large and complex*

*metagenomics assembly via succinct de Bruijn graph.* Bioinformatics, 2015. **31**(10): p.

1674-6.

66.  Bankevich, A., et al., *SPAdes: a new genome assembly algorithm and its applications to*

*single-cell sequencing.* J Comput Biol, 2012. **19**(5): p. 455-77.

67.  Boisvert, S., F. Laviolette, and J. Corbeil, *Ray: simultaneous assembly of reads from a*

*mix of high-throughput sequencing technologies.* J Comput Biol, 2010. **17**(11): p. 1519-

33.

68.  Zerbino, D.R. and E. Birney, *Velvet: algorithms for de novo short read assembly using de*

*Bruijn graphs.* Genome Res, 2008. **18**(5): p. 821-9.

69.    Narzisi, G. and B. Mishra, *Comparing De Novo Genome Assembly: The Long and Short*

424           *of It.* PLOS ONE, 2011. **6**(4): p. e19175.

425    70.    Thrash, A., F. Hoffmann, and A. Perkins, *Toward a more holistic method of genome*

426           *assembly assessment.* BMC Bioinformatics, 2020. **21**(4): p. 249.

427    71.    Simakov, O., et al., *Insights into bilaterian evolution from three spiralian genomes.*

428           Nature, 2013. **493**(7433): p. 526-31.

429    72.    Martin-Duran, J.M., et al., *Conservative route to genome compaction in a miniature*

430           *annelid.* Nat Ecol Evol, 2021. **5**(2): p. 231-242.

431    73.    Shikuma, N.J., et al., *Stepwise metamorphosis of the tubeworm Hydroides elegans is*

432           *mediated by a bacterial inducer and MAPK signaling.* Proc Natl Acad Sci U S A, 2016.

433           **113**(36): p. 10097-102.

434    74.    Cunha, L., et al., *A large set of microsatellites for the highly invasive earthworm*

435           *Amynthas corticis predicted from low coverage genomes.* Applied Soil Ecology, 2017.

436           **119**: p. 152-155.

437    75.    Li, Y., et al., *Genomic adaptations to chemosymbiosis in the deep-sea seep-dwelling*

438           *tubeworm Lamellibrachia luymesi.* BMC Biol, 2019. **17**(1): p. 91.

439    76.    Bhambri, A., et al., *Large scale changes in the transcriptome of Eisenia fetida during*

440           *regeneration.* PLoS One, 2018. **13**(9): p. e0204234.

441    77.    Price, M.N., P.S. Dehal, and A.P. Arkin, *FastBLAST: homology relationships for millions*

442           *of proteins.* PLoS One, 2008. **3**(10): p. e3589.

443    78.    O'Leary, N.A., et al., *Reference sequence (RefSeq) database at NCBI: current status,*

444           *taxonomic expansion, and functional annotation.* Nucleic Acids Res, 2016. **44**(D1): p.

445           D733-45.

446    79.    Harris, M.A., et al., *The Gene Ontology (GO) database and informatics resource.*

447           Nucleic Acids Res, 2004. **32**(Database issue): p. D258-61.

448    80.    Harley, C.M. and D.A. Wagenaar, *Scanning behavior in the medicinal leech Hirudo*

449           *verbana.* PLoS One, 2014. **9**(1): p. e86120.

450    81.    Yazdani, N., et al., *Expression of a dominant negative mutant innexin in identified*

451           *neurons and glial cells reveals selective interactions among gap junctional proteins.* Dev

452           Neurobiol, 2013. **73**(8): p. 571-86.

453    82.    Yuan, S. and B.D. Burrell, *Endocannabinoid-dependent long-term depression in a*

454           *nociceptive synapse requires coordinated presynaptic and postsynaptic transcription and*

455           *translation.* J Neurosci, 2013. **33**(10): p. 4349-58.

456    83.    Baranzini, N., et al., *Antimicrobial Role of RNASET2 Protein During Innate Immune*

457           *Response in the Medicinal Leech Hirudo verbana.* Front Immunol, 2020. **11**: p. 370.

458    84.    Grey, K.B., B.L. Moss, and B.D. Burrell, *Molecular identification and expression of the*

459           *NMDA receptor NR1 subunit in the leech.* Invert Neurosci, 2009. **9**(1): p. 11-20.

460    85.    Kabeiseman, E., R. Paulsen, and B.D. Burrell, *Characterization of a monoacylglycerol*

461           *lipase in the medicinal leech, Hirudo verbana.* Comp Biochem Physiol B Biochem Mol

462           Biol, 2020. **243-244**: p. 110433.

## Figure Legends

**Figure 1. *H. verbana* draft genome gene ontology distribution.** Displayed are the number of sequences present corresponding to the top 20 most frequent annotations for each classification. Gene ontologies shown are grouped into the major categories of: biological process (BP, green), molecular function (MF, blue), and cellular component (CC, yellow) at a gene ontology level of 5.

**Figure 2. Alignment of *H. verbana* draft genome to *C. elegans* and *H. robusta*.** (A) Alignment (>100 bp) of top 600 *H. verbana* draft genome contigs to the chromosomes (I-X) of *C. elegans*. (B) Alignment (>100 bp) of the top 600 *H. verbana* draft genome contigs to the scaffolds of the *H. robusta* genome assembly. *H. verbana* contigs are in black and abbreviated as "Hv," the *C. elegans* chromosomes are abbreviated "ce," and the *H. robusta* scaffolds are abbreviated "he." Both the corresponding chromosomes and scaffolds are colored, and the projections are anchored to different contigs from the draft genome.

**Figure 3. Genomic and mitochondrial phylogenetic relationships.** (A) Reconstructed phylogenic tree based on orthologous gene families. (B) Phylogeny of our draft genome in relation to 4 leech species by comparing the presence and similarity of mitochondrial contigs through the Maximum Likelihood method via the Tamura 3-parameter model. NCBI accession numbers are provided next to each given taxa name.

# Supplementary Figure/Tables

**Supplementary Figure 1.** Quality Control from Sequencing to Final Assembly Stage. (A)

Integrated quality assessment for short sequencing reads generated by FastQC and MultiQC (B)

GC content for final assembly and (C) TruSPAdes synthetic long reads after barcode assembly.

**Supplementary Table 1. Annelid Genome Comparison**. Comparison of select assembly

statistics at the contig level (number of contigs, genome length, N50, and GC content) for

annelid genomes. Accession numbers are provided, and the mode of sequencing technology and

assembly software are also included for comparison.

**Supplementary Table 2. Repeat sequence and transposable element composition in the**

**draft genome of *H. verbana*.** Identification of repetitive elements in the *H. verbana* draft

genome using RepeatMasker.

**Supplementary Table 3. Chromosome rearrangement of *C. elegans* onto *H. verbana* contigs.**

Mapping of *H. verbana* draft genome contigs onto the chromosome scaffolds of *C. elegans*.

# GO Distribution by Level (5) – Top 20