Short title: Vacuolar hexose transport and xylem development

* Corresponding author: Rozenn Le Hir

Telephone: +33 1 30 83 30 57

Fax: +33 1 30 83 30 96

Email : rozenn.le-hir@inrae.fr

ORCID ID: 0000-0001-6076-5863

A vacuolar hexose transport is required for xylem development in the inflorescence stem
of Arabidopsis.

Emilie Aubry[1,2], Beate Hoffmann[1], Françoise Vilaine[1], Françoise Gilard[3], Patrick A.W.
Klemens[4], Florence Guérard[3], Bertrand Gakière[3], H. Ekkehard Neuhaus[4], Catherine Bellini[1,5],
Sylvie Dinant[1] and Rozenn Le Hir[1,*]

[1] Institut Jean-Pierre Bourgin, INRAE, AgroParisTech, Université Paris-Saclay, 78000
Versailles, France

[2] Ecole Doctorale 567 Sciences du Végétal, Univ Paris-Sud, Univ Paris-Saclay, bat 360,
91405 Orsay Cedex, France

[3] Plateforme Métabolisme-Métabolome, Institute of Plant Sciences Paris-Saclay IPS2, CNRS,
INRAE, Univ Paris Sud, Univ Evry, Univ Paris-Diderot, Sorbonne Paris-Cité, Université
Paris-Saclay, Bâtiment 360, Rue de Noetzlin, 91192 Gif sur Yvette, France

[4] Universität Kaiserslautern, Pflanzenphysiologie, Postfach 3049, D-67653 Kaiserslautern,
Germany

[5] Umeå Plant Science Centre, Department of Plant Physiology, Umeå University, 90187
Umeå, Sweden

* Author for correspondence

One-sentence summary: A control of cytosolic sugar availability, mediated by SWEET16
and SWEET17, is required to sustain xylem development and secondary cell wall formation.

Author contributions

E.A. performed most of the research and made a first analysis of the data; P.K. performed the BIFC experiment under the supervision of H.E.N.; F.Gi., F.Gu and B.G. supervised and helped E.A. for the GC-MS analysis; F.V. cloned the complemented lines; B.H. provided technical assistance to E.A.; S.D., C.B. and R.LH reviewed and edited the manuscript; R. LH conceived the project, designed the research and wrote the article with contributions of all the authors.

Funding information:

This work has benefited from the support of IJPB's Plant Observatory technological platforms and from a French State grant (Saclay Plant Sciences, reference ANR-17-EUR-0007, EUR SPS-GSR) managed by the French National Research Agency under an Investments for the Future program (reference ANR-11-IDEX-0003-02) through PhD funding to E.A.

ABSTRACT

In higher plants, the development of the vascular system is controlled by a complex network of transcription factors. However, how nutrient availability in the vascular cells affects their development remains to be addressed. At the cellular level, cytosolic sugar availability is regulated mainly by sugar exchanges at the tonoplast through active and/or facilitated transport. In *Arabidopsis thaliana*, among the tonoplastic transporters, SWEET16 and SWEET17 have been previously localized in the vascular system. Here, using a reverse genetic approach, we propose that sugar exchanges at the tonoplast, mediated by SWEET16, are important for xylem cell division as revealed in particular by the decreased number of xylem cells in the *swt16* mutant and the expression of *SWEET16* at the procambium-xylem boundary. In addition, we demonstrate that transport of hexoses mediated by SWEET16 and/or SWEET17 is required to sustain the formation of the xylem secondary cell wall. This result is in line with a defect in the xylem cell wall composition as measured by FTIR in the *swt16swt17* double mutant and by upregulation of several genes involved in secondary cell wall synthesis. Our work therefore supports a model in which xylem development is partially dependent on the exchange of hexoses at the tonoplast of xylem-forming cells.

Keywords: Vacuole, tonoplast, sugar, transport, xylem, development, Arabidopsis, inflorescence stem

INTRODUCTION

The plant vasculature, composed of phloem, procambium/cambium, and xylem, is an elaborate system responsible for the transport of most biological compounds throughout the plant (Lucas et al., 2013). At the molecular level, vasculature development is governed by a complex network of transcription factors that are under the control of several signals, including hormones, peptides, and miRNAs (Fukuda and Ohashi-Ito, 2019; Smit et al., 2019). However, within this well-organized framework, a certain plasticity is required to adjust to cellular variations in terms of the availability of nutrients (i.e. sugars and amino acids).

Sugars, which represent the main source of energy, are required for metabolic activities, and they serve as a carbon reserve, and as intermediates for the synthesis of cell wall polysaccharides. Additionally, they have morphogenetic activity and act as primary messengers in signal transduction pathways (Sakr et al., 2018). It is therefore logical that modifications of sugar metabolism, transport or signaling can lead to multiple defects in plant growth and development (Eveland and Jackson, 2012). However, despite this central role, the role of sugar availability in the development of the vascular system in general and more specifically in heterotrophic tissues such as cambium and xylem is still elusive.

In these tissues, it has been suggested that lateral transport of sugars, coming from leakages from phloem sieve tubes, provides the sugars needed for vascular cell development (Minchin and McNaughton, 1987; Sibout et al., 2008; Spicer, 2014; Furze et al., 2018). Lateral transport is especially crucial for xylem secondary cell wall formation, since sugars are intermediate compounds in the synthesis of the cell wall polysaccharides which represent 80 % of the secondary cell wall (Marriott et al., 2016; Verbančič et al., 2018). The xylem tissue thus represents a strong sink for sugars that must be imported from surrounding tissues to serve as the source of carbon and energy. This is supported by the fact that perturbations in sugar transport at the plasma membrane of vascular cells, *via* SWEET or SUT transporters, affect the composition of the xylem secondary cell wall both in aspen and in Arabidopsis inflorescence stems (Mahboubi et al., 2013; Le Hir et al., 2015). Furthermore, in the Arabidopsis inflorescence stem, it has been suggested that movements of sucrose and/or hexoses towards the apoplast, mediated by SWEET11 and SWEET12, occur between the vascular parenchyma cells and the developing conducting cells to drive cell wall formation in a cell-specific manner (Dinant et al., 2019). Intercellular sugar availability seems, therefore, to play an important role in xylem development. However, the question remains open as to whether modification of sugar partitioning within the vasculature cells is also of importance.

The vacuole represents the main storage compartment for numerous primary and specialized metabolites including sugars (Martinoia, 2018). In tobacco leaves, up to 98% of hexoses are found in the vacuole (Heineke et al., 1994), whereas in Arabidopsis leaves, sucrose is mostly present in the cytosol and vacuoles contain half of the glucose and fructose (Weiszmann et al., 2018). Sugar exchanges between the vacuole and the cytosol are therefore required for dynamic adjustment of the quantity of sugar needed for metabolic and signaling pathways. In herbaceous and ligneous plants, few sugar transporters have been functionally characterized at the tonoplast (Hedrich et al., 2015), and localization in the cells of the vascular system has been shown only for *SUC4/SUT4*, *ESL1*, *SWEET16* and *SWEET17* (Yamada et al., 2010; Payyavula et al., 2011; Chardon et al., 2013; Klemens et al., 2013). In *Populus*, sugar export mediated by the tonoplastic sucrose symporter PttSUT4 is required for carbon partitioning between the source leaves and the lateral sinks (e.g. xylem). In Arabidopsis, we previously showed that the *SWEET16* promoter is active in the xylem parenchyma cells (Klemens et al., 2013), while the *SWEET17* promoter is active in the xylem parenchyma cells and young xylem cells of the Arabidopsis inflorescence stem (Chardon et al., 2013). Moreover high levels of *SWEET17* transcripts have been measured in the inflorescence stem, compared to other organs, after 7 to 8 weeks of growth (Guo et al., 2014). *SWEET16* and *SWEET17* are therefore good candidates with which to assess whether the maintenance of sugar homeostasis between the cytosol and the vacuole influences xylem development in Arabidopsis.

In the present work, through a reverse genetic approach, we demonstrate that SWEET16 and SWEET17 have specific and overlapping roles during xylem development. In particular, we suggest that sugar exchanges across the procambium-xylem boundary mediated by SWEET16 are important for xylem cell proliferation. By using infrared spectroscopy and gene expression analysis, we also show that both *SWEET16* and *SWEET17* are required for correct development of the secondary cell wall of xylem cells. Finally, since glucose and fructose accumulation are observed in the inflorescence stem of the double mutant, we suggest that maintenance of hexose homeostasis through the action of SWEET16 and/or SWEET17 is important at different stages of xylem development.

RESULTS

Radial growth of the inflorescence stem is altered in the *swt16swt17* double mutant

To explore to what extent mutations in *SWEET16* and/or *SWEET17* impact inflorescence stem development in Arabidopsis, we used the previously described *sweet17-1* (hereafter called *swt17*) mutant line (Chardon et al., 2013) and identified two new T-DNA insertion lines in the *SWEET16* gene. The mutants *sweet16-3* (SM_3.1827) and *sweet16-4* (SM_3.30075) possess an insertion in the promoter region and in the first exon of the *SWEET16* gene respectively (Supplemental Figure 1A). They were named after the previous alleles already published (Guo et al., 2014). A full-length *SWEET16* transcript was detected by RT-PCR in the *sweet16-3* allele, while no full-length transcript could be detected in the *sweet16-4* allele (Supplemental Figure 1B and 1C). The *sweet16-4* allele (hereafter called *swt16*) was therefore deemed to be a null allele. We generated the double mutant *sweet16sweet17* (hereafter called *swt16swt17*) and confirmed by RT-PCR that both genes full length were absent in this double mutant (Supplemental Figure 1C).

Analysis of the inflorescence stems of the *swt16*, *swt17* and *swt16swt17* mutants showed that the area of stem cross-sections was significantly smaller compared to that of the wild-type (Figure 1A and B). More precisely, the stem of all mutant lines contained less xylem tissue compared with the wild type, while only the stem of *swt16swt17* displayed significantly less phloem tissue (Figure 1C-D). Additionally, the proportion of xylem or phloem per stem was calculated (Figure 1E-F). While no change in the proportion of phloem was observed in the mutants compared to the wild type (Figure 1E), a significant reduction in the xylem proportion was observed in the double *swt16swt17* mutant (Figure 1F). Both *SWEET16* and *SWEET17* are therefore required for proper xylem development and hence for proper radial growth of the stem.

SWEET16 but not SWEET17 is required for proliferation of xylem cells

The phenotype of the xylem tissue was further analyzed on an independent set of plants (Figure 2). We first checked the robustness of the inflorescence stem phenotype and confirmed that the *swt16* and *swt17* single mutants and the *swt16swt17* double mutant consistently displayed a significantly thinner inflorescence stem compared to the wild type (Figure 2A). In addition, we confirmed our previous results that showed a significantly shorter

inflorescence stem in the *swt17* mutant compared to the wild-type (Figure 2B) (Chardon et al., 2013). Interestingly, we did not observed any alteration in the inflorescence stem height in *swt16* or *swt16swt17* compared to the wild type (Figure 2B).

The xylem phenotype was then studied in more detail by counting the number of xylary fibers (cells with an area of between 5-150 µm²) and xylem vessels (cells with an area greater than 150 µm²) as well as measuring the individual cross-sectional areas within each vascular bundle (Figure 2C-I). In the vascular bundles of the *swt16* and *swt16swt17* mutants, but not *swt17*, the area occupied by the xylem tissue was significantly smaller than in the wild type (Figure 2C). These changes could result from modification of either cell size or cell number. While no changes in the size of the xylary fibers or the xylem vessels were observed in any of the genotypes analyzed (Figure 2D-E), the total number of xylem cells per vascular bundle was significantly reduced, by about 20%, in the single mutant *swt16* and the double mutant *swt16swt17* but not in *swt17* (Figure 2F). The numbers of xylary fibers and xylem vessels per vascular bundle were significantly reduced in the stem of the *swt16* single and *swt16swt17* double mutant but not in the *swt17* mutant (Figure 2G-H). The decreased number of xylary fibers was proportional to that of xylem vessels since the vessels-to-fibers ratio was the same in the wild type and the *swt16*, *swt17* and *swt16swt17* mutant lines (Figure 2I).

Overall, these results show that the single *swt16* mutant and the *swt16swt17* double mutant have the same phenotype (Figure 2 and Supplemental Table 1) and suggest that the expression of *SWEET16*, but not that of *SWEET17*, is required for correct division of xylem cells.

SWEET16 and SWEET17 are required for normal secondary cell wall composition and development in xylem cells

To explore whether modifications in the vacuolar transport of sugars impact the formation of the xylem cell wall, we first exploited the transcriptomic dataset obtained from plants overexpressing a dexamethasone (DEX)-inducible version of the *VASCULAR RELATED NAC-DOMAIN PROTEIN 7* (*VND7*) gene, the master secondary wall-inducing transcription factor (Li et al., 2016). These plants allow the transcriptional and metabolic changes occurring during secondary cell wall formation to be studied. From the RNA-seq dataset, we extracted information related to the expression of the family of *SWEET* genes at

different time points after induction of *VND7* expression (Supplemental Figure 2). Out of the 17 *SWEET* genes identified in Arabidopsis, 7 were differentially expressed during secondary cell wall formation (Supplementary Figure 2). Most interestingly, the vacuolar *SWEET2* and *SWEET17* were significantly upregulated 3 hours after DEX induction while *SWEET16* expression was upregulated 12 hours after DEX induction (Supplementary Figure 2). In contrast, the expression of genes encoding the plasma membrane localized SWEET transporters (e.g. *SWEET1*, *SWEET3*, *SWEET11* and *SWEET12*) was significantly downregulated during secondary cell wall formation (Supplementary Figure 2). Additional analysis of the dataset showed that *SWEET2* and *SWEET17* were co-regulated with genes related to cell wall synthesis (*CESA*, *SND2*, *SDN3*, *MYB46*) as well as those encoding other sugar transporters localized at the tonoplast (*ESL1*) or at the plasma membrane (*STP1*, *STP13* and *PMT4*) (Supplementary Table 2). These results support the fact that sugar export from the vacuole to the cytosol is ongoing during secondary cell wall formation, most probably to provide sugars to be used as intermediates for cell wall formation.

To assess whether *SWEET16* and *SWEET17* are indeed functionally involved in xylem secondary cell wall formation, we performed a targeted gene expression analysis including genes that are known to be part of the transcriptional network involved in stem cell proliferation/organization (*PXY*, *WOX4*) (Etchells et al., 2013), xylem cell identity (*ATHB8*) (Smetana et al., 2019), and secondary cell wall biosynthesis in vessels/fibers (*CESA4*, *CESA7*, *CESA8*, *KNAT7, MYB4, MYB43, MYB46, MYB52*, *MYB54*, *MYB58*, *MYB63*, *MYB83*, *MYB103*, *NST1*, *VND6*, *VND7*, *SND1*/*NST3*, *SND3, VNI2* and *XND1*) (Hussey et al., 2013) (Figure 3A-I). When looking at the overall transcriptional profile of the wild type and the *swt16*, *swt17* and *swt16swt17* mutants, two clusters can be identified (Figure 3A). The first cluster contains the wild type, the *swt16* and *swt17* single mutants, whereas the second includes only the *swt16swt17* double mutant. Only a subset of genes shows significantly increased expression among the different genotypes, namely *SND3*, *MYB103*, *MYB4*, *VNI2*, *SND1*, *MYB83*, *MYB54* and *MYB46* (Figure 3A), though a tendency, albeit not significant, is observed for *MYB43* (P=0.053) and *KNAT7* (P=0.091) (Figure 3A). Interestingly, all these genes belong to the transcriptional network involved in secondary cell wall biosynthesis in xylem vessels and/or in xylary fibers (for review Hussey et al., 2013). A Student's *t*-test was then performed to compare each mutant line to the wild-type plants (Figure 3B-I). On average, a 2-fold increase in expression was measured for the genes *SND1*, *MYB46*, *VNI2*, *MYB83* and *MYB54* in the *swt16swt17* double mutant compared to the wild type (Figure 3B,

C, D, F and G), while a similar tendency was observed for *MYB4*, *SND3* and *MYB103*

expression (Figure 3E, H and I). Overall, these results show that in the *swt16swt17* double

mutant neither stem cell maintenance nor xylem identity genes are affected, whereas

secondary cell wall biosynthesis genes are deregulated.

Next, we tested whether this transcriptional deregulation was accompanied by

modifications in the cell wall composition. We used Fourier-transformed infrared

spectroscopy (FTIR) on inflorescence stem cross-sections to analyze the xylem secondary cell

wall composition as previously described in Le Hir et al. (2015) (Figure 4). The average

spectra for all three mutants showed several differences compared to the wild-type spectra in

fingerprint regions associated with cellulose, hemicelluloses and lignin (Figure 4A). The *t*-

values, plotted against each wavenumber of the spectrum, showed that the mutant lines

exhibited several significant negative and positive peaks (higher or lower absorbance than in

the wild type) at wavenumbers associated with cellulosic and hemicellulosic polysaccharides

(898 cm^{-1}, 995-1120 cm^{-1}, 1187 cm^{-1}, 1295 cm^{-1}, 1373 cm^{-1}, 1401 cm^{-1}, 1423 cm^{-1}, 1430 cm^{-1},

1440 cm^{-1}, and 1485 cm^{-1}) (Åkerholm and Salmén, 2001; Kačuráková et al., 2002; Lahlali et

al., 2015) (Figure 4A and B). More precisely, wavenumbers at 898 cm^{-1}, associated with the

amorphous region of cellulose (Kačuráková et al., 2002), and at 1430 cm^{-1}, associated with

crystalline cellulose (Åkerholm and Salmén, 2001), showed opposite and significant

differences (Figure 4B). This suggests a potential defect in cellulose organization in the xylem

secondary cell wall. Measurements of the cellulose C-O vibrations at a peak height at 1050

cm^{-1} (Lahlali et al., 2015) further indicate modifications of the cellulose composition in the

cell wall of all the mutant lines (Figure 4C).

The *swt16* and *swt16swt17* mutant also displayed significant variations compared to

the wild type at 1740 cm^{-1} (band specific for acetylated xylan; Gou et al. 2003) and 1369 cm^{-1}

(deformation of C-H linkages in the methyl group *O*-acetyl moieties; Mohebby, 2008)

suggesting modifications in xylan acetylation (Figure 4A and B). Furthermore, the

hemicellulose peak height at 1740 cm^{-1} (C-O and C-C bond stretching) was significantly

smaller in the *swt16* mutant suggesting less acetylated xylan (Figure 3D). Although the *swt17*

single mutant was not distinguishable from the wild type, the *swt16swt17* double mutant had

significantly fewer acetylated xylans than the wild type and the *swt16* single mutant (Figure

4D). The lignin-associated bands at 1510 cm^{-1} (Faix, 1991), 1520 cm^{-1} (Faix, 1991; Gou et al.,

2008) and 1595 cm^{-1} also exhibited significant differences in the single and double mutants

compared to the wild type plants (Figure 4B). Measurements of the lignin peak height ratio

(1510/1595 cm^{-1}) showed that the secondary cell wall of the *swt16* single mutant contains

significantly more G-type lignin than that of the wild type, while only a tendency was

measured in the cases of the *swt17* and the *swt16swt17* mutants (Figure 4E).

Overall, these results suggest that sugar export towards the cytosol mediated by

SWEET16 and/or SWEET17 is required to provide the intermediates needed for the synthesis

of cellulosic and hemicellulosic polysaccharides.

The hexose content is modified in the inflorescence stem of the *swt16swt17* double

mutant

Assuming that SWEET16 and SWEET17 are sugar carriers, we wondered what would

be the metabolic status of the inflorescence stem in the *swt16swt17* double mutant. We

therefore used GC-MS to explore the global metabolomic profiles of the wild-type and the

double *swt16swt17* mutant, identifying a total of 158 metabolites. In order to identify the

subset of metabolites that best discriminate between the genotypes, we performed a sPLS-DA

(Figure 5A). The resulting score plot clearly allows the two genotypes to be separated by the

first dimension (Figure 5A). Among the metabolites selected by the sPLS-DA analysis, a

subsequent *t*-test identified nine that were significantly different between wild type and the

swt16swt17 double mutant: allothreonine, benzoic acid, citraconic acid, cysteinylglycine,

fructose, fumaric acid, glucose-6-phosphate, phytol and valine (Figure 5B and Supplemental

Table 3). The relative quantities of benzoic acid, citraconic acid and fumaric acid (a

component of the tricarboxylic cycle) were significantly reduced in the double mutant

compared to the wild type (Supplemental Figure 3A, B and F). On the other hand, significant

accumulation of cysteinylglycine (an intermediate in glutathione biosynthesis;(Hasanuzzaman

et al., 2017), hexoses and hexose-phosphates (e.g. glucose-6-phosphate and fructose), amino

acids (e.g. allothreonine and valine) and phytol (a chlorophyll component; (Gutbrod et al.,

2019) was measured in the *swt16swt17* mutant compared to the wild-type stems

(Supplemental Figure 3C, D, E, G, H and I). We further quantified the soluble sugars and

starch content in both genotypes by enzymatic methods. Consistent with the metabolomics

results, a significant accumulation of fructose in the double mutant was confirmed (Figure

5C). In addition, the glucose content was significantly increased in the stem of the double

mutant (Figure 5C), while no variation in the sucrose and starch contents was observed

(Figure 5C). Interestingly, the inflorescence stem of the *swt16swt17* double mutant

accumulated mostly hexoses while no significant changes in glucose or sucrose were observed in the stem of the *swt16* and *swt17* single mutants (Supplemental Figure 4A). Although it was not significant, a tendency to accumulate glucose was observed in the single mutants (Supplemental Figure 4B). A significant increase in fructose content was measured only in the *swt17* mutant compared to the wild type (Supplemental Figure 4C).

SWEET16 and SWEET17 proteins interact physically and are expressed in the xylem during its early development

To localize SWEET16 and SWEET17 in the inflorescence stem, we generated translational fusions between GFP and the N-terminus of *SWEET16* or *SWEET17* coding sequences under the control of their native promoters. The translational fusions were transferred into the respective knockout mutants in order to check their functionalities. We then looked at the radial growth of the inflorescence stem in the transgenic plants. *pSWT17:GFP-SWT17* successfully complemented the phenotype of the *swt17* mutant (Supplemental Figure 5), while *pSWT16:GFP-SWT16* only partially complemented the stem phenotype of the *swt16* mutant (Supplementary Figure 5). However, full complementation of the double mutant *swt16swt17* was achieved when both translational GFP fusions were expressed (Supplemental Figure 5). Unfortunately, despite the phenotype complementation, we could not detect any GFP signal in these lines. Previously, using lines expressing *pSWT:GUS* translational fusions, we showed that SWEET16 and SWEET17 were expressed in the xylem tissue of petioles and inflorescence stems (Chardon et al., 2013; Klemens et al., 2013). Here we completed this analysis by studying the expression patterns of SWEET16 and SWEET17 in cross sections from three different zones of the stem: (1) a stem region where the growth was rapid, (2) a stem region where elongation growth had finished but where further thickening of the secondary cell wall was still ongoing and (3) the base of the stem, which corresponds to a mature zone (Hall and Ellis, 2013) (Figure 6). Interestingly, the expression pattern of *SWEET16* and *SWEET17* varied depending on the developmental stages. The *SWEET16* expression pattern fluctuated more depending on the stage of inflorescence stem development (Figure 6A-C). In the region where the stem was growing rapidly (Figure 6A and inset) and in the region where secondary cell wall thickening was still ongoing (Figure 6B and inset), expression of *SWEET16* was observed in the interfascicular fibers, across the procambium-xylem boundary and in the young xylem vessels (Figure 6A-B and insets). Faint

GUS staining was also observed in the phloem cells at these stages (Figure 6A-B). In the
mature stem, *SWEET16* expression was restricted to the xylary parenchyma cells (Figure 6C).
In contrast, *SWEET17* expression was restricted mostly to the xylem parenchyma cells present
in the distal part of the vascular bundle (Figure 6D-F). Expression of *SWEET17* was also
observed in the young xylem cells before extensive cell wall lignification had occurred, as
shown by the weaker phloroglucinol cell wall staining (Figure 6E, 6F and inset). Here, it is
worth noting that we use the general term "young xylem cells" because it is not possible to
know whether the GUS signal is present in developing xylary fibers and/or in developing
xylem vessels. In conclusion, *SWEET16* and *SWEET17* expression patterns overlap in the
young xylem cells as well as in the xylary parenchyma cells.

It has been established that plant sugar SWEET transporters require homo or hetero-
oligomerization to gain functionality (Xuan et al., 2013). Because *SWEET16* and *SWEET17*
expression patterns overlap in young xylem cells and xylary parenchyma cells, we
investigated whether these proteins could interact. Xuan et al. (2013) previously showed that
SWEET16 and SWEET17 can form homodimers as well as heterodimers in a split ubiquitin
yeast two-hybrid assay. We confirmed that SWEET16 and SWEET17 can form a heterodimer
at the vacuolar membrane in a bimolecular fluorescence complementation assay in
Arabidopsis mesophyll protoplasts (Figure 6G-I).

DISCUSSION

To efficiently control carbon homeostasis within a cell and to fuel the different
metabolic and signaling pathways, dynamic sugar storage in the plant vacuole is critical. Over
the past years, several vacuolar transporters have been identified at the molecular level (for
review see Hedrich et al., 2015). Among them, SWEET16 and SWEET17 have been
characterized as bidirectional tonoplastic sugar facilitators and shown to be involved in seed
germination, root growth and stress tolerance (Chardon et al., 2013; Klemens et al., 2013;
Guo et al., 2014). In addition, the expression of both genes has been shown in the
inflorescence stem's vascular parenchyma cells, but this had not previously been explored
further. In this work, we ask whether facilitated sugar transport (*via* SWEET16 and
SWEET17) across the vacuolar membrane limits vascular tissue development in the
inflorescence stem of Arabidopsis.

First, our data highlight modifications of hexose homeostasis in the inflorescence stem of the different mutant lines. Although a tendency to accumulate glucose was observed in the *swt16* mutant, a significant increase in fructose was measured in the *swt17* mutant stem. Furthermore, mutations in both *SWEET16* and *SWEET17* induced somewhat specific accumulation of glucose, glucose-6-phosphate and fructose in the inflorescence stem. It has been previously shown that defects in the expression of vacuolar sugar transporters alter carbon partitioning and allocation in different organs, which is in line with our findings for the inflorescence stem (Wingenter et al., 2010; Yamada et al., 2010; Poschet et al., 2011; Chardon et al., 2013; Klemens et al., 2013; Guo et al., 2014; Klemens et al., 2014). Knowing that SWEET proteins are sugar facilitators working along the concentration gradient (Chen et al., 2010) and that at least half of the hexoses are present in the plant vacuole (Heineke et al., 1994; Weiszmann et al., 2018), we can reasonably propose that some of the hexoses are trapped inside the vacuole in the single *swt16* and *swt17* mutants and the *swt16swt17* double mutant. As a consequence, modifications in the distribution of hexose concentrations between the vacuole and the cytosol, which would impact the availability of hexoses for subsequent metabolic and signaling purposes, could be expected. Hexoses are known to favor cell division and expansion, while sucrose favors differentiation and maturation (Koch, 2004). In addition, after metabolization, hexoses and hexoses-phosphates constitute the building blocks for the synthesis of cell wall polysaccharides (Verbančič et al., 2018). Since *SWEET16* and/or *SWEET17* are expressed in the xylem initials, in young xylem cells and in xylem parenchyma cells, we propose that enhanced storage of vacuolar hexoses in these cells will affect different stages of xylem development.

(Pro)cambium and xylem tissues can be regarded as sinks because they rely mostly on the supply of carbohydrates from the surrounding cells to sustain their development (Sibout et al., 2008; Spicer, 2014). In aspen stem, a gradual increase in sucrose and reducing sugars, together with a rise in the activities of sugar metabolism enzymes, are observed across the cambium-xylem tissues (Roach et al., 2017). In addition, in tomato, modification of fructose phosphorylation by the fructokinase SlFRK2 leads to a defect in cambium activity (Damari-Weissler et al., 2009). Taken together, these results support the need for maintenance of sugar homeostasis in the (pro)cambium to respond to the high metabolic activity required during cell division. Our work identified SWEET16 as a player in the dividing xylem cells, acting to balance the tradeoffs between the need for sugars in the cytosol and their storage in the vacuole (Figure 7). This conclusion is supported by the fact that *SWEET16* is expressed across

the procambium-xylem boundary and that a mutation in *SWEET16* leads to defects in the number of xylem cells and in radial growth of the inflorescence stem. Furthermore, the expression of the gene coding for the WUSCHEL RELATED HOMEOBOX 4 (WOX4) transcription factor (Etchells et al., 2013), which is involved in cellular proliferation, was unchanged in both *swt16* and *sw16swt17* mutants. This suggest that the defects in xylem cell division result mostly from reduced availability of energy and matter resources due to a reduction in sugar transport, rather than from the signaling role of sugars. Finally, the partial complementation of the *swt16* mutant could also suggest that other vacuolar sugar transporters, yet to be identified, might be at play in this process.

The reduced number of xylem vessels in the *swt16swt17* double mutant's vascular bundle could be explained by the upregulation of *VND-INTERACTING 2* (*VNI2*), which is a repressor of the activity of the master regulator of xylem vessel differentiation VND7 (Zhong et al., 2008; Yamaguchi et al., 2010). On the other hand, the overexpression in the double mutant *swt16swt17* of SECONDARY WALL-ASSOCIATED NAC DOMAIN 1 (SND1), the master switch for fiber differentiation, would be expected to result in a shift towards increased differentiation of xylary fibers (Zhong et al., 2006), which is not consistent with the fewer fibers observed in the double mutant. Based on these results, we can assume that the increase in storage of vacuolar hexoses in the double mutant also affects xylem cell differentiation. Against this hypothesis, our results show that both xylary fibers and xylem vessels number decreased proportionately, since no change in the xylem vessels/xylary fibers ratio was measured. Consistent with this observation, the expression of the gene coding for the PHLOEM INTERCALATED WITH XYLEM (PXY) receptor, which is involved in xylem cell differentiation (Etchells et al., 2016), was not modified. Despite the upregulation of *VNI2* and *SND1* expression, which could be due to a feedback mechanism yet to be identified, these results therefore tend to suggest that no disequilibrium is occurring in xylem cell differentiation in the *swt16swt17* mutant stem. The enhanced storage of hexoses in the vacuole of the double mutant is therefore affecting the overall pattern of xylem cell division rather than xylem cell differentiation.

After cell division and differentiation, xylem cells undergo a developmental program that includes secondary cell wall formation, lignification and programmed cell death, to produce functional xylem fibers and vessels (Schuetz et al., 2012) (Figure 7). Along with the overexpression of *SDN1*, the overexpression of genes encoding its downstream targets,

namely MYB DOMAIN PROTEIN 46 and 83 (MYB46 and MYB83), was observed in the *swt16swt17* double mutant. Furthermore, the targets of the MYB46/MYB83 node, which positively regulates SND3 (SECONDARY WALL-ASSOCIATED NAC DOMAIN PROTEIN 3), KNAT7 (KNOTTED-LIKE HOMEOBOX OF ARABIDOPSIS THALIANA 7), MYB43 and MYB54 or/and negatively regulates MYB103, KNAT7 and MYB4, all of which are involved in the formation of the xylem secondary cell wall, are also upregulated (Hussey et al., 2013). KNAT7 directly or indirectly represses cellulose, hemicellulose and lignin biosynthetic genes (Li et al., 2012), while MYB54 is related to cellulose synthesis (Zheng et al., 2019). In Arabidopsis, MYB43 along with other MYB transcription factors regulates lignin biosynthesis (Geng et al., 2020), while its ortholog in rice is involved in the regulation of cellulose deposition (Ambavaram et al., 2011). Finally, upregulation of MYB4 in Arabidopsis results in downregulation of the lignin pathway (Jin et al., 2000), while a role for MYB103 in lignin biosynthesis has been shown in Arabidopsis stem (Öhman et al., 2013). In the *swt16wt17* double mutant, these transcriptional changes were accompanied by modifications of the secondary cell wall in terms of cellulose and hemicellulose composition. However, a single mutation in *SWEET16* or *SWEET17* was sufficient to modify the composition of the xylem cell wall without any alteration in the expression of genes involved in secondary cell wall synthesis. Our data further show that the *SWEET16* and *SWEET17* expression patterns overlap in xylem cells that are building a secondary cell wall and that they form a heterodimer in Arabidopsis mesophyll protoplasts. We therefore postulate that the intermediate sugars required for the synthesis of cell wall polysaccharides come in part from the vacuole unloading mediated by SWEET16 and SWEET17 homo- and heterodimers (Figure 7). Previously, it has been shown that genes encoding vacuolar sugar facilitators are up-regulated during secondary cell wall formation in xylem vessels, while sugar facilitators expressed at the plasma membrane are down-regulated (Supplementary Figure 2) (Li et al., 2016), supporting the idea that secondary cell wall formation relies on sugar export from the vacuole. In the current model of cell wall synthesis, the cytosolic catabolism of sucrose is thought to be the main source of nucleotide sugars (e.g. UDP-glucose, UDP-galactose, GDP-mannose) that act as precursors for cellulose and hemicellulose synthesis (Verbančič et al., 2018). Our data support the existence of a more complex system in which the export of vacuolar hexoses also represents a source for the synthesis of nucleotide sugars and subsequent cell wall formation (Figure 7).

Because *SWEET16* and *SWEET17* are also expressed in the xylary parenchyma cells, we postulate that the maintenance of sugar homeostasis within this cell type, is important and could contribute to the provision of carbon skeletons for secondary cell wall synthesis after the disappearance of the vacuole from the maturing xylem cells (Figure 7). Within this scheme, the export of sugars in the apoplastic space between parenchyma cells and developing conducting cells could be carried out by the plasmalemmal SWEET11 and SWEET12 transporters which also expressed in the xylary parenchyma cells (Figure 7 and (Le Hir et al., 2015). Such cooperation between parenchyma cells and developing conducting cells was previously described as the "good neighbor" hypothesis in the context of H_2O_2 and monolignol transport (Barcelo, 2005; Smith et al., 2013; Smith et al., 2017). To further explore the importance of sugar transport between xylary parenchyma cells and developing xylem cells, more experiments, such as cell-specific complementation of the *sweet* mutant lines, will be needed in order to better comprehend the role of xylary parenchyma cells in xylem development.

In conclusion, our work shows that exchange of intracellular hexoses, mediated by SWEET16 and/or SWEET17 at the tonoplast, contributes to xylem development by regulating the amounts of sugars that will be made available to the different cellular processes. However, how the cell is prioritizing the distribution of sugars among the different processes remains an open question. Although these technologies are challenging, the use of non-aqueous fractionation and metabolomics approaches (Fürtauer et al., 2019) could help in resolving subcellular sugar metabolism in a cell-specific context in mutant lines affected in sugar metabolism, transport and signaling.

MATERIALS AND METHODS

Plant material and growth conditions

Seeds of T-DNA insertion lines homozygous for *SWEET17* (*sweet17-1*) and *SWEET16* (*sweet16-3* and *sweet16-4*) were gifts from Dr. F. Chardon and Pr. E. Neuhaus respectively. The *sweet17-1* line was previously reported to be a knock-out by Chardon et al. (2013). The *sweet16-3* (SM_3_1827) and *sweet16-4* (SM_3_30075) lines were numbered following the *sweet16-1* and *sweet16-2* lines already published by Guo et al. (2014). To verify whether *sweet16-3* and *sweet16-4* were knock-out mutants we performed RT-PCR with specific primers to amplify the full-length *SWEET16* cDNA (Supplemental Figure 1 and Supplemental Table 4). Since only the *sweet16-4* mutant turned to be a knock-out (Supplemental Figure 1B), we crossed it with *swt17-1* to obtained the double mutant *sweet16-4sweet17-1* (hereafter referred as *swt16swt17*). Homozygous plants were genotyped using gene-specific primers in combination with a specific primer for the left border of the T-DNA insertion (Supplemental Table 4).

To synchronize germination, seeds were stratified at 4°C for 48 hours and sown in soil in a growth chamber in long day conditions (16 hours day/8 hours night and 150 μE m^{-2} s^{-1}) at 22/18°C (day/night temperature) with 35% relative humidity. Plants were watered with Plant-Prod nutrient solution twice a week (Fertil, https://www.fertil.fr/). For all experiments, the main inflorescence stems (after removal of lateral inflorescence stems, flowers and siliques) were harvested from seven-week old plants.

Inflorescence stem sample preparation

For each plant, the main inflorescence stem height was measured with a ruler before harvesting a 1 to 2 cm segment taken at the bottom part of the stem. The stem segments were embedded in 8% agarose solution and sectioned with a VT100 S vibratome (Leica, https://www.leica-microsystems.com/). Some of the cross-sections were used for FT-IR analysis and the others were stained with a FASGA staining solution prepared as described in Tolivia and Tolivia (1987) for morphometric analysis of the xylem.

Morphometric analysis of the xylem

Previously stained inflorescence stem cross-sections were imaged under an Axio Zoom V16
microscope equipped with a Plan-Neofluar Z 2.3/0.57 FWD 10.6 objective (Zeiss,
https://www.zeiss.fr/microscopie/). For each section, the diameter of the inflorescence stem
was measured using the Image J software package (https://imagej.nih.gov/ij/). For the same
sections all the vascular bundles were photographed individually using a confocal laser
scanning microscope and morphological analysis of the xylem were performed as described in
Le Hir et al. (2015). For each vascular bundle, the morphological segmentation made it
possible to find the number of xylem cells (xylary fibers and xylem vessels) as well as their
cross-sectional areas. Cells with a cross-sectional area of between 5 to 150 µm² were
considered to be xylary fibers and cells with a cross-sectional area greater than 150 µm² were
considered to be xylem vessels. The sum of all xylem cell cross-sectional areas was then
calculated to give the total xylem cross-sectional area. The average xylary fiber and xylem
vessel area was calculated by dividing the total xylem cross-sectional area by the number of
each cell type.

FT-IR analysis of the xylem secondary cell wall

The composition of the secondary cell wall of the xylem tissue was determined by Fourier
Transformed Infra-red spectroscopy using an FT-IR Nicolet^{TM} iN^{TM} (Thermo Fisher
Scientific, https://www.thermofisher.com). Spectral acquisitions were done in transmission
mode on a 30 µm by 30 µm acquisition area targeting the xylem tissue (xylem vessels and
xylary fibers) as described in Le Hir et al. (2015). Between 10 to 15 acquisition points
sweeping the xylem tissue homogeneously were performed on each vascular bundle within a
stem section. Three individual inflorescence stems were analyzed for each genotype. After
sorting the spectra and correcting the baseline, the spectra were area-normalized and the
different genotypes were compared as described in Le Hir et al. (2015). The absorbance
values (maximum height) of the major cellulose, lignin and hemicellulose bands in the
fingerprint region (1800–800 cm^{-1}) were collected using TQ Analyst EZ edition (Thermo
Fisher Scientific, https://www.thermofisher.com).

Metabolomic analysis

The inflorescence stems of the wild-type and the *swt16swt17* double mutant were harvested in the middle of the day (8 hours after the beginning of the light period). Metabolites were extracted from 4.5 mg of lyophilized stem powder from eight individual plants and analyzed by GC-MS as described in Cañas et al. (2020). Relative concentrations of metabolites were determined relative to the internal standard ribitol, which was added after grinding the lyophilized material. Differential accumulation of metabolites was determined by one-way analysis of variance (ANOVA) and *post hoc* Tukey tests ($P<0.05$).

Quantification of soluble sugars and starch

The main inflorescence stems of the wild type, and the *swt16*, *swt17* and *swt16swt17* mutants, were harvested in the middle of the day (8 hours after the beginning of the light period), frozen in liquid nitrogen and ground with a mortar and a pestle. Soluble sugars and starch were extracted from 50 mg of powder from an individual stem as described in Sellami et al. (2019). Depending on the experiment, 4 to 9 biological replicates were analyzed.

RNA isolation and cDNA synthesis

RNAs were prepared from the main inflorescence stem from four 7-week-old individual plants grown as described above. Samples were frozen in liquid nitrogen before being ground with a mortar and a pestle. Powders were stored at -80°C until use. Total RNA was extracted from frozen tissue using TRIzol® reagent (Thermo Fisher Scientific, 15595-026, https://www.thermofisher.com) and treated with DNase I, RNase-free (Thermo Fisher Scientific, EN0521, https://www.thermofisher.com). cDNA was synthetized by reverse transcribing 1 µg of total RNA using RevertAid H minus reverse transcriptase (Thermo Fisher Scientific, EP0452, https://www.thermofisher.com) with 1 µl of oligo(dT)18 primer (100 pmoles) according to the manufacturer's instructions. The reaction was stopped by incubation at 70 °C for 10 min.

Quantitative qPCR experiment

Transcript levels were assessed for four independent biological replicates in assays with

triplicate reaction mixtures by using specific primers either designed with the Primer3

software (http://bioinfo.ut.ee/primer3-0.4.0/primer3/) or taken from the literature

(Supplemental Table 5). qPCR reactions were performed in a 96-well transparent plate on a

Bio-Rad CFX96 Real-Time PCR machine (Bio-Rad) in 10 µl mixtures each containing 5 µl

of Takyon™ ROX SYBR$^{®}$ MasterMix dTTP Blue (Eurogentec, UF-RSMT-B0710,

https://www.eurogentec.com/), 0.3 µl forward and reverse primer (30 µM each), 2.2 µl sterile

water and 2.5 µl of a 1/30 dilution of cDNA. The following qPCR program was applied:

initial denaturation at 95°C for 5 min, followed by thirty-nine cycles of 95°C for 10 sec, 60°C

for 20 sec, 72°C for 30 sec. Melting curves were derived after each amplification by

increasing the temperature in 0.5°C increments from 65°C to 95°C. The Cq values for each

sample were acquired using the Bio-Rad CFX Manager 3.0 software package. The specificity

of amplification was assessed for each gene, using dissociation curve analysis, by the

precision of a unique dissociation peak. If one of the Cq values differed from the other two

replicates by > 0.5, it was removed from the analysis. The amplification efficiencies of each

primer pair were calculated from a 10-fold serial dilution series of cDNA (Supplemental

Table 5). Four genes were tested as potential reference genes: *APT1* (At1g27450), *TIP41*

(At4g34270), *EF1α* (At5g60390) and *UBQ5* (At3g62250). The geNorm algorithm

(Vandesompele et al., 2002) was used to determine the gene most stably expressed among the

different genotypes analyzed, namely *UBQ5* in this study. The relative expression level for

each genotype was calculated according to the ΔCt method using the following formula:

average $E_t^{-Cq(of\ target\ gene\ in\ A)}/E_r^{-Cq(of\ reference\ gene\ in\ A)}$, where E_t is the amplification efficiency of

the target gene primers, E_r is the reference gene primer efficiency, A represents one of the

genotypes analyzed.

Tagged protein constructs

For complementation of the single *sweet16-4* and *sweet17-1* and the double *sweet16-*

4sweet17-1 mutants, N terminal fusions with GFP were constructed as follow. First, the

coding sequence of eGFP was amplified from the pKGWFS7 plasmid (Karimi et al., 2002)

without a stop codon and then introduced into a modified donor pENTR vector to produce

pENT-GFP (w/o stop). To make the N terminal translational GFP fusions (pSWEET16:GFP-

SWEET16 and pSWEET17:GFP-SWEET17), the promoters (1295 bp for *SWEET16* and 2004

bp for *SWEET17*) and genomic sequences (1863 bp for *SWEET16* and 2601 bp for *SWEET17*)

were amplified separately and then cloned on each side of the GFP gene in the intermediary

vector pENT-GFP (w/o stop) by taking advantage of the restriction sites generated by PCR.

All the PCR reactions were performed using Phusion High-Fidelity DNA Polymerase

(Thermo Fisher Scientific, F-530S, https://www.thermofisher.com) with the primers listed in

Supplemental Table 4. Donor vectors created in this way were analyzed by sequencing in

order to check the reading frame of the translational fusions and the integrity of the whole

genomic sequences. Destination binary vectors were then obtained by recombination, using

Gateway® LR Clonase II Enzyme Mix (Thermo Fisher Scientific, 11791-100,

https://www.thermofisher.com), between pENTR donor vectors and pMDC123 for N terminal

fusions (Curtis and Grossniklaus, 2003). All binary vectors were introduced into

Agrobacterium tumefaciens C58pMP90 (Koncz and Schell, 1986) by electroporation.

Arabidopsis single mutants *swt16-4* and *swt17-1* as well as the double mutant *sweet16-*

4sweet17-1 plants were transformed by the floral dip method (Clough and Bent, 1998).

Transformants were selected on hygromycin (15 mg/L) for pMDC99 constructs and/or Basta

(7.5 mg/L) for pMDC123 constructs. For all constructs, three independent transgenic lines

were analyzed and one representative line was selected for subsequent studies.

For the bimolecular fluorescence complementation (BiFC) assay, the full-length ORFs of

SWEET16 and *SWEET17* were amplified from cDNA with the primers given in Supplemental

Table 4, either with or without their stop codons, depending on the final vector used. The

ORFs were further sub-cloned into pBlueScript II SK, blunt end cut with EcoRV. The

resulting vectors were checked for errors and orientation of the insert by sequencing with T3

and T7 primers. Subsequently, positive clones in the T7 orientation and the corresponding

pSAT1 vectors (Lee et al., 2008) were cut with EcoRI and XhoI. *SWEET16* including the stop

codon was ligated into pSAT1-cCFP-C, and *SWEET17* without the stop codon into pSAT1-

nCerulean-N. Plasmid DNA of the final constructs was isolated with a PureLink™ HiPure

Plasmid Filter Midiprep Kit (Invitrogen™ /Thermo Fisher Scientific) according to the

manufacturer's manual. Isolation and transfection of Arabidopsis mesophyll protoplasts were

performed as described by Yoo et al. (2007). For imaging protoplasts, a Leica TCS SP5 II

confocal laser scanning microscope (http://www.leica-microsystems.com) was used. All

pictures were taken using a Leica HCX PL APO 63·/1.20 w motCORR CS objective with a

VIS☐Argon laser suitable for constructs with CFP (Cerulean, 458 nm/480-520 nm) derivates.

Statistical analysis

Differences between genotypes were assessed by a Student's *t*-test for comparison between wild-type plants and mutant lines or by using one-way analysis of variance (ANOVA) with a Tukey HSD post-hoc test. The sPLS-DA analysis was performed according to Jiang et al. (2014) and Lê Cao et al. (2011). Irrelevant variables were removed using lasso (least absolute shrinkage and selection operator) penalizations and 20 variables were selected in each dimension. The 'mixOmics' package (Rohart et al., 2017) was used to perform sPLS-DA. All the statistical analysis and graph production were done in RStudio (version 1.1.456) (Rstudio Team, 2015), which incorporates the R software package (version 3.5.1) (R Core Team, 2017) using 'ggplot2' (Wickham, 2016), 'ggthemes' (Arnold, 2019), 'cowplot' (Wilke, 2019), 'hyperSpec' (Beleites and Sergo, 2020) and 'multcompView'(Graves et al., 2015).

ACCESSION NUMBERS

Sequence data from this article can be found in the GenBank/EMBL data libraries under the following accession numbers: *SWEET16* (AT3G16690), *SWEET17* (AT4G15920).

ACKNOWLEDGMENTS

The authors would like to thank Dr. Fabien Chardon and Dr. Anne Krapp (IJPB, INRAE Versailles, France) for providing seeds of the *sweet17-1* mutant. The authors also thank Dr. Grégory Mouille (IJPB, INRAE Versailles, France) for advice on FTIR dataset analysis.

FIGURE LEGENDS

Figure 1. Altered development of the inflorescence stem in the *swt16swt17* double
mutant.

(A) Transverse sections of the basal part of the inflorescence stem of 7-week-old plants
stained with FASGA solution. Bars = 200 µm.
(B-F) Boxplots showing the inflorescence stem cross-sectional area (B), the area occupied by
xylem tissue (C) or phloem tissue (D) within a stem section, the ratio of xylem area to stem
area (E) and the ratio of phloem area to stem area (F).
The box and whisker plots represent values from 7 to 8 independent plants. The lines
represent median values, the tops and bottoms of the boxes represent the first and third
quartiles respectively, and whisker extremities represent maximum and minimum data points.
A one-way analysis of variance combined with the Tukey's comparison post-hoc test were
performed. Values marked with the same letter were not significantly different from each
other, whereas different letters indicate significant differences ($P < 0.05$).

Figure 2. Knockout of *SWEET16* gene expression impacts the proliferation of xylem
cells.

(A to I) Boxplots showing the inflorescence stem height (A) and diameter (B), the ratio of
vessel number to fiber number (C), the cross-sectional area occupied by xylem tissue per
vascular bundle (D), the average cross-sectional area of a xylary fiber (E) or of a xylem vessel
(F), and the average number of xylem cells (G), of xylary fiber vessels (H) and of xylem
vessels (I) per vascular bundle.
The box and whisker plots represent values from 5 to 7 independent plants (A and B) or from
71, 53, 41 and 50 individual vascular bundles from wild type, *swt16*, *swt17* and *swt16swt17*
respectively taken from 5-7 independent plants for each genotype (C-I). The lines represent
median values, the tops and bottoms of the boxes represent the first and third quartiles
respectively, and whisker extremities represent maximum and minimum data points. Stars

denote significant differences between the mutant lines compared to the wild type according
to a Student's *t*-test (* *P*<0.05).

Figure 3. Genes involved in the development of xylem secondary cell wall are
deregulated in the *swt16swt17* double mutant.
(A to I) The mRNAs were extracted from total inflorescence stems collected from 7-week-old
wild-type, *swt16*, *swt17* and *swt16swt17* plants. The mRNA contents are expressed relative to
those of the reference gene *UBQ5*.

(A) Heatmap of expression of candidate genes involved in xylem development and secondary
cell wall biosynthesis in the inflorescence stem of the wild type, *swt16*, *swt17* and *swt16swt17*
mutants. The values used to build the heatmap are the mean accumulation of transcripts (*n*=4)
normalized by the median value of each gene and expressed as $\log_2$ values. For each gene, the
result of one-way ANOVA is displayed beside the heatmap. Those *p*-values below the
significance threshold (*P*<0.05) are in red.

(B to I) Boxplots showing the relative expression of *SND1* (B), *MYB46* (C), *VNI2* (D), *MYB4*
(E), *MYB83* (F), *MYB54* (G), *SND3* (H) and *MYB103* (I). The box-and-whisker plots represent
values from 4 biological replicates. The lines represent median values, the tops and bottoms
of the boxes represent the first and third quartiles respectively, and the ends of the whiskers
represent maximum and minimum data points. Black dots are outliers. Stars denote significant
differences between the mutant line compared to the wild type according to a Student's *t*-test
(* *P*<0.05; ** *P*<0.01; *** *P*<0.001). The experiment was repeated twice and gave similar
results.

Figure 4. The composition of the xylem secondary cell wall is altered in the *swt16swt17*
double mutant.
FTIR spectra were acquired on xylem tissue from sections of the basal part of the
inflorescence stem. All spectra were baseline-corrected and area-normalized in the range
1800-800 cm^{-1}.
(A) Average FTIR spectra were generated from 266, 170, 123 and 170 spectra for the wild
type, *swt16*, *swt17* and *swt16swt17* respectively, obtained using three independent plants for
each genotype.

(B) Comparison of FTIR spectra obtained from xylem cells of the *swt16*, *swt17*, and

swt16swt17 mutants. A Student's *t*-test was performed to compare the absorbances for the

wild type, single and double mutants. The results were plotted against the corresponding

wavenumbers. T-values (vertical axis) between -2 and +2 correspond to non-significant

differences (*p*-value < 0.05) between the genotypes tested (n=3). T-values above +2 or below

-2 correspond to, respectively, significantly weaker or stronger absorbances in the mutant

spectra relative to the wild type.

(C to E) Boxplots of the cellulose (C-O vibration band at 1050 cm^{-1}) (C), hemicellulose (C-O

and C-C bond stretching at 1740 cm^{-1}) (D) peak height and lignin peak height ratio

(1510/1595 cm^{-1}) (E). The lines represent median values, the tops and bottoms of the boxes

represent the first and third quartiles respectively, and the ends of the whiskers represent

maximum and minimum data points. The boxplots represent values (shown as colored dots)

from 266, 170, 123 and 170 spectra from the wild type, *swt16*, *swt17* and *swt16swt17*

respectively, obtained from three independent plants for each genotype. A one-way analysis

of variance combined with the Tukey's comparison post-hoc test were performed. Values

marked with the same letter were not significantly different from each other, whereas

different letters indicate significant differences (*P* < 0.05).

Figure 5. Hexoses accumulate in the inflorescence stem of the *swt16swt17* double mutant.

(A and B) Multivariate analysis of the metabolomic datasets obtained from wild-type and

swt16swt17 inflorescence stems. Metabolites were extracted from eight individual plants for

each genotype and analyzed by GC-MS. Plants were grown under long-day conditions for

seven weeks. (A) sPLS-DA score plot for wild-type (purple) and *swt16swt17* (green) samples.

The variable plot of the sPLS-DA is presented in (B) and metabolites in red are significantly

different between the wild type and the *swt16swt17* mutant according to Student's *t*-test

(*P*<0.05) (Supplemental Table 3). ADP: adenosine-5-diphosphate; G-6-P: glucose-6-

phosphate.

(C) Boxplots showing the sucrose, glucose, fructose and starch contents in the inflorescence

stems of the wild type (in purple) and the *swt16swt17* (in green) mutant grown under long-day

conditions for seven weeks. The box-and-whisker plots represent values from 9 biological

replicates coming from plants grown at two separated times. The lines represent median

values, the tops and bottoms of the boxes represent the first and third quartiles respectively,

and the ends of the whiskers represent maximum and minimum data points. Black dots are
outliers. Asterisks above the boxes indicate statistical differences between genotypes
according to Student's *t*-test ($P<0.05$).

Figure 6. *SWEET16* and *SWEET17* expression patterns vary during inflorescence stem
development and the two proteins interact physically.

(A to C) *pSWEET16:GUS* expression pattern in sections taken at different positions in the
inflorescence stem of 8-week-old plants. (D to F) *pSWEET17:GUS* expression pattern in
sections taken at different positions in the inflorescence stem section of 8-week-old plants.
Sections were taken in a stem region where the growth was still rapid (A, D and inset), in a
stem region where elongation growth had finished but where thickening of the secondary cell
wall was still ongoing (B, E and inset), and at the bottom of the stem, a region that
corresponds to a mature stem (C, F and inset). Arrows point to cells showing blue GUS
staining and asterisks indicate xylary parenchyma cells. Lignin is colored pink after
phloroglucinol staining. The intensity of the pink color is correlated with the stage of
lignification of the xylary vessels. ep: epidermis; co: cortex; iff: interfascicular fibers; ph:
phloem; xy: xylem.

(G to I) Fluorescence in an Arabidopsis mesophyll protoplast expressing *SWEET16:cCFP-C*
and *SWEET17:nCerulean-C* revealed by false color cyan (G). Chloroplast auto-fluorescence
is in false color red (H). Merge of (G) and (H) and bright field image (I). Invaginations around
the chloroplasts in (G) and (I) indicate that SWEET16 and SWEET17 interact at the vacuolar
membrane. Scale bar = 17 µm.

Figure 7. Model for the role of SWEET transporters during xylem development in
Arabidopsis inflorescence stems.
This model is based on the results presented in this work on SWEET16 and SWEET17 and
those previously published on SWEET11, SWEET12 and SUC2 transporters (Truernit and
Sauer, 1995; Chen et al., 2012; Gould et al., 2012; Le Hir et al., 2015). In the phloem tissue,
sucrose and hexoses present in the phloem parenchyma cells (PPC) are exported into the

apoplastic space between PPC and companion cells (CC) by the sugar transporters SWEET11 and SWEET12 (fuchsia circles). Apoplastic sucrose is then imported into the CC cytosol *via* the SUC2 transporter (yellow circles) before entering the phloem sieve tubes (SE) and being transported over long distances (light green arrows). A part of these sugars leaks from the SE, most probably through plasmodesmata (orange arrow), and reaches axial sinks (e.g. procambium and xylem) while another part of the sugars is reimported inside the SE, mostly through the action of SUC2 (1). In the cells at the cambium-xylem boundary, soluble sugars are probably exported by SWEET16 (light blue circles) into the cytosol in order to sustain the division of xylem cells (2). Given the high cytosolic sugar demand required to sustain the secondary cell wall (SCW) deposition process (3), sugars stored in the vacuole are likely exported into the cytosol through the action of SWEET16 (purple circles) and/or SWEET17 (green circles). Interaction between SWEET16 and SWEET17 is shown as bicolor circles. After the completion of programmed cell death (PCD) and the disintegration of the vacuole (4), the SCW is still being reinforced (5) and we can assume that the sugar demand is still high. At this stage, the sugars stored in the vacuole of the xylary parenchyma cells (XPC) are likely released by SWEET16 and/or SWEET17 and then exported into the apoplastic space by SWEET11 and SWEET12. Whether it is the sugars themselves or more complex cell wall sugar-derived molecules that reach the dead xylem cells remains an open question.

REFERENCES

Åkerholm M, Salmén L (2001) Interactions between wood polymers studied by dynamic FT-IR spectroscopy. Polymer (Guildf) 42: 963–969

Ambavaram MMR, Krishnan A, Trijatmiko KR, Pereira A (2011) Coordinated activation of cellulose and repression of lignin biosynthesis pathways in rice. Plant Physiol 155: 916–931

Arnold JB (2019) Ggthemes: extra themes, scales and geoms for "ggplot2."

Barcelo RA (2005) Xylem parenchyma cells deliver the H2O2 necessary for lignification in differentiating xylem vessels. Planta 220: 747–756

Beleites C, Sergo V (2020) hyperSpec: a package to handle hyperspectral data sets in R. CRAN Packag.

Cañas RA, Yesbergenova-Cuny Z, Belanger L, Rouster J, Brulé L, Gilard F, Quilleré I, Sallaud C, Hirel B (2020) NADH-GOGAT overexpression does not improve maize (Zea mays L.) performance even when pyramiding with NAD-IDH, GDH and GS. Plants. doi: 10.3390/plants9020130

Chardon F, Bedu M, Calenge F, Klemens PAW, Spinner L, Clement G, Chietera G, Léran S, Ferrand M, Lacombe B, et al (2013) Leaf fructose content is controlled by the vacuolar transporter SWEET17 in *Arabidopsis*. Curr Biol 23: 697–702

Chen L-Q, Hou B-H, Lalonde S, Takanaga H, Hartung ML, Qu X-Q, Guo W-J, Kim J-G, Underwood W, Chaudhuri B, et al (2010) Sugar transporters for intercellular exchange and nutrition of pathogens. Nature 468: 527–532

Chen L-Q, Qu X-Q, Hou B-H, Sosso D, Osorio S, Fernie AR, Frommer WB (2012) Sucrose efflux mediated by SWEET proteins as a key step for phloem transport. Science (80-) 335: 207–211

Clough SJ, Bent AF (1998) Floral dip: A simplified method for Agrobacterium-mediated transformation of *Arabidopsis thaliana*. Plant J 16: 735–743

Curtis MD, Grossniklaus U (2003) A gateway cloning vectors set for high-throughput functional analysis of genes in planta. Plant Physiol 133: 462–469

Damari-Weissler H, Rachamilevitch S, Aloni R, German MA, Cohen S, Zwieniecki MA,
Holbrook NM, Granot D (2009) LeFRK2 is required for phloem and xylem
differentiation and the transport of both sugar and water. Planta 230: 795–805

Dinant S, Wolff N, De Marco F, Vilaine F, Gissot L, Aubry E, Sandt C, Bellini C, Le Hir R
(2019) Synchrotron FTIR and Raman spectroscopy provide unique spectral fingerprints
for *Arabidopsis* floral stem vascular tissues. J Exp Bot 70: 871–884

Etchells JP, Provost CM, Mishr L, Turner SR (2013) WOX4 and WOX14 act downstream of
the PXY receptor kinase to regulate plant vascular proliferation independently of any
role in vascular organisation. Dev 140: 2224–2234

Etchells JP, Smit ME, Gaudinier A, Williams CJ, Brady SM (2016) A brief history of the
TDIF-PXY signalling module: Balancing meristem identity and differentiation during
vascular development. New Phytol 209: 474–484

Eveland AL, Jackson DP (2012) Sugars, signalling, and plant development. J Exp Bot 63:
3367–3377

Faix O (1991) Classification of lignins from different botanical origins by FT-IR
spectroscopy. Holzforschung-International J Biol Chem Phys Technol Wood 45: 21–27

Fukuda H, Ohashi-Ito K (2019) Vascular tissue development in plants, 1st ed. Curr Top Dev
Biol. doi: 10.1016/bs.ctdb.2018.10.005

Fürtauer L, Küstner L, Weckwerth W, Heyer AG, Nägele T (2019) Resolving subcellular
plant metabolism. Plant J 0–3

Furze ME, Trumbore S, Hartmann H (2018) Detours on the phloem sugar highway: stem
carbon storage and remobilization. Curr Opin Plant Biol 43: 89–95

Geng P, Zhang S, Liu J, Zhao C, Wu J, Cao Y, Fu C, Han X, He H, Zhao Q (2020) MYB20,
MYB42, MYB43, and MYB85 regulate phenylalanine and lignin biosynthesis during
secondary cell wall formation1[OPEN]. Plant Physiol 182: 1272–1283

Gou JY, Park S, Yu XH, Miller LM, Liu CJ (2008) Compositional characterization and
imaging of "wall-bound" acylesters of Populus trichocarpa reveal differential
accumulation of acyl molecules in normal and reactive woods. Planta 229: 15–24

Gould N, Thorpe MR, Pritchard J, Christeller JT, Williams LE, Roeb G, Schurr U, Minchin

PEH (2012) AtSUC2 has a role for sucrose retrieval along the phloem pathway: Evidence from carbon-11 tracer studies. Plant Sci 188–189: 97–101

Graves S, Piepho H-P, Selzer L, Dorai-Raj S (2015) multcompView: Visualizations of paired comparisons.

Guo W-J, Nagy R, Chen H-Y, Pfrunder S, Yu Y-C, Santelia D, Frommer WB, Martinoia E (2014) SWEET17, a facilitative transporter, mediates fructose transport across the tonoplast of Arabidopsis roots and leaves. Plant Physiol 164: 777–789

Gutbrod K, Romer J, Dörmann P (2019) Phytol metabolism in plants. Prog Lipid Res 74: 1–17

Hall H, Ellis B (2013) Transcriptional programming during cell wall maturation in the expanding Arabidopsis stem. BMC Plant Biol 13: 14

Hasanuzzaman M, Nahar K, Anee TI, Fujita M (2017) Glutathione in plants: biosynthesis and physiological role in environmental stress tolerance. Physiol Mol Biol Plants 23: 249–268

Hedrich R, Sauer N, Neuhaus HE (2015) Sugar transport across the plant vacuolar membrane: Nature and regulation of carrier proteins. Curr Opin Plant Biol. doi: 10.1016/j.pbi.2015.04.008

Heineke D, Wildenberger K, Sonnewald U, Willmitzer L, Heldt HW (1994) Accumulation of hexoses in leaf vacuoles: Studies with transgenic tobacco plants expressing yeast-derived invertase in the cytosol, vacuole or apoplasm. Planta 194: 29–33

Le Hir R, Spinner L, Klemens PAW, Chakraborti D, De Marco F, Vilaine F, Wolff N, Lemoine R, Porcheron B, Géry C, et al (2015) Disruption of the Sugar Transporters AtSWEET11 and AtSWEET12 Affects Vascular Development and Freezing Tolerance in *Arabidopsis*. Mol Plant 8: 1687–1690

Hussey SG, Mizrachi E, Creux NM, Myburg AA (2013) Navigating the transcriptional roadmap regulating plant secondary cell wall deposition. Front Plant Sci 4: 1–21

Jiang M, Wang C, Zhang Y, Feng Y, Wang Y, Zhu Y (2014) Sparse partial-least-squares discriminant analysis for different geographical origins of salvia miltiorrhiza by 1h-nmr-based metabolomics. Phytochem Anal 25: 50–58

Jin H, Cominelli E, Bailey P, Parr A, Mehrtens F, Jones J, Tonelli C, Wiesshaar B, Martin C
(2000) Transcriptional repression by AtMYB4 controls production of UV-protecting
sunscreens in Arabidopsis. EMBO J 19: 6150–6161

Kačuráková M, Smith AC, Gidley MJ, Wilson RH (2002) Molecular interactions in bacterial
cellulose composites studied by 1D FT-IR and dynamic 2D FT-IR spectroscopy.
Carbohydr Res 337: 1145–1153

Karimi M, Inze D, Depicker A (2002) GATEWAY((TM)) vectors for Agrobacterium-
mediated plant transformation. Trends Plant Sci 7: 193–195

Klemens PAW, Patzke K, Deitmer J, Spinner L, Le Hir R, Bellini C, Bedu M, Chardon F,
Krapp A, Neuhaus HE (2013) Overexpression of the vacuolar sugar carrier *AtSWEET16*
modifies germination, growth, and stress tolerance in Arabidopsis. Plant Physiol 163:
1338–1352

Klemens PAW, Patzke K, Trentmann O, Poschet G, Büttner M, Schulz A, Marten I, Hedrich
R, Neuhaus HE (2014) Overexpression of a proton-coupled vacuolar glucose exporter
impairs freezing tolerance and seed germination. New Phytol 202: 188–197

Koch KE (2004) Sucrose metabolism: regulatory mechanisms and pivotal roles in suger
sensing and plant development. Curr Opin Plant Biol 7: 235–246

Koncz C, Schell J (1986) The promoter of TL-DNA gene 5 controls the tissue-specific
expression of chimaeric genes carried by a novel type of Agrobacterium binary vector.
Mol Gen Genet 204: 383–396

Lahlali R, Karunakaran C, Wang L, Willick I, Schmidt M, Liu X, Borondics F, Forseille L,
Fobert PR, Tanino K, et al (2015) Synchrotron based phase contrast X-ray imaging
combined with FTIR spectroscopy reveals structural and biomolecular differences in
spikelets play a significant role in resistance to Fusarium in wheat. BMC Plant Biol. doi:
10.1186/s12870-014-0357-5

Lê Cao KA, Boitard S, Besse P (2011) Sparse PLS discriminant analysis: Biologically
relevant feature selection and graphical displays for multiclass problems. BMC
Bioinformatics. doi: 10.1186/1471-2105-12-253

Lee LY, Fang MJ, Kuang LY, Gelvin SB (2008) Vectors for multi-color bimolecular

fluorescence complementation to investigate protein-protein interactions in living plant
cells. Plant Methods 4: 1–11

Li E, Bhargava A, Qiang W, Friedmann MC, Forneris N, Savidge RA, Johnson LA,
Mansfield SD, Ellis BE, Douglas CJ (2012) The Class II KNOX gene KNAT7
negatively regulates secondary wall formation in Arabidopsis and is functionally
conserved in Populus. New Phytol 194: 102–115

Li Z, Omranian N, Neumetzler L, wang ting, Herter T, Usadel B, Demura T, Giavalisco P,
Nikoloski Z, Persson S (2016) A Transcriptional and Metabolic Framework for
Secondary Wall Formation in Arabidopsis. Plant Physiol pp.01100.2016

Lucas WJ, Groover A, Lichtenberger R, Furuta K, Yadav SR, Helariutta Y, He XQ, Fukuda
H, Kang J, Brady SM, et al (2013) The Plant Vascular System: Evolution, Development
and Functions. J Integr Plant Biol 55: 294–388

Mahboubi A, Ratke C, Gorzsás A, Kumar M, Mellerowicz EJ, Niittylä T (2013) Aspen
SUCROSE TRANSPORTER3 allocates carbon into wood fibers. Plant Physiol 163:
1729–1740

Marriott PE, Gómez LD, Mcqueen-Mason SJ (2016) Unlocking the potential of
lignocellulosic biomass through plant science. New Phytol 209: 1366–1381

Martinoia E (2018) Vacuolar transporters - Companions on a longtime journey. Plant Physiol
176: pp.01481.2017

Minchin PEH, McNaughton GS (1987) Xylem transport of recently fixed carbon within lupin.
Aust. J. Plant Physiol. 14:

Mohebby B (2008) Application of ATR infrared spectroscopy in wood acetylation. J Agric
Sci Technol 10: 253–259

Öhman D, Demedts B, Kumar M, Gerber L, Gorzsás A, Goeminne G, Hedenström M, Ellis B,
Boerjan W, Sundberg B (2013) MYB103 is required for FERULATE-5-
HYDROXYLASE expression and syringyl lignin biosynthesis in Arabidopsis stems.
Plant J 73: 63–76

Payyavula RS, Tay KHC, Tsai C-J, Harding SA (2011) The sucrose transporter family in
Populus: the importance of a tonoplast PtaSUT4 to biomass and carbon partitioning.

Plant J 65: 757–770

Poschet G, Hannich B, Raab S, Jungkunz I, Klemens PAW, Krueger S, Wic S, Neuhaus HE,
Buttner M (2011) A Novel Arabidopsis Vacuolar Glucose Exporter Is Involved in
Cellular Sugar Homeostasis and Affects the Composition of Seed Storage Compounds.
Plant Physiol 157: 1664–1676

R Core Team (2017) R: a language and environment for statistical computing.

Roach M, Arrivault S, Mahboubi A, Krohn N, Sulpice R, Stitt M, Niittylä T (2017) Spatially
resolved metabolic analysis reveals a central role for transcriptional control in carbon
allocation to wood. J Exp Bot 68: 3529–3539

Rohart F, Gautier B, Singh A, Lê Cao KA (2017) mixOmics: An R package for 'omics feature
selection and multiple data integration. PLoS Comput Biol 13: 1–19

Rstudio Team (2015) RStudio: Integrated Development for R.

Sakr S, Wang M, Dédaldéchamp F, Perez-Garcia M-D, Ogé L, Hamama L, Atanassova R
(2018) The Sugar-Signaling Hub: Overview of Regulators and Interaction with the
Hormonal and Metabolic Network. Int J Mol Sci 19: 2506

Schuetz M, Smith R, Ellis B (2012) Xylem tissue specification, patterning, and differentiation
mechanisms. J Exp Bot 64: 11–31

Sellami S, Le Hir R, Thorpe MR, Vilaine F, Wolff N, Brini F, Dinant S (2019) Salinity
effects on sugar homeostasis and vascular anatomy in the stem of the *Arabidopsis*
thaliana inflorescence. Int J Mol Sci 20: 3167

Sibout R, Plantegenet S, Hardtke CS (2008) Flowering as a Condition for Xylem Expansion
in *Arabidopsis* Hypocotyl and Root. Curr Biol 18: 458–463

Smetana O, Mäkilä R, Lyu M, Amiryousefi A, Rodríguez FS, Wu M, Solé-gil A, Gavarrón
ML, Siligato R, Miyashima S, et al (2019) High levels of auxin signalling define the
stem-cell organizer of the vascular cambium. Nature 565: 485–491

Smit M, McGregor S, Sun H, Gough C, Bågman A-M, Soyars CL, Kroon JTM, Gaudinier A,
Williams CJ, Yang X, et al (2019) A PXY-Mediated Transcriptional Network Integrates
Signaling Mechanisms to Control Vascular Development in Arabidopsis. Plant Cell
tpc.00562.2019

Smith RA, Schuetz M, Karlen SD, Bird D, Tokunaga N, Sato Y, Mansfield SD, Ralph J,
Samuels AL (2017) Defining the diverse cell populations contributing to lignification in
Arabidopsis thaliana stems. Plant Physiol 174: 1028–1036

Smith RA, Schuetz M, Roach M, Mansfield SD, Ellis B, Samuels L (2013) Neighboring
parenchyma cells contribute to *Arabidopsis* xylem lignification, while lignification of
interfascicular fibers is cell autonomous. Plant Cell 25: 3988–3999

Spicer R (2014) Symplasmic networks in secondary vascular tissues: Parenchyma distribution
and activity supporting long-distance transport. J Exp Bot 65: 1829–1848

Tolivia D, Tolivia J (1987) Fasga: A new polychromatic method for simultaneous and
differential staining of plant tissues. J Microsc 148: 113–117

Truernit E, Sauer N (1995) The promoter of the Arabidopsis thaliana SUC2 sucrose-H+
symporter gene directs expression of B-glucuronidase to the phloem: evidence for
phloem loading and unloading by SUC2. Planta 196: 564–570

Vandesompele J, De Preter K, Pattyn F, Poppe B, Van Roy N, De Paepe A, F. S (2002)
Accurate normalization of real-time quantitative RT-PCR data by geometric averaging of
multiple internal control genes. Genome Biol 3: research0034.1-0034.11

Verbančič J, Lunn JE, Stitt M, Persson S (2018) Carbon supply and the regulation of cell wall
synthesis. Mol Plant 11: 75–94

Weiszmann J, Fürtauer L, Weckwerth W, Nägele T (2018) Vacuolar sucrose cleavage
prevents limitation of cytosolic carbohydrate metabolism and stabilizes photosynthesis
under abiotic stress. FEBS J. doi: 10.1111/febs.14656

Wickham H (2016) ggplot2: Elegant Graphics for Data Analysis. Springer-Verlag, New York

Wilke CO (2019) Cowplot: streamlined plot theme and plot annotation for "ggplot2."

Wingenter K, Schulz A, Wormit A, Wic S, Trentmann O, Hoermiller II, Heyer AG, Marten I,
Hedrich R, Neuhaus HE (2010) Increased Activity of the Vacuolar Monosaccharide
Transporter TMT1 Alters Cellular Sugar Partitioning, Sugar Signaling, and Seed Yield in
Arabidopsis. Plant Physiol 154: 665–677

Xuan YH, Hu YB, Chen L-Q, Sosso D, Ducat DC, Hou B-H, Frommer WB (2013)
Functional role of oligomerization for bacterial and plant SWEET sugar transporter

family. Proc Natl Acad Sci 110: E3685–E3694

Yamada K, Osakabe Y, Mizoi J, Nakashima K, Fujita Y, Shinozaki K, Yamaguchi-Shinozaki
K (2010) Functional analysis of an Arabidopsis thaliana abiotic stress-inducible
facilitated diffusion transporter for monosaccharides. J Biol Chem 285: 1138–1146

Yamaguchi M, Ohtani M, Mitsuda N, Kubo M, Ohme-Takagi M, Fukuda H, Demura T
(2010) VND-INTERACTING2, a NAC Domain Transcription Factor, Negatively
Regulates Xylem Vessel Formation in Arabidopsis. Plant Cell 22: 1249–1263

Yoo SD, Cho YH, Sheen J (2007) Arabidopsis mesophyll protoplasts: A versatile cell system
for transient gene expression analysis. Nat Protoc 2: 1565–1572

Zheng M, Liu X, Lin J, Liu X, Wang Z, Xin M, Yao Y, Peng H, Zhou DX, Ni Z, et al (2019)
Histone acetyltransferase GCN5 contributes to cell wall integrity and salt stress tolerance
by altering the expression of cellulose synthesis genes. Plant J 97: 587–602

Zhong R, Demura T, Ye Z-H (2006) SND1, a NAC Domain Transcription Factor, Is a Key
Regulator of Secondary Wall Synthesis in Fibers of Arabidopsis. Plant Cell Online 18:
3158–3170

Zhong R, Lee C, Zhou J, McCarthy RL, Ye Z-H (2008) A Battery of Transcription Factors
Involved in the Regulation of Secondary Cell Wall Biosynthesis in Arabidopsis. Plant
Cell 20: 2763–2782

1027

1028

1029

Figure 1. Altered development of the inflorescence stem in the *swt16swt17* double mutant.

(A) Transverse sections of the basal part of the inflorescence stem of 7-week-old plants stained with FASGA solution. Bars = 200 µm.

(B to F) Boxplots showing the inflorescence stem cross-sectional area (B), the area occupied by xylem tissue (C) or phloem tissue (D) within a stem section, the ratio of xylem area to stem area (E) and the ratio of phloem area to stem area (F).

The box and whisker plots represent values from 7 to 8 independent plants. The lines represent median values, the tops and bottoms of the boxes represent the first and third quartiles respectively, and whisker extremities represent maximum and minimum data points. A one-way analysis of variance combined with the Tukey's comparison post-hoc test were performed. The values marked with the same letter were not significantly different from each other, whereas different letters indicate significant differences ($P < 0.05$).

Figure 2. Knockout of the *SWEET16* gene expression impacts the proliferation of xylem cells.

(A to I) Boxplots showing the inflorescence stem height (A) and diameter (B), the ratio of vessel number to fiber number (C), the cross-sectional area occupied by xylem tissue per vascular bundle (D), the average cross-sectional area of a xylary fiber (E) or of a xylem vessel (F), and the average number of xylem cells (G), of xylary fiber vessels (H) and of xylem vessels (I) per vascular bundle.

The box and whisker plots represent values from 5 to 7 independent plants (A and B) or from 71, 53, 41 and 50 individual vascular bundles from wild type, *swt16*, *swt17* and *swt16swt17* respectively coming from 5-7 independent plants for each genotype (C-I). The lines represent median values, the tops and bottoms of the boxes represent the first and third quartiles respectively, and whisker extremities represent maximum and minimum data points. Stars denote significant differences between the mutant lines compared to the wild type according to a Student's *t*-test (* *P*<0.05).

1059

1060

1061

Figure 3. Genes involved in the development of xylem secondary cell wall are deregulated in the *swt16swt17* double mutant.

(A to I) mRNAs were extracted from total inflorescence stems collected from 7-week-old wild-type, *swt16*, *swt17* and *swt16swt17* plants. The mRNA contents are expressed relative to those of the reference gene *UBQ5*.

(A) Heatmap of expression of candidate genes involved in xylem development and secondary cell wall biosynthesis in the inflorescence stem of the wild type, *swt16*, *swt17* and *swt16swt17* mutants. The values used to build the heatmap are the mean accumulation of transcripts (*n*=4) normalized by the median value of each gene and expressed as $\log_2$ values. For each gene, the result of one-way ANOVA is displayed beside the heatmap. Those *p*-values below the significance threshold ($P<0.05$) are in red.

(B to I) Boxplots showing the relative expression of *SND1* (B), *MYB46* (C), *VNI2* (D), *MYB4* (E), *MYB83* (F), *MYB54* (G), *SND3* (H) and *MYB103* (I). The box-and-whisker plots represent values from 4 biological replicates. The lines represent median values, the tops and bottoms

1076 of the boxes represent the first and third quartiles respectively, and the ends of the whiskers
1077 represent maximum and minimum data points. Black dots are outliers. Stars denote significant
1078 differences between the mutant line compared to the wild type according to a Student's t-test
1079 (* $P<0.05$; ** $P<0.01$; *** $P<0.001$). The experiment was repeated twice and gave similar
1080 results.

1081

1082

1083

Figure 4. The composition of the xylem secondary cell wall is impaired in the *swt16swt17*
double mutant.

FTIR spectra were acquired on xylem tissue from sections of the basal part of the
inflorescence stem. All spectra were baseline-corrected and area-normalized in the range
1800-800 cm^{-1}.

(A) Average FTIR spectra were generated from 266, 170, 123 and 170 spectra for the wild
type, *swt16*, *swt17* and *swt16swt17* respectively obtained using three independent plants for
each genotype.

(B) Comparison of FTIR spectra obtained from xylem cells of the *swt16*, *swt17*, and
swt16swt17 mutants. A Student's *t*-test was performed to compare the absorbances for the
wild type, single and double mutants. The results were plotted against the corresponding
wavenumbers. T-values (vertical axis) between -2 and +2 correspond to non-significant

differences (p-value < 0.05) between the tested genotypes (n=3). T-values above +2 or below

-2 correspond to, respectively, significantly weaker or stronger absorbances in the mutant

spectra relative to the wild type.

(C to E) Boxplots of the cellulose (C-O vibration band at 1050 cm^{-1}) (C), hemicellulose (C-O

and C-C bond stretching at 1740 cm^{-1}) (D) peak height and lignin peak height ratio

(1510/1595 cm^{-1}) (E). The lines represent median values, the tops and bottoms of the boxes

represent the first and third quartiles respectively, and the ends of the whiskers represent

maximum and minimum data points. The boxplots represent values (shown as colored dots)

from 266, 170, 123 and 170 spectra from the wild type, *swt16*, *swt17* and *swt16swt17*

respectively obtained from three independent plants for each genotype. A one-way analysis of

variance combined with the Tukey's comparison post-hoc test were performed. Values

marked with the same letter were not significantly different from each other, whereas

different letters indicate significant differences ($P < 0.05$).

1109

Figure 5. Hexoses accumulate in the inflorescence stem of the *swt16swt17* double mutant.

(A and B) Multivariate analysis of the metabolomic datasets obtained from wild-type and *swt16swt17* inflorescence stems. Metabolites were extracted and analyzed by GC-MS from eight individual plants for each genotype. Plants were grown under long-day conditions for seven weeks. (A) sPLS-DA score plot for wild-type (purple) and *swt16swt17* (green) samples. The variable plot for the sPLS-DA is presented in (C) and metabolites in red are significantly different between the wild type and the *swt16swt17* mutant according to Student's *t*-test

(*P*<0.05) (Supplemental Table 3). ADP: adenosine-5-diphosphate; G-6-P: glucose-6-

phosphate.

(C) Boxplots showing the sucrose, glucose, fructose and starch contents of the inflorescence

stems of the wild type (in purple) and the *swt16swt17* (in green) mutant grown under long-day

conditions for seven weeks. The box-and-whisker plots represent values from 9 biological

replicates coming from plants grown at two separate times The lines represent median values,

the tops and bottoms of the boxes represent the first and third quartiles respectively, and the

ends of the whiskers represent maximum and minimum data points. Black dots are outliers.

Asterisks above the boxes indicate statistical differences between genotypes according to

Student's *t*-test (*P*<0.05).

1127

1128

1129

Figure 6. *SWEET16* and *SWEET17* **expression patterns vary during the inflorescence stem development and the two proteins interact physically.**

(A to C) *pSWEET16:GUS* expression pattern in sections taken at different positions in the inflorescence stem of 8-week-old plants. (D to F) *pSWEET17:GUS* expression pattern in sections taken at different positions in the inflorescence stem section of 8-week-old plants. Sections were taken in a stem region where growth was still rapid (A, D and inset), in a stem region where elongation growth had finished but where thickening of the secondary cell wall

was still ongoing (B, E and inset), and at the bottom of the stem, a region that corresponds to

a mature stem (C, F and inset). Arrows point to cells showing blue GUS staining and asterisks

indicate xylary parenchyma cells. Lignin is colored pink after phloroglucinol staining. The

intensity of the pink color is correlated with the stage of lignification of the xylary vessels. ep:

epidermis; co: cortex; iff: interfascicular fibers; ph: phloem; xy: xylem.

(G to I) Fluorescence in an Arabidopsis mesophyll protoplast expressing *SWEET16:cCFP-C*

and *SWEET17:nCerulean-C* revealed by false color cyan (G). Chloroplast auto-fluorescence

is in false color red (H). Merge of (G) and (H) and bright field image (I). Invaginations around

the chloroplasts in (G) and (I) indicate that SWEET16 and SWEET17 interact at the vacuolar

membrane. Scale bar = 17 µm.

1147

1148

1149

Figure 7. Model for the role of SWEET transporters during xylem development in the Arabidopsis inflorescence stem.

This model is based on the results presented in this work on SWEET16 and SWEET17 and those previously published on SWEET11, SWEET12 and SUC2 transporters (Truernit and Sauer, 1995; Chen et al., 2012; Gould et al., 2012; Le Hir et al., 2015). In the phloem tissue, sucrose and hexoses present in the phloem parenchyma cells (PPC) are exported into the apoplastic space between PPC and companion cells (CC) by the sugar transporters SWEET11 and SWEET12 (fuchsia circles). Apoplastic sucrose is then imported into the CC cytosol, *via* the SUC2 transporter (yellow circles), before entering the phloem sieve tubes (SE) and being transported over long distances (light green arrows). A part of these sugars leaks from the SE, most probably through plasmodesmata (orange arrow), and reaches axial sinks (e.g. procambium and xylem) while another part of the sugars is reimported inside the SE, mostly through the action of SUC2 (1). In the cells at the cambium-xylem boundary, soluble sugars are probably exported by SWEET16 (light blue circles) into the cytosol in order to sustain the division of xylem calls (2). Given the high cytosolic sugar demand required to sustain the secondary cell wall (SCW) deposition process (3), sugars stored in the vacuole are likely exported into the cytosol through the action of SWEET16 (purple circles) and/or SWEET17 (green circles). Interaction between SWEET16 and SWEET17 is shown as bicolor circles. After the completion of programmed cell death (PCD) and the disintegration of the vacuole (4), the SCW is still being reinforced (5) and we can assume that the sugar demand is still high. At this stage, the sugars stored in the vacuole of the xylary parenchyma cells (XPC) are likely released by SWEET16 and/or SWEET17 and then exported into the apoplastic space by

1174 SWEET11 and SWEET12. Whether it is the sugars themselves or more complex cell wall

1175 sugar-derived molecules that reach the dead xylem cells remains an open question.