**Global query on bacterial sirtuin CobB interactants reveals crosstalk with PRPP synthase Prs**

Beata M. Walter[1], Joanna Morcinek-Orłowska[1], Aneta Szulc[1], Andrew L. Lovering[2], Manuel Banzhaf[2], Monika Glinkowska[1] ✉

1- Department of Bacterial Molecular Genetics, Faculty of Biology, University of Gdansk, Gdansk, Poland

2 - Institute of Microbiology & Infection and School of Biosciences, University of Birmingham, UK;

✉ - to whom correspondence should be addressed;  monika.glinkowska@ug.edu.pl

## Abstract

Protein lysine acetylation regulates a wide range of cellular functions. It is controlled by a family of NAD-dependent protein deacetylases called sirtuins. In eukaryotes, sirtuins activity is coupled to spatiotemporally-controlled NAD+ level, whereas the mechanism of their regulation in bacteria is less clear. *E. coli* possesses a single sirtuin – CobB. However, it is unclear how CobB activity is coupled to NAD+ metabolism. In this work we show that this coordination is achieved in *E. coli* cells through a CobB interaction with PRPP synthase Prs, an enzyme necessary for NAD+ synthesis. Employing global analysis of protein-protein interactions formed by CobB, we demonstrate that it forms a stable complex with Prs. This assembly stimulates CobB deacetylase activity and partially protects it from inhibition by nicotinamide. We provide evidence that Prs acetylation is not necessary for CobB binding but affects the global acetylome *in vivo*. Our results show that CobB ameliorates Prs activity under conditions of Prs cofactors deficiency. Therefore, we propose that CobB-Prs crosstalk orchestrates the NAD+ metabolism and protein acetylation in response to environmental cues.

**Key words:** Protein acetylation/ sirtuins/ NAD metabolism/ phosporibosyl pyrophosphate

## Intoduction

Lysine acetylation is a post-translational protein modification regulating a wide range of protein functions. It has been investigated thoroughly in eukaryotic cells and recently was discovered to be important for protein regulation in prokaryotes as well (Christensen *et al*, 2019; Bernal *et al*, 2014). In bacteria, the level of protein acetylation is a result of two counterbalancing processes. On one hand, proteins become acetylated enzymatically or chemically, by lysine transacetylases or acetyl phosphate, respectively (Ma & Wood, 2011; Weinert *et al*, 2013; Kuhn *et al*, 2014). In the opposing process, acetyl groups are removed from acetyl-lysine residues by deacetylases, most of which are NAD+-dependent homologs of eukaryotic sirtuins (Greiss & Gartner, 2009; Imai & Guarente, 2010).

CobB is a highly conserved protein lysine deacetylase among bacteria (Landry *et al*, 2000; Tsang & Escalante-Semerena, 1998). It removes the acetyl group from acetyl-lysines, utilizing NAD+ and producing nicotinamide (NAM) and 2''-O-acetyl-ADP-ribose as byproducts. CobB is also the only sirtuin-like deacetylase identified in *E. coli* so far (Zhao *et al*, 2004). Its activity regulates global protein acetylation level (Choudhary *et al*, 2014; Castaño-Cerezo *et al*, 2014; Weinert *et al*, 2017, 2013; Kuhn *et al*, 2014) and thus affects many cellular functions, including gene expression (Thao *et al*, 2010; Lima *et al*, 2011; Qin *et al*, 2016), the cell cycle (Zhang *et al*, 2016), metabolism (Castaño-Cerezo *et al*, 2011; Venkat *et al*, 2017; Castaño-Cerezo *et al*, 2014), stress response[3,12], motility and pathogenicity (Liu *et al*, 2018). In addition, CobB can also act as desuccinylase, de-2-hydroxyisobutyrylase and lipoamidase (Colak *et al*, 2013; Dong *et al*, 2019; Rowland *et al*, 2017). Despite its important role, factors affecting protein deacetylation rate in bacteria are poorly understood. The levels of CobB reaction product NAM (Gallego-Jara *et al*, 2017), as well as NADH and c-di-GMP (Xu *et al*) have been implicated in regulation of CobB activity so far.

In eukaryotic cells, interplay between activity of sirtuins and NAD+ metabolism is well established, showing that NAD+ is a crucial factor in controlling chromatin structure, DNA repair, lifespan and circadian rhythm (Imai & Guarente, 2016; James Theoga Raj & Lin, 2019). This makes NAD+ not only an enzyme cofactor in various redox reactions but also an important signaling molecule. Bacterial sirtuins on the other hand, have low $K_m$'s for NAD+

whereas its intracellular concentration is high (Guan *et al*, 2014). This may jointly result in
lower sensitivity of bacterial sirtuins to regulation by NAD+ than their eukaryotic
homologues. *In vivo*, the level of NAD+ is maintained by *de novo* synthesis and salvage
pathways (Fig. 1). Both pathways require the pivotal metabolite phosphoribosyl
pyrophosphate (PRPP) that is produced by the evolutionary conserved PRPP synthase – Prs
(Gazzaniga *et al*, 2009).
In this work we propose that coordination between NAD+ metabolism and protein acetylation
is exerted in *E.coli* by crosstalk of the CobB deacetylase with the PRPP synthase Prs. Namely,
we provide evidence that CobB forms a stable complex with Prs affecting its deacetylase
activity. Formation of this complex is not dependent on Prs acetylation status *in vitro* but
modification of Prs by lysine acetylation affects CobB function *in vivo*. We also show that,
under certain conditions, the rate of PRPP synthesis by Prs may be influenced by CobB.
Based on those observations, we propose a model how this mutual dependence links NAD+
metabolism and cellular energetics to protein acetylation.
107
108

## Results

**CobB interacts with phosphoribosyl phosphate synthase *in vivo* and *in vitro*.** Numerous
acetylated proteins have been suggested to be CobB targets *in vivo*. Some of them were found
in global acetylome studies showing differential acetylation of proteins in wild-type and
*ΔcobB* mutants (Castaño-Cerezo *et al*, 2014; Weinert *et al*, 2017). Others were identified as
CobB interactants with the use of a protein microarray consisting of the majority of the *E. coli*
proteome (~4000 protein, non-acetylated) (Liu *et al*, 2014). In order to investigate the
regulation of CobB-mediated protein deacetylation we first aimed to identify its interaction
partners under physiologically relevant conditions. For this, we performed an
immunoprecipitation pull-down, dubbed sequential peptide affinity purification (SPA)(Babu
*et al*, 2009). In this approach, proteins are labeled with a double-affinity tag consisting of
triple FLAG epitope and calmodulin binding protein (CBP), separated by a protease cleavage
site. After immunoprecipitation from cell lysates with anti-FLAG antibodies, protein
complexes are released from the resin by protease digest and re-purified using a calmodulin
sepharose column. As a bait, we used chromosomally expressed (from the native promoter),
C-terminally SPA-tagged CobB. Subsequently to the pull-down, interaction partners were
identified by liquid chromatography coupled to tandem mass spectrometry (LC-MS/MS). The
experiments have been carried out using cells grown to stationary phase in a rich undefined
medium (LB) and minimal medium containing acetate as a carbon source, both conditions
supporting high protein acetylation level. Under those conditions, the major component of the
130
131

purified CobB complex was the phosphoribosyl phosphate synthase Prs (Supplementary
Table 1). This interaction has been previously described by others in a large-scale study of
protein-protein interactions, confirming validity of our results (Butland *et al*, 2005). We
further compared our data with the interactions previously described in a CobB protein
microarray study (Liu *et al*, 2014). We found the fatty acids synthesis proteins FabB and
FabG are significantly enriched in the presence of CobB, in comparison to control samples
analyzed by LC-MS/MS. However, other interactants found in this microarray study were not
present or enriched in our data set, suggesting that those interactions are of lower affinity or
more transient than the one between CobB and Prs (Supplementary Table 1).
We further confirmed the Prs and CobB interaction performing the pull-down experiment in a
reversed set-up, where SPA-tagged Prs was used as a bait (Supplementary Table 1).

Consistently, we observed binding of purified His-tagged Prs (bait) and CobB (prey) in a pull-
down experiment (Fig. 2A).
Overall, these results indicate that CobB forms a stable complex with Prs in *E. coli*.
146
**CobB-Prs interaction is not affected by the Prs acetylation state.** Prs has been previously
reported as one of the proteins acetylated *in vivo* at widely-conserved positions K182, K194,
K231(Kuhn *et al*, 2014; Castaño-Cerezo *et al*, 2014; Weinert *et al*, 2017) (Supplementary Fig.
1). Therefore, we investigated whether Prs interaction with the CobB is affected by
acetylation of its lysine residues. To achieve this, we repeated the pull-down experiment using
acetylated His-tagged Prs and CobB. Prs was acetylated *in vitro* by reacting with acetyl
phosphate, as described before by others (Kuhn *et al*, 2014; Qin *et al*, 2016). Subsequently,
successful
modification of lysine residues was confirmed by western blot (Fig. 2B). In addition, we
verified by mass spectrometry that acetylated Prs lysine residues are mainly those previously
shown to become modified this way *in vivo* (K182, K194, K231)(Table 1). Chemically
acetylated Prs interacted with CobB in a manner indistinguishable from the unmodified
protein (Fig. 2C). It is worth noting that the pull-down reactions did not contain NAD+,
disabling deacetylation of Prs by CobB during the course of reaction. Next, we confirmed that
Prs becomes deacetylated by CobB *in vitro* (Fig. 2D, Table 1). Our experiments revealed also
that CobB likewise becomes acetylated by acetyl phosphate in a system consisting of purified
proteins (Fig. 2B) and acetylation has a moderate influence on its deacetylase activity (Fig.
2D). Acetylated lysine residues of the sirtuin were subsequently identified by MS and are
presented in Supplementary Table 2. Moreover, our result also showed that CobB
undergoes auto-deacetylation *in vitro* (Fig. 2D, Supplementary Table 1). It remains unknown
if CobB acetylation is of physiological importance. However, neither Prs nor CobB
acetylation had any impact on their protein-protein interaction compared to the wt and non-
acetylated protein (Fig. 2C). This suggests that acetylated lysine residues
in Prs do not take part in its interaction with the CobB deacetylase. We further probed this by
repeating the pull downs with purified Prs variants where the acetylable lysines K182, K194
and K231 were substituted by alanines. All three variants were still able to interact with CobB
(Fig. 2EF). Corroborating those results, we found that CobB was pulled-down by Prs in *ΔpatZ*
and *Δpta* strains (Supplementary Table 1). The former strain is devoid of protein lysine
acetyltranferase (Ma & Wood, 2011), the latter lacks phosphate acetylase (Wolfe, 2005)
which synthesizes acetyl phosphate. The two strains are characterized by decreased level of
enzymatic and chemical acetylation, respectively (Schilling *et al*, 2015; Weinert *et al*, 2013).
In summary, our results have shown that Prs and CobB form a complex and their interaction
does not rely on Prs acetylation, nor is it influenced by the acetylation state of both proteins *in*
*vitro*.
181
**Prs stimulates the deacetylase activity of CobB and partially protects it from the**
**inhibitory effect of nicotinamide.** Next, we investigated the consequences of complex
formation between Prs and CobB on their catalytic activity. Prs produces one of the key
metabolites for the synthesis of NAD+(Hove-Jensen *et al*, 2017) (Fig. 1) which is a substrate
of CobB in deacetylation. Therefore, a possible role of their interaction could be to orchestrate
the sirtuin activity of the NAD+ metabolism. First, we investigated the effect of Prs-CobB
assembly on CobB-mediated removal of acetyl groups from modified lysines. To elucidate
this, we measured deacetylation rate of an artificial fluorogenic substrate for Zn2+ and NAD+
dependent deacetylases – MAL (BOC-Ac-Lys-AMC)(Heltweg *et al*, 2003). In the presence of
Prs, CobB activity in deacetylating MAL was stimulated by about 50% (Fig. 3A). The degree
of stimulation was dependent on Prs concentration and reached maximum in 6:1 molar ratio

of Prs to CobB (Prs monomer : CobB monomer). A further increase of Prs concentration had no effect on CobB activity, suggesting that Prs hexamers stimulate a single catalytic center of CobB. As expected, CobB deacetylase activity was dependent on NAD+ concentration and Prs enhanced it in a wide range of NAD+ concentrations, even in those lower than the physiological ones (Bennett *et al*, 2009) (Fig. 3B). Neither Prs substrates (ribose 5-phosphate, ATP) nor the products of its catalytic activity (PRPP, AMP) and allosteric inhibitor ADP, had significant impact on its biding to CobB or ameliorating deacetylation of MAL substrate (Supplementary Fig. 2). CobB, as other sirtuins, was shown to be sensitive to feedback inhibition by deacetylation reaction byproduct – nicotinamide (NAM), with $IC_{50}$ value estimated at approximately 52 µM (Gallego-Jara *et al*, 2017) (Fig. 3C). NAM is also one of the metabolites of the *E. coli* NAD+ salvage pathway I and its intracellular level has been shown to fluctuate dependent on bacterial growth conditions (Gallego-Jara *et al*, 2017). This makes NAM a likely candidate for a regulator of CobB activity *in vivo*. Therefore, we tested how interaction with Prs affects NAM-mediated inhibition of CobB deacetylase activity. In the presence of Prs, deacetylation of MAL substrate by CobB was more effective even at very high NAM concentrations, indicating that Prs partially protects CobB from inhibition by NAM (Fig. 3D). We also confirmed using pull-down assay that CobB forms a complex with Prs in the presence of NAM (Supplementary Fig. 2).

Moreover, it has been shown previously that several metabolites of the NAD salvage pathways, like nicotinamide mononucleotide (NMN), can act as weak inhibitors of Sir2 family deacetylases *in vitro* (Schmidt *et al*, 2004). Corroborating those results, we observed that NMN, as well as nicotinic acid adenine dinucleotide (NaAD), NADH and NADP negatively influence CobB activity at high concentrations (Fig. 3C). As in the case of NAM, Prs increased CobB activity in the presence of those metabolites (Fig. 3D, Supplementary Fig. 3).

Those results suggest that Prs-CobB interaction influences catalytic activity of CobB by increasing its efficiency as a deacetylase. Formation of a complex with Prs also partially protects CobB from inhibition by NAM and other NAD+ metabolites, further suggesting that the interaction exerts an impact on the catalytic center of CobB.

**Proteome acetylation is altered in strains producing of Prs variants K182A and K231A.**
PRPP, produced by Prs, is a necessary metabolite for *de novo* synthesis of nucleotides, cofactors, histidine and tryptophan (Hove-Jensen *et al*, 2017), making Prs an essential enzyme under standard bacterial growth conditions (Baba *et al*, 2006). Therefore, testing Prs influence on CobB activity in *E. coli* cells is challenging. However, we found that mutations in the *prs* gene, affecting acetylation of the PRPP synthase, result in profound changes in the global proteome acetylation level. To assess those alterations, we measured protein acetylation in *E. coli* cells, using Western blot and anti-N(ε)-acetyl lysine antibody. Cells were sampled in middle-exponential phase, when overall acetylation is low, and also in early stationary phase, when protein acetylation increases (Schilling *et al*, 2015; Weinert *et al*, 2013). In those experiments, cells that chromosomally express *prs* variants K182A and K231A showed higher protein acetylation level than their wild-type counterpart upon entry to stationary phase (Fig. 4A). Since Prs produces a precursor for NAD+ synthesis changes in the enzyme's activity could affect intracellular NAD+ concentration, ultimately leading to differences in CobB-mediated protein deacetylation rate. Therefore, product synthesis by both Prs variants was measured *in vitro*, showing that Prs K231A had a similar activity as wild-type Prswhereas the activity of Prs K182A was slightly decreased (Fig. 4B). The latter result was also consistent with a slower growth rate of the cognate strain. Moreover, both variants stimulated CobB-mediated deacetylation of MAL substrate *in vitro* (Fig. 4C), confirming that they retain the activity of the wild-type protein. PRPP formation by Prs acetylated *in vitro* by acetyl

phosphate was lower by about 15%, whereas the main lysine residue acetylated under those
conditions is K182 (Supplementary Fig. 4). Together, those results suggest that the increased
protein lysine acetylation in strains producing Prs that does not undergo acetylation at
positions 182 or 231 is unlikely to be a mere result of altered PRPP concentration. Further
evidence that higher proteome acetylation level in those strains is connected to cooperation
between Prs and CobB is provided by Western blot results obtained with double mutants *prs*
(K182A or K231A) *ΔcobB* (Fig. 4A). Protein acetylation profile of the double mutant, upon
cell entry to the stationary phase, was practically indistinguishable from that of the single
*ΔcobB* mutant, particularly for the strain *prs* (K231A) *ΔcobB,* suggesting an epistatic effect
(Fig. 4A). Mass spectrometry analysis of gel bands showing high acetylation signal after
probing with anti-N(ε)-acetyl lysine antibody, revealed that their most prominent components
were Lpd, Pgk and also GadA-GadB. All those proteins have been previously reported to be
acetylated *in vivo* and their acetylation level was affected in a *ΔcobB* mutant (Zhang *et al*,
2013; Castaño-Cerezo *et al*, 2014). Thus, our data corroborate previous findings of other
groups and suggest that although Prs acetylation is not required for Prs-CobB interaction (Fig.
2CEF), it may play a role *in vivo* in sustaining CobB-mediated deacetylation rate during the
stationary phase.
Lpd (lipoamide dehydrogenase) is a component of pyruvate dehydrogenase, α-ketoglutarate
dehydrogenase and the glycine cleavage system, taking part in glycolysis, TCA cycle and
glycine catabolism, respectively (Carothers *et al*, 1989). Pgk (phosphoglycerate kinase) on the
other hand, participates in glycolysis and gluconeogenesis (Nellemann *et al*, 1989), whereas
GadA and GadB are homologous decarboxylases playing a major role in glutamate-dependent
acid resistance system (Foster, 2004). Considering the role of identified proteins, our results
also imply that Prs-CobB interplay is important for regulation of cellular energetics and the
stress response.
268
269
**CobB stimulates Prs activity *in vitro* under suboptimal conditions and impacts NAD+**
**metabolism *in vivo*.** The results presented so far strongly suggest that formation of the Prs-
CobB complex impacts the rate of acetyl group removal from acetylated lysine residues. We
next asked, whether assembly with CobB reciprocally influences Prs activity. Prs synthesizes
PRPP from ribose 5-phosphate and ATP, producing AMP as a byproduct (Hove-Jensen *et al*,
2017). It utilizes magnesium and phosphate ions as a cofactor and allosteric activator,
respectively (Willemoes *et al*, 2000). To assess Prs catalytic activity, we monitored AMP
formation using a luciferase based assay that produces a luminescent signal proportional to
AMP present in the samples.
Results of this assay showed that addition of CobB alone, CobB together with its substrate
NAD+ or product NAM had little impact on Prs activity, when sufficient amount of
magnesium and phosphate ions was provided (Supplementary Fig. 5). Conversely, when
assays were carried out under conditions of low magnesium or phosphate concentration,
CobB significantly stimulated Prs activity (Fig. 5A). This suggests that under suboptimal
conditions for the Prs function,
assembly with CobB may balance PRPP synthesis by Prs. The intracellular magnesium level
has been previously implicated as one of the factors regulating protein acetylation in *E. coli*
cells. Namely, protein acetylation was higher in cells grown in magnesium-limited
media(Christensen *et al*, 2017). This effect could be due to insufficient CobB stimulation by
Prs. However, at least *in vitro*, magnesium ions concentration had no impact on the
enhancement of CobB-mediated deacetylation rate by Prs (Supplementary Fig. 6). Interaction
with CobB also slightly enhanced feedback inhibition of Prs activity by PRPP (Fig. 5B).

Together, those result show that, at least under certain conditions, formation of the complex
with CobB may also influence Prs function.
NAD+ is a key cofactor of many enzymes and its concentration in *E. coli* cells is high,
within millimolar range(Bennett *et al*, 2009). It has been demonstrated that in eukaryotic
cells, NAD+ -consuming enzymes, including sirtuins, substantially contribute to NAD+
expenditure and impact NAD homeostasis (Strømland *et al*, 2019). This relationship in *E. coli*
has not been investigated, whereas its existence would support the need of coupling between
deacetylation and NAD+ biosynthesis and explain reciprocal effect of Prs and CobB on their
activity. Therefore, we were interested whether the absence of CobB function will influence
the level of NAD+ metabolites. We employed metabolomics to quantitatively assess the
intracellular levels of those compounds in wild-type and *ΔcobB* strains. Indeed, we observed
that in absence of CobB, concentration of all metabolites of the NAD synthesis and salvage
pathways were raised by about 25% (Fig. 6), suggesting that CobB is an important player in
NAD+ turnover.
306
**Discussion**
In this work we have provided evidence that the CobB deacetylase and the PRPP
synthase form a complex in *E. coli*. (Supplementary Table 1, Fig. 2A) Within this complex,
both proteins can affect each other's activity (Fig. 3, 5). Why would cells need such an ally?
Lysine acetylation is a protein modification that affects protein activity. It allows cells to
quickly adjust the available pool of active proteins in response to environmental cues,
avoiding energy consuming degradation and synthesis. To be efficient as a fast, adaptable
regulatory system, acetylation needs to be reversible. This is provided by the action of
deacetylases, which in bacteria are mostly NAD+ -dependent. This makes NAD+
concentration a possible mean to regulate the rate of protein deacetylation. In eukaryotic cells,
a clear link has been shown between sirtuins activity and processes influencing the level of
NAD+ and its metabolites, like NAM (Lu & Lin, 2010; Zhang & Sauve, 2018; Anderson *et*
*al*, 2017; Imai & Guarente, 2016). Sirtuins activation has also been observed under various
stress conditions, including low energy level manifested by ATP depletion (Kang *et al*, 2017;
Nogueiras *et al*, 2012). Bacterial sirtuins have low $K_m$ values for NAD+ and prokaryotic cells
lack compartmentalization that could ensure spatial regulation of NAD+ level. In addition,
NADH is rather a weak inhibitor of CobB activity, whereas intracellular NADH concentration
is around two orders of magnitude lower than that of NAD+ (Bennett *et al*, 2009). All those
factors may limit the possibility of a direct regulation of CobB activity by NAD+ or
NAD/NADH ratio in bacterial cells. Dynamics of NAD+ synthesis under various growth
conditions is poorly understood. Nevertheless, NAD+ synthesis *de novo* and through salvage
pathways requires nicotinamide moiety, ribose phosphate, provided by PRPP, and adenine
nucleotide. The two latter metabolites both link NAD+ and cellular energetics to Prs.
330
In the experiments described in this work we have not found conditions that would
break the Prs-CobB complex or restrain its formation. We have also not identified factors that
would disable Prs-mediated stimulation of CobB activity *in vitro*, and could therefore serve as
regulators of the interaction. The known Prs inhibitors - ADP or PRPP had little effect on the
cooperation between Prs and CobB (Supplementary Fig. 2). However, while the *cobB* gene is
constitutively expressed (Castaño-Cerezo *et al*, 2011), *prs* is regulated at transcriptional level
and activated by low purine and pyrimidine nucleotides concentration (He *et al*, 1993; White
*et al*, 1971). Hence, the amount of Prs-CobB complexes formed in the cell and overall CobB
deacetylation rate would depend on the demand to synthesize new purine nucleotides,
including ATP precursors (Fig. 7). Moreover, several purine synthesis enzymes, like PurA
and Adk were found among proteins acetylated in *E. coli* cells (Weinert *et al*, 2013; Zhang *et*

*al*, 2013). The effect of acetylation on those enzymes has not been determined so far but acetylation has usually an inhibitory effect. Thus, transcriptional activation of *prs* could ultimately lead to more effective protein deacetylation by CobB and, among other effects, to an increase in the activity of purine synthesis pathway. At the same time, assembly with Prs would partially overcome inhibition of CobB by accumulating deacetylation product – NAM (Fig. 3D). In the stationary phase, Prs acetylation at lysines K182 and K231 could stimulate CobB activity (Fig. 4A) or have an effect on Prs stability, enabling an efficient interaction between the two proteins. In another direction, CobB stimulates Prs when magnesium or phosphate level fall to meet NAD+ demands (Fig. 5A).

In summary, the crosstalk between Prs and CobB allows *E. coli* cells to integrate various cues and adjust protein acetylation level to NAD+ metabolism.

## Methods

### Materials, reagents and strains

Primers used in this study were synthesized by Sigma/Merck. List of primers, vectors and strains is available in Supplementary table 3. Polymerases and enzymes used for cloning were purchased from Thermo Scientific or New England Biolabs. Reagents used for buffers were purchased from either Carl Roth or Bioshop Life Science. $NAD^+$ salvage pathway substrates were purchased from Sigma/Merck except for nicotinamide (NAM) which was purchased from Bioshop Life Science.

### Cloning, expression and purification of recombinant proteins

His-Prs and catalytically inactive His-PrsK194A were expressed from pET28a-His-*prs* and pET28a-His-*prs*K194A vectors in *E. coli* Rosetta (DE3) as we described earlier (Walter *et al*, 2020). Additional point mutations replacing acetylated lysines K182 and K231 for alanines were introduced into pET28a-His-*prs* vector with phosphorylated primers (Walter *et al*, 2020). *E. coli* Rosetta (DE3) was transformed with pET28a-His-*prs*K231A, grown to $OD_{600}$ between 0.8 and 1.0 and induced with 100 µM Isopropyl β-D-1-thiogalactopyranoside (IPTG) at 37°C for 5 h in 2xYT medium (BioShop). *E. coli* BL21-DE3-pLysE was transformed with pET28A-His-*prs*K182A and induced at $OD_{600}$ between 0.8 and 1.0 with 1 mM IPTG at 37°C for 3 h in 2xYT medium.

The *cobB* sequence, amplified on *E. coli* MG1655 genomic DNA was N-terminally cloned with RF-cloning (Bond & Naus, 2012) into modified pET28a-TEV vector (Walter *et al*, 2020). The *E. coli* Rosetta (DE3) was transformed with vector pET28a-His-TEV-*cobB*, grown in Terrific Broth (BioShop) to $OD_{600}$ between 0.8 and 1.0 and induced with 200 µM IPTG at 37°C for 5 h.

His-Prs protein and its variants were purified at 20°C, as described earlier (Walter *et al*, 2020) with buffer A (50 mM potassium phosphate pH 7.5, 10 % glycerol, 500 mM NaCl, 20 mM imidazole pH 7.8); buffer B (50 mM potassium phosphate pH 7.5, 10 % glycerol, 500 mM NaCl, 300 mM imidazole pH 7.8, 0.5 mM tris(2-carboxyethyl)phosphine (TCEP)) and dialysis SEC1 buffer (50 mM potassium phosphate pH 8.2, 10 % glycerol, 500 mM NaCl).

Non-acetylated CobB was purified on His-Trap columns (GE healthcare) as described above however, the lysate was loaded on the column and protein eluted on ice. Further, fractions containing protein were combined and diluted 1:1 with buffer A-0 (50 mM potassium phosphate pH 8.2, 10 % glycerol) and supplemented with Dithiothreitol (DTT) and ethylenediaminetetraacetic acid (EDTA) to a final concentration 1 mM and 0.5 mM respectively. Aliquots of 20 ml were made. TEV protease (0.2 mg) purified in-house[54] was added and incubated at 24 °C for 3 h with no shaking or rotation. After incubation, NaCl concentration was readjusted to 500 mM by spiking with 5M NaCl stock solution and protein

solution was dialyzed overnight into SEC1 buffer at 4$^{o}$C followed by protein separation from His tag on a His-Trap column (GE Healthcare), concentration with an Amicon Ultra 15 MWCO10kDa (Millipore) filter and size-exclusion chromatography at 4 $^{o}$C on a Superdex 200 10/300 GL gel filtration column (GE Healthcare). Purity of the proteins was evaluated on SDS-PAGE gel. Concentrated to 10-15 mg ml$^{-1}$ purified proteins were snap-frozen in liquid nitrogen and stored at -70$^{o}$C.

**Acetylation of His-Prs and CobB proteins during protein purification**
His-Prs was purified on His-Trap columns (GE Healthcare) as described above and dialyzed overnight at 20 $^{o}$C into 1 x acetylation buffer (100 mM Tris-HCl pH 7.5, 10 % glycerol, 150 mM NaCl) [modified from (Kuhn *et al*, 2014; Qin *et al*, 2016)]. Further, it was concentrated using the Amicon Ultra 15 MWCO30kDa (Millipore) filter at 18 $^{o}$C to 4 mg ml$^{-1}$ and acetylated with acetyl phosphate (lithium potassium salt, Sigma) at final concentration 20 mM at 24 $^{o}$C for 3 h. Following acetylation, His-Prs was concentrated at 18 $^{o}$C (Amicon Ultra 15 MWCO30kDa) to approximately 3 – 4 ml and further purified by size-exclusion chromatography at 18 $^{o}$C on Superdex 200 10/300 GL gel filtration column (GE Healthcare). His-TEV-CobB, prior to acetylation, was purified and washed on His-Trap columns (GE Healthcare) with buffer Ac (50 mM Tris pH 7.9, 500 mM NaCl, 20 mM imidazole pH 7.8, 10 % glycerol) and eluted on ice with buffer Bc (50 mM Tris pH 7.9, 500 mM NaCl, 300 mM imidazole pH 7.8, 10 % glycerol). Fractions containing protein were combined, diluted 1:1 with buffer A-0 and supplemented with DTT (1 mM), EDTA (0.5 mM) and TEV protease (0.2 mg). Further, fractions were incubated for 3 h at 20$^{o}$C as above, followed by overnight dialysis at 4$^{o}$C into acetylation dialysis buffer (100 mM Tris, pH 8.2, 300 mM NaCl, 10 % glycerol). The CobB protein was separated from His tag on His-Trap column (GE Healthcare), concentrated at 4 $^{o}$C to 6 mg ml$^{-1}$, diluted 1 : 1 in 0 x acetylation buffer (100 mM Tris-HCl pH 7.5, 10 % glycerol) and acetylated with acetyl phosphate at final concentration 20 mM at 24 $^{o}$C for 3 h. Protein solution was further dialyzed overnight into SEC1 buffer at 4 $^{o}$C.
Purity of proteins was evaluated on 10 % SDS-PAGE gel. Purified proteins, concentrated to 10-15 mg ml$^{-1}$, were snap-frozen in liquid nitrogen and stored at -70$^{o}$C. Acetylation was evaluated with Western blot probed with anti-acetyl lysine antibodies (1:800 dilution) (Sigma/Merck). Mass spectrometry outsourced to Mass Spectrometry Laboratory IBB PAN, Warsaw, Poland. Two independent batches of both His-PrsAc and CobBAc subjected to acetyl phosphate treatments were purified and acetylation was assessed.

**CobB pull-down on His-Prs and its variants**.
His-Prs or its variants (10 µg) were bound to 4 µl of Ni-NTA Magnetic Agarose Beads (Qiagen) in 50 µl interaction buffer (50 mM Tris pH 9.0, 30 mM imidazole pH 7.8, 300 mM NaCl, 0.2 % Tween 20, 10 % glycerol) for 1 h on vibrating mixer at 20$^{o}$C. Following triple wash with interaction buffer, 30 µg of CobB was added to His-Prs bound to beads in 50 µl fresh interaction buffer and incubated for 2 h on vibrating mixer at 20$^{o}$C. Beads were washed 4 times with interaction buffer and resuspended in 20 µl 1 x SDS loading dye (31.25 mM Tris pH 6.8, 10 % glycerol, 1 % SDS, bromophenol blue). Proteins were resolved by electrophoresis in 10 % SDS-PAGE gel and visualized with Coomassie Brilliant Blue R250 staining. 4 µg of His-Prs and CobB proteins were loaded on gel separately as control. The experiments were performed in 3 independent repeats and representative gels were shown in figures.

**CobB deacetylase activity on MAL (BOC-Ac-Lys-AMC) substrate**

CobB deacetylase activity was measured as described earlier (Heltweg *et al*, 2003, 2005) as
deacetylation of MAL substrate (Sigma/Merck), an artificial fluorogenic substrate for $Zn^{2+}$
and $NAD^+$ -dependent deacetylases. Briefly, deacetylation of 8 nmol MAL substrate by 320
pmol of CobB was performed for 1 h at 24 $^o$C in 50 µl sirtuin buffer (50 mM Tris-HCl pH 8.5,
137 mM NaCl, 2.7 mM KCl, 1 mM $MgCl_2$) [modified from (Heltweg *et al*, 2003)] in
presence of $NAD^+$ at final concentration 400 µM. The $NAD^+$ titration was performed in
sirtuin buffer supplemented with $NAD^+$ independently at final concentrations 10 µM, 20 µM,
50 µM, 100 µM, 250 µM, 500 µM, 750 µM and 1 mM. The $MgCl_2$ titration was performed in
sirtuin buffer containing 400 µM $NAD^+$ and $MgCl_2$ was supplemented independently at final
concentrations 10 µM, 20 µM, 50 µM, 100 µM, 250 µM, 500 µM, 750 µM, 1 mM, 5 mM and
10 mM. The effect of $NAD^+$ salvage pathway substrates was evaluated by adding 5 µl of 50
mM β-nicotinamide mononucleotide (NMN), nicotinamide (NAM), nicotinic acid (NA),
nicotinic acid mononucleotide (NaMN), nicotinic acid adenine dinucleotide (NaAD) or 10
mM NADH, NADP, NADPH to the reaction. The effect of Prs influence on CobB activity
was measured in sirtuin buffer by addition of 150 pmol, 300 pmol and 600 pmol of Prs
hexamer. The effect of Prs influence on CobB activity in NAD+ titration assay, $MgCl_2$
titration assay and in presence of $NAD^+$ salvage pathway substrates was measured by addition
of 150 pmol of Prs hexamer. All reactions were stopped by addition of 400 µl of 1 M HCl and
fluorescent substrate was extracted as an upper phase after vortexing for 30 s with 800 µl
ethyl acetate and centrifugation (9 500 x g, 5 min). Ethyl acetate was evaporated under the
hood at 65 $^o$C and the residue was dissolved in 600 µl acetonitrile buffer (39.6 % acetonitrile,
5 µM $KH_2PO_4$, 4.6 µM NaOH). The fluorescence (2 x 250 µl) was measured at 330/390 nm
in black, flat bottom 96 well plate (Heltweg *et al*, 2005). The experiments were repeated in at
least 3 independent replicates.

**CobB deacetylase activity on His-PrsAc**

CobB driven deacetylation of His-PrsAc was analyzed by Western blot with anti-acetyl lysine
antibodies (Sigma/Merck) and mass spectrometry. Briefly, CobB driven deacetylation of His-
PrsAc was performed in sirtuin buffer (as above) for 1 h at 24$^o$C in the presence of 400 µM
$NAD^+$. Deacetylation of 147 pmol Prs hexamer (30 µg of protein) was performed in 50 µl
reaction with 650 pmol or 1940 pmol CobB or CobBAc (20 µg and 60 µg of protein
respectively). The reaction was stopped by dilution in 4 x SDS loading dye and 5 minutes
incubation at 100$^o$C. Samples were resolved by electrophoresis in 10 % SDS-PAGE gel in
duplicates with one gel visualized with Coomassie Brilliant Blue R250 staining and the other
transferred to PVDF membrane, probed with anti-acetyl lysine antibodies (1:800 dilution) and
visualized. Selected bands were excised from the gel and outsourced for identification of
protein lysine acetylation by mass spectrometry to Mass Spectrometry Laboratory IBB PAN,
Warsaw, Poland. The experiments were repeated in at least 3 independent replicas and
representative gels are shown in figures.

**Measurement of Prs catalytic activity**

The activity of Prs and its variants was measured as a luminescence signal from AMP formed
during catalytic formation of PRPP from ribose-5-phosphate (60 µM) and ATP (60 µM) for
15 min at 37$^o$C. The assay was carried out with His-Prs protein and its variants His-Prs-
K194A, His-Prs-K182A and His-Prs-K231A, in 100 µl $MgCl_2$ rich reaction buffer (50 mM
Tris pH 8.0, 100 mM KCl, 13 mM $MgCl_2$, 0.5 mM K-phosphate pH 8.0, 0.5 mM DTT, 0.1
mg mL$^{-1}$ BSA) at final Prs hexamer concentration 7 nM (0.7 pmol of hexamer per reaction,
142.8 ng of protein per reaction).
The catalytic activity of His-Prs protein in the presence of CobB at various $MgCl_2$ and
potassium phosphate (pH 8.0) concentrations was measured in 100 µl reaction buffer base (50

mM Tris pH 8.0, 100 mM KCl, 0.5 mM DTT, 0.1 mg mL$^{-1}$ BSA) at final Prs hexamer concentration 7 nM and CobB monomer concentration 40 nM (4 pmol of monomer per reaction, 124 ng of protein per reaction). Reactions were supplemented with 1 mM or 3mM potassium phosphate pH 8.0 and / or 1 mM or 3 mM MgCl$_2$.

The activity of His-Prs in presence of various PRPP concentrations was measured in Prs reaction buffer (50 mM Tris pH 8.0, 100 mM KCl, 1 mM MgCl$_2$, 0.5 mM K-phosphate pH 8.0, 0.5 mM DTT, 0.1 mg ml$^{-1}$ BSA) at final Prs hexamer concentration 7 nM and CobB monomer concentration 40 nM.

Reactions were ceased by placing samples immediately after incubation on iced water. The concentration of AMP was measured with AMP-Glo™ Assay (Promega) according to manufacturer`s protocol. Briefly, the luminescence was measured for 10 µl subsample after adding 10 µl of AMP-Glo™ Reagent I which stopped the reaction, removed remaining ATP (1 h incubation at 24$^o$C) and converted AMP produced into ADP, followed by conversion of ADP to ATP through luciferase reaction with the AMP-Glo™ reagent II (1 h incubation at 24$^o$C). The luminescence was measured in white 96 well plate and the concentration of AMP in experimental samples was calculated based on AMP standard curve measured in triplicates. The catalytic activity of His-Prs protein variants was measured in two independent repeats while the activity in other conditions was measured in at least 3 independent experiments.

**Acetylation level in Prs mutants *in-vivo***

Mutagenesis of MG1655 wild-type strain was performed with lambda Red recombineering as described earlier (Baba *et al*, 2006).

Strains with single mutations replacing acetylated lysine for alanine in *prs* gene: MG1655:*prs*K182A, MG1655:*prs*K231A; strain with *cobB* deletion MG1655:Δ*cobB;* and double mutant strains MG1655:*prs*K182A:Δ*cobBKanR*, MG1655:*prs*K231A:*ΔcobBKanR* were constructed by recombineering (Baba *et al*, 2006).

The acetylation level of the whole *E. coli* proteome was measured in TP7.3 medium (10 g l$^{-1}$ tryptone, buffered at pH 7.3 with 100 mM potassium phosphate) supplemented with 0.2 % glucose at mid logarithmic phase (6 h) and early stationary phase (12 h). Bacterial cells (bacterial culture volume = 15/OD$_{600}$) were harvested and the pellet was snap-frozen in liquid nitrogen. Frozen pellet was resuspended in 450 µl lysis buffer (30 mM Tris pH 6.8, 10 % glycerol, 100 mM NaCl) followed by 35 s sonication with pulsing on ice. Samples were centrifuged 10 min at 16 000 x g, 4$^o$C and protein concentration was measured with Bradford assay. Samples (6 µg) were resolved by electrophoresis in 10 % SDS-PAGE gel in duplicates, with first gel visualized with Coomassie brilliant blue and second transferred to PVDF membrane, probed with anti-acetyl lysine antibodies (1:800 dilution) and visualized. The representative gels and Western blots were repeated in at least 3 independent experiments. Selected bands were excised from gel and outsourced for mass spectrometry identification to Mass Spectrometry Laboratory IBB PAN, Warsaw, Poland.

**NAD+ metabolomics**

Metabolomics was performed by Creative Proteomics. Briefly, MG1655 strain and its *ΔcobB* strain were grown in LB medium with aeration at 37°C to OD$_{600}$ ~0.3. At this point 1ml of sample was withdrawn, cells pelleted and flash-frozen. For the measurements, the cell samples were thawed on ice and each sample (ca. 25 µL) was added with 25 µL of an internal standard (IS) solution containing isotope-labeled NAD, NADH, NA and NAM. Cells were lysed with the aid of two 3-mm metal balls on a MM400 mill mixer for 1 min at 30 Hz. 75 µL of acetonitrile was then added. After vortexing for 10 s and sonication in an ice-water bath for 30 s, the samples were centrifuged at 4$^o$C to pellet protein. Clear supernatants were collected and the pellets were used for protein assay using a standard BCA procedure (Pierce). Standard

solutions: a mixed standard solution containing all the targeted compounds were prepared at 20 nmol/mL in a mixture of IS solution - 50% acetonitrile (1:4). This solution was serially diluted at 1 to 4 (v/v) with the same solution. A Waters Acquity UPLC system coupled to a 4000 QTRAP MS instrument was operated in the mode of multiple-reaction monitoring (MRM)/MS.

*Quantitation of NAD, NADH, NADP, NADPH, NaMN, NMN and NA*

40 μL of each supernatant or each standard solution was diluted 3 fold with water. 20-μL aliquots of the resulting solutions were injected to run LC-MRM/MS with negative-ion detection on a C18 column (2.1x100 mm, 1.8μm) and with an ammonium formate buffer (A) and methanol (B) as the mobile phase for gradient elution (efficient gradient 5% to 50% B in 10 min) at 0.3 mL/min and 40°C.

*Quantitation of NAM*

Mix 25 μL of each supernatant or each standard solution with 4 volumes of 60% acetonitrile. 10 μL aliquots of the resulting solutions were injected to run LC-MRM/MS with positive-ion detection on a HILIC column (2.1x100 mm, 1.7μm) and with the use of 0.1% formic acid (A) and acetonitrile (B) as the mobile phase for gradient elution (efficient gradient 90% to 15% B in 12.5 min) at 0.3 mL/min and 30 °C.

*Analytical Results*

Concentrations of detected compounds were calculated from the constructed linear-regression curve of each compound with internal standard calibration using the analyte-to-IS peak ratios measured from sample solutions.

## Acknowledgements

The authors thank dr Ian Cadby and Krzysztof Sitko for assistance in some of the presented experiments. This work was supported by the National Science Center (Poland) [No. UMO-2014/13/B/NZ2/01139 to M.G].

## Authors contribution

B.W. performed majority of the experimental work, contributed to work conceptualization and preparation of the manuscript; J.M-O. performed the MS analysis of protein complexes and reviewed the manuscript; A.S. performed a part of the biochemical assays; M.B. contributed to project conceptualization and manuscript writing; A.L. contributed to establishment of protein assays and reviewed the manuscript; M.G. contributed to project conceptualization, data analysis and manuscript writing.

## Conflict of interest

The authors declare no conflict of interest

## References

Anderson KA, Madsen AS, Olsen CA & Hirschey MD (2017) Metabolic control by sirtuins and other enzymes that sense NAD+, NADH, or their ratio. *Biochim Biophys Acta - Bioenerg* 1858: 991–998

Baba T, Ara T, Hasegawa M, Takai Y, Okumura Y, Baba M, Datsenko KA, Tomita M, Wanner BL & Mori H (2006) Construction of Escherichia coli K-12 in-frame, single-gene knockout mutants: The Keio collection. *Mol Syst Biol* 2

Babu M, Butland G, Pogoutse O, Li J, Greenblatt JF & Emili A (2009) Sequential peptide affinity purification system for the systematic isolation and identification of protein complexes from Escherichia coli. *Methods Mol Biol* 564: 373–400

Bennett BD, Kimball EH, Gao M, Osterhout R, Van Dien SJ & Rabinowitz JD (2009) Absolute metabolite concentrations and implied enzyme active site occupancy in Escherichia coli. *Nat Chem Biol* 5: 593–599

Bernal V, Castaño-Cerezo S, Gallego-Jara J, Écija-Conesa A, de Diego T, Iborra JL & Cánovas M (2014) Regulation of bacterial physiology by lysine acetylation of proteins. *N Biotechnol* 31: 586–595

Bond SR & Naus CC (2012) RF-Cloning.org: An online tool for the design of restriction-free cloning projects. *Nucleic Acids Res* 40: 209–213

Butland G, Peregrin-Alvarez JM, Li J, Yang W, Yang X, Canadien V, Starostine A, Richards D, Beattie B, Krogan N, *et al* (2005) Interaction network containing conserved and essential protein complexes in Escherichia coli. *Nature* 433: 531–537

Carothers DJ, Pons G & Patel MS (1989) Dihydrolipoamide dehydrogenase: functional similarities and divergent evolution of the pyridine nucleotide-disulfide oxidoreductases. *Arch Biochem Biophys* 268: 409–425

Castaño-Cerezo S, Bernal V, Blanco-Catalá J, Iborra JL & Cánovas M (2011) cAMP-CRP co-ordinates the expression of the protein acetylation pathway with central metabolism in Escherichia coli. *Mol Microbiol* 82: 1110–1128

Castaño-Cerezo S, Bernal V, Post H, Fuhrer T, Cappadona S, Sánchez-Díaz NC, Sauer U, Heck AJ, Altelaar AM & Cánovas M (2014) Protein acetylation affects acetate metabolism, motility and acid stress response in Escherichia coli . *Mol Syst Biol* 10: 762

Choudhary C, Weinert BT, Nishida Y, Verdin E & Mann M (2014) The growing landscape of lysine acetylation links metabolism and cell signalling. *Nat Rev Mol Cell Biol* 15: 536–550

Christensen DG, Orr JS, Rao C V & Wolfe AJ (2017) Increasing Growth Yield and Decreasing Acetylation in Escherichia coli by Optimizing the Carbon-to-Magnesium Ratio in Peptide-Based Media. *Appl Environ Microbiol* 83

Christensen DG, Xie X, Basisty N, Byrnes J, McSweeney S, Schilling B & Wolfe AJ (2019) Post-translational Protein Acetylation: An elegant mechanism for bacteria to dynamically regulate metabolic functions. *Front Microbiol* 10: 1–22

Colak G, Xie Z, Zhu AY, Dai L, Lu Z, Zhang Y, Wan X, Chen Y, Cha YH, Lin H, *et al* (2013) Identification of lysine succinylation substrates and the succinylation regulatory enzyme CobB in escherichia coli. *Mol Cell Proteomics* 12: 3509–3520

Dong H, Zhai G, Chen C, Bai X, Tian S, Hu D, Fan E & Zhang K (2019) Protein lysine de-2-hydroxyisobutyrylation by CobB in prokaryotes. *Sci Adv* 5: eaaw6703

Foster JW (2004) Escherichia coli acid resistance: Tales of an amateur acidophile. *Nat Rev Microbiol* 2: 898–907

Gallego-Jara J, Conesa AÉ, Puente T de D, Terol GL & Díaz MC (2017) Characterization of CobB kinetics and inhibition by nicotinamide. *PLoS One* 12: 1–19

Gazzaniga F, Stebbins R, Chang SZ, McPeek MA & Brenner C (2009) Microbial NAD Metabolism: Lessons from Comparative Genomics. *Microbiol Mol Biol Rev* 73: 529–541

Greiss S & Gartner A (2009) Sirtuin/Sir2 phylogeny, evolutionary considerations and structural conservation. *Mol Cells* 28: 407–415

Guan X, Lin P, Knoll E & Chakrabarti R (2014) Mechanism of inhibition of the human sirtuin enzyme SIRT3 by nicotinamide: Computational and experimental studies. *PLoS One* 9

He B, Choi KY & Zalkin H (1993) Regulation of Escherichia coli glnB, prsA, and speA by
the purine repressor. *J Bacteriol* 175: 3598 LP – 3606
Heltweg B, Dequiedt F, Verdin E & Jung M (2003) Nonisotopic substrate for assaying both
human zinc and NAD+-dependent histone deacetylases. *Anal Biochem* 319: 42–48
Heltweg B, Trapp J & Jung M (2005) In vitro assays for the determination of histone
deacetylase activity. *Methods* 36: 332–337
Hove-Jensen B, Andersen KR, Kilstrup M, Martinussen J, Switzer RL & Willemoes M
(2017) Phosphoribosyl Diphosphate (PRPP): Biosynthesis, Enzymology, Utilization, and
Metabolic Significance. *Microbiol Mol Biol Rev* 81
Imai S & Guarente L (2010) Ten years of NAD-dependent SIR2 family deacetylases:
implications for metabolic  diseases. *Trends Pharmacol Sci* 31: 212–220
Imai SI & Guarente L (2016) It takes two to tango: Nad+ and sirtuins in aging/longevity
control. *npj Aging Mech Dis* 2: 1–6
James Theoga Raj C & Lin SJ (2019) Cross-talk in NAD+ metabolism: insights from
Saccharomyces cerevisiae. *Curr Genet* 65: 1113–1119
Kang H, Oka S, Lee D-Y, Park J, Aponte AM, Jung Y-S, Bitterman J, Zhai P, He Y,
Kooshapur H, *et al* (2017) Sirt1 carboxyl-domain is an ATP-repressible domain that is
transferrable to other proteins. *Nat Commun* 8: 15560
Kuhn ML, Zemaitaitis B, Hu LI, Sahu A, Sorensen D, Minasov G, Lima BP, Scholle M,
Mrksich M, Anderson WF, *et al* (2014) Structural, kinetic and proteomic
characterization of acetyl phosphate-dependent bacterial protein acetylation. *PLoS One* 9
Landry J, Sutton A, Tafrov ST, Heller RC, Stebbins J, Pillus L & Sternglanz R (2000) The
silencing protein SIR2 and its homologs are NAD-dependent protein deacetylases. *Proc
Natl Acad Sci U S A* 97: 5807–5811
Lima BP, Antelmann H, Gronau K, Chi BK, Becher D, Brinsmade SR & Wolfe AJ (2011)
Involvement of protein acetylation in glucose-induced transcription of a stress-
responsive promoter. *Mol Microbiol* 81: 1190–1204
Liu CX, Wu FL, Jiang HW, He X, Guo SJ & Tao SC (2014) Global identification of CobB
interactors by an Escherichia coli proteome microarray. *Acta Biochim Biophys Sin
(Shanghai)* 46: 548–555
Liu W, Tan Y, Cao S, Zhao H, Fang H, Yang X, Wang T, Zhou Y, Yan Y, Han Y, *et al*
(2018) Protein Acetylation Mediated by YfiQ and CobB Is Involved in the Virulence and
Stress Response of <span class="named-content genus-species"
id="named-content-1">Yersinia pestis</span> *Infect Immun* 86:
e00224-18
Lu SP & Lin SJ (2010) Regulation of yeast sirtuins by NAD+ metabolism and calorie
restriction. *Biochim Biophys Acta - Proteins Proteomics* 1804: 1567–1575
Ma Q & Wood TK (2011) Protein acetylation in prokaryotes increases stress resistance.
*Biochem Biophys Res Commun* 410: 846–851
Nellemann LJ, Holm F, Atlung T & Hansen FG (1989) Cloning and characterization of the
Escherichia coli phosphoglycerate kinase (pgk)  gene. *Gene* 77: 185–191
Nogueiras R, Habegger KM, Chaudhary N, Finan B, Banks AS, Dietrich MO, Horvath TL,
Sinclair DA, Pfluger PT & Tschöp MH (2012) Sirtuin 1 and sirtuin 3: physiological
modulators of metabolism. *Physiol Rev* 92: 1479–1514
Qin R, Sang Y, Ren J, Zhang Q, Li S, Cui Z & Yao Y-F (2016) The Bacterial Two-Hybrid
System Uncovers the Involvement of Acetylation in Regulating of Lrp Activity in
Salmonella Typhimurium  . *Front Microbiol*  7: 1864
(https://www.frontiersin.org/article/10.3389/fmicb.2016.01864) [PREPRINT]
Rowland EA, Greco TM, Snowden CK, McCabe AL, Silhavy TJ & Cristea IM (2017) Sirtuin
Lipoamidase Activity Is Conserved in Bacteria as a Regulator of Metabolic  Enzyme

Complexes. *MBio* 8

Schilling B, Christensen D, Davis R, Sahu AK, Hu LI, Walker-Peddakotla A, Sorensen DJ, Zemaitaitis B, Gibson BW & Wolfe AJ (2015) Protein acetylation dynamics in response to carbon overflow in Escherichia coli. *Mol Microbiol* 98: 847–863

Schmidt MT, Smith BC, Jackson MD & Denu JM (2004) Coenzyme specificity of Sir2 protein deacetylases. Implications for physiological regulation. *J Biol Chem* 279: 40122–40129

Strømland Ø, Niere M, Nikiforov AA, VanLinden MR, Heiland I & Ziegler M (2019) Keeping the balance in NAD metabolism. *Biochem Soc Trans* 47: 119–130

Thao S, Chen CS, Zhu H & Escalante-Semerena JC (2010) Nε-lysine acetylation of a bacterial transcription factor inhibits its DNA-binding activity. *PLoS One* 5

Tsang AW & Escalante-Semerena JC (1998) CobB, a new member of the SIR2 family of eucaryotic regulatory proteins, is required to compensate for the lack of nicotinate mononucleotide:5,6-dimethylbenzimidazole phosphoribosyltransferase activity in cobT mutants during cobalamin biosynthesis in Sal. *J Biol Chem* 273: 31788–31794

Venkat S, Gregory C, Sturges J, Gan Q & Fan C (2017) Studying the Lysine Acetylation of Malate Dehydrogenase. *J Mol Biol* 429: 1396–1405

Walter BM, Szulc A & Glinkowska MK (2020) Reliable method for high quality His-tagged and untagged E. coli phosphoribosyl phosphate synthase (Prs) purification. *Protein Expr Purif* 169

Weinert BT, Iesmantavicius V, Wagner SA, Schölz C, Gummesson B, Beli P, Nyström T & Choudhary C (2013) Acetyl-Phosphate is a critical determinant of Lysine Acetylation in E.coli. *Mol Cell* 51: 265–272

Weinert BT, Satpathy S, Hansen BK, Lyon D, Jensen LJ & Choudhary C (2017) Accurate quantification of site-specific acetylation stoichiometry reveals the impact of Sirtuin deacetylase CobB on the E. coli acetylome. *Mol Cell Proteomics* 16: 759–769

White MN, Olszowy J & Switzer RL (1971) Regulation and mechanism of phosphoribosylpyrophosphate synthetase: repression by end products. *J Bacteriol* 108: 122–131

Willemoes M, Hove-Jensen B & Larsen S (2000) Steady state kinetic model for the binding of substrates and allosteric effectors to Escherichia coli phosphoribosyl-diphosphate synthase. *J Biol Chem* 275: 35408–35412

Wolfe AJ (2005) The acetate switch. *Microbiol Mol Biol Rev* 69: 12–50

Xu Z, Zhang H, Zhang X, Jiang H & Liu C Interplay between the protein deacetylase CobB and second messenger c-di-GMP signaling.

Zhang K, Zheng S, Yang JS, Chen Y & Cheng Z (2013) Comprehensive Profiling of Protein Lysine Acetylation in Escherichia coli. *J Proteome Res* 12: 844–851

Zhang N & Sauve AA (2018) Regulatory Effects of NAD+ Metabolic Pathways on Sirtuin Activity. *Prog Mol Biol Transl Sci* 154: 71–104

Zhang Q, Zhou A, Li S, Ni J, Tao J, Lu J, Wan B, Li S, Zhang J, Zhao S, *et al* (2016) Reversible lysine acetylation is involved in DNA replication initiation by regulating activities of initiator DnaA in Escherichia coli. *Sci Rep* 6: 1–23

Zhao K, Chai X & Marmorstein R (2004) Structure and Substrate Binding Properties of cobB, a Sir2 Homolog Protein Deacetylase from Escherichia coli. *J Mol Biol* 337: 731–741

**Figure legends**

**Figure 1. NAD+ producing and consuming pathways in *E. coli*.** NAD+ de novo synthesis from aspartate (thin black arrows) and salvage pathways in *E. coli*:, I (represented jointly by dark and light gray arrows), II (shown by dark gray arrows), IV (from nicotinamide riboside, depicted by thick black arrows). Names of the enzymes engaged in NAD+ metabolism were provided next to the arrows depicting respective reactions. CobB and Prs role in NAD+ transformations is shown. Prs catalyzes PRPP biosynthesis which serves as a substrate for NAD+ formation de-novo from aspartic acid and in NAD+ salvage pathways. NAD+ is essential for CobB mediated protein deacetylation. NAM is a byproduct of this reaction. NAD - nicotinamide adenine dinucleotide, NMN – nicotinamide mononucleotide; NAM – nicotinamide, NA – nicotinic acid, NAMN - nicotinic acid mononucleotide, NaAD – nicotinate adenine dinucleotide, NR – nicotinamide ribonucleotide, R5P – ribose 5-phospahte, PRPP – phosphoribosyl pyrophosphate, Asp – aspartate

**Figure 2. CobB and Prs form a complex *in vitro* irrespective of their acetylation status.**

(A) His$_6$-Prs interacts with CobB in a pull-down assay. His$_6$-Prs was pre-bound to Ni$^{2+}$-coated magnetic beads. After washing off the excess of Prs, the beads were incubated with CobB. Unbound protein was removed by washing and the beads were resuspended in a loading dye and separated in 10% SDS-PAGE gel. Lane 3 shows the extent of CobB binding to magnetic beads in the absence of the bait.

(B) Prs and CobB can be acetylated *in vitro* with acetyl phosphate. Prs and CobB were acetylated in the presence of 20 mM acetyl phosphate and excess of the reagent was dialyzed. Acetylation was confirmed by Western blot with anti-acetyl lysine antibodies.

(C) Chemically acetylated Prs interacts with CobB . Interaction of acetylated proteins was tested *in vitro* in a pull-down assay, as above. The presence of acetylated protein in a reaction was marked with Ac.

(D) Acetylated Prs (Ac) becomes deacetylated *in vitro* by CobB. Acetylated Prs and CobB were incubated in the presence of 400 µM NAD+ (1h). Proteins were separated in 10% SDS-PAGE gel and Western blot with anti-acetyl lysine antibodies was subsequently performed. Numbers below the blot image represent the results of densitometric analysis of bands intensity from three independent blots.

(E) Prs variants with key acetylable lysine residues substituted by alanines interact with CobB and (F) acetylated CobB (Ac). Protein binding was assessed in a pull-down assay, as described above.

**Figure 3. Prs stimulates the CobB deacetylase activity and partially overcomes NAM-mediated inhibition.**

(A) Prs stimulates deacetylation of MAL substrate by CobB. Deacetylation of MAL substrate (8 nmol) by CobB (320 pmol) in the presence and absence of indicated amounts of Prs (as a hexamer) was performed for 1h. Fluorescent substrate was extracted with ethyl acetate and fluorescence was measured at 330/390 nm in a plate reader. Error bars represent SD between 3 independent experiments.

(B) Prs stimulates CobB activity in a wide range of NAD+ concentrations. The assay of MAL deacetylation was performed as described above in the presence of Prs (6 µM) and indicated amounts of NAD+. Mean values of 3 independent experiments were presented. Error bars show SD between the repeats.

(C) The effect of NAD+ metabolites on CobB activity. Deacetylation of MAL substrate by CobB was measured as described above, in the presence of indicated metabolites at 5 mM (NMN, NAM, NaMN, NaAD) or 1mM concentration (NADH, NADP, NADPH). Mean values of 3 independent experiments were presented. Error bars show SD between repeats.

789 (D) Prs partially overcomes inhibition of CobB by NAM. MAL deacetylation was performed
790 as described above in the presence of indicated amounts of NAM and NMN and Prs (3 μM).
791 Mean values of 3 independent experiments were presented. Error bars show SD between the
792 repeats.
793

**Figure 4. Prs variants K182A and K231A affect protein acetylation level *in vivo*.**
(A) Global protein acetylation was assessed in MG1655 strain and its *ΔcobB* derivative,
producing wild-type Prs or its K182A and K231A variant from the *prs* gene in its native
chromosomal position. Overnight cultures of the strains were diluted in TP7.3 medium with
0.2% glucose and samples were collected after 6 and 12 hrs of cell growth. After preparation
of whole cell lysates, 6 μg of proteins from each of them was separated in 10% SDS-PAGE
gel, followed by Western blot with anti-acetyl lysine antibodies.
(B) Prs variants K182A and K231A are active in PRPP synthesis *in vitro*. Prs activity was
assessed by measuring AMP formation from ribose 5-phospate (60 μM) and ATP (60 μM).
After termination of Prs reaction, AMP was converted to ADP and the residual ATP was
removed. Next, ATP was produced from ADP and utilized in luciferase reaction. Reactions
were performed in 96-well plates and luminescence was measured using plate reader. Mean
values of 2 independent experiments were presented. Error bars show SE between the repeats.
(C) Prs variants K182A, K194A and K231A stimulate CobB deacetylase activity *in vitro*.
Deacetylation of MAL substrate (8 nmol) by CobB (320 pmol) in the presence and absence of
Prs (6 μM as a hexamer) was performed for 1h. Fluorescent substrate was extracted with ethyl
acetate and fluorescence was measured at 330/390 nm in a plate reader. Mean values of 3
independent experiments were presented. Error bars show SD between the repeats.
812

**Figure 5. CobB stimulates Prs activity *in vitro* under conditions of low phosphate or**
**magnesium ion concentration.**
(A) Prs reaction was performed in the presence of CobB (1:1 hexamer : monomer ratio) and
$Mg2+$ and $PO4-$ at indicated concentrations. Prs activity was assessed by measuring AMP
formation from ribose 5-phospate (60 μM) and ATP (60 μM). After termination of Prs
reaction, AMP was converted to ADP and the residual ATP was removed. Next, ATP was
produced from ADP and utilized in luciferase reaction. Reactions were performed in 96-well
plates and luminescence was measured using plate reader. Mean values of 3 independent
experiments were presented. Error bars show SD between the repeats.
(B) CobB slightly increases Prs sensitivity to feedback inhibition by PRPP. Prs activity was
measured as described above in the presence of indicated concentrations of PRPP. All
measurements were taken in at least 3 independent repeats. Error bars show SD between the
repeats.
826

**Figure 6. Deletion of the *cobB* gene increases the level of NAD+ metabolites.** MG1655
wild-type strain and its *ΔcobB* derivative were grown in LB medium at 37°C with aeration. 1
ml of the cultures was collected at $OD_{600}$ ~0.3 and disrupted in the presence of isotope-labeled
standards. Metabolites were extracted with acetonitrile (1:3) and detected with LC-MRM/MS.
Concentrations of detected compounds were calculated from the constructed linear-regression
curve of each compound with internal standard (IS) calibration using the analyte-to-IS peak
ratios measured from sample solutions. Protein concentration was measured with bicinchronic
acid kit and protein amount in the sample was taken as proxy of cellular mass present in the
sample.
836

837 **Figure 7. Overview of the interplay between CobB deacetylase and the PRPP synthase**
838 **Prs in *E. coli* cells.** Prs and CobB form a complex in *E. coli* cells which stimulates CobB

deacetylase activity. The number of Prs-CobB complexes depends on transcriptional activity of the *prs* gene, which can be stimulated under conditions of low purine and pyrimidine nucleotides. Prs acetylation in the stationary phase is required for CobB stimulation. CobB in turn protects Prs form loss of activity under conditions of low magnesium or phosphate ion concentration.

**Table 1**. Acetylated lysines in Prs and their deacetylation by CobB. Chemically acetylated lysines K182, K194, K231 are deacetylated *in-vitro* by CobB deacetylase.

Samples of PrsAc and PrsAc deacetylated by CobB (Fig. 2D) were excised from 2 independent Coomassie stained SDS-Polyacrylamide gels and analyzed with Mass Spectrometry. All identified peptides carrying lysines of interest were summed up (second value in brackets) and acetylated lysines were counted (first value in brackets). The acetylation percentage was calculated and average values are shown in the table. A PrsAc protein sample aliquot was additionally analyzed with mass spectrometry where 108, 82 and 73 peptides with acetylated lysines K182, K194 and K231 respectively were identified suggesting those lysines were acetylated *in-vitro*.

| Lysine | Protein sequence | PrsAc | | PrsAc after CobB deacetylation | |
|---|---|---|---|---|---|
| K182 | VVRARAIA**K**LLNDTDMA | sample1 (6/8) | 75% | sample3 (2/34) | 6% |
| | | sample2 (16/19) | 84% | sample4 (5/54) | 9% |
| | | average | 80% | average | 8% |
| K194 | DTDMAIID**K**RRPRANVSQ | sample1 (11/69) | 16% | sample3 (3/57) | 5% |
| | | sample2 (27/251) | 11% | sample4 (0/86) | 0% |
| | | average | 14% | average | NA |
| K231 | IDTGGTLC**K**AAEALKERG | sample1 (2/24) | 8% | sample3 (0/19) | 0% |
| | | sample2 (0/34) | 0% | sample4 (0/19) | 0% |
| | | average | NA | average | NA |

938                                      Fig. 1

Fig. 2

983                                                   Fig. 3

1000

1001                                             Fig. 4

1002

1003

1004

1005

1006

1007

1008

1009

1010

1011

1012

1013

1014

1015

1016

1017

1018

1019

1020

1021

1022

1023

1024

1025

1026

1027

1028

1029

1030

1031

1032

Fig. 5

1033

1034

1035

1036

1037

1038

1039

1040

1041

1042

1043

1044

1045

1046

1047

1048

1049

1050                                                Fig. 6

1051

1052

1053

1054

1055

1056

1057

1058

1059

1060

1061

1062

1063

1064

1065

1066

1067

1068

1069

1070

1071

1072

1073

1074

1075

1076

1077

1078                                                      Fig. 7

1079

1080