Increased prefrontal top-down control in older adults predicts motor performance and age-group association

Philipp Alexander Loehrer[1,2*], Felix Sebastian Nettersheim[3], Carina Oehrn[1,2,4], Fabienne Homberg[1], Marc Tittgemeyer[5], Lars Timmermann[1,2,4], Immo Weber[2,4]

[1]*Department of Neurology, University Hospital Gießen and Marburg, Marburg, Germany*

[2]*Department of Neurology, Philipps-University Marburg, Marburg, Germany*

[3]*Department of Cardiology, University Hospital Cologne, Cologne, Germany*

[4]*Center for Mind, Brain and Behavior (CMBB), Philipps-University Marburg, Marburg, Germany*

[5]*Max Planck Institute for Metabolism Research, Cologne, Germany*

*Corresponding author:

Philipp Loehrer, Department of Neurology, University Hospital Marburg, Baldinger Str., 35043, Marburg, Tel: +49 6421 5866419, Email: philipp.loehrer@uk-gm.de

Abstract

Bimanual motor control declines during healthy ageing, affecting the ability of older adults to maintain independence. Age-related changes in neural dynamics of movement preparation that underlie reduced bimanual motor control, however, remain largely elusive. Here, we assessed effective connectivity within a prefrontal-premotor network of 19 young and 20 older adults during the preparatory phase of complex bimanual movements using high-density electroencephalography and Dynamic Causal Modelling of evoked responses. We found that excitatory interhemispheric prefrontal to premotor coupling in older adults predicted age-group association and poor motor-performance. In contrast, excitatory intrahemispheric prefrontal to premotor coupling enabled older adults to maintain motor-performance at the cost of lower movement speed. Our results disentangle the complex interplay in the prefrontal-premotor network underlying reduced bimanual control and the well-known speed-accuracy trade-off seen in older adults. We anticipate these coupling patterns to be potential biomarkers for healthy ageing and a possible future target of non-invasive brain-stimulation.

1. Introduction

Physiological ageing is associated with a decline in fine motor control[1]. This decline becomes particularly apparent during the performance of bimanual movements affecting activities of daily living and the ability to maintain independence[1,2]. Hence, in an ageing society, it is of great interest to understand the mechanisms of physiological ageing and associated deterioration of motor functions.

Deficits in motor performance in older adults (OA) comprise several aspects such as coordination problems, movement variability, as well as an overall slowing of movements[1]. Due to temporal and spatial coordination problems, OA particularly face difficulties performing bimanual tasks[3,4]. Furthermore, older adults are often overchallenged by complex tasks that require a high amount of flexibility[5]. In contrast, the ability of OA to perform learned and frequently practiced actions as well as automatized movements is not impaired[6].

An important underlying factor of age-related motor performance deficits is a loss of brain volume[1]. Several studies have linked reduced gray matter volume as well as decreased white matter integrity in OA to reduced behavioural performance (for a review see Seidler et al.[1]). Here, frontal areas, including the prefrontal cortex, seem to be particularly susceptible to atrophy[7]. Counterintuitively, a wealth of imaging studies have demonstrated an enhanced and more widespread blood-oxygen-level-dependent-(BOLD)-profile in older adults while performing unimanual[8,9] and bimanual motor tasks[10,11], which was interpreted as over-activation. This over-activation was particularly present in frontal[10,11,12] and parietal areas[9,10,11]. Over time, two theories have emerged to explain over-activation in the ageing brain. The de-differentiation hypothesis states that the increased activation pattern represents a non-selective and inefficient recruitment of brain areas[13]. In contrast, the more frequently invoked compensation hypothesis states that over-activation compensates for a reduction in the brain's structural integrity and is associated with better motor performance[8]. The described pattern of

over-activation in OA during movement is paralleled by distinct connectivity changes in the motor system[9,14,15,16]. Overall, an increase in connectivity metrics, such as connectivity strength and network efficiency, measured by functional magnetic resonance imaging (fMRI) and Dynamic Causal Modelling as well as graph-theoretical network analysis, was associated with healthy ageing[14,15]. On the other hand, distinct connections were associated with reduced connection strengths in older adults during the execution of unimanual finger tapping (left PFC to left premotor cortex)[9] and unilateral foot moving (right M1 to right premotor cortex)[16], supporting the notion that the influence of age depends on the connection studied.

The wealth of studies investigating age-related changes in movement execution is accompanied by a dearth of studies addressing the effects of healthy ageing on movement preparation. During the preparatory phase of a movement, young adults (YA) show a lateralized activation over the hemisphere contralateral to the moving hand[17]. This lateralization diminishes in OA[17], while the amplitude of distinct event-related potentials (lateralized readiness potential, P300, contingent negative variation) increases, i.e. becomes more positive, in OA[17,18]. Hinder and colleagues showed that interhemispheric inhibition (IHI) between premotor and motor areas is reduced (i.e. more facilitatory) during motor preparation of unimanual finger movements in OA[19], supporting the idea that increased amplitudes in OA arise from a reduction in IHI[20]. Furthermore, the pattern of prefrontal over-activation revealed during movement execution could be observed during movement preparation also. To reach accuracy levels comparable to those of YA, OA prepared simple and complex unimanual finger movements with higher prefrontal activation[21]. On the other hand, older adults were not able to modulate interhemispheric projections from dorsolateral prefrontal cortex to contralateral primary motor cortex during the preparation of a bimanual coordination task, which was associated with performance deficits and reduced white matter microstructure[22]. To date, no studies have addressed age-related effects on the causal interplay within the motor system during movement preparation that might be related to a decline in bimanual coordination and functional

independence. Furthermore, no literature exists with regard to the individual electrophysiological differences between older adults with preserved and reduced bimanual coordination. To address these issues, we assessed effective connectivity in young and older adults during the preparatory phase of bimanual movements. Participants performed temporally coupled and spatially uncoupled movements while a 128-channel electroencephalography (EEG) was recorded. We used Dynamic Causal Modelling for evoked responses (DCM for ERP) to infer effective connectivity between prefrontal cortex (PFC), lateral premotor cortex (lPM), and supplementary motor area (SMA), as these structures are highly involved in bimanual movements[23]. DCM for ERP is a statistical approach employing neuronal mass models to infer effective connectivity between distinct neuronal sources based on measured EEG responses[24,25]. In the present study, we employed the Parametric Empirical Bayes (PEB) framework to delineate differences between age groups and relate effective connectivity during movement preparation to behavioural parameters[26,27]. We hypothesized that (1) OA exhibit increased connection strengths compared to young adults, (2) increased connectivity strengths are related to differences in behavioural parameters, and (3) older adults with preserved bimanual coordination show connectivity patterns resembling those of young adults.

2. Results

2.1 Behavioral results

Two-tailed independent samples t-tests, comparing all older adults (n=20) with young adults (n=19), revealed that older adults made more mistakes compared to young adults, however, this difference was not significant ($t(37) = 1.278$, $p = .209$). Furthermore, older adults needed more time to perform the task compared to young adults ($t(37) = 3.452$, $p = .001$). To evaluate differences in top performing older adults (n=10) compared to low performers (n=10) and young adults (n=19), we used repeated measures ANOVA. For the variable *error rate*, an overall between-subject effect of "group" ($F(2,36) = 7.658$, $p = .002$) was revealed. Post hoc

two-tailed t-tests, Bonferroni-corrected, demonstrated that old – low performers made more mistakes compared to old – top performers (p = .002, old – low performer: 38.86±1.96, old – top performer: 17.8±1.28, mean±SEM) and compared to young adults (p = .009, young: 22.87±1.19). No difference could be observed between old – top performers and young adults (p = .883, Figure 1.A). Moreover, there was a trend toward a significant main effect of complexity (F(4, 142) = 2.38 , p = .055). No further main effects or interactions were revealed for *error rate* (all p > .108). Repeated measures ANOVA of *performance time* similarly revealed an overall between-subject effect of "group" (F(2,36) = 7.837 , p = .001). Post hoc two-tailed t-tests, Bonferroni-corrected, showed that old – low performers needed more time compared to young adults (p = .001; old – low performer: 3.34±0.07, young: 2.34±0.04, mean±SEM). No difference could be observed between old – top performers and young adults (p = .199; old top – performer: 2.83±0.08) as well as old – low performers and old – top performers (p = .257, see Figure 1.B). There were main effects of hand (F(1, 36) = 14.765, p < .001; subjects needed more time when the learned sequence was tapped with the right hand) and complexity (F(3.5, 126) = 11.212 , p < .001). Moreover, there was an interaction between the factors hand and complexity for *performance time* (F(4,146) = 2.892, p = .024). No further significant main effects or interactions were revealed for *performance time* (all p > .085).

Figure 1.A. Mean error rates and standard error of the mean (SEM) for young (n=19) and older adults (n=20). Old – low performers (n=10) made more mistakes compared to young adults (Bonferroni corrected two-sided t-test, p = .009) and older adults in the top-performing group (n=10; Bonferroni corrected two-sided t-test, p = .002). No

difference was observed between YA and Old – top-performer (Bonferroni corrected two-sided t-test, p = .883).

B. Mean performance time and SEM for young and older adults. Old – low-performers needed more time to complete the task compared to YA (Bonferroni corrected two-sided t-test, p < .001). No difference could be observed between old – top performers and YA (Bonferroni corrected two-sided t-test, p = .199). Asterisks indicate significant differences (** p ≤ .01, *** p ≤ .001).

2.2 Event Related Potentials - Sensor Space Analysis

In both conditions older adults (n=20) exhibited more positive ERPs than young adults (n=19). If the learned sequence was tapped with the right hand, statistical analysis of ERPs using a one-sided permutation test revealed a group-difference between 517 and 758 ms post-stimulus (p = .003). Furthermore, we found differences in ERPs between 793 and 1000 ms post-stimulus intermittently. Differences were located in the sensors above the right hemisphere over the postcentral sulcus and the angular gyrus (Figure 2.A and C). If the learned rule was executed with the left hand, however, differences were found in the left hemisphere, more anteriorly above the frontal gyrus in a latency range of 313 to 570 ms (p = .043, Figure 2.B and D). ERP-analysis of YA compared to OA stratified by performance yielded similar results (cf. Supplementary Figure 1 and 2). Here, differences between YA and OA – top performers (n=10) showed differences in the same latency ranges over the same areas, whereas differences between YA and OA – low performers (n=10) were non-significant for condition 2 and showed a trend toward significance for condition 1.

In summary, differences were located on the hemisphere contralateral to the hand tapping the new number sequence in both conditions.

Figure 2: Scalp topography of EEG sensors showing significant differences between grand-mean ERPs for young and older adults.

A. Scalp topography plotted for correctly performed trials at 688 ms after stimulus onset when the learned rule was performed with the right hand and the new number sequence with the left hand. The cluster with a p-value = .003 is denoted with two starlets (one-sided permutation test, YA: n=19, OA: n=20). The red line shows the time point for which the scalp topography is plotted. **B.** Scalp topography for correctly performed trials when the learned rule was performed with the left hand and the new number sequence with the right hand for 440 ms after stimulus onset. The cluster with a p-value of .043 is denoted with a cross (one-sided permutation test, YA: n=19, OA: n=20). The red line shows the time point for which the scalp topography is plotted. The dark green line denotes 300 ms post "go"-stimulus. **C.** ERP waveforms averaged time-locked to the "go"-stimulus: Grand-mean ERP of young (blue curve) and older (orange curve) adults recorded with electrode P4 (right hand rule). Significant differences are shaded in light blue color. **D.** ERP waveforms averaged time locked to the "go"-stimulus: Grand-mean ERP of young and older adults recorded with electrode FCC1H (left hand rule). Significant differences are shaded in light blue color.

2.3 Dynamic Causal Modelling – Source Space Analysis

Random effects Bayesian model comparison between three fully connected models with either forward, backward, or lateral connections between lPM and SMA favored a model featuring forward connections from SMA to lPM and backward connections from lPM to SMA (Model

Exceedance Probability: 0.70; see Supplementary Figure 3). Ensuing PEB models were based on this model structure. Here, we took intrinsic as well as extrinsic connectivity strengths to the group level and used Bayesian Model Reduction (BMR) to exclude connections that did not contribute to the model evidence. Commonalities between participants as well as differences due to age-group, performance-group, *error rate*, and *performance time* were determined for parameters of A- and B-Matrix separately.

2.3.1 A-Matrix Parameters – Commonalities

Bayesian model averages of the A-Matrix parameters (average effective connectivity across conditions) that were in common over subjects showed a consistent inhibitory connectivity profile between premotor and prefrontal areas (Figure 3.A-B). Here, forward connections from left lPM to left PFC and right lPM to right PFC were of inhibitory nature. Similarly, forward connections from SMA to left and right PFC were inhibitory. Spearman correlation analysis revealed that weaker inhibitory influence of lateral premotor areas on prefrontal Cortex positively correlated with higher *error rates* (left lPM to left PFC: YA: $r = .99$, $p < .001$, OA: $r = .98$, $p < .001$; right lPM to right PFC: YA: $r = .90$, $p = < .001$, OA: $r = -.21$, $p = .45$, Figure 3.C-D). Weaker inhibitory influence of SMA on PFC, on the other hand, was correlated with higher performance time in young adults (SMA to left PFC: $r = .57$, $p = .024$; SMA to right PFC: $r = .69$, $p = .004$, Figure 3.E-F). In older adults, this association was not significant (SMA to left PFC: $r = .03$, $p = .95$; SMA to right PFC: $r = .10$, $p = .78$). When differentiating between old top- and low-performers, however, this association reached statistical significance at the .05-level (SMA to left PFC: old top-performer: $r = .92$, $p = .001$, old low-performer: $r = .50$, $p = .22$; SMA to right PFC: old top-performer: $r = .98$, $p < .001$, old low-performer: $r = .84$, $p = .009$, Figure 3.E-F).

Figure 3.A. Bayesian model averages of the commonalities between groups thresholded at posterior probability >95% for clarity. The schematic in **B** illustrates the parameters that survived thresholding. Red arrows denote inhibitory connections. **C-D.** Error rates plotted against individual coupling parameters of lateral premotor (lPM) to prefrontal cortex (PFC) coupling (YA: young adults, n=19; OA: older adults, n=20). **E-F.** Performance time plotted against individual coupling parameters of supplementary motor area (SMA) to PFC coupling (OA Top: older adults – top performer (n=10), OA Low: older adults – low performer (n=10)). The schematic was visualized with BrainNet Viewer[28].

2.3.2 A-Matrix Parameters – Age-Group Differences

Bayesian model averages of parameters related to differences in age-group showed a consistent pattern of being more excitatory in older adults. Here, interhemispheric backward connections from left PFC to right lPM as well as from right PFC to left lPM were more excitatory in OA (Figure 4.A-B). Spearman correlation analysis revealed that a weaker inhibitory influence of left PFC on right lPM in YA and a stronger excitatory influence in OA correlated with higher *error rates* (left PFC to right lPM: YA: *r* = .85, *p* < .001, OA: *r* = .93, *p* < .001, Figure 4.D).

On the other hand, weaker inhibitory influence of right PFC on left lPM in young adults and a stronger excitatory influence in older adults correlated with higher *performance time* (right PFC to left lPM: YA: *r* = .79, *p* < .001, OA: *r* = .95, *p* < .001, Figure 4.F). A post-hoc ANOVA revealed that OA in the top-performing group had lower connection strengths compared to subjects in the low-performing group for both interhemispheric connections. Here, the connection strength from left PFC to right lPM was significantly lower (*p* < .001) and connection strength from right PFC to left lPM showed a trend toward significance for being lower in the top-performing group (*p* = .06). Thus, old top-performers had connectivity profiles more similar to young adults.

Figure 4.A. Bayesian model averages of the group differences thresholded at posterior probability >95% for clarity. The four parameters displayed are the only parameters that survived thresholding. The schematic in **B** illustrates the interhemispheric parameters that survived thresholding. Green arrows denote more excitatory connections in older adults compared to young adults. **C and E.** Box-plots for coupling parameters of connections left PFC to right lPM (**C**) and right PFC to left lPM (**E**, YA: young adults (n=19), OA: older adults (n=20)). Center line indicates the median, box limits represent upper and lower quartiles and whiskers indicate most extreme data points not considered outliers **D.** Error rates plotted against individual coupling parameters of left PFC to right

lPM coupling. **E.** Performance time plotted against individual coupling parameters of right PFC to left lPM. **G.** Mutual information quotients for coupling parameters that survived thresholding. Interhemispheric connections were selected to reduce dimensionality and increase classification accuracy. **H.** Results of leave-one-out cross validation. Actual subject group-parameter (YA = -1, OA = 1) plotted against the expected value of the estimated group-parameter.

Furthermore, backward connections from right PFC to right lPM and from right PFC to SMA were more excitatory in older adults. Spearman correlation analysis revealed that a stronger inhibitory influence of right PFC on right lPM in young adults was associated with higher *error rates* (right PFC to right lPM: young adults: $r = -.91$, $p <.001$, Figure 5.D). In older adults, this association was not significant (old adults: $r = -.35$, $p = .21$). When differentiating between old top- and low-performers, however, the association between weaker excitatory influence and higher error rates reached statistical significance at the .05-level (old top-performer: $r = -.84$, $p = .009$, old low-performer: $r = -.87$, $p = .006$, Figure 5.D). Additionally, a weaker inhibitory influence of right PFC on SMA in young adults and a stronger excitatory influence in older adults was related to higher *performance time* (young adults: $r = .87$, $p <.001$, older adults: $r = .89$, $p <.001$, Figure 5.F). Here, no differences in connection strengths between top- and low-performing older adults could be revealed by the post-hoc ANOVA.

Figure 5.A. Bayesian model averages of the group differences thresholded at posterior probability >95% for clarity. The four parameters displayed are the only parameters that survived thresholding. The schematic in **B** illustrates the intrahemispheric parameters that survived thresholding. Green arrows denote more excitatory connections in older adults compared to young adults. **C and E.** Box-plots for coupling parameters of connections right PFC to right lPM (**C**) and right PFC to SMA (**E**, YA: young adults (n=19), OA: older adults (n=20)). Center line indicates the median, box limits represent upper and lower quartiles and whiskers indicate most extreme data points not considered outliers. **D.** Error rates plotted against individual coupling parameters of right PFC to right lPM coupling (OA Top: older adults – top performer (n=10), OA Low: older adults – low performer (n=10)). **F.** Performance time plotted against individual coupling parameters of right PFC to SMA coupling (YA: young adults (n=19), OA: older adults (n=20)).

2.3.3 B-Matrix Parameters – Commonalities and Age Group Differences

To evaluate the influence of age on the differences between hands, we used the PEB framework to model the B-Matrix parameters. Here, Bayesian model averages of parameters related to commonalities between subjects showed that the backward connection from left PFC to SMA was decreased in condition 1 (learned sequence tapped with right hand) compared to condition

2 (learned sequence tapped with left hand). Furthermore, connections from right lPM to SMA and from left lPM to left PFC were increased, whereas the influence of left lPM on itself was reduced in condition 1 compared to condition 2. Spearman correlation analysis revealed that a larger difference in PFC to SMA connectivity strength between both conditions correlated with a larger difference in *performance time* in all subjects (left PFC to SMA: $r = .70$, $p < .001$). Weaker PFC to SMA connectivity in condition 1 was hence associated with an increased *performance time*. There were no other correlations between connectivity parameters related to commonalities and performance differences. Furthermore, there were no difference in the B-Matrix parameters due to age-group.

2.4 Mutual Information Quotient and Leave-One-Out Cross Validation

To assess the predictive validity of the A-matrix connectivity parameters, we employed leave-one-out cross validation. MIQ feature selection algorithm selected both interhemispheric connections (Figure 4.G). Thus, we used these connections for leave-one-out-analysis (LOO). Here, the estimated effect sizes of interhemispheric connections were large enough to predict age-group affiliation (Pearson's correlation coefficient: 0.53, $p < .001$, Figure 4.H).

3. Discussion

Using Dynamic Causal Modelling for evoked responses and Bayesian Model Averaging, we show for the first time that excitatory interhemispheric top-down control from prefrontal to lateral premotor cortex predicts age-group association and motor performance. Thus, this parameter could be used as a biomarker for healthy ageing and as a potential target for non-invasive interventions such as transcranial magnetic stimulation (TMS). Furthermore, we report that right-hemispheric top-down control from prefrontal cortex to premotor areas underlies the commonly observed finding of preserved accuracy at the cost of increased performance time in older adults. Our results disentangle the complex interplay between prefrontal and premotor

areas during movement preparation and describe distinct connectivity profiles that predict poor as well as preserved motor performance in older adults.

3.1 Preserved accuracy at the cost of increased performance time in older adults

It is a common finding that older adults can match the accuracy of young adults in simple tasks[14,29]. When task complexity rises, however, OA have to compensate for the increased task demands at the cost of execution time and eventually are unable to parallel the accuracy of YA[21,30]. Conforming to these findings, our results showed that OA achieved comparable *error rates*, whereas *performance time* was markedly increased. Although in line with the literature, this was a somewhat surprising finding, as we expected increased *error rates* for OA due to the substantial complexity of our paradigm. These results can be explained by the performance of a top-performing subgroup of OA. Within this group (N=10), OA paralleled and occasionally outperformed the accuracy of YA drawing the overall group mean of *error rates* towards the mean of YA. The main effect of hand and the interaction between complexity and hand for the factor *performance time* suggest that, regardless of age, subjects had difficulty to tap a new number sequence with the non-dominant left hand.

3.2 Increased and spatially extended activity in older adults

Statistical analysis of ERP data consistently demonstrated enhanced positive potentials in older adults compared to young adults. In both experimental conditions, OA showed these enhanced positive potentials in distributed clusters located mainly above the hemisphere contralateral to the hand tapping a new sequence. This finding is in concordance with a study by Wu and Hallett who found that OA showed an increased BOLD-signal, interpreted as enhanced cortical activity, and activated additional sources while performing complex automatic movements at the same accuracy as YA[6]. Since the present analysis included only correct trials, our findings support the hypothesis that older adults rely on the recruitment of spatially extended brain areas to match the performance of YA during complex movements.

Analysis of the stimulus-locked ERP components showed no differences between both groups during early phases of the stimulus analysis, whereas pronounced differences could be revealed for the later phase. This is in accordance with previous ERP studies showing that age-related differences particularly emerge during later cognitive components, i.e. the P300, over central and parietal regions[21,29]. In the present study, YA showed a marked positive deflection at 450 ms post "go"-stimulus, which can be interpreted as P300 as the deflection lies within the typical P300 latency range (250-500 ms)[31]. In contrast, a sustained positive deflection was observed in OA. The P300 is an ERP component elicited when attention has to be drawn to integrate sensory information with working memory[32]. Thus, the P300 seems to reflect the processing of a stimulus as well as the overall level of attention[31]. The amplitude of the P300 has been associated with memory processes but it is also thought to represent the amount of attentional resources engaged[31]. Although evidence exists that P300 amplitude declines with age when elicited by visual or auditory stimuli[33], Sterr and Dean showed increased P300 amplitudes in OA during unimanual movement preparation[17]. Our results support the notion, that amplitudes of later cognitive components of the ERP such as the P300 are increased in OA compared to YA during preparation of movements. We speculate that this increase in amplitude along with the sustained positivity represents the higher cognitive demand OA have to engage to match the performance of YA. This is also supported by our supplementary analysis were we compared YA with low and high performing OA and could replicate our findings. In conclusion, our results extend the findings of Wu and Hallett as well as Sterr and Dean in that a broader activation profile as well as an increased amplitude of cognitive ERP components is also present during the preparation of bimanual movements.

3.3 Absence of M1 activity and hierarchy in premotor areas

In the present study, we found no evoked source activity in M1 in either of the two conditions or groups (cf. 5.8 Source Specification). Although subsets of neurons in the primary motor area are active during movement preparation[34], there is evidence for gating mechanisms which

inhibit M1 neurons during movement preparation to prevent premature action[35]. It is therefore likely, that we were not able to detect this complex interplay of inhibition and facilitation within M1 when averaging ERPs time-locked to the "go"-stimulus. By choosing to average time-locked to the "go"-stimulus, we focused on the cognitive and preparatory part of the movement and not execution per se. Furthermore, the electrophysiological activity time-locked to the movement itself is very likely to vary inter-individually in relation to the "go"-stimulus and therefore averages out over several trials.

Random effects Bayesian model comparison revealed evidence for a top-down influence from lPM to SMA during preparation of complex bimanual movements. Numerous studies have shown the importance of both supplementary motor area[23,36] and premotor Cortex[36,37] for the coordination of bimanual movements. Clear evidence for a hierarchy between SMA and PM, however, could not be detected in anatomical studies[38]. On the other hand, studies investigating the task-related involvement of both areas showed that SMA and PM are of importance for different subsets of bimanual movements. Here, SMA seems to be responsible for the temporal sequencing of movements and the parallel coding of actions of multiple effectors[23]. By contrast, the premotor cortex is engaged when movements depend on external stimuli and becomes particularly active when task complexity rises[37]. Since our paradigm involved complex bimanual movements an increased involvement of PM seems to be likely. In conjunction with the studies named above, our results support the notion that the bimanual coordination network is not a rigid entity. Rather, it is a dynamic system in which top-down control can change when adapting to task demands[23].

3.4 Inhibitory feedback from premotor to prefrontal areas might represent conveyance of efference copies

Bayesian model averages of the A-matrix parameters common across all subjects consistently showed inhibitory bottom-up connectivity from premotor to prefrontal areas. It is widely

acknowledged that the prediction of a movement-outcome relies on so-called forward models, which are simulations of the motor system about the consequences of an action[39,40]. The forward model uses a copy of the motor command, the efference copy, which is thought to be generated in the supplementary motor area as well as the premotor cortex[41,42]. We speculate that the bottom-up feedback from premotor areas could represent the conveyance of efference copies providing information about the expected consequences of movements. Since the PFC is engaged in online movement monitoring and online error correction[43,44], the inhibition of PFC could represent a reduced need for top-down error correction when the motor system expects a correct execution (only correct trials were used for analysis). This notion is supported by the findings that reduced inhibition of PFC from lPM was associated with higher error rates, whereby reduced inhibition of PFC from SMA was associated with higher performance time. This interpretation requires further investigation.

3.5 The distinct role of the PFC in increased intra- and interhemispheric coupling in older adults

There is a dominant view that the prefrontal cortex plays an eminent role in the cognitive control of action[45]. It has been suggested that activity in the dorsolateral prefrontal cortex reflects "attention to action" and that specific subregions within the PFC play a crucial role in online movement monitoring and working memory[43,44,46]. Furthermore, functions, ascribed to the PFC, seem to show a hemispheric predominance. Left PFC is supposed to be implicated in task setting and switching, whereas right PFC seems to be particularly involved in online movement monitoring and memory retrieval[44]. In the present study, high levels of attention were required to perform the complex bimanual movements correctly. In fact, in two previous studies from our group as well as another study from Meister et al. it was demonstrated that precise prefrontal to premotor coupling is important for the execution of complex bimanual movements and that distinct coupling changes are associated with healthy ageing and Parkinson's disease[2,47,48]. The

influence of prefrontal to premotor coupling during movement planning, however, has not been investigated hitherto and we will discuss these interactions in the subsequent sections.

3.5.1 Increased interhemispheric coupling in older adults

Converging evidence from TMS studies suggests that interhemispheric inhibition (IHI) between both primary motor cortices as well as premotor and motor areas is reduced (i.e. more facilitatory) in OA compared to YA[19,49,50]. Little evidence, however, exists regarding the influence of age on interhemispheric prefrontal to premotor connectivity. In this study, Bayesian Model Averaging (BMA) revealed that interhemispheric connections from PFC to contralateral lPM were excitatory in OA whereas they were inhibitory in YA. This increased excitatory coupling conforms to fMRI studies which have shown over-activation of prefrontal areas in OA during the performance of unimanual and bimanual movements, which was associated with better motor performance[10,11,12]. The translation of this overactivation into altered functional connectivity and motor performance, however, is poorly understood. We argue that the increased prefrontal-premotor-coupling is an expression of prefrontal overactivity and show for the first time that this coupling pattern is already present during movement preparation. The importance of interhemispheric prefrontal to premotor top-down control during movement preparation is underscored by the finding that leave-one-out cross validation predicted the group association based on the two interhemispheric connections. Therefore, prefrontal to premotor interhemispheric interactions might represent a valid biomarker for healthy ageing and future TMS studies. Furthermore, the correlation analysis of our study revealed a complex association between increased coupling patterns and motor performance. Here, increased excitatory interhemispheric PFC to lPM connectivity was associated with poorer motor performance, arguing for de-differentiation in the prefrontal-premotor system. On the other hand, right-sided intrahemispheric connectivity was associated with better motor performance, giving evidence for a compensatory mechanism within this

system. Interestingly, older adults in the top-performing group, who matched the performance of young adults, showed excitatory left PFC to right lPM coupling that was lower than the coupling of older adults in the low-performing group. Therefore, we argue that weak excitatory top-down influence from prefrontal to premotor areas can serve as a compensatory mechanism but impairs bimanual movements as excitatory top-down control increases. In summary, we show, for the first time, that excitatory top-down control from prefrontal to lateral premotor areas is present in older adults during movement preparation. Influence of these interhemispheric top-down connections predicted age-group whereby connection strength predicted motor outcome.

3.5.2 Increased intrahemispheric coupling in older adults

The age-related prefrontal over-activation described in imaging studies was also detected during unimanual movement preparation in an electrophysiological analysis. Employing a reaction time task Berchicci and colleagues could demonstrate increased prefrontal activation in OA during a simple as well as a complex condition which was related to increased movement accuracy[21]. The increased right-sided prefrontal to premotor coupling in OA in the present study can be interpreted as elevated prefrontal activation but further delineates how prefrontal activity influences motor output. Importantly, the described coupling was associated with reduced *error rates* but increased *performance time*. Evidence exists that rising complexity of bimanual movements intensifies activity of premotor cortex and that this increased activity, particularly of the right premotor cortex, is important for bimanual control[37,51]. Our results show that the increased recruitment of the lateral premotor cortex in OA is mediated via top-down control of the right PFC. This increased recruitment was associated with reduced errors whereas increased activation of SMA was correlated with increased performance time. We suppose that the excitatory top-down influences from right PFC to SMA and lPM are an important underlying

factor for the frequently described phenomenon that OA can match the performance of YA at the cost of reduced movement speed.

3.6 Differences between the performance of both hands

Using Bayesian model averages of B-matrix parameters we could investigate the neural dynamics underlying the performance difference between both hands, which was revealed in our behavioural analysis. Here, in common across subjects, left PFC to SMA coupling was weaker in condition 1 compared to condition 2 which was associated with increased *performance time*. This association suggests that performance differences between hands, i.e. longer *performance time* for trials in which the new sequences had to be tapped with the non-dominant left hand, is at least partly due to a reduced left PFC to SMA coupling. It is noteworthy, that left prefrontal cortex is particularly important for task switching. In this regard, several studies could link left PFC activity to tasks in which subjects had to constantly update and switch the task set[52,53]. In the present study, subjects had to tap a new sequence in every trial, requiring participants to update the task continuously. This seemed to pose a difficulty for young and older adults when the new sequence was tapped with the non-dominant left hand. The associated reduction in left PFC to SMA connectivity may represent reduced left prefrontal control and seems to be an important underlying factor for performance differences between hands.

3.7 Limitations

The present study has several limitations. First, we pursued a region of interest approach and included predefined cortical sources based on their activity in previous studies. Although it is likely that additional cortical and subcortical sources such as the basal ganglia, the cerebellum, and the anterior cingulate cortex play a role in bimanual movement preparation, we did not include these areas to reduce data dimensionality and to account for the limited capability of electroencephalography to model subcortical sources. Second, Bayesian model selection

chooses the model that describes the data best within a given model space[54]. This entails that models with an architecture not incorporated in our model space might be equally or more probable. Third, the cross-sectional design of our study enabled us to investigate age-related differences in effective connectivity. To investigate age-related and subject specific changes in effective connectivity, longitudinal studies are necessary.

4. Conclusion

In the present study, we investigated the influence of healthy ageing on the neural dynamics within the prefrontal-premotor network during the preparation of a complex bimanual task. We could show that distinct connectivity profiles were common across age-groups, whereas others differed between young and older adults. Hence, several conclusions can be drawn from our results. First, OA seem to rely on an increased recruitment of spatially extended areas and engage more attentional resources as represented by more positive amplitudes of cognitive ERP-components to match the performance of YA. Second, correct and rapid execution of bimanual movements appears to require inhibitory bottom-up connections from prefrontal to premotor areas, which might represent the conveyance of efference copies. Third, interhemispheric inhibition from prefrontal to premotor areas was reduced in OA, whereby weak excitatory interhemispheric connectivity might represent a compensatory mechanism in top-performing OA but eventually leads to poorer performance as excitatory influences increase. Furthermore, interhemispheric prefrontal to premotor connections might represent a biomarker for healthy ageing as they predicted age-group association. Fourth, increased intrahemispheric right PFC to premotor coupling in OA was associated with increased performance time but reduced error rates, suggesting an important involvement in the frequently seen phenomenon that OA are able to match the performance of YA but require more time to perform the movement. Fifth, participants had difficulties when performing a new sequence with the non-dominant left hand, which was associated with reduced left PFC to SMA

coupling. Hence, reduced left PFC influence might underlie performance differences between right and left hand when constant task monitoring and updating is required.

5. Materials and Methods

5.1 Participants

28 young and 32 old healthy volunteers participated in this study upon written informed consent. Following EEG-preprocessing, the datasets of 19 young (mean age ± SD: 25.79 ± 4.29 years; 5 female) and 20 old (mean age ± SD: 59.65 ± 6.82 years; 8 female) subjects were included for further analysis (see "Data Acquisition" and "EEG Preprocessing"). All participants were right-handed according to the Edinburgh Handedness Inventory, did not play an instrument for more than five hours per month, and had normal neuropsychological test scores (Mini-Mental State Examination, cut-off $\leq$ 25; DemTect, cut-off $\leq$ 13, and Beck's Depression Inventory, cut-off $\geq$ 13). The experimental procedures were conducted in accordance with the Declaration of Helsinki and approved by the local ethics committee (study number: 13-394). The same data set has been previously published in another study[2].

5.2 Experimental Paradigm

Experimental procedures were performed in an electrically and acoustically shielded room. Participants were seated comfortably in front of a computer screen with their fingers placed on a response pad (Cedrus, San Pedro, USA). The response pad consisted of eight buttons, four buttons for each hand, whereby a number as well as a corresponding button were allocated to each finger (1 for left and right thumb; 2 for left and right index finger; 3 for left and right middle finger and 4 for left and right ring finger). A comprehensive description of the experimental conditions and the behavioural paradigm is found in Loehrer and Nettersheim et al.[2]. In short, participants were instructed to learn a sequence of four button presses for one hand (e.g. left hand 1, 2, 3, 4). The learned sequence had to be executed with the respective

hand while, simultaneously, the other hand had to tap a different sequence that was presented on the screen (see Figure 1.A). Participants were instructed to focus on correct trial execution rather than speed. Learned sequences were performed with the right hand (condition 1) and left hand (condition 2), whereas the starting hand was counter balanced across subjects. The behavioral task comprised six different tapping sequences of varying complexity which had to be learned and each of which had to be executed 20 times for both hands. At the beginning of each trial, the upcoming sequence of numbers was shown in red and signaled the subject to prepare for movement initiation. The switch from red numbers to green numbers indicated to start tapping ("go"-signal). A red cross marked the end of one trial, followed by a short break of 5 seconds. Trials in which subjects pressed the learned as well as the new sequences in the correct order were included for further analysis only. In addition to the condition analyzed here, participants completed a task in which they had to tap the same sequences with both hands.

Figure 6. Instructions presented on a screen (left) and demanded button presses (right) in chronological order. In this example, participants had to learn a sequence for their right hand (condition 1) and tap a new sequence with their left hand. Hence, the first demanded pair of buttons was right thumb and left index finger.

5.3 Analysis of Behavioural Data

Error rate denotes the ratio of correct trials to overall trials. *Performance time* denotes the time between the go-signal and the last button press. Statistical analysis was performed using SPSS 22.0 (IBM, Armonk, USA) for Windows 10. The Shapiro-Wilk test revealed a non-Gaussian

distribution for values of *error rate* and *performance time*, which prompted us to employ square root transformed data for further analysis. We used independent samples t-tests (two-tailed) to investigate overall differences between young (n=19) and older adults (n=20) in *error rate* and *performance time*. To evaluate differences in top performing old subjects compared to low performers and young adults, we conducted a median split for the variable *error rate*. *Error rate* and *performance time* were entered separately into a mixed-design analysis of variance (ANOVA) with the repeated measures "complexity" (levels 1-6) and "hand" (learned sequence performed with left or right hand) as well as the independent variable "group" (Young (n=19), Old – top performer (n=10), Old – low performer (n=10)) using Greenhouse-Geisser correction for non-sphericity when appropriate. Post-hoc Bonferroni-corrected t-tests were conducted comparing the three groups.

5.4 Data Acquisition

EEG-data were acquired using a 128-channel system (Brain Products GmbH, Gilching, Germany) in an electrically shielded room. EEG-caps were placed upon subjects head and correct positioning was ensured by measuring the electrodes distances to nasion, inion, as well as left and right preauricular fiducials. After assuring that electrode impedances were below 10 kΩ, EEG-signals were recorded with Brainvision recorder (Brain Products GmbH, Gilching, Germany). Signals were amplified, band-pass filtered from 0.1 to 1000 Hz, and digitized at a sampling rate of 5000 Hz. Prior to EEG recordings, individual T1-weighted magnetic resonance images (MRI) of the head were acquired on a 3-Tesla Trio scanner (Siemens, Erlangen, Germany) using a 3D Modified Driven Equilibrium Fourier Transform sequence (repetition-time = 1930 ms, echo-time = 5.8 ms, flip-angle = 18°, slice-thickness = 1.25 mm). Three young and one older adult discontinued MRI acquisition and subsequent EEG recordings due to claustrophobia.

5.5 EEG Preprocessing

Data pre-processing was performed with BESA Research 6.0 ("brain electrical source analysis", BESA, Gräfelfing, Germany). During offline analysis, the data was average-referenced and artefact-corrected. The amplitude threshold for excluding a channel from the analysis was below 0.07 μV, above 120 μV, or a gradient larger than 75 μV to adjacent channels. Noisy trials were excluded from further analysis. Eyeblink removal was performed with the automatic artefact removal tool provided by BESA (threshold for horizontal and vertical eye movements: 150 μV and 250 μV) or, if the method failed to remove all eyeblinks, with an additional independent component analysis (ICA). For calculating event-related activity, data was filtered (Butterworth filter: 0.53 – 200 Hz, notch filter: 50 Hz), downsampled to 1000 Hz, and averaged time-locked to the presentation of the green numbers ("go"-stimulus) for correctly performed trials. Subjects with too many noisy channels (> 60), too little correct trials (< 10), or without clear evoked activity were excluded from the analysis (6 young subjects, 11 old subjects). For DCM analysis, event related potential data was transformed to SPM data format (SPM12, update revision number 7487, Wellcome Trust Centre for Neuroimaging, London, UK), bandpass filtered between 1 and 48 Hz and downsampled to 250 Hz.

5.6 Statistical Analysis of Event-Related Potentials

Statistical analysis of event related potentials (ERP) was based on permutation testing to avoid alpha-error accumulation and was performed with BESA statistics 1.0. As a first step, for every sample of the ERPs a two-sided unpaired t-Test was calculated and significant differences between groups were defined (threshold: $\alpha = .05$). Clusters were specified by selecting connected data points on the basis of temporal and spatial adjacency[55]. Subsequently, for each cluster, a cluster value was derived consisting of the sum of all t-values in the cluster. Then, the data of subjects were permutated, i.e. systematically interchanged. For each of the permutations a new t-test was computed and a new cluster-value derived. Thereby, a new distribution of

cluster-values was obtained for each of the initial clusters. Based on the new cluster values resulting from the permuted data, it was possible to determine the significance of the initial cluster-value. If less than 5 % of the new cluster values were larger than the initial cluster, the initial cluster was considered significant with a p-value of .05.

5.7 Dynamic Causal Modelling

Dynamic causal modelling is a statistical approach to model the underlying neuronal architecture that gives rise to measured hemodynamic or electromagnetic time series data[25]. Employing neural mass models and Bayesian statistics, DCM allows one to infer the effective connectivity between brain regions and how these parameters are influenced by experimental factors[25]. Here, we used DCM for evoked responses to model the neuronal dynamics underlying movement preparation of a bimanual task in young and old subjects.

DCM for evoked responses rests on neural mass models with three subpopulations of neurons, as described by Jansen and Rit, to model source activity[24,56]. Within this model, each subpopulation is assigned to one of three cortical layers: inhibitory interneurons in the supra-granular layer, excitatory neurons (spiny stellate cells) in the granular layer, and deep pyramidal cells in the infra-granular layer. Within a source, the three different layers are connected via excitatory and inhibitory interneurons. Connectivity between sources is established via pyramidal cells in the infra-granular layer. Here, bottom-up processing (forward connections) is represented by connections terminating in the granular layer, whereas top-down processing (backward connections) is mediated via connections terminating in the infra- and supra-granular layers. Lateral connections originate in the infra-granular layer also and terminate in all layers[24]. Experimental perturbations of the connectivity within a network is implemented via an exogenous input u, which has the same characteristics as a forward connection, hence terminating in the granular layer[24]. This architecture is based on the hierarchical organization

described in Fellemann and Van Essen and allows one to build models with multiple interconnected sources[24,38].

The response of each subpopulation to an input is characterized by a set of differential equations. These equations describe the average response of a subpopulation in terms of firing rate-dependent current and voltage changes[24,57]. The change in neuronal state of a source ($\dot{x}$) represents the average depolarization of pyramidal cells and can be described as a function of input (u) and its previous state (x):

$$\dot{x} = f(x, u, \theta) \tag{1}$$

where θ represents the parameters and modulations of forward, backward, and lateral connections. These parameters are also called "hidden neuronal states" and have to be estimated. An electromagnetic forward model is used to estimate the change of neuronal state of a source (x_0) based on a measured ERP signal (y):

$$y = L(\theta)x_0 + \varepsilon \tag{2}$$

where $L(\theta)$ represents the parameterized lead field and ε observation noise (equations (1) and (2) are adopted from Garrido et al.[57,58]). The combination of neural mass models with a spatiotemporal forward model results in a generative model that describes how model parameters and perturbations give rise to the measured ERP signal[25].

Bayesian inversion of this generative model minimizes the Variational Free Energy under the Laplace Approximation[58,59]. The employed procedure can be regarded as an Expectation-Maximization algorithm and provides an estimate of the posterior mean values of the model parameters θ and the log-model evidence. This model evidence is composed of the accuracy of a model, describing the data fit, and a complexity term, which penalizes models with increasing numbers of parameters[58,60]. The model evidence can then be used to compare different models[54].

5.8 Source Specification

In the framework of DCM, the specification of sources can be made using prior knowledge of the relevant sources based on previous studies[61]. In this study, activity of cortical sources included in subsequent dynamic causal models was confirmed in a source reconstruction of ERPs using a multiple sparse priors algorithm[62]. First, we defined a core motor network based on the MNI coordinates reported by Haslinger et al. and Herz et al. which were used in previous studies[2,47,63,64]. Sources included in this network were dorsolateral prefrontal Cortex (PFC; MNI coordinates: left: [-37 48 22], right: [37 48 22]), lateral premotor Cortex (lPM; MNI coordinates: left: [-52 14 42], right: [52 14 42]), supplementary motor area (SMA; MNI coordinates: [0 -5 70]), and primary motor cortex (M1; MNI coordinates: left: [-41 -26 66], right: [41 -26 66]). Due to the spatial adjacency of bilateral SMA and the poor spatial resolution of EEG, we included SMA as a single source.

To evaluate whether these sources were indeed active during bimanual task performance, individual MRIs were coregistered with EEG electrode locations and a BEM (boundary element model) head model, employing a cortical mesh with 8196 dipoles, was calculated. Source activity for correct trials was inverted for both groups at 0-758 ms post-stimulus using the multiple sparse priors algorithm as implemented in SPM[62]. Length of this time interval was defined by the significant difference in ERP magnitude between young and old subjects (see "Results"). T-statistics were used to contrast inversion results to a 100 ms prestimulus baseline using family wise error (FWE) correction thresholded at <.05 at the peak and cluster level. To account for the low spatial resolution of EEG, we allowed for a deviation of 10 mm from our ROIs[64]. We could not find consistent activity in M1, which prompted us to exclude this source from our model space (cf. 4.3 Absence of M1 activity and hierarchy in premotor areas). Activity was confirmed in our other ROIs during the respective time interval (see Supplementary Figure 4).

5.9 Model Specification

Based on evidence from anatomical studies, we assumed reciprocal connections between lPM and SMA[38] as well as reciprocal connections between PFC and lPM and PFC and SMA[38]. This connectivity structure was applied to the interhemispheric connectivity between prefrontal and premotor areas also. With respect to the employed hierarchy, i.e. the nature of reciprocal connections, we refer to a study of Felleman and Van Essen[38] that suggests ascending (i.e. forward) connections going from lPM to PFC, and descending (i.e. backward) connections from PFC to lPM. Regarding lPM and SMA, there is no evidence for hierarchical differences. Therefore, three fully connected models with either forward, backward, or lateral connections between lPM and SMA were inverted per subject. Model evidence was compared using a random effects Bayesian model comparison (see "Results")[65]. In addition, lateral connections were used for interconnecting left and right PFC as well as lPM of both hemispheres. The input to the network was set to the PFC of both hemispheres, assuming prefrontal areas to drive the motor output as described in Herz et al.[66].

5.10 Implementation of DCM in this Study

Dynamic causal modelling for evoked potentials was used to infer effective connectivity between five sources in 39 subjects in this study. A forward model was calculated for each subject as stated above (see "Source Specification"). ERPs at 0-758 ms post-stimulus were fitted to a fully connected model with backward connections from lPM to SMA and forward connections from SMA to lPM. This interval was necessary to capture the significant differences between young and old subjects in ERPs in both conditions (see "Results"). Activity of cortical sources was modeled using the equivalent current dipole (ECD) method as implemented in SPM[58]. To infer the effect of age on the average connectivity over conditions and on the difference between hands, we contrasted both conditions (condition 1: learned sequence performed with the right hand; condition 2 learned sequence performed with the left

hand). Reduction of data dimensionality was achieved by singular value decomposition and the first eight principal components were used for inversion, which corresponds to the SPM default settings (see Supplementary Figure 5 for the frequency modes of a representative patient). A bandwidth from 4 to 48 Hz was used to avoid a 50 Hz electric current artifact.

5.11 Parametric Empirical Bayes and Bayesian Model Reduction

Parametric Empirical Bayes (PEB) describes the inversion of hierarchical models in which the posterior density of model parameters are constrained by the posterior density of the hierarchy level above[26]. The priors, i.e. the constraints, used for this inversion are derived from empirical data – contrary to standard Bayesian methods, for which the prior distribution is fixed – and are therefore called empirical priors[26]. This procedure involves inversion of a parent model (e.g. a fully connected model) for every subject at the first level. The parameters of interest (e.g. connectivity parameters) are then taken to the second level (i.e. group level). Here, a Bayesian General Linear Model is specified by collating the mean and the covariance matrices of the parameters of interest as well as a design matrix containing experimental effects called regressors (e.g. group affiliation, age, handedness)[27]. By integrating the covariance matrix into the GLM, modelling the parameters of interest at the group level takes into account the individual differences of each parameter. This approach, therefore, differentiates between hypothesized group-level effects and uninteresting random effects[27].

In an ensuing Bayesian Model Comparison, PEB models with combinations of different parameters of interest can be compared to identify the model best explaining the differences and commonalities due to the entered regressors. Here, large numbers of models can be compared using Bayesian Model Reduction. BMR derives the parameters and model evidence of nested PEB models by systematically removing one parameter or subsets of parameters, i.e. setting the prior mean of the respective parameter(s) to zero[26]. If one assumes that all nested models are equally probable a priori, BMR can be used to perform an automated search over

the model space and prune away parameters that do not contribute to the model evidence (an approach that we pursued here)[27].

In the present study, the coupling parameters were assessed within the PEB framework to characterize the commonalities between subjects and the differences due to age-group (YA: n=19, OA n=20), performance-group (top-performer (n=10)/low-performer (n=10), error rate, and performance time. In a first step, the A-matrix parameters, which represent the average effective connectivity across experimental conditions, were entered into a PEB model with the mean-centered regressors mentioned above. BMR was used to perform an exhaustive comparison of different PEB models as described above. Subsequently, mean connectivity parameter estimates were derived by averaging the parameters across models weighed by the models' posterior probability, a process called Bayesian Model Averaging[67]. BMA was thresholded at 95 % posterior probability to focus the results on parameters with strong evidence of being non-zero[27]. This procedure was repeated for the B-matrix parameters, which - in our study - represent the effective connectivity necessary to model the difference between both conditions. It is noteworthy that the PEB model for the B-matrix parameters comprised age-group (YA: n=19, OA n=20), performance-group (top-performer (n=10)/low-performer (n=10), as well as the difference between both conditions in *error rate* and *performance time* as mean-centered regressors. PEB models with additional regressors (such as age or handedness) were not superior to the selected PEB model as revealed by free energy comparison. To disentangle the interaction between coupling strength, age-group, and performance parameters, coupling parameters were correlated with performance parameters partitioned by age-group using spearman-correlation. We controlled the false discovery rate using the Benjamini-Hochberg method to account for multiple comparisons and report corrected p-values[68]. In a last step, we assessed the predictive validity of estimated connectivity parameters using leave-one-out cross validation[27]. First, we defined connections that were most relevant for age-group association by

employing the mutual information quotient (MIQ) as described in Weber[69]. Subsequently, we sought to ascertain whether the effect size of the selected connectivity parameters was large enough to predict a subject's group affiliation.

6. Funding

PAL was supported by the SUCCESS-Program of the Philipps-University of Marburg and the 'Stiftung zur Förderung junger Neurowissenschaftler'.

7. Acknowledgement

The authors would like to thank the participants for their active engagement in this study.

8. Author contributions

Study design: PAL, FSN, LT, and IW; Conceptualization: PAL and IW; Methodology: PAL FSN, FJ, CO, and IW; Formal analysis: PAL, FSN, FJ and IW; Investigation: PAL, FSN, FJ, CO, MT, and IW; Resources: MT and LT; Writing - original draft: PAL; Writing - review & editing, PAL, FSN, CO, LT and IW; Supervision: LT

9. Data availability

The data that support the findings of this study are available upon reasonable request from the corresponding author (PAL). MRI data are not publicly available since they contain information that could compromise research participant privacy/consent. A source data file is provided for Figures 1.A, 1.B, 3, 4, 5, and Supplementary Figure 3.

10. Code availability

We used the commercial software BESA research 6.0 (BESA, Gräfelfing, Germany) for preprocessing of EEG data and BESA statistics 1.0 (BESA, Gräfelfing, Germany) for statistical

analysis of ERP data. Analysis of behavioural data was performed using the commercial software SPSS 22.0 (IBM, Armonk, USA). Auxiliary scripts and code used for further analysis was written and developed in MATLAB (The MathWorks, Inc., MA, USA; commercial software), predominantly on version 2016b and 2018a, and are available at https://github.com/Loehrerph/BiSyT. For DCM analysis we used SPM12 (update revision number 7487, Wellcome Trust Centre for Neuroimaging, London, UK) freely available at https://www.fil.ion.ucl.ac.uk/spm/software/spm12/. The mutual information quotient feature selection algorithm is freely available within the NoLiTiA-toolbox at https://www.nolitia.com/download.html. Furthermore, the MATLAB plugin BrainNet Viewer, freely available at https://www.nitrc.org/projects/bnv/, was used to visualize the schematics in Figures 3 to 5.

11. References

1. Seidler RD*, et al.* Motor control and aging: links to age-related brain structural, functional, and biochemical effects. *Neuroscience and biobehavioral reviews* **34**, 721-733 (2010).

2. Loehrer PA*, et al.* Ageing changes effective connectivity of motor networks during bimanual finger coordination. *NeuroImage* **143**, 325-342 (2016).

3. Stelmach GE, Amrhein PC, Goggin NL. Age differences in bimanual coordination. *Journal of gerontology* **43**, P18-23 (1988).

4. Wishart LR, Lee TD, Murdoch JE, Hodges NJ. Effects of aging on automatic and effortful processes in bimanual coordination. *J Gerontol B Psychol Sci Soc Sci* **55**, P85-94 (2000).

5. Glisky EL. Frontiers in Neuroscience Changes in Cognitive Function in Human Aging. In: *Brain Aging: Models, Methods, and Mechanisms* (ed^(eds Riddle DR). CRC Press/Taylor & Francis Copyright © 2007, Taylor & Francis Group, LLC. (2007).

6. Wu T, Hallett M. The influence of normal human ageing on automatic movements. *The Journal of physiology* **562**, 605-615 (2005).

7.   Raz N*, et al.* Selective aging of the human cerebral cortex observed in vivo: differential vulnerability of the prefrontal gray matter. *Cerebral cortex (New York, NY : 1991)* **7**, 268-282 (1997).

8.   Mattay VS*, et al.* Neurophysiological correlates of age-related changes in human motor function. *Neurology* **58**, 630-635 (2002).

9.   Rowe JB*, et al.* Aging is associated with contrasting changes in local and distant cortical connectivity in the human motor system. *NeuroImage* **32**, 747-760 (2006).

10.   Coxon JP, Goble DJ, Van Impe A, De Vos J, Wenderoth N, Swinnen SP. Reduced Basal Ganglia Function When Elderly Switch between Coordinated Movement Patterns. *Cerebral Cortex* **20**, 2368-2379 (2010).

11.   Goble DJ, Coxon JP, Van Impe A, De Vos J, Wenderoth N, Swinnen SP. The neural control of bimanual movements in the elderly: Brain regions exhibiting age-related increases in activity, frequency-induced neural modulation, and task-specific compensatory recruitment. *Human brain mapping* **31**, 1281-1295 (2010).

12.   Heuninckx S, Wenderoth N, Swinnen SP. Systems Neuroplasticity in the Aging Brain: Recruiting Additional Neural Resources for Successful Motor Performance in Elderly Persons. *The Journal of Neuroscience* **28**, 91 (2008).

13.   Li S-C, Lindenberger U. Cross-level unification: A computational exploration of the link between deterioration of neurotransmitter systems and dedifferentiation of cognitive abilities in old age. In: *Cognitive neuroscience of memory* (ed^(eds). Hogrefe & Huber (1999).

14.   Heitger MH*, et al.* Bimanual motor coordination in older adults is associated with increased functional brain connectivity--a graph-theoretical analysis. *PloS one* **8**, e62133 (2013).

15.   Michely J*, et al.* Network connectivity of motor control in the ageing brain. *NeuroImage Clinical* **18**, 443-455 (2018).

16.   Spedden ME*, et al.* Directed connectivity between primary and premotor areas underlying ankle force control in young and older adults. *NeuroImage* **218**, 116982 (2020).

17.   Sterr A, Dean P. Neural correlates of movement preparation in healthy ageing. *The European journal of neuroscience* **27**, 254-260 (2008).

18.   Roggeveen AB, Prime DJ, Ward LM. Lateralized readiness potentials reveal motor slowing in the aging brain. *The Journals of Gerontology Series B: Psychological Sciences and Social Sciences* **62**, P78-P84 (2007).

19.   Hinder MR, Fujiyama H, Summers JJ. Premotor-motor interhemispheric inhibition is released during movement initiation in older but not young adults. *PloS one* **7**, e52573 (2012).

20. Levin O, Fujiyama H, Boisgontier MP, Swinnen SP, Summers JJ. Aging and motor inhibition: A converging perspective provided by brain stimulation and imaging approaches. *Neuroscience & Biobehavioral Reviews* **43**, 100-117 (2014).

21. Berchicci M, Lucci G, Pesce C, Spinelli D, Di Russo F. Prefrontal hyperactivity in older people during motor planning. *NeuroImage* **62**, 1750-1760 (2012).

22. Fujiyama H*, et al.* Age-Related Changes in Frontal Network Structural and Functional Connectivity in Relation to Bimanual Movement Control. *The Journal of neuroscience : the official journal of the Society for Neuroscience* **36**, 1808-1822 (2016).

23. Swinnen SP, Wenderoth N. Two hands, one brain: cognitive neuroscience of bimanual skill. *Trends in cognitive sciences* **8**, 18-25 (2004).

24. David O, Harrison L, Friston KJ. Modelling event-related responses in the brain. *NeuroImage* **25**, 756-770 (2005).

25. Friston KJ, Harrison L, Penny W. Dynamic causal modelling. *NeuroImage* **19**, 1273-1302 (2003).

26. Friston KJ*, et al.* Bayesian model reduction and empirical Bayes for group (DCM) studies. *NeuroImage* **128**, 413-431 (2016).

27. Zeidman P*, et al.* A guide to group effective connectivity analysis, part 2: Second level analysis with PEB. *NeuroImage* **200**, 12-25 (2019).

28. Xia M, Wang J, He Y. BrainNet Viewer: a network visualization tool for human brain connectomics. *PloS one* **8**, e68910 (2013).

29. Kolev V, Falkenstein M, Yordanova J. Motor-response generation as a source of aging-related behavioural slowing in choice-reaction tasks. *Neurobiology of Aging* **27**, 1719-1730 (2006).

30. Seidler-Dobrin RD, He J, Stelmach GE. Coactivation to reduce variability in the elderly. *Motor control* **2**, 314-330 (1998).

31. Polich J. Updating P300: an integrative theory of P3a and P3b. *Clin Neurophysiol* **118**, 2128-2148 (2007).

32. Rossini PM, Rossi S, Babiloni C, Polich J. Clinical neurophysiology of aging brain: From normal aging to neurodegeneration. *Progress in Neurobiology* **83**, 375-400 (2007).

33. Polich J*, et al.* P300 topography of amplitude/latency correlations. *Brain topography* **9**, 275-282 (1997).

34. Tanji J, Evarts EV. Anticipatory activity of motor cortex neurons in relation to direction of an intended movement. *Journal of Neurophysiology* **39**, 1062-1068 (1976).

35. Soteropoulos DS. Corticospinal gating during action preparation and movement in the primate motor cortex. *J Neurophysiol* **119**, 1538-1555 (2018).

36. Tanji J, Okano K, Sato KC. Neuronal activity in cortical motor areas related to ipsilateral, contralateral, and bilateral digit movements of the monkey. *J Neurophysiol* **60**, 325-343 (1988).

37. Debaere F, Wenderoth N, Sunaert S, Van Hecke P, Swinnen SP. Cerebellar and premotor function in bimanual coordination: parametric neural responses to spatiotemporal complexity and cycling frequency. *NeuroImage* **21**, 1416-1427 (2004).

38. Felleman DJ, Van Essen DC. Distributed hierarchical processing in the primate cerebral cortex. *Cerebral cortex (New York, NY : 1991)* **1**, 1-47 (1991).

39. Wolpert DM, Flanagan JR. Motor prediction. *Current Biology* **11**, R729-R732 (2001).

40. Wolpert DM, Ghahramani Z, Jordan MI. An internal model for sensorimotor integration. *Science (New York, NY)* **269**, 1880-1882 (1995).

41. Ellaway PH, Prochazka A, Chan M, Gauthier MJ. The sense of movement elicited by transcranial magnetic stimulation in humans is due to sensory feedback. *The Journal of physiology* **556**, 651-660 (2004).

42. Haggard P, Whitford B. Supplementary motor area provides an efferent signal for sensory suppression. *Brain research Cognitive brain research* **19**, 52-58 (2004).

43. Jueptner M, Stephan KM, Frith CD, Brooks DJ, Frackowiak RSJ, Passingham RE. Anatomy of Motor Learning. I. Frontal Cortex and Attention to Action. *Journal of Neurophysiology* **77**, 1313-1324 (1997).

44. Stuss DT, Alexander MP. Is there a dysexecutive syndrome? *Philosophical Transactions of the Royal Society B: Biological Sciences* **362**, 901-915 (2007).

45. Cieslik EC, Zilles K, Grefkes C, Eickhoff SB. Dynamic interactions in the fronto-parietal network during a manual stimulus–response compatibility task. *NeuroImage* **58**, 860-869 (2011).

46. Rowe J, Friston K, Frackowiak R, Passingham R. Attention to action: specific modulation of corticocortical interactions in humans. *NeuroImage* **17**, 988-998 (2002).

47. Nettersheim FS*, et al.* Dopamine substitution alters effective connectivity of cortical prefrontal, premotor, and motor regions during complex bimanual finger movements in Parkinson's disease. *NeuroImage* **190**, 118-132 (2019).

48. Meister IG, Foltys H, Gallea C, Hallett M. How the brain handles temporally uncoupled bimanual movements. *Cerebral cortex (New York, NY : 1991)* **20**, 2996-3004 (2010).

49. Gooijers J, Swinnen SP. Interactions between brain structure and behavior: The corpus callosum and bimanual coordination. *Neuroscience & Biobehavioral Reviews* **43**, 1-19 (2014).

50. Talelli P, Waddingham W, Ewas A, Rothwell JC, Ward NS. The effect of age on task-related modulation of interhemispheric balance. *Experimental Brain Research* **186**, 59-66 (2008).

51. Liuzzi G, Hörniss V, Zimerman M, Gerloff C, Hummel FC. Coordination of uncoupled bimanual movements by strictly timed interhemispheric connectivity. *The Journal of neuroscience : the official journal of the Society for Neuroscience* **31**, 9111-9117 (2011).

52. Derrfuss J, Brass M, Neumann J, von Cramon DY. Involvement of the inferior frontal junction in cognitive control: Meta-analyses of switching and Stroop studies. *Human brain mapping* **25**, 22-34 (2005).

53. Kim C, Cilles SE, Johnson NF, Gold BT. Domain general and domain preferential brain regions associated with different types of task switching: A Meta-Analysis. *Human brain mapping* **33**, 130-142 (2012).

54. Penny WD, Stephan KE, Mechelli A, Friston KJ. Comparing dynamic causal models. *NeuroImage* **22**, 1157-1172 (2004).

55. Maris E, Oostenveld R. Nonparametric statistical testing of EEG- and MEG-data. *Journal of neuroscience methods* **164**, 177-190 (2007).

56. Jansen BH, Rit VG. Electroencephalogram and visual evoked potential generation in a mathematical model of coupled cortical columns. *Biological cybernetics* **73**, 357-366 (1995).

57. Garrido MI, Kilner JM, Kiebel SJ, Stephan KE, Friston KJ. Dynamic causal modelling of evoked potentials: a reproducibility study. *NeuroImage* **36**, 571-580 (2007).

58. Kiebel SJ, David O, Friston KJ. Dynamic causal modelling of evoked responses in EEG/MEG with lead field parameterization. *NeuroImage* **30**, 1273-1284 (2006).

59. Friston K, Mattout J, Trujillo-Barreto N, Ashburner J, Penny W. Variational free energy and the Laplace approximation. *NeuroImage* **34**, 220-234 (2007).

60. Friston KJ, Penny W, Phillips C, Kiebel S, Hinton G, Ashburner J. Classical and Bayesian inference in neuroimaging: theory. *NeuroImage* **16**, 465-483 (2002).

61.   Stephan KE, Penny WD, Moran RJ, den Ouden HE, Daunizeau J, Friston KJ. Ten simple rules for dynamic causal modeling. *NeuroImage* **49**, 3099-3109 (2010).

62.   Friston K*, et al.* Multiple sparse priors for the M/EEG inverse problem. *NeuroImage* **39**, 1104-1120 (2008).

63.   Haslinger B*, et al.* Event-related functional magnetic resonance imaging in Parkinson's disease before and after levodopa. *Brain* **124**, 558-570 (2001).

64.   Herz DM*, et al.* Task-specific modulation of effective connectivity during two simple unimanual motor tasks: a 122-channel EEG study. *NeuroImage* **59**, 3187-3193 (2012).

65.   Penny WD*, et al.* Comparing Families of Dynamic Causal Models. *PLOS Computational Biology* **6**, e1000709 (2010).

66.   Herz DM, Siebner HR, Hulme OJ, Florin E, Christensen MS, Timmermann L. Levodopa reinstates connectivity from prefrontal to premotor cortex during externally paced movement in Parkinson's disease. *Neuroimage* **90**, 15-23 (2014).

67.   Hoeting JA, Madigan D, Raftery AE, Volinsky CT. Bayesian model averaging: a tutorial. *Statistical science*, 382-401 (1999).

68.   Benjamini Y, Hochberg Y. Controlling the false discovery rate: a practical and powerful approach to multiple testing. *Journal of the Royal statistical society: series B (Methodological)* **57**, 289-300 (1995).

69.   Weber I. NoLiTiA: An Open-Source Toolbox for Nonlinear Time Series Analysis (Version 1.1) [Software]. Available from http://www.nolitia.com/download.html . (2018).