Effect of host genetics on the gut microbiome in 7,738 participants of the Dutch Microbiome Project

Authors

E.A. Lopera-Maya[1,§], A. Kurilshikov[1,§], A. van der Graaf[1,§], S. Hu[1,2,§], S. Andreu-Sánchez[1,3], L. Chen[1,3], A. Vich Vila[1,2], R. Gacesa[1,2], T. Sinha[2], V. Collij[1,2], M.A.Y. Klaassen[1,2], L.A. Bolte[1,2], M.F. Brandao Gois[1], P.B.T. Neerincx[1,4], M.A. Swertz[1,4], LifeLines Cohort Study*, H.J.M. Harmsen[5], C. Wijmenga[1], J. Fu[1,3], R.K. Weersma[2], A. Zhernakova[1,#], S. Sanna[1,6,#]

[§] shared first authors: EALM, AK, AvdG, SH

[#] shared last authors: AZ, SS

Correspondence to:
Dr. A. Zhernakova: sasha.zhernakova@gmail.com
Dr. S. Sanna, email: serena.sanna@irgb.cnr.it

Author affiliations

[1] University of Groningen and University Medical Center Groningen, Department of Genetics, Groningen, the Netherlands

[2] University of Groningen and University Medical Center Groningen, Department of Gastroenterology and Hepatology, Groningen, the Netherlands

[3] Department of Pediatrics, University of Groningen and University Medical Center Groningen, Groningen, the Netherlands

[4] Genomics Coordination Center, University of Groningen, University Medical Center Groningen, Groningen, the Netherlands

[5] Department of Medical Microbiology and Infection prevention: University of Groningen and University Medical Center Groningen, Groningen, The Netherlands

[6] Institute for Genetic and Biomedical Research (IRGB), National Research Council (CNR), Cagliari, Italy

* A list of authors and their affiliations appears at the end of the paper

Abstract

Host genetics are known to influence the gut microbiome, yet their role remains poorly understood. To robustly characterize these effects, we performed a genome-wide association study of 207 taxa and 205 pathways representing microbial composition and function within the Dutch Microbiome Project, a population cohort of 7,738 individuals from the northern Netherlands. Two robust, study-wide significant ($p<1.89\times10^{-10}$) signals near the *LCT* and *ABO* genes were found to be associated with multiple microbial taxa and pathways and were replicated in two independent cohorts. The *LCT* locus associations seemed modulated by lactose intake, while those at *ABO* could be explained by participant secretor status determined by their *FUT2* genotype. Twenty-two other loci showed suggestive evidence ($p<5\times10^{-8}$) of association with microbial taxa and pathways. At a more lenient threshold, the number of loci we identified strongly correlated with trait heritability, suggesting that much larger sample sizes are needed to elucidate the remaining effects of host genetics on the gut microbiome.

Introduction

The human intestinal microbial community contains trillions of microorganisms that play an important role in maintaining normal gut function and immune homeostasis[1]. Emerging evidence has shown that alterations in gut microbial composition are associated with the pathogenesis of many human diseases, including gastrointestinal disorders, metabolic syndrome, cardiovascular diseases and other conditions[2,3].

The gut microbiome is influenced by many factors, such as environmental factors including diet and medication usage[4,5], but also by host genetics. Heritability studies in twins and families have highlighted that human genetics contributes to gut microbial variation, showing heritability estimates ranging from 1.9% to 8.1%[6,7]. This observation drove the first efforts to identify genomic loci that influence gut microbiota through genome-wide association studies (GWAS). These early gut microbiome GWAS identified several microbial quantitative trait loci (mbQTL) located in genes related to the intestinal mucosal barrier, immune response and drug and food metabolism[8–11]. However, the reproducibility of these findings has been limited by differences in data processing methodologies, modest sample sizes and strong environmental effects, which, taken together, limit the detection of robust host genetic associations[12]. A recent large-scale genome-wide meta-analysis of 24 cohorts replicated the association between *Bifidobacterium* abundance and the lactase (*LCT*) gene locus[13], which had previously only been reported in single-cohort studies[6,14]. Other suggestive mbQTLs identified in this broad meta-analysis were proportional to heritability estimates from independent twin studies, indicating that additional loci found at lenient levels of significance are likely to be real but larger sample sizes are needed to reach sufficient statistical power[13]. Nonetheless, meta-analyses of mbQTL studies still lack resolution due to the high levels of heterogeneity between cohorts, which can reduce power in meta-analysis. On top of this, many existing cohorts rely on 16S rRNA measurements, which do not allow for bacterial identification at species-level resolution or for identification of bacterial pathway abundances. Indeed, measuring both species- and pathway-level abundances is essential for a further understanding of the function of an individual's microbiome. Two closely related bacterial species can have very different functions in the gut microbiome, yet pathways may be shared across distant microbial species and have roughly the same biological effect[15]. mbQTL studies using shotgun metagenomic sequencing in large population cohorts are therefore needed to overcome the high variability in microbiome definition and reveal more robust and refined associations.

To enable a broader and deeper understanding of host☐microbiota interactions, we used
shotgun metagenomic sequencing on feces from 7,738 individuals of the Dutch Microbiome
Project (DMP)[16] and matched their imputed genotypes to differences in taxa and pathway
abundances. By comparing our results with summary statistics from other independent
studies[13,17,18], we identify novel host☐microbiota interactions. Furthermore, we explore the
impact of potential confounding factors in modulating these genetic effects and identify potential
diet-dependent host☐microbiota interactions. We further assess the potential causal
relationships between the gut microbiome and dietary habits, biomarkers and disease using
Mendelian randomization. Finally, we carry out a power analysis that shows how microbiome
studies, even at the current sample size, are not well powered to reveal the complex genetic
architecture by which host genetics can regulate the gut microbiome.
90
91

Results

Genome-wide association identifies known and novel associations with several bacterial taxonomies and bacterial pathways

We investigated 5.5 million common (minor allele frequency (MAF) > 0.05) genetic variants on all autosomes and the X chromosome, using linear mixed models[19] to test their association and following an additive genetic model, with 207 taxa and 205 bacterial pathways in 7,738 individuals from the DMP cohort (**Methods, Supplementary Table 1**)[19]. Our quality control steps did not detect evidence for test statistic inflation (median genomic lambda 1.002 (range 0.75–1.03) for taxa and 1.004 (range 0.87–1.04) for pathways). We identified 37 SNP☐trait associations at 24 independent loci at a genome-wide p-value threshold of $5x10^{-8}$ (**Figure 1, Supplementary Table 2**). Genetic variants at two loci passed the more stringent study-wide threshold of $1.89x10^{-10}$ that accounts for the number of independent tests performed (**Methods**). The other 22 loci were associated at p-values between $5x10^{-8}$ and $1.89x10^{-10}$, with four associated to taxa and 18 to pathways. We refer to these 22 loci as "suggestive" and discuss them below.

The strongest signal was seen for rs182549 located in an intron of *MCM6*, a perfect proxy of rs4988235 ($r^2 = 1$, 1000 Genome European populations), one of the variants known to regulate the *LCT* gene and responsible for lactase persistence in adults (ClinVar accession RCV000008124). The T allele of rs182549, which confers lactase persistence through a dominant model of inheritance, was found to be associated with decreased abundances of the species *Bifidobacterium adolescentis* ($p = 7.6x10^{-14}$) and *Bifidobacterium longum* ($p = 3.2x10^{-08}$), as well as decreased abundances of higher-level taxa (phyla Actinobacteria $p = 2.3x10^{-13}$; class Actinobacteria, $p = 2.3x10^{-13}$; order Bifidobacteriales, $p = 5.9x10^{-15}$; family Bifidobacteriaceae, $p = 5.86x10^{-15}$; genus *Bifidobacterium, p = 8.59x10^{-15}$). Associations at this locus were also seen for other taxa of the same genus, but at lower levels of significance (*Bifidobacterium catenulatum, p = 3.9x10^{-5}$), and for species of the *Collinsella* genus (**Supplementary Figure 1**). The genetic association at the *LCT* locus has been previously described, albeit only at the genus level, in Dutch, UK and US cohorts[6,8,14], as well as in a recent large-scale meta-analysis[13]. The presence of the *LCT* association across several taxa suggests that this locus has a wide-ranging effect on microbiome composition.

The second locus that passed study-wide significance consists of genetic variants near the *ABO* gene. *ABO* encodes the BGAT protein, a histo-blood group ABO system transferase.

Associations found at this locus include species *B. bifidum* (rs8176645, *p* = 5.5x10^{-15}) and
Collinsella aerofaciens (rs550057, *p* = 2.0x10^{-8}, *r*^{2} = 0.59 with rs8176645 in 1000 Genomes
Europeans) and higher order taxa (rs550057, genus *Collinsella, p* = 9.3x10^{-11}; family
Coriobacteriaceae, *p* = 3.01x10^{-9}; order Coriobacteriales, *p* = 3.03x10^{-9}) (**Supplementary**
Figure 1). Interestingly, the metabolic pathway representing the bacterial degradation of lactose
and galactose was also associated to the *ABO* locus (Metacyc ID: "*LACTOSECAT-PWY:*
lactose and galactose degradation I", rs507666, *p* = 5.38x10^{-15}). Associations of this locus with
the genus *Collinsella* and the metabolic pathway *LACTOSECAT-PWY* have been recently
described[18,20]. To our knowledge, the associations with *B. bifidum* and *C. aerofaciens* are novel.
134

Association at *LCT* reflects the lactase persistence inheritance model and affects
multiple taxonomic levels and bacterial pathways
Given that lactose tolerance is inherited in a dominant fashion, we tested the associations found
in this locus using a dominant model for the alternative allele at SNP rs182549 and thereby
compared lactase persistent (LP) to lactose intolerant (LI) individuals. Indeed, all seven taxa
associated to the *LCT* locus at genome-wide significance showed a stronger association signal
when we used a dominant model (all associations *p* < 2x10^{-27}), with decreased taxa abundance
in LP individuals (**Supplementary Table 3**). The associations seen at the family level could
mostly be accounted for by species *B. adolescentis* (no significant difference in effect size,
Cochran's Q p values>0.05), while lower effects were seen for species *B. longum* and *B.*
bifidum (Cochran's Q p values when comparing effect sizes with those observed for *B.*
adolescentis: 0.018 and 0.003). Moreover, the association with these species remained
unchanged when adding *B. adolescentis* to the association models, indicating that the
associations are independent and not driven by species correlation.

We further tested the other 200 taxa for this SNP and the dominant model. Intriguingly, we also
observed suggestive association (*p* < 1x10^{-4}) with rs182549 for taxa that were associated to
ABO locus in our GWAS analysis (*Collinsella* genus and species *B. bifidum* and *C. aerofaciens)*
and for the species *Roseburia inulinivorans* of the family Lachnospiraceae (**Supplementary**
Table 3). For all but *Roseburia inulinivorans,* there was a consistent direction of effect across
the associated taxa, with increased abundance in LI compared to LP individuals (**Figure 2** and
Supplementary Figure 2).

Finally, when comparing the abundance of bacterial pathways between LI and LP groups, we
also observed a higher abundance of the *LACTOSECAT-PWY* in LI individuals (effect = +0.300

in standard deviation (SD) units, standard error (SE) = 0.49, p = 1.02x10^{-9}). This is not

surprising given that in our dataset this pathway correlates mostly with class Actinobacteria and

species *B. adolescentis (*Spearman correlations r_S: 0.73 and 0.69, respectively)*,* which are both

associated with SNPs at the *LCT* locus.

Associations at *ABO* are dependent on secretor status

To further understand the mechanisms underlying the association signals at the *ABO* locus, we

derived blood-group types based on the genotype status of three genetic variants (see

Methods). The majority of the individuals were either type A (40%) or type O (48%), reflecting

estimates from previous studies[21]. Genetic associations at this locus could be explained by

differences between non-O blood type individuals and O blood type individuals. We observed

higher abundances of *Collinsella* and the metabolic pathway *LACTOSECAT-PWY* in blood type

A or AB individuals compared to individuals with blood type O (Wilcoxon test for *Collinsella*,

blood type O vs. A: p = 2.8x10^{-9}, O vs. AB: p = 0.11; Wilcoxon test for metabolic pathway, O vs.

A: p = 5x10^{-14}, O vs. AB: p = 0.038). In contrast, blood type O individuals had the highest

observed abundance of *B. bifidum* compared to other blood type groups (Wilcoxon test blood

type, A vs. O: p = 2.3x10^{-14}, B vs. O: p = 0.007, AB vs. O: p = 0.006) (**Figure 3**).

The histo-blood group ABO system is determined by two genes: *FUT1*, which encodes a

fucosyltransferase that assemblies the H antigen, and *ABO*, which encodes a

glycosyltransferase that assemblies A or B antigens by adding a different terminal sugar to the

H antigen precursor. Therefore, the presence of A, B or H antigens on the surface of presenting

cells determines the blood type (A, B, or O, respectively) of an individual[22]. The gene *FUT2*,

encodes for a soluble form of the H antigen, that allows these antigens to be expressed on

intestinal mucosa and in bodily fluids[23]. However, individuals carrying two copies of the

alternative allele A at the enzyme-inactivating nonsense mutation rs601338 (OMIM 182100)

located in the *FUT2* gene are unable to express blood antigens on the intestinal mucosa (non-

secretors)[24]. We observed that the associations of taxa and bacterial pathways with blood types

were present in secretors and absent in non-secretors (**Figure 3**). This indicates that the

associations at the *ABO* locus are likely to be driven by the exposure of A/B antigens to gut

bacteria. These results recapitulate recent evidence for an mbQTL association at the *ABO* locus

depending on *FUT2* genotypes with the *Collinsella* genus and with another bacteria reported in

a Finnish population cohort[18]. Associations of a functional variant in the *FUT2* gene to other

bacterial taxa were observed in a recent meta-analysis, but none of these bacterial taxa or
pathways showed significant association for this locus in our cohort[13].

These novel associations with species levels at *ABO* are intriguing. To our knowledge, this is
the first time *B. bifidum* has been associated with predicted blood type. Early genomic analyses
suggest that, among *Bifidobacterium*, *B. bifidum* is particularly adapted to the human
gastrointestinal mucosa due to a unique set of genes encoding for enzymes involved in the
degradation and utilization of mucin, the main component of gastrointestinal mucosa[25,26]. The
degradation activity of the mucosa, where antigens are secreted, could explain the association
with differences in *B. bifidum* abundance at *ABO* modulated by *FUT2*. This mechanism has
been proposed for the association with *Collinsella* genus[18], and it is likely to also apply to the
novel associations we identified with *C. areofaciens* and LACTOSECAT-PWY pathway
abundances.
203
Genetic associations may be modulated by diet
Gut microbiome composition and function is known to be affected by several factors including
sex, body-mass index (BMI), diet and medication usage[4]. None of our genome-wide significant
associations were attenuated when including BMI, medication usage, stool frequency or stool
consistency as covariates (**Methods**, all comparisons, Cochran's Q for difference in effect size *p*
> 0.05), indicating that the genetic associations are independent and not confounded by these
factors (**Supplementary Table 4a**). Furthermore, we did not find any sex-specific associations
for the 37 SNP☐trait associations, although five associations did exhibit a smaller genetic effect
in females compared to males (**Supplementary Table 4b**).
We also investigated the effect of diet at the *LCT* and *ABO* loci, taking into account the
dominant inheritance model at *LCT* and the observed dependence on secretor status at *ABO*.
Based on 21 dietary factors derived from a food questionnaire recorded 4 years prior
microbiome collection, we considered dietary factors previously associated (FDR < 0.05) with
microbial taxa and pathway abundances that show genome-wide association signals in the *ABO*
and *LCT* loci (see **Methods**)[16]. In an analysis that included age, sex, genetic and dietary factors,
the dietary factors did not significantly attenuate the effect of the genetic components,
suggesting that diet is not a source of bias in these genetic associations (**Supplementary Table
5**). Nonetheless, diet remained an important factor after correction for genetic factors. Four taxa
and one pathway associated to *LCT* and *ABO* SNPs were statistically associated with at least
one dietary factor (*p* < 0.05) (**Supplementary Table 5**), with a maximum of 16 factors found for
B. longum. We further tested these associated dietary (44 diet☐microbiome pairs) factors for

interaction with genetics and detected evidence for a gene☐diet interaction for only one bacteria

at the *LCT* locus. Specifically, we observed an increased abundance of the Bifidobacteriaceae

family in LI individuals who consumed larger amounts of lactose or dairy (interaction term *p* =

0.03) (**Figure 3, Supplementary Table 6**), a finding that is consistent with previous reports[8,18].

In contrast, there was no evidence for interaction with diet at the *ABO* locus for the *B. bifidum*

species, the *Collinsella* genus and the bacterial pathway *LACTOSECAT-PWY* (interaction term

p > 0.05) (**Supplementary Table 6)**. This could be attributable to the limited accuracy of our

diet scores, information that was recorded 4 years prior to microbiome collection. Interaction

between fiber intake and genetic variants at this locus, when it is associated with *Collinsella*

genus, have been reported in other populations[18].

Association signals reveal a polygenic architecture of microbiome taxa and pathways

and replicate in other independent cohorts

Apart from the study-wide significant associations at the *LCT* and *ABO* gene regions, 22 other

loci showed suggestive association at $p < 5 \times 10^{-8}$ (**Supplementary Table 2**), none of which were

reported previously. The majority (18) were associated with bacterial pathways that could not

have been directly quantified in studies using 16S rRNA data, the methodology predominantly

used in microbiome genetic studies to date. These associations included several genomic loci

located in genes associated with human immune or metabolic disorders and mainly affected

microbial amino acids (5), vitamin (5), nucleotide (4), TCA cycle (2), carbohydrate (1) and

enterobactin toxin (1) pathways. These SNP☐pathway associations provide insights into

potential relationships between host phenotypes and gut microbial metabolic functions. For

instance, SNP rs9927590 in 16q23.1 (in the *WWOX* gene), which has been reported to be

involved in stenosis of Crohn's disease[27,28], was associated with a metabolic pathway involved

in short chain fatty acid (SCFA) production (*PWY-5088*, $p = 9.2 \times 10^{-9}$). SCFAs have been shown

to boost intestinal barrier functions, reduce local inflammation and decrease the risk of immune

and metabolic diseases, including type 1 and type 2 diabetes[17,29]. At another locus, SNP

rs11242123 (5q31.1) was associated with a decreased abundance of nucleotide biosynthesis

(*PWY-7209*, $p = 4.89 \times 10^{-8}$) and was previously reported to affect expression of the nearby genes

IL4 and *KIF3A*, two genes involved in asthma onset[30,31].

Interestingly, two suggestive genome-wide associations point towards a role in Vitamin B9

biosynthesis. A variant (rs2714053) in 11q24.1 (in the *GRAMD1B* gene) was associated with

lower abundance of bacterial genes involved in methyl diphosphate biosynthesis (*PWY-6147*, p

$= 4.58 \times 10^{-8}$), and this gene was previously linked to ulcerative colitis in a Polish cohort[32].

Notably, methyl diphosphate is a precursor of folate (Vitamin B9) that is required for the survival
of regulatory T cells in the human intestine[33,34]. In addition, another SNP, rs17824268 (14q23.2)
was found to be associated with decreased abundance of folate (PWY-6612, p = 2.63×10^{-8}).
This SNP was reported to be associated with human blood HDL, LDL and a variety of
metabolites[35,36]. Therefore, these two SNP-pathway associations point to a potential
dysregulation of the host immune and metabolic system through a decrease of folate production
from the gut microbiota.

In other loci, associations also point to immune and metabolic function. For example, two SNPs,
rs12137024 (1p21.3, close to gene ALG14, p = 3.53x10-8) and rs78778914 (1q21.2, close to
gene *RPRD2*, p = 4.34×10^{-8}), were associated with the L-histidine processing pathway. L-
histidine is usually catabolized by microbiota into glutamate, which is considered as an
"immunonutrient" serving as major fuel for immune cells including lymphocytes and
macrophages [37,38]. Intriguingly, the gene *ALG14* has been related to pneumonia [39], while
RPRD2 has been associated with immune diseases like asthma. Furthermore, another
example is the association of SNP rs1584586 (3q25.1), near the type 2 diabetes associated
gene *TSC22D2*[40], with the purine processing pathway (*PWY-841*, *p* = 3.47×10^{-8}) and of SNP
rs59657730 (6p12.3) within metabolic health related gene *RCAN2*, with microbial glycogen
degradation (GLYCOCAT.PWY, p = 3.50x10-8) [39]support a role for host–microbiome interaction
in regulating metabolic function. Four additional suggestive associations for taxa were identified,
including signals in 8p22 (in the *SGCZ* gene related to diabetes)[41] and in Xp21.1 (in the *DMD*
gene), although little is known about the function of the associated species.

We sought to replicate our genome-wide significant signals using summary statistics from other
independent cohorts in which microbiome data was characterized using either 16S rRNA (the
MiBioGen study)[13] or metagenomic sequencing (the LL-DEEP cohort)[17]. In the MiBioGen study,
a meta-analysis of 24 cohorts comprising up to 18,340 individuals, the 16S rRNA
measurements do not allow for the evaluation of the abundance of bacterial species and
pathways and the X chromosome was not analyzed. Consequently, only 10 of our 18 SNP taxa
pairs could be tested in MiBioGen and no pathways were testable. In LL-DEEP, a genome-wide
microbiome association study on 952 individuals, we extracted information for the majority of the
SNP taxa pairs (14/18) and SNP-pathway associations (18/19). Unfortunately, the power to
replicate the associations in LL-DEEP was limited due to the small sample size. In both studies,
we observed significant replication of the study-wide significant loci, *LCT* and *ABO,* using a
conservative Bonferroni threshold of p < 0.0015 (equivalent to 32 SNP-trait pairs tested). All

seven taxa associated with SNPs near *LCT* were replicated with consistent allelic effect
directions (all *p* < 3.7x10^{-6}). For the *ABO* locus, we found significant replication for the
Collinsella genus (*p* < 2x10^{-5} in MiBioGen) and replication at only nominal significance for the
metabolic pathway *LACTOSECAT-PWY* and *B. bifidum* species (*p* < 0.05 in LL-DEEP)
(**Supplementary Table 7** and **Supplementary Table 8**). The association at *ABO* with *B.*
bifidum has not been reported previously and it does not reach the multiple-testing adjusted
threshold for replication. Therefore, while the consistent direction of effects is encouraging, we
cannot exclude the possibility for this signal to be a false positive. None of the other SNP☐taxa
or pathway pairs were replicated in MiBioGen or LL-DEEP. Interestingly, another independent
SNP in the *COL23A1* gene (rs11958296; r^{2} = 0.1 with rs10447306 from our study) shows
association in MiBioGen to the abundance of the same taxa: family Rikenellaceae (*p* = 2.4x10^{-5})
and genus *Alistipes* (*p* = 9.3x10^{-6}).

To explore if the association signals at lower levels of significance are enriched in heritable
bacteria, which would indicate if it were possible to detect more genome-wide significant
mbQTLs by further increasing the sample size, we investigated the correlation of taxa and
pathway heritability estimations from family-based analysis with the number of suggestively
associated loci for each taxon and pathway. Here we observed a positive and significant
correlation for both taxonomic (r_S = 0.248, *p* = 3.1x10^{-4}) and pathway (r_S = 0.263, *p* = 1.3x10^{-4})
heritability with the number of suggestive (*p* < 1x10^{-5}) loci identified in our GWAS (**Figure 4**).
The correlation for pathways remained significant when increasing the mbQTL threshold to *p* <
5x10^{-4} (r_S = 0.541, *p* = 6.3x10^{-17}) and when further removing the *LCT* and *ABO* loci from the
analyses (r_S = 0.540, *p* = 6.3x10^{-17}).

We also evaluated if we could replicate any of the genome-wide significant signals outside the
LCT and *ABO* loci that were reported in a recent and similarly-sized Finnish population study[18].
After extracting all associations with *p* < 1x10^{-4} in our dataset, we identified 3 out of 451
genome-wide significant SNPs from the Finnish study using direct or proxy (r^2 > 0.8) information
(279 of the SNPs reported in this study were not included in our dataset because their MAF was
< 0.05). For one SNP, rs642387, we identified an association with consistent allelic effect for
similar taxa: the minor allele at this variant (through proxy SNP rs632222) was associated with a
decreased abundance of the *Desulfovibrio* genus (*p* = 8.7x10^{-5}) in our study. This corroborates
the allelic effects seen in the Finnish study for the abundances of phylum Desulfobacterota A,
class Desulfovibrionia, order Desulfovibrionales and family Desulfovibrionaceae. For the other

taxa, we observed consistent direction in our dataset, although the signals did not reach
nominal significance ($p = 0.14$) (**Supplementary Table 9**).

Mendelian randomization suggests salt intake and triglycerides are controlled by
microbiome abundance

To investigate the causal relationships between microbiome composition/function and complex
traits, and *vice-versa*, we used publicly available summary statistics in conjunction with our
mbQTL results to perform two-sample Mendelian randomization (MR) analyses (see **Methods**).
We focused on 78 phenotypes representing autoimmune diseases, cardiometabolic diseases,
and related risk factors, as well as food preferences (**Supplementary Table 10**), and on 37
microbiome features that were associated with at least one variant at genome-wide significance
($p < 5 \times 10^{-8}$). None were significant at FDR < 0.05. At FDR < 0.1 (corresponding to a $p = 2.7 \times 10^{-5}$
obtained from the inverse-variance weighted (IVW) MR test), we observed three causal
relationships in the direction from microbiome to phenotypes. Two of the three are based on the
same SNPs used as genetic instruments: genetic predispositions to a higher abundance of
family Rikenellaceae and genus *Alistipes* are linked to lower consumption of salt ('Salt added to
food' ordinal questionnaire phenotype, both traits: IVW $p = 2.4 \times 10^{-5}$, causal effect = -0.06 SDs of
salt intake for each SD increase in Rikenellaceae or *Alistipes* abundance, based on three
genetic instruments located on different chromosomes) (**Supplementary Table 11**)
(**Supplementary Figure 3a,b**). This correlation was not significant in the opposite direction
(IVW $p = 0.32$ for both), suggesting that Rikenellaceae abundance may have an effect on the
increased consumption of salt without being tagged by a shared causal factor. The other
potential causal relationship that passed our FDR < 0.1 threshold was between the genus
Collinsella and triglyceride levels. We found that a genetic predisposition to a one SD increase
in the relative abundance of *Collinsella* was associated with a 0.12 SD decrease in triglyceride
levels ($p = 5.7 \times 10$-5, FDR = 0.084) (**Supplementary Table 11**) (**Supplementary Figure 3**). All
of these causal estimates were consistent when using other MR tests (weighted median and
MR Egger) (**Methods** and **Supplementary Table 11**). Furthermore, there was no evidence for
the causal effect being affected by pleiotropy (Egger intercept $p = 0.74$ for Rikenellaceae and
Alistipes and 0.755 for *Collinsella*) (**Supplementary Table 11**). The results found in the two-
sample MR were very similar when we performed a polygenic risk score analysis in the UK
Biobank cohort (**Supplementary Notes**).

While all these potential causal effects are intriguing and exhibit robustness to sensitivity
analyses, we recognize that the limited availability of genetic variants and the high FDR
threshold dictate caution in their interpretation.

Power analysis indicates that larger sample sizes are necessary to identify host genetic
effects for microbial features with low prevalence

One of the complicating factors in any genome-wide association study of the gut microbiome is
the variable detection rate of the pathways and taxa measured, which reduces the effective
sample size and consequently statistical power. In our study, the additive effect of associated
variants at *LCT* explains 0.8% of the variance for species *B. longum* and *B. adolescentis*, which
are present in >80% of the samples, and this allowed us to have ~70% power to detect the
association at $p < 1\times10^{-10}$ (**Figure 5**). In contrast, we had no power (<5%) to detect the
association for species *B. bifidum* and *B. catenulatum* at this level of significance, as they were
present in only 14% and 26% of samples, respectively. For *B. bifidum*, despite it being a rare
species, we were able to detect a study-wide significant association at *ABO* because the effect
was substantial (2.5% of the variance explained by SNP rs8176645, power = 83%), but we were
underpowered to see the association at this locus with species *C. aerofaciens*, which was
present in only 45% of the samples, and for which the additive effect of *ABO* variants was
smaller (0.5% variance explained) (**Figure 5**). In a power analysis, we estimated that our
sample size (7,738 individuals) is underpowered to detect genetic effects for taxa that are
present in <80% of samples when considering an effect size comparable to or smaller than the
effect of *LCT* for *B. adolescentis.* This is striking, considering that 93% of the taxa we identified
in this cohort are present in <80% of the samples (**Methods**). We estimate that to find a similar
effect size for at least 20% of gut microbiome composition (approximately 20% of the taxa are
present in at least 20% of the samples in our cohort) would require a cohort of ~50,000
individuals.

Discussion

We carried out the largest genome-wide association study of gut microbiome composition and
function in a single population by analyzing metagenomic sequencing data in 7,738 volunteers
from the northern Netherlands. We recapitulated genetic associations at two known loci, *LCT*
and *ABO,* and the resolution of our metagenomic sequencing allowed us to pinpoint novel
associations with species (*B. adolescentis* at *LCT*, *B. bifidum* and *C. aerofaciens* at *ABO*) and
bacterial pathways (*LACTOSECAT-PWY* at *ABO*) in these loci. Furthermore, we identified

associations ($p < 5 \times 10^{-8}$) at 22 other loci for four taxa and 18 bacterial pathways. None of these
associations were affected by major confounders of the gut microbiome such as medication
usage, diet and BMI. Finally, we used an MR approach to pinpoint potential causal links
between gut microbiome composition, complex traits and food intake habits.

The strongest association with the gut microbiome was found at the *LCT* locus, which remains
the most robust genetic association with the gut microbiome identified to date. Associations at
this locus with *Bifidobacterium* have been consistently reported in studies of different ethnicities,
across a range of sample sizes, and in studies using different technologies and protocols for gut
microbiome characterization[6,8,13,14,42]. We also recapitulate that an increase of *Bifidobacterium*
was more evident in LI individuals who were consuming milk or milk-derived products[13,43]. In
addition, given that the resolution of metagenomic sequencing allows for species-level
characterization of microbiome profiles, we could show that this effect was mainly attributable to
the species *B. longum, B. adolescentis, B. catenulatum* and *B. bifidum*, an observation that was
corroborated by two recent studies in Finnish and US Hispanic/Latino populations[18,44].

We found another study-wide association in the *ABO* gene locus with changes in abundance of
several taxa and pathways. Associations with microbiome composition and blood types were
observed in previous experimental studies[45,46]. Genetic associations at the *ABO* locus have also
been reported previously for populations of different ethnicities and in non-human species. For
example, associations with *Bacteroides* and *Faecalibacterium* were reported in a study of five
German cohorts that used 16S rRNA sequencing for gut microbiome characterization[42] and with
the microbial pathway of lactose and galactose degradation in a cohort of 3,432 Chinese
individuals[47]. The importance of *ABO* in determining host☐microbiome interaction has also been
recently reported in pigs[48]. A deletion at this locus that inactivates the ABO acetyl-
galactosaminyl-transferase has been shown to change the porcine microbiome composition by
altering intestinal N-acetyl-galactosamine concentrations and consequently reducing the
abundance of *Erysipelotrichaceae* strains, which have the capacity to import and catabolize N-
acetyl-galactosamine[49]. In our analysis, the strongest associations were with *B. bifidum*, which
has not been reported before, and with the *Collinsella* genus, which was recently reported[18],
and at genome-wide significance with *C. areofaciens*. We did not detect any evidence of an
interaction with diet at this locus, although this could be due to limitations in available
information as the recording of dietary information was done at different times than the stool
collection. We did, however, find that associations at this locus depend on secretor status that is
determined by a nonsense mutation at the *FUT2* gene[42] and thus on the host's ability to

incorporate antigens into bodily fluids that are released in the gut. Intriguingly, we observed that
taxa associated with the *ABO* locus also showed evidence of association at the *LCT* locus, this
association being independent of blood type (interaction pvalue >0.05 for all taxa), indicating a
common, independent action of these two loci in contributing to the growth of these bacteria.
The most compelling hypothesis is that the availability of sugars in the gut, via undigested
lactose in LI individuals or secretion of antigens with accessible glycans in non-O blood type
secretors, provides direct energy sources for most of these bacteria. This is further supported by
the observation that LI individuals and non-O blood type secretors were both associated with
the increased abundance of a bacterial pathway for lactose and galactose degradation.
However, this mechanism would not fully explain the opposite direction of the association at
ABO seen for *B. bifidum,* which, under this hypothesis and considering its adaptation to normal
gastrointestinal mucosa (apparently independent of H-antigen secretion, as shown in **Figure 4**),
would be subjected to competition in the environment, such as in the case of non-O blood
secretors subjects. Of note, a similar pattern of complex association at this locus was found in a
recent study of German cohorts[42], where a branch of the Bacteroides genus represented by
OTU97_12, OTU99_12, and TestASV_13, showed association with an inverse relationship
between their prevalence and the non-O blood type group and instead a positive relationship
with prevalence of OTU97_27. While these opposite associations could be explained by antigen
degradation activity of certain species and consequent environment competition for others,
more studies are needed to clarify the complex mechanisms involved

We acknowledge that anachronistic diet information is a limitation of our study. While we have
shown that in general the microbiome remains fairly stable in an individual after 4 year, with
inter-individual differences being larger than inter-individual differences[50] , short-term changes in
diet, especially those introducing drastic shifts, can perturbate both microbiome composition and
function. These could not be taken in account by our analyses. Capturing these shifts would not
be easy even with frequency food questionnaires recorded at the time of sample collection;
ideally, real-time extensive recording in weeks preceding microbiome collection should be
implemented in future biobanks.

The strongest mbQTLs we identified reside in genes under selective pressure. The *LCT* gene is
a highly differentiated gene among human populations due to positive selection conferred by the
lactase persistence phenotype. It has been estimated that strong selection occurred within the
past 5,000–10,000 years, consistent with there being an advantage to lactase persistence and
the ability to digest milk in the setting of dairy farming[51]. Variants at this locus have been linked

through GWAS to not only food habits and metabolic phenotypes, but also to immune cell
populations[52]. The *ABO* locus is evolutionarily highly differentiated; it has been shown to have
experienced balancing selection for the last 3 million years in many primate species[53]. Several
evolutionary sources of selective pressure have been proposed, including via infections by
pathogens such as malaria[54] and cholera[55]. *ABO* variants have also been linked to
cardiometabolic traits, white and red blood cell levels and cytokine levels[56,57]. Therefore,
host‑microbiome interactions are likely to be shaped by human‑microbe co-evolution and
survival, probably through a balance between food availability for gut bacteria and enhanced
immune response of the host. A better understanding of these interactions will expand our
current knowledge of human evolution[58].

From this perspective, it will be crucial to compare genetic studies of the gut microbiome in
diverse populations with different genetic backgrounds in order to understand the complexity of
host‑microbe interactions. This requires community efforts to standardize the definition of
taxonomies and to standardize measurement methodologies in order to facilitate comparison
between cohorts. For example, in our attempt to replicate the findings from the Finnish
population cohort, only a limited number of taxa could be directly matched or connected through
the Genome Taxonomy Database[59].

To explore the causality of the relations of the microbiome with complex traits and food
preferences, we performed bi-directional MR analysis using 56 dietary traits, 16 diseases and 5
biomarkers. At FDR < 0.1, we observed three causal relationships. Although the limited impact
of genetic variants on both microbiome composition and dietary preferences requires caution
when interpreting causality estimation by MR[60,61], it is intriguing to observe a causal role for
microbial composition on food preferences. Here we observed that an increase in the
abundance of genus *Alistipes* and its family Rikenellaceae led to decreased consumption of
salt. Although multiple studies have shown that dietary changes, such as variations in salt and
meat intake, have a strong effect on microbiome composition[62], it is intriguing to suggest that
genetically determined variations in the microbiome might affect individual food preferences.
This would be supported by bacterial genetic variations in salt tolerance[63] and by the
established knowledge that the composition of gut microbiome can accurately predict the effect
of food items on host metabolism[64]. There was additional evidence from our results to support a
role for the microbiome in influencing food preferences. A perfect proxy of rs642387 (rs503397,
$r^2 = 0.99$ in 1000 Genomes Europeans), which was recently reported to be associated with the
family Desulfovibrionaceae and related taxa in a Finnish population and replicated in our study,

had previously been associated to bitter alcoholic beverage consumption in an independent
cohort[65]. Another study found an increase in family Desulfovibrionaceae in individuals with high
alcohol consumption[66], and family Desulfovibrionaceae, genus *Desulfovibrio* and other related
taxa were also associated with increased consumption of alcohol in our DMP cohort[16],
supporting a pleiotropic effect of this locus on both microbiome and alcohol intake. These
studies, together with our findings, suggest that gut microbiota could indirectly influence an
individual's food preferences by mediating the downstream effect of the consumption of different
products.

Understanding the differences in microbial pathway abundance is crucial for deeply
understanding the underlying function of the gut microbiome. Microbiome measurements that
target single organisms or taxonomic groups may be inefficient because two strains of one
species may differ in their functional capacities. It is therefore essential to evaluate not only the
taxonomic but also the functional composition of the microbiome. To our knowledge, we report
the largest single study to investigate host genetic effects on the human gut microbiome
function using metagenomics sequencing and thereby perform genome-wide association
analysis of microbial pathways. Our analyses revealed several loci associated with microbial
pathways, although only one passed stringent study-wide significance. Functional pathway
abundances are also an approximate indicator of metabolite expression. For example, we report
an association with PWY-5088, which is involved in propionate production. Although we do not
have SCFA levels in our samples for direct validation, we have measured propionate levels in
feces in the LL-DEEP cohort and using summary statistics from our previous study[17], we see
consistent directions of effect for the association with PWY-5088 (allele T was associated with
decreased levels of propionate), but the association was not significant, perhaps due to the
small sample size (N=897, *p* value = 0.08). It also should be noted that pathway abundance,
even if not necessarily representing the expression, might represent the gene load and
abundance of the bacterial taxa that are able to execute this metabolic pathway, which by itself
is an important indicator for the metabolic capacity of a given gut microbiomal community. We
believe more metabolite screening should be undertaken once more robust and consistent
replications are detected for microbial pathways by other genetic studies.

In addition to the two study-wide significant loci, we also observed 22 loci at genome-wide
significance and many more at a more lenient threshold. The correlation we observe between
the heritability of microbial taxa and pathways and the number of suggestively associated loci
indicates that mbQTLs with smaller effects are likely to exist. Those loci remained under the

detection limit in the current study, and the sample size would need to be increased by orders of
magnitude to discover their role. In fact, gut microbiome composition is characterized by high
inter-individual variation. Analysis of core microbiota in different populations indicates that only a
few bacteria are present in >95% of studied individuals[5,13,16], drastically reducing the effective
sample sizes for analyses. We performed power calculations taking into account the inter-
individual variations in microbiome composition and concluded that a sample size comparable
to our study (~8,000 participants) is only sufficient to identify associations to taxa present in
>80% individuals, which comprise only 7% of all identified taxa in the cohort. For bacteria
present in >20% of the samples, >50,000 participants would be necessary to identify an effect
size similar to that of *LCT* and *ABO*. Much larger cohorts are therefore required to identify
genetic contribution to rarer bacteria. We conclude that joint efforts that combine tens of
thousands of individuals, combined with harmonized methodology to reduce technical bias, will
be needed to characterize more than a few major loci, as has also been the case for genetic
studies of much more heritable quantitative traits such as BMI (heritability ~40%), height
(heritability ~80%) and other human phenotypes[67–69].
538
539
540
541   **Methods**
542
Cohort description
Lifelines is a multi-disciplinary prospective population-based cohort study with a unique three-
generation design that is examining the health and health-related behaviors of 167,729 people
living in the north of the Netherlands. Lifelines employs a broad range of investigative
procedures to assess the biomedical, socio-demographic, behavioral, physical and
psychological factors that contribute to the health and disease of the general population, with a
special focus on multi-morbidity and complex genetics[70,71]. During the first follow-up visit, all
participants were invited to participate in a parallel project, The Dutch Microbiome Project
(DMP), on a voluntary basis. The goal of this project is to evaluate the impact of different
exposures and life-styles on gut microbiota composition[16]. A subset of 10,000 Lifelines
participants were included, for 8,719 of them both feces and phenotype information were
collected. Of these, 8,208 were retained for downstream analysis after stringent quality control[16]
. Distribution of age, gender and location within the three north provinces was similar to that
observed in the total Lifelines cohort[71] .

The Lifelines study was approved by the medical ethical committee from the University Medical
Center Groningen (METc number: 2017/152). Additional written consents were signed by the
DMP participants or legal representatives for children aged under 18.

Genome characterization
Genotyping of 38,030 Lifelines participants was carried out using the Infinium Global Screening
array® (GSA) MultiEthnic Diseases version, following the manufacturer's protocols, at the
Rotterdam Genotyping Center and the Department of Genetics of the University Medical Center
Groningen. Here, we used available quality-controlled genotyping data imputed with Haplotype
Reference Consortium (HRC) panel v1.1[72], as described elsewhere[73]. To avoid population
stratification, we analyzed only European samples. We selected 43,587 genetic markers by
applying linkage disequilibrium (LD) pruning on the genotyped data (sliding window of 1Mb,
linkage disequilibrium r^2<0.2, step = 5) and used them for a Principal Component Analysis
(PCA) projecting the 1000 Genomes (all populations) and GoNL cohorts a population
references[74,75]. Europeans were then identified as samples clustering together with European
populations according to the first two principal components or <3 standard deviations away from
the most distant European samples in the reference. In total, 35 non-European participants
were removed. Quality-controlled genotype information was obtained for 7,738 of the DMP
participants for whom quality-controlled microbiome data and BMI were also available. Of these,
58.1% were females and had ages ranging from 8 to 84 (mean of 48.5 years). The mean BMI
value was 25.58 (range from 13.10 to 63.70).

Microbiome characterization
The gut microbiome was characterized from stool samples as described in Gacesa et al[16]. In
brief, stool samples were collected by participants and frozen within 15 minutes after production
and then transported on dry ice to the Lifelines facility to be stored at -80°C. Microbial DNA was
extracted using the QIAamp Fast DNA Stool Mini Kit (Qiagen, Germany), following the
manufacturer's instructions. Samples with a total DNA yield lower than 200 ng (as determined
by Qubit 4 Fluorometer) were prepared using NEBNext® Ultra™ DNA Library Prep Kit for
Illumina, and samples with higher DNA yield were prepared using NEBNext® Ultra™ II DNA
Library Prep Kit for Illumina®. Shotgun metagenomic sequencing was carried out using the
Illumina HiSeq 2000 platform at Novogene, China. Metagenomic sequencing data was profiled
following methods previously used in other cohorts, as described in[16,76]. Low quality reads

(PHRED quality ≤ 30), adapters and host sequences were removed using KneadData tools

v0.5.1. Taxonomic composition was determined with MetaPhlAn2 v2.7.2 [77]. Characterization of

biochemical pathways was performed with the HUMAnN2 pipeline v0.11.1[78], integrated with the

UniRef90 v0.1.1 protein database[79], the ChocoPhlAn pangenome database and the DIAMOND

alignment tool v0.8.22[80]. After quality control (samples with eukaryotic or viral abundance ≤ 25%

and total read depth ≥ 10 million were retained), we had information on 950 microbial taxa and

559 functional pathways. For this study, we focused only on bacterial taxa and functional

pathways with mean relative abundance >0.001% across all samples and present in at least

1000 of the 7,738 participants, which resulted in a list of 207 taxonomies (5 phyla, 10 classes,

13 orders, 26 families, 48 genera and 105 species) and 328 bacterial pathways. Furthermore,

we removed redundant pathways by discarding one pathway among pairs that were highly

correlated (r_S > 0.95), as well as pathways not previously described in bacteria that could thus

be coming from sources other than bacteria, resulting in 205 pathways for genetic analyses.

Diet phenotypes definition

Dietary habits were assessed using a semi-quantitative Food Frequency Questionnaire (FFQ)

designed and validated by the division of Human Nutrition of Wageningen University as

described before in Siebelink et al and Gacesa et al.[16,81]. The FFQ data was collected 4 years

prior to fecal sampling, and supplementary questionnaires were collected concurrent with fecal

sampling, with the stability of long-term dietary habits between time points assessed as

described in Gacesa et al.[16]. We analyzed the dietary factors that were previously found to be

associated (FDR < 0.05) to the microbiome features in our study that had a genome-wide

significant signal in the *ABO* and *LCT* loci. We also analyzed lactose intake for the species

Bifidobacterium longum and *Bifidobacterium adolescentis*, given their association with the *LCT*

region in our study. Participants with an implausible caloric intake (< 800 or > 3934 kcal/day for

males, < 500 or > 2906 kcal/day for females)[82] were not included in these analyses.

GWAS analysis method

Genome-wide association analysis was performed in 7,738 European samples for 412 features

(205 functional pathways and 207 microbial taxa), investigating genetic additive effects using

allele dosages for 5,584,686 genetic variants with MAF > 0.05 and information score > 0.4 on

the autosomes (chromosomes 1-22) and the X chromosome. We focused on the quantitative

dimensions of relative bacterial and pathway abundances, treating all zero values as missing

data. We used natural-log-transformed abundances and regressed these in a linear mixed

model using SAIGE v.0.38[19], with age, sex and the genetic relationship matrix (GRM) among

participants as covariates. We used the standard settings of SAIGE, which applies inverse-rank

normalization to the traits prior to the association analyses. The GRM was built with SAIGE

using a set of 54,565 SNPs selected from the total set of quality-controlled SNPs directly

genotyped and filtered for allele frequency and redundancy (MAF $\geq$ 0.05, r^2 < 0.2, sliding

window = 500Kb).

Definition of the study-wide significant threshold

To estimate the number of independent phenotypes assessed, we used PCA on the matrix of

412 microbiome features (207 taxonomies and 205 pathways) available for GWAS analysis to

decompose variability in independent components (axes). We estimated that 264 components

are needed to explain 90% of the microbiome variance. We then defined our study-wide

significant p-value threshold by correcting the genome-wide significance threshold for this factor

($5 \times 10^{-8}/264 = 1.89 \times 10^{-10}$).

Association using dominant model

To evaluate association using a dominant model on SNP rs182549 at the *LCT* locus, we used

best-guess genotypes and converted T/C to T/T. Association analysis was then run for all taxa

as well as the *LACTOSECAT-PWY* pathway using SAIGEgds[83] and the same covariates and

transformation used for the GWAS analysis.

Inference of blood groups

We estimated blood groups from genotyped and imputed data following the scheme of

Ellinghaus et al.[84]. Specifically, we used the absence of the rs8176719 insertion to define blood-

type allele O1, the T allele of rs41302905 to define blood-type allele O2 and the T allele of

rs8176746 to define the blood-type allele B (instead of rs8176747). Diploid individuals O1O1,
O2O2 and O1O2 were considered blood type O. Diploid individuals O1B, O2B and BB were
considered blood type B. Absence of the alleles mentioned above was used to define blood-
type allele A. To evaluate differences across blood types, we compared the mean relative
abundance of microbiome features in individuals with A, B and AB blood type to that in
individuals with the O blood type using a two-sided Wilcoxon test. To evaluate the interaction
with the rs601338 *FUT2* (secretor/non-secretor) locus, we grouped individuals into two groups
(non-O blood type and blood type O) to distinguish production or non-production of antigens,
and compared pairs using a two-sided Wilcoxon test. All analyses were done using base R
v3.6.1 (https://www.R-project.org/).

Effects of potential confounders on significantly associated loci
We evaluated the robustness of genome-wide associated signals by incorporating the following
potential confounders into our statistical model: medication usage, anthropometric data and
stool frequency and consistency data (collection and processing was described in Gacesa et
al.[16]). We analyzed the effects of the following medication groups: proton pump inhibitors (ATC
A02BC, N = 130), laxatives (osmotic ATC A06AD, N = 44; volume increasing ATC A06AC, N =
77), one group of antibacterials (ATC J01, N = 24) and other anti-infectives (ATC J, N = 39).
The other group of medication considered was antibiotic use in the 3 months prior to stool
collection (N = 450). For each of these medications, we created dichotomous variables for all
participants coded as 0 (non-user) or 1 (user). The other factors included were BMI, stool
frequency and stool consistency (mean Bristol stool scale). All models also incorporated age
and sex as covariates and were run only for the genome-wide☐significant SNP☐trait pairs
(**Supplementary Table 1**) using the same software used for GWAS (SAIGE, Zhou et al.[19]). To
evaluate the impact of these covariates on the genetic signals, we used Cochran's Q
heterogeneity test to compare the effect size obtained by the covariate-inclusive model and the
basic model (that only includes age, sex and the genetic variant). To evaluate the impact of sex,
we ran the SNP☐association analysis in SAIGE separately for males and females using only
age as a covariate. For each genetic variant, differences in effect size in males and females
were tested using Cochran's Q heterogeneity test.

Interaction analyses
We used a three-step procedure to evaluate gene☐diet interactions for all the taxa and
pathways associated with SNPs at the *LCT* and *ABO* loci. First, we extracted the variables

representing dietary habits that had previously shown significant association with these
microbial traits at FDR<0.05[16]. For the genus *Collinsella* and pathway *LACTOSECAT-PWY*, no
dietary factors were found at this FDR threshold. We therefore considered the same dietary
factors associated with B. *bifidum* in the analyses, given that they showed similar patterns of
genetic association. Next, we added these variables to the basic genetic model (feature ~ age +
sex + genetic variant) to confirm their association at (at least) nominal significance level ($p <$
0.05) while accounting for the associated genetic variant(s). Finally, for the dietary variables
showing nominal significance, we evaluated the interaction with the genetic variant(s) by
including an interaction term into the association model. For the *LCT* locus, we considered a
binary variable to distinguish two groups of genotypes at SNP rs182549 (C/C vs. C/T and T/T)
according to the dominant inheritance model at this locus. For the *ABO* locus, we used a binary
definition of blood type (blood type O vs A/B/AB) and also considered the effect of the rs601338
genotype in the *FUT2* gene (defining secretor/non-secretor individuals). All microbiome features
were inverse-rank normalized before analyses, and age and sex were added as covariates as in
the main GWAS analysis. All models were fit using the lm() function from base R v3.6.1, other
statistical tests were as implemented in packages *rstatix* v0.5.0 and *ggpubr* v0.3.0 and package
RNOmni v0.7.1 was used for the inverse-rank normalization.

Replication in other cohorts and data sets
We looked for replication of our results using summary statistics from two independent studies:
a genome-wide meta-analysis of 16S rRNA data from 24 cohorts (the MiBioGen consortium)
and a genome-wide study on metagenomics data in the LL-DEEP cohort, another subset of the
Lifelines cohort with data generated 4 years before the DMP and in which 255 participants were
also later enrolled in DAG3[13,17]. In this study, a different DNA isolation procedure (AllPrep Kit)
was used and taxonomies were defined using the Bracken pipeline, which may explain why C.
areofaciens was not identified. In the MiBioGen study, we could not look at SNPs associated
with species or pathways, as these microbiome features cannot be defined using 16S data, or at
SNPs on the X chromosome, as they were not analyzed in this study. In the second study, we
searched for the exact same taxonomy or pathway, but similarly to MiBioGen, X chromosomal
variants were not tested and some taxonomies were not defined due to the differences in
metagenomic data processing pipelines.

Next to the replication of our findings, we also evaluated whether the genome-wide signals
reported in a recent genome-wide study of microbiome taxa from a Finnish cohort were
replicable in our data[18]. We searched for all SNPs outside the *LCT* and *ABO* loci or any proxy (r^2

> 0.8) in our dataset and selected all SNP☐taxonomy pairs that showed a $p < 1\times10^{-4}$ with at
least one taxonomy in our cohort (**Supplementary Table 9**). We then looked at these
associated taxa in the respective cohorts and compared them visually and with the aid of the
Genome Taxonomy Database (https://gtdb.ecogenomic.org/) to determine if they were the same
bacterial taxa or taxa from the same taxonomic branch.

Correlation between heritability estimates and the number of associated loci
To analyze the correlation between family-based heritability and the number of suggestive
mbQTLs, we used narrow-sense heritability estimates for taxa and pathways that accounted for
household environment sharing and previously derived for this cohort[16]. We then calculated the
number of independent mbQTLs per microbial trait by performing LD pruning ($r^2 < 0.1$ in our
data set, window size 1Mb using Plink v1.9[85]) for all SNPs at the three different thresholds: $p <$
5×10^{-4}, $< 1\times10^{-4}$ and $< 5\times10^{-5}$. The association of heritability and the number of mbQTLs was
calculated in R v.4.0.3 using a weighted Spearman correlation from the `wCorr` v1.9.1 package,
with each taxon or pathway treated as a data point. The weights used in calculating the
correlation were inversely proportional to the Z scores calculated from heritability pvalue
estimates. The regression lines in **Figure 4** were fit using the *loess* function (locally estimated
scatterplot smoothing, base R package v4.0.3) with span and degree parameters set to 1.

Mendelian randomization analysis
To evaluate potential causal relationships between the gut microbiome and other common traits,
we performed Mendelian randomization (MR) analyses that combined the summary statistics of
the microbiome with publicly available summary statistics on food preferences, autoimmune and
cardiovascular diseases and other cardio-metabolic traits. We analyzed the 37 microbiome
features (pathways and taxa) with at least one variant passing the $p < 5\times10^{-8}$ threshold
(**Supplementary Table 2**) and combined these with 78 publicly available summary statistic
datasets retrieved using the IEU GWAS DATABASE[86].

We performed a bi-directional MR analysis, first testing if microbiome traits causally affect a
phenotype and then testing if phenotypes can causally affect the microbiome traits. For each
comparison, we intersected the microbiome variants (MAF > 0.05) by rsID, position and alleles
with the publicly available summary statistic variants. We then selected instruments using the
'clump_data()' function of the TwoSampleMR package (v0.5.5)[87]. The publicly available
summary statistics were clumped using a *p*-value threshold of $< 5\times10^{-8}$ and otherwise standard

settings ($r^2 < 0.001$, 10mb window size). Due to the limited statistical significance of the

microbiome traits, we performed *p*-value clumping at a less stringent $p < 5\times10^{-6}$ threshold. If

fewer than three variants were clumped, we removed the trait combination from analysis.

The MR analysis was done using the TwoSampleMR v0.5.5 package. We first selected trait

combinations that passed the Benjamini-Hochberg FDR threshold of 0.1 ($p = 2.805\times10^{-5}$) in the

inverse-variance weighting (IVW) test, resulting in one suggestively causal trait combination. We

further checked that the significant trait combinations were unlikely to be driven by pleiotropy

based on two criteria: i) the Egger regression intercept was nominally significant (indicating the

presence of horizontal pleiotropy) and ii) the weighted median results were not nominally

significant (indicating that no single variant influences the result)[88,89].

Power analysis

We calculated the variance explained at our loci using the formula described in Teslovich et

al.[90], which takes into account MAF, effect size, standard error and sample size. We then

performed a power analysis based on a linear model of association, considering different

genetic   effect   sizes   (variance   explained)   and   sample   sizes   (N)

<https://genome.sph.umich.edu/wiki/Power_Calculations:_Quantitative_Traits>. We   performed

a sample size calculation by doing a grid search in the sample size sequence: [1000, 1050, … ,

50000] and kept the lowest sample size that had power > 80%.

Data availability

Raw sequencing microbiome data is available at EGA (accession number EGAS00001005027**).**

Genotyping data and participant metadata are not publicly available to protect participants'

privacy and research agreements in the informed consent. The data can be accessed by all

bona-fide researchers with a scientific proposal by contacting the Lifelines Biobank (instructions

at: https://www.lifelines.nl/researcher). The full summary statistics are available for download at

https://dutchmicrobiomeproject.molgeniscloud.org. The processed microbiome data (taxonomy

and pathway abundance per individual) can also be downloaded after filling in a request form

available at the same website and after signing a data access agreement. The authors declare

that all other data supporting the findings of this study are available within the paper and its

Supplementary Information files.

LifeLines Cohort Study - group authors genetics

Raul Aguirre-Gamboa[1], Patrick Deelen[1], Lude Franke[1], Jan A Kuivenhoven[3], Esteban A Lopera
Maya[1], Ilja M Nolte[7], Serena Sanna[1,6], Harold Snieder[7], Morris A Swertz[1,4], Judith M Vonk[7],
Cisca Wijmenga[1]

[7] *Department of Epidemiology, University of Groningen, University Medical Center Groningen, the Netherlands*

Acknowledgements

The authors wish to acknowledge the services of the Lifelines Cohort Study, the contributing
research centers delivering data to Lifelines and all the study participants. The Lifelines initiative
has been made possible by subsidy from the Dutch Ministry of Health, Welfare and Sport; the
Dutch Ministry of Economic Affairs; the University Medical Center Groningen (UMCG); the
University of Groningen (UG) and the Provinces of the North of the Netherlands (Drenthe,
Friesland and Groningen). This project was carried out under Lifelines Project number
OV18_0464. We thank Mathieu Plateel and Jody Geelderloos-Arends for their contribution in
genotyping the Lifelines samples, Kate McIntyre for help developing the manuscript, Marije van
der Geest for setting up the website for sharing summary statistics and Patrick Deelen for
discussion of results. We also thank the UMCG Genomics Coordination Center, the UG Center
for Information Technology and their sponsors BBMRI-NL & TarGet for storage and
computational infrastructure, and the company Novogene for providing gut metagenome
sequencing of all DMP samples. Finally, we thank the UK Biobank for making their resource
available. Analyses of UK Biobank data described in this work were carried out under project
number 48548 to C.W.

The generation and management of genotype data for the Lifelines Cohort Study was supported
by the UMCG Genetics Lifelines Initiative (UGLI). Genotyping QC was supported by UMCG
HAP grant CD017.0031/ ronde 2017-2/ nr 324. Metagenomics sequencing of the cohort was
mainly funded by the CardioVasculair Onderzoek Nederland (CVON) grant (CVON 2012-03) to
Marten Hofken (who passed away in 2016), JF and AZ, as well as by other grants to RKW and
CW listed below. RG and RKW are supported by the collaborative TIMID project (LSHM18057-
SGF) financed by the PPP allowance made available by Top Sector Life Sciences & Health to
Samenwerkende Gezondheidsfondsen (SGF) to stimulate public-private partnerships and co-
financing by health foundations that are part of the SGF. RKW is supported by the Seerave
Foundation. AZ is supported by a European Research Council (ERC) Starting Grant 715772, a
Netherlands Organization for Scientific Research (NWO) VIDI grant 016.178.056, a CVON grant
2018-27 and the NWO Gravitation grant ExposomeNL 024.004.017. JF is supported by the

NWO Gravitation grant The Netherlands Organ-on-Chip Initiative (024.003.001), a CVON grant

2018-27, a NWO VICI grant VI.C.202.022 and ERC Consolidator grant 101001678. CW is

further supported by an ERC advanced grant (ERC-671274) and an NWO Spinoza award

(NWO SPI 92-266). HJMHd is supported by the INTERREG V A (202085) funded project

EurHealth-1Health. LC is supported by a joint fellowship from the UMCG and China Scholarship

Council (CSC201708320268) and a Foundation De Cock-Hadders grant (20:20-13). ELM is

supported by Colciencias fellowship ed.783.

Author Contributions

ELM, AK and SH performed genetic analyses.

AvdG performed Mendelian randomization and power analyses.

AK, SH, LC, AVV and RG processed microbiome data.

SAS, TS, VC, MAYK, LAB and MBG processed samples meta-data.

PBTN and MAS provided resources on the HP computing cluster and assisted with data

management.

HJMH, CW, JF, RKW and AZ provided funding resources and designed the DMP.

AZ and SS supervised the study.

ELM, AK, AvdG, SH, SAS, AZ and SS drafted the manuscript.

All authors were involved in data interpretation and provided critical input to the manuscript

draft.

Competing Interests statement

837   The authors declare no conflicts of interest.

References

1. Valdes, A. M., Walter, J., Segal, E. & Spector, T. D. Role of the gut microbiota in nutrition and health. *BMJ* **361**, k2179 (2018).

2. Hall, A. B., Tolonen, A. C. & Xavier, R. J. Human genetic variation and the gut microbiome in disease. *Nat. Rev. Genet.* **18**, 690–699 (2017).

3. Fan, Y. & Pedersen, O. Gut microbiota in human metabolic health and disease. *Nat. Rev. Microbiol.* (2020) doi:10.1038/s41579-020-0433-9.

4. Zhernakova, A. *et al.* Population-based metagenomics analysis reveals markers for gut microbiome composition and diversity. *Science* **352**, 565–569 (2016).

5. Falony, G. *et al.* Population-level analysis of gut microbiome variation. *Science* **352**, 560–564 (2016).

6. Goodrich, J. K. *et al.* Genetic Determinants of the Gut Microbiome in UK Twins. *Cell Host Microbe* **19**, 731–743 (2016).

7. Rothschild, D. *et al.* Environment dominates over host genetics in shaping human gut microbiota. *Nature* **555**, 210–215 (2018).

8. Bonder, M. J. *et al.* The effect of host genetics on the gut microbiome. *Nat. Genet.* **48**, 1407–1412 (2016).

9. Wang, J. *et al.* Genome-wide association analysis identifies variation in vitamin D receptor and other host factors influencing the gut microbiota. *Nat. Genet.* **48**, 1396–1406 (2016).

10. Turpin, W. *et al.* Association of host genome with intestinal microbial composition in a large healthy cohort. *Nat. Genet.* **48**, 1413–1417 (2016).

11. Hughes, D. A. *et al.* Genome-wide associations of human gut microbiome variation and implications for causal inference analyses. *Nat. Microbiol.* **5**, 1079–1087 (2020).

12. Kurilshikov, A., Wijmenga, C., Fu, J. & Zhernakova, A. Host Genetics and Gut Microbiome: Challenges and Perspectives. *Trends Immunol.* **38**, 633–647 (2017).

13. Kurilshikov, A. *et al.* Large-scale association analyses identify host factors influencing

human gut microbiome composition. *Nat. Genet.* **53**, 156–165 (2021).

14. Blekhman, R. *et al.* Host genetic variation impacts microbiome composition across human

body sites. *Genome Biol.* **16**, 191 (2015).

15. Vieira-Silva, S. *et al.* Species–function relationships shape ecological properties of the

human gut microbiome. *Nat. Microbiol.* **1**, 1–8 (2016).

16. Gacesa, R. *et al.* The Dutch Microbiome Project defines factors that shape the healthy gut

microbiome. *bioRxiv* 2020.11.27.401125 (2020) doi:10.1101/2020.11.27.401125.

17. Sanna, S. *et al.* Causal relationships among the gut microbiome, short-chain fatty acids and

metabolic diseases. *Nat. Genet.* **51**, 600–605 (2019).

18. Qin, Y. *et al.* Combined effects of host genetics and diet on human gut microbiota and

incident disease in a single population cohort. *medRxiv* 2020.09.12.20193045 (2020)

doi:10.1101/2020.09.12.20193045.

19. Zhou, W. *et al.* Efficiently controlling for case-control imbalance and sample relatedness in

large-scale genetic association studies. *Nat. Genet.* **50**, 1335–1341 (2018).

20. Liu, X. *et al.* M-GWAS for the gut microbiome in Chinese adults illuminates on complex

diseases. *bioRxiv* 736413 (2019) doi:10.1101/736413.

21. Van Der Heide, H. M., Magnee, W. & Van Loghem, J. J. Blood group frequencies in the

Netherlands. *Am. J. Hum. Genet.* **3**, 344–347 (1951).

22. Oriol, R., Le Pendu, J. & Mollicone, R. Genetics of ABO, H, Lewis, X and related antigens.

Vox Sang. **51**, 161—171 (1986).

23. Kelly, R. J., Rouquier, S., Giorgi, D., Lennon, G. G. & Lowe, J. B. Sequence and expression

of a candidate for the human Secretor blood group alpha(1,2)fucosyltransferase gene

(FUT2). Homozygosity for an enzyme-inactivating nonsense mutation commonly correlates

with the non-secretor phenotype. *J. Biol. Chem.* **270**, 4640–4649 (1995).

24. Hazra, A. *et al.* Common variants of FUT2 are associated with plasma vitamin B12 levels.

Nat. Genet. **40**, 1160–1162 (2008).

25. Turroni, F. *et al.* Bifidobacterium bifidum as an example of a specialized human gut

commensal. *Front. Microbiol.* **5**, (2014).

26. Turroni, F., Milani, C., Sinderen, D. van & Ventura, M. Genetic strategies for mucin

metabolism in Bifidobacterium bifidum PRL2010: An example of possible human-microbe

co-evolution. *Gut Microbes* **2**, 183–189 (2011).

27. Visschedijk, M. C. *et al.* Genomic and Expression Analyses Identify a Disease-Modifying

Variant for Fibrostenotic Crohn's Disease. *J. Crohns Colitis* **12**, 582–588 (2018).

28. Wang, K. *et al.* A genome-wide association study on obesity and obesity-related traits. *PloS*

One **6**, e18939 (2011).

29. Kim, C. H. Microbiota or short-chain fatty acids: which regulates diabetes? *Cell. Mol.*

Immunol. **15**, 88–91 (2018).

30. Pividori, M., Schoettler, N., Nicolae, D. L., Ober, C. & Im, H. K. Shared and distinct genetic

risk factors for childhood-onset and adult-onset asthma: genome-wide and transcriptome-

wide studies. *Lancet Respir. Med.* **7**, 509–522 (2019).

31. Consortium, T. Gte. The GTEx Consortium atlas of genetic regulatory effects across human

tissues. *Science* **369**, 1318–1330 (2020).

32. Ostrowski, J. *et al.* Genetic architecture differences between pediatric and adult-onset

inflammatory bowel diseases in the Polish population. *Sci. Rep.* **6**, 39831 (2016).

33. Kunisawa, J. & Kiyono, H. Vitamin-Mediated Regulation of Intestinal Immunity. *Front.*

Immunol. **4**, (2013).

34. Kunisawa, J., Hashimoto, E., Ishikawa, I. & Kiyono, H. A Pivotal Role of Vitamin B9 in the

Maintenance of Regulatory T Cells In Vitro and In Vivo. *PLOS ONE* **7**, e32094 (2012).

35. Shin, S.-Y. *et al.* An atlas of genetic influences on human blood metabolites. *Nat. Genet.* **46**,

543–550 (2014).

36. Draisma, H. H. M. *et al.* Genome-wide association study identifies novel genetic variants

contributing to variation in blood metabolite levels. *Nat. Commun.* **6**, 7208 (2015).

37. Calder, P. C. & Yaqoob, P. Glutamine and the immune system. *Amino Acids* **17**, 227–241

(1999).

38. Newsholme, P. Why Is L-Glutamine Metabolism Important to Cells of the Immune System in

Health, Postinjury, Surgery or Infection? *J. Nutr.* **131**, 2515S-2522S (2001).

39. Tian, C. *et al.* Genome-wide association and HLA region fine-mapping studies identify

susceptibility loci for multiple common infections. *Nat. Commun.* **8**, 599 (2017).

40. Xue, A. *et al.* Genome-wide association analyses identify 143 risk variants and putative

regulatory mechanisms for type 2 diabetes. *Nat. Commun.* **9**, 2941 (2018).

41. Vujkovic, M. *et al.* Discovery of 318 new risk loci for type 2 diabetes and related vascular

outcomes among 1.4 million participants in a multi-ancestry meta-analysis. *Nat. Genet.* **52**,

680–691 (2020).

42. Rühlemann, M. C. *et al.* Genome-wide association study in 8,956 German individuals

identifies influence of ABO histo-blood groups on gut microbiome. *Nat. Genet.* **53**, 147–155

(2021).

43. Bonder, M. J. *et al.* Genetic and epigenetic regulation of gene expression in fetal and adult

human livers. *BMC Genomics* **15**, 1–13 (2014).

44. Moon Jee-Young *et al.* Abstract P459: Milk Intake, Host LCT Genotype and Gut

Bifidobacteria in Relation to Obesity: Results From the Hispanic Community Health

Study/Study of Latinos (HCHS/SOL). *Circulation* **141**, AP459–AP459 (2020).

45. Arnolds, K. L., Martin, C. G. & Lozupone, C. A. Blood type and the microbiome- untangling a

complex relationship with lessons from pathogens. *Curr. Opin. Microbiol.* **56**, 59–66 (2020).

46. Mäkivuokko, H. *et al.* Association between the ABO blood group and the human intestinal

microbiota composition. *BMC Microbiol.* **12**, 94 (2012).

47. Liu, X. *et al.* Inter-determination of blood metabolite levels and gut microbiome supported by

Mendelian randomization. *bioRxiv* 2020.06.30.181438 (2020)

doi:10.1101/2020.06.30.181438.

48. Motta, V., Luise, D., Bosi, P. & Trevisi, P. Faecal microbiota shift during weaning transition

in piglets and evaluation of AO blood types as shaping factor for the bacterial community

profile. *PloS One* **14**, e0217001 (2019).

49. An ancient deletion in the ABO gene affects the composition of the porcine microbiome by

altering intestinal N-acetyl-galactosamine concentrations | bioRxiv.

https://www.biorxiv.org/content/10.1101/2020.07.16.206219v1.

50. Chen, L. *et al.* The long-term genetic stability and individual specificity of the human gut

microbiome. *Cell* **184**, 2302-2315.e12 (2021).

51. Bersaglieri, T. *et al.* Genetic Signatures of Strong Recent Positive Selection at the Lactase

Gene. *Am. J. Hum. Genet.* **74**, 1111–1120 (2004).

52. Auer, P. L. *et al.* Imputation of exome sequence variants into population- based samples

and blood-cell-trait-associated loci in African Americans: NHLBI GO Exome Sequencing

Project. *Am. J. Hum. Genet.* **91**, 794–808 (2012).

53. Ségurel, L. *et al.* The ABO blood group is a trans-species polymorphism in primates. *Proc.*

Natl. Acad. Sci. **109**, 18493–18498 (2012).

54. Band, G. *et al.* Insights into malaria susceptibility using genome-wide data on 17,000

individuals from Africa, Asia and Oceania. *Nat. Commun.* **10**, 5732 (2019).

55. Barua, D. & Paguio, A. S. ABO blood groups and cholera. *Ann. Hum. Biol.* **4**, 489–492

(1977).

56. Watanabe, K. *et al.* A global overview of pleiotropy and genetic architecture in complex

traits. *Nat. Genet.* **51**, 1339–1348 (2019).

57. Naitza, S. *et al.* A Genome-Wide Association Scan on the Levels of Markers of Inflammation

in Sardinians Reveals Associations That Underpin Its Complex Regulation. *PLOS Genet.* **8**,

e1002480 (2012).

58. Suzuki, T. A. & Ley, R. E. The role of the microbiota in human genetic adaptation. *Science*

370, (2020).

59. Parks, D. H. *et al.* A complete domain-to-species taxonomy for Bacteria and Archaea. *Nat.*

Biotechnol. **38**, 1079–1086 (2020).

60. Wade, K. H. & Hall, L. J. Improving causality in microbiome research: can human genetic

epidemiology help? *Wellcome Open Res.* **4**, 199 (2020).

61. Pirastu, N. *et al.* Using genetics to disentangle the complex relationship between food

choices and health status. *bioRxiv* 829952 (2019) doi:10.1101/829952.

62. Wang, C. *et al.* High-Salt Diet Has a Certain Impact on Protein Digestion and Gut

Microbiota: A Sequencing and Proteome Combined Study. *Front. Microbiol.* **8**, (2017).

63. Culligan, E. P., Marchesi, J. R., Hill, C. & Sleator, R. D. Combined metagenomic and

phenomic approaches identify a novel salt tolerance gene from the human gut microbiome.

Front. Microbiol. **5**, (2014).

64. Zeevi, D. *et al.* Personalized Nutrition by Prediction of Glycemic Responses. *Cell* **163**,

1079–1094 (2015).

65. Zhong, V. W. *et al.* A genome-wide association study of bitter and sweet beverage

consumption. *Hum. Mol. Genet.* **28**, 2449–2457 (2019).

66. Bjørkhaug, S. T. *et al.* Characterization of gut microbiota composition and functions in

patients with chronic alcohol overconsumption. *Gut Microbes* **10**, 663–675 (2019).

67. Sanna, S. *et al.* Common variants in the GDF5-UQCC region are associated with variation

in human height. *Nat. Genet.* **40**, 198–203 (2008).

68. Weedon, M. N. *et al.* A common variant of HMGA2 is associated with adult and childhood

height in the general population. *Nat. Genet.* **39**, 1245–1250 (2007).

69. Loos, R. J. F. *et al.* Common variants near MC4R are associated with fat mass, weight and

risk of obesity. *Nat. Genet.* **40**, 768–775 (2008).

70. Tigchelaar, E. F. *et al.* Cohort profile: LifeLines DEEP, a prospective, general population

cohort study in the northern Netherlands: study design and baseline characteristics. *BMJ*

Open **5**, e006772 (2015).

71. Scholtens, S. *et al.* Cohort Profile: LifeLines, a three-generation cohort study and biobank.

Int. J. Epidemiol. **44**, 1172–1180 (2015).

72. the Haplotype Reference Consortium *et al.* A reference panel of 64,976 haplotypes for

genotype imputation. *Nat. Genet.* **48**, 1279–1283 (2016).

73. Lopera Maya, E. A. *et al.* Lack of Association Between Genetic Variants at ACE2 and

TMPRSS2 Genes Involved in SARS-CoV-2 Infection and Human Quantitative Phenotypes.

Front. Genet. **11**, (2020).

74. The 1000 Genomes Project Consortium. A global reference for human genetic variation.

Nature **526**, 68–74 (2015).

75. Francioli, L. C. *et al.* Whole-genome sequence variation, population structure and

demographic history of the Dutch population. *Nat. Genet.* **46**, 818–825 (2014).

76. Tigchelaar, E. F. *et al.* An introduction to LifeLines DEEP: study design and baseline

characteristics. *bioRxiv* 009217 (2014) doi:10.1101/009217.

77. Truong, D. T. *et al.* MetaPhlAn2 for enhanced metagenomic taxonomic profiling. *Nat.*

Methods **12**, 902–903 (2015).

78. Franzosa, E. A. *et al.* Species-level functional profiling of metagenomes and

metatranscriptomes. *Nat. Methods* **15**, 962–968 (2018).

79. Suzek, B. E. *et al.* UniRef clusters: a comprehensive and scalable alternative for improving

sequence similarity searches. *Bioinforma. Oxf. Engl.* **31**, 926–932 (2015).

80. Buchfink, B., Xie, C. & Huson, D. H. Fast and sensitive protein alignment using DIAMOND.

Nat. Methods **12**, 59–60 (2015).

81. Siebelink, E., Geelen, A. & de Vries, J. H. M. Self-reported energy intake by FFQ compared

with actual energy intake to maintain body weight in 516 adults. *Br. J. Nutr.* **106**, 274–281

(2011).

82. Willett, W. C. *Nutritional Epidemiology. Nutritional Epidemiology* (Oxford University Press).

83. Zheng, X. & Davis, J. W. SAIGEgds—an efficient statistical tool for large-scale PheWAS

with mixed models. *Bioinformatics* doi:10.1093/bioinformatics/btaa731.

84. Genomewide Association Study of Severe Covid-19 with Respiratory Failure. *N. Engl. J.*

Med. **383**, 1522–1534 (2020).

85. Chang, C. C. *et al.* Second-generation PLINK: rising to the challenge of larger and richer

datasets. *GigaScience* **4**, 7 (2015).

86. Elsworth, B. *et al.* The MRC IEU OpenGWAS data infrastructure. *bioRxiv*

2020.08.10.244293 (2020) doi:10.1101/2020.08.10.244293.

87. Hemani, G. *et al.* The MR-Base platform supports systematic causal inference across the

human phenome. *eLife* **7**, e34408 (2018).

88. Burgess, S. & Thompson, S. G. *Mendelian Randomization Methods for Using Genetic*

Variants in Causal Estimation. (CRC Press, 2015).

89. Bowden, J., Davey Smith, G. & Burgess, S. Mendelian randomization with invalid

instruments: effect estimation and bias detection through Egger regression. *Int. J.*

Epidemiol. **44**, 512–525 (2015).

90. Teslovich, T. M. *et al.* Biological, clinical and population relevance of 95 loci for blood lipids.

Nature **466**, 707–713 (2010).

1038

Figures

Figure 1. Genome-wide association scan results.

Manhattan plot of host genomic associations to bacterial taxa and bacterial pathway
abundances with at least one genome-wide significant association ($p < 5\text{x}10^{-8}$). Y-axis shows
the -log10 transformation of the association p-value observed at each tested variant. X-axis
shows the genomic position of variants. The thresholds of study-wide ($p = 1.89\text{x}10^{-10}$) and
genome-wide ($p = 5\text{x}10^{-8}$) significance are shown as horizontal red and blue lines, respectively.
Independent SNP☐trait associations reaching genome-wide significance are listed in the tables
and labelled on the Manhattan plot. The colors of associated hits indicate whether they
represent an association with taxonomy (orange) or a bacterial pathway (blue).

1049

1050

1051

1052

Figure 2. Association at the LCT locus and interaction with lactose intake.

Comparison of f. Bifidobacteriaceae relative abundance a) between groups of Lactase

Persistent (LP, rs182549 C/T or T/T) and Lactose Intolerant (LI, rs182549 C/C) participants and

b) stratified among individuals with low or high daily lactose intake levels. Lactose intake was

corrected for daily calorie consumption. Relative abundance was natural log–transformed and

adjusted by age and sex. Distribution is displayed with violin plots, where each inner box

represents the first and third quartiles of the distribution and the middle line represents the

median. Lactose intake levels were defined as low if < first quartile and high if ≥ first quartile.

Lactose intake was only available for 5,801 of the 6,809 LP samples and for 376 of the 443 LI

samples. *P*-values were obtained with a two-sided Wilcoxon rank test. Species in the same

family shared similar distributions, although the difference in distribution within the LI group is

not significant (**Supplementary Figure 2**).

1065

1066

1067

Figure 3. Association with blood types and interaction with *FUT2*.

Relative abundances of microbiome features found to be associated with SNPs at the ABO

locus are compared between the inferred blood types (O, A, B and AB) with O as reference

group (panels **a**, **b** and **c**) and between secretors (light blue) and non-secretors (dark blue)

stratified again by the blood types grouped by the presence (A/B/AB) or absence (O) of terminal

sugars (panels **d**, **e** and **f**). The natural log–transformed relative abundance, adjusted by age

and sex, and is shown using a violin plot in which the inner box represents the first and third

quartiles of the distribution and the middle line represents the median. In the top panels, the

distribution across all blood types is also significantly different for all features (ANOVA: $p =$

6.7×10^{-14} for species *B. Bifidum*, $p = 5.4 \times 10^{-9}$ for *LACTOSECAT-PWY* and $p = 1.4 \times 10^{-13}$ for

genus *Collinsella*). *P*-values are obtained from a two-sided Wilcoxon ranked test. S.: secretors,

N.s.: non-secretors, ns: $p > 0.05$, * $p \leq 0.05$, ** $p \leq 0.01$, *** $p \leq 0.001$, **** $p \leq 0.0001$.

1080

1081

1082

Figure 4. Weighted Spearman correlation between estimated heritability and number of suggestive loci.

Each point represents one taxon or pathway. Bold red line represents the locally estimated scatterplot smoothing fit, with orange shaded bands indicating the 95% regression confidence intervals. The number of independent suggestive mbQTLs was obtained using LD pruning (**Methods**). **a)** Weighted Spearman correlation of number of suggestive mbQTLs and family-based heritability for microbiome taxa. **b)** Weighted Spearman correlation of number of suggestive mbQTLs and family-based heritability for microbiome functional pathways.

1095

1096

1097

Figure 5. Power analysis taking into account bacterial presence levels

a) Statistical power (at alpha = 1×10^{-10}) in our dataset for SNPs at the *ABO* and *LCT* loci,

considering the associated taxonomies. 80% detection power is shown by the red horizontal

line. **b**) Power (y-axis) to detect an effect at significance level alpha = 1×10^{-10} depending on the

variance explained by a genetic variant (x-axis). Colored lines distinguish power under different

levels of bacterial or pathway prevalence, which reduces effective sample size. **c**) Sample size

needed to identify an association at 80% power level across different bacterial presence levels

and alpha = 1×10^{-10}. Horizontal dashed line indicates the sample size of the DMP cohort.

1106

1107

Locus ID	Genes (Variant function)	id	chr	pos	pval	⬤ Trait
1	*MCM6 (intronic);*	rs182549	2	136616754	5.86E-15	family Bifidobacteriaceae
1	*MCM6 (intronic);*	rs182549	2	136616754	5.91E-15	order Bifidobacteriales
1	*MCM6 (intronic);*	rs182549	2	136616754	8.59E-15	genus Bifidobacterium
2	*ABO (intronic)*	rs8176645	9	136149098	5.54E-14	species Bifidobacterium bifidum
1	*DARS1 (intronic)*	rs6754311	2	136707982	7.06E-14	species Bifidobacterium adolescentis
1	*MCM6 (intronic);*	rs182549	2	136616754	2.32E-13	class Actinobacteria
1	*MCM6 (intronic);*	rs182549	2	136616754	2.32E-13	phylum Actinobacteria
2	*ABO (intronic)*	rs550057	9	136146597	9.34E-11	genus Collinsella
2	*ABO (intronic)*	rs550057	9	136146597	3.01E-09	family Coriobacteriaceae
2	*ABO (intronic)*	rs550057	9	136146597	3.03E-09	order Coriobacteriales
2	*ABO (intronic)*	rs550057	9	136146597	2.05E-08	species Collinsella aerofaciens
3	*SGCZ (intron)*	rs28450200	8	15070111	1.37E-08	sècoes Alistipes sp. AP11
4	*COL23A1 (intron)*	rs73344869	5	177746345	2.10E-08	family Rikenellaceae
4	*COL23A1 (intron)*	rs73344869	5	177746345	2.14E-08	genus Alistipes
1	*RH3DM1 (missense)*	rs1446585	2	136407479	3.18E-08	species Bifidobacterium longum
5	*TNFAIP2 (intronic); LINC0067 (intronic)*	rs2282037	14	103588515	4.10E-08	class Negativicutes
5	*TNFAIP2 (intronic); LINC0067 (intronic)*	rs2282037	14	103588515	4.10E-08	order Selenomonadales
6	DMD (intron)	rs921896	X	32213962	4.15E-08	species Escherichia unclassified

Locus ID	Genes (Variant function)	id	chr	pos	pval	⬤ Trait
2	*ABO (intronic)*	rs507666	9	136149399	5.38E-15	LACTOSECAT-PWY: lactose and galactose degradation I
7	intergenic	rs7513344	1	4220095	1.79E-09	PWY-6151: S-adenosyl L methionine cycle I
8	*AC105942.1 (intronic)*	rs10159356	1	48103139	7.52E-09	PWY-REDCITCYC: TCA cycle VIII (helicobacter)
9	*WWOX (intron)*	rs9927590	16	78318743	9.28E-09	PWY-5088: L-glutamate degradation VIII to propanoate
10	*SALL4 (intron)*	rs79676117	20	50415961	1.16E-08	PWY-7234: inosine-5'-phosphate biosynthesis III
11	*AC010149.1 (intron)*	rs11898768	2	231419306	2.00E-08	PWY-6708: ubiquinol-8 biosynthesis
12	closest genes: INSIG and BLACE	rs36182340	7	155118470	2.00E-08	P105-PWY: TCA cycle IV (2-oxoglutarate decarboxylase)
13	*LINC02851 (upstream)*	rs820496	9	6703997	2.43E-08	ASPASN-PWY: superpathway of L-aspartate and L-asparagine biosynthesis
14	*SYNE2 (intronic)*	rs17824268	14	64467727	2.63E-08	PWY-6612: superpathway of tetrahydrofolate biosynthesis
15	*NUMB (intronic)*	rs148280750	14	73763641	2.93E-08	PYRIDNUCSYN-PWY: NAD biosynthesis I from aspartate
16	closest gene: TSC22D2	rs1584586	3	150194351	3.47E-08	PWY-841: superpathway of purine nucleotides de-novo biosynthesis I
17	RCAN2 (intron)	rs59657730	6	46422725	3.50E-08	GLYCOCAT.PWY: glycogen degradation I (bacterial)
18	*closest genes: CNN3 and ALG14*	rs12137024	1	95409858	3.53E-08	HISDEG-PWY: L-histidine degradation I
19	*AL451048.1 (intron)*	rs11795855	X	140027569	3.59E-08	ENTBACSYN-PWY: enterobactin biosynthesis
20	CDK11B	rs11486024	1	1655474	3.89E-08	TRPSYN PWY: L-tryptophan biosynthesis
21	SLC25A37 (intron)	rs2137304	8	23415240	4.26E-08	PWY-6353: purine nucleotides degradation II (aerobic)
22	closest gene: RPRD2	rs78778914	1	150449701	4.34E-08	HISTSYN PWY: L-histidine biosynthesis
23	GRAMD1B	rs2714053	11	123312063	4.85E-08	PWY-6147: 6-hydroxymethyl dihydropterin diphosphate biosynthesis I
24	closest genes: IL4 and KIF3A (eQTL)	rs11242123	5	132025487	4.89E-08	PWY-7209: superpathway of pyrimidine ribonucleosides degradation