Detecting *de novo* mitochondrial mutations in angiosperms with highly divergent evolutionary rates

Amanda K. Broz*[,1], Gus Waneka*[,1], Zhiqiang Wu[†,*,1], Matheus Fernandes Gyorfy*, Daniel B. Sloan*[,2]

*Department of Biology, Colorado State University, Fort Collins, CO 80523

[†]Shenzhen Branch, Guangdong Laboratory for Lingnan Modern Agriculture, Genome Analysis Laboratory of the Ministry of Agriculture, Agricultural Genomics Institute at Shenzhen, Chinese Academy of Agricultural Sciences, 518120 Shenzhen, China

[1]These authors contributed equally to this work.

[2]Author for Correspondence:
Daniel B. Sloan: dan.sloan@colostate.edu
Colorado State University
1878 Campus Delivery
Fort Collins, CO 80523
970.491.2256

ABSTRACT

Although plant mitochondrial genomes typically show low rates of sequence evolution, levels of divergence in certain angiosperm lineages suggest anomalously high mitochondrial mutation rates. However, *de novo* mutations have never been directly analyzed in such lineages. Recent advances in high-fidelity DNA sequencing technologies have enabled detection of mitochondrial mutations when still present at low heteroplasmic frequencies. To date, these approaches have only been performed on a single plant species (*Arabidopsis thaliana*). Here, we apply a high-fidelity technique (Duplex Sequencing) to multiple angiosperms from the genus *Silene*, which exhibits extreme heterogeneity in rates of mitochondrial sequence evolution among close relatives. Consistent with phylogenetic evidence, we found that *S. latifolia* maintains low mitochondrial variant frequencies that are comparable to previous measurements in *Arabidopsis*. *Silene noctiflora* also exhibited low variant frequencies despite high levels of historical sequence divergence, which supports other lines of evidence that this species has reverted to lower mitochondrial mutation rates after a past episode of acceleration. In contrast, *S. conica* showed much higher variant frequencies in mitochondrial (but not in plastid) DNA, consistent with an ongoing bout of elevated mitochondrial mutation rates. Moreover, we found an altered mutational spectrum in *S. conica* heavily biased towards AT→GC transitions. We also observed an unusually low number of mitochondrial genome copies per cell in *S. conica*, potentially pointing to reduced opportunities for homologous recombination to accurately repair mismatches in this species. Overall, these results suggest that historical fluctuations in mutation rates are driving extreme variation in rates of plant mitochondrial sequence evolution.

INTRODUCTION

Plant mitochondrial genomes exhibit dramatic variation in rates of nucleotide substitution. Early molecular evolution studies (WOLFE *et al.* 1987; PALMER AND HERBON 1988) established that mitochondrial rates in most angiosperms are about an order of magnitude lower than in the nucleus (DROUIN *et al.* 2008), which contrasts with the rapid evolution of mitochondrial DNA (mtDNA) in many other eukaryotic lineages (BROWN *et al.* 1979; SMITH AND KEELING 2015; LAVROV AND PETT 2016). However, subsequent phylogenetic surveys have identified a number of angiosperms with exceptionally high levels of mtDNA sequence divergence (PALMER *et al.* 2000; CHO *et al.* 2004; PARKINSON *et al.* 2005; MOWER *et al.* 2007; SLOAN *et al.* 2009; SKIPPINGTON *et al.* 2015; ZERVAS *et al.* 2019). As such, despite the relatively recent origin and diversification of angiosperms (BARBA-MONTOYA *et al.* 2018), mitochondrial substitution rates are estimated to span a remarkable 5000-fold range across this group (RICHARDSON *et al.* 2013). At one extreme, *Magnolia stellata* has a rate of only ~0.01 synonymous substitutions per site per billion years (SSB), while certain *Plantago* and *Silene* species have estimated rates of >50 SSB (MOWER *et al.* 2007; SLOAN *et al.* 2012a; RICHARDSON *et al.* 2013). In some cases, rate accelerations appear to be short-lived with bursts of sequence divergence inferred for internal branches on phylogenetic trees followed by reversions to slower rates of sequence evolution (CHO *et al.* 2004; PARKINSON *et al.* 2005; SLOAN *et al.* 2009; SKIPPINGTON *et al.* 2017).

The angiosperm genus *Silene* (Caryophyllaceae) is a particularly interesting model for the study of mitochondrial genome evolution and substitution rate variation. This large genus comprises approximately 850 species (JAFARI *et al.* 2020) and exhibits rate accelerations that rival the magnitude of other extreme cases in genera such as *Plantago*, *Pelargonium*, and *Viscum* (MOWER *et al.* 2007; SKIPPINGTON *et al.* 2015). Moreover, *Silene* is distinct because the observed rate accelerations appear to have occurred very recently (<10 Mya), such that close relatives within the genus exhibit radically different substitution rates (Figure 1) (MOWER *et al.* 2007; SLOAN *et al.* 2009; RAUTENBERG *et al.* 2012). For example, *S. latifolia* (section *Melandrium*) has retained a substitution rate that is roughly in line with the low levels of sequence divergence found in most angiosperms. In contrast, species such as *S. noctiflora* and *S. conica* represent lineages (sections *Elisanthe* and *Conoimorpha*, respectively) that are within the same subgenus as section *Melandrium* but have highly divergent mtDNA sequence. These large differences among close relatives in *Silene* have enabled comparative approaches to investigate the evolutionary consequences of accelerated substitution rates for mitochondrial genome architecture (SLOAN *et al.* 2012a), RNA editing (SLOAN *et al.* 2010), mitonuclear coevolution (SLOAN *et al.* 2014; HAVIRD *et al.* 2015; HAVIRD *et al.* 2017), and mitochondrial physiology (HAVIRD *et al.* 2019; WEAVER *et al.* 2020). Notably, accelerated species such as *S. noctiflora* and *S. conica* also exhibit massively expanded mitochondrial genomes that

58 have been fragmented into dozens of circularly mapping chromosomes (SLOAN *et al.* 2012a).

59 However, the mechanisms that cause increased mitochondrial substitution rates in some *Silene*

60 species remain unclear (HAVIRD *et al.* 2017).

61   Previous studies on *Silene* and other angiosperms have generally concluded that elevated

62 rates of mitochondrial sequence evolution reflect increased mutation rates rather than changes in

63 selection pressure (CHO *et al.* 2004; PARKINSON *et al.* 2005; MOWER *et al.* 2007; SLOAN *et al.* 2009;

64 SLOAN *et al.* 2012a; SKIPPINGTON *et al.* 2015). Increases are especially pronounced at synonymous

65 sites, which are thought to experience very limited effects of selection in plant mtDNA (SLOAN AND

66 TAYLOR 2010; WYNN AND CHRISTENSEN 2015). Likewise, the ratio of nonsynonymous to synonymous

67 substitutions (d_N/d_S) is still low in accelerated species (SLOAN *et al.* 2012a), indicating that purifying

68 selection on mitochondrial genes remains strong. In *Silene* species with rapidly evolving mtDNA,

69 there does not appear to be a genome-wide increase in synonymous substitution rates in the

70 nucleus or plastids (RAUTENBERG *et al.* 2012; SLOAN *et al.* 2012b), suggesting that the accelerated

71 point mutation rates are largely specific to the mitochondria. A number of mechanisms have been

72 hypothesized to explain cases of increased mitochondrial mutation rates. However, thus far, all

73 inferences about variation in plant mitochondrial divergence are based on long-term patterns of

74 sequence divergence across species rather than direct detection and analysis of *de novo* mutations,

75 making it difficult to investigate the actual role of mutation.

76   Recent advances in high-fidelity DNA sequencing have improved our ability to distinguish

77 signal from noise and detect *de novo* mutations (SALK *et al.* 2018; SLOAN *et al.* 2018). In particular,

78 Duplex Sequencing (SCHMITT *et al.* 2012; KENNEDY *et al.* 2014) has been used to obtain error rates

79 as low as ~2 × 10^{-8} per bp, allowing for accurate identification of mitochondrial variants that are still

80 present at low heteroplasmic frequencies within tissue samples (KENNEDY *et al.* 2013; AHN *et al.*

81 2016; HOEKSTRA *et al.* 2016; ARBEITHUBER *et al.* 2020; WU *et al.* 2020a). Duplex Sequencing works

82 by attaching random barcodes to each original DNA fragment and independently sequencing the two

83 complementary DNA strands from that fragment multiple times each to produce a highly accurate

84 double-stranded consensus. To date, *Arabidopsis thaliana* is the only plant system to have been

85 analyzed with this method (WU *et al.* 2020a). Here, we apply Duplex Sequencing to detect *de novo*

86 mitochondrial and plastid mutations in three *Silene* species previously inferred to have dramatically

87 different histories of mitochondrial mutation.

90 **MATERIALS AND METHODS**

92 **Plant lines and growth conditions**

A single family of full siblings was grown for each of three *Silene* species. Families were taken from

lines with previously sequenced mitochondrial and plastid genomes: *S. latifolia* UK2600, *S. noctiflora*

OSR, and *S. conica* ABR (SLOAN *et al.* 2012a; SLOAN *et al.* 2012b). The latter two species are

hermaphroditic and readily self-fertilize, so families were derived from selfed parents. In contrast, *S.*

latifolia is dioecious and exhibits substantial inbreeding depression from full-sib crosses (TEIXEIRA *et*

al. 2009). Therefore, we crossed a female from the UK2600 line with a male from a different line

(Kew 32982, obtained from the Kew Gardens Millenium Seed Bank) to produce the full-sib family

used in this study. Seeds were germinated on ProMix BX soil mix and grown in a growth room under

short-day conditions (10-hr/14-hr light/dark cycle). All three species were grown for approximately

seven weeks to produce sufficiently large rosettes for organelle DNA extractions.

Organelle DNA extractions and Duplex Sequencing

Each full-sib family was divided into three biological replicates, consisting of approximately 30-40

individual plants per replicate. A total of 35 g of rosette tissue was harvested from each replicate and

used for simultaneous extraction of enriched mtDNA and cpDNA as described previously (WU *et al.*

2020a; WU *et al.* 2020b). Mitochondria were isolated by differential centrifugation followed by DNase

I treatment to remove contaminating DNA not protected by intact mitochondrial membranes.

Chloroplasts were isolated on discontinuous sucrose gradients. Following DNA extraction, Duplex

Sequencing libraries were constructed as described previously (WU *et al.* 2020a). These libraries

were multiplexed and sequenced with 2×150 bp reads on an Illumina NovaSeq 6000 S4 Lane

(NovoGene).

Shotgun Illumina sequencing of total-cellular DNA samples for *k*-mer database construction

Detection of low-frequency mitochondrial heteroplasmies presents a number of technical challenges

that can lead to false positives. In particular, plant nuclear genomes harbor numerous insertions of

mtDNA and cpDNA fragments (which are known as "numts" and "nupts", respectively) that can differ

in sequence from the mitochondrial or plastid genomes because they accumulate mutations over

time (HUANG *et al.* 2005; NOUTSOS *et al.* 2005; LOUGH *et al.* 2008; HAZKANI-COVO *et al.* 2010;

MICHALOVOVÁ *et al.* 2013). Therefore, contaminating nuclear DNA in mtDNA and cpDNA samples

can mimic low-frequency *de novo* mitochondrial and plastid mutations. Because numts and nupts

often closely resemble the mitochondrial and plastid genome sequence, they can be problematic to

accurately assemble in nuclear genome sequencing projects even in high quality reference

assemblies (STUPAR *et al.* 2001). As such, it can be difficult to identify and filter out numt- and nupt-

derived variants based only on reference genome assemblies. We have found that an effective

alternative is to use raw reads from total-cellular sequencing datasets to generate a database of

counts of all sequences of a specified length *k*, which are referred to as *k*-mers (WU *et al.* 2020a).

The premise of this approach is that variants associated with numts or nupts should be abundant in

total-cellular sequencing datasets (as quantified by counting corresponding k-mers in the raw reads)

and match the level expected for other nuclear sequences. However, for reasons discussed below

(see Results), we expect this filtering approach to be more reliable for numts than for nupts.

To generate a k-mer database for each species we extracted total-cellular DNA from mature

leaf tissue using the Qiagen DNeasy kit. For *S. noctiflora* and *S. conica*, the sampled individuals

were from the same inbred line as the full-sib family used for Duplex Sequencing but separated by at

least three generations. In contrast to the inbred history of the *S. noctiflora* and *S. conica* lines, *S.*

latifolia is expected to be highly heterozygous, including for numt and nupt alleles. Therefore, in

order to capture all numt and nupt alleles that might be segregating in the *S. latifolia* full-sib family,

we generated total-cellular samples for both parents that were crossed to generate the family.

Sampling the male parent in this *S. latifolia* cross also provides reads from its actual mitochondrial

genome sequence, which can identify potential false positives resulting from low-frequency paternal

transmission of mtDNA. Such "paternal leakage" has been identified in outcrossing *Silene* species

(MCCAULEY *et al.* 2005; BENTLEY *et al.* 2010).

Illumina libraries were generated from each total-cellular DNA sample using the New

England Biolabs NEB FS II Ultra kit with approximately 100 ng of input DNA, a 20 min fragmentation

time, and eight cycles of PCR. The *S. latifolia* and *S. noctiflora* libraries were multiplexed and

sequenced on the same NovaSeq 6000 lane as the above Duplex Sequencing libraries*.* The *S.*

conica library was sequenced separately with 2×150 bp reads on an Illumina HiSeq 4000 lane. The

resulting raw reads were used to generate databases of k-mer counts (k = 39 bp) with KMC v3.0.0

(KOKOT *et al.* 2017).

Duplex Sequencing data analysis and variant calling

Duplex Sequencing datasets were processed with a previously published pipeline

(https://github.com/dbsloan/duplexseq) (WU *et al.* 2020a). This pipeline uses Duplex Sequencing

barcodes (i.e., unique molecular identifiers) to group raw reads into families corresponding to the two

complementary strands of an original DNA fragment, requiring a minimum of three reads for each

strand. After trimming barcodes and linker sequences, the consensus sequences for each double-

stranded family were mapped to the reference mitochondrial and plastid genomes for the

corresponding species. Because of known sequencing artefacts associated with end repair and

adapter ligation (KENNEDY *et al.* 2014), the 10 bp at the end of each read were excluded when

identifying variants and calculating sequencing coverage.

Reads that contained a single mismatch relative to the reference sequences were used to

identify single nucleotide variants (SNVs). Variants with a *k-mer* count of 10 or greater in the

corresponding k-mer database were excluded as likely numts or nupts. This k-mer filtering also

ensured elimination of false positives due to paternal leakage from the *S. latifolia* male parent or

errors in the published reference sequence. The latter is an important concern because the

published *S. conica* mitochondrial genome sequence is a draft assembly due to its extreme

repetitiveness (SLOAN *et al.* 2012a). Accordingly, we also excluded variants if the corresponding

reference sequence was not detected in the *k*-mer database to account for sites with ambiguities

(Ns) or possible errors in the reference. Variants were also filtered using the pipeline's

--recomb_check option, which identifies SNVs that can be explained by recombination between non-

identical repeat sequences within the mitochondrial genome rather than *de novo* point mutations

(DAVILA *et al.* 2011). Finally, the pipeline's --contam_check option was used to provide the reference

genome sequences from the other *Silene* species for filtering of variants arising from contamination

between multiplexed libraries.

To calculate SNV frequencies for each library, the number of reads containing an SNV (after

filtering) was divided by the total bp of duplex consensus sequence mapped to the genome.

Differences in SNV frequency among species were tested with a one-way ANOVA implemented,

using the aov function in R v3.6.3 followed by *post hoc* pairwise comparisons with the TukeyHSD

function.

For calling mitochondrial SNVs, we were able to use the cpDNA Duplex Sequencing libraries

to increase our mitochondrial genome coverage because they contained a substantial amount of

contaminating mtDNA reads resulting from incomplete enrichment of cpDNA. Importantly, being able

to use mitochondrial reads from the cpDNA library for *S. conica* biological replicate 2 was key

because sequencing of the mtDNA library for that replicate failed (see Results). However, we did not

do the reverse (use mtDNA libraries to supplement plastid datasets) because the mtDNA libraries

were expected to have a far higher ratio of nuclear to plastid DNA than the cpDNA libraries,

exacerbating the challenges associated with filtering nupts.

Analysis of organelle genome copy number

To estimate the copy number of mitochondrial and plastid genomes based on the total-cellular

sequencing datasets, raw reads were trimmed with cutadapt v1.16 (MARTIN 2011) to remove adapter

sequences with an error tolerance of 0.15 and to trim low-quality ends with a q20 threshold. Read

pairs with a minimum length of 50 bp each after trimming were retained. Trimmed reads were then

mapped to reference mitochondrial and plastid genomes for the corresponding species using bowtie

v2.2.23 (LANGMEAD AND SALZBERG 2012), and position-specific coverage data were extracted from

the resulting alignment (BAM/SAM) files. Average coverage was summarized for 2-kb windows tiled

across each organelle genome. To estimate the number of mitochondrial and plastid genomes per

nuclear genome copy, we assumed that the remaining unmapped reads were all nuclear. We used

the total length of these unmapped reads divided by the nuclear genome size of the corresponding

species to estimate the average nuclear genome coverage and obtain ratios of organelle to nuclear

coverage. Nuclear genome size estimates of 2.67, 2.78, and 0.93 Gb were used for *S. latifolia*, *S.*

noctiflora, and *S. conica,* respectively (WILLIAMS *et al.* 2020).

The above analysis of genome copy number identified a surprisingly low number of

mitochondrial genome copies in *S. conica*. To validate this unexpected result and compare

stoichiometry of mitochondrial, plastid and nuclear genomes across multiple tissue samples in *S.*

conica, we performed a droplet digital PCR (ddPCR) analysis. We used the same *S. conica* total-

cellular DNA extraction that was used for the Illumina shotgun sequencing. In addition, we harvested

leaf tissue from three individuals from a different batch of *S. conica* ABR plants. For both the original

sample and the newer samples, tissue was harvested from the largest rosette leaves. The original

sample was grown in a growth chamber under long-day conditions (16-hr/8-hr light/dark cycle) and

harvested after 28 days, whereas the newer samples were grown on light racks in a growth room

under short-day conditions (10-hr/14-hr light/dark cycle) and harvested after 40 days. The tissue

sampling also differed in that the entire rosette leaf was sampled for the newer replicates, whereas

only the distal half of the leaf was taken for the original DNA extraction. In both cases, DNA

extractions were performed with a Qiagen DNeasy Kit. A total of six ddPCR markers were

developed, with two each targeting the mitochondrial, plastid and, nuclear genomes (Table S1).

Each ddPCR reaction was set up in a 20 μl volume, containing Bio-Rad QX200 ddPCR EvaGreen

Supermix and 0.2 μM of each primer. For mitochondrial and nuclear markers, 5 ng of total-cellular

DNA was used as template. To avoid saturation with the higher-copy-number plastid makers, a 200-

fold dilution (25 pg) of the template DNA was used. The reaction volumes were then emulsified in

Bio-Rad QX200 Droplet Generation Oil for EvaGreen, using the Bio-Rad QX200 Droplet Generator.

PCR amplification was performed on a Bio-Rad C1000 Touch thermal cycler with an initial

incubation at 95 °C for 5 min, 40 cycles of 90 °C for 30 s and 60 °C for 1 min, followed by a 4 °C

incubation for 5 min and a final 90 °C incubation for 5 min before holding at 4 °C. After amplification,

droplets were analyzed on a Bio-Rad QX200 Droplet Reader, and the absolute copy number of each

PCR target was estimated based on a Poisson distribution in the Bio-Rad QuantaSoft package.

Mitochondrial:nuclear and plastid:nuclear ratios were calculated by dividing organellar marker copy

numbers by the mean of the estimates for the two nuclear markers for a given sample.

Data Availability

All duplex sequencing and shotgun Illumina sequencing reads were deposited to the NCBI

Sequence Read Archive (SRA) under BioProject PRJNA682809 (Table S2).

RESULTS

Duplex Sequencing yield and read mapping

Each of the Duplex Sequencing libraries produced between 88 M and 142 M read pairs (Table S2), with the exception of the *S. conica* mtDNA library for biological replicate 2, which was not sequenced due to an apparent loading error. Nevertheless, we were still able to calculate mitochondrial SNV frequencies for *S. conica* replicate 2 by taking advantage of contaminating mitochondrial reads in the enriched cpDNA library for that replicate. The large number of reads per library translated into sizeable single-stranded read families for construction double-stranded consensus sequences, with modal values of >10 reads per family for most libraries (Figure S1). The three species differed in their degree of enrichment in the mtDNA libraries. For both *S. noctiflora* and *S. conica*, an average of 79% of the sequences in the mtDNA libraries could be mapped to the reference mitochondrial genome, whereas only 30% of the sequences in the *S. latifolia* mitochondrial libraries mapped, indicating a much lower level of mtDNA enrichment (Table S2). For all three species, between 64% and 67% of sequences from the cpDNA libraries mapped to the plastid genome, with a substantial number of contaminating sequences mapping to the mitochondrial genome (8%, 27%, and 31% for *S. latifolia*, *S. noctiflora*, and *S. conica*, respectively). To make use of as much mitochondrial data as possible, we combined all mitochondrial-derived sequences from both sample types. After collapsing raw reads into duplex consensus sequences and mapping them to the reference genome, we obtained between 85 and 382 Mb of mapped mitochondrial sequence data for each replicate (based on combing coverage from both mtDNA and cpDNA libraries; Table S2). For plastid genome coverage, we relied solely on the cpDNA libraries, which yielded between 203 and 319 Mb of coverage per replicate (Table S2).

Three *Silene* species differ in their frequencies of mitochondrial SNVs but show little variation in plastid SNV frequencies

Using the variant calls from the duplex consensus sequences (File S1), we calculated the frequency of mitochondrial SNVs per mapped bp for each *Silene* replicate and compared those values to previously published estimates from *A. thaliana* (Wu *et al.* 2020a). After applying filtering criteria to exclude false positives associated with numts, low-frequency paternal transmission, chimeric recombination products, or errors in the reference sequence (see Materials and Methods), we found significant variation in (log-transformed) mitochondrial SNV frequency among species (one-way ANOVA, $p = 5.3 \times 10^{-6}$; Figure 2, Table S3).

The three biological replicates of *S. latifolia* had a mean mitochondrial SNV frequency of 1.7×10^{-7} per bp. *Silene latifolia* was selected for this study because it exhibited very little mitochondrial

rate acceleration in previous phylogenetic analyses, suggesting that it has retained the slow rate of

sequence evolution that is characteristic of most plant mitochondrial genomes (MOWER *et al.* 2007;

SLOAN *et al.* 2009). Accordingly, the *S. latifolia* estimate was very similar to our previous estimate of

1.8×10^{-7} per bp for the mitochondrial SNV frequency in wild type *A. thaliana* Col-0 (WU *et al.*

2020a), another plant species with a typically low rate of mitochondrial sequence evolution (MOWER

et al. 2007). In contrast to the low historical substitution rates in *S. latifolia* and *Arabidopsis*, *S.*

noctiflora exhibits highly accelerated mitochondrial sequence evolution since diverging from other

major lineages within the genus *Silene* (Figure 1). However, we did not find an elevated frequency of

mitochondrial SNVs in our Duplex Sequencing analysis of *S. noctiflora*, suggesting that this species

may have experienced a reversion to lower mutation rates. In fact, the mean SNV frequency in *S.*

noctiflora was 6.1×10^{-8} per bp, which was approximately 3-fold lower than in *S. latifolia* or *A.*

thaliana (Tukey's HSD *post hoc* test, $p = 0.01$ for both comparisons). The most noteworthy variation

among species was based on observed SNV frequencies in *S. conica*, representing another *Silene*

lineage with a history of rapid mitochondrial sequence divergence (Figure 1). The mean

mitochondrial SNV frequency in *S. conica* was 1.7×10^{-6} per bp, which was 9-fold to 27-fold higher

than in *A. thaliana* and the other two *Silene* species (Tukey's HSD post hoc test, $p < 0.0001$ for all

three comparisons). All of these SNV frequencies substantially exceed the noise threshold of ~$2 \times$

10^{-8} that we previously estimated for this protocol using *Escherichia coli* samples derived from single

colonies as "negative controls" (WU *et al.* 2020a).

Silene conica was also distinct in that a large proportion of the identified mitochondrial SNVs

(31.7%) were found in two or more biological replicates. Because our biological replicates

represented sets of individuals from the same full-sib family, variants that are shared among

replicates likely indicate SNVs that were already heteroplasmic in the parent and then inherited by

the offspring. In contrast, none of the identified mitochondrial SNVs in either *S. latifolia* or *S.*

noctiflora were present in multiple biological replicates. There is reason to expect that our *k*-mer

filtering may have introduced bias against detection of inherited heteroplasmies in *S. latifolia* (see

Discussion). Nevertheless, even without this filtering, only 3.7% of *S. latifolia* SNVs (and only 1.8%

of *S. noctiflora* SNVs) would be present in two or more libraries. Therefore, the filtering does not

appear to explain this observed difference between *S. conica* and the other *Silene* species.

Unlike in the mitochondrial genome, we found no evidence that *S. conica* has an elevated

frequency of SNVs in its plastid genome, as the three *Silene* species all exhibited similarly low

plastid SNV frequencies (Figure 2). We recommend that the estimates of the *Silene* plastid SNV

frequencies and spectra be interpreted cautiously because of the nuclear contamination in these

libraries and the fact that nupts are more difficult to reliably filter with our *k*-mer database than

numts. The challenge that nupts pose relates to the high coverage of true plastid DNA in our total-

cellular libraries (>10,000×). As such, even rare sequencing errors in total-cellular libraries have the

potential to occur repeatedly and exhibit sizeable counts in our *k*-mer database, which could lead to

exclusion of variant calls that are actually true *de novo* mutations. Nevertheless, we can confidently

conclude that *S. conica* does not exhibit a major increase in plastid SNV frequency. Even if we

performed no *k*-mer filtering whatsoever on the *S. conica* plastid samples, SNV frequencies would

only increase by 55% on average (Table S4), leaving them at a level that is still similar to that of the

other *Silene* species and more than an order of magnitude lower than the (filtered) mitochondrial

SNV frequencies in *S. conica* (Figure 2).

Variation in mitochondrial mutation spectra among *Silene* species and extreme GC bias in *S.

conica*

Previous analysis of low-frequency mitochondrial SNVs in rosette tissue from wild type *A. thaliana*

Col-0 (WU *et al.* 2020a) identified a mutation spectrum dominated by GC→AT transitions (Figure 2).

The slowly evolving *S. latifolia* mitochondrial genome exhibited a bias in this same direction with

57% of all observed SNVs being GC→AT transitions (Table S3). In contrast, *S. noctiflora* did not

show a similar bias. The low overall SNV frequency in *S. noctiflora* makes it difficult to precisely

estimate the mutation spectrum, but no single substitution type dominated, as GC→AT transitions,

AT→GC transitions, and GC→CG transversions all had similar frequencies in the observed set of

SNVs (Figure S2, Table S3). Once again, *S. conica* exhibited the most extreme departure from the

other species. Notably, the high rate in *S. conica* was not driven by an increased frequency of the

GC→AT transitions that dominate the spectrum of *A. thaliana* and *S. latifolia*. In fact, the frequency

of GC→AT transitions in *S. conica* was lower than in either of those species. Instead, the high

overall SNV frequency was largely the result of a massive increase in the frequency AT→GC

transitions, which account for 77% of the observed SNVs in *S. conica* (Figure 2; Table S3). This

species also exhibited a substantial increase in the frequency of AT→CG transversions (11% of

observed SNVs). As such, both of the dominant types of substitutions in the *S. conica* mitochondrial

mutation spectrum increase GC content, which is unusual because mutation spectra are generally

AT-biased (HERSHBERG AND PETROV 2010; HILDEBRAND *et al.* 2010; SLOAN AND WU 2014).

Unusually low mitochondrial genome copy number in *Silene conica* rosette tissue

By performing deep sequencing of total-cellular DNA to generate a *k*-mer database for variant

filtering, we were also able to estimate the relative copy number of mitochondrial, plastid, and

nuclear genomes in each of the three *Silene* species (Figures 3 and S3-S5, Table S5). We found

similar plastid genome copy numbers across species, with mean estimates of 378, 263, and 275

plastid genome copies per nuclear genome copy for *S. latifolia*, *S. noctiflora*, and *S. conica*,

respectively. If we assume that each cell is diploid and has not yet undergone DNA replication, the

plastid genome copy numbers per cell would be double those values, but that may be an

underestimate because many plants undergo extensive endoreduplication, in which the nuclear

genome replicates without subsequent cell divisions, resulting in cells with higher nuclear ploidies

(JOUBES AND CHEVALIER 2000). The number of mitochondrial genome copies was surprisingly low in

S. conica, with an average of only 0.38 copies per nuclear genome copy. In contrast, *S. latifolia* and

S. noctiflora had an average of 47.7 and 9.7 mitochondrial genome copies per nuclear genome

copy, respectively, which is more consistent with estimates from other plants (PREUTEN *et al.* 2010;

OLDENBURG *et al.* 2013; SHEN *et al.* 2019).

To validate the finding of extremely low mitochondrial genome copy number in *S. conica*, we

performed a ddPCR analysis with two markers each for the mitochondrial, plastid, and nuclear

genomes. We first analyzed DNA from the same extraction that was originally used for the total-

cellular Illumina shotgun sequencing, obtaining an estimate of 0.42 mitochondrial genomes per

nuclear genome, in close agreement with our estimate of 0.38 from the sequencing data. Based on

this ddPCR analysis, we also estimated that there were 737 copies of the plastid genome per

nuclear genome copy, which was 2.7-fold higher than our original estimate (Figure 3), possibly

indicating that our sequencing estimate was downwardly biased for plastid genome copy number.

We then analyzed leaf DNA collected from three new *S. conica* plants that were grown separately

from the originally sampled plant to assess whether the original DNA extraction was anomalous in

some way. These three new samples also produced extremely low estimates for the number of

mitochondrial genomes copies with a mean of 1.02 per nuclear genome copy (Figure 3 and Table

S6). Therefore, these new *S. conica* samples exhibited a small increase in the mitochondrial:nuclear

ratio relative to our original sample but still fell well below typical observations for plant cells.

DISCUSSION

The challenges of detecting *de novo* mutations in plant organelle genomes

High-fidelity techniques such as Duplex Sequencing (SCHMITT *et al.* 2012) have been key

innovations to address the challenge of detecting and quantifying rare mutations (SALK *et al.* 2018;

SLOAN *et al.* 2018), but some sources of false positives remain. The potential misidentification of

numts and nupts as *de novo* mutations was a particular concern in this study. High quality nuclear

genome assemblies are not available for *Silene*, so it is not possible to use a reference genome to

identify and filter numt- and nupt-associated variants. Moreover, our mitochondrial and plastid DNA

preparations only reached moderate levels of enrichment, leaving substantial amounts of

contaminating nuclear DNA (Table S2). Our approach to avoid erroneous numt and nupt variant

calls was based on filtering using a *k*-mer database derived from total-cellular sequencing (see

Materials and Methods), but there are some concerns about balancing false positives and false
negatives that should be considered.

In particular, there is a risk that *k*-mer filtering may exclude true heteroplasmies if they are
shared between the total-cellular sample used to generate the *k*-mer database and the family used
for Duplex Sequencing. We reduced the risk of this in *S. conica* and *S. noctiflora* by using individuals
separated by at least three generations for constructing the *k*-mer database. Therefore, low-
frequency heteroplasmies would have to have been maintained across multiple generations to lead
to improper exclusion of true mitochondrial variants. Although transmission of heteroplasmies across
generations certainly occurs (MCCAULEY 2013; ZHANG *et al.* 2018; MANDEL *et al.* 2020), the
segregational loss of low-frequency variants should greatly reduce the magnitude of this problem. In
contrast, for the outcrossing species *S. latifolia*, we used both parents of the full-sib family to
construct total-cellular *k*-mer databases in order to screen for possible numts and nupts in all
contributing nuclear haplotypes. As such, variants that were heteroplasmic in the *S. latifolia* mother
and transmitted to her offspring might have been improperly filtered because of their presence in the
total-cellular *k*-mer database, resulting in a potential downward bias on our estimate of the overall
frequency of SNVs in *S. latifolia*.

Despite the uncertainty that this introduces into SNV frequency estimates, we feel that the
major qualitative conclusions of this study are robust to the challenges of numt and nupt artefacts.
For example, the finding that *S. noctiflora* appears to have "reverted" to a low SNV frequency is not
sensitive to *k*-mer filtering. Even if no *k*-mer filtering whatsoever were performed for *S. noctiflora*, it
would still exhibit an SNV frequency lower than the filtered values from the other three species
(Table S3). Likewise, even if we did not perform any numt filtering on the *S. latifolia* SNVs (which
would almost certainly lead to a substantial overestimation of true mitochondrial mutations), the SNV
frequency for *S. latifolia* would still not reach the highly elevated levels in *S. conica*. Therefore, the
key distinctions among the three species in mitochondrial SNV frequency appear to hold even
though some caution is warranted in interpreting the specific frequency estimate in *S. latifolia*.
Furthermore, as noted in the Results, the conclusion that plastid SNV frequencies remain low in *S.*
conica is not sensitive to *k*-mer filtering, as removing this filtering step only produces a modest
increase in the estimate for *S. conica* plastid mutations (Table S4).

Heteroplasmic SNV frequencies support a history of mitochondrial mutation rate acceleration
and reversion in *Silene*
The high frequency of mitochondrial SNVs captured by Duplex Sequencing in *S. conica* tissue
(Figure 2) is consistent with previous interpretations that increased mutation rates are driving
accelerated mitochondrial genome evolution in this and other atypical plant species (CHO *et al.* 2004;
PARKINSON *et al.* 2005; MOWER *et al.* 2007; SLOAN *et al.* 2009; SLOAN *et al.* 2012a; SKIPPINGTON *et*

414 *al.* 2015). This view has become the consensus because accelerations are evident at relatively

415 neutral positions like synonymous sites over phylogenetic timescales, but *de novo* mitochondrial

416 mutations have never been directly investigated in these high-rate plant lineages until now. The

417 huge increase in AT→GC transitions that dominated the *S. conica* mutation spectrum (Figure 2)

418 coincides with the most common type of misincorporation observed in steady-state kinetic analysis

419 of the *Arabidopsis* mitochondrial DNA polymerases, which appear to be prone to misincorporating

420 Gs opposite Ts (AYALA-GARCÍA *et al.* 2018). Therefore, it is possible that the increased mitochondrial

421 substitution rate and biased mutation spectrum in *S. conica* reflect a reduced ability to repair

422 mismatches created by polymerase misincorporations. A disproportionate increase in AT→GC

423 transitions was also observed in *Arabidopsis* mutants lacking a functional copy of *MSH1* (WU *et al.*

424 2020a), a gene that may be involved in repair of mismatches via homologous recombination

425 (CHRISTENSEN 2014; WYNN *et al.* 2020). An intact and transcribed copy of *MSH1* is retained in *S.*

426 *conica* (HAVIRD et al. 2017), but its function and expression patterns have not been investigated.

427 Alternatively, GC-biased gene conversion has also been hypothesized as a mechanism to explain

428 skewed substitution patterns in some plant mitochondrial genomes (LIU *et al.* 2020).

429   The extreme bias towards AT→GC transitions in *S. conica* mitochondrial SNVs (Figure 2) is

430 not entirely consistent with longer-term patterns of mitochondrial sequence divergence in this

431 species. The genome-wide GC content in *S. conica* (43.1%) is only slightly higher than in congeners

432 like *S. latifolia* (42.6%), *S. noctiflora* (40.8%), and *S. vulgaris* (41.8%) (SLOAN *et al.* 2012a). An

433 analysis that used plastid DNA insertions in mitochondrial genomes as neutral markers did find that

434 *S. conica* had unusually high transition:transversion ratio compared to other angiosperms (SLOAN

435 AND WU 2014), which is consistent with the Duplex Sequencing results. However, it did not detect

436 the strong GC bias that we observed in the current study.

437   These discrepancies between phylogenetic patterns and our duplex sequencing data raise

438 two obvious possibilities. First, the observed SNVs in this study may not reflect the spectrum of

439 heritable mutations because they are measured from rosette tissue and thus are expected to include

440 some *de novo* mutations that occurred in vegetative tissues and were not inherited from the mother

441 or transmitted to future generations. Our choice to sample whole rosettes (as opposed to more

442 targeted "germline" tissue such as dissected meristems) reflected the practical need to obtain

443 sufficient quantities of mtDNA and cpDNA for Duplex Sequencing library construction. Although it is

444 important to recognize the observed SNVs include some mutations that are not heritable, we do not

445 believe that this is likely to be the primary explanation for inferred differences in mitochondrial

446 mutation spectra. A large proportion of the *S. conica* SNVs were shared across more than one

447 biological replicate, implying that they were inherited from a heteroplasmic mother and thus

448 transmitted across generations. Furthermore, these shared variants were even more skewed

449 towards AT→GC transitions than variants that were only detected in a single replicate (Table S7),

suggesting that heritable mutations do indeed exhibit a very strong GC bias. Relatedly, the fact that

S. conica had such a large number of shared SNVs compared to the other two *Silene* species (see

Results and Table S7) supports the conclusion that the higher overall SNV frequency in *S. conica* is

not solely due to a higher mutation rate in vegetative tissue and indeed reflects an elevated rate of

heritable mutations.

Second, it is possible that the mitochondrial mutation spectrum in *S. conica* is unstable and

has changed over time such that the "snapshot" from Duplex Sequencing of heteroplasmic SNVs

does not match the average spectrum over the past millions of years in this lineage. A recent

analysis of another genus with accelerated mitochondrial sequence divergence (*Ajuga*) found large

increases in GC content (LIU *et al.* 2020), suggesting that bouts of accelerated and GC-biased

evolution can occur in angiosperm mitochondrial genomes.

In contrast to the findings in *S. conica*, we did not observe elevated SNV frequencies in *S.*

noctiflora (Figure 2) despite a comparable history of accelerated sequence evolution (Figure 1). The

low SNV frequencies in *S. noctiflora* suggest that it has reverted to lower mutation rates after a past

episode of acceleration. This type of reversion has been inferred based on phylogenetic evidence in

other accelerated lineages such as *Plantago* and *Pelargonium* (CHO *et al.* 2004; PARKINSON *et al.*

2005). We also have previously speculated that *S. noctiflora* no longer has a high mitochondrial

mutation rate based on its reduced rate of sequence divergence from close relatives *S. turkestanica*

and *S. undulata* and its extremely low amount of intraspecific sequence polymorphism (SLOAN *et al.*

2009; SLOAN *et al.* 2012a; WU *et al.* 2015; WU AND SLOAN 2019). However, if a mutation rate

reversion has occurred in this lineage, it may not have simply reversed the process that led to the

initial rate increase. Notably, *S. noctiflora* had a different mitochondrial mutation spectrum than the

two species that have maintained low mitochondrial substitution rates throughout their histories (*A.*

thaliana and *S. latifolia*). It also retains a mitochondrial genome that is radically altered in size,

structure, and apparent recombinational activity (SLOAN *et al.* 2012a). Therefore, the mechanisms

responsible for mitochondrial DNA replication and maintenance in *S. noctiflora* may still be quite

different than in typical angiosperms despite the apparent reversion to ancestral-like rates in this

species.

The above interpretations are largely based on the premise that the abundance of

heteroplasmic SNVs is correlated with the mutation rate. Although it is probably a reasonable

assumption that these two features are correlated, the amount of heteroplasmic genetic variation

that is maintained will also depend on the (effective) population size of mitochondrial genome

copies. Therefore, we cannot rule out the possibility that some of the observed differences in SNV

frequency among species could be related to variation in traits such as the extent of the mtDNA

transmission bottleneck during reproduction (STEWART AND CHINNERY 2015; ZHANG *et al.* 2018;

JOHNSTON 2019). Likewise, analysis of mitochondrial SNVs in *Arabidopsis* leaf tissue has indicated

that variant frequencies may be affected by features such as plant age and development (WYNN *et*

al. 2020). Therefore, future studies to separate effects of mutation and population size will be useful.

One possibility is that heteroplasmic SNVs identified by Duplex Sequencing could then be tracked

across generations with allele-specific ddPCR. Quantifying variance in levels of inherited

heteroplasmies can serve as an effective means to quantify the effective number of transmitted

genome copies (JOHNSTON 2019). However, this may be more challenging with species such as *S.*

latifolia and *S. noctiflora* where inherited heteroplasmies appear to be rare.

Mitochondrial genome copy number and recombinational repair

One unexpected finding from total-cellular shotgun sequencing was the remarkably low copy number

of the mitochondrial genome is *S. conica* (Figure 3). The ratio of mitochondrial to nuclear genome

copies in the analyzed leaf tissue samples implies that there is only about one to two mitochondrial

genome copies per cell, under the assumption that most cells are diploid. However, this would

depend on the extent of endoreduplication in *S. conica*. Species with smaller nuclear genomes have

been found to undergo a greater amount of endoreduplication on average (BAROW AND MEISTER

2003), so it is possible that the ratio of organellar to nuclear genome copies is skewed downward in

S. conica for this reason. Although there is evidence that plant cells can have far more mitochondria

than mitochondrial genome copies (PREUTEN *et al.* 2010; SHEN *et al.* 2019), it is difficult to imagine

how mitochondrial function could be maintained throughout development with only one or two

mitochondrial genome copies per cell. The sequencing and ddPCR datasets used to generate copy-

number estimates were derived from mature rosette leaf tissue. Therefore, it is possible that this is a

case of mitochondrial genome "abandonment" in tissue that is not destined for further growth or

reproduction (BENDICH 2013; OLDENBURG *et al.* 2013; WYNN *et al.* 2020). Previous studies have

suggested that some plants undergo a major decline in plastid genome copy number in mature

leaves (SHAVER *et al.* 2006; ROWAN *et al.* 2009), although this conclusion has been the subject to

substantial criticism and debate (GOLCZYK *et al.* 2014; GREINER *et al.* 2020). We found that all three

Silene species retained hundreds of plastid genome copies per cell, but the dramatic differences in

mitochondrial genome copy number across species have intriguing implications. An important future

direction will be to characterize variation in *S. conica* mitochondrial genome copy numbers

throughout development, especially in meristematic and reproductive tissues.

Even under the likely scenario that other *S. conica* tissues harbor higher mitochondrial

genome copy numbers than observed in our analysis, it is possible that such values will still be

unusually low compared to most plants and other eukaryotes. *Silene conica* is distinct in having one

of the largest and most fragmented mitochondrial genomes ever identified (SLOAN *et al.* 2012a).

Such genome size and architecture might pose a challenge for mtDNA maintenance in this species.

Notably, mtDNA accounted for a similar proportion of total-cellular DNA in *S. conica* and *S. latifolia*

despite the ~100-fold difference in mitochondrial genome copy number between these samples

because the *S. conica* mitochondrial genome is ~45-fold larger than in *S. latifolia*, and the *S. conica*

nuclear genome is ~3-fold smaller than in *S. latifolia*. Nevertheless, the low mitochondrial genome

copy number in *S. conica* means that any given region of the mitochondrial genome, including key

functional content such as protein-coding sequence, has an unusually low stoichiometry relative to

the nucleus.

We hypothesize that low mitochondrial genome copy number may be a cause of the high

mutation rates in *S. conica*. It is thought that the typically low mutation rates in plant organelle

genomes can be attributed to accurate DNA repair via homologous recombination (KHAKHLOVA AND

BOCK 2006; CHRISTENSEN 2014; AYALA-GARCÍA *et al.* 2018; CHEVIGNY *et al.* 2020; WU *et al.* 2020a).

As such, the ability to maintain low rates would be sensitive to the availability of templates for

recombinational repair and, thus, the number of genome copies. Notably, we did not observe

elevated SNV frequencies in the *S. conica* plastid genome (Figure 2), which appears to retain a

typical copy number, unlike the *S. conica* mitochondrial genome (Figure 3). This hypothesized role of

copy number in recombinational repair is consistent with the observation that nucleotide substitution

rates are lower in large repeats than in single-copy regions within plant organelle genomes (WOLFE

et al. 1987; DAVILA *et al.* 2011; ZHU *et al.* 2016). Therefore, if the population of mitochondrial genome

copies is too sparsely distributed in the cells of *S. conica*, it may be unable to make full use of

recombinational repair and instead rely on less accurate forms of repair or leave some DNA damage

and mismatches entirely unrepaired. In dissertation research conducted with Jeffrey Mower, Wenhu

Guo (2014) arrived at a similar hypothesis after observing an unusually low mitochondrial genome

copy number in *Plantago media*, another angiosperm with extremely elevated rates of mitochondrial

sequence evolution (CHO *et al.* 2004). Therefore, it appears possible that correlated changes in

mitochondrial mutation rate and genome copy number may have occurred many times

independently in plants. Testing this hypothesized relationship between mitochondrial genome copy

number and mutation rate should provide insight into the causes of extreme variation in rates of

mitochondrial sequence evolution observed across angiosperms.

ACKNOWLEDGEMENTS

We thank Justin Havird for helpful discussion and providing *S. conica* full-sib seeds and Jessica

Warren for assistance with DNA extraction and figure preparation. We also thank two anonymous

reviewers for insightful comments that improved the manuscript. This work was supported by a grant

from the NIH (R01 GM118046) and an NSF graduate fellowship (DGE-1450032).

REFERENCES

Ahn, E. H., S. H. Lee, J. Y. Kim, C.-C. Chang and L. A. Loeb, 2016 Decreased mitochondrial mutagenesis during transformation of human breast stem cells into tumorigenic cells. Cancer Research 76: 4569-4578.

Arbeithuber, B., J. Hester, M. A. Cremona, N. Stoler, A. Zaidi *et al.*, 2020 Age-related accumulation of de novo mitochondrial mutations in mammalian oocytes and somatic tissues. PLoS Biology 18: e3000745.

Ayala-García, V. M., N. Baruch-Torres, P. L. García-Medel and L. G. Brieba, 2018 Plant organellar DNA polymerases paralogs exhibit dissimilar nucleotide incorporation fidelity. The FEBS Journal 285: 4005-4018.

Barba-Montoya, J., M. dos Reis, H. Schneider, P. C. Donoghue and Z. Yang, 2018 Constraining uncertainty in the timescale of angiosperm evolution and the veracity of a Cretaceous Terrestrial Revolution. New Phytologist 218: 819-834.

Barow, M., and A. Meister, 2003 Endopolyploidy in seed plants is differently correlated to systematics, organ, life strategy and genome size. Plant, Cell & Environment 26: 571-584.

Bendich, A. J., 2013 DNA abandonment and the mechanisms of uniparental inheritance of mitochondria and chloroplasts. Chromosome Research 21: 287-296.

Bentley, K. E., J. R. Mandel and D. E. McCauley, 2010 Paternal leakage and heteroplasmy of mitochondrial genomes in Silene vulgaris: evidence from experimental crosses. Genetics 185: 961-968.

Brown, W. M., M. George and A. C. Wilson, 1979 Rapid evolution of animal mitochondrial DNA. Proceedings of the National Academy of Sciences 76: 1967-1971.

Chevigny, N., D. Schatz-Daas, F. Lotfi and J. M. Gualberto, 2020 DNA repair and the stability of the plant mitochondrial genome. International Journal of Molecular Sciences 21: 328.

Cho, Y., J. P. Mower, Y. L. Qiu and J. D. Palmer, 2004 Mitochondrial substitution rates are extraordinarily elevated and variable in a genus of flowering plants. Proceedings of the National Academy of Sciences of the United States of America 101: 17741-17746.

Christensen, A. C., 2014 Genes and junk in plant mitochondria-repair mechanisms and selection. Genome Biology and Evolution 6: 1448-1453.

Davila, J. I., M. P. Arrieta-Montiel, Y. Wamboldt, J. Cao, J. Hagmann *et al.*, 2011 Double-strand break repair processes drive evolution of the mitochondrial genome in Arabidopsis. BMC Biology 9: 64.

Drouin, G., H. Daoud and J. Xia, 2008 Relative rates of synonymous substitutions in the mitochondrial, chloroplast and nuclear genomes of seed plants. Molecular Phylogenetics and Evolution 49: 827-831.

Golczyk, H., S. Greiner, G. Wanner, A. Weihe, R. Bock *et al.*, 2014 Chloroplast DNA in mature and senescing leaves: a reappraisal. Plant Cell 26: 847-854.

Greiner, S., H. Golczyk, I. Malinova, T. Pellizzer, R. Bock *et al.*, 2020 Chloroplast nucleoids are highly dynamic in ploidy, number, and structure during angiosperm leaf development. The Plant Journal 102: 730-746.

Guo, W., 2014 Evolution of organellar genome architecture in seed plants: The role of intracellular gene transfer, recombination and mutation, pp. 107-144. University of Nebraska, Lincoln, Nebraska.

Havird, J. C., G. R. Noe, L. Link, A. Torres, D. C. Logan *et al.*, 2019 Do angiosperms with highly divergent mitochondrial genomes have altered mitochondrial function? Mitochondrion 49: 1-11.

Havird, J. C., P. Trapp, C. Miller, I. Bazos and D. B. Sloan, 2017 Causes and consequences of rapidly evolving mtDNA in a plant lineage. Genome Biology and Evolution 9: 323-336.

Havird, J. C., N. S. Whitehill, C. D. Snow and D. B. Sloan, 2015 Conservative and compensatory evolution in oxidative phosphorylation complexes of angiosperms with highly divergent rates of mitochondrial genome evolution. Evolution 69: 3069-3081.

Hazkani-Covo, E., R. M. Zeller and W. Martin, 2010 Molecular poltergeists: mitochondrial DNA copies (numts) in sequenced nuclear genomes. PLoS Genetics 6**:** e1000834.

Hershberg, R., and D. A. Petrov, 2010 Evidence that mutation is universally biased towards AT in bacteria. PLoS Genetics 6**:** e1001115.

Hildebrand, F., A. Meyer and A. Eyre-Walker, 2010 Evidence of selection upon genomic GC-content in bacteria. PLoS Genetics 6**:** e1001107.

Hoekstra, J. G., M. J. Hipp, T. J. Montine and S. R. Kennedy, 2016 Mitochondrial DNA mutations increase in early stage Alzheimer disease and are inconsistent with oxidative damage. Annals of Neurology 80**:** 301-306.

Huang, C. Y., N. Grunheit, N. Ahmadinejad, J. N. Timmis and W. Martin, 2005 Mutational decay and age of chloroplast and mitochondrial genomes transferred recently to angiosperm nuclear chromosomes. Plant Physiology 138**:** 1723-1733.

Jafari, F., S. Zarre, A. Gholipour, F. Eggens, R. K. Rabeler *et al.*, 2020 A new taxonomic backbone for the infrageneric classification of the species-rich genus Silene (Caryophyllaceae). Taxon 69**:** 337-368.

Johnston, I. G., 2019 Varied mechanisms and models for the varying mitochondrial bottleneck. Frontiers in Cell and Developmental Biology 7**:** 294.

Joubes, J., and C. Chevalier, 2000 Endoreduplication in higher plants, pp. 191-201 in *The Plant Cell Cycle*. Springer.

Kennedy, S. R., J. J. Salk, M. W. Schmitt and L. A. Loeb, 2013 Ultra-sensitive sequencing reveals an age-related increase in somatic mitochondrial mutations that are inconsistent with oxidative damage. PLoS Genetics 9**:** e1003794.

Kennedy, S. R., M. W. Schmitt, E. J. Fox, B. F. Kohrn, J. J. Salk *et al.*, 2014 Detecting ultralow-frequency mutations by Duplex Sequencing. Nature Protocols 9**:** 2586-2606.

Khakhlova, O., and R. Bock, 2006 Elimination of deleterious mutations in plastid genomes by gene conversion. Plant Journal 46**:** 85-94.

Kokot, M., M. Długosz and S. Deorowicz, 2017 KMC 3: counting and manipulating k-mer statistics. Bioinformatics 33**:** 2759-2761.

Langmead, B., and S. L. Salzberg, 2012 Fast gapped-read alignment with Bowtie 2. Nature Methods 9**:** 357-359.

Lavrov, D. V., and W. Pett, 2016 Animal mitochondrial DNA as we don't know it: mt-genome organization and evolution in non-bilaterian lineages. Genome Biology and Evolution 8**:** 2896-2913.

Liu, F., W. Fan, J. B. Yang, C. L. Xiang, J. P. Mower *et al.*, 2020 Episodic and guanine–cytosine-biased bursts of intragenomic and interspecific synonymous divergence in Ajugoideae (Lamiaceae) mitogenomes. New Phytologist 228**:** 1107-1114.

Lough, A. N., L. M. Roark, A. Kato, T. S. Ream, J. C. Lamb *et al.*, 2008 Mitochondrial DNA transfer to the nucleus generates extensive insertion site variation in maize. Genetics 178**:** 47-55.

Mandel, J. R., A. J. Ramsey, J. M. Holley, V. A. Scott, D. Mody *et al.*, 2020 Disentangling complex inheritance patterns of plant organellar genomes: an example from carrot. Journal of Heredity In Press.

Martin, M., 2011 Cutadapt removes adapter sequences from high-throughput sequencing reads. EMBnet.journal 17**:** 10-12.

McCauley, D. E., 2013 Paternal leakage, heteroplasmy, and the evolution of plant mitochondrial genomes. New Phytologist 200**:** 966-977.

McCauley, D. E., M. F. Bailey, N. A. Sherman and M. Z. Darnell, 2005 Evidence for paternal transmission and heteroplasmy in the mitochondrial genome of Silene vulgaris, a gynodioecious plant. Heredity 95**:** 50-58.

Michalovová, M., B. Vyskot and E. Kejnovsky, 2013 Analysis of plastid and mitochondrial DNA insertions in the nucleus (NUPTs and NUMTs) of six plant species: size, relative age and chromosomal localization. Heredity 111**:** 314-320.

Mower, J. P., P. Touzet, J. S. Gummow, L. F. Delph and J. D. Palmer, 2007 Extensive variation in synonymous substitution rates in mitochondrial genes of seed plants. BMC Evolutionary Biology 7**:** 135.

Noutsos, C., E. Richly and D. Leister, 2005 Generation and evolutionary fate of insertions of organelle DNA in the nuclear genomes of flowering plants. Genome Research 15**:** 616-628.

Oldenburg, D. J., R. A. Kumar and A. J. Bendich, 2013 The amount and integrity of mtDNA in maize decline with development. Planta 237**:** 603-617.

Palmer, J. D., K. L. Adams, Y. Cho, C. L. Parkinson, Y.-L. Qiu *et al.*, 2000 Dynamic evolution of plant mitochondrial genomes: mobile genes and introns and highly variable mutation rates. Proceedings of the National Academy of Sciences 97**:** 6960-6966.

Palmer, J. D., and L. A. Herbon, 1988 Plant mitochondrial DNA evolves rapidly in structure, but slowly in sequence. Journal of Molecular Evolution 28**:** 87-97.

Parkinson, C. L., J. P. Mower, Y. L. Qiu, A. J. Shirk, K. Song *et al.*, 2005 Multiple major increases and decreases in mitochondrial substitution rates in the plant family Geraniaceae. BMC evolutionary biology 5**:** 73.

Preuten, T., E. Cincu, J. Fuchs, R. Zoschke, K. Liere *et al.*, 2010 Fewer genes than organelles: extremely low and variable gene copy numbers in mitochondria of somatic plant cells. Plant Journal 64**:** 948-959.

Rautenberg, A., D. B. Sloan, V. Aldén and B. Oxelman, 2012 Phylogenetic relationships of Silene multinervia and Silene section Conoimorpha (Caryophyllaceae). Systematic Botany 37**:** 226-237.

Richardson, A. O., D. W. Rice, G. J. Young, A. J. Alverson and J. D. Palmer, 2013 The "fossilized" mitochondrial genome of Liriodendron tulipifera: ancestral gene content and order, ancestral editing sites, and extraordinarily low mutation rate. BMC Biology 11**:** 29.

Rowan, B. A., D. J. Oldenburg and A. J. Bendich, 2009 A multiple-method approach reveals a declining amount of chloroplast DNA during development in Arabidopsis. BMC Plant Biology 9**:** 3.

Salk, J. J., M. W. Schmitt and L. A. Loeb, 2018 Enhancing the accuracy of next-generation sequencing for detecting rare and subclonal mutations. Nature Reviews Genetics 19**:** 269-285.

Schmitt, M. W., S. R. Kennedy, J. J. Salk, E. J. Fox, J. B. Hiatt *et al.*, 2012 Detection of ultra-rare mutations by next-generation sequencing. Proceedings of the National Academy of Sciences of the United States of America 109**:** 14508-14513.

Shaver, J. M., D. J. Oldenburg and A. J. Bendich, 2006 Changes in chloroplast DNA during development in tobacco, Medicago truncatula, pea, and maize. Planta 224**:** 72-82.

Shen, J., Y. Zhang, M. J. Havey and W. Shou, 2019 Copy numbers of mitochondrial genes change during melon leaf development and are lower than the numbers of mitochondria. Horticulture Research 6**:** 95.

Skippington, E., T. J. Barkman, D. W. Rice and J. D. Palmer, 2015 Miniaturized mitogenome of the parasitic plant Viscum scurruloideum is extremely divergent and dynamic and has lost all nad genes. Proceedings of the National Academy of Sciences In Press.

Skippington, E., T. J. Barkman, D. W. Rice and J. D. Palmer, 2017 Comparative mitogenomics indicates respiratory competence in parasitic Viscum despite loss of complex I and extreme sequence divergence, and reveals horizontal gene transfer and remarkable variation in genome size. BMC Plant Biology 17**:** 49.

Sloan, D. B., A. J. Alverson, J. P. Chuckalovcak, M. Wu, D. E. McCauley *et al.*, 2012a Rapid evolution of enormous, multichromosomal genomes in flowering plant mitochondria with exceptionally high mutation rates. PLoS Biology 10**:** e1001241.

Sloan, D. B., A. J. Alverson, M. Wu, J. D. Palmer and D. R. Taylor, 2012b Recent acceleration of plastid sequence and structural evolution coincides with extreme mitochondrial divergence in the angiosperm genus Silene. Genome Biology and Evolution 4**:** 294-306.

Sloan, D. B., A. K. Broz, J. Sharbrough and Z. Wu, 2018 Detecting rare mutations and DNA damage with sequencing-based methods. Trends in Biotechnology 36**:** 729-740.

Sloan, D. B., A. H. MacQueen, A. J. Alverson, J. D. Palmer and D. R. Taylor, 2010 Extensive loss of RNA editing sites in rapidly evolving Silene mitochondrial genomes: Selection vs. retroprocessing as the driving force. Genetics 185**:** 1369-1380.

Sloan, D. B., B. Oxelman, A. Rautenberg and D. R. Taylor, 2009 Phylogenetic analysis of mitochondrial substitution rate variation in the angiosperm tribe *Sileneae* (Caryophyllaceae). BMC Evolutionary Biology 9**:** 260.

Sloan, D. B., and D. R. Taylor, 2010 Testing for selection on synonymous sites in plant mitochondrial DNA: the role of codon bias and RNA editing. Journal of Molecular Evolution 70**:** 479-491.

Sloan, D. B., D. A. Triant, M. Wu and D. R. Taylor, 2014 Cytonuclear interactions and relaxed selection accelerate sequence evolution in organelle ribosomes. Molecular Biology and Evolution 31**:** 673-682.

Sloan, D. B., and Z. Wu, 2014 History of plastid DNA insertions reveals weak deletion and AT mutation biases in angiosperm mitochondrial genomes. Genome Biology and Evolution 6**:** 3210-3221.

Smith, D. R., and P. J. Keeling, 2015 Mitochondrial and plastid genome architecture: Reoccurring themes, but significant differences at the extremes. Proceedings of the National Academy of Sciences of the United States of America 112**:** 10177-10184.

Stewart, J. B., and P. F. Chinnery, 2015 The dynamics of mitochondrial DNA heteroplasmy: implications for human health and disease. Nature Reviews Genetics 16**:** 530-542.

Stupar, R. M., J. W. Lilly, C. D. Town, Z. Cheng, S. Kaul *et al.*, 2001 Complex mtDNA constitutes an approximate 620-kb insertion on Arabidopsis thaliana chromosome 2: implication of potential sequencing errors caused by large-unit repeats. Proceedings of the National Academy of Sciences of the United States of America 98**:** 5099-5103.

Teixeira, S., K. Foerster and G. Bernasconi, 2009 Evidence for inbreeding depression and post-pollination selection against inbreeding in the dioecious plant Silene latifolia. Heredity 102**:** 101-112.

Weaver, R. J., G. Carrion, R. Nix, G. P. Maeda, S. Rabinowitz *et al.*, 2020 High mitochondrial mutation rates in Silene are associated with nuclear-mediated changes in mitochondrial physiology. Biology Letters 16**:** 20200450.

Williams, A. M., M. W. Itgen, A. Lambert, A. K. Broz, R. L. Mueller *et al.*, 2020 Long-read transcriptome and other genomic resources for the angiosperm Silene noctiflora. bioRxiv.

Wolfe, K. H., W. H. Li and P. M. Sharp, 1987 Rates of nucleotide substitution vary greatly among plant mitochondrial, chloroplast, and nuclear DNAs. Proceedings of the National Academy of Sciences 84**:** 9054-9058.

Wu, Z., J. M. Cuthbert, D. R. Taylor and D. B. Sloan, 2015 The massive mitochondrial genome of the angiosperm Silene noctiflora is evolving by gain or loss of entire chromosomes. Proceedings of the National Academy of Sciences 112**:** 10185-10191.

Wu, Z., and D. B. Sloan, 2019 Recombination and intraspecific polymorphism for the presence and absence of entire chromosomes in mitochondrial genomes. Heredity 122**:** 647-659.

Wu, Z., G. Waneka, A. K. Broz, C. R. King and D. B. Sloan, 2020a MSH1 is required for maintenance of the low mutation rates in plant mitochondrial and plastid genomes. Proceedings of the National Academy of Sciences In Press.

Wu, Z., G. Waneka and D. B. Sloan, 2020b The tempo and mode of angiosperm mitochondrial genome divergence inferred from intraspecific variation in Arabidopsis thaliana. G3: Genes, Genomes, Genetics 10**:** 1077-1086.

Wynn, E., E. Purfeerst and A. Christensen, 2020 Mitochondrial DNA repair in an *Arabidopsis thaliana* uracil N-glycosylase mutant. Plants 9**:** 261.

Wynn, E. L., and A. C. Christensen, 2015 Are synonymous substitutions in flowering plant mitochondria neutral? Journal of Molecular Evolution 81**:** 131-135.

Yang, Z., 2007 PAML 4: Phylogenetic Analysis by Maximum Likelihood. Molecular Biology and Evolution 24**:** 1586-1591.

Zervas, A., G. Petersen and O. Seberg, 2019 Mitochondrial genome evolution in parasitic plants. BMC Evolutionary Biology 19**:** 87.

Zhang, H., S. P. Burr and P. F. Chinnery, 2018 The mitochondrial DNA genetic bottleneck: inheritance and beyond. Essays in Biochemistry 62**:** 225-234.

Zhu, A., W. Guo, S. Gupta, W. Fan and J. P. Mower, 2016 Evolutionary dynamics of the plastid inverted repeat: the effects of expansion, contraction, and loss on substitution rates. New Phytologist 209**:** 1747-1756.

Figure 1. Mitochondrial substitution rate variation among *Silene* species. Branch lengths are scaled based on synonymous substitutions per site (d_S) in a concatenation of three mitochondrial protein-coding genes (*atp1*, *cox3*, and *nad9*) used in previous analyses (SLOAN *et al.* 2009; RAUTENBERG *et al.* 2012). Branch lengths were estimated with PAML v4.9j (YANG 2007), using a constrained topology. Note that the two accelerated clades (section Conoimorpha [represented by *S. conica*] and section Elisanthe [represented by *S. noctiflora*]) were set as sister groups for this analysis, but there is only weak and inconsistent evidence for that relationship (RAUTENBERG *et al.* 2012; HAVIRD *et al.* 2017; JAFARI *et al.* 2020). Images are shown for each of the three focal species in this study (*S. latifolia*, *S. conica*, and *S. noctiflora*).

Figure 2. Variation in mitochondrial and plastid SNV frequencies and spectra across *Arabidopsis* and *Silene* species. SNV frequencies are calculated as the total number of observed single-nucleotide mismatches in duplex consensus sequence datasets dividided by the total base-pairs of mitochondrial or plastid genome coverage in those datasets (i.e., GC coverage, AT coverage, or total coverage depending on the SNV type). Three biological replicates (circles) are shown for each species, with the mean of those replicates indicated with a horizontal line. The *Arabidopsis* data points are taken from Wu et al. (2020). The same data are plotted on a log scale in Figure S2.

Figure 3. Variation in mitochondrial genome copy number among *Silene* species. (A) Average mitochondrial, plastid, and nuclear genome copy number were estimated from total-cellular Illuimina shotgun sequencing of leaf tissue from each species. Boxplots show median and interquartile ranges for the ratio of organelle genome copy number to nuclear genome copy number based on scanning the organelle genomes in 2-kb windows. Green and gold boxplots correspond to plastid:nuclear and mitochondrial:nuclear ratios, respectively. (B) ddPCR analysis of mitochondrial and plastid genome copies per nuclear genome copy in *S. conica*. Points are shown for two mitochondrial markers (*matR* and *nad9*) and two plastid markers (*petA* and *rpoB*). Estimates for copy number ratios were generated by dividing each mitochondrial or plastid value by the average copy number of two nuclear markers for the corresponding sample (Table S1). The triangles indicate ddPCR estimates for the sample taken from the same DNA extraction used in the original sequencing analysis in part (A). The circles represent the three new samples collected for this ddPCR analysis.

Figure S1. Summary of family sizes for single-stranded consensus sequences used for generation of double-stranded consensus sequences for each Duplex Sequencing library. The analysis pipeline required a minimum family size of 3 reads for each of the two complementary single-stranded families. The mitochondrial DNA library for the second biological replicate from *S. conica* was not sequenced because of an apparent library loading error.

Figure S2. Variation in mitochondrial and plastid SNV frequencies and spectra across *Arabidopsis* and *Silene* species. The same data shown in Figure 2 are plotted on a log scale here.

Figure S3. Summary of mitochondrial genome coverage in the *S. latifolia* UK2600 total-cellular sequencing library. Coverage is estimated based on reads mapping with a maximum of one mismatch and no indels in 2-kb windows.

Figure S4. Summary of mitochondrial genome coverage in the *S. noctiflora* OSR total-cellular sequencing library. Coverage is estimated based on reads mapping with a maximum of one mismatch and no indels in 2-kb windows. Each panel represents a different chromosome within the multichromosomal genome.

Figure S5a. Summary of coverage for mitochondrial chromosomes 1 through 64 in the *S. conica* ABR total-cellular sequencing library. Coverage is estimated based on reads mapping with a maximum of one mismatch and no indels in 2-kb windows. Note that four of the data points on chromosome 20 and two of the data points on chromosome 32 exceeded a coverage of 250× and are not shown to improve readability of the plots.

Figure S5b. Summary of coverage for mitochondrial chromosomes 65 through 128 in the *S. conica* ABR total-cellular sequencing library. Coverage is estimated based on reads mapping with a maximum of one mismatch and no indels in 2-kb windows. Note that four of the data points on chromosome 121 exceeded a coverage of 250× and are not shown to improve readability of the plots.

Table S1. Primers used in the *S. conica* ddPCR analysis. The two nuclear markers are referred to by the identifiers for the homologs in *Arabidopsis thaliana*.

Gene	Genome	Forward Primer	Reverse Primer
nad9	mitochondrial	CAAGAAGTGGGTCAAAAGAATG	CGAGAATTAAAACGAGTACTCAG
matR	mitochondrial	GATCAGAATGGTACCCGAATC	CCTTGAAGATTGCTAGGAGTTG
rpoB	plastid	CGGGAGATGCAAGAACAAAC	CTGATCTTCCTCCCCAATCTG
petA	plastid	CTATCCCATTTTTGCCCAGC	CCCCTTTCTTGCCGTTAGC
AT1G06040	nuclear	CCTTTATCTTCTGTGTGGAGG	GGCTGTGCACTTTGGTTGG
AT1G07630	nuclear	CAGAAAAGTGGAATGATGCTC	CAATGAGATGAAGTGCTCTACC

Table S2. Summary of library sequencing and mapping. Reported coverage values and mapping percentages are for duplex consensus sequences.

Library	Library Type	SRA Accession	Raw Read Pairs	Mito Coverage (bp)	Plastid Coverage (bp)	Mito Mapping (%)	Plastid Mapping (%)	Unmapped (%)
Silene conica chloroplast 1	Chloroplast duplex	SAMN17011097	1.15E+08	9.25E+07	2.03E+08	29.49%	64.88%	5.63%
Silene conica chloroplast 2	Chloroplast duplex	SAMN17011098	1.15E+08	1.33E+08	2.72E+08	31.48%	64.13%	4.39%
Silene conica chloroplast 3	Chloroplast duplex	SAMN17011099	1.42E+08	1.37E+08	2.70E+08	31.72%	62.55%	5.73%
Silene conica mito 1	Mitochondrial duplex	SAMN17011100	1.01E+08	2.18E+08	1.77E+07	80.87%	6.58%	12.55%
Silene conica mito 3	Mitochondrial duplex	SAMN17011101	1.14E+08	2.16E+08	2.01E+07	77.21%	7.16%	15.63%
Silene latifolia chloroplast 1	Chloroplast duplex	SAMN17011085	1.33E+08	2.89E+07	2.83E+08	7.39%	72.35%	20.26%
Silene latifolia chloroplast 2	Chloroplast duplex	SAMN17011086	1.28E+08	3.54E+07	2.41E+08	8.82%	60.10%	31.08%
Silene latifolia chloroplast 3	Chloroplast duplex	SAMN17011087	1.01E+08	3.68E+07	2.73E+08	9.13%	67.73%	23.14%
Silene latifolia mito 1	Mitochondrial duplex	SAMN17011088	1.13E+08	1.16E+08	2.67E+07	38.26%	8.83%	52.91%
Silene latifolia mito 2	Mitochondrial duplex	SAMN17011089	9.79E+07	4.93E+07	1.63E+07	24.87%	8.21%	66.92%
Silene latifolia mito 3	Mitochondrial duplex	SAMN17011090	8.83E+07	6.56E+07	2.23E+07	27.94%	9.48%	62.58%
Silene noctiflora chloroplast 1	Chloroplast duplex	SAMN17011091	1.37E+08	1.06E+08	2.55E+08	27.63%	66.35%	6.02%
Silene noctiflora chloroplast 2	Chloroplast duplex	SAMN17011092	1.25E+08	9.39E+07	2.20E+08	28.18%	66.12%	5.70%
Silene noctiflora chloroplast 3	Chloroplast duplex	SAMN17011093	1.37E+08	1.25E+08	3.19E+08	25.69%	65.36%	8.95%
Silene noctiflora mito 1	Mitochondrial duplex	SAMN17011094	1.32E+08	2.76E+08	1.56E+07	83.03%	4.68%	12.29%
Silene noctiflora mito 2	Mitochondrial duplex	SAMN17011095	1.06E+08	1.88E+08	1.22E+07	81.54%	5.28%	13.18%
Silene noctiflora mito 3	Mitochondrial duplex	SAMN17011096	1.03E+08	1.93E+08	2.39E+07	71.53%	8.84%	19.63%
Silene conica ABR	Total cellular shotgun	SAMN17011105	4.98E+08	3.53E+08	1.68E+09	0.49%	2.31%	97.20%
Silene noctiflora OSR	Total cellular shotgun	SAMN17011104	5.76E+08	3.45E+09	2.10E+09	2.51%	1.53%	95.96%
Silene latifolia UK2600	Total cellular shotgun	SAMN17011102	2.81E+08	5.95E+08	5.51E+09	0.50%	4.64%	94.86%
Silene latifolia Kew 32982	Total cellular shotgun	SAMN17011103	3.04E+08					

Table S3. *Silene* mitochondrial SNV count and frequency data with and without *k*-mer filtering

Species	Rep	k-mer Filt	SNV Counts							SNV Frequencies						
			A>C	A>G	A>T	C>A	C>G	C>T	Total	A>C	A>G	A>T	C>A	C>G	C>T	Total
Silene conica	1	10	53	356	26	13	6	18	472	3.0E-07	2.0E-06	1.5E-07	9.7E-08	4.5E-08	1.3E-07	1.5E-06
Silene conica	2	10	27	176	15	4	9	6	237	3.6E-07	2.3E-06	2.0E-07	6.9E-08	1.6E-07	1.0E-07	1.8E-06
Silene conica	3	10	67	462	17	13	6	13	578	3.3E-07	2.3E-06	8.5E-08	8.5E-08	3.9E-08	8.5E-08	1.6E-06
Silene conica	1	None	150	483	58	49	12	217	969	8.5E-07	2.7E-06	3.3E-07	3.6E-07	8.9E-08	1.6E-06	3.1E-06
Silene conica	2	None	64	238	24	20	12	98	456	8.5E-07	3.2E-06	3.2E-07	3.5E-07	2.1E-07	1.7E-06	3.4E-06
Silene conica	3	None	158	610	50	59	17	231	1125	7.9E-07	3.0E-06	2.5E-07	3.9E-07	1.1E-07	1.5E-06	3.2E-06
Silene latifolia	1	10	1	1	0	4	3	18	27	5.7E-09	5.7E-09	0.0E+00	3.0E-08	2.2E-08	1.3E-07	8.7E-08
Silene latifolia	2	10	1	0	1	2	2	5	11	1.3E-08	0.0E+00	1.3E-08	3.5E-08	3.5E-08	8.6E-08	8.2E-08
Silene latifolia	3	10	1	2	6	1	0	10	20	5.0E-09	1.0E-08	3.0E-08	6.5E-09	0.0E+00	6.5E-08	5.7E-08
Silene latifolia	1	None	9	8	6	17	5	42	87	1.1E-07	9.7E-08	7.3E-08	2.7E-07	8.1E-08	6.8E-07	6.0E-07
Silene latifolia	2	None	5	8	14	17	6	58	108	1.0E-07	1.7E-07	2.9E-07	4.7E-07	1.7E-07	1.6E-06	1.3E-06
Silene latifolia	3	None	9	16	18	19	10	52	124	1.5E-07	2.7E-07	3.1E-07	4.3E-07	2.3E-07	1.2E-06	1.2E-06
Silene noctiflora	1	10	3	5	1	0	4	3	16	1.3E-08	2.2E-08	4.4E-09	0.0E+00	2.6E-08	1.9E-08	4.2E-08
Silene noctiflora	2	10	1	8	0	0	7	5	21	6.0E-09	4.8E-08	0.0E+00	0.0E+00	6.1E-08	4.3E-08	7.4E-08
Silene noctiflora	3	10	2	7	0	2	4	6	21	1.1E-08	3.7E-08	0.0E+00	1.5E-08	3.1E-08	4.6E-08	6.6E-08
Silene noctiflora	1	None	3	8	3	4	6	12	36	1.3E-08	3.5E-08	1.3E-08	2.6E-08	3.8E-08	7.7E-08	9.4E-08
Silene noctiflora	2	None	2	11	1	2	13	9	38	1.2E-08	6.6E-08	6.0E-09	1.7E-08	1.1E-07	7.8E-08	1.3E-07
Silene noctiflora	3	None	5	13	2	11	9	11	51	2.7E-08	6.9E-08	1.1E-08	8.4E-08	6.9E-08	8.4E-08	1.6E-07

Table S4. *Silene* plastid SNV count and frequency data with and without *k*-mer filtering

Species	Rep	k-mer Filt	SNV Counts							SNV Frequencies						
			A>C	A>G	A>T	C>A	C>G	C>T	Total	A>C	A>G	A>T	C>A	C>G	C>T	Total
Silene conica	1	10	0	2	1	3	4	8	18	0.0E+00	1.6E-08	7.9E-09	3.9E-08	5.2E-08	1.0E-07	8.8E-08
Silene conica	2	10	0	0	0	4	7	10	21	0.0E+00	0.0E+00	0.0E+00	3.9E-08	6.8E-08	9.7E-08	7.7E-08
Silene conica	3	10	0	1	0	0	2	19	22	0.0E+00	6.0E-09	0.0E+00	0.0E+00	2.0E-08	1.9E-07	8.2E-08
Silene conica	1	None	1	4	1	3	4	12	25	7.9E-09	3.2E-08	7.9E-09	3.9E-08	5.2E-08	1.6E-07	1.2E-07
Silene conica	2	None	0	0	0	6	8	20	34	0.0E+00	0.0E+00	0.0E+00	5.8E-08	7.7E-08	1.9E-07	1.3E-07
Silene conica	3	None	0	1	1	2	4	26	34	0.0E+00	6.0E-09	6.0E-09	2.0E-08	3.9E-08	2.5E-07	1.3E-07
Silene latifolia	1	10	4	2	0	1	2	10	19	2.3E-08	1.1E-08	0.0E+00	9.4E-09	1.9E-08	9.4E-08	6.7E-08
Silene latifolia	2	10	4	0	3	0	7	13	27	2.7E-08	0.0E+00	2.0E-08	0.0E+00	7.7E-08	1.4E-07	1.1E-07
Silene latifolia	3	10	4	2	0	0	7	19	32	2.4E-08	1.2E-08	0.0E+00	0.0E+00	6.7E-08	1.8E-07	1.2E-07
Silene latifolia	1	None	8	6	4	2	3	31	54	4.5E-08	3.4E-08	2.3E-08	1.9E-08	2.8E-08	2.9E-07	1.9E-07
Silene latifolia	2	None	10	4	11	10	10	38	83	6.7E-08	2.7E-08	7.3E-08	1.1E-07	1.1E-07	4.2E-07	3.4E-07
Silene latifolia	3	None	7	12	6	6	9	30	70	4.1E-08	7.1E-08	3.5E-08	5.8E-08	8.7E-08	2.9E-07	2.6E-07
Silene noctiflora	1	10	3	0	1	0	5	6	15	1.9E-08	0.0E+00	6.3E-09	0.0E+00	5.2E-08	6.2E-08	5.9E-08
Silene noctiflora	2	10	5	0	0	2	14	11	32	3.6E-08	0.0E+00	0.0E+00	2.4E-08	1.7E-07	1.3E-07	1.5E-07
Silene noctiflora	3	10	11	0	1	0	14	8	34	5.6E-08	0.0E+00	5.1E-09	0.0E+00	1.2E-07	6.6E-08	1.1E-07
Silene noctiflora	1	None	5	1	1	1	5	9	22	3.1E-08	6.3E-09	6.3E-09	1.0E-08	5.2E-08	9.3E-08	8.6E-08
Silene noctiflora	2	None	6	1	0	2	14	19	42	4.4E-08	7.3E-09	0.0E+00	2.4E-08	1.7E-07	2.3E-07	1.9E-07
Silene noctiflora	3	None	12	0	2	2	14	17	47	6.1E-08	0.0E+00	1.0E-08	1.7E-08	1.2E-07	1.4E-07	1.5E-07

Table S5. Genome copy number estimates

Species	Nuclear (=Unmapped) Coverage (bp)	Mito Coverage (bp)	Plastid Coverage (bp)	Nuclear Genome Size (bp)	Mito Genome Size (bp)	Plastid Genome Size (bp)	Nuclear Cov.	Mito Cov.	Plastid Cov.	Mito per Nuclear	Plastid per Nuclear
Silene latifolia	7.58E+10	3.53E+08	1.68E+09	2.67E+09	2.53E+05	1.52E+05	29.22	1394.70	11047.66	47.72	378.02
Silene noctiflora	1.41E+11	3.45E+09	2.10E+09	2.78E+09	6.73E+06	1.52E+05	52.70	512.25	13838.01	9.72	262.59
Silene conica	1.20E+11	5.95E+08	5.51E+09	9.30E+08	1.13E+07	1.47E+05	136.49	52.55	37486.69	0.38	274.64

Table S6. ddPCR droplet counts and copy number calculations

Sample	Genome	Marker	Positive droplets	Negative droplets	Copies per reaction	Dilution Factor	Copies per nuclear copy
New1	Mito	matR	1244	6955	3967	1	0.98
New2	Mito	matR	2413	10431	4900	1	1.09
New3	Mito	matR	1987	11250	3820	1	0.91
Original	Mito	matR	2356	13893	3680	1	0.42
New1	Mito	nad9	1842	10249	3880	1	0.96
New2	Mito	nad9	1202	4781	5280	1	1.18
New3	Mito	nad9	990	5106	4160	1	1.00
Original	Mito	nad9	2268	13951	3540	1	0.41
New1	Plastid	petA	6502	7514	14660	200	727.54
New2	Plastid	petA	6692	6199	17220	200	767.04
New3	Plastid	petA	6392	8264	13480	200	644.98
Original	Plastid	petA	5601	1651	34820	200	804.16
New1	Plastid	rpoB	5868	6456	15220	200	755.33
New2	Plastid	rpoB	6803	6015	17800	200	792.87
New3	Plastid	rpoB	6409	8078	13740	200	657.42
Original	Plastid	rpoB	10798	4448	28980	200	669.28
New1	Nuclear	AT1G06040	1524	7912	4140	1	
New2	Nuclear	AT1G06040	1751	8889	4240	1	
New3	Nuclear	AT1G06040	2070	10630	4180	1	
Original	Nuclear	AT1G06040	3982	10165	7780	1	
New1	Nuclear	AT1G07630	1756	9820	3920	1	
New2	Nuclear	AT1G07630	2242	10039	4740	1	
New3	Nuclear	AT1G07630	2118	10890	4180	1	
Original	Nuclear	AT1G07630	5315	10629	9540	1	

Table S7. Detection of mitochondrial SNVs in multiple biological replicates

Species	SNV Detection	A>C	A>G	A>T	C>A	C>G	C>T	Total
Silene conica	Shared among replicates	17	100	5	3	1	2	128
Silene conica	Unique to one replicate	32	157	28	19	19	21	276
Silene latifolia	Shared among replicates**	0	0	0	0	0	0	0
Silene latifolia	Unique to one replicate	3	3	7	7	5	33	58
Silene noctiflora	Shared among replicates	0	0	0	0	0	0	0
Silene noctiflora	Unique to one replicate	6	20	1	2	15	14	58

**The use of parentals for *k*-mer filtering may have biased against detection of shared SNVs in *S. latifolia* (see main text).