Directed differentiation of Dental Pulp Pluripotent-like Stem Cells into Hepatocyte-like Cells

Carlos Gil-Recio[1], Sheyla Montori[1], Cámara Vallejo M[2], Saddam Al Demour[3], Eduard Ferrés-Padró[4], Miguel Barajas[5], Carles Martin[2], Ashraf Al Madhoun[6*] and Maher Atari[1,7*]

[1]Regenerative Medicine Research Institute, UIC Barcelona, Barcelona, Spain
[2]Oral and Maxillofacial Surgery Department, Hospital Clinico de Barcelona, Barcelona, Spain
[3]Department of Special Surgery/Division of Urology, The University of Jordan, School of Medicine, Amman, Jordan.
[4]Oral and Maxillofacial Surgery Department, Fundació Hospital de Nens de Barcelona, Barcelona, Spain
[5]Biochemistry and Molecular Biology Department, Universidad Pública de Navarra, Pamplona, Spain
[6]Functional Genomic Unit, Dasman Diabetes Institute, Kuwait
[7]Surgery and Oral Implantology Department, College of Dentistry Universitat Internacional de Catalunya, Spain

* Correspondence: matari@biointelligentsl.com and ashraf.madhoun@dasmaninstitute.org

Running Title: Hepatocyte generation from DPPSCs

Carlos Gil-Recio, carlosg@uic.es
Sheyla Montori, smontori@uic.es
Saddam H. Al Demour, saldemour@ju.edu.jo
Mera A. Ababneh, mababneh@just.edu.jo
Eduard Ferrés Padró, eduard.fp@institutferresamat.com
Carles Martin, martipages.c@gmail.com
Elvira Ferrés-Amat, elvira.fa@institutferresamat.com
Miguel Barajas, miguel.barajas@unavarra.es
Ashraf Madhoun, ashraf.madhoun@dasmaninstitute.org
Maher Atari, matari@biointelligentsl.com

Abstract: Liver diseases are a major cause of morbidity and mortality. Dental pulp pluripotent- like stem cells (DPPSC) are a considerable promise in tissue engineering and regenerative medicine as a source of tissue-specific cells, therefore this study aims to demonstrate their ability to generate functional hepatocyte-like cells *in vitro*. Cells were differentiated on a collagen scaffolds in serum free media supplemented with growth factors and cytokines to recapitulate liver development. At day 5, the differentiated DPPSC cells expressed the endodermal markers FOXA1 and FOXA2. Then, the cells were derived into the hepatic lineage generating hepatocyte-like cells. In addition to the associated morphological changes, the cells expressed the hepatic genes HNF6, AFP. The terminally differentiated hepatocyte-like cells expressed the liver functional proteins albumin and CYP3A4. In this study, we report an efficient serum-free protocol to differentiate DPPSCs into functional hepatocyte-like cells. Our approach promotes the use of DPPSCs as a new source of adult stem cells for prospective use in liver regenerative medicine.

Keywords: hepatocyte differentiation; adult stem cells; dental pulp pluripotent stem cells; DPPSCs

1. Introduction

The liver is the largest internal organ providing essential metabolic, exocrine, and endocrine metabolites. Hepatocytes and cholangiocytes, the bile duct epithelial cells, are parenchymal cells which comprise approximately 70% of the adult liver mass. Both cell types are derived from the embryonic definitive endoderm [1]. The liver is the main body homeostasis regulator; therefore, liver diseases cause high morbidity and mortality rates. The incidence of end-stage liver disease (ESLD) is increasing worldwide [2] due to alcohol induced liver disease, nonalcoholic fatty liver disease, or hepatitis. Orthotopic liver transplantation (OLT) is the current optimal treatment with more than 70% of the overall 5-year survival rate for certain ESLD cases [3]. OLT obstacles include donor availability, surgical risk, high costs, and the requirement for life-long immunosuppressors [4; 5; 6]. Alternatively, hepatocyte transplantation (HT) is a promising procedure, less invasive, can be performed repeatedly, used mainly to replace an inborn deficient metabolic function [7] [8]. Yet, HT hurtles include donor availability, low cell engraftment, and longevity [9]. Therefore, the scientific community has put an effort to improve cell-based hepatocyte generation.

Dental pulp stem cells (DPSCs) are a multipotent heterogeneous cell population embedded within the pulp cavity of impacted third molars. DPSCs were initially isolated and characterized by Gronthos *et. al.* [10]. Subsequently, several investigators reported DPSCs' isolation, characterization, differentiation, and banking [11; 12]. In comparison with embryonic and other adult stem cells, DPSCs are obtained from disposable dental pulp after occlusion management, which is considered as a medical waste, noninvasive isolation procedure, not associated with ethical constraints, and no risk of teratoma formation. Furthermore, DPSCs do not compromise their stemness, viability, proliferation, or differentiating capabilities after cryopreservation [13; 14]. Therefore, DPSCs have the potential to be a promising personalized pa-

tient-specific stem cell source for regenerative therapy. Nevertheless, a major concern is that DPSCs comprise progenitor cell populations that are marked by diverse characteristics, such as clonal heterogeneity, multilineage differentiation, self-renewal capacity and phenotypic complexity.

In the context of cell-based hepatogenesis, DPSCs-mediated hepatocyte-like generations has advanced rapidly because of the previously accumulated differentiation protocols applied for human embryonic stem cells (hESCs) and bone marrow mesenchymal cells (BM-MSCs) as prospective sources for regenerative hepatocytes [15; 16; 17; 18; 19; 20]. Implementing defined, serum-free and stepwise differentiation protocols that mimic hepatocytes' developmental stages during embryogenesis are sufficient for inducing hepatogenesis. Using this approach, Ishkitiev *et. al.* were the pioneers to demonstrate the hepatogenic differentiation potential of DPSCs. Initially, Ishkitiev *et. al.* developed two staged conditioned media containing a low percentage of fetal bovine serum (FBS) but enriched with essential hepatogenic inducers [21]. Later, they used serum-free conditioned media to generate hepatocyte-like cells from CD117$^+$ DPSCs subpopulation [22]. The latter hepatogenesis protocol utilized three developmental stages, i.e., cell specification, differentiation, and maturation, to generate cells with morphological, phenotypical, and functional characteristics like hepatocytes. This approach was further improved and implemented in later studies [11; 23]. Nevertheless, the Heterogeneity nature of the DPSCs influences their differentiation capacity and yield, therefore, we focused our efforts to identify and characterize the differentiation potential of a defined subpopulation of DPSCs with pluripotent-like phenotype [24; 25; 26; 27; 28].

Dental pulp pluripotent stem cells (DPPSCs) are a unique sub-population of DPSCs [25], expressing the pluripotency markers Oct4A, NANOG, and SOX2 [24; 29; 30]. DPPSCs exhibits a normal human karyotype with no aneuploidy, polyploidy, or any chromosomal abnormality

during the metaphase stage even after more than 65 passages [24; 31]. Unlike other adult stem cells, DPPSCs form teratoma-like structures when injected subcutaneously in immunodeficient mice, and generate embryonic bodies like in *in vitro* cultures, which are exclusive phenotypes of hESCs and induced pluripotent stem cells (iPSCs) iPSCs [24; 32; 33; 34]. Furthermore, DPPSCs can be differentiated into cells from each of the three germ-embryonic layers, including osteocytes and bone- [25; 26], endothelial- and neural- like cells [25; 26], smooth and skeletal muscles [27].

In this study, we evaluated the adult DPPSCs- mediated hepatogenic capacity using a novel serum free- directed- step wide differentiation protocol and collagen scaffolds to induce 3-dimentaial (3D) culture conditions. DPPSCs were first differentiated into definitive endoderm (DE), which were then directed to the hepatogenic lineage. We observed that treatment of DPPSCs, seeded on scaffolds with Activin A (Act A), Wnt Family Member 3A (Wnt3A), and knockout serum replacement (KOSR) conditioned media, enhances their commitment capacity toward DE- lineage. Subsequently, treatment with fibroblast growth factor 4 (FGF4) and hepatocyte growth factor (HGF) induced gut tube formation, early hepatogenic markers, and finally, functional hepatocyte-like cells were generated in conditional differentiation medium supplemented with HGF, FGF4, dexamethasone (Dex) and oncostatin M (OSM). Preliminary results were previously published in a preprint [35].

2. Materials and methods

2.1. Ethics Approval and Statement

All research in the present study was conducted in accordance with the code of ethics of the World Medical Association for experiments involving humans (Helsinki Declaration of 1975) and the guidelines on human stem cell research issued by the Committee of Bioethics at the Universitat International de Catalunya, Spain.

2.2. DPPSCs Culture and Maintenance

DPPSCs clones were previously isolated and characterized by our group as previously de-

scribed [24; 31]. In Brief, immediately after extraction, the third molars were vigorously

washed with 70% ethanol and sterile distilled water. The molar pulp tissues were extracted,

and cells were isolated by digesting the pulp tissue with collagenase type I (3 mg/ml, Sig-

ma-Aldrich, Germany) for 60 minutes at 37°C. Then, cells were separated mechanically with

an insulin syringe and centrifuged for 10 minutes at 1800 rpm. Primary cell lines were estab-

lished, the medium was changed every 4 days, and cell density was maintained at a low den-

sity of 80-100 cells/cm^2. DPPSCs were cultured in precoated flasks with 100 ng/ml fibronectin

(Life Technologies, Waltham, MA, USA) in a medium consisting of 60% DMEM-low glucose

(Life Technologies) and 40% MCDB-201 (Sigma-Aldrich); supplemented with 1X SITE Liquid

Media Supplement (Sigma-Aldrich); 1X linoleic acid-bovine serum albumin (LA-BSA, Sig-

ma-Aldrich); 10^{-4} M L-ascorbic acid 2-phosphate (Sigma-Aldrich); 1X Penicillin-Streptomycin

(Life Technologies); 2% fetal bovine serum (FBS, Sigma-Aldrich); 10 ng/ml hPDGF-BB (Abcam,

Waltham, MA, USA); 10 ng/ml EGF (Sigma-Aldrich); 1000 U/ml LIF (Millipore, USA); chemically

defined Lipid Concentrate (Life Technologies); 0.8 mg/ml BSA (Sigma-Aldrich) and 55 µM

β-mercaptoethanol (Sigma-Aldrich).

2.3. Definitive Endoderm Induction using Growth Factors

For the DE- induction, $5x10^4$ cells/cm^2 were seeded on 6-well plates (Cell Coat, UK) and

cultured in RPMI medium (Mediatech, Los Altos, CA, USA) containing GlutaMAX (Life Tech-

nologies), penicillin/streptomycin, 0.5% defined fetal bovine serum (FBS, HyClone, Cytiva Eu-

rope GmbH, Spain) and supplemented with each one of the following combinations of growth

factors: (i) 100 ng/ml Act A (R&D Systems, UK), (ii) 100 ng/ml Act A + 50 ng/ml Wnt3A (R&D

Systems), (iii) 100 ng/ml Act A + 50 ng/ml bone morphogenetic protein-4 (BMP4, R&D Sys-

tems), (iv) 100 ng/ml Act A + 50 ng/ml Wnt3A + 50 ng/ml BMP4, (v) 100 ng/ml Act A + 10 ng/ml FGF4 (R&D Systems) or (vi) 100 ng/ml Act A + 10 ng/ml basic-FGF (bFGF, Life Technologies). Three days post-induction, the medium was changed to the same RPMI-based medium and supplements except for the replacement of FBS with 2% KOSR (Life Technologies) and the incubation was maintained for another 2 days.

2.4. Definitive Endoderm Induction Using Different Biomaterials

Application of biomaterials to cell culture has been shown to enhance cellular programming and differentiation. In this study, we used collagen I or fibronectin, the most abundant components of the extracellular matrix that provide a structural network for the cells microenvironment and were reported to enhance DE- differentiation *in vitro* [36; 37]. Using the sample induction media described above, cells were seeded on plates coated with collagen type I, 100 ng/ml fibronectin, or were co-cultured with HepG2 cells (20,000 cell/cm^2), in addition to the controls. 2.5. Hepatogenic differentiation

After setting the optimal conditions for DE-induction; we aimed to generate stepwise functional hepatocyte-like cell. The gut tube and hepatocyte specifications were initiated using RPMI medium supplemented with 2% KOSR, 10 ng/ml FGF4, and 10 ng/ml HGF (R&D Systems). Three days later, RPMI was replaced with the enriched minimum essential medium (MEM, Sigma-Aldrich) supplemented with 1% BSA, 10 ng/ml FGF4, and 10 ng/ml HGF. Three days later, hepatocyte maturation was induced using complete hepatocyte culture medium (HCM, Single-quote, Lonza, USA) with all supplements according to the manufacturer recommendations; containing 10 ng/ml FGF4, 10 ng/ml HGF, 10 ng/ml OSM and 10^{-7} M Dex (Sigma-Aldrich). The hepatocyte maturation was proceeded for nine days as previously reported by Ishkitiev *et. al.* [21; 22]. The differentiation protocol was performed for a total of 22 days, and the media were refreshed every 2-3 days.

2.6. Immunofluorescence microscope imaging

Immunofluorescence was performed as described previously [38; 39]. Briefly, cells were cultured overnight on glass coverslips coated with collagen I, fixed with 4% paraformaldehyde for 15 min, and permeabilized with 10% Triton X-100 for 30 min. Then, cells were blocked with 5% BSA in PBS for 30 min and incubated for one hour with the primary antibodies: alpha-1 antitrypsin (AAT; Abcam) or albumin (Alb; Abcam). Followed by one-hour incubation with prospective secondary antibodies (Abcam). Between each step, cells were washed with 1% BSA in PBS. Fluorescent images were captured using a fluorescence microscopy Olympus AX70 (Olympus Optical, Tokyo, Japan) as previously described [40; 41].

2.7. Immunocytochemistry

Cells were fixed with Thin Prep-CytoLyt solution for 30 min at room temperature and centrifuged at 5,000 rpm for 5 min. Cell's pellet was solubilized in Thin prep-CytoLyt solution and then resuspended in PreservCyt solution (Thinprep) for 15 min and processed on a ThinPrep 2000 processor. The microscope slides harboring the processed cells were fixed with 96% ethanol, washed twice with distilled water, and blocked in 0.5% hydrogen peroxide/methanol for 10 min. Immunostainings were performed using a Leica Bond-MAX automated IHC and ISH *Stainer* in accordance with the manufacturer's instructions. Briefly, the slides were washed with bond wash solution (Biosystems, USA) and the antigens retrieval procedure was performed in accordance with Bond heat standard protocol (ER1) using citric buffer, pH 6, for 30 min at 95°C. Slides are washed and treated with post-primary solution for 8 min. Next, the slides were incubated in Polymer AP for 8 min and washed. Then, the mixed Diaminobenzidine Refine solution was applied for 10 min. Samples were counterstained with water hematoxylin for 5 min. Non-immune immunoglobulins of the same isotype as the primary antibodies were used as a control for each experiment.

2.8. RNA Extraction, cDNA Synthesis and qRT-PCR Reactions

Total RNA was extracted from cells using the total RNA purification Trizol Reagent (Life Technologies) in accordance with the manufacturer's protocol. RNA was quantified using a NanoDrop 2000c spectrophotometer (Thermo Fisher Scientific, Waltham, MA, USA) and RNA integrity was evaluated by 2% agarose gel electrophoresis [42]. First strand cDNA was synthesized from 200 ng RNA by reverse transcription using QuantiTect Reverse Transcription Kit (Qiagen Inc., Germantown, MD, USA). qRT-PCR reactions were performed as previously described [43; 44]. Primer pairs with equivalent efficiencies (Table 1) were selected from Primer Bank [45] or designed using primer-BLAST tools (http://www.ncbi.nlm.nih.gov/tools/primer-blast/) [46]. qRT-PCR was performed on the ABI7900 system (Applied Biosystems, USA) using SDS software. Relative gene expression was calculated using a comparative Ct method as previously described [47; 48]. Results were normalized to GAPDH CT-values, and the relative expressions were determined relative to control or Day 0 undifferentiated cells. Data show Means ± standard deviation. Total liver RNA (Life Technologies) was used as a positive control.

2.9. Hepatic Biochemical Analysis of Supernatants

Hepatic enzymatic profiles for aspartate transaminase (AST), alanine aminotransferase (ALT), alkaline phosphatase (ALP) and gamma-glutamyl transferase (GGT) were analyzed from supernatant extracts of the different samples. The activity was measured by specific colorimetric detection kit (Linear Chemicals, Spain) at 25°C in accordance with the manufacturer's instructions. The specific compounds used for the following kinetics of the reaction were as the following: for AST and ALT, the oxidation of NADH was measured at light *absorbance* at 340 nm; for ALP, the formation of 4-Nitrophenol was measured at

light *absorbance* at 405 nm; for GGT, the catalysis of γ-glutamyl-3-carboxy-4-nitroanilide was measured at light *absorbance* at 410 nm using spectrophotometer (Thermo Fisher Scientific).

2.10. Cytochrome P450 3A4 Metabolic Activity Assay

Cytochrome P450 3A4 (CYP3A4) enzyme activity assays were assessed by measurement of luciferase activity using P450-Glo CYP3A4 assay kit (Promega, Madison, WI, USA) in accordance with the manufacturer's instructions. Differentiated and undifferentiated control DPPSC cells were treated with 25 µM Rifampicin for 48 hrs. Then, treated cells were incubated with a fresh serum-free medium containing 50 µM Luciferin PFBE with or without 5 µM Erythromycin (an antagonist) for 30 min at 37°C. Then, 50 µL of the medium was transferred into a 96-well plate and mixed with 50 µL luciferin detection reagent to initiate the luminescent reaction. After 20 min incubation at room temperature, the luminescence was measured with a luminometer (biotek, USA).

2.11. Albumin Assay

Alb secretion was measured by using an Albumin Fluorescence Assay Kit (Fluka, USA) in accordance with the manufacturer's instructions. Briefly, a calibration curve was generated using the kit- supplied standard human Alb concentrations. These calibration samples were mixed with albumin Blue 580 reagent pre-diluted in buffer solution. The fluorescence was measured in a spectrophotometer (absorbance wavelength peal, λex = 600 nm, and λem = 630 nm. Differentiated and undifferentiated control DPPSCs were mixed with the same reagents and the fluorescence was measured under the same conditions. Results are extrapolated to the calibration curve. Conditioned medium was collected from equivalent numbers of cells.

2.12. Periodic acid–Schiff staining for glycogen accumulation

At differentiation day 22, differentiated and undifferentiated control DPPSCs were fixed in 4% formaldehyde for 15 min at room temperature. After two washing steps with PBS, cells were incubated in 1% periodic acid for 5 min, then washed with distilled water. The treated cells were incubated with Schiff's reagent (Sigma-Aldrich) for 15 min. After a 10 min wash in distilled water, hematoxylin counterstain solution was applied for 90 sec and washed with distilled water.

2.13. Statistical analysis

Results are reported as mean ± standard deviation (see Figure legend for specific details regarding the number of biological replicates, independent experiments, and technical replicates). Statistics were performed using the Statgraphics XVI software. The methods used were two-tailed Student's t-test and ANOVA for multiple factors. Values with $p < 0.05$ were considered statistically significant.

3. Results

3.1. Definitive Endoderm Induction

Using hESCs, iPSCs, and human umbilical cord Wharton's jelly mesenchymal stem cells, studies have shown that Act A is the main DE induction factor [49; 50; 51; 52]. Therefore, we tested this concept on DPPSCs using Act A with or without other signaling factors to delineate the optimal conditional media sufficient to mediate DE- differentiation. Quantitative real-time PCR (qRT-PCR) analyses demonstrated a differential induction of the DE genes Forkhead Boxes A1 and A2 (FoxA1 and FoxA2) at day 5. FoxA1 transcripts elevated significantly in conditional media supplemented with Act A and Wnt3A, 14-fold relative to undifferentiated cells (Fig 1A). Similar induction levels were also observed in cells treated with Act A alone or Act A and BMP4, but albeit to a lower degree (6- to 8-fold) and notable variations between differ-

ent experiments. Alternatively, FoxA2 expression levels were comparable in all tested condi-

tional media, with a statistically significant 20-fold induction in media supplemented with Act

A alone, and with Wnt3A or BMP4. Remarkably, conditional media supplemented with Act A

and bFGF or FGF4 failed to significantly upregulate the studied DE-genes. Moreover, qRT-PCR

analyses indicated an elevation in the transcript levels of GATA4, with a 40 to 60-fold increase

relative to undifferentiated cells, post-treatment with Act A alone or in the presence of

Wnt3A or BMP4 (Fig. 1A). In contrary, no GATA4 transcripts were detected in conditional me-

dia containing the three signaling factors together (Fig. 1A). Noticeably, Act A failed to induce

GATA4 in media supplemented with either bFGF or FGF4 (Fig. 1A).

Worthy to note that at day 5, no significant upregulation of visceral endoderm and the early hepatogenic marker alpha-fetoprotein (AFP) in cells treated with the studied conditional media, suggesting that Act A and other signaling molecules do not support visceral endoderm lineage differentiation and are not sufficient for hepatogenic specification.

Taking together, the increase in expression levels of bona-fide DE- markers in conditional media supplemented with Act A alone or with Wnt3A suggests that our approaches using chemically defined- differentiation conditions has successfully enriched the DE- lineage of DPPSCs. Due to the experimental variations in the transcript levels of the DE- genes in cells treated with conditional media supplemented with containing Act A alone, therefore, the fol-lowing experiments were carried out using Act A and WNT3A as the main induction cytokines.

It is well documented that of biomaterials in cell cultures supports viability, differentia-tion, maturation, and long-term function [36; 53]. To further improve the differentiation pro-tocol and study the influence of biomaterials on DPPSCs mediated DE- indication, we perused DE- differentiation under three different conditions: cells were cultured on plates precoated with collagen I, fibronectin, or co-cultured with HepG2 cells for 5 days.

Interestingly, the DE-marker gene expression was significantly elevated in DPPSCs differ-entiated on the scaffolds (Fig. 1B) versus on no scaffolds (Fig. 1A). Relative to DPPSCs differ-entiated on plain plates (Fig. 1A), FOXA1 transcript levels in cells differentiated on collagen I or fibronectin were 3- to 4-folds higher, respectively. Similarly, the expression levels of FOXA2 mRNA were elevated by 1.5- to 3-fold when cells were differentiated on the scaffolds (Fig. 1B). qRT-PCR analyses of GATA4 transcripts were significantly enriched in cells differentiated with collagen I or fibronectin, 120-fold increase relative to undifferentiation cells, at day 5 (Fig 1B). No DE-marker gene expressions were detected in DPPSCs co-cultured with HepG2, sug-gesting a differentiation suppression effect of hepatic cells (Fig. 1B). Furthermore, the used

biomaterials did not facilitate prospective generation of visceral endoderm cells and did not

support early hepatogenic specification, since AFP transcripts were not detected. In summary,

biomaterial usage enhanced DPPSCs- mediated DE-differentiation. Since collagen I is an af-

fordable, inexpensive substrate, and commonly used in stem cell studies, we use it as the

scaffold to induce hepatogenic differentiation.

3.2. Hepatic Specification and Maturation.

We investigated the developmental progress in the hepatogenic differentiation potential

of DPPSCs using two-step protocol, which is schematically illustrated and presented in Fig. 2.

At day 5 post DE-induction, the conditional media were supplemented with FGF4 and HGF, for

8 days, to enrich for hepatic cells specification, followed by 8 days culture in HCM supple-

mented with hepatogenic inducers such as OSM and Dex in addition to the growth factors

(Fig. 2).

As we have previously reported, DPPSCs grow as a flat monolayer and exhibit a small

spindle-shaped fibroblast-like morphology when cultured on polystyrene tissue culture plates

or scaffolds (Fig. 3A) [25]. During the differentiation induction period, DPPSCs morphology

further changed to a round or epithelioid shape, a characteristic of DE- cells Fig 3 B) [54]. At

day 7, the cells became flat and tightly packed; and by day 22, they adopted a hepatocyte-like

morphology (Fig 3 C-E). In contrast, undifferentiated cells cultured in regular growth medium

containing FBS, leaned toward an elongated spindle-shaped fibroblastic morphology (Fig. 3F).

At the molecular level, gene expression studies revealed a distinct early posterior foregut

signature at day 7 of DPPSCs differentiation. The transcript levels of hepatic nuclear factor 6

(HNF6) and cytokeratin 19 (CK19) were notably elevated at day 7, relative to undifferenced

cells (Fig. 4), indicating a significant commitment toward the foregut tube. Although the tran-

scripts of GATA4 were upregulated at day 5 (Fig. 1B), simultaneously with the DE- markers, its

elevated expression levels were maintained until day 7. Unlike, FoxA1 mRNA levels which were significantly reduced at day 7. This observation is likely due to a possible commitment of a small cell fraction to DE- lineage and/or a significant profile of these genes in the early posterior foregut (Fig. 4) [52; 55].

Next, the developmental progress of differentiated DPPSCs toward hepatogenic specification was investigated at day 14 of treatment with conditional media. A notable reduction in the expression levels of the foregut tube markers was observed, concomitant with a significant increase in AFP transcripts (Fig. 4), which was not observed in earlier stages of differentiation, indicating a progression toward early hepatic specification. In addition, moderate levels of hepatic markers' transcripts were noticed at day 14 (Fig. 4), including hepatic nuclear factor 4α (HNF4α), glucose 6-phosphate (G6P), and Cholesterol 7 alpha-hydroxylase (CYP7A1).

Simultaneously, eight days treatment of these committed DPSSCs with HCM supplemented hepatic induction molecules resulted in a significantly exaggeration in the expression levels of the hepatic markers at differentiation day 22. Relative to undifferentiated DPPSCs, the HNF4α and Alb transcripts were 40-and 15-fold higher, AAT and CYP7A1 transcripts were 20-fold higher, and G6P mRNA levels were 100-fold higher at day 22 (Fig. 4). Taking together, we implemented an optimized 3-stage protocol that directed a stepwise differentiation of DPPSCs to the hepatocyte-like cell populations.

3.3. DPPSCs-derived hepatocyte-like cells demonstrate hepatic function:

Next, to corroborate the qRT-PCR analysis and evaluate the differentiation efficiency, we ascertained the protein expression of the hepatogenic markers by immuno-localization and functional analysis. Immunofluorescence with specific antibodies directed against AAT and Alb proteins revealed a significant expression of these hepatic markers at day 22 of DPPSCs differentiation (Fig. 5A-B).

Mature hepatocytes are characterized by their ability to store glycogen [56]. Interestingly, most of the differentiated DPPSCs showed a significant positive periodic acid Schiff (PAS) staining in the cytoplasm, albeit to a different degree, at Day 22 (Fig. 5C), which is indicative for glycogen storage, and the generation of mature hepatocyte-like cells. Alternatively, no PAS staining was detected in undifferentiated cells (Fig. 5D).

Then, we investigated the activity of hepatic metabolic enzymes. Time course studies performed on protein extracts from differentiated DPPSCs for GGT, AST, ALP, and ALT activities revealed a time-dependent upregulation of liver metabolic enzymes which beaked at day 18 of differentiation (Fig. 6A-D, respectively). Furthermore, the enzymatic activity of CYP3A4 was significantly induced in differentiated DPPSCs post treatment with rifampicin, 5-fold relative to untreated cells. Co-treatment with erythromycin, a CYP3A4 inhibitor, significantly down-regulated the observed enzymatic action against rifampicin (Fig. 6E).

Alb secretion was assayed at the time course of the differentiation process. As shown in Fig. 6F, at earlier stages of hepatogenesis, day 16 of differentiation, a limited amount of secreted Alb was detected in the media. However, at differentiation day 19 and 22, the amount of secreted albumin was triplicated into 3.5 ug/mg protein (Fig. 6F). Taking these observations together; the described differentiation protocol (Fig. 2) efficiently induced the generation of DPPSCs mediated functional hepatocyte-like cells and indicates that DPPSCs are a useful cell models for hepatogenesis.

4. Discussion

The current strategy in stem cell differentiation protocols is to mimic cellular signaling events associated with the embryonic developmental process for the lineage of interest [8, 47, 48]. However, *in vitro* cell differentiation gives a broad range of outcome lineages, thus, the development of a protocol that significantly improves a targeted cell type is of particular

interest [49]. Targeted differentiation requires culturing the cells in chemically well-defined media, which regulate cell signaling pathways, resulting in the generation of cell type of interest. This approach has been successfully applied to hESCs and iPSCs [51–53]; but has yet to be applied to other pluripotent cell types, including adult DPPSCs.

In this study, we applied an optimized 3D- 3-stages protocol directed toward a stepwise differentiation of DPPSCs to hepatocyte-like cells. For efficient differentiation, cells were checked at each developmental stage using lineage-specific markers. Initially, DPPSCs were cultured for 5 days in differentiation media supplemented with Act A, with or without Wnt3a, BMP4, or the growth factors bFGF and FGF4. Interestingly, Act A alone was sufficient to induce DE- differentiation. Our results are in accordance with the reported role of Act A inducing DE from pluripotent stem cells [55; 57]. It is well known that nodal signaling is required for endoderm specification during the gastrulation [58]. Furthermore, DE- induction depends on the duration of exposure to the stimulation factors; Agarwal *et. al.* reported that 5-days incubations are sufficient to induce DE [49].

Media supplemented with Act A and Wnt3A were superior in inducing DE- lineage [57; 59]. Using hESCs, Mathew *et. al.* observed an enhanced DE induction using Act A and Wnt3A supplemented media compared to media contained Act A alone [51]. Like hESC, in this study, we observed an improved induction of DE- markers in DPPSCs cultured with Act A and Wnt3A. Alternatively, the growth factors bFGF or FGF4 suppressed Act A- mediated DE induction [60]. Noteworthy, FGF family members are multifunctional factors. They can maintain hESC pluripotency [61] or DE- mediator [62], depending on the associated conditions.

During liver development, BMPs signaling is crucial; however, their functional role during *in vitro*- adult stem cell differentiation is not clear. Cultural media supplemented with BMP2 and BMP4 fail to improve hepatogenesis from adipose tissue stem cell [63]. In accordance

with the other studies, our results confirmed BMP4 signaling has no functional role during

DPPSCs differentiation into DE reconciled by Act A induction. Otherwise, studies have shown

that a combination of Act A and BMP4 signaling induce mesoderm formation [64]. Notably,

DE induction was further enhanced using three-dimensional (3D) scaffolds. Treatment with

either collagen or fibronectin upregulated DE markers indicated that 3D enhances the cellular

programming potential, which is in accordance with previously reported observations using

ESCs, iPSCs and other mesenchymal stem cells [36; 37; 65].

The use of minimal amount of FBS in differentiation media has been previously reported

to inhibit PI3K signaling [66], which is necessary to enhance ACT A-mediated DE generation.

Using 0.5% FBS during the first 5 days of DE induction, D'Amour and her team observed an

enhancement in the expression of Sox17 or FoxA2 in hESC [57]. In our protocol, we used 0.5%

FBS-media for the first 3 days of differentiation. Then, it was replaced with 2% KOSR for an

extra 4 days, a serum that has been proven to be effective and suitable with defined compo-

nents [49].At differentiation days 7-22 , KOSR was removed and hepatogenic lineage induc-

tion was carried on with serum-free conditional culture media. As described in schematic Fig.

2, the growth factors FGF4 and HGF were used to induce hepatogenesis, followed by the us-

age of OSM with Dex for the hepatocytes' maturation. The role of FGF4 and HGF in hepato-

cyte specification has been previously reported for both Wharton's jelly- and bone marrow

mesenchymal stem cells [67; 68; 69]. In accordance, both FGF4 and HGF were required to

mediate hepatic specification in DPPSCs. We observed an improved expression of several he-

patic markers including immature and mature markers at day 13 of differentiation.

Studies have shown that the interleukine-6 family cytokine (OSM) is required for hepato-

cyte maturation in combination with glucocorticoids, such as Dex [70; 71]. Additionally, it is

well documented that HGF, EGF, and OSM have decisive effects on the maintenance of pri-

mary human hepatocytes *in vitro* [72]. The combination of HGF, OSM, and Dex is widely used in protocols to differentiate stem cells into hepatocytes [49; 73]. In our protocol, we applied the maturation factors OSM and Dex at day 13 posting commitment to the hepatic fate. Consequently, we documented hepatocyte-like cell maturation using cellular and molecular techniques. After 15 days exposure to media containing OSM and Dex, we observed changes in cell morphology with hepatocyte- like structures, which was associated with Alb secretion, and the activation of hepatic specific enzymes.

After the establishment of the most optimal protocol used for DPPSC, we performed the tests that were available for us to prove the efficacy of the differentiation of DPPSC into hepatocyte-like cells. At the molecular level, differentiated DPPSC expressed AAT, Alb, and G6P transcripts, genes that are markers for mature hepatocytes. Moreover, the protein expression of these markers was also documented in our study. Cellular morphological changes were also detected, and mature cells acquired polygonal-shape typical for hepatocytes [74]. In addition, DPPSCs mediated hepatocyte-like cells were functional. These cells stored glycogen, secreted Alb, and other metabolic enzymes as efficient as other stem cell models used for hepatogenesis including MSCs [39] and hESCs[75; 76]. Furthermore, differentiated DPPSCs have shown an improved Cyp3A4 enzymatic activity, suggesting that the generated hepatocyte-like cells can be used as a model for drug metabolism, which would be useful for the pharmaceutical industry [77].

5. Conclusions

We designed a novel protocol to differentiate DPPSCs into functional hepatocyte-like cells. Our protocol is a directed differentiation protocol mimicking the stepwise process observed during embryonic liver development. Using DPPSCs as a pluripotent-like stem cell

model suggesting their potential usage for liver regenerative medicine and prospective treatments to be developed in the future. However, for clinical applications, there are still many studies that are needed to be improved, yet the generated functional DPPSCs-hepatocyte-like cells can be used as a model for drug screening, hepatic metabolism studies, and hepatic disease applications.

Data Availability: All data are available.

Conflicts of Interest: The authors declare no conflicts of interest.

Funding Statement: This study was funded by the Universitat International de Catalunya (UIC) and the Agència de Gestió d'ajuts Universitaris de Recerca, Generalitat de Catalunya project number SGR 1060 for MA, and Kuwait Foundation for the Advancement of Sciences (KFAS) and Dasman Diabetes Institute under projects number RA-2013-009 for AAM. CGR, EMS, and RNT were funded by the predoctoral grant Junior Faculty award from The Obra Social, La Caixa, and UIC.

Acknowledgment: Authors would like to thank Dr. Cámara Vallejo at Oral and Maxillofacial Surgery Department, Hospital Clinico de Barcelona, Barcelona, Spain, for patient's recruitment and molar extractions.

Abbreviations: AST: Aspartate aminotransferase, BAL: Bioartificial liver, CGH: Chromosome genomic, hybridization, DEX: Dexamethasone, DPMSC: Dental pulp mesenchymal stem cells, DPPSC: Dental pulp pluripotent stem cells, EB: Embryoid bodies, ECM: Extracellular matrix, ESDL: End-stage liver disease , GGT: Gammaglutamyl transferase, HB: Hepatoblasts, HCC:

Hepatocellular carcinoma, HT: Hepatocyte transplantation, MSC: Mesenchymal stromal cells,

OLT: Orthotopic liver transplantation , OSM: Oncostatin M

Authors' contribution: C.G.R., Collection and assembly of data and first manuscript draft;

S.M. and S.A.D., Data analysis and interpretation; M.A.A. and C.M., Statistic analysis; E.F.P.

and E.F.A, Provision of study patients and cells isolation; M.B., A.A.M and M.A., Conception

and design, collection and assembly of data, data analysis and Interpretation, manuscript

writing, and final approval of the Manuscript.

Figures Legend:

Figure 1. Molecular evaluation of different signaling factors and biomaterials on the genera-
tion of definitive endoderm lineage. qRT-PCR was performed for the indicated genes at day 5
of DPPSCs differentiation into DE lineage. A. Cells treated with differentiation conditional me-
dia supplemented with indicated signaling factors. B. Cells were seeded on precoated plates
with/without the described scaffolds or co-cultured with HepG2 cells. Results were normal-
ized to GAPDH and expressed relative to undifferentiated cells at day. Data are shown as
mean ± SD (n = 4). *p < 0.05.

Figure 2. The experimental protocol for DPPSCs mediated- hepatocyte-like cell generation.
Schematic representation of the differentiation protocol including the key manipulated sig-
naling pathways and the different molecular and functional studies performed.

Figure 3. Changes of cellular morphological during the time course of DPPSCs- mediated
hepatocyte-like. Phage contrast representative images of DPPSCs at different stages of dif-
ferentiation from day 0 to day 22 and undifferentiated cells at day 22. Scale A. 100X magnifi-
cation. B - F. 200X magnification.

Figure 4. Molecular and cellular evaluation of DPPSCs- mediated hepatocyte-like cells. Time
course qRT-PCR analysis for lineage-specific gene expression. The differentiation was per-
formed using the previously described experimental protocol. Total RNA was harvested from
the cells at the indicated days for gene expression. Results were normalized to GAPDH and
expressed relative to DPPSCs at day 0, *p < 0.05. # p < 0.05 relative to Day 7. Data are shown
as mean ± SD (n = 4).

Figure 5. Protein expression and glycogen storage of DPPSCs- mediated hepatocyte-like
cells. A. and B. Representative fluorescence microscopy images of DPPSCS mediated hepato-
cyte-like cells at day 22, immunoassayed with antibodies directed against A. AAT (AF594, red)
and B. ALB (FITC, green) proteins. Hoechst nuclear marker in blue. 200X magnification. C. and

D. Representative light microscope images of differentiated and undifferentiated DPPSCs at day 22, respectively, stained with periodic acid Schiff (PAS) for glycogen storage detection. 100X magnification.

Figure 6. Functional analysis of the generated DPPSCs- mediated hepatocyte-like cells. Time course hepatocyte enzymatic activity of A. GGT, B. AST, C. APF, D. ALT were performed for the differentiated DPPSCs. A notable induction was observed at day 18. E. CYP3A4 activity was measured by luminescence in DPPSCs at day 0 and day 22 of differentiation using rifampicin in the presence or absence of the antagonist erythromycin. F. Alb secretion assay was performed at a time course differentiated DPPSCs, elevated level of secreted Alb was detected at day 19 and later. All measurements are relative to DPPSCs at Day 0, * $p < 0.05$. # $p < 0.05$ relative to rifampicin treatment alone. Data are shown as mean ± SD (n = 4).

References

[1] A.M. Zorn, Liver development, StemBook, Cambridge (MA), 2008.

[2] M. Blachier, H. Leleu, M. Peck-Radosavljevic, D.C. Valla, and F. Roudot-Thoraval, The burden of liver disease in Europe: a review of available epidemiological data. Journal of hepatology 58 (2013) 593-608.

[3] M.F. Dawwas, A.E. Gimson, J.D. Lewsey, L.P. Copley, and J.H. van der Meulen, Survival after liver transplantation in the United Kingdom and Ireland compared with the United States. Gut 56 (2007) 1606-13.

[4] J. Neuberger, An update on liver transplantation: A critical review. J Autoimmun 66 (2016) 51-9.

[5] V. Iansante, R.R. Mitry, C. Filippi, E. Fitzpatrick, and A. Dhawan, Human hepatocyte transplantation for liver disease: current status and future perspectives. Pediatr Res 83 (2018) 232-240.

[6] J.A. Wertheim, H. Petrowsky, S. Saab, J.W. Kupiec-Weglinski, and R.W. Busuttil, Major challenges limiting liver transplantation in the United States. American journal of transplantation : official journal of the American Society of Transplantation and the American Society of Transplant Surgeons 11 (2011) 1773-84.

[7] A. Dhawan, R.R. Mitry, and R.D. Hughes, Hepatocyte transplantation for liver-based metabolic disorders. J Inherit Metab Dis 29 (2006) 431-5.

[8] F. Wang, L. Zhou, X. Ma, W. Ma, C. Wang, Y. Lu, Y. Chen, L. An, W. An, and Y. Yang, Monitoring of intrasplenic hepatocyte transplantation for acute-on-chronic liver failure: a prospective five-year follow-up study. Transplant Proc 46 (2014) 192-8.

[9] M. Ott, and J.V. Castell, Hepatocyte transplantation, a step forward? Journal of hepatology 70 (2019) 1049-1050.

[10] S. Gronthos, M. Mankani, J. Brahim, P.G. Robey, and S. Shi, Postnatal human dental pulp stem cells (DPSCs) in vitro and in vivo. Proc Natl Acad Sci U S A 97 (2000) 13625-30.

[11] F. Ferro, R. Spelat, A.P. Beltrami, D. Cesselli, and F. Curcio, Isolation and characterization of human dental pulp derived stem cells by using media containing low human serum percentage as clinical grade substitutes for bovine serum. PLoS One 7 (2012) e48945.

[12] V. Tirino, F. Paino, A. De Rosa, and G. Papaccio, Identification, isolation, characterization, and banking of human
dental pulp stem cells. Methods Mol Biol 879 (2012) 443-63.
[13] W. Zhang, X.F. Walboomers, S. Shi, M. Fan, and J.A. Jansen, Multilineage differentiation potential of stem cells
derived from human dental pulp after cryopreservation. Tissue Eng 12 (2006) 2813-23.
[14] N. Pilbauerova, T. Soukup, T. Suchankova Kleplova, J. Schmidt, and J. Suchanek, The Effect of Cultivation
Passaging on the Relative Telomere Length and Proliferation Capacity of Dental Pulp Stem Cells.
Biomolecules 11 (2021).
[15] L. Rambhatla, C.P. Chiu, P. Kundu, Y. Peng, and M.K. Carpenter, Generation of hepatocyte-like cells from human
embryonic stem cells. Cell transplantation 12 (2003) 1-11.
[16] Y.F. Chen, C.Y. Tseng, H.W. Wang, H.C. Kuo, V.W. Yang, and O.K. Lee, Rapid generation of mature
hepatocyte-like cells from human induced pluripotent stem cells by an efficient three-step protocol.
Hepatology 55 (2012) 1193-203.
[17] K. Si-Tayeb, F.K. Noto, M. Nagaoka, J. Li, M.A. Battle, C. Duris, P.E. North, S. Dalton, and S.A. Duncan, Highly
efficient generation of human hepatocyte-like cells from induced pluripotent stem cells. Hepatology 51
(2010) 297-305.
[18] K.D. Lee, T.K. Kuo, J. Whang-Peng, Y.F. Chung, C.T. Lin, S.H. Chou, J.R. Chen, Y.P. Chen, and O.K. Lee, In vitro
hepatic differentiation of human mesenchymal stem cells. Hepatology 40 (2004) 1275-84.
[19] A. Banas, T. Teratani, Y. Yamamoto, M. Tokuhara, F. Takeshita, G. Quinn, H. Okochi, and T. Ochiya, Adipose
tissue-derived mesenchymal stem cells as a source of human hepatocytes. Hepatology 46 (2007) 219-28.
[20] M.J. Seo, S.Y. Suh, Y.C. Bae, and J.S. Jung, Differentiation of human adipose stromal cells into hepatic lineage in
vitro and in vivo. Biochemical and biophysical research communications 328 (2005) 258-64.
[21] N. Ishkitiev, K. Yaegaki, B. Calenic, T. Nakahara, H. Ishikawa, V. Mitiev, and M. Haapasalo, Deciduous and
permanent dental pulp mesenchymal cells acquire hepatic morphologic and functional features in vitro. J
Endod 36 (2010) 469-74.
[22] N. Ishkitiev, K. Yaegaki, T. Imai, T. Tanaka, T. Nakahara, H. Ishikawa, V. Mitev, and M. Haapasalo, High-purity
hepatic lineage differentiated from dental pulp stem cells in serum-free medium. Journal of endodontics 38
(2012) 475-80.
[23] A. Kumar, V. Kumar, V. Rattan, V. Jha, A. Pal, and S. Bhattacharyya, Molecular spectrum of secretome regulates
the relative hepatogenic potential of mesenchymal stem cells from bone marrow and dental tissue. Sci Rep 7
(2017) 15015.
[24] M. Atari, C. Gil-Recio, M. Fabregat, D. Garcia-Fernandez, M. Barajas, M.A. Carrasco, H.S. Jung, F.H. Alfaro, N.
Casals, F. Prosper, E. Ferres-Padro, and L. Giner, Dental pulp of the third molar: a new source of
pluripotent-like stem cells. J Cell Sci 125 (2012) 3343-56.
[25] M. Atari, M. Barajas, F. Hernandez-Alfaro, C. Gil, M. Fabregat, E. Ferres Padro, L. Giner, and N. Casals, Isolation
of pluripotent stem cells from human third molar dental pulp. Histology and histopathology 26 (2011)
1057-70.
[26] M. Atari, J. Caballe-Serrano, C. Gil-Recio, C. Giner-Delgado, E. Martinez-Sarra, D.A. Garcia-Fernandez, M. Barajas,
F. Hernandez-Alfaro, E. Ferres-Padro, and L. Giner-Tarrida, The enhancement of osteogenesis through the
use of dental pulp pluripotent stem cells in 3D. Bone 50 (2012) 930-41.
[27] E. Martinez-Sarra, S. Montori, C. Gil-Recio, R. Nunez-Toldra, D. Costamagna, A. Rotini, M. Atari, A. Luttun, and
M. Sampaolesi, Human dental pulp pluripotent-like stem cells promote wound healing and muscle
regeneration. Stem Cell Res Ther 8 (2017) 175.

[28] R. Nunez-Toldra, P. Dosta, S. Montori, V. Ramos, M. Atari, and S. Borros, Improvement of osteogenesis in dental
pulp pluripotent-like stem cells by oligopeptide-modified poly(beta-amino ester)s. Acta Biomater 53 (2017)
152-164.
[29] F. Ferro, R. Spelat, F. D'Aurizio, E. Puppato, M. Pandolfi, A.P. Beltrami, D. Cesselli, G. Falini, C.A. Beltrami, and F.
Curcio, Dental pulp stem cells differentiation reveals new insights in Oct4A dynamics. PLoS One 7 (2012)
e41774.
[30] F.N. Faruqu, S. Zhou, N. Sami, F. Gheidari, H. Lu, and K.T. Al-Jamal, Three-dimensional culture of dental pulp
pluripotent-like stem cells (DPPSCs) enhances Nanog expression and provides a serum-free condition for
exosome isolation. FASEB Bioadv 2 (2020) 419-433.
[31] R. Nunez-Toldra, E. Martinez-Sarra, C. Gil-Recio, M.A. Carrasco, A. Al Madhoun, S. Montori, and M. Atari, Dental
pulp pluripotent-like stem cells (DPPSC), a new stem cell population with chromosomal stability and
osteogenic capacity for biomaterials evaluation. BMC Cell Biol 18 (2017) 21.
[32] R.V. Nelakanti, N.G. Kooreman, and J.C. Wu, Teratoma formation: a tool for monitoring pluripotency in stem cell
research. Curr Protoc Stem Cell Biol 32 (2015) 4A.8.1-4A.8.17.
[33] H. Hentze, P.L. Soong, S.T. Wang, B.W. Phillips, T.C. Putti, and N.R. Dunn, Teratoma formation by human
embryonic stem cells: evaluation of essential parameters for future safety studies. Stem Cell Res 2 (2009)
198-210.
[34] K. Zeevaert, M.H. Elsafi Mabrouk, W. Wagner, and R. Goetzke, Cell Mechanics in Embryoid Bodies. Cells 9
(2020).
[35] C. Gil-Recio, S. Montori, C. Vallejo, S.A. Demour, E. Ferrés-Padró, M. Barajas, C. Martin, A.A. Madhoun, and M.
Atari, Direct differentiation of Dental Pulp Pluripotent-like Stem Cells differentiation into Hepatocyte-like
Cells. bioRxiv (2020) 2020.12.09.418780.
[36] S. Nakai, I. Shibata, T. Shitamichi, H. Yamaguchi, N. Takagi, T. Inoue, T. Nakagawa, J. Kiyokawa, S. Wakabayashi,
T. Miyoshi, E. Higashi, S. Ishida, N. Shiraki, and S. Kume, Collagen vitrigel promotes hepatocytic
differentiation of induced pluripotent stem cells into functional hepatocyte-like cells. Biol Open 8 (2019).
[37] C.H. Rasmussen, D.R. Petersen, J.B. Moeller, M. Hansson, and M. Dufva, Collagen Type I Improves the
Differentiation of Human Embryonic Stem Cells towards Definitive Endoderm. PLoS One 10 (2015) e0145389.
[38] A. Al Madhoun, S.K. Marafie, D. Haddad, M. Melhem, M. Abu-Farha, H. Ali, S. Sindhu, M. Atari, and F. Al-Mulla,
Comparative Proteomic Analysis Identifies EphA2 as a Specific Cell Surface Marker for Wharton's
Jelly-Derived Mesenchymal Stem Cells. Int J Mol Sci 21 (2020).
[39] F. Al-Rashed, S. Sindhu, H. Arefanian, A. Al Madhoun, S. Kochumon, R. Thomas, S. Al-Kandari, A. Alghaith, T.
Jacob, F. Al-Mulla, and R. Ahmad, Repetitive Intermittent Hyperglycemia Drives the M1 Polarization and
Inflammatory Responses in THP-1 Macrophages Through the Mechanism Involving the TLR4-IRF5 Pathway.
Cells 9 (2020) 1892.
[40] A. Al Madhoun, S. Alkandari, H. Ali, N. Carrio, M. Atari, M.S. Bitar, and F. Al-Mulla, Chemically Defined Conditions
Mediate an Efficient Induction of Mesodermal Lineage from Human Umbilical Cord- and Bone Marrow-
Mesenchymal Stem Cells and Dental Pulp Pluripotent-Like Stem Cells. Cell Reprogram 20 (2018) 9-16.
[41] A. Khadir, A. Tiss, J. Abubaker, M. Abu-Farha, I. Al-Khairi, P. Cherian, J. John, S. Kavalakatt, S. Warsame, A.
Al-Madhoun, F. Al-Ghimlas, N. Elkum, K. Behbehani, S. Dermime, and M. Dehbi, MAP kinase phosphatase
DUSP1 is overexpressed in obese humans and modulated by physical exercise. Am J Physiol Endocrinol
Metab 308 (2015) E71-83.
[42] S. Kochumon, A.A. Madhoun, F. Al-Rashed, R. Azim, E. Al-Ozairi, F. Al-Mulla, and R. Ahmad, Adipose tissue gene
expression of CXCL10 and CXCL11 modulates inflammatory markers in obesity: implications for metabolic
inflammation and insulin resistance. Ther Adv Endocrinol Metab 11 (2020) 2042018820930902.

[43] A. Maher, R. Nunez-Toldra, N. Carrio, E. Ferres-Padro, H. Ali, S. Montori, and A. Al Madhoun, The Effect of Commercially Available Endodontic Cements and Biomaterials on Osteogenic Differentiation of Dental Pulp Pluripotent-Like Stem Cells. Dent J (Basel) 6 (2018).

[44] S. Kochumon, A. Al Madhoun, F. Al-Rashed, R. Thomas, S. Sindhu, E. Al-Ozairi, F. Al-Mulla, and R. Ahmad, Elevated adipose tissue associated IL-2 expression in obesity correlates with metabolic inflammation and insulin resistance. Sci Rep 10 (2020) 16364.

[45] X.W. Wang, and B. Seed, A PCR primer bank for quantitative gene expression analysis. Nucleic Acids Res 31 (2003).

[46] J. Ye, G. Coulouris, I. Zaretskaya, I. Cutcutache, S. Rozen, and T.L. Madden, Primer-BLAST: a tool to design target-specific primers for polymerase chain reaction. BMC Bioinformatics 13 (2012) 134.

[47] A. Voronova, E. Coyne, A. Al Madhoun, J.V. Fair, N. Bosiljcic, C. St-Louis, G. Li, S. Thurig, V.A. Wallace, N. Wiper-Bergeron, and I.S. Skerjanc, Hedgehog Signaling Regulates MyoD Expression and Activity. Journal of Biological Chemistry 288 (2013) 4389-4404.

[48] A. Voronova, A. Fischer, T. Ryan, A. Al Madhoun, and I.S. Skerjanc, Ascl1/Mash1 Is a Novel Target of Gli2 during Gli2-Induced Neurogenesis in P19 EC Cells. PLoS One 6 (2011).

[49] S. Agarwal, K.L. Holton, and R. Lanza, Efficient differentiation of functional hepatocytes from human embryonic stem cells. Stem Cells 26 (2008) 1117-27.

[50] J. Cai, Y. Zhao, Y. Liu, F. Ye, Z. Song, H. Qin, S. Meng, Y. Chen, R. Zhou, X. Song, Y. Guo, M. Ding, and H. Deng, Directed differentiation of human embryonic stem cells into functional hepatic cells. Hepatology 45 (2007) 1229-39.

[51] S. Mathew, M. Jaramillo, X. Zhang, L.A. Zhang, A. Soto-Gutierrez, and I. Banerjee, Analysis of alternative signaling pathways of endoderm induction of human embryonic stem cells identifies context specific differences. BMC systems biology 6 (2012) 154.

[52] A. Al Madhoun, H. Ali, S. AlKandari, V.L. Atizado, N. Akhter, F. Al-Mulla, and M. Atari, Defined three-dimensional culture conditions mediate efficient induction of definitive endoderm lineage from human umbilical cord Wharton's jelly mesenchymal stem cells. Stem Cell Res Ther 7 (2016) 165.

[53] T. Takebe, N. Koike, K. Sekine, R. Fujiwara, T. Amiya, Y.W. Zheng, and H. Taniguchi, Engineering of human hepatic tissue with functional vascular networks. Organogenesis 10 (2014) 260-7.

[54] E.L. Calderon-Gierszal, and G.S. Prins, Directed Differentiation of Human Embryonic Stem Cells into Prostate Organoids In Vitro and its Perturbation by Low-Dose Bisphenol A Exposure. PLoS One 10 (2015) e0133238.

[55] P. Wang, K.D. McKnight, D.J. Wong, R.T. Rodriguez, T. Sugiyama, X. Gu, A. Ghodasara, K. Qu, H.Y. Chang, and S.K. Kim, A molecular signature for purified definitive endoderm guides differentiation and isolation of endoderm from mouse and human embryonic stem cells. Stem cells and development 21 (2012) 2273-87.

[56] C.M. Chinnici, F. Timoneri, G. Amico, G. Pietrosi, G. Vizzini, M. Spada, D. Pagano, B. Gridelli, and P.G. Conaldi, Characterization of Liver-Specific Functions of Human Fetal Hepatocytes in Culture. Cell transplantation 24 (2015) 1139-53.

[57] K.A. D'Amour, A.D. Agulnick, S. Eliazer, O.G. Kelly, E. Kroon, and E.E. Baetge, Efficient differentiation of human embryonic stem cells to definitive endoderm. Nature biotechnology 23 (2005) 1534-41.

[58] J.G. Robert Lanza, Brigid Hogan, Douglas Melton, Roger Pedersen, E. Donnall Thomas, James Thomson, Ian Wilmut, Essentials of Stem Cell Biology, 2009.

[59] S. Toivonen, K. Lundin, D. Balboa, J. Ustinov, K. Tamminen, J. Palgi, R. Trokovic, T. Tuuri, and T. Otonkoski, Activin A and Wnt-dependent specification of human definitive endoderm cells. Experimental cell research 319 (2013) 2535-44.

[60] R.E. Schwartz, J.L. Linehan, M.S. Painschab, W.S. Hu, C.M. Verfaillie, and D.S. Kaufman, Defined conditions for
development of functional hepatic cells from human embryonic stem cells. Stem cells and development 14
(2005) 643-55.

[61] S. Lotz, S. Goderie, N. Tokas, S.E. Hirsch, F. Ahmad, B. Corneo, S. Le, A. Banerjee, R.S. Kane, J.H. Stern, S. Temple,
and C.A. Fasano, Sustained levels of FGF2 maintain undifferentiated stem cell cultures with biweekly feeding.
PLoS One 8 (2013) e56289.

[62] N. Shiraki, T. Yoshida, K. Araki, A. Umezawa, Y. Higuchi, H. Goto, K. Kume, and S. Kume, Guided differentiation of
embryonic stem cells into Pdx1-expressing regional-specific definitive endoderm. Stem Cells 26 (2008)
874-85.

[63] A. Bonora-Centelles, R. Jover, V. Mirabet, A. Lahoz, F. Carbonell, J.V. Castell, and M.J. Gomez-Lechon, Sequential
hepatogenic transdifferentiation of adipose tissue-derived stem cells: relevance of different extracellular
signaling molecules, transcription factors involved, and expression of new key marker genes. Cell
transplantation 18 (2009) 1319-40.

[64] M.A. Laflamme, K.Y. Chen, A.V. Naumova, V. Muskheli, J.A. Fugate, S.K. Dupras, H. Reinecke, C. Xu, M.
Hassanipour, S. Police, C. O'Sullivan, L. Collins, Y. Chen, E. Minami, E.A. Gill, S. Ueno, C. Yuan, J. Gold, and C.E.
Murry, Cardiomyocytes derived from human embryonic stem cells in pro-survival factors enhance function
of infarcted rat hearts. Nature biotechnology 25 (2007) 1015-24.

[65] H.Y. Lin, C.C. Tsai, L.L. Chen, S.H. Chiou, Y.J. Wang, and S.C. Hung, Fibronectin and laminin promote
differentiation of human mesenchymal stem cells into insulin producing cells through activating Akt and ERK.
J Biomed Sci 17 (2010) 56.

[66] A.B. McLean, K.A. D'Amour, K.L. Jones, M. Krishnamoorthy, M.J. Kulik, D.M. Reynolds, A.M. Sheppard, H. Liu, Y.
Xu, E.E. Baetge, and S. Dalton, Activin a efficiently specifies definitive endoderm from human embryonic
stem cells only when phosphatidylinositol 3-kinase signaling is suppressed. Stem Cells 25 (2007) 29-38.

[67] X.Q. Kang, W.J. Zang, L.J. Bao, D.L. Li, T.S. Song, X.L. Xu, and X.J. Yu, Fibroblast growth factor-4 and hepatocyte
growth factor induce differentiation of human umbilical cord blood-derived mesenchymal stem cells into
hepatocytes. World journal of gastroenterology : WJG 11 (2005) 7461-5.

[68] R.E. Schwartz, M. Reyes, L. Koodie, Y. Jiang, M. Blackstad, T. Lund, T. Lenvik, S. Johnson, W.S. Hu, and C.M.
Verfaillie, Multipotent adult progenitor cells from bone marrow differentiate into functional hepatocyte-like
cells. The Journal of clinical investigation 109 (2002) 1291-302.

[69] C. Schmidt, F. Bladt, S. Goedecke, V. Brinkmann, W. Zschiesche, M. Sharpe, E. Gherardi, and C. Birchmeier,
Scatter factor/hepatocyte growth factor is essential for liver development. Nature 373 (1995) 699-702.

[70] A. Kamiya, T. Kinoshita, and A. Miyajima, Oncostatin M and hepatocyte growth factor induce hepatic maturation
via distinct signaling pathways. FEBS letters 492 (2001) 90-4.

[71] S. Hanada, H. Kayano, J. Jiang, N. Kojima, A. Miyajima, A. Sakoda, and Y. Sakai, Enhanced in vitro maturation of
subcultivated fetal human hepatocytes in three dimensional culture using poly-L-lactic acid scaffolds in the
presence of oncostatin M. The International journal of artificial organs 26 (2003) 943-51.

[72] J. Dong, C.F. Mandenius, M. Lubberstedt, T. Urbaniak, A.K. Nussler, D. Knobeloch, J.C. Gerlach, and K. Zeilinger,
Evaluation and optimization of hepatocyte culture media factors by design of experiments (DoE)
methodology. Cytotechnology 57 (2008) 251-61.

[73] X. Li, J. Yuan, W. Li, S. Liu, M. Hua, X. Lu, and H. Zhang, Direct differentiation of homogeneous human adipose
stem cells into functional hepatocytes by mimicking liver embryogenesis. Journal of cellular physiology 229
(2014) 801-12.

[74] P. Godoy, N.J. Hewitt, U. Albrecht, M.E. Andersen, N. Ansari, S. Bhattacharya, J.G. Bode, J. Bolleyn, C. Borner, J.
Bottger, A. Braeuning, R.A. Budinsky, B. Burkhardt, N.R. Cameron, G. Camussi, C.S. Cho, Y.J. Choi, J. Craig

Rowlands, U. Dahmen, G. Damm, O. Dirsch, M.T. Donato, J. Dong, S. Dooley, D. Drasdo, R. Eakins, K.S.
Ferreira, V. Fonsato, J. Fraczek, R. Gebhardt, A. Gibson, M. Glanemann, C.E. Goldring, M.J. Gomez-Lechon,
G.M. Groothuis, L. Gustavsson, C. Guyot, D. Hallifax, S. Hammad, A. Hayward, D. Haussinger, C. Hellerbrand,
P. Hewitt, S. Hoehme, H.G. Holzhutter, J.B. Houston, J. Hrach, K. Ito, H. Jaeschke, V. Keitel, J.M. Kelm, B.
Kevin Park, C. Kordes, G.A. Kullak-Ublick, E.L. LeCluyse, P. Lu, J. Luebke-Wheeler, A. Lutz, D.J. Maltman, M.
Matz-Soja, P. McMullen, I. Merfort, S. Messner, C. Meyer, J. Mwinyi, D.J. Naisbitt, A.K. Nussler, P. Olinga, F.
Pampaloni, J. Pi, L. Pluta, S.A. Przyborski, A. Ramachandran, V. Rogiers, C. Rowe, C. Schelcher, K. Schmich, M.
Schwarz, B. Singh, E.H. Stelzer, B. Stieger, R. Stober, Y. Sugiyama, C. Tetta, W.E. Thasler, T. Vanhaecke, M.
Vinken, T.S. Weiss, A. Widera, C.G. Woods, J.J. Xu, K.M. Yarborough, and J.G. Hengstler, Recent advances in
2D and 3D in vitro systems using primary hepatocytes, alternative hepatocyte sources and non-parenchymal
liver cells and their use in investigating mechanisms of hepatotoxicity, cell signaling and ADME. Archives of
toxicology 87 (2013) 1315-530.

[75] R. Ji, N. Zhang, N. You, Q. Li, W. Liu, N. Jiang, J. Liu, H. Zhang, D. Wang, K. Tao, and K. Dou, The differentiation of
MSCs into functional hepatocyte-like cells in a liver biomatrix scaffold and their transplantation into
liver-fibrotic mice. Biomaterials 33 (2012) 8995-9008.

[76] K. Subramanian, D.J. Owens, R. Raju, M. Firpo, T.D. O'Brien, C.M. Verfaillie, and W.S. Hu, Spheroid culture for
enhanced differentiation of human embryonic stem cells to hepatocyte-like cells. Stem cells and
development 23 (2014) 124-31.

[77] J. Jozefczuk, A. Prigione, L. Chavez, and J. Adjaye, Comparative analysis of human embryonic stem cell and
induced pluripotent stem cell-derived hepatocyte-like cells reveals current drawbacks and possible strategies
for improved differentiation. Stem cells and development 20 (2011) 1259-75.

Table 1. Oligonucleotide sequences of primers utilized for real-time qRT-PCR.

Gene	Forward primer (5'-3')	Reverse primer (5'-3')
GAPDH	TGCACCACCAACTGCTTAGC	GGCATGGACTGTGGTCATGAG
FoxA1	GCCTACTCCTCCGTCCCGGT	CCGGGGTCATGTTGCCGCTC
FoxA2	GCGACCCCAAGACCTACAG	GGTTCTGCCGGTAGAAGGG
Gata4	TCCCTCTTCCCTCCTCAAAT	TTCCCCTAACCAGATTGTCG
AFP	AGGGTGTAGCGCTGCAAACGA	TGTTCCTGGCCTTGGCAGCAT
Oct4A	GACAGGGGGAGGGGAGGAGCTAGG	CTTCCCTCCAACCAGTTGCCCCAAAC
HNF6	CTTAGCAGCATGCAAAAGGA	TGCGTTCATGAAGAAGTTGC
CK19	GAGGAAATCAGTACGCTGAG	GTTTCTGCCAGTGTGTCTTC
HNF4α	ACTACATCAACGACCGCCAGT	ATCTGCTCGATCATCTGCCAG
Alb	CCTTGGTGTTGATTGCCTTTGCTC	CATCACATCAACCTCTGGTCTCACC
G6P	TCAGCTCAGGTGGTCCTCTT	CCTCCTTAGGCAGCCTTCTT
AAT	TCGCTACAGCCTTTGCAATG	TTGAGGGTACGGAGGAGTTCC
CYP7A1	AGGACGGTTCCTACAACATC	CGATCCAAAGGGCATGTAGT

738   Figure 1

Figure 2

Figure 3

Figure 4

Figure 5

761   Figure 6

764