A single dose, BCG-adjuvanted COVID-19 vaccine provides sterilizing immunity against SARS-CoV-2 infection in mice

Claudio Counoupas[1,2#], Matt D. Johansen[3#], Alberto O. Stella[4], Duc H. Nguyen[3], Angela L. Ferguson[1,2] Anupriya Aggarwal[4], Nayan D. Bhattacharyya[2], Alice Grey[5], Karishma Patel[6], Rezwan Siddiquee[6], Erica L. Stewart[1], Carl G. Feng[2], Nicole G. Hansbro[3], Umaimainthan Palendira[1], Megan C. Steain[2], Bernadette M. Saunders[3], Jason K. K. Low[6], Joel P. Mackay[6], Anthony D. Kelleher[4], Warwick J. Britton[2,5], Stuart G Turville[4], Philip M. Hansbro[3*], James A. Triccas[1,7*]

[1]School of Medical Sciences, Faculty of Medicine and Health, The University of Sydney, Camperdown, NSW, Australia. [2]Tuberculosis Research Program at the Centenary Institute, The University of Sydney, Sydney, NSW, Australia. [3]Centre for Inflammation, Centenary Institute and University of Technology Sydney, Faculty of Science, School of Life Sciences, Sydney, NSW, Australia. [4]Kirby Institute, University of New South Wales, Sydney, NSW, Australia. [5]Department of Clinical Immunology, Royal Prince Alfred Hospital, Sydney, NSW, Australia. [6]School of Life and Environmental Sciences, The University of Sydney, Sydney, NSW 2006. [7]Charles Perkins Centre and Marie Bashir Institute for Infectious Diseases and Biosecurity, The University of Sydney, Camperdown, NSW, Australia.

[#]Authors contributed equally

*Correspondence to James A. Triccas (jamie.triccas@sydney.edu.au); Philip Hansbro (Philip.Hansbro@uts.edu.au)

ABSTRACT

Global control of COVID-19 requires broadly accessible vaccines that are effective against SARS-CoV-2 variants. In this report, we exploit the immunostimulatory properties of bacille Calmette-Guérin (BCG), the existing tuberculosis vaccine, to deliver a vaccination regimen with potent SARS-CoV-2-specific protective immunity. Combination of BCG with a stabilized, trimeric form of SARS-CoV-2 spike antigen promoted rapid development of virus-specific IgG antibodies in the blood of vaccinated mice, that was further augmented by the addition of alum. This vaccine formulation, BCG:CoVac, induced high-titre SARS-CoV-2 neutralizing antibodies (NAbs) and Th1-biased cytokine release by vaccine-specific T cells, which correlated with the early emergence of T follicular helper cells in local lymph nodes and heightened levels of antigen-specific plasma B cells after vaccination. Vaccination of K18-hACE2 mice with a single dose of BCG:CoVac almost completely abrogated disease after SARS-CoV-2 challenge, with minimal inflammation and no detectable virus in the lungs of infected animals. Boosting BCG:CoVac-primed mice with a heterologous vaccine further increased SARS-CoV-2-specific antibody responses, which effectively neutralized B.1.1.7 and B.1.351 SARS-CoV-2 variants of concern. These findings demonstrate the potential for BCG-based vaccination to protect against major SARS-CoV-2 variants circulating globally.

INTRODUCTION

The world has entered a critical stage in the continuing fight against COVD-19. The deployment of effective vaccines has had a profound impact in reducing cases and SARS-CoV-2 transmission in countries with high vaccine coverage[1,2]. However global cases are again on the rise, driven predominantly by a surge of infections in Europe, South America and the subcontinent due to the emergence of new SARS-CoV-2 variants. Two of the dominant variants of concern (VOCs), B.1.1.7 and B.1.351, both have increased transmissibility[3,4] and antibodies from convalescent patients and vaccinees have a reduced capacity to neutralize B.1.351[5,6]. Vaccination with one of the most widely deployed vaccines, ChAdOx1 nCoV-19, does not protect against mild-to-moderate COVID-19 due to the B.1.351 variant in South Africa[7]. A critical issue is ensuring the adequate supply of vaccines to the hardest-hit low and middle income countries, particularly due to complex logistical requirements (e.g. storage at low temperature for mRNA vaccines). The requirement for multiple doses for most approved vaccines is a barrier to rapid, mass vaccination and has necessitated changes in dosing schedules in some countries to ensure sufficient vaccine coverage[8]. Thus, ensuring the global supply of vaccines effective against emerging variants will be necessary to control the global COVID-19 pandemic.

One unique strategy is to 'repurpose' existing licensed vaccines for use against COVID-19. Significant interest has focussed on *Mycobacterium bovis* bacille Calmette-Guerin (BCG), the tuberculosis (TB) vaccine. Considerable data has been accumulated to show that BCG has beneficial, non-specific effects on immunity that affords protection against other pathogens, particularly respiratory infections[9]. Most recently, BCG vaccination was shown to protect against viral respiratory tract infections in the elderly (greater than 65 years old) with no significant adverse events[10]. This non-specific protective effect is attributed to the ability of BCG to induce 'trained immunity' i.e. reprogramming of innate immune responses to provide

heterologous protection against disease. For these reasons, a Phase 3, randomised controlled

trial in healthcare workers has commenced to determine if BCG vaccination can reduce the

incidence and severity of COVID-19 (the BRACE Trial)[9]. While that trial will determine if

BCG can reduce the impact on COVID-19 during the current pandemic, BCG does not express

SARS-CoV-2 specific antigens and thus, would not induce long-term immune memory.

Here, we have exploited the immunostimulatory properties of BCG to develop a SARS-

CoV-2 vaccine, BCG:CoVac, that combines a stabilized, trimeric form of the spike protein

with the alum adjuvant. BCG:CoVac stimulated SARS-CoV-2-specific antibody and T cell

responses in mice after a single vaccination, including the elicitation of high-titre NAbs.

Critically, a single dose was shown to protect mice against severe SARS-CoV-2, demonstrating

that BCG:CoVac is a highly immunogenic and promising vaccine candidate.

RESULTS

BCG vaccination promotes SARS-CoV-2 specific antibody and T cell

responses in mice

The immunostimulatory properties of BCG[11] led to us to test if the vaccine could serve as the

backbone for a unique vaccine platform against COVID-19. This was also supported by our

observation that prior BCG immunization could augment anti-spike IgG responses after

boosting with SpK formulated in Alhydrogel/alum (Alm^{SpK}) (Fig. S1). To determine if this

property of BCG could be used in a single vaccine formulation, we subcutaneously (s.c)

vaccinated mice with a single dose of BCG formulated with a stabilized, trimeric form of the

SARS-CoV-2 spike protein[12] and the titre of IgG2c or IgG1 anti-SpK antibodies was

determined at various timepoints post-immunization (Fig. 1a). While BCG vaccination resulted

in background levels of anti-SpK antibodies, titres were approximately 100-fold higher for both

antibody isotypes after BCG^{Spk} vaccination, and similar to those levels achieved with Alm^{SpK}

(Fig. 1b, 1c). Addition of alum to BCG^{Spk} (termed BCG:CoVac) further increased antibodies

titres, particularly IgG2c, which were significantly greater after BCG:CoVac vaccination

compared to mice immunized with either BCG or Alm^{SpK}, at all timepoints examined (Fig. 1b,

1c).

The IgG2c Ab isotype correlates with Th1-like immunity in C57BL/6 mice[13], and such

responses are considered necessary for effective protection against SARS-CoV-2 infection[14].

We therefore examined the frequency of IFN-γ-expressing T cells after a single dose of

BCG:CoVac at 2 weeks post-vaccination. BCG^{SpK} and BCG:CoVac induced the generation of

SpK-specific CD4$^+$ and CD8$^+$ T cells secreting IFN-γ (Fig. 1d, 1e), consistent with Th1

immunity observed after BCG vaccination[15]. The greatest response was observed after

vaccination with BCG:CoVac, with the numbers of IFN-γ-secreting T cells significantly

increased compared to vaccination with either BCG or Alum^{SpK}. Low levels of the

inflammatory cytokines IL-17 and TNF were observed after BCG:CoVac vaccination (Fig. 1e).

We further dissected vaccine-induced immunity by defining the cellular composition in

draining lymph nodes 7 days after vaccination. Both Alum^{SpK} and BCG:CoVac induced

appreciable   expansion   of   SpK-specific   germinal   centre   (GC)   B   cells

(CD19$^+$MHCII$^+$GL7$^+$CD38$^-$; Fig. 2a) and plasma B cells (CD19$^+$MHCII$^+$CD138$^+$; Fig. 2b).

Cells with a T follicular helper cell (Tfh) phenotype (CD4$^+$CXCR5$^+$BCL6$^+$) were apparent after

vaccination with Alum^{SpK} or BCG:CoVac, with Tfh frequency greatest in the latter group (Fig.

2c). The total numbers of GC B cells (Fig. 2d), plasma B cells (Fig. 2e) and Tfh cells (Fig. 2f)

were all significantly increased in BCG:CoVac-vaccinated mice compared to immunization

with Alum^{SpK}.

Overall, these data show that co-delivery of trimeric SpK antigen with BCG vaccination

promotes early and pronounced anti-SARS-CoV-2 immunity, and this is further enhanced with

the addition of alum.

High-titre, SARS-CoV-2 neutralizing antibodies after a single immunization with BCG:CoVac

The elicitation of GC B cell and Tfh responses after immunization with experimental SARS-CoV-2 vaccines correlate strongly with the induction of neutralizing antibodies (NAbs)[16]. Such NAbs are a key determinant of protection induced by current vaccines used in humans[17]. We therefore measured NAb levels after a single dose of BCG:CoVac. No NAbs were detected in the plasma of mice vaccinated with BCG (Fig. 3a). Surprisingly, NAb titres were at near background levels for mice vaccinated with BCG^{SpK} (Fig. 3a), despite the high levels of IgG Ab isotypes detected in these same animals (Fig. 1). High NAb titres were detected as early as 2 weeks post-immunization upon vaccination with BCG:CoVac, and titres were significantly increased compared to vaccination with Alum^{SpK} (approximate 10-fold increase). The mean NAb titres in the plasma of BCG:CoVac-vaccinated mice were approximately 10-fold greater than those seen in SARS-CoV-2 infected humans (Fig. 3a). Although the levels of NAbs peaked at 2 weeks post-vaccination with BCG:CoVac, they remained significantly elevated up to day 42 post-immunization unlike those in the other immunized groups.

Since previous work suggests that the level of IgG antibody correlates with NAb titres after SARS-CoV-2 infection[18], we examined whether a similar phenomenon was observed after vaccination with BCG:CoVac. Strong correlation (r> 0.9) was observed between IgG2c isotype and NAbs in groups vaccinated with BCG:CoVac or Alum^{SpK} (Fig. 3b), with a significant yet less robust correlation between IgG1 and NAbs for these groups (Fig. 3c). There was no correlation between NAbs and either IgG1 or IgG2c Ab for mice vaccinated with BCG^{SpK} alone (Fig. 3d, 3e).

These data suggest that alum is required for the optimal generation of NAbs after BCG:CoVac vaccination. This is a significant advantage for implementation of this vaccine candidate, due to the low cost and long standing safety record of alum[19,20]. Importantly, the

147   potential risk of vaccine-associated enhanced respiratory disease (VAERD) caused by the

148   selective induction of Th2 T cell responses by alum is offset by the strong Th1 immunity

149   induced by BCG:CoVac, which in turn is driven by BCG (Fig. 1e).

151   BCG:CoVac affords sterilizing immunity against SARS-CoV-2 infection in

152   K18-hACE2 mice

Wild-type mice are not permissive to SARS-CoV-2 infection, owing to incompatibility in the

receptor binding domain of the viral spike protein with the murine angiotensin-converting

enzyme 2 (ACE2) [21]. Transgenic mice expressing the human (h)ACE2 such as the K18-hACE2

mouse, are highly susceptible to SARS-CoV-2 infection, succumbing to lethal infection within

7 days post-infection[22]. We therefore assessed the protective role of BCG or BCG:CoVac

vaccination in SARS-CoV-2 infection in K18-hACE2 mice. Mice were vaccinated 21 days

prior to inoculation with 10^3 PFU SARS-CoV-2 (Fig. 4a). Mice sham vaccinated with PBS

succumbed to infection within 6 days with substantial deterioration in their condition with high

clinical scores (Fig. 4b) and 20% weight loss (Fig. 4c). This outcome was associated with high

viral titres in the airways (bronchoalveolar lavage fluid, BALF) (Fig. 4d) and lung tissues (Fig.

4e). These events led to extensive lung inflammation with substantial increases in inflammatory

cells in the airways (Fig. 4f) and lung tissue (Fig. 4g), and the levels of the pro-inflammatory

cytokine, IL-6, and chemokines KC (murine equivalent of IL-8) and MCP-1, in the lung tissues

(Fig. 4h) and airways (Fig. S2). MCP-1 was also increased in serum (Fig. S2). These are the

archetypal cytokines associated with severe human COVID-19[23]. Vaccination with BCG

showed some beneficial effects and partially protected against weight loss (~10%) and lung

IL-6 and KC responses but not in other disease features. Remarkably, vaccination with

BCG:CoVac 21 days prior to infection completely protected against infection, with no

observable weight loss or any clinical scores throughout the duration of the experiment (Fig.

4b, 4c). These mice had no detectable virus in the airways or lungs (Fig. 4D, 4E). They had

few signs of lung inflammation with moderate levels on inflammatory cells in the airways and

virtually none in the lung tissue (Fig. 4g), and only baseline levels of all pro-inflammatory

cytokines in the airways, lung and serum (Fig. 4h, S2). Importantly, combination of the spike

protein and alum with BCG did not alter the protective efficacy of the BCG vaccine against

aerosol *M. tuberculosis* in mice (Fig 4i).

Collectively, these findings demonstrate that single dose administration of BCG:CoVac is

sufficient to completely protect mice from the development of COVID-19 disease

manifestations, and to neutralize infectious SARS-CoV-2 and prevent pathogenic

inflammation in the lung.

183 Enhancing BCG:CoVac immunity against SARS-CoV-2 by heterologous
184 vaccine boosting

COVID-19 subunit vaccines typically display poor immunity after a single dose and require a

booster to induce sufficient generation of NAbs[24]. Whilst we observed high-titre NAbs as early

as two weeks post-BCG:CoVac vaccination (Fig. 3), we sought to determine if responses could

be further augmented by boosting with a prototype subunit vaccine (Alum^SpK) (Fig. 5a). At 7

days post-boost (day 28), IgG2c titres in plasma from mice primed either with BCG^SpK or

BCG:CoVac were increased and remained elevated up to day 42 (Fig. 5b). Corresponding

augmentation of NAbs was also seen in these boosted groups, with significantly elevated

responses in BCG:CoVac primed mice boosted with Alum^SpK (Fig. 5c). Boosting Alum^SpK

vaccination with a second dose led to a greater than 10-fold increase in NAbs in boosted mice;

however, responses were significantly higher in those with the BCG:CoVac-prime, Alum^SpK-

boost combination (Fig. 5c). Strikingly, plasma from BCG:CoVac-vaccinated mice was able

to neutralize both the B.1.1.7 variant (1.3-fold decrease compared to wild-type virus) and

B.1.351 variant (2.7-fold decrease) (Fig. 5d). Neutralization capacity against B.1.1.7 and

B.1.351 was maintained to some extent after prime-boost with Alum^{SpK} only, however titres

were approximately 10-fold less than those following the BCG:CoVac prime, Alum^{SpK}

combination (Fig. 5e).

Taken together, these data indicate that the antigen-specific immunity imparted by

BCG:CoVac can be further enhanced by heterologous boosting with a second SARS-CoV-2

vaccine, with this vaccination regime able to induce antibodies that can neutralize key VOCs.

DISCUSSION

Global vaccine access and distribution to low- and middle-income countries is critical for the

control of the COVID-19 pandemic. Vaccines must offer effective protective immunity yet

should be cheap to manufacture and have feasible cold chain management requirements. This

study describes a COVID-19 vaccine formulation, BCG:CoVac, that when delivered as a single

dose induces potent SARS-CoV-2 specific immunity in mice, particularly the generation of

high-titre, anti-viral neutralizing antibodies. Encouragingly, the level of immune responses

observed (particularly the generation of neutralizing antibodies) is equivalent to or exceeds

immunity elicited by approved COVID-19 vaccines, when these candidate vaccines were tested

in the murine model[24-26]. BCG:CoVac may have the additional advantage of inducing

protection against other respiratory infections for which BCG is known to induce some level

of protective immunity, including future pandemic viruses[11]. In addition, the possibility that

prior BCG exposure may impart protection against severe COVID-19[27], which is currently

under evaluation through numerous randomised control studies[9], raises the possibility that a

BCG-based vaccine could afford protection against SARS-CoV-2 escape mutants or new

pandemic coronavirus that may emerge. Indeed our data demonstrate that BCG:CoVac can

neutralize two of the key VOCs that are circulating globally, namely B.1.1.7 and B.1351.

BCG:CoVac could also provide additional benefit in countries where BCG is part of childhood

immunization programs for the control of TB, based on recent findings that repeat BCG

vaccination significantly reduced rates of *M. tuberculosis* infection[28].

An advantage of our vaccine approach is the use of alum to potentiate immune responses,

particularly the generation of NAbs after vaccination. Alum is a low cost, globally accessible

vaccine adjuvant with an excellent safety record in humans[19]. The relative paucity of IFN-γ-

secreting T cells observed after Alum[SpK] vaccination corresponds with that previously seen

with alum-precipitated vaccines using the spike protein[29] and is consistent with the known

preferential priming of Th2-type immunity by alum-based adjuvants[30]. The ability of

BCG:CoVac to induce strong Th1 immunity is due to the adjuvant effect of BCG components

to induce such responses[31]. This has clear importance, as T cell responses in recovering

COVID-19 patients are predominately Th1-driven[32], expression of IFN-γ was lower in severe

COVID-19 cases compared to mild ones[33] and the development of Th2 immunity correlated

with VAERD[34].We also observed only background levels of the inflammatory cytokines IL-17

and TNF after BCG:CoVac delivery, suggesting reduced levels of potentially deleterious,

circulating inflammatory cytokines. Heightened expression of IL-17 correlates with severe

COVID-19 disease[35], while neutralizing IL-17 has been suggested as a possible therapy to treat

acute respiratory distress syndrome in SARS-CoV-2-infected individuals[36]. In addition, the

development of VAERD is also associated with Th17 immunity[37].

Our study contributes to defining correlates of immunity in animal models that could be

applied to fast-track the development of next-generation COVID-19 vaccines. A single dose of

BCG:CoVac was sufficient to clear infectious SARS-CoV-2 from the lungs of K18-hACE2

mice, with no signs of clinical disease during the infection time-course, which is otherwise

lethal (Fig. 4). Our findings are in agreement with previous reports that demonstrate that these

mice succumb rapidly to infection, indicating that the level of NAbs elicited by BCG:CoVac

is sufficient to clear SARS-CoV-2 infection in this model[21,38]. While NAb levels are known to

correlate with efficacy of COVID-9 vaccines in humans[17], other immune parameters may play

an important role. Accordingly, we observed that high NAb levels in blood corelated with

strong induction of GC and plasma B cells in lymph nodes, as well as heightened levels of Tfh

cells. In COVID-19 convalescent individuals, the presence of memory B cells and Tfh are

closely associated with NAb activity[38,39]. Further, although BCG vaccination alone did not

reduce lung viral load, mice immunized with this vaccine did show an intermediate level of

both weight loss and cytokine/chemokine induction during SARS-CoV-2 infection, suggesting

a possible 'dampening' of host inflammatory responses (Fig. 4 and Fig. S2). Such an effect

was seen in a human challenge model using the live-attenuated yellow fever vaccine, where

BCG reduced circulating pro-inflammatory cytokines[40]. It will be of particular interest to see

the outcome of ongoing clinical trials to determine if BCG vaccination can reduce COVID-19

incidence and severity[9].

In conclusion, we describe a COVID-19 vaccine strategy based on the existing BCG

vaccine, that would be broadly applicable for all populations susceptible to SARS-CoV-2

infection. Of particular note, this strategy could be readily incorporated into current vaccine

schedules in countries where BCG is currently used. Further assessment in humans will

determine if BCG:CoVac can impart protective immunity against not only SARS-CoV-2, but

also other respiratory infections for which BCG has known efficacy.

METHODS

Bacterial culture

M. bovis BCG (strain Pasteur) was grown at 37°C in Middlebrook 7H9 media (Becton

Dickinson, BD, New Jersey, USA) supplemented with 0.5% glycerol, 0.02% Tyloxapol, and

10% albumin-dextrose-catalase (ADC) or on solid Middlebrook 7H11 media (BD)

supplemented with oleic acid–ADC. To prepare single cell suspensions, cultures in exponential

phase (OD_{600}=0.6) were washed in PBS, passaged 10 times through a 27G syringe, briefly

sonicated and centrifuged at low speed for 10 min to remove residual bacterial clumps. BCG

suspensions were frozen at -80° C in PBS supplemented with 20% glycerol, and colony

forming units (CFU) for vaccination enumerated on supplemented Middlebrook 7H11 agar

plates.

Ethics statement

All mouse experiments were performed according to ethical guidelines as set out by the Sydney

Local Health District (SLHD) Animal Ethics and Welfare Committee, which adhere to the

Australian Code for the Care and Use of Animals for Scientific Purposes (2013) as set out by

the National Health and Medical Research Council of Australia. SARS-CoV-2 mouse infection

experiments were approved by the SLHD Institutional Biosafety Committee. COVID-19

patients were recruited through Royal Prince Alfred Hospital (RPA) Virtual, a virtual care

system enabling remote monitoring of patients. The study protocol was approved by the RPA

ethics committee (Human ethics number X20-0117 and 2020/ETH00770) and by the

participants' written consent. All associated procedures were performed in accordance with

approved guidelines.

Immunization

Female C57BL/6 (6-8 weeks of age) purchased from Australian BioResources (Moss Vale,

Australia) or hemizygous male K18-hACE2 mice bred in-house[41] were housed at the Centenary

Institute in specific pathogen-free conditions. SARS-CoV-2 full-length spike stabilized,

trimeric protein (SpK) was expressed in EXPI293F™ cells and purified as described

previously[42]. Mice (n=3-4) were vaccinated subcutaneously in the footpad (s.c) with 5×10^5

CFU of BCG alone, 5 μg of SpK combined with either BCG (BCG[SpK]) or 100 μg of Alhydrogel

(Alum) (Invivogen, California, USA, Alum[SpK]), or a combination of BCG ($5x10^5$ CFU), SpK

(5 μg) and Alyhydrogel (100 μg) (BCG:CoVac). Some mice were boosted three weeks after

the first vaccination with 5 μg of SpK combined with 100 μg of Alhyhdrogel. Mice were bled

fortnightly after the first immunization (collected in 10 μl of Heparin 50000 U/ml). Plasma was

collected after centrifugation at 300 x *g* for 10 min and remaining blood was resuspended in 1

mL of PBS Heparin 20 U/mL, stratified on top of Histopaque 10831 (Sigma-Aldrich, Missouri,

USA) and the PBMC layer collected after gradient centrifugation.

Flow cytometry assays

Popliteal lymph nodes were collected at day 7 post immunization, and single cell suspensions

were prepared by passing them through a 70 μm sieve. To assess specific B cell responses,

$2x10^6$ cells were surface stained with Fixable Blue Dead Cell Stain (Life Technologies) and

Spike-AF647 (1 μg), rat anti-mouse GL7-AF488 (clone GL7, 1:200, Biolegend cat#144612),

rat anti-mouse MHC-II-AF700 (clone M5/114.15.2, 1:150, cat#107622), rat anti-mouse IgD-

PerCP5.5 (clone 11-26c.2a, 1:200, BD, cat#564273), rat anti-mouse IgM-BV421 (clone

RMM-1, 1:200, Biolegend, cat#406518), rat anti-mouse CD138-BV605 (clone 281-2, 1:200,

Biolegend, cat#142516), rat anti-mouse CD19-BV785 (clone 1D3, 1:200, BD, cat#563333),

rat anti-mouse CD38-APCy7 (clone 90, 1:200, Biolegend, cat#102728). To assess T cell

responses, $2x10^6$ lymph node cells were stained with the following monoclonal antibodies: rat

anti-mouse CXCR5-biotin (clone 2G8, 1:100, BD, cat#551960), streptavidin PECy7, rat anti-

mouse CD4-AF700 (clone RM4-5, 1:200, BD cat#557956), rat anti-mouse CD8-APCy7 (clone

53-6.7, 1:200, BD cat#557654), rat anti-mouse CD44-BV605 (clone IM7, 1:300, BD

cat#563058). Cells were then fixed and permeabilized using the eBioscience

fixation/permeabilization kit (ThermoFischer) according to the manufacturer's protocol and

intracellular staining was performed using anti-BCL-6-AF647 (clone K112-91, 1:100, BD,

cat#561525).

To assess SpK-specific cytokine induction by T cells, murine PBMCs were stimulated for

4 hrs with SpK (5 μg/mL) and then supplemented with Protein Transport Inhibitor cocktail

(Life Technologies, California, USA) for a further 10-12 hrs. Cells were surface stained with

Fixable Blue Dead Cell Stain (Life Technologies) and the marker-specific fluorochrome-

labeled antibodies rat anti-mouse CD4-AF700 (clone RM4-5, 1:200, BD cat#557956), rat anti-

mouse CD8-APCy7 (clone 53-6.7, 1:200, BD cat#557654), rat anti-mouse CD44-FITC (clone

IM7, 1:300, BD cat#561859). Cells were then fixed and permeabilized using the BD

Cytofix/Cytoperm^{TM} kit according to the manufacturer's protocol. Intracellular staining was

performed using rat anti-mouse IFN-γ-PECy7 (clone XMG1-2, 1:300, BD cat#557649), rat

anti-mouse IL-2-PE (clone JES6-5H4, 1:200, BD cat#554428), rat anti-mouse IL-17-PB (clone

TC11-18H10.1, 1:200, cat#506918, BioLegend California, USA), rat anti-mouse TNF-

PErCPCy5.5 (clone MP6-XT22, 1:200, BD cat#560659). All samples were acquired on a BD

LSR-Fortessa (BD) or a BD-LSRII and assessed using FlowJo^{TM} analysis software v10.6

(Treestar, USA).

Antibody ELISA

Microtitration plates (Corning, New York, USA) were incubated overnight with 1 μg/mL SpK

at room temperature (RT), blocked with 3% BSA and serially diluted plasma samples were

added for 1 hour at 37˚C. Plates were washed and biotinylated polyclonal goat anti-mouse IgG1

(1:50,000, abcam Cambridge, UK, cat#ab97238), polyclonal goat anti-mouse IgG2c (1:10,000,

Abcam, cat# ab97253), or polyclonal goat anti-mouse IgG (1:350,000, clone abcam

cat#ab6788) added for 1 hour at RT. After incubation with streptavidin-HRP (1:30,000, abcam,

cat#405210) for 30 min at RT, binding was visualized by addition of tetramethyl benzene

(Sigma-Aldrich). The reaction was stopped with the addition of 2N H_2SO_4 and absorbances

were measured at 450 nm using a M1000 pro plate reader (Tecan, Männedorf, Switzerland).

End point titres were calculated as the dilution of the sample that reached the average of the

control serum ± 3 standard deviations.

High content live SARS-CoV-2 neutralization assay

High-content fluorescence microscopy was used to assess the ability of sera/plasma to inhibit

SARS-CoV-2 infection and the resulting cytopathic effect in live permissive cells (VeroE6).

Sera were serially diluted and mixed in duplicate with an equal volume of $1.5x10^3$ $TCID_{50}$/mL

virus solution (B.1.319) or $1.25x10^4$ $TCID_{50}$/mL virus solution (A2.2, B.1.1.7, B.1.351). After

1 hour of virus-serum coincubation at 37°C, 40 µL were added to equal volume of freshly-

trypsinised VeroE6 cells in 384-well plates ($5x10^3$/well). After 72 hrs, cells were stained with

NucBlue (Invitrogen, USA) and the entire well surface was imaged with InCell Analyzer 2500

(Cytiva). Nuclei counts were obtained for each well with InCarta software (Cytiva), as proxy

for cell death and cytopathic effect resulting from viral infection. Counts were compared

between convalescent sera, mock controls (defined as 100% neutralization), and infected

controls (defined as 0% neutralization) using the formula; % viral neutralization = (D-(1-

Q))x100/D, where Q = nuclei count of sample normalized to mock controls, and D = 1-Q for

average of infection controls. The cut-off for determining the neutralization endpoint titre of

diluted serum samples was set to ≥50% neutralization.

SARS-CoV-2 challenge experiments

Male hemizygous K18-hACE2 mice were transported to the PC3 facility in the Centenary

Institute for SARS-CoV-2 infection. Mice were anaesthetised with isoflurane followed by

intranasal challenge with 10^3 PFU SARS-CoV-2 (VIC01/2020) in a 30 µL volume. Following

infection, mice were housed in the IsoCage N biocontainment system (Tecniplast, Italy) and

were given access to standard rodent chow and water *ad libitum*. Mice were weighed and

monitored daily, with increased frequency of monitoring when mice developed symptoms. At

day 6 post-infection, mice were euthanised with intraperitoneal overdose of pentobarbitone

(Virbac, Australia). Blood was collected *via* heart bleed, allowed to coagulate at RT and

centrifuged (10,000 *g*, 10 min) to collect serum. Multi-lobe lungs were tied off and BALF was

collected from the single lobe *via* lung lavage with 1 mL HANKS solution using a blunted 19-

gauge needle inserted into the trachea. BALF was centrifuged (300 *g*, 4°C, 7 min), and

supernatants collected and snap frozen. Cell pellets were treated with 200 µL Red Blood Cell

Lysis Buffer (ThermoFisher, USA) for 5 min, followed by addition of 700 µL HANKS solution

to inactivate the reaction and then centrifuged again. Cell pellets were resuspended in 160 µL

HANKS solution and enumerated using a haemocytometer (Sigma-Aldrich, USA). Multi-lobe

lungs were collected and cut into equal thirds, before snap freezing on dry ice. Lung

homogenates were prepared fresh, with multi-lobe lungs placed into a gentleMACS C-tube

(Miltenyi Biotec, Australia) containing 2 mL HANKS solution. Tissue was homogenised using

a gentleMACS tissue homogeniser, after which homogenates were centrifuged (300 *g*, 7 min)

to pellet cells, followed by collection of supernatants for plaque assays and cytokine/chemokine

measurements. The single lobe lung was perfused with 0.9% NaCl solution *via* the heart,

followed by inflation with 0.5 mL 10% neutral buffered formalin through the trachea, and

placed into a tube containing 10% neutral buffered formalin. Following fixation for at least 2

weeks, single lobes were transported to a PC2 facility where they were paraffin-embedded,

sections cut to 3 µm thickness using a Leica microtome (Leica, Germany) and then stained

using Quick Dip Stain Kit (Modified Giemsa Stain) protocol as per manufacturer's instructions

(POCD Scientific, Australia). Inflammatory cells in single lobe lungs were counted using a

Zeiss Axio Imager.Z2 microscope with a 40X objective (Zeiss, Germany).

Plaque assays

VeroE6 cells (CellBank Australia, Australia) were grown in Dulbecco's Modified Eagles Medium (Gibco, USA) supplemented with 10% heat-inactivated foetal bovine serum (Sigma-Aldrich, USA) at 37°C/5% CO_2. For plaque assays, cells were placed into a 24-well plate at 1.5×10^5 cells/well and allowed to adhere overnight. The following day, virus-containing samples were serially diluted in Modified Eagles Medium (MEM), cell culture supernatants removed from the VeroE6 cells and 250 µL of virus-containing samples was added to cell monolayers. Plates were incubated and gently rocked every 15 min to facilitate viral adhesion. After 1 hr, 250 µL of 0.6% agar/MEM solution was gently overlaid onto samples and placed back into the incubator. At 72 hrs post-infection, each well was fixed with an equal volume of 8% paraformaldehyde solution (4% final solution) for 30 min at RT, followed by several washes with PBS and incubation with 0.025% crystal violet solution for 5 min at RT to reveal viral plaques.

Cytometric bead arrays (CBAs)

CBAs were performed as per the manufacturer's instructions (Becton Dickinson, USA). Briefly, a standard curve for each analyte was generated using a known standard supplied with each CBA Flex kit. For each sample, 10 µL was added to a well in a 96-well plate, followed by incubation with 1 µL of capture bead for each analyte (1 hr, RT, in the dark). Following capture, 1 µL of detection bead for each analyte was added to each well, followed by incubation (2 hrs, RT, in the dark). Samples were then fixed overnight in an equal volume of 8% paraformaldehyde solution (4% final solution). The following day, samples were transferred to a new 96-well plate and then transported to the PC2 facility for a second round of fixation. Samples were examined using a BD LSR Fortessa equipped with a High-Throughput Sampler (HTS) plate reader.

Mycobacterium tuberculosis aerosol challenge

Eight weeks after the last vaccination mice were infected with *M. tuberculosis* H37Rv via the aerosol route using a Middlebrook airborne infection apparatus (Glas-Col, IN, USA) with an infective dose of ~100 viable bacilli. Four weeks later, the lungs and spleen were harvested, homogenized, and plated after serial dilution on supplemented Middlebrook 7H11 agar plates. Colonies forming units (CFU) were determined 3 weeks later and expressed as $\log_{10}$ CFU.

Statistical analysis

The significance of differences between experimental groups was evaluated by one-way analysis of variance (ANOVA), with pairwise comparison of multi-grouped data sets achieved using Tukey's or Dunnett's *post-hoc* test. Differences were considered statistically significant when $p \leq 0.05$.

ACKNOWLEDGMENTS

This work was supported by MRFF COVID-19 Vaccine Candidate Research Grant 2007221 (JAT, CC, PMH, SGT, MCG, ALF, WJB). We thank Florian Krammer of the Icahn School of Medicine, Mt Sinai for provision of the pCAGGS vector containing the SARS-CoV-2 Wuhan-Hu-1 Spike Glycoprotein Gene. We acknowledge the support of the University of Sydney Advanced Cytometry Facility and the animal facility at the Centenary Institute. We thank Sunil David (ViroVax LLC) and Wolfgang Leitner (NIAID, NIH) for helpful discussions. PMH is funded by a Fellowship and grants from the National Health and Medical Research Council (NHMRC) of Australia (1175134) and by UTS. JAT, BMS and WJB are supported by the NHMRC Centre of Research Excellence in Tuberculosis Control (1153493). The establishment of the PC3 COVID-19 facility was supported by UTS and the Rainbow Foundation.

AUTHOR CONTRIBUTIONS

448 CC, MDJ, AOS, SGT, PHM and JAT designed the study. CC, MDJ, AOS, DHN, ALF, AA,

449 RS, NDB, AG, KP, BMS, JKKL, SGT performed the experiments. All authors contributed to

450 data analysis/interpretation. CC, MDJ, PHM and JAT wrote the first manuscript draft and all

451 authors provided revision to the scientific content of the final manuscript.

REFERENCES

Dagan, N. *et al.* BNT162b2 mRNA Covid-19 Vaccine in a Nationwide Mass Vaccination Setting. *N Engl J Med* **384**, 1412-1423 (2021).

Pritchard, E. *et al.* Impact of vaccination on SARS-CoV-2 cases in the community: a population-based study using the UK's COVID-19 Infection Survey. *medRxiv* (2021).

Davies, N. G. *et al.* Increased mortality in community-tested cases of SARS-CoV-2 lineage B.1.1.7. *Nature* (2021).

Tegally, H. *et al.* Detection of a SARS-CoV-2 variant of concern in South Africa. *Nature* **592**, 438-443 (2021).

Wang, P. *et al.* Antibody resistance of SARS-CoV-2 variants B.1.351 and B.1.1.7. *Nature* **593**, 130-135 (2021).

Wibmer, C. K. *et al.* SARS-CoV-2 501Y.V2 escapes neutralization by South African COVID-19 donor plasma. *Nat Med* **27**, 622-625 (2021).

Madhi, S. A. *et al.* Efficacy of the ChAdOx1 nCoV-19 Covid-19 Vaccine against the B.1.351 Variant. *N Engl J Med* **384**, 1885-1898 (2021).

Iacobucci, G. & Mahase, E. Covid-19 vaccination: What's the evidence for extending the dosing interval? *BMJ* **372** (2021).

Netea, M. G. *et al.* Trained Immunity: a Tool for Reducing Susceptibility to and the Severity of SARS-CoV-2 Infection. *Cell* **181**, 969-977 (2020).

Giamarellos-Bourboulis, E. J. *et al.* Activate: Randomized Clinical Trial of BCG Vaccination against Infection in the Elderly. *Cell* **183**, 315-323 e319 (2020).

Covian, C. *et al.* BCG-Induced Cross-Protection and Development of Trained Immunity: Implication for Vaccine Design. *Front Immunol* **10**, 2806 (2019).

Amanat, F. *et al.* A serological assay to detect SARS-CoV-2 seroconversion in humans. Nature Medicine **26**, 1033-1036 (2020).

13 Nazeri, S., Zakeri, S., Mehrizi, A. A., Sardari, S. & Djadid, N. D. Measuring of IgG2c

isotype instead of IgG2a in immunized C57BL/6 mice with *Plasmodium vivax* TRAP as a

subunit vaccine candidate in order to correct interpretation of Th1 versus Th2 immune

response. *Exp Parasitol* **216**, 107944 (2020).

14 Sauer, K. & Harris, T. An Effective COVID-19 Vaccine Needs to Engage T Cells. *Front*

Immunol **11**, 581807 (2020).

15 Counoupas, C. & Triccas, J. A. The generation of T-cell memory to protect against

tuberculosis. *Immunol Cell Biol* **97** (2019).

16 Lederer, K. *et al.* SARS-CoV-2 mRNA Vaccines Foster Potent Antigen-Specific Germinal

Center Responses Associated with Neutralizing Antibody Generation. *Immunity* **53**, 1281-

1295 e1285 (2020).

17 Khoury, D. S. *et al.* Neutralizing antibody levels are highly predictive of immune

protection from symptomatic SARS-CoV-2 infection. *Nature Medicine* (2021).

18 Suthar, M. S. *et al.* Rapid Generation of Neutralizing Antibody Responses in COVID-19

Patients. *Cell Rep Med* **1**, 100040 (2020).

19 Hotez, P. J., Corry, D. B., Strych, U. & Bottazzi, M. E. COVID-19 vaccines: neutralizing

antibodies and the alum advantage. *Nat Rev Immunol* **20**, 399-400 (2020).

20 Lancet Covid-19 Commissioners, T. F. C. & Commission, S. Lancet COVID-19

Commission Statement on the occasion of the 75th session of the UN General Assembly.

Lancet **396**, 1102-1124 (2020).

21 Johansen, M. D. *et al.* Animal and translational models of SARS-CoV-2 infection and

COVID-19. *Mucosal Immunol* **13**, 877-891 (2020).

22 Winkler, E. S. *et al.* SARS-CoV-2 infection of human ACE2-transgenic mice causes

severe lung inflammation and impaired function. *Nat Immunol* **21**, 1327-1335 (2020).

23 Yang, Y. *et al.* Plasma IP-10 and MCP-3 levels are highly associated with disease severity

and predict the progression of COVID-19. *J Allergy Clin Immunol* **146**, 119-127 e114

(2020).

24 J.-H. Tian *et al.* SARS-CoV-2 spike glycoprotein vaccine candidate NVX-CoV2373

immunogenicity in baboons and protection in mice *Nat Commun* **12,** 372 (2021).

25 Corbett, K. S. *et al.* SARS-CoV-2 mRNA vaccine design enabled by prototype pathogen

preparedness. *Nature* **586**, 567-571 (2020).

26 Graham, S. P. *et al.* Evaluation of the immunogenicity of prime-boost vaccination with the

replication-deficient viral vectored COVID-19 vaccine candidate ChAdOx1 nCoV-19.

NPJ Vaccines **5**, 69 (2020).

27 Escobar, L. E., Molina-Cruz, A. & Barillas-Mury, C. BCG vaccine protection from severe

coronavirus disease 2019 (COVID-19). *Proc Natl Acad Sci U S A* **117**, 17720-17726

(2020).

28 Nemes, E. *et al.* Prevention of *M. tuberculosis* Infection with H4:IC31 Vaccine or BCG

Revaccination. *N Engl J Med* **379**, 138-149 (2018).

29 Kuo, T. Y. *et al.* Development of CpG-adjuvanted stable prefusion SARS-CoV-2 spike

antigen as a subunit vaccine against COVID-19. *Sci Rep* **10**, 20085 (2020).

30 HogenEsch, H., O'Hagan, D. T. & Fox, C. B. Optimizing the utilization of aluminum

adjuvants in vaccines: you might just get what you want. *NPJ Vaccines* **3**, 51 (2018).

31 Uthayakumar, D. *et al.* Non-specific Effects of Vaccines Illustrated Through the BCG

Example: From Observations to Demonstrations. *Front Immunol* **9**, 2869 (2018).

32 Grifoni, A. *et al.* Targets of T Cell Responses to SARS-CoV-2 Coronavirus in Humans

with COVID-19 Disease and Unexposed Individuals. *Cell* **181**, 1489-1501 e1415 (2020).

33 Chen, G. *et al.* Clinical and immunological features of severe and moderate coronavirus

disease 2019. *J Clin Invest* **130**, 2620-2629 (2020).

34 Jeyanathan, M. *et al.* Immunological considerations for COVID-19 vaccine strategies. *Nat*

Rev Immunol **20**, 615-632 (2020).

35 Xu, Z. *et al.* Pathological findings of COVID-19 associated with acute respiratory distress

syndrome. *Lancet Respir Med* **8**, 420-422 (2020).

36 Pacha, O., Sallman, M. A. & Evans, S. E. COVID-19: a case for inhibiting IL-17? *Nat Rev*

Immunol **20**, 345-346 (2020).

37 Hotez, P. J., Corry, D. B. & Bottazzi, M. E. COVID-19 vaccine design: the Janus face of

immune enhancement. *Nat Rev Immunol* **20**, 347-348 (2020).

38 Zheng, J. *et al.* COVID-19 treatments and pathogenesis including anosmia in K18-hACE2

mice. *Nature* **589**, 603-607 (2021).

39 Wheatley, A. K. *et al.* Evolution of immune responses to SARS-CoV-2 in mild-moderate

COVID-19. *Nat Commun* **12**, 1162 (2021).

40 Arts, R. J. W. *et al.* BCG Vaccination Protects against Experimental Viral Infection in

Humans through the Induction of Cytokines Associated with Trained Immunity. *Cell Host*

Microbe **23**, 89-100 e105 (2018).

41 McCray, P. B., Jr. *et al.* Lethal infection of K18-hACE2 mice infected with severe acute

respiratory syndrome coronavirus. *J Virol* **81**, 813-821 (2007).

42 Xi, C. R. *et al.* A Novel Purification Procedure for Active Recombinant Human DPP4 and

the Inability of DPP4 to Bind SARS-CoV-2. *Molecules* **25** (2020).

     FIGURE LEGENDS

Fig. 1 | Single immunization with BCG:CoVac vaccine induces rapid development of anti-

SARS-CoV-2 spike antibodies and IFN-γ-secreting T cells. a, C57BL/6 mice were

vaccinated subcutaneously with PBS, BCG, BCG^{SpK}, Alum^{SpK} or BCG:CoVac and whole

blood collected at day 14, 28 and 42. **b,c**, Spike-specific IgG1 and IgG2c titres in plasma were

determined by ELISA and estimated by the sigmoidal curve of each sample interpolated with

the threshold of the negative sample ± 3 standard deviations. The dotted line shows the limit

of detection. **d**, At 14 days post-vaccination PBMCs were restimulated *ex vivo* with 5 μg/mL

of SARS-CoV-2 spike and cytokine production determined by flow cytometry. Representative

dot plots of CD44$^+$ CD4$^+$ T cells and CD44$^+$ CD8$^+$ T cells expressing IFN-γ. **e**, Numbers of

circulating CD4$^+$ and CD8$^+$ T cells expressing IFN-γ or CD4$^+$ T cells expressing IL-17 or TNF.

Data presented as mean ± s.d. Significant differences between groups compared to BCG^{Spk}

*p<0.05, **p<0.01, ***p<0.01 or Alum^{Spk} †p<0.05, ††p<0.01, †††p<0.001 were determined

using one-way ANOVA.

Fig. 2 | BCG:CoVac vaccination promotes expansion of T follicular helper cells and spike-

specific B cells in mice. a-c, C57BL/6 mice were vaccinated subcutaneously with PBS, BCG,

BCG^{SpK}, Alum^{SpK} or BCG:CoVac and 7 days after immunization B and T cell response

assessed by multicolour flow cytometry in the draining lymph node. Shown are representative

dot plots of spike-specific germinal centre B cells (**a**, CD19$^+$MHCII$^+$GL7$^+$CD38$^-$), plasma B

cells (**b**, CD19$^+$MHCII$^+$CD138$^+$) and T follicular helper T cells (**c**, CXCR5$^+$BCL6$^+$). **d-f**, The

total number of **d**, spike$^+$ GC B cells, **e**, spike$^+$ plasma cells and **f**, T follicular helper cells. Data

presented as mean ± s.d. Significant differences between groups *p<0.05, **p<0.01,

***p<0.01 were determined by one-way ANOVA.

Fig. 3 | BCG:CoVac induces high titre neutralizing antibodies against live SARS-CoV-2 that correlate with the production of antigen-specific IgG2c. a-e, Plasma from vaccinated mice (from Fig. 1) were tested for neutralizing activity against live SARS-CoV-2 infection of VeroE6 cells. **a**, Neutralizing antibody (NAb) titres (IC_{50}) were calculated as the highest dilution of plasma that still retained at least 50% inhibition of infection compared to controls. NAb titres from PCR confirmed SARS-CoV-2-infected individuals (COVID) were determined using the same method. **b,c**, Spearman correlations of spike-specific IgG2c or IgG1 titres and NAbs after $Alum^{SpK}$ or BCG:CoVac vaccination. **d,e**, Correlation of IgG2c or IgG1 titres and NAbs after vaccination with BCG^{SpK}. The dotted line shows the limit of detection. Data presented as mean ± s.d. Significant differences between groups compared to BCG^{Spk} **p<0.01, ***p<0.0 or $Alum^{Spk}$ †p<0.05, †††p<0.001 was determined by one-way ANOVA.

Fig. 4 | A single dose of BCG:CoVac protects against severe SARS-CoV-2 infection. a, Mice were immunized with sham (PBS), BCG or BCG:CoVac 21 days prior to challenge with 10^3 PFU SARS-CoV-2. Disease outcomes were assessed 6 days later. **b**, Clinical scores at day 6 post-infection. **c**, Percentage of initial body weight loss in hemizygous male K18-hACE2 mice (n=4/group). Viral titres in lung homogenates (**d**) or bronchoalveolar lavage fluid (BALF) (**e**) were determined using plaque assay. The dotted line represents the limit of detection. **f**, Total inflammatory cells in bronchoalveolar lavage fluid (BALF). **g**, Total number of inflammatory cells in stained histological sections of lungs. **h**, Cytokine/chemokine quantification in lung homogenates. **i**, Six weeks after immunization mice were challenged with *M. tuberculosis* H37Rv by aerosol (~100 CFU) and four weeks later the bacterial load was assessed in the lungs and presented as log_{10} of the mean CFU ± SEM. Significant differences between groups *p<0.05, **p<0.01 were determined by one-way ANOVA.

Fig. 5 | Heterologous boosting of BCG:CoVac-primed mice results in augmented SARS-

CoV-2-specific IgG2c titres and neutralizing antibodies. a, C57BL/6 mice were vaccinated

(as in Fig. 1) and at day 21 mice were boosted with Alum^{Spk}. **b**, Spike-specific IgG2c titres in

plasma were determined by ELISA estimated from the sigmoidal curve of each sample

interpolated with the threshold of the negative sample $\pm$ 3 standard deviations. **c**, Neutralizing

antibody (NAb) titres (IC$_{50}$) were calculated as the highest dilution of plasma for all groups

that still retained at least 50% inhibition of infection compared to controls. The dotted line

shows the limit of detection. **d,e**, NAb titres against the B.1.1.7 or B.1.351 SARS-CoV-2

variants were also determined using plasma from either **d**, Alum^{SpK} or **e**, BCG:CoVac-

vaccinated mice. Data presented as mean $\pm$ s.d. Significant differences between groups

compared to BCG^{Spk} *$p<0.05$, **$p<0.01$, ***$p<0.01$ or Alum^{Spk} †$p<0.05$, ††$p<0.01$,

†††$p<0.001$ were determined by one-way ANOVA.

Figure 1

Figure 2

Figure 3

Figure 4

Figure 5

Figure S1

Fig. S1 | BCG promotes spike-specifica antibody reponses. Mice were vaccinated s.c with BCG (5x10[5] CFU) and 12 week later were vaccinated twice, 3 weeks apart with s.c with SARS-CoV-2 spike protein (5 μg) formulated in alum (100 μg; Alum^{SpK}). At the indicated timepoints the titre of spike-specific IgG2c in sera was determined by ELISA.

Figure S2

Fig. S2 | Single dose BCG:CoVac prevents the development of clinical disease in SARS-CoV-2 infected K18-hACE2 mice. a, Mice were immunised with sham (PBS), BCG or BCG:CoVac 21 days prior to challenge with 10^3 PFU SARS-CoV-2. Cytokine/chemokine levels were determined in BALF (**a**) and serum (**b**) by cytometric bead array.