The combined treatment of Molnupiravir and Favipiravir results in a marked potentiation of antiviral

efficacy in a SARS-CoV-2 hamster infection model

Rana Abdelnabi[1#*], Caroline S. Foo[1#], Suzanne J. F. Kaptein[1], Xin Zhang[1], Lana Langendries[1], Laura

Vangeel[1], Judith Breuer[2], Juanita Pang[2], Rachel Williams[2], Valentijn Vergote[1], Elisabeth Heylen[1], Pieter

Leyssen[1], Kai Dallmeier[1], Lotte Coelmont[1], Dirk Jochmans[1], Arnab K. Chatterjee[3], Steven De Jonghe[1],

Birgit Weynand[4], Johan Neyts[1,5*].

1. KU Leuven Department of Microbiology, Immunology and Transplantation, Rega Institute for

Medical Research, Laboratory of Virology and Chemotherapy, B-3000 Leuven, Belgium.

2. UCL Institute of Child Health, 30 Guilford Street, London, WC1N 1EH, United Kingdom.

3. Calibr at Scripps Research, La Jolla, CA, USA.

4. KU Leuven Department of Imaging and Pathology, Translational Cell and Tissue Research, B-3000

Leuven, Belgium; Division of Translational Cell and Tissue Research.

5. Global Virus Network, GVN.

#R.A. and C.S.F. contributed equally to this work

*To whom correspondence may be addressed. Email: rana.abdelnabi@kuleuven.be and

johan.neyts@kuleuven.be.

Abstract

Favipiravir and Molnupiravir, orally available antivirals, have been reported to exert antiviral activity

against SARS-CoV2. In recent days preliminary efficacy data have been reported in COVID-19 patients.

We here studied the combined antiviral effect of the drugs in the SARS-CoV2 hamster infection model.

We first demonstrate that Molnupiravir can reduce infectious virus titers in lungs of infected animals

in a dose-dependent manner by up to 3.5 $\log_{10}$ which is associated with a marked improvement of

virus-induced lung pathology. When animals are treated with a combination of suboptimal doses of

Molnupiravir and Favipiravir (that each alone result in respectively a 1.3 $\log_{10}$ and 1.1 $\log_{10}$ reduction

of infectious virus titers in the lungs), a marked combined potency is observed. Infectious virus titers

in the lungs of animals treated with the combo are on average reduced by 4.5 $\log_{10}$ and infectious virus

are no longer detected in the lungs of 60% of treated infected animals. Both drugs result in an

increased mutation frequency of the remaining viral RNA recovered from the lungs. In the combo-

treated hamsters an increased frequency of C-to-T and G-to-A mutations in the viral RNA is observed

as compared to the single treatment groups which may explain the pronounced antiviral potency of

the combination. Our findings may lay the basis for the design of clinical studies to test the efficacy of

the combination of Molnupiravir and Favipiravir in the treatment of COVID-19.

Keywords

SARS-CoV-2; Antivirals; Molnupiravir; Favipiravir, hamsters, coronavirus

Main text

The severe acute respiratory syndrome coronavirus 2 (SARS-CoV-2) was first identified in Wuhan, China in December 2019 [1]. Since then, the virus rapidly spread around the globe with more than 115 million cases and 2.5 million deaths reported until 3rd March 2021 [www.covid19.who.int]. Infection with SARS-CoV-2 results in coronavirus-induced disease (COVID-19) which is characterized by a wide range of symptoms including fever, dry cough, muscle and/or joint pain, headache, decreased sense of taste and smell and diarrhea. The disease can progress into severe complications such as acute respiratory distress syndrome (ARDS), respiratory failure, septic shock as well as multi-organ failure, which are mainly attributed to a massive cytokine storm and exaggerated immune response [2].

To date, there are no approved, selective coronavirus antivirals to treat or prevent infections. Even those vaccinated may not all be protected against infection and disease, in particular following infection with variants that are less susceptible to the current vaccines. Antivirals against SARS-CoV-2 will, at least when given early enough after a positive test or after onset of symptoms, reduce the chance to progress to (more) severe disease. In addition, such antiviral drugs will be useful to protect for example healthcare workers and high-risk patients in a prophylactic setting. Such drugs are also needed for the treatment of immunodeficient patients who do not mount a (sufficiently robust) immune response following vaccination. Since the *de novo* development and approval of (a) specific, highly potent antiviral(s) for SARS-CoV-2 may require years, the main focus for COVID-19 treatment in the current pandemic is to repurpose drugs that have been approved or in clinical trials for other diseases [3].

We recently demonstrated that the anti-influenza drug Favipiravir results in a pronounced antiviral activity in SARS-CoV-2-infected hamsters [4]. The efficacy of the drug is currently being explored in multiple phase II clinical studies. Interim data from a multicenter, open-labeled study of Avifavir (Favipiravir) in patients with COVID-19, reveals a faster virological response, a shorter time to clinical symptoms and reduced mortality rates compared to patients receiving the Standard of Care (Corritori

et al., Science Spotlight[TM], Conference on Retroviruses and Opportunistic Infections [CROI] 2021). The

ribonucleoside analogue, N4-hydroxycytidine (NHC, EIDD-1931), was initially developed as an influenza

inhibitor, but exerts also broader-spectrum antiviral activity against multiple viruses belonging to

different families of RNA viruses. Activity against SARS-CoV-2 has been reported in cell lines and

primary human airway epithelial cell cultures [5].

Both Favipiravir and NHC act through lethal mutagenesis. Incorporation into viral RNA results in the

accumulation of deleterious transition mutations beyond a permissible error threshold to sustain the

virus population, leading to error catastrophe [6,7]. The orally bioavailable, pro-drug counterpart of NHC

[8], Molnupiravir (EIDD-2801, MK-4482) was shown to result in an antiviral effect against SARS-CoV-2 in

a Syrian hamster model [9], in a mouse model [10] and in ferrets [11]. Data from a first-in-human, phase 1,

randomized, double-blind, placebo-controlled study in healthy volunteers indicate that the drug is well

tolerated and that plasma exposures exceed the expected efficacious doses based on scaling from

animal models [12]. The drug is currently being assessed for its potential as an antiviral treatment of

SARS-CoV-2 infection in Phase 2 clinical trials of infected patients (NCT04405570, NCT04405739). Very

recently interim data (on one secondary objective) were reported in 202 non-hospitalized adults who

had signs or symptoms of COVID-19 and active confirmed SARS-CoV2 infection. At day 5, a reduction

in positive viral culture from nasopharyngeal swabs was noted in subjects who received molnupiravir

(0/47) as compared to placebo (6/25). No safety signals were identified in the 202 participants (Painter

et al., Science Spotlight[TM], CROI 2021).

We here report on the pronounced combined activity of Favipiravir and Molnupiravir in the hamster

infection model.

Results

Dose-response efficacy of Molnupiravir against SARS-CoV-2 in Syrian hamsters

We first evaluated the (dose-response) effect of Molnupiravir (EIDD-2801) in SARS-CoV-2-infected

hamsters to select an appropriate dose for the combination study. Briefly, 6-8 weeks female SG

hamsters were treated orally with Molnupiravir (either 75, 150, or 200 mg/kg, BID) or the vehicle (i.e.

the control group) for four consecutive days starting one hour before intranasal infection with SARS-

CoV-2. At day four post-infection (pi), the animals were euthanized and lungs were collected for

quantification of viral RNA, infectious virus titers and lung histopathology as described before[4] (Fig.

1A). Molnupiravir treatment resulted in a dose-dependent reduction in the viral RNA copies per mg of

lung tissue with 1.3 (P=0.002), 1.9 (P<0.0001) and a 3.3 $\log_{10}$ (P<0.0001) reduction was noted in the

groups that had been treated BID with 75, 150 or 200 mg/kg, respectively (Fig. 1B). A similar pattern

was observed for the infectious virus load in the lungs whereas the intermediate and high doses, but

not the 75 mg/kg dose BID, significantly reduced infectious virus lung titers (Fig. 1C). The reduction in

infectious virus titers ($TCID_{50}$/mg tissue) in the lungs of hamsters treated BID with 150 and 200 mg/kg

was 1.3 (P=0.0002) and 3.5 (P<0.0001) $\log_{10}$, respectively (Fig. 1C). Treatment with 75, 150 and 200

mg/kg Molnupiravir BID significantly reduced the histological lung disease score (P=0.0025, P=0.005,

P<0.0001, respectively) (Fig. 1D). All the doses studied were well tolerated without significant weight

loss or any obvious adverse effects (Fig. 1E).

The combined treatment of Molnupiravir and Favipiravir results in a marked potentiation of efficacy.

Next, we studied what the combined efficacy is of suboptimal doses of Molnupiravir and Favipiravir

(Fig. 2A). Treatment with Favipiravir alone (300 mg/kg, BID) reduced viral RNA and infectious virus

loads in the lungs of infected animals by 0.7 (P=0.0009) and 1.2 (P=0.0002) $\log_{10}$/mg tissue, respectively

(Fig. 2B/C). Treatment with Molnupiravir (150 mg/kg BID) only resulted in 1.9 (P<0.0001) and 1.3

(P=0.0002) $\log_{10}$/mg reduction in viral RNA and infectious virus loads respectively. The combined

treatment resulted in a reduction of 2.7 $\log_{10}$ viral of lung viral RNA titers (Fig. 2B), but interestingly, in

a markedly enhanced reduction in infectious virus titers (4.5 $\log_{10}$ $TCID_{50}$ per mg lung, P=0.02, P=0.0005

as compared to Molnupiravir and Favipiravir alone, respectively) (Fig. 2C). Notably, there was no

detectable infectious virus in the lungs of six out of ten hamsters in the combined treatment group

(Fig. 2C). A marked improvement in the histological lung pathology scores was also observed in the

combined treatment group (Fig. 2D) and no significant weight loss or toxicity signs were observed (Supplementary Fig. S1).

Molnupiravir is known to increase the mutation frequency of MERS-CoV viral RNA in infected mice [5]. To test whether this is also the case in SARS-CoV-2-infected hamsters, we used Illumina deep sequencing to determine the SARS-CoV-2 mutations rate in remaining viral RNA in lung samples of hamsters after treatment. A dose-dependent increase in the mutation count (in particular, C-to-T and G-to-A transitions) in samples from Molnupiravir treated hamsters was observed as compared to the vehicle control group (Supplementary Fig. S2). The Molnupiravir (150mg/kg) + Favipiravir (300 mg/kg) combination resulted in a markedly higher number of C-to-T and G-to-A mutations (68 and 50, respectively), as compared to the single dose groups [150 mg/kg Molnupiravir group (33 and 31, respectively) and 300 mg/kg Favipiravir (14 and 21, respectively)] (Fig. 2E). The C-to-T and G-to-A mutation count in the combination group was also markedly higher than in the highest dose group of Molnupiravir (45 and 39, respectively) (Supplementary Fig. S2). These results may at least partially explain the markedly enhanced reduction in infectious viral loads observed in the combination treatment group.

Discussion

Remdesivir (Veklury), is the first drug to have received FDA approval for use in hospitalised COVID19 patients, although the World Health Organisation has issued a conditional recommendation against its use on 20 Nov 2020 (www.who.int). Remdesivir needs to be administrated intravenously which precludes its use in the early stages of the infection/disease or even a prophylactic use. We previously demonstrated that treatment of SARS-CoV-2-infected hamsters with a high doses of Favipiravir largely reduces infectious virus titers in the animals and results as a consequence in a markedly improved lung pathology [4]. Favipiravir is currently being studied for the treatment of COVID-19 in clinical trials in several countries and was very recently reported to result in COVID-19 patients in a faster virological response, a shorter time to clinical symptoms and reduced mortality rates (Corritori et al., Science

SpotlightTM, CROI 2021). Also Molnupiravir has been reported to exert therapeutic and prophylactic

activity against SARS-CoV-2 in several animal models [9–11]. Importantly very recent interim data from

phase II clinical studies in COVID-19 patients, revealed a reduction in the time required to reach

negative isolation of infectious virus from the nasopharyngeal swabs from participants with

symptomatic SARS-CoV-2 infection (Painter et al., Science Spotlight[TM], CROI 2021). Both Molnupiravir

and Favipiravir have a high barrier to resistance, resistant variants have a loss in fitness and induce

lethal viral mutagenesis [6,7].

We here demonstrate that the combination of suboptimal doses of Molnupiravir and Favipiravir results

in a marked antiviral activity in the hamster model. Infectious virus titers were reduced to undetectable

levels in 6 out of 10 treated animals. A median reduction of 4.5 $\log_{10}$ $TCID_{50}$/mg lung tissue was

achieved, which is markedly more pronounced than what could be expected from an additive activity

of either Molnupiravir (1.3 $\log_{10}$) or Favipiravir (1.1 $\log_{10}$) when dosed alone. This pronounced efficacy

of the combination may partially or even entirely be explained by the increased total mutation count

in viral RNA collected from the lungs of combo-treated hamsters as compared to the single treatment

groups.

In conclusion, the combination of Molnupiravir and Favipiravir (two oral drugs with a high barrier to

resistance for which there is very recent initial evidence that they exert antiviral activity in COVID-19

patients), is particularly effective in the treatment of SARS-CoV2 infections in hamsters. This efficacy

may be explained by an enhanced accumulation of mutations as compared to monotherapy. Our

findings may lay the basis for the design of clinical studies to test the efficacy of the combination of

Molnupiravir and Favipiravir in the treatment of COVID-19.

Methods

SARS-CoV-2

The SARS-CoV-2 strain used in this study, BetaCov/Belgium/GHB-03021/2020 (EPI ISL 109

407976|2020-02-03), was recovered from a nasopharyngeal swab taken from an RT-qPCR confirmed

asymptomatic patient who returned from Wuhan, China in the beginning of February 2020. A close

relation with the prototypic Wuhan-Hu-1 2019-nCoV (GenBank accession 112 number MN908947.3)

strain was confirmed by phylogenetic analysis. Infectious virus was isolated by serial passaging on

HuH7 and Vero E6 cells [4]; passage 6 virus was used for the study described here. Live virus-related

work was conducted in the high-containment A3 and BSL3+ facilities of the KU Leuven Rega Institute

(3CAPS) under licenses AMV 30112018 SBB 219 2018 0892 and AMV 23102017 SBB 219 20170589

according to institutional guidelines.

Cells

Vero E6 cells (African green monkey kidney, ATCC CRL-1586) were cultured in minimal essential

medium (Gibco) supplemented with 10% fetal bovine serum (Integro), 1% L- glutamine (Gibco) and 1%

bicarbonate (Gibco). End-point titrations were performed with medium containing 2% fetal bovine

serum instead of 10%.

Compounds

For the first pilot experiment, EIDD-2801 was kindly provided by Calibr at Scripps Research (USA). For

further studies, Molnupiravir (EIDD-2801) was purchased from Excenen Pharmatech Co., Ltd (China)

and was formulated as 50 or 100 mg/ml (for groups with the highest dose) stocks in a vehicle

containing 10%PEG400 and 2.5% Kolliphor-EL in water. Favipiravir was purchased from BOC Sciences

(USA) and was formulated as a 50 mg/mL stock in 3% sodium bicarbonate.

SARS-CoV-2 infection model in hamsters

The hamster infection model of SARS-CoV-2 has been described before [4,13]. Housing conditions and

experimental procedures were approved by the ethics committee of animal experimentation of KU

Leuven (license P065-2020). For infection, female hamsters of 6-8 weeks old were anesthetized with

ketamine/xylazine/atropine and inoculated intranasally with 50 μL containing 2×10^6 TCID50 SARS-CoV-

2 (day 0).

Treatment regimen

For dose-response treatment, animals were treated twice daily with 75, 150 or 200 mg/kg of EIDD-

2801 by oral gavage just before infection with SARS-CoV-2. For combination therapy, hamsters were

treated from day0 with 150 mg/kg EIDD-2801 (oral gavage) and 300 mg/kg Favipiravir (intraperitoneal,

i.p.) twice daily. All the treatments continued until day 3 pi. Hamsters were monitored for appearance,

behavior and weight. At day 4 pi, hamsters were euthanized by i.p. injection of 500 μL Dolethal

(200mg/mL sodium pentobarbital, Vétoquinol SA). Lungs were collected and viral RNA and infectious

virus were quantified by RT-qPCR and end-point virus titration, respectively.

SARS-CoV-2 RT-qPCR

196   Hamster lung tissues were collected after sacrifice and were homogenized using bead disruption

197   (Precellys) in 350 μL TRK lysis buffer (E.Z.N.A.® Total RNA Kit, Omega Bio-tek) and centrifuged (10.000

198   rpm, 5 min) to pellet the cell debris. RNA was extracted according to the manufacturer's instructions.

199   RT-qPCR was performed on a LightCycler96 platform (Roche) as described before [4].

End-point virus titrations

201   Lung tissues were homogenized using bead disruption (Precellys) in 350 μL minimal essential medium

202   and centrifuged (10,000 rpm, 5min, 4°C) to pellet the cell debris. To quantify infectious SARS-CoV-2

203   particles, endpoint titrations were performed as described before [4].

Histology

For histological examination, the lungs were fixed overnight in 4% formaldehyde and embedded in

paraffin. Tissue sections (5 μm) were analyzed after staining with hematoxylin and eosin and scored

blindly for lung damage by an expert pathologist. The scored parameters, to which a cumulative score

of 1 to 3 was attributed, were the following: congestion, intra-alveolar hemorrhagic, apoptotic bodies

in bronchus wall, necrotizing bronchiolitis, perivascular edema, bronchopneumonia, perivascular

inflammation, peribronchial inflammation and vasculitis.

Deep sequencing and analysis of whole genome sequences

Genomic sequences from all samples were obtained using SureSelect[XT] target enrichment and Illumina

sequencing. Reads generated were trimmed with Trim Galore

(https://github.com/FelixKrueger/TrimGalore). Duplicated reads were removed using Picard

(http://broadinstitute.github.io/picard). Reads from the inoculation sample were mapped to the SARS-

CoV-2 reference genome (NC_045512) from GenBank using BWA-MEM [14]. The mapping quality was

checked using Qualimap and the consensus whole genome sequence was generated using QUASR [15,16].

Reads from the lung samples were mapped to this unique reference sequence. Genomes with less than

less than a 100 read depth were excluded. Variants above 1% and with a minimum of 2 supporting

reads per strand were identified at sites with a read depth of ≥ 10 using VarScan [17].

Statistics

GraphPad Prism (GraphPad Software, Inc.) was used to perform statistical analysis. Statistical

significance was determined using the non-parametric Mann Whitney U-test. P-values of ≤0.05 were

considered significant.

Data Availability
All of the data generated or analysed during this study are included in this published article.

Acknowledgments

We thank Carolien De Keyzer, Lindsey Bervoets, Thibault Francken, Elke Maas, Jasper Rymenants, Birgit

Voeten, Dagmar Buyst, Niels Cremers, Bo Corbeels and Kathleen Van den Eynde for excellent technical

assistance. We are grateful to Piet Maes for kindly providing the SARS-CoV-2 strain used in this study.

We thank Prof. Jef Arnout and Dr. Annelies Sterckx (KU Leuven Faculty of Medicine, Biomedical

Sciences Group Management) and Animalia and Biosafety Departments of KU Leuven for facilitating

the animal studies. This project has received funding from the Covid-19-Fund KU Leuven/UZ Leuven

and the COVID-19 call of FWO (G0G4820N), the European Union's Horizon 2020 research and

innovation program under grant agreements No 101003627 (SCORE project) and Bill & Melinda Gates

Foundation (BGMF) under grant agreement INV-00636. Sequencing was performed by the Pathogen

Genomics Unit at UCL. R.A., C.S.F. and L.L. were supported by a KU Leuven internal project fund. X.Z.

received funding of the China Scholarship Council (grant No.201906170033). J.B. receives funding from

the NIHR UCL/UCLH Biomedical Research Centre. J.P. is funded by the Rosetrees Trust.

Author Contributions

R.A., C.S.F. and J.N. designed the studies; R.A., C.S.F., S.J.F.K., X.Z., J.B., J.P. and L.L. performed the

studies ; R.A., C.S.F., J.B, J.P. and B.W. analyzed data; D.J. and J.N. provided advice on the interpretation

of data; R.A., C.S.F. and J.N. wrote the paper with input from co-authors; A.K.C. and S.D.J provided

essential reagents; V.G. and E.H. provided and facilitated access to essential infrastructure; R.A., C.S.F.,

R.W. and J.N. supervised the study; L.V., L.C., P.L., J.N. and K.D. acquired funding.

Competing Interest Statement: None to declare.

References

1.   Zhu, N. *et al.* A Novel Coronavirus from Patients with Pneumonia in China, 2019. *N. Engl. J.*

Med. **382,** 727–733 (2020).

2.   Tay, M. Z., Poh, C. M., Rénia, L., MacAry, P. A. & Ng, L. F. P. The trinity of COVID-19: immunity,

inflammation and intervention. *Nature Reviews Immunology* **20,** 363–374 (2020).

3.   Delang, L. & Neyts, J. Medical treatment options for COVID-19. *Eur. Hear. J. Acute Cardiovasc.*

Care 204887262092279 (2020). doi:10.1177/2048872620922790

4.   Kaptein, S. J. F. *et al.* Favipiravir at high doses has potent antiviral activity in SARS-CoV-

2−infected hamsters, whereas hydroxychloroquine lacks activity. *Proc. Natl. Acad. Sci. U. S. A.*

117, 26955–26965 (2020).

5.   Sheahan, T. P. *et al.* An orally bioavailable broad-spectrum antiviral inhibits SARS-CoV-2 in

human airway epithelial cell cultures and multiple coronaviruses in mice. *Sci. Transl. Med.* **12,**

(2020).

6.   Urakova, N. *et al.* β-D- N 4 -Hydroxycytidine Is a Potent Anti-alphavirus Compound That

Induces a High Level of Mutations in the Viral Genome . *J. Virol.* **92,** (2017).

7.   Delang, L., Abdelnabi, R. & Neyts, J. Favipiravir as a potential countermeasure against

neglected and emerging RNA viruses. *Antiviral Research* **153,** 85–94 (2018).

8.   Toots, M. *et al.* Characterization of orally efficacious influenza drug with high resistance

barrier in ferrets and human airway epithelia. *Sci. Transl. Med.* **11,** (2019).

9.   Rosenke, K. *et al.* Orally delivered MK-4482 inhibits SARS-CoV-2 replication in the Syrian

hamster model. *Res. Sq.* (2020).

10. Wahl, A. *et al.* Acute SARS-CoV-2 Infection is Highly Cytopathic, Elicits a Robust Innate

Immune Response and is Efficiently Prevented by EIDD-2801. *Res. Sq.* (2020).

11.   Cox, R. M., Wolf, J. D. & Plemper, R. K. Therapeutically administered ribonucleoside analogue

MK-4482/EIDD-2801 blocks SARS-CoV-2 transmission in ferrets. *Nat. Microbiol.* (2020).

12.   Painter, W. P. *et al.* Human Safety, Tolerability, and Pharmacokinetics of Molnupiravir, a Novel

Broad-Spectrum Oral Antiviral Agent with Activity Against SARS-CoV-2. *Antimicrob. Agents*

Chemother. (2021). doi:10.1128/AAC.02428-20

13.   Boudewijns, R. *et al.* STAT2 signaling restricts viral dissemination but drives severe pneumonia

in SARS-CoV-2 infected hamsters. *Nat. Commun.* **11,** 5838 (2020).

14.   Li, H. & Durbin, R. Fast and accurate short read alignment with Burrows-Wheeler transform.

Bioinformatics **25,** 1754–1760 (2009).

15.   Okonechnikov, K., Conesa, A. & García-Alcalde, F. Qualimap 2: Advanced multi-sample quality

control for high-throughput sequencing data. *Bioinformatics* **32,** 292–294 (2016).

16.   Watson, S. J. *et al.* Viral population analysis and minority-variant detection using short read

next-generation sequencing. *Philos. Trans. R. Soc. B Biol. Sci.* **368,** (2013).

17.   Koboldt, D. C. *et al.* VarScan 2: Somatic mutation and copy number alteration discovery in

cancer by exome sequencing. *Genome Res.* **22,** 568–576 (2012).

Fig.1. Dose-response efficacy of Molnupiravir (EIDD-2801) against SARS-CoV-2 in a hamster infection model.
(A) Set-up of the study. (B) Viral RNA levels in the lungs of control (vehicle-treated) and EIDD-2801-treated (75, 150 or 200 mg/kg, BID) SARS-CoV-2−infected hamsters at day 4 post-infection (pi) are expressed as $\log_{10}$ SARS-CoV-2 RNA copies per mg lung tissue. Individual data and median values are presented. (C) Infectious viral loads in the lungs of control (vehicle-treated) and EIDD-2801-treated SARS-CoV-2−infected hamsters at day 4 pi are expressed as $\log_{10}$ TCID$_{50}$ per mg lung tissue Individual data and median values are presented. (D) Cumulative severity score from H&E stained slides of lungs from control (vehicle-treated) and EIDD-2801-treated SARS-CoV-2−infected hamsters. Individual data and median values are presented and the dotted line represents the median score of untreated non-infected hamsters. (E) Weight change at day 4 pi in percentage, normalized to the body weight at the time of infection. Bars represent means ± SD. Data were analyzed with the Mann−Whitney U test. *$P < 0.05$, **$P < 0.01$, ***$P < 0.001$, ****$P < 0.0001$, ns=non-significant. All data (panels B, C, D, E) are from two independent experiments except for the 75 mg/kg group. The number of animals were 15, 6, 10, and 10 for respectively the vehicle 75, 150, and 200 mg/kg condition.

Fig.2. Combined efficacy of Favipiravir and Molnupiravir (EIDD-2801) against SARS-CoV-2 in a hamster infection model. (A) Set-up of the study. (B) Viral RNA levels in the lungs of control (vehicle-treated), Favipiravir-treated (300 mg/kg, BID), EIDD-2801-treated (150 mg/kg, BID) and combination-treated (Favipiravir+EIDD-2801 at 300+150 mg/kg, BID, respectively) SARS-CoV-2−infected hamsters at day 4 post-infection (pi) are expressed as $\log_{10}$ SARS-CoV-2 RNA copies per mg lung tissue. Individual data and median values are presented. (C) Infectious viral loads in the lungs of control (vehicle-treated), Favipiravir-treated, EIDD-2801-treated and combination-treated (Favipiravir+EIDD-2801) SARS-CoV-2−infected hamsters at day 4 pi are expressed as $\log_{10}$ $TCID_{50}$ per mg lung tissue. Individual data and median values are presented. (D) Cumulative severity score from H&E stained slides of lungs from control (vehicle-treated), Favipiravir-treated, EIDD-2801-treated and combination-treated (Favipiravir+EIDD-2801) SARS-CoV-2−infected hamsters. Individual data and median values are presented and the dotted line represents the median score of untreated non-infected hamsters. (E) Mean mutation count (per the whole genome) in the viral RNA isolated from the lungs of control (vehicle-treated), Favipiravir-treated (300 mg/kg, BID), EIDD-2801-treated (150 mg/kg, BID) and combination-treated (Favipiravir+EIDD-2801 at 300+150 mg/kg, BID, respectively) SARS-CoV-2−infected hamsters at day 4 post-infection (pi). Data were analyzed with the Mann−Whitney U test. *P < 0.05, **P < 0.01, ***P < 0.001, ****P < 0.0001, ns=non-significant. Favi=Favipiravir, EIDD=EIDD-2801. All data (panels B, C, D) are from two independent experiments with 15, 10, 10 and 10 animals for respectively the vehicle, Favipiravir 300 mg/kg, EIDD-2801 150 mg/kg and Favipiravir+EIDD-2801 condition.

Fig. S1. Tolerability of combined treatment with Favipiravir and EIDD-2801 in SARS-CoV-2-infected hamsters.
Weight change at day 4 post-infection in percentage, normalized to the body weight at the time of infection.
Bars represent means ± SD.

Fig. S2. Dose-dependent increase in mutation count in SARS-CoV-2 viral RNA by Molnupiravir (EIDD-2801).
Mean mutation count (per the whole genome) in the viral RNA isolated from the lungs of control (vehicle-treated), EIDD-2801-treated (75, 150 or 200 mg/kg, BID) and combination-treated (Favipiravir+EIDD-2801 at 300+150 mg/kg, BID, respectively) SARS-CoV-2–infected hamsters at day 4 post-infection (pi).