Metabolic reprogramming and host-immune response against Crimean-Congo Hemorrhagic Fever Viruses

Short title: Metabolic reprogramming in CCHFV

Authors

Ujjwal Neogi[1,2, #*], Nazif Elaldi[3#], Sofia Appelberg[4], Anoop T. Ambikan[1], Emma Kennedy[5,10], Stuart Dowall[5], Binnur K. Bagci[6], Soham Gupta[1], Jimmy E. Rodriguez[7], Sara Svensson-Akusjärvi[1], Vanessa M. Monteil[1], Ákos Végvári[7], Friedemann Weber[8], Roger Hewson[5,10,11], Ali Mirazimi[1,4,9*]

Affiliations

[1]Division of Clinical Microbiology, Department of Laboratory Medicine, Karolinska Institute, ANA Futura, Campus Flemingsberg, Stockholm, Sweden.

[2]Department of Molecular Microbiology and Immunology and the Bond Life Science Center, University of Missouri, Columbia, MO 65211, USA

[3]Deptartment of Infectious Diseases and Clinical Microbiology, Medical Faculty, Cumhuriyet University, Sivas, Turkey

[4]Public Health Agency of Sweden, Solna, Sweden

[5]Public Health England, Porton Down, Salisbury, Wiltshire SP4 0JG, England.

[6]Department of Nutrition and Dietetics, Faculty of Health Sciences, Sivas Cumhuriyet University, Sivas, Turkey

[7]Division of Chemistry I, Department of Medical Biochemistry and Biophysics, Karolinska Institutet, Stockholm, Sweden

[8]Institute for Virology, FB10-Veterinary Medicine, Justus-Liebig University, Giessen, Germany

[9]National Veterinary Institute, Uppsala, Sweden

[10]Oxford Brookes University, Headington Road, Oxford OX3 0BP

[11]Faculty of Infectious & Tropical Diseases, London School of Hygiene and Tropical Medicine, Keppel Street, London, WC1E 7HT

[#]These authors contributed equally to this work

*Corresponding Authors: Ali Mirazimi (ali.mirazimi@folkhalsomyndigheten.se), Ujjwal Neogi (ujjwal.neogi@ki.se)

Abstract

The pathogenesis and host-viral interactions of the Crimean–Congo hemorrhagic fever virus (CCHFV) are convoluted and have not been evaluated previously. To understand the host immune responses against CCHFV, we have performed a global transcriptomic analysis of peripheral blood mononuclear cells from a longitudinal cohort of CCHF patients who survived and temporal untargeted proteomics analysis of CCHFV infected cells. Our results indicate that during the acute phase of CCHFV infection, the host's metabolic reprogramming towards central carbon metabolism including glycolysis/gluconeogenesis occurs. This could be regulated by the PI3K/Akt, HIF-1, FoxO, and AMPK signaling pathways and play a central role in viral replication. Moreover, key interferon stimulating genes (ISGs; ISG12, ISG15, ISG20 and MXs; Mx1 and Mx2) are activated during infection, suggesting a role for type I and II interferon-mediated antiviral mechanisms. Targeting type I interferon response through metabolic perturbation could be an attractive therapeutic intervention for CCHFV as well as aid as antiviral strategies.

Introduction

Crimean–Congo hemorrhagic fever orthonairovirus (CCHFV) is considered to be a major emerging viral threat causing an increasing number of outbreaks in several parts of the globe. CCHFV is a negative-sense RNA virus belonging to the *Nairoviridae* virus family, which causes a mild to severe viral hemorrhagic fever. Crimean–Congo hemorrhagic fever (CCHF) poses a substantial threat to public health due to its high mortality rate in humans (3–40%), modes of transmission (tick-to-human/animal, animal-to-human and human-to-human) and geographical distribution (WHO, 2020). The Ixodid ticks, especially those of the genus, *Hyalomma*, are both a vector and a reservoir for CCHFV and are highly ubiquitous with its presence in more than 40 countries (ECDC, 2020). CCHF is endemic in almost 30 countries that are widely distributed throughout sub-Saharan Africa, South-Eastern Europe, the Middle-East, Central Asia down to Pakistan, and as far as east as the Xinjiang province of Northwest China(Bente, Forrester et al., 2013, Zivcec, Scholte et al., 2016). In recent years, CCHFV outbreaks have become more frequent and have spread to new geographical areas; this has been attributed to climate changes and the spread of infected ticks by birds and the livestock trade. The presence of the CCHFV vector in Portugal, Spain, Germany, and even Sweden (Grandi, Chitimia-Dobler et al., 2020) and England (McGinley, Hansford et al., 2020) highlights the need for stricter surveillance due to the possibility of a future epidemic(Estrada-Peña, Sánchez et al., 2012).

Type-I interferon (IFN-I) is one of the first-line of antiviral innate immune defenses against viruses. Following infection, viral RNA or its replication products can induce secretion of IFN-I that subsequently stimulates interferon-stimulated genes (ISGs) in a tightly controlled signaling cascade. Thus, components of the IFN-I signaling pathway are often targeted by viruses to evade the antiviral immune response. It is known that CCHFV replication is sensitive to IFN-I (Andersson, Lundkvist et al., 2006). However, the virus is also able to delay the induction of IFN, and IFN treatment has been shown to be ineffective following establishment of infection, suggesting that CCHFV has developed mechanisms to block innate immune responses(Andersson, Karlberg et al., 2008). In a limited number of studies, CCHF patients were shown to have elevated innate immune response-associated cytokines and chemokines, of which IL-6 and TNF-α were often associated with mortality (Kaya, Elaldi et al., 2014, Korva, Rus et al., 2019, Weber & Mirazimi, 2008). CCHF was also observed to be associated with increased secretion of IFN-α2 and IFN-γ, irrespective of severity and disease fatality(Korva et al., 2019, Papa, Bino et al., 2006). The existing knowledge from in vitro studies, autopsies of CCHF patients and clinical findings suggests a regulatory mechanism of the IFN-response during

CCHFV infection and indicates that IFN stimulated genes may offer protection by inhibiting
CCHFV. However, it is not known which among the more than 1000 ISGs are responsible for the
IFN effect against CCHFV or how the strength of viral anti-IFN mechanisms and cytokine
profiles might correlate with the virulence level of particular CCHFV strains.

Because of the sporadic nature of CCHF outbreaks in humans in the endemic regions, a lack of
infrastructure, and the absence of structured studies, little is known about the pathogenesis and
host-virus interactions during the acute phase of CCHF disease and associated sequelae after
recovery. Understanding the host response to CCHFV is necessary to design better therapeutic
and containment strategies for CCHF. Recently, systems biology studies using -omics approaches
on patient material and infected cells have offered a route to elucidate potential mechanisms of
the host immune response against invading viruses and disease pathogenesis, as recently reported
by us in case of SARS-CoV-2 infection (Appelberg, Gupta et al., 2020).

In this study, we have applied transcriptomics and proteomics analysis to understand the host
immune and cellular responses against CCHFV. We have used longitudinal blood samples
collected during both the acute phase of CCHFV infection and the convalescent-phase (nearly
after a year of recovery), as well as an *in vitro* infection model in Huh7 cells,. Our study is the
first to provide a comprehensive understanding of the fundamental innate immune responses that
modulate successful CCHFV infection.

Results

Samples and clinical data

In this study, 18 samples were collected during the acute phase of the disease with a median time
of 4 days (range 1-6 days) after the onset of symptoms. By using SGS, 33% (6/18) patients were
grouped into severity group 1, 61% (11/18) patients into severity group 2 and, 6% (1/18) patients
into severity group 3. The median age of the patients was 49 (range: 18–79) years, and 12
(66.7%) of the patients were male. A 79-year-old male patient in severity group 3 died on the
third day of hospitalization. The case-fatality-rate (CFR) for the cohort was 5.6%. Follow-up
samples were collected from 12 individuals after a median duration of 54 weeks (range: 46-57
weeks) (Fig. 1a). The CCHF patient characteristics are summarized individually in
Supplementary Table S1.

The impact of CCHF severity on peripheral blood transcriptome during acute infection

We first considered the effect of the SGS disease severity group on the gene expression profile at

the mRNA level. The UMAP analysis using all of the genes did not show distinct differences

between samples of different severity groups (Figure 1b). The heterogenous gene-expression

profile was further refined by hierarchical clustering analysis of the top 500 high variable genes

(Supplementary Figure S1). To specifically investigate the genes that were significantly

associated with disease severity, the samples were grouped into either severity group 1 or severity

groups 2 and 3. The differential gene expression (DGE) was plotted as an MA -plot, as a measure

of the number of genes whose expression was consistently altered between these two groups

(Figure 1c). There were 12 genes (ERG, PROM1, HP, HBD, AHSP, CTSG, PPARG, TIMP4,

SMIM10, RNASE1, VSIG4, CMBL, MT1G) that were significantly upregulated in patients in

severity groups 2 and 3 compared to severity group 1 (Figure 1c). However, no obvious links

between these genes were noted.

Cytokine levels in severity group 1 and severity group 2 during acute CCHFV infection

Serum samples collected during the acute phase of the disease from severity group 1 (n=6) and

severity group 2 (n=11) were analyzed using Luminex assay to assess the soluble marker. Of the

22 markers used for analysis, only IL-8 and GM-CSF were significantly higher in severity group

2, and a near significant upregulation was also observed with TNF-α and IL-10 compared to

severity group 1 (Figure 1d). The levels of all of the rest 18 biomarkers are shown in

Supplementary Figure S2. Cumulatively, the transcriptomics and cytokine array data suggest that

there were no significant inflammatory differences with severity in patients who survived CCHF

infection.

Transcriptomics changes in the acute phase and post-CCHF recovery

Due to the natural heterogeneity in human cohorts, we used longitudinal samples to perform

differential expression analyses for each infected patient between the time of infection and

approx. 1 year post-recovery (Range: 46 – 57 weeks). There were 12 patients (severity group 1:

n=5; severity group 2: n= 7) available for which sampling was done both during acute infection

and post-recovery. We observed a clear separation between the acute phase and the recovered

phase samples using UMAP clustering (Supplementary Figure S3).

The DGE profile for the acute phase compared to the recovered phase in all patients showed an

upregulation of 2891 genes and a downregulation of 2738. The complete profile is given in

Supplementary Data File 1. The pathways that are significantly modulated by the upregulated

genes (FDR<0.05) are mainly related to metabolic pathways like glycolysis/gluconeogenesis, N-
glycan biosynthesis and oxidative phosphorylation (OXPHOS), and antiviral pathways like the
NOD-like receptor signaling pathway (Figure 2b and Supplementary Data File 2). However, the
pathways related to the down-regulated genes were mainly related to antiviral defense
mechanism-associated pathways including innate immune responses like Th1, Th2, and Th17,
cell differentiation, the NF-kB pathways, chemokine signaling pathway, etc. (Figure 2c).
Interestingly, although the HIF-1 signaling pathway was downregulated, the expression of the
HIF-1 targeted genes that play a role in glycolysis/gluconeogenesis and coding for the glycolytic
enzymes, Enolase 1 (ENO1) and Phosphoglucomutase 2 (PGM2) was significantly upregulated,
while hexokinase 1 (HK1) were downregulated (Supplementary Data File 1). However, when we
compared the acute phase with the recovered phase in severity group 1, and severity group 2
separately, there was a distinct DEG profile. In severity group 1 the differentially expressed genes
were significantly fewer (n=1617, upregulated: 954 and downregulated: 663) compared to those
in severity group 2 (n=4256, upregulated: 2182 and downregulated: 2074) (Figure 2d and 2e).
This indicates that the disease severity had a significant effect on gene expression profiling during
the acute phase. There were 1451 overlapping genes between severity group 1 and severity group
2 that were differentially up-regulated (n=882) and downregulated (n=569). Using gene ontology
(GO) analysis after removal of the redundant terms using REVIGO (Supek, Bošnjak et al., 2011),
the top two GO terms that were significantly upregulated were the type I interferon signaling
pathway (GO:0060337) and the regulation of viral genome replication (GO:0045069) (Figure 2f).

Interferon signaling pathways in CCHFV infection:

We observed significant differences in gene expression relating to interferon signaling in both
severity group 1 and severity group 2, and thus we further explored the alteration of gene
expression in all the interferon signaling related pathways. To this end, we used supervised
analysis with the genes that are part of 13 GO terms and are part of both type-I and type-II
interferon signaling pathways (Figure 3a). Of the IFN-regulated genes, IFI27 (ISG12) showed the
most robust upregulation (Figure 3b). This was further supported by the RNAscope analysis
targeting the IFI27 transcript in the SW13 cell line infected with CCHFV strain IbAr10200
(Figure 3c). Among the pathways, the maximum significantly differentially expressed genes were
part of cellular response to ITN-γ (GO:0071346) (n=57) followed by the Type I interferon (IFN-I)
signaling pathway (GO:0060337) (n=35) (Figure 3a) and the majority of genes responsible for
IFN-I signaling (89%; 31/35) were upregulated in the acute phase (Figure 3d). IFI27 has been
previously shown to be upregulated specifically during virus infection and has the ability to

restrict several viral infections. Apart from this ISG20, ISG15, Mx1, Mx2 and several IFN-α-

inducible proteins (IRFs) showed upregulation in the acute phase.

181 Quantitative temporal proteomics analysis of CCHFV infection in Huh7 cell lines:

Our transcriptomics analysis of CCHF patient samples revealed the role of interferon signaling

pathways in the pathogenesis of CCHFV, with several ISGs being elevated during the acute phase

of infection. To understand the global changes in the host interferome (cluster of interferon genes)

during CCHFV infection, we infected Huh7 cells with CCHFV and used a time-course

experiment for 24 and 48 hpi. We then performed TMT-based mass-spectrometric analysis

(Figure 4a). The global proteome analysis revealed changes in several pathways that were related

to metabolism and the organismal system (Figure 4b). This was in line with the changes we

observed in the patient transcriptomics analysis (Figure 2b and 2c). Overall, 16 pathways were

observed both in patients' transcriptomics and *in vitro* infection assays in Huh7 cells

(Supplementary Figure S4).

To identify changes in the interferon landscape, we used the GO terms for the cellular response to

IFN-γ (GO:0071346) and the Type I interferon signaling pathway (GO:0060337) that showed

significant changes in expression in patient samples during acute infection. The overall changes in

the interferome over time are represented as a heat-map in Figure 5a. The significant changes in

the protein levels at different time points are represented as volcano plots (Figure 5b-c). At 24hpi,

several ISGs, like Mx1, ISG20 and IFI16, that were upregulated in the acute-phase patient

samples were also significantly elevated in infected Huh7 cells (Figure 5b). Mx1 retained a

sustained elevation 48 hpi, and ISG15 was also significantly elevated (Figure 5c). The most

interesting ISG was Mx2 that showed a suppression at 24hpi and a significant elevation at 48hpi

(Figure 5d).

We have previously shown the anti-CCHF role of Mx1 and the interplay between ISG15 and

CCHFV is well documented by us (Andersson, Bladh et al., 2004) and others(Scholte, Zivcec et

al., 2017). We also observed induction of ISG20, a 3′-to-5′ exonuclease that showed elevation in

both infected patients and infected cell lines. ISG20 has been reported to have anti-viral activity

against a number of bunyaviruses(Feng, Wickenhagen et al., 2018). In order to ascertain its effect

on CCHFV replication, we used doxycycline inducible stable cell lines that expressed human

ISG20 and CAT protein as a control. Following infection with CCHFV at an MOI of 1 for 48 h,

there was some degree of inhibition noted in the doxycycline-induced ISG20 cell line (p=0.26)

and no inhibition was observed in the CAT control cell line (Figure 5e). Though not significant,

the results are suggestive of an antiviral property of ISG20 that is dependent on the enzymatic

activity and this requires further investigation. Furthermore, there is also a possibility that the
virus has evolved measures to counteract these antiviral responses. During the full course of
infection, there was significant suppression of IRF3, the transcription factor for IFNβ production,
indicating viral mediated evasion or delay in activation of the immune response. The time-
dependent stimulation of different ISGs by CCHFV suggests differential regulation of the IFN
pathway and does not rule out an IFN-I independent mechanism of activation of ISGs.

Discussion

Metabolic reprogramming during innate antiviral immune responses changes the subsequent phases of disease progression and influences the adaptive immune response. In this study, using the genome-wide transcriptomic analysis of a longitudinal patient cohort, and temporal proteomics from *in vitro* infection assays in Huh7 cells, we showed that metabolic reprogramming could facilitate optimal CCHFV replication and the antiviral mechanism of the host. Our data indicated that during the acute phase of CCHFV infection there is metabolic reprogramming of host cells towards central carbon metabolism (e.g. glycolysis/gluconeogenesis, pyruvate metabolism, citrate cycle/TCA cycle, pentose phosphate pathway etc) that could potentially be regulated by the PI3K/Akt, HIF-1, FoxO, and AMPK signaling pathways and play a central role in viral replication (Figure 6 and Supplementary Figure S5). Moreover, type I and II interferon-mediated antiviral mechanisms were also activated with elevated key antiviral ISGs (ISG12, ISG15, ISG20), and MXs (Mx1 and Mx2).

Both DNA and RNA viruses have been known to alter the host cell metabolism to promote their replication. Such metabolic changes follow similar patterns that include an alteration of the central carbon metabolism and a redirection of the carbon supply to metabolic pathways that are crucial for replication, indeed several virus-specific signatures have been observed in previous studies(Mayer, Stöckl et al., 2019). However, due to the lack of patient cohorts and the high-level of containment required for CCHFV, knowledge about host-virus interaction has been limited(Papa, Tsergouli et al., 2017) until now. The transcriptomics data on patient materials and *in vitro* cell culture assays indicated transient dysregulation of glycolysis/gluconeogenesis after CCHFV infection, which is known to promote replication in several other RNA viruses including HIV-1, rubella virus, dengue virus, rhinovirus, hepatitis C virus (HCV), influenza virus etc (Eisenreich, Rudel et al., 2019, Fontaine, Sanchez et al., 2015, Mayer et al., 2019, Smallwood, Duan et al., 2017, Thaker, Ch'ng et al., 2019). The upregulation of glycolysis/gluconeogenesis in CCHFV infection could be mediated by the HIF-1 signaling pathway, as the HIF-targeted genes coding for the key glycolytic enzymes ENO1 and PGM2 were significantly upregulated in CCHFV infected patients during the acute phase. We therefore hypothesize that during the CCHFV infection, the stabilization and subsequent activation of HIF-1 regulates the glycolysis/gluconeogenesis that is required for virus replication. This is further supported by our previous study in which exogenous nitric oxide (NO) inhibited CCHFV *in vitro* (Simon, Falk et al., 2006) NO regulates the HIF-1 via the PI3K/Akt pathway under normoxic conditions (Sandau, Faus et al., 2000). Interestingly we have observed a similar interplay of HIF-1 signaling and

PI3K/Akt signaling in SARS-CoV-2(Appelberg et al., 2020), where NO is able to suppress SARS-CoV-2 replication *in vitro*(Akaberi, Krambrich et al., 2020). The growing interest in hypoxia-induced signaling in emerging and re-emerging viruses and its relation with NO biology is expected to lead to further insights into the complex roles of NO, and to HIF signaling as a crucial feature of NO-mediated events during acute viral diseases. Several genes were significantly downregulated in the PI3K/Akt pathway during acute CCHFV infection, indicating the dysregulation of apathway that is critical for the cellular defense against invading pathogens.

Interestingly, an *in vitro* study in another *Bunyaviridae* virus, Rift valley fever virus (RVFV), identified the inhibition of the PI3K/Akt pathway by dephosphorylation of the AKT and Forkhead box protein O1 (FoxO1) (Popova, Turell et al., 2010). In our study we observed that the majority of the coding genes in the FoxO signaling pathway were downregulated by CCHFV, including the FOXO transcription factors FOXO1 and FOXO3, that can act as negative feedback regulators of the innate cellular antiviral response(Lei, Zhang et al., 2013). This could be due to the regulation of PI3K signaling as observed in the Epstein-Barr virus (EBV) infection (Shore, White et al., 2006). FoxO1 and FoxO3 also play an essential role in the immunometabolic dynamics and are important targets for glycolysis and gluconeogenesis (Lundell, Massart et al., 2019, Puigserver, Rhee et al., 2003).

Another critical signaling pathway that was downregulated during the CCHFV infection was AMP-activated protein kinase (AMPK) signaling that has an antiviral mechanism in addition to its role in regulating glycolysis/gluconeogenesis and FoxO1 transcriptional activity (Saline, Badertscher et al., 2019). Earlier *in vitro* studies showed that HCV inhibits the AMPK signaling pathway to promote viral replication, resulting in the downregulation of IFN-I signaling (Mankouri, Tedbury et al., 2010, Tsai, Chang et al., 2017). Interestingly, activation of the AMPK signaling pathway has been shown to inhibit the replication *in vitro* of the bunyavirus RVFV by inhibiting fatty acid synthesis (Moser, Schieffer et al., 2012). This indicates that modulation of the AMPK signaling could be an attractive therapeutic intervention for CCHFV infection through metabolic reprogramming or modulation of the IFN-I response.

The interferon signaling pathway plays a central role in innate immune responses against viruses and mediates stimulation of antiviral ISGs and secretion of pro-inflammatory cytokines. RIG-I has been recognized as the pathogen recognition receptor that recognizes CCHFV RNA, subsequently initiating a signaling cascade leading to secretion of IFN-I, mainly IFN-α and IFN-β (Spengler, Patel et al., 2015). The secreted IFN-I is further identified by the type I interferon (IFN-α/β) receptor (IFNAR) which initiates JAK-STAT signaling, leading to the stimulation of

ISGs. The protective role of IFN-I against CCHFV has been exemplified in mouse models, in which IFNAR$^{-/-}$ or STAT-1$^{-/-}$ mice (Bente, Alimonti et al., 2010, Zivcec, Safronetz et al., 2013) or STAT2$^{-/-}$ hamsters (Ranadheera, Valcourt et al., 2020) showed enhanced susceptibility to CCHFV infection. Even in *in-vitro* experiments, pre-treatment of cells with IFN-α was found to be inhibitory to CCHFV (Andersson et al., 2008). Though CCHFV is inhibited by the IFN - response, not many ISGs with anti CCHFV activity have been identified appart from MxA although, ISG20 and PKR have been proposed (Andersson et al., 2004) to have anti-CCHFV activity. Nevertheless, the anti-CCHFV activity of ISGs has not been characterized. Furthermore, in the STAT-1$^{-/-}$ mice model, CCHFV induced several IFN-stimulated genes in a delayed fashion (Zivcec et al., 2013). However, there are limited data on the effect of CCHFV on ISG stimulation in human samples. In humans, the role of the IFN response in CCHFV infection is noted by the fact that Toll-like receptor (8/9 or 3) polymorphism was shown to be associated with increased infection and fatality (Engin, Arslan et al., 2010). In the present study we mapped the ISGs induced during the acute phase of CCHFV infection compared to the recovered phase and observed that several ISGs related to the IFN-I pathway were upregulated (Figure 3a and 3b). We also observed that the ISGs with known or proposed anti-CCHFV activity, i.e. Mx1, ISG15 and ISG20 were upregulated.

Additionally, we observed significant upregulation of IFI27 (ISG12), IFITs, OASs and ISG15 (Figure 3b). IFI27 has been shown to have an antiviral effect on West-Nile virus (Lucas, Richner et al., 2015) but its impact on other viruses has not been well-studied. IFI27 was shown to distinguish between influenza and bacteria in respiratory infections (Tang, Shojaei et al., 2017) and is present at high levels in severe cases of Enterovirus 71 (EV71)-induced severe hand foot and mouth disease (Min, Ye et al., 2020), indicating its potential to be used as a biomarker. Though statistical significance was not reached, IFI27 twas 1.4-fold higher in severity groups 2 and 3 compared to severity group 1. Our study observed that IFI27 was highly differentially upregulated in CCHF infection; its role in controlling the infection therefore and its possibile use as a biomarker requires interrogation.

We further observed changes in the induction of ISGs during infection in a time-dependent manner *in vitro* by infecting the Huh7 cell line with CCHFV and observed several ISGs to be upregulated. Though we did not identify IFI27 in proteomics data, several ISGs including Mx1, OAS3, ISG20, ISG15, IFIT3, IFITM3, IFI16 that were upregulated in the acute phase CCHFV patient samples were observed to be also upregulated in the cell-infection model (Figure 5a-d). Among these, ISG20 has been shown to have antiviral activity against related-Bunyaviruses

318 (Feng et al., 2018), and we observed some degree of inhibitory activity of ISG20 on CCHFV
319 replication (Figure 5e). Notably, Mx2 was differentially regulated in a time-dependent manner. At
320 24 hpi, Mx2 was significantly downregulated but it was upregulated by 48 hpi (Figure 4a, b and
321 c), suggestive of a suppressive mechanism by the virus during early infection. The virus has
322 evolved several mechanisms of evading the antiviral effects of the ISGs as observed in the case of
323 ISG20 in our study, and one such mechanism is the de-ubiquitinase and de-ISGylase activity of
324 the ovarian tumor (OTU)-protease encoded by the L1 protein (Scholte et al., 2017) that could
325 directly inhibit ISG15 activity.

326 Similar to IFN-I, proinflammatory cytokines are important for effective innate immunity. CCHFV
327 patients are known to have higher levels of plasma cytokine and chemokine compared to healthy
328 people. Elevated levels of IL-6, IL-10 and TNF-α are most commonly associated with fatal
329 outcomes(Ergonul, Tuncbilek et al., 2006, Korva et al., 2019, Saksida, Duh et al., 2010). A few
330 studies have suggested that the co-elevation of IL-6 and TNF-α are strongly associated with
331 fatality (Ergonul et al., 2006). Our study did not include samples from fatal cases and thus we
332 performed a cytokine array and compared moderate and severe cases. Our results further
333 confirmed the roles of TNF-α, IL-10, IL-8 and GM-CSF since these showed significant or near-
334 significant elevation in severe cases. Korva et al., also observed significant differences in IL-10
335 and GM-CSF between moderate and severe cases(Korva et al., 2019). In another study by Hulva
336 et al., changes in IL-6 and TNF-α were noted between moderate and severe cases, but no
337 significant change in IL-8 was observed (Hülya, Yilmaz et al., 2017). Altough IL-6 is often
338 associated with disease severity and fatality, we did not observe any significant difference in IL-6
339 levels.

340 In conclusion, our study provides a comprehensive description of the host-immune response
341 against CCHFV that can explain viral pathogenesis. The interplay of the metabolic
342 reprogramming and the IFN-mediated host antiviral mechanism could provide attractive options
343 for therapeutic intervention of CCHF. The specific role of the different ISGs that were regulated
344 during the course of infections need to be studied further to provide important clues for the
345 development of antivirals or vaccines.

346 **Materials and Methods**

347 **Ethics statement and biosafety**

This study was approved by the Local Research Ethics Committee of the Ankara Numune Education and Research Hospital, Turkey (Protocol # 17-1338) and Regional Ethics Committee, Stockholm (Dnr. 2017-/1712-31/2). All patients and/or their relatives were informed about the purpose of the study and signed a consent form before collection. All the procedures were performed according to the local authority biosafety guidelines and biosafety level 4 (BSL4) practice.

Study design, patients, and sample collection:

We enrolled 18 adult patients (≥18 years) diagnosed with CCHF who were followed up by the clinical service of Infectious Diseases and Clinical Microbiology of Sivas Cumhuriyet University Hospital, Sivas, Turkey. Information about the patients was collected, including age, gender, date of symptoms onset, date of hospitalization, date of discharge/death, Severity Grading System (SGS) scores and severity groups, and outcome data (Supplementary Table 1). The CCHF patients were divided into three groups using the SGS scores of 1, 2 and 3(Bakir, Engin et al., 2012). Blood samples were collected on the admission day (acute stage) and from the survivors one year after their recovery. The samples were used for diagnosis and were sent to the CCHF Reference Laboratory, Ankara, Turkey (Public Health of Turkey). For the detection of CCHFV nucleic acid, a commercial real-time RT-PCR test (Altona Diagnostics®, Hamburg, Germany) was used, and for the serological diagnosis, a commercial IgM indirect immunofluorescence antibody (IFA) assay (Euroimmun®, Luebeck, Germany) was used. Patients having positive test results by PCR and/or serologic analyses were diagnosed with CCHF and such patients were included in the study. Peripheral blood mononuclear cells (PBMCs) were isolated by the modified density gradient centrifugation method. PBMCs and plasma were sent to the Public Health Agency of Sweden, Stockholm, Sweden. Serum cytokine profiling targeting 22 cytokines/chemokines was performed by Public Health England using methods as previously described (Dowall et al)(Dowall, Graham et al., 2019) Briefly, a 22 -plex customised luminex kit (Merck Millipore, Darmstadt, Germany) was designed to quantitate: Eotaxin, Granulocyte-Colony Stimulating Factor (G-CSF), Granulocyte Macrophage-Colony Stimulating Factor (GM-CSF), Interferon-alpha 2 (IFN-a2), Interferon-gamma (IFN-γ), Interleukin (IL)-10, IL-12, IL-15, IL-17a, IL-1a, IL-9, IL-1b, IL-2, IL-4, IL-5, IL-6, IL-8, IFN-γ-Inducible Protein 10 (IP-10), Monocyte Chemotactic Protein 1 (MCP-1), Macrophage Inhibitory Protein (MIP)-1a, MIP-1b and Tumour Necrosis Factor alpha (TNF-a). Two quality control samples and a standard included with the kit were prepared as per the manufacturers instructions. 25µl of each QC and standard concentration and 25µl of serum matrix solution were added to relevant wells of a 96 well plate in

duplicate. 25µl of each patient plasma sample and 25µl of assay buffer was added to relevant wells in duplicate. Custom pre-mixed beads were sonicated and vortexed and 25µl added to each test and standard/QC well. Plates were incubated overnight at 4°C. Following incubation, plates were washed twice with 200µl wash buffer using a hand held magnet. 25µl of detection antibody was added to each well and the plates incubated at room temperature for 1hr. 25µl streptavidin-phycoerythrin was added to each well for 30 minutes at room temperature. Each well was washed twice with 200µl wash buffer using a hand-held magnet and 100µl of 10% formalin added to each well: 40% w/v formaldehyde solution (Scientific Laboratory Supplies, Nottingham England) diluted 1:9 v/v with phosphate buffered saline solution (Thermo Fisher, Loughborough, England). The plate was subjected to formaldehyde fumigation prior to removal from Biosafety level (BSL)-4 facilities. At BSL2, the plates were washed twice with 200µl wash buffer and once with 200µl sheath buffer using a hand held magnet to remove formalin, prior to addition of 150µl sheath fluid to resuspend beads. A Luminex MAGPIX instrument using Exponent software (Invitrogen, Paisley, UK) was used to obtain results in the form of MFI (median fluorescence intensity), which was converted to concentration (pg/ml) using the standard curve prepared. Plate data was accepted based on negative background wells and QC data within the expected ranges.

RNA extraction and sequencing (Illumina RNAseq):

Total RNA was extracted from Trizol-treated PBMC using the Direct-zol RNA Miniprep (Zymo Research, CA, USA) according to the manufacturer's protocol, and the RNA quality was determined by Agilent RNA 6000 Pico kit on an Agilent 2100 Bioanalyzer (Agilent Technologies, Waldbronn, Germany). RNA-Seq was performed at the National Genomics Infrastructure, Science for Life Laboratory, Stockholm, Sweden, as described by us previously(Zhang, Ambikan et al., 2018).

Data processing and transcriptomics analysis:

The raw reads were first checked for quality and adapter content using the FastQC v0.11.8 (https://github.com/s-andrews/FastQC) tool. The Illumina adapter sequences found in the raw reads were trimmed using the tool Cutadapt v1.18 (Martin, 2011). The mean quality score across each base position in the cleaned reads was above Q30. The cleaned reads were then aligned against the human reference genome version GRCh38. The reference genome was obtained from the Illumina iGenomes repository. RNASeq aligner STAR v2.6.1a_08-27 was used for the alignment (Dobin, Davis et al., 2013). A gene-level read count table was generated from the alignment using featureCounts v1.6.3 (Liao, Smyth et al., 2014). The read count table was further

split into protein-coding genes and non-coding genes. There were 22,022 protein-coding genes.

All the downstream analysis was performed on the protein-coding genes.

The raw read count data were first normalized to log2 transformed counts per million (CPM) to

remove the library size effect from the R/Bioconductor package edgeR v3.28.1(Robinson,

McCarthy et al., 2010). The distribution of all samples was then visualized after reducing the

dimension of the data by applying the Uniform Manifold Approximation and Projection for

Dimension Reduction (UMAP) technique using R package umap v0.2.6.0 (Leland, John et al.,

2018). The reduced dimensions of the data were plotted in 2D space using R package ggplot2

v3.3.2 (Wickham, 2009). Differential gene expression analysis was performed using raw read

counts of remaining samples using the R/Bioconductor package DESeq2 v1.26.0 (Love, Huber et

al., 2014). Genes with adjusted p-values of less than 0.05 were considered as significantly

regulated. Functional analysis of the significantly regulated genes was performed using the

enrichr module of the python package GSEAPY v 0.9.16 (https://pypi.org/project/gseapy/)

(Kuleshov, Jones et al., 2016, Subramanian, Tamayo et al., 2005). The total number of protein-

coding genes as background and KEGG gene-set library downloaded from Enrichr web resources

were used for the enrichment test for molecular pathway analysis. Pathways having adjusted p-

values of less than 0.05 were regarded as significant. Gene ontology (GO) enrichment analysis

was performed using the Enrichr web resource (Kuleshov et al., 2016), and the GO biological

process 2018 category was selected. GO terms with adjusted p-values of less than 0.05 were

considered as significant. Redundant GO terms were removed using the online tool REVIGO

(Supek et al., 2011) using the following settings: allowed similarity 0.5, GO database *Homo*

sapiens, and semantic similarity measure SimRel. Heatmaps were generated using the

R/Bioconductor package ComplexHeatmap v2.2.0 (Gu, Eils et al., 2016). Bubble plots, MA plots,

Volcanoplots, violin plots and bar plots were created using R package ggplot2 v3.3.2. Network

visualization was performed using Cytoscape v3.6.1 (Shannon, Markiel et al., 2003). Venn

diagrams were constructed using the online tool InteractiVenn (Heberle, Meirelles et al., 2015).

Cells and viruses:

The CCHFV strain IbAr10200 (originally isolated from *Hyalomma excavatum* ticks from Sokoto,

Nigeria, in 1966) was used in this study. The SW13- ATCC®-CCL-105[TM] and human hepatocyte-

derived cellular carcinoma cell line Huh7 was obtained from Marburg Virology Laboratory,

(Philipps-Universität Marburg, Marburg, Germany) and matched the STR reference profile of

Huh7(Rohde, Becker et al., 2019). CCHFV was propagated and titered in SW13 cells. The virus

was serially diluted 10-fold and 100ul of each dilution was added to SW13 cells in a 96 well

plate. At 48 h post-infection (hpi), cells were fixed in ice-cold acetone-methanol (1:1) and stained
using a rabbit polyclonal anti-CCHFV nucleocapsid antibody followed by a fluorescein
isothiocyanate (FITC)-conjugated anti-rabbit antibody (Thermo Fisher Scientific, US) and DAPI
(Roche, US). The number of fluorescent foci in each well was counted using a fluorescence
microscope and the titer was determined. Human 293 FLP-IN TRex cells expressing CAT
or ISG20 under the control of a Tet-on promoter were kindly provided by Ju-Tao Guo (Drexel
University College of Medicine, Pennsylvania , USA).(Jiang, Weidner et al., 2010) The cells were
propagated in DMEM supplemented with 5% Tet-negative FCS (PAA Lab, US).

***In vitro* infection assay in Huh7 cells and tandem mass tag (TMTpro) labelled reverse phase**
liquid chromatography mass-spectrometric (RPLC-MS/MS) analysis:

Huh7 cells were infected with the CCHFV in triplicate, as described by us previously (Appelberg
et al., 2020). Briefly, Huh7 cells were infected with CCHFV IbAr10200 at a multiplicity of
infection (MOI) of 1. After 1 h of incubation (37°C, 5% CO_2) the inoculum was removed, the
cells were washed with PBS, and 2 ml DMEM supplemented with 5% heat-inactivated FBS was
added to each well. Samples were collected in triplicate at 24 and 48 hpi along with controls. The
medium was removed, cells were washed carefully with PBS and lyzed in lysis buffer, (10 mM
Tris, 150 mM NaCl, 10% SDS, and protease inhibitor). After that, 4x sample buffer
(ThermoFisher, US) was added and the samples were boiled at 99°C for 10 min. Protein digestion
was performed in S-Trap microcolumns (Protifi, Huntington, NY) and the resulting peptides were
labeled with TMTpro tags. Labeled peptides were fractionated by high pH (HpH) reversed-phase
chromatography, and each fraction was analyzed on an Ultimate 3000 UHPLC (Thermo
Scientific, San Jose, CA) in a 120 min linear gradient. Data were acquired on a Q Exactive HF-X
mass spectrometer (Thermo Scientific, San Jose, CA) in data-dependent acquisition (DDA) mode
collecting full mass spectra at 120,000 resolution in a mass range of 350 – 1400 *m/z* for a
maximum injection time (IT) of 50 ms. Tandem mass spectra of the top 20 precursors were
acquired with high collision energy (HCD) of 34%, resolution of 45,000 and maximum IT of 86
ms, isolating precursors with 0.7 *m/z* and applying 45s dynamic exclusion. Proteins were searched
against the SwissProt human database using the search engine Mascot v2.5.1 (MatrixScience Ltd,
UK) in Proteome Discoverer v2.4 (Thermo Scientific) software allowing up to two missed
cleavages. The oxidation of methionine, deamidation of asparagine and glutamine, TMTpro at
lysine and N-termini were set as variable modifications; while carbamidomethylation of cysteine
was set as a fixed modification. The false discovery rate (FDR) was set to 1%.

Proteomics data analysis:

The raw data were first filtered to remove missing data. Proteins detected in all samples were
retained for analysis resulting in 8,501 proteins in the filtered dataset. The filtered data was then
normalized by applying eight different methods using R/Bioconductor package NormalyzerDE
v1.4.0 (Willforss, Chawade et al., 2019). The quantile normalization was found superior to other
methods and was selected for further use. Differential protein expression analysis was performed
using R/Bioconductor package limma v3.42.2 (Ritchie, Phipson et al., 2015). Proteins with
adjusted p-values of less than 0.05 were regarded as significant. KEGG pathway enrichment
analysis of significantly regulated proteins was performed using the enrichr module of python
package GSEAPY v 0.9.16.

ISG20 inhibition assay:

Tet-inducible HEK-293 FLP-IN TRex cells containing a CAT or ISG20 expression cassette were
seeded on in six-well dishes and treated with doxycycline at a final concentration of 2 μg/ml and
incubated for 36 h. Cells were infected with CCHFV strain IbAr10200 at an MOI of 1, and
supernatants were titrated at 48 hpi by enumeration of fluorescent foci and calculation of progeny
virus titers as described by us previously(Andersson et al., 2006).

RNAscope targeting IFI27 (ISG12) and CCHF:

The *RNAscope*® ISH Assays (ACD Bioscience,US) targeting IFI27 (440111, ACD Bioscience,
US) and CCHFV (510621, ACD Bioscience, US) were perfomed as described
previously.(Zhang, Svensson Akusjarvi et al., 2018) SW13 cells were infected with CCHFV (Ibar
10200 strain) at MOI of 0.1. After 1h, inoculum was replenished with fresh Leitbovitz medium
containing 5% FBS and incubated 48h. After infection cells were fixed 30 min using ice-cold
acetone. Cells were permeabilized using 0.1% Triton x-100 for 10 min at RT prior probe
incubation at 40°C of IFI27 and CCHFV probes, Amp-1, Amp-2, Amp-3 and Amp-4 Alt C for
2h, 30min, 15 min, 30 min and 15 min, respectively. Cells were further stained using DAPI and
coverslip attached using ProLong Gold Antifade reagent (P10144, Thermofisher). Images were
acquired using Nikon Single Point scanning confocal with 60/1.4 oil objective.

Acknowledgments: The authors would like to acknowledge support from the National Genomics
Infrastructure (NGI), Science for Life Laboratory, for RNAseq and Proteomics Biomedicum; and
Karolinska Institute, Solna, for LC-MS/MS analysis. The computations were performed using
resources provided by SNIC through the Uppsala Multidisciplinary Center for Advanced
Computational Science (UPPMAX) under Project SNIC2017-550. The microscopy part of the
study was performed at the Live Cell Imaging Facility, Karolinska Institute, Sweden, supported

by grants from the Knut and Alice Wallenberg Foundation, the Swedish Research Council, the Centre for Innovative Medicine, and the Jonasson Center at the Royal Institute of Technology, Sweden. The study is funded by Swedish Research Council Grants 2018-05766 and 2017-03126 to AM and 2017-01330 to UN, PHE Grant In Aid 109509 (inc. PhD studentship[EK]) and EU-H2020 CCHFVaccine to RH.

Author contributions: Conceptualization: U.N. and A.M., Methodology: U.N. N.E. A.M., A.V., F.W., and R.H., Formal analysis: A.T.A., S.A.,E.K., S.D., B.B., S.G., J.E.R, V.M.M. and S.S., Resources: U.N., A.M., F.W., and R.H., Writing (original draft): U.N., A.T.A, and S.G., Writing (review and editing): N.E., S.A., A.V., F.W., R.H., and A.M., Visualization: U.N., A.T.A, and S.S., Supervision: U.N., A.M., A.V., N.E., F.W., R.H., Project administration: U.N., N.E. and A.M., Funding acquisition: U.N., A.M., F.W., and R.H. All authors discussed the results, commented and approved the final version of the manuscript.

Conflict of interests: The authors declare no competing interests.

The Paper Explained: The pathogenesis and host-viral interactions of the Crimean–Congo hemorrhagic fever virus (CCHFV). We have performed a global transcriptomic analysis of peripheral blood mononuclear cells from a longitudinal cohort of CCHF patients who survived and temporal untargeted proteomics analysis of CCHFV infected cells. Our results indicate that during the acute phase of CCHFV infection, the host's metabolic reprogramming towards central carbon metabolism including glycolysis/gluconeogenesis occurs. This could be regulated by the anti-viral signaling pathways that play a central role in viral replication. Therefore targeting type I interferon response through metabolic perturbation could be an attractive therapeutic intervention for CCHFV as well as aid as antiviral strategies.

References

Akaberi D, Krambrich J, Ling J, Luni C, Hedenstierna G, Järhult JD, Lennerstrand J, Lundkvist Å (2020) Mitigation of the replication of SARS-CoV-2 by nitric oxide in vitro. Redox biology 37: 101734

Andersson I, Bladh L, Mousavi-Jazi M, Magnusson KE, Lundkvist A, Haller O, Mirazimi A
(2004) Human MxA protein inhibits the replication of Crimean-Congo hemorrhagic fever virus.
Journal of virology 78: 4323-9
Andersson I, Karlberg H, Mousavi-Jazi M, Martínez-Sobrido L, Weber F, Mirazimi A (2008)
Crimean-Congo hemorrhagic fever virus delays activation of the innate immune response. Journal
of medical virology 80: 1397-404
Andersson I, Lundkvist A, Haller O, Mirazimi A (2006) Type I interferon inhibits Crimean-
Congo hemorrhagic fever virus in human target cells. Journal of medical virology 78: 216-22
Appelberg S, Gupta S, Svensson Akusjarvi S, Ambikan AT, Mikaeloff F, Saccon E, Vegvari A,
Benfeitas R, Sperk M, Stahlberg M, Krishnan S, Singh K, Penninger JM, Mirazimi A, Neogi U
(2020) Dysregulation in Akt/mTOR/HIF-1 signaling identified by proteo-transcriptomics of
SARS-CoV-2 infected cells. Emerging microbes & infections 9: 1748-1760
Bakir M, Engin A, Gozel MG, Elaldi N, Kilickap S, Cinar Z (2012) A new perspective to
determine the severity of cases with Crimean-Congo hemorrhagic fever. Journal of vector borne
diseases 49: 105-10
Bente DA, Alimonti JB, Shieh WJ, Camus G, Ströher U, Zaki S, Jones SM (2010) Pathogenesis
and immune response of Crimean-Congo hemorrhagic fever virus in a STAT-1 knockout mouse
model. Journal of virology 84: 11089-100
Bente DA, Forrester NL, Watts DM, McAuley AJ, Whitehouse CA, Bray M (2013) Crimean-
Congo hemorrhagic fever: history, epidemiology, pathogenesis, clinical syndrome and genetic
diversity. Antiviral research 100: 159-89
Dobin A, Davis CA, Schlesinger F, Drenkow J, Zaleski C, Jha S, Batut P, Chaisson M, Gingeras
TR (2013) STAR: ultrafast universal RNA-seq aligner. Bioinformatics (Oxford, England) 29: 15-
21
Dowall SD, Graham VA, Fletcher T, Hewson R (2019) Use and reliability if multiplex bead-
based assays (Luminex) at Containment Level 4. Methods (San Diego, Calif) 158: 17-21
ECDC (2020) Hyalomma marginatum - Factsheet for experts, European Centre for Disease
Prevention and Control. In
Eisenreich W, Rudel T, Heesemann J, Goebel W (2019) How Viral and Intracellular Bacterial
Pathogens Reprogram the Metabolism of Host Cells to Allow Their Intracellular Replication.
Frontiers in cellular and infection microbiology 9: 42
Engin A, Arslan S, Kizildag S, Oztürk H, Elaldi N, Dökmetas I, Bakir M (2010) Toll-like
receptor 8 and 9 polymorphisms in Crimean-Congo hemorrhagic fever. Microbes and infection
12: 1071-8
Ergonul O, Tuncbilek S, Baykam N, Celikbas A, Dokuzoguz B (2006) Evaluation of serum levels
of interleukin (IL)-6, IL-10, and tumor necrosis factor-alpha in patients with Crimean-Congo
hemorrhagic fever. The Journal of infectious diseases 193: 941-4
Estrada-Peña A, Sánchez N, Estrada-Sánchez A (2012) An assessment of the distribution and
spread of the tick Hyalomma marginatum in the western Palearctic under different climate
scenarios. Vector borne and zoonotic diseases (Larchmont, NY) 12: 758-68
Feng J, Wickenhagen A, Turnbull ML, Rezelj VV, Kreher F, Tilston-Lunel NL, Slack GS,
Brennan B, Koudriakova E, Shaw AE, Rihn SJ, Rice CM, Bieniasz PD, Elliott RM, Shi X,
Wilson SJ (2018) Interferon-Stimulated Gene (ISG)-Expression Screening Reveals the Specific
Antibunyaviral Activity of ISG20. Journal of virology 92
Fontaine KA, Sanchez EL, Camarda R, Lagunoff M (2015) Dengue virus induces and requires
glycolysis for optimal replication. Journal of virology 89: 2358-66
Grandi G, Chitimia-Dobler L, Choklikitumnuey P, Strube C, Springer A, Albihn A, Jaenson TGT,
Omazic A (2020) First records of adult Hyalomma marginatum and H. rufipes ticks (Acari:
Ixodidae) in Sweden. Ticks and tick-borne diseases 11: 101403

Gu Z, Eils R, Schlesner M (2016) Complex heatmaps reveal patterns and correlations in
multidimensional genomic data. Bioinformatics (Oxford, England) 32: 2847-9
Heberle H, Meirelles GV, da Silva FR, Telles GP, Minghim R (2015) InteractiVenn: a web-based
tool for the analysis of sets through Venn diagrams. BMC bioinformatics 16: 169
Hülya Y, Yilmaz G, Kostakoglu U, Arslan M, Karahan SC, Koksal I (2017) Evaluation of Serum
TNF-alpha, IL-6, IL-10, and IFN-gamma Levels in Patients with Crimean–Congo Hemorrhagic
Fever. Open Forum Infectious Diseases 4: S312-S313
Jiang D, Weidner JM, Qing M, Pan XB, Guo H, Xu C, Zhang X, Birk A, Chang J, Shi PY, Block
TM, Guo JT (2010) Identification of five interferon-induced cellular proteins that inhibit west nile
virus and dengue virus infections. Journal of virology 84: 8332-41
Kaya S, Elaldi N, Kubar A, Gursoy N, Yilmaz M, Karakus G, Gunes T, Polat Z, Gozel MG,
Engin A, Dokmetas I, Bakir M, Yilmaz N, Sencan M (2014) Sequential determination of serum
viral titers, virus-specific IgG antibodies, and TNF-α, IL-6, IL-10, and IFN-γ levels in patients
with Crimean-Congo hemorrhagic fever. BMC infectious diseases 14: 416
Korva M, Rus KR, Pavletič M, Saksida A, Knap N, Jelovšek M, Smrdel KS, Jakupi X, Humolli I,
Dedushaj J, Petrovec M, Avšič-Županc T (2019) Characterization of Biomarker Levels in
Crimean-Congo Hemorrhagic Fever and Hantavirus Fever with Renal Syndrome. Viruses 11
Kuleshov MV, Jones MR, Rouillard AD, Fernandez NF, Duan Q, Wang Z, Koplev S, Jenkins SL,
Jagodnik KM, Lachmann A, McDermott MG, Monteiro CD, Gundersen GW, Ma'ayan A (2016)
Enrichr: a comprehensive gene set enrichment analysis web server 2016 update. Nucleic acids
research 44: W90-7
Lei CQ, Zhang Y, Xia T, Jiang LQ, Zhong B, Shu HB (2013) FoxO1 negatively regulates cellular
antiviral response by promoting degradation of IRF3. The Journal of biological chemistry 288:
12596-604
Leland M, John H, Nathaniel S, Lukas G (2018) UMAP: uniform manifold approximation and
projection. Journal of Open Source Software 3: 861
Liao Y, Smyth GK, Shi W (2014) featureCounts: an efficient general purpose program for
assigning sequence reads to genomic features. Bioinformatics (Oxford, England) 30: 923-30
Love MI, Huber W, Anders S (2014) Moderated estimation of fold change and dispersion for
RNA-seq data with DESeq2. Genome biology 15: 550
Lucas TM, Richner JM, Diamond MS (2015) The Interferon-Stimulated Gene Ifi27l2a Restricts
West Nile Virus Infection and Pathogenesis in a Cell-Type- and Region-Specific Manner. Journal
of virology 90: 2600-15
Lundell LS, Massart J, Altıntaş A, Krook A, Zierath JR (2019) Regulation of glucose uptake and
inflammation markers by FOXO1 and FOXO3 in skeletal muscle. Molecular metabolism 20: 79-
88
Mankouri J, Tedbury PR, Gretton S, Hughes ME, Griffin SD, Dallas ML, Green KA, Hardie DG,
Peers C, Harris M (2010) Enhanced hepatitis C virus genome replication and lipid accumulation
mediated by inhibition of AMP-activated protein kinase. Proceedings of the National Academy of
Sciences of the United States of America 107: 11549-54
Martin M (2011) Cutadapt removes adapter sequences from high-throughput sequencing reads.
EMBnet journal 17: pp. 10-12
Mayer KA, Stöckl J, Zlabinger GJ, Gualdoni GA (2019) Hijacking the Supplies: Metabolism as a
Novel Facet of Virus-Host Interaction. Frontiers in immunology 10: 1533
McGinley L, Hansford KM, Cull B, Gillingham EL, Carter DP, Chamberlain JF, Hernandez-
Triana LM, Phipps LP, Medlock JM (2020) First report of human exposure to Hyalomma
marginatum in England: Further evidence of a Hyalomma moulting event in north-western
Europe? Ticks and tick-borne diseases 12: 101541
Min Z, Ye Z, Gang L, Boyu D, Xueyan X (2020) IFI27 as a potential indicator for severe
Enterovirus 71-infected hand foot and mouth disease. Virus research 289: 198149

Moser TS, Schieffer D, Cherry S (2012) AMP-activated kinase restricts Rift Valley fever virus infection by inhibiting fatty acid synthesis. PLoS pathogens 8: e1002661

Papa A, Bino S, Velo E, Harxhi A, Kota M, Antoniadis A (2006) Cytokine levels in Crimean-Congo hemorrhagic fever. Journal of clinical virology : the official publication of the Pan American Society for Clinical Virology 36: 272-6

Papa A, Tsergouli K, Tsioka K, Mirazimi A (2017) Crimean-Congo Hemorrhagic Fever: Tick-Host-Virus Interactions. Frontiers in cellular and infection microbiology 7: 213

Popova TG, Turell MJ, Espina V, Kehn-Hall K, Kidd J, Narayanan A, Liotta L, Petricoin EF, 3rd, Kashanchi F, Bailey C, Popov SG (2010) Reverse-phase phosphoproteome analysis of signaling pathways induced by Rift valley fever virus in human small airway epithelial cells. PloS one 5: e13805

Puigserver P, Rhee J, Donovan J, Walkey CJ, Yoon JC, Oriente F, Kitamura Y, Altomonte J, Dong H, Accili D, Spiegelman BM (2003) Insulin-regulated hepatic gluconeogenesis through FOXO1-PGC-1alpha interaction. Nature 423: 550-5

Ranadheera C, Valcourt EJ, Warner BM, Poliquin G, Rosenke K, Frost K, Tierney K, Saturday G, Miao J, Westover JB, Gowen BB, Booth S, Feldmann H, Wang Z, Safronetz D (2020) Characterization of a novel STAT 2 knock-out hamster model of Crimean-Congo hemorrhagic fever virus pathogenesis. Sci Rep 10: 12378

Ritchie ME, Phipson B, Wu D, Hu Y, Law CW, Shi W, Smyth GK (2015) limma powers differential expression analyses for RNA-sequencing and microarray studies. Nucleic acids research 43: e47

Robinson MD, McCarthy DJ, Smyth GK (2010) edgeR: a Bioconductor package for differential expression analysis of digital gene expression data. Bioinformatics (Oxford, England) 26: 139-40

Rohde C, Becker S, Krahling V (2019) Marburg virus regulates the IRE1/XBP1-dependent unfolded protein response to ensure efficient viral replication. Emerging microbes & infections 8: 1300-1313

Saksida A, Duh D, Wraber B, Dedushaj I, Ahmeti S, Avsic-Zupanc T (2010) Interacting roles of immune mechanisms and viral load in the pathogenesis of crimean-congo hemorrhagic fever. Clinical and vaccine immunology : CVI 17: 1086-93

Saline M, Badertscher L, Wolter M, Lau R, Gunnarsson A, Jacso T, Norris T, Ottmann C, Snijder A (2019) AMPK and AKT protein kinases hierarchically phosphorylate the N-terminus of the FOXO1 transcription factor, modulating interactions with 14-3-3 proteins. The Journal of biological chemistry 294: 13106-13116

Sandau KB, Faus HG, Brüne B (2000) Induction of hypoxia-inducible-factor 1 by nitric oxide is mediated via the PI 3K pathway. Biochemical and biophysical research communications 278: 263-7

Scholte FEM, Zivcec M, Dzimianski JV, Deaton MK, Spengler JR, Welch SR, Nichol ST, Pegan SD, Spiropoulou CF, Bergeron É (2017) Crimean-Congo Hemorrhagic Fever Virus Suppresses Innate Immune Responses via a Ubiquitin and ISG15 Specific Protease. Cell Rep 20: 2396-2407

Shannon P, Markiel A, Ozier O, Baliga NS, Wang JT, Ramage D, Amin N, Schwikowski B, Ideker T (2003) Cytoscape: a software environment for integrated models of biomolecular interaction networks. Genome research 13: 2498-504

Shore AM, White PC, Hui RC, Essafi A, Lam EW, Rowe M, Brennan P (2006) Epstein-Barr virus represses the FoxO1 transcription factor through latent membrane protein 1 and latent membrane protein 2A. Journal of virology 80: 11191-9

Simon M, Falk KI, Lundkvist A, Mirazimi A (2006) Exogenous nitric oxide inhibits Crimean Congo hemorrhagic fever virus. Virus research 120: 184-90

Smallwood HS, Duan S, Morfouace M, Rezinciuc S, Shulkin BL, Shelat A, Zink EE, Milasta S, Bajracharya R, Oluwaseum AJ, Roussel MF, Green DR, Pasa-Tolic L, Thomas PG (2017)

Targeting Metabolic Reprogramming by Influenza Infection for Therapeutic Intervention. Cell
Rep 19: 1640-1653
Spengler JR, Patel JR, Chakrabarti AK, Zivcec M, García-Sastre A, Spiropoulou CF, Bergeron É
(2015) RIG-I Mediates an Antiviral Response to Crimean-Congo Hemorrhagic Fever Virus.
Journal of virology 89: 10219-29
Subramanian A, Tamayo P, Mootha VK, Mukherjee S, Ebert BL, Gillette MA, Paulovich A,
Pomeroy SL, Golub TR, Lander ES, Mesirov JP (2005) Gene set enrichment analysis: a
knowledge-based approach for interpreting genome-wide expression profiles. Proceedings of the
National Academy of Sciences of the United States of America 102: 15545-50
Supek F, Bošnjak M, Škunca N, Šmuc T (2011) REVIGO summarizes and visualizes long lists of
gene ontology terms. PloS one 6: e21800
Tang BM, Shojaei M, Parnell GP, Huang S, Nalos M, Teoh S, O'Connor K, Schibeci S, Phu AL,
Kumar A, Ho J, Meyers AFA, Keynan Y, Ball T, Pisipati A, Kumar A, Moore E, Eisen D, Lai K,
Gillett M et al. (2017) A novel immune biomarker IFI27 discriminates between influenza and
bacteria in patients with suspected respiratory infection. The European respiratory journal 49
Thaker SK, Ch'ng J, Christofk HR (2019) Viral hijacking of cellular metabolism. BMC biology
17: 59
Tsai WL, Chang TH, Sun WC, Chan HH, Wu CC, Hsu PI, Cheng JS, Yu ML (2017) Metformin
activates type I interferon signaling against HCV via activation of adenosine monophosphate-
activated protein kinase. Oncotarget 8: 91928-91937
Weber F, Mirazimi A (2008) Interferon and cytokine responses to Crimean Congo hemorrhagic
fever virus; an emerging and neglected viral zonoosis. Cytokine & growth factor reviews 19: 395-
404
WHO (2020) Fact Sheet: Crimean Congo Haemorrhagic Fever. In World Health Organization
Regional Office for Africa
Wickham H (2009) ggplot2: Elegant Graphics for Data Analysis. Springer International
Publishing, New York
Willforss J, Chawade A, Levander F (2019) NormalyzerDE: Online Tool for Improved
Normalization of Omics Expression Data and High-Sensitivity Differential Expression Analysis.
Journal of proteome research 18: 732-740
Zhang W, Ambikan AT, Sperk M, van Domselaar R, Nowak P, Noyan K, Russom A, Sonnerborg
A, Neogi U (2018) Transcriptomics and Targeted Proteomics Analysis to Gain Insights Into the
Immune-control Mechanisms of HIV-1 Infected Elite Controllers. EBioMedicine 27: 40-50
Zhang W, Svensson Akusjarvi S, Sonnerborg A, Neogi U (2018) Characterization of Inducible
Transcription and Translation-Competent HIV-1 Using the RNAscope ISH Technology at a
Single-Cell Resolution. Frontiers in microbiology 9: 2358
Zivcec M, Safronetz D, Scott D, Robertson S, Ebihara H, Feldmann H (2013) Lethal Crimean-
Congo hemorrhagic fever virus infection in interferon α/β receptor knockout mice is associated
with high viral loads, proinflammatory responses, and coagulopathy. The Journal of infectious
diseases 207: 1909-21
Zivcec M, Scholte FE, Spiropoulou CF, Spengler JR, Bergeron É (2016) Molecular Insights into
Crimean-Congo Hemorrhagic Fever Virus. Viruses 8: 106

Figures

Fig. 1. Study design and severity group association with gene expression and cytokine profiling. (a) Overall work-flow of the data generation and analysis. **(b)** Sample distribution during the acute phase of infection in different severity groups as reported. **(c)** MA plot of differentially regulated genes during the acute phase between samples of severity group 1 and severity group 2 or 3. **(d)** Violin plot of significantly changed proteins as determined from ELISA assays.

Fig. 2. Differential gene expression and pathway analysis between acute and recovery phases. (a) Heatmap of Z-score transformed expression values of significantly regulated genes in the pair-wise comparisons namely Recovered vs Acute (overall), Recovered vs Acute (severity group 1), Recovered vs Acute (severity group 2). The columns represent the patient samples and their corresponding severity groups at different time points. The rows represent genes that are hierarchically clustered based on Euclidean distance. **(b)** KEGG pathway enrichment analysis results of genes belonging to cluster 1 of the heatmap. The bubble size and colour gradient correspond to the number of genes and adjusted P value of the enrichment test, respectively. **(c)** KEGG pathway enrichment results of genes belonging to cluster 2 of the heatmap. The bubble size and color gradient correspond to the number of genes and adjusted P value of the enrichment test, respectively. **(d)** Venn diagram of significantly up-regulated genes in Recovered vs Acute (severity group 1) and Recovered vs Acute (severity group 2) phases **(e)** Venn diagram of significantly down-regulated genes in Recovered vs Acute (severity group 1) and Recovered vs Acute (severity group 2) phases. **(f)** Gene ontology (biological process) enrichment analysis results of commonly regulated genes (882 upregulated and 569 down-regulated) from (d) and (e). the colour gradient and bubble size correspond to the gene ratio of each GO term and the adjusted P value of the enrichment test, respectively. The adjacent bar graph represents the percentage of genes upregulated or downregulated in each gene ontology term.

Fig. 3. Differentially expressed genes in interferon (IFN) signaling pathways. (a) Heatmap visualizing the expression pattern of IFN-signaling genes (including ISGs) that are significantly different between the recovered and acute phases. The columns represent the patient samples and

their corresponding severity groups at different time points. The rows represent genes hierarchically clustered based on Euclidean distance. The row annotation indicates the differential expression of genes (DEG) and presence or absence of genes in the selected GO terms. The names of genes involved in the cellular response to IFN-γ (GO:0071346) and the IFN-Isignaling (GO:0060337) are printed. **(b)** MA plot of differentially regulated genes between the recovered and acute phases. ISGs are marked. **(c)** RNAscope analysis targeting IFI27 genes in infected and non-infected cells. **(d)** Network type visualization of genes belonging to GO:0071346 and GO:0060337. The edges indicate the presence of the corresponding gene in the gene-ontology term. The node size and colour gradient corrospond to the adjusted P value of differential expression analysis and the log2 fold change, respectively.

Fig. 4. LC-MS/MS based proteomics analysis in CCHFV infected Huh7 cells. (a) General work-flow showing the sample preparation and data generation for the proteomics experiment. **(b)** Significant pathways identified by mapping proteins regulated over time to the KEGG database. The pathways are grouped into two: the first 29 pathways are metabolism related and the last 16 pathways are related to organismal systems and environmental information processing. The bubble size and colour gradient correspond to the number of proteins and adjusted P value of the enrichment test, respectively. The bar graphs represent the percentage of proteins upregulated or downregulated in each pathway as per the corresponding comparisons.

Fig. 5. Temporal dynamics of interferon stimulating genes (ISGs). (a) Heatmap of Z-score transformed expression values of proteins belonging to the cellular response to IHN-γ (GO:0071346) and the IFN-I interferon signaling pathway (GO:0060337) gene ontology terms. The columns represent the different sampls and their corresponding treatment conditions and time points. The rows represent proteins hierarchically clustered based on Euclidean distance. The row annotation indicates the presence or absence of proteins in the corresponding GO terms and log 2-fold change values in the proteomics and transcriptomics (recovered -vs- acute) experiments. **(b)** Volcano plot of ISGs visualizing the expression status of 24 h Mock and Infected (24 hpi) samples. The size and colour gradients of the dots correspond to the adjusted P values of differential expression analysis and the log2 fold change, respectively. **(c)** Volcano plot of ISGs visualizing the expression status of 48 h Mock and Infected (48 hpi). The size and colour gradient of the dots correspond to the adjusted P values of differential expression analysis and log2 fold change. respectively. **(d)** Volcano plot of ISGs visualizing the expression status of 24 hpi and 48 hpi. The size and colour gradient of the dots correspond to the adjusted P value of differential expression analysis and the log2 fold change, respectively. **(e)** HEK293-FLP-IN T-

Rex cells that conditionally express ISG20 or human CAT protein were treated with doxycycline or left untreated for 36 h, followed by infection with CCHFV at an MOI 1 for 48hours. Cell-free culture supernatants were harvested and progeny viral titers were determined by fluorescent focus-forming assay. Columns show means ± the standard deviations ($n = 2$). A two-tailed paired Student *t* test was performed to compare titers of untreated and doxycycline-treated cells.

Fig. 6. Potential host-cellular, immune response and metabolic reprogramming in CCHFV infection. The figure is created with BioRender.com.

Supplementary Materials

Supplementary Table S1. The CCHF patient characteristics.

Supplementary Figure S1: The heterogenous gene-expression profile was further refined by hierarchical clustering analysis of the top 500 high variable genes

Supplementary Figure S2. The levels of all of the rest 18 soluble marker.

Supplementary Figure S3: UMAP clustering of the caute phase and recovered phase samples.

Supplementary Figure S4: Overlap of the pathways identified between patients based transcriptomics data and in vitro cellular data.

Supplementary figure S5. Potential pathways regulated by CCHFV during acute phase of infection

Supplementary Data File 1: The DGE profile for the acute phase compared to the recovered phase in all patients showed an upregulation of 2891 genes and a downregulation of 2738

Supplementary Data File 2: The pathways that are significantly modulated by the upregulated and downregulated genes.

a

Severity group 1, 2, and 3

Follow-up
Median (range)
54 (46-57) Week

Death (n=1)
Low quality samples (n=5)

Acute Phase
(n=18)

After Recovery
(n=12)

PBMC
isolation

RNA
isolation

Illumina
NovaSeq6000

Bioinformatics
Analysis

b.

UMAP 2

○ Severity group 1
● Severity group 2
● Severity group 3

UMAP 1

c.

Log2 fold change

AHSP

PROM1   HBD
TIMP4   CA1
CMBL CTSG
            ERG   MT1G
      SMIM10   VSIG4
          PPARG   RNASE1

Log2 mean expression

d.

GM-CSF
p=0.049

TNF-α
p=0.055

IL-8
p=0.04

IL-10
p=0.055

log2(pg/ml)

☐ Severity score 1   ☐ Severity score 2

a.

Group
- Acute
- Recovered

SeverityScore
- Severity group 1
- Severity group 2

Comparisions
- Recovered vs Acute (Overall)
- Recovered vs Acute (Severity group 1)
- Recovered vs Acute (Severity group 2)

Z-Score
4
2
0
-2
-4

LFC
4
2
0
-2
-4

b.
- N–Glycan biosynthesis
- Citrate cycle (TCA cycle)
- Thermogenesis
- Oxidative phosphorylation
- Terpenoid backbone biosynthesis
- Amino sugar and nucleotide sugar metabolism
- One carbon pool by folate
- Valine, leucine and isoleucine degradation
- Propanoate metabolism
- Glyoxylate and dicarboxylate metabolism
- Pyruvate metabolism
- Pyrimidine metabolism
- Alanine, aspartate and glutamate metabolism
- Glycolysis / Gluconeogenesis
- Glycine, serine and threonine metabolism
- NOD–like receptor signaling pathway
- Arginine and proline metabolism

Genes
20 30 40 50 60

Adj.Pvalue
3e-10 3e-05 1e-06 1e-04 1e-02

c.
- Hematopoietic cell lineage
- Th17 cell differentiation
- Th1 and Th2 cell differentiation
- MAPK signaling pathway
- Cell adhesion molecules (CAMs)
- Antigen processing and presentation
- T cell receptor signaling pathway
- Ras signaling pathway
- Natural killer cell mediated cytotoxicity
- Intestinal immune network for IgA production
- Cytokine–cytokine receptor interaction
- Parathyroid hormone synthesis, secretion and action
- NF-kappa B signaling pathway
- FoxO signaling pathway
- Leukocyte transendothelial migration
- Fc gamma R–mediated phagocytosis
- Phospholipase D signaling pathway
- Rap1 signaling pathway
- HIF–1 signaling pathway
- Chemokine signaling pathway
- Phosphatidylinositol signaling system
- Notch signaling pathway
- PI3K–Akt signaling pathway
- Sphingolipid signaling pathway

Genes
20 30 40 50 60 70

Adj.Pvalue
3e-10 3e-05 1e-05 1e-04 1e-02

d.
Severity group 1 (954)   Severity group 2 (2182)
72   882   1300

e.
Severity group 1 (663)   Severity group 2 (2074)
94   569   1505

f.
- Type I interferon signaling pathway
- Regulation of viral genome replication
- DNA replication
- G1/S transition of mitotic cell cycle
- Protein N–linked glycosylation
- DNA damage response, signal transduction by p53 class mediator
- Antigen processing and presentation of exogenous peptide antigen
- DNA metabolic process
- Cellular response to DNA damage stimulus
- Cellular macromolecule biosynthetic process

% Genes
0 10 20 30

Gene-ratio(%)
20 28 32

-log10(FDR)
4 8 12

- Up-regulated
- Down-regulated