The endosymbiotic origins of the apicoplast link fever-survival and artemisinin-

resistance in the malaria parasite

Min Zhang[a,1], Chengqi Wang[a,1], Jenna Oberstaller[a,1], Phaedra Thomas[a], Thomas D.

Otto[b,c], Debora Casandra[a], Sandhya Boyapalle[a], Swamy R. Adapa[a], Shulin Xu[a], Katrina

Button-Simons[d], Matthew Mayho[b], Julian C. Rayner[b,e], Michael T. Ferdig[e], Rays H. Y.

Jiang[a], John H. Adams[a,2]

[a]Center for Global Health and Infectious Diseases Research and USF Genomics

Program, College of Public Health, University of South Florida, 3720 Spectrum Blvd,

Suite 404, Tampa, Florida 33612, USA

[b]Wellcome Sanger Institute, Wellcome Genome Campus, Hinxton Cambridgeshire,

CB10 1SA United Kingdom

[c]Institute of Infection, Immunity and Inflammation, MVLS, University of Glasgow,

Glasgow G12 8TA United Kingdom

[d]Eck Institute for Global Health, Department of Biological Sciences, University of Notre

Dame, Notre Dame, IN 46556

[e]Cambridge Institute for Medical Research, University of Cambridge, Cambridge

Biomedical Campus, The Keith Peters Building, Hills Road, Cambridge, Cambridgeshire,

CB2 0XY United Kingdom

[1] M.Z., C.W. and J.O. contributed equally to this work.

[2]To whom correspondence should be addressed. Email: ja2@usf.edu

ABSTRACT

Background: The emergence and spread of *Plasmodium falciparum* parasites resistant to front-line antimalarial artemisinin-combination therapies (ACT) threatens to erase the considerable gains against the disease of the last decade. We developed a new large-scale phenotypic screening pipeline and used it to carry out the first large-scale forward-genetic phenotype screen in *P. falciparum* to identify genes that allow parasites to survive febrile temperatures.

Results: Screening identified more than 200 *P. falciparum* mutants with differential responses to increased temperature. These mutants were more likely to be sensitive to artemisinin derivatives as well as to heightened oxidative stress. Major processes critical for *P. falciparum* tolerance to febrile temperatures and artemisinin included highly essential, conserved pathways associated with protein-folding, heat-shock and proteasome-mediated degradation, and unexpectedly, isoprenoid biosynthesis, which originated from the parasite's algal endosymbiont-derived plastid, the apicoplast. Apicoplast-targeted genes in general were up-regulated in response to heat shock, as were other *Plasmodium* genes with orthologs in plant and algal genomes.

Conclusions*: Plasmodium falciparum* parasites appear to exploit their innate febrile-response mechanisms to mediate resistance to artemisinin. Both responses depend on endosymbiotic cyanobacterium-related ancestral genes in the parasite's genome, suggesting a link to the evolutionary origins of *Plasmodium* parasites in free-living ancestors.

Running title: Plastid metabolism enables malaria parasites to survive fever and artemisinin

INTRODUCTION

Malaria remains a leading infectious disease causing >200 million clinical cases and a

half-million deaths every year. *Plasmodium falciparum* is the deadliest malaria parasite

by far, with growing parasite resistance to front-line antimalarial artemisinin-combination

therapies (ACT) threatening to erase the considerable gains against the disease of the

last decade. Alarmingly, data indicate that for the first time since 2010, progress in

reducing global burden of malaria cases and fatalities nearly flatlined between 2015 and

2017 [1]. New therapies, ideally informed by an understanding of basic parasite biology,

are needed to confront these urgent threats to global malaria control. The study of

malaria-parasite biology and gene-function has traditionally been limited, because

targeted gene-by-gene approaches are laborious and fraught with difficulty due to an

AT-rich (~82%) genome that limits scalability of specific targeted gene-editing methods

(such as CRISPR). Despite the considerable knowledge gene-by-gene studies have

enabled, and the ~two decades that have passed since the *P. falciparum* genome was

completed [2], the limited throughput of targeted gene-editing strategies combined with

evolutionary distance of *P. falciparum* from classical model eukaryotes has left >90% of

genes untouched experimentally, and ~35% of the parasite's ~5474 genes without

meaningful functional annotation (www.plasmodb.org) [3]. High-throughput methods for

functionally profiling the malaria-parasite genome can hasten development of effective

interventions to control a parasite proven to be an adaptable foe.

Parasite-specific processes essential for parasite survival are naturally attractive as

potential drug-targets, given the decreased likelihood of deleterious off-target effects to

the host. One such process ripe for interrogation is the parasite's survival-response to

the extreme conditions of the host's malarial fever. Repeating fever is a hallmark of all

types of malaria and the cyclical patterns serve as key diagnostic features of infections.

In malignant tertian malaria caused by *P. falciparum,* the 48-hour cycle corresponds to

the parasite's asexual intraerythrocytic-stage life-cycle, wherein parasites invade,

develop, asexually replicate and then rupture their host red blood cell (RBC) to begin the

destructive blood-stage cycle anew. Host fever is triggered by a Type I shock-like

response of the innate immune-system exposure to extracellular parasite debris

released when infected RBCs are lysed during parasite egress. Malarial fever

concomitantly attenuates and synchronizes development of blood-stage *P. falciparum*

infections, as it is lethal to all parasite stages except for early intraerythrocytic ring

stages. However, parasite tolerance of febrile temperatures is crucial for its successful

propagation in human populations as well as a fundamental aspect of malaria

pathogenesis. Previous research suggests parasite-specific factors play a role in

modulating this tolerance for febrile temperatures, though the identities of many of these

factors or the mechanisms by which they operate remain uncertain [4, 5].

We previously used random *piggyBac*-transposon insertional mutagenesis to uncover

genes essential for *P. falciparum* blood-stage survival, generating a saturation-level *P.*

falciparum mutant library containing ~38,000 single-disruption mutants [6]. We defined

2680 genes as essential for asexual blood-stage growth, including ~1000 *Plasmodium*-

conserved genes of unknown function. Here we demonstrate the potential of this

piggyBac-mutant (*pB*-mutant) library to systematically assign functional annotation to the

P. falciparum genome by genome-wide phenotypic screens. In this study, we present the

first large-scale forward-genetic functional screen in *P. falciparum* to identify factors

linked to parasite survival of febrile temperatures. Importantly, we functionally annotate

hundreds of parasite genes as critical for the parasite's response to heat shock (HS) but

dispensable under ideal growth-conditions, ~26% of which were previously unannotated

with no known function. Expression-profiling the HS-responses in two different heat

shock-sensitive (HS-Sensitive) *pB*-mutant clones vs. the wildtype parent NF54 via

RNAseq revealed concordance between (1) genes regulated in the parasite's innate

response to HS, (2) the processes dysregulated in these mutants vs. wildtype responses

to HS, and (3) those mutants we identified as HS-Sensitive in our pooled screens.

Together these analyses identify genes and pathways essential in the HS-response,

implicating oxidative stress and protein-damage responses, host-cell remodelling, and

unexpectedly, apicoplast isoprenoid biosynthesis. Apicoplast-associated genes in

general were up-regulated in response to HS, as were other *Plasmodium* genes with

orthologs in plant and algal genomes. Finally, parallel phenotyping of a mutant library

revealed a significant overlap between parasite pathways underlying the response to

febrile temperatures and those implicated in the artemisinin mechanism of action (MOA),

including oxidative stress, protein-damage responses, and apicoplast-mediated vesicular

trafficking [7, 8]. Mutants in known protein-targets of artemisinin tended to be sensitive to

HS [9], and expression-data from recent field-isolates directly correlates artemisinin-

resistance with HS tolerance in our pooled screen [10]. Further, we found the K13-

associated parasite endocytosis pathway responsible linked to artemisinins resistance

[11, 12] is also downregulated in response to HS. Together these data identify an

unexpected link between artemisinin MOA, HS-survival, and algal origins of the

apicoplast, suggesting the parasite exploits its innate fever-response mechanisms to

gain resistance to artemisinin. This study creates a blueprint for developing a large-scale

phenotypic screening pipeline of the *P. falciparum pB*-mutant library to enable high-

throughput interrogation of phenotypes of interest to hasten further biological insight that

can be weaponized against the parasite.

RESULTS

Pooled screens of an extensively characterized pB-mutant clone-library allow

robust identification of heat-shock phenotypes

To interrogate pathways and processes associated with parasite survival at febrile

temperatures, we developed a large-scale phenotypic screening pipeline to analyze the

phenotypes in pooled *pB*-mutant parasites exposed to HS-induced stress (Fig. S1). We

previously demonstrated using individual clonal *pB*-mutant parasite lines that mutant

growth-phenotypes can be detected and differentiated in pooled screening utilizing

QIseq—"Sensitive" mutants with disruptions in genes/genomic features important for

growth have lower QIseq reads, while "Neutral" disruptions in features not vital for

growth under the same conditions have higher reads [13]. We therefore reasoned that

mutants with mutations in genes underlying the HS-response would grow poorly in

response to HS compared to mutants in genes not contributing to HS-survival.

We used a pool of 128 unique, extensively characterized *P. falciparum pB*-mutant clones

reflecting disruptions in genes spanning a range of functional categories, as well as

many genes without existing functional information, as a "pilot library" for initial

phenotypic screen-development ([13, 14]; Methods). An *in vitro* HS-screen of this pilot-

library, adapted from a phenotype- screen of many *pB*-mutant-clones comprising the

pilot-library [15], defined *pB*-mutant HS-response phenotypes to fever-like temperatures

(Fig. 1A-E, Table S1, Methods). We next calculated a measure of fitness for each

mutant in response to HS while also taking into account inherent differences in mutant-

growth in ideal conditions, which we termed the Phenotypic-Fitness Score in response to

HS (PFS_{HS}; Methods). We classified 28 mutants of the pilot-library as HS-Sensitive (Fig.

1E-F, indicated in red; Table S1). Fourteen mutants performed poorly in both the

Growth- and HS-Screens (Fig. 1E-F, yellow). We classified 28 mutants displaying a

slight growth advantage in response to HS (Fig. 1E-F, green) as "HS-Tolerant". Mutants

exhibiting neither sensitivity nor tolerance to HS were classified as HS-Neutral (n = 49).

QIseq-data resulting from the HS- and Growth-screens allowed robust assignment of

mutant-phenotypes for both (see Methods). We primarily classified mutants sensitive to

heat-shock alone as HS-Sensitive to avoid possible over-interpretation of generally-sick

Growth-Sensitive mutants (Fig. 1E-G).

Pooled phenotypic screens scaled up to a 1K *pB*-mutant library enable

identification of processes driving the *P. falciparum* heat-shock response

We next scaled our pooled HS-screen to a mutant library of 922 functionally

uncharacterized mutants using the methods we established in our pilot-library screens.

This 1K-library comprised mutant-pools randomly selected from our saturation library,

covering genes annotated to diverse GO-categories, as well as many genes of unknown

function. Mutants were ranked by fold-change growth in response to HS from HS-

Sensitive to HS-Tolerant, as per cut-offs determined from our pilot-library screens. Our

analysis distinguished 149 mutants growing well in ideal growth conditions but poorly in

response to HS as HS-Sensitive (FC-HS < 0.5 and PFS_{HS} < 0.25; Fig. 2A, red box),

while 91 mutants performed poorly in both the Growth- and HS-screens (FC-HS < 0.5

and PFS_{HS} > 0.25; yellow). Of the remaining mutants, 139 HS-Tolerant mutants had

slightly better growth in HS than ideal growth-conditions (FC-HS > 1.5; green box), while

543 classified as HS-Neutral were neither sensitive nor tolerant (taupe).

This larger scale of screening allowed us to assess gene functional-enrichment in HS-

Sensitive and Growth-Sensitive phenotypic categories vs all other mutants in the ~1K-

library. HS-Sensitive mutants were enriched in GO terms associated with HS-response

such as protein-folding, response to DNA-damage, DNA-repair, and regulation of

vesicle-mediated transport, broadly in agreement with processes identified to underlie

the HS-response by more conventional gene expression-based methods [4, 5]. Growth-

Sensitive mutants tended to be enriched for more general categories broadly important

for survival in all conditions, such as translation- or mRNA-metabolism-related terms

(Fig. 2B), as might be expected given the high essentiality of these processes in ideal

growth [6, 16].

Increased transcription of the unfolded protein response (UPR), organelle-

targeted stress-response pathways and host-cell remodeling characterize the

parasite HS-response

We characterized the wildtype parent-NF54 transcriptome in response to HS to establish

a baseline for comparison using an experimental design similar to a prior study to

assessing transcriptional changes in response to febrile temperatures via microarray [5].

The HS assay-design mimicking parasite exposure to malarial fever was modelled after

conditions we established for our pooled-screens (Methods). RNAseq was performed on

heat-shocked parasites vs. a non-heat-shocked control. Genes identified as differentially

expressed in response to febrile temperatures vs. 37°C were classified into three

different categories based on direction of response: (1) upregulated in response to HS;

(2) downregulated in HS, and (3) neutral in HS (Fig. 3A-B, Table S3A). The majority of

genes expressed above threshold in our analysis were HS-neutral (1541 genes out of

2567, or ~60%) and were enriched for genes involved in general housekeeping functions

such as the proteasome core complex (ubiquitin-proteasome system), the ubiquitin-

dependent ERAD-pathway, and regulators thereof), RNA metabolism (RNA-binding,

mRNA-splicing) and transport functions (e.g. protein import into nucleus, vesicle-

mediated transport). We primarily considered genes upregulated in HS as drivers of the

HS-response.

Genes upregulated in HS (n = 415) tended to be enriched for processes such as protein-

folding, unfolded protein-binding, response to heat, mitochondrial processes, and host-

cell remodelling-associated exported proteins localizing to the Maurer's clefts (Fig. 3B,

Table S3C-D). Genes downregulated in HS (n = 611) tended to be enriched for

pathogenesis-related functions and components of the parasite invasion machinery,

such as entry/exit from the host cell and cell-cell adhesion, and organelles including the

inner-membrane pellicle complex, micronemes, and rhoptries. These data are in general

agreement with previously-reported processes expected to drive the parasite HS-

response [4, 5].

The unfolded protein response, organelle-targeted pathways are dysregulated in

HS-Sensitive mutants

We reasoned that genes dysregulated in HS-Sensitive mutants compared to wildtype

underlie the HS-response. We chose two individual HS-Sensitive mutant clones for

additional profiling via RNAseq to identify dysregulated genes responsible for this

sensitivity. Both mutant lines have a single disruption in the coding region of a gene not

previously implicated in the HS-response: ΔDHC (dynein heavy-chain gene

PF3D7_1122900), and ΔLRR5 (leucine-rich repeat protein PF3D7_1432400).

The 1298 genes which could be classified into HS-response categories across all three

parasites were analyzed for functional-enrichment (Table S3B). The majority of genes

were HS-neutral across all three parasites and were enriched for essential

housekeeping functions (n = 615; Table S3B-D). We reasoned these non-HS-regulated

genes have functions too important for basic survival to tolerate large stress-associated

expression-changes, and these genes were not considered drivers of the HS-response.

We identified 91 genes significantly upregulated in HS across all three parasites, which

were functionally enriched for protein-folding, chaperone-related processes, and other

processes related to heat-stress and the UPR, in agreement with previous expression-

based studies [5], as well as enrichment-results from HS-Sensitive mutants in our pooled

screening, indicating the parasite increases production of heat-shock proteins (HSPs)

and associated chaperones to repair the glut of proteins damaged/misfolded by heat-

stress (Table S3B-D). Energy-producing processes (gluconeogenesis, glycolysis) were

also upregulated, suggesting the parasite reroutes anabolic metabolism to increase

energy production to support ATP-dependent processes such as protein-refolding to

correct heat-damaged proteins. Genes upregulated in HS were further enriched for

processes involved in host-cell remodeling, including genes targeted to the Maurer's

clefts, the host cell, and intracellular vesicles—all known to be important for parasite-

remodeling of the host-cell to promote structural reinforcement against heat-shock

damage to ensure its own survival [4, 5]. Organellar targeting to the mitochondria and

apicoplast are also enriched in upregulated HS-responsive genes. The parasite's

increased utilization of mitochondrial stress-response pathways may aid in degrading

heat-damaged proteins that cannot be correctly refolded. Increased activity in the food

digestive vacuole may allow the parasite to phagocytose and eliminate toxic misfolded

protein-aggregates. The apicoplast involvement, particularly the isoprenoid biosynthesis

pathway, has not been previously implicated in the HS-response.

Genes downregulated in all three parasites in response to HS (n = 205) were enriched

for virulence-factor and invasion-machinery-associated GO terms, suggesting the

parasite decreases production of transcripts associated with pathogenesis, invasion and

egress, lengthening its intracellular recovery-time to address global protein-damage.

Both HS-Sensitive mutants share many characteristic features of the wildtype response

to febrile temperatures, which likely enabled their survival (Fig. 3A-B, red, blue; Table

S3C-D). We identified two primary expression categories of genes dysregulated in the

HS-Senstive mutants: (1) genes upregulated in the wildtype HS-response that were

otherwise dysregulated in the HS-Sensitive mutants, which we interpreted as loss-of-

function changes, and (2) genes that were not regulated in response to HS in the

wildtype but were upregulated in the HS-Sensitive mutants, presumably equivalent to

dominant-negative gain-of-function changes (Fig. 3A-B, ochre and tan, respectively).

This first category of mutant-dysregulated genes was enriched for the UPR, as well as

mitochondrial and apicoplast-localized pathways (cytochrome oxidase-assembly and

fatty-acid biosynthesis, respectively). Several apicoplast isoprenoid biosynthesis-related

genes upregulated in the wildtype HS-response were additionally dysregulated in one or

both HS-Sensitive *pB*-mutant clones (Fig. 3C). The second category of mutant-

dysregulated genes, those that are not HS-responsive in wildtype, were enriched for

translation-associated processes.

These data taken together suggest underlying mechanisms responsible for the HS-

response. Critically, HS-Sensitive mutants fail to upregulate mitochondrial and apicoplast

stress-response pathways, as well as signal peptide-processing pathways that might

enable appropriate activation of those pathways. Mutants do not increase production of

transcripts associated with responding to unfolded proteins. HS-Sensitive mutants

additionally upregulate translation-related processes in response to HS when translation

should be paused or neutral. This increase may overwhelm the parasite's capacity to

repair or degrade heat-damaged proteins, exacerbating the formation/accumulation of

toxic misfolded-protein aggregates that increase parasite sensitivity to HS.

Apicoplast isoprenoid biosynthesis is critical for *P. falciparum* survival of febrile

temperatures

We examined our RNAseq data more closely to discern contributions of the apicoplast to

HS-survival. We found that apicoplast-targeted genes tended to be increased in

response to HS as compared to all non-apicoplast-targeted genes (Fig. 4A), were more

likely to be essential during ideal blood-stage growth conditions (Fig. 4B), and were

enriched for stress-response processes such as the UPR and oxidative-stress, and less

expectedly, isoprenoid biosynthesis (Fig. 4C). As a major function of isoprenoid

biosynthesis is in protein-prenylation—an important post-translational modification that

regulates protein-targeting and function throughout the cell—we hypothesized that

mutants in known-prenylated proteins [17, 18] would also have a phenotype in HS. We

examined our 1K mutant-library for representation of isoprenoid biosynthesis, its

immediate upstream-regulators (proteins responsible for modulation and import of

glycolytic intermediates that serve as pathway substrates), and immediate downstream-

effector proteins, and found that all eight isoprenoid biosynthesis-related *pB*-mutants

included in the pooled screen were indeed HS-Sensitive (Fig. 4D).

Based on these data we further hypothesized that proteins or pathways allowing *P.*

falciparum survival of febrile temperatures would be absent or otherwise divergent in

Plasmodium species whose hosts do not mount fever-responses. We therefore

compared the apicoplast isoprenoid biosynthesis pathway between *P. falciparum* and

two rodent-infective species, *P. berghei* and *P. yoelii*. We found key thiamine-synthesis

enzymes directly upstream of the pathway missing in the rodent-infective malaria

parasites, including hydroxy-ethylthiazole kinase (ThzK); ThzK is up-regulated in the

canonical parasite response to febrile temperatures and dysregulated in HS-Sensitive

mutants (Fig. 4E). Perhaps most importantly, DOXP-Synthase (DXS), the critical

enzyme marking the first step in isoprenoid biosynthesis, is upregulated in HS,

dysregulated in HS-Sensitive mutants, and was HS-Sensitive in pooled screening, as

were all four members of the prenylated blood-stage proteome represented in our

screen (Fig. 4E). These data taken together strongly implicate isoprenoid biosynthesis in

the HS-response.

Though the apicoplast has not previously been implicated in parasite survival of febrile

temperatures, there is extensive literature on the ability of plants to mount effective

defenses against heat as well as other external stressors, particularly critical for non-

motile organisms at the mercy of their environments. We investigated the relationship

between the parasite's HS-response and "plant-like" stress-responses by evaluating

phyletic distribution of parasite HS-response genes in representative plant and algal

genomes. *P. falciparum* genes with plant orthologs indicating potential endosymbiont-

ancestry tended to be increased in response to HS vs. genes that do not have plant

orthologs (Fig. 4F). These lines of evidence considered together present an evolutionary

explanation that endosymbiosis of the apicoplast's algal progenitor enabled parasite-

survival of extreme temperatures.

Processes enabling parasites to survive fever also drive resistance to artemisinin

We noted similarities between processes we identified to be driving the parasite HS-

response and those implicated in parasite-resistance to artemisinin [7, 8, 10]. Therefore,

we did a series of parallel phenotype- screens of our *pB*-mutant pilot-library using

sublethal concentrations of two artemisinin compounds (dihydroartemisinin, DHA;

artesunate, AS), heightened conditions of oxidative stress of RBCs, and exposure to a

proteasome inhibitor (Bortezomib; BTZ) to investigate the possible relationship between

HS-response and artemisinin MOA. HS-Sensitive mutants tended to be sensitive to both

artemisinin derivatives and H_2O_2-induced oxidative stress, while HS-Tolerant mutants

were less sensitive to either condition (Fig. 5A, Table S4). Also, HS-Sensitive mutants

shared an increased sensitivity to the proteasome inhibitor BTZ, consistent with

laboratory observations connecting artemisinin MOA to the proteasome and clinical data

that proteasome-inhibitors act synergistically with artemisinins [8, 19-21]. Overall,

correlation of mutant phenotypic profiles across screens varied, with 16-45% having

correlating phenotypes in at least one additional screen (Fig. 5B, Fig S2A-B).

We next assessed whether these laboratory-based experimental findings corresponded

to 'real world' changes associated with *P. falciparum* in artemisinin-resistant (ART-R)

clinical isolates [10]. Consistent with our laboratory findings linking HS- sensitivity and

ART- sensitivity, the genes associated with the HS-Sensitive mutants were significantly

correlated with genes linked to ART-R in recent field isolates (Fig. 5C, Table S4). We

also found clear GO relationships between HS phenotype-categories and ART-R in field-

isolates, with genes driving the HS-response most positively correlated with reduced

parasite clearance-rates (Fig. 5D, Table S4).

Artemisinin is activated by degradation of host hemoglobin. Recent evidence has

suggested two key, temporally-distinct ART-R mechanisms: (1) a multi-functional protein

long associated with resistance in field-isolates, *kelch13* (K13) confers resistance

upstream of hemoglobin degradation by modulating an associated endocytosis pathway;

and (2) downstream of hemoglobin degradation through the ubiquitin-proteasome

system (UPS), where K13 may function as or regulate a ubiquitin ligase [10-12, 22-25].

In upstream-resistance, endocytotic transport of hemoglobin to the digestive vacuole

(DV) is down-regulated as this is the key process through which the parasite ingests,

degrades, and then releases hemoglobin. K13 mutant-isolates appear to downregulate

processes along this endocytosis pathway, decreasing parasite hemoglobin digestion

and release of heme to activate artemisinin, thereby increasing parasite survival (Fig.

S2C). We found that K13-defined endocytosis is also downregulated in response to HS

(Fig. 5E). As the K13-mediated endocytosis pathway culminates in host haemoglobin-

cargo being degraded in the DV, we further assessed our 1k HS-screen for DV-

associated proteins. We found DV-associated proteins did tend to be sensitive to heat-

shock, including key DV resident-proteases (Plasmepsin I, M1-family alanyl

aminopeptidase; Fig. S3A) [26]. We next evaluated our 1K HS-sScreen for direct K13-

interacting partner-proteins recently identified via immunoprecipitation [25], and found

that mutants in 10 of the 24 unique putative K13-partner-proteins represented in the

screen were sensitive to HS. Further, 5 of 7 known alkylation-targets of artemisinin

represented in our screen had sensitivity to HS [9, 26] (Fig. S3B). We noted significant

overlap in each of these categories of ART MOA-related genes and isoprenoid

biosynthesis-related genes (Fig. S3C).

In a second downstream step post-activation of artemisinin, the parasite engages the

UPS to further mitigate artemisinin-induced damage. Artemisinins mount a multi-pronged

attack against the parasite by causing a global, non-specific accumulation of damaged

parasite proteins, which are then polyubiquitinated/marked for degradation, while also

inhibiting proteasome-function. These poly-ubiquitinated proteins ultimately overwhelm

the parasite's decreased capacity for UPS-mediated protein-degradation [8]. Key

ubiquitinating components of this system, including E2/E3 ligases and K13, are

downregulated in response to HS, while key components of the UPR and protein folding

are increased (Fig. 5E). In contrast, components of the core proteasome were

universally increased in response to HS when considered in aggregate, although the

change did not meet our fold-change criteria for being HS-regulated (Fig. S4).

Synthesizing these data, we present a model for the relationship between DHA MOA

described recently [8] and HS-response (Fig. 5F). The canonical parasite-response to

fever is to increase protein-folding and UPR while inhibiting ubiquitination to prevent

accumulation of toxic, polyubiquitinated protein-aggregates. The parasite simultaneously

increases its capacity for proteasome-mediated degradation—ultimately enabling it to

resolve HS-instigated stress and thus survive febrile temperatures. Artemisinins kill by

overwhelming these same pathways: damaging and unfolding proteins, preventing

folding of newly synthesized proteins and inhibiting the proteasome, while at the same

time activating ubiquitination-machinery to ensure the accumulation of toxic

polyubiquitinated proteins that eventually cause cell-death. ART-R-associated mutations

allow the parasite to constitutively activate unfolded-protein response mechanisms which

increase its capacity for refolding or degrading those toxic proteins [27]. The overall

increase in damaged-protein degradation-capacity allows ART-R parasites to keep up

with the influx of artemisinin-induced protein-damage, clearing the waste and enabling

parasite survival. This direct inverse relationship in activation of endocytosis, the

ubiquitin-proteasome system and other pathways underlying DHA-mediated killing and

febrile-temperature survival, supports a shared mechanism for artemisinin-resistance

and HS-response, suggesting that ART-R parasites evolved to harness canonical HS-

survival mechanisms to survive artemisinin.

Discussion

Our data indicate that the parasite crisis-response to HS is multi-faceted to relieve the

build-up of heat-damaged proteins before it is overwhelmed by toxic, misfolded-protein

aggregates. Responding to or perhaps preventing a build-up of potentially toxic heat-

damaged proteins, the parasite upregulates expression of chaperones to stabilize and

detoxify them, downregulating ubiquitinating enzymes to discourage their aggregation

while upregulating the core proteasome and vesicular trafficking to degrade and

eliminate proteins which can't be repaired. Equally important in the survival-response are

changes in redox homeostasis, lipid metabolism, cellular transport, and metabolic

processes associated with the endosymbiont-derived organelles. The parasite requires

increased energy to mount this febrile response, which it provides by redirecting its own

internal biosynthetic pathways to produce glucose. Interestingly, we confirm the

parasite's protective response-mechanisms include proteins exported into the

erythrocyte, suggesting that the parasite's metabolic processes exported to remodelled

cytoplasm of the parasitized host cell are equally vulnerable and vital to malaria parasite

survival.

A high proportion of essential genes that are upregulated in response to heat stress are

targeted to the apicoplast. The apicoplast isoprenoid biosynthesis pathway's critical

involvement in survival of febrile temperatures is nevertheless a surprise, as it has not

been implicated before in the *Plasmodium* HS-response. Isoprenoids are required for

myriad functions across the tree of life—plant chloroplasts, algae, some parasitic-

protozoa and bacterial pathogens utilize a specialized form of this pathway absent from

all metazoans (called the non-mevalonate, MEP, or DOXP pathway), which has made

isoprenoid biosynthesis an attractive target for intervention against a range of pathogens

[28, 29]. Most studied organisms make wide use of protein-prenylation and have large

prenylated proteomes; malaria parasites, in contrast, have a very small prenylated

blood-stage proteome (~20 proteins) consisting primarily of vesicular trafficking proteins,

notably the Rab-family GTPases [17, 18]. Recent studies indicate the key essential

function of isoprenoids in the parasite blood-stage is in their roles as substrate for

protein-prenylation—specifically, in prenylating proteins driving vesicular transport to the

digestive vacuole [30, 31]. In the absence of prenylation, Rab5 trafficking is disrupted,

which leads to digestive vacuole-destabilization and parasite death [31]. Notably,

artemisinin also disrupts digestive vacuole-morphology, resulting in a very similar

phenotype as a consequence of its activation via hemoglobin digestion [32, 33].

Intriguingly, recent data confirm the association of key resistance-mediator K13 with

Rab-GTPases [25], adding to the repertoire of proteins comprising K13-mediated

endocytic vesicles, and by extension supporting the role of prenylation in K13-mediated

processes associated with ART MOA.

Another key parasite-defense against oxidative stress induced by pro-oxidant

compounds (such as artemisinin) includes increased vitamin E biosynthesis—another

exclusive function of the MEP isoprenoid-biosynthesis pathway, whose stress-related

regulation has been extensively studied in plants [34, 35]. Further insights to the role

isoprenoids play in the HS-response may be gleaned from plants and pathogenic

bacteria, where research suggests key branchpoint-enzyme DXS, which catalyzes the

first and rate-determining step of the MEP pathway [36], has a role in sensing and then

facilitating adaptation to ever-changing environmental conditions, including temperature,

light-exposure, chemical compounds, and oxidative stress (for example [37, 38]).

Elevated levels of isoprenoids have been found to correlate with plant exposure to

drought and other stressors and are considered a key component of plant-defenses

against abiotic stress [39]. The DXS ortholog may play a similar role in *P. falciparum*,

enabling the parasite to mount quick responses to unfavorable conditions in the host-

environment, such as fever.

Interestingly, concurrent studies now provide mechanistic insights illuminating the

biochemical relationship between apicoplast isoprenoid biosynthesis and the parasite

febrile-temperature survival response (co-submission by Mathews et al.). Farnesylation

of HSP40 (PF3D7_1437900), a type of prenylation mediated by the MEP pathway, is

critical for *P. falciparum* survival of thermal stress. In this study Inhibition of isoprenoid

biosynthesis ultimately resulted in reduced association of HSP40 with critical

components of the cytoskeleton, protein-export, and vesicular transport pathways—

without which *P. falciparum* could survive neither heat nor cold stress. Suppression of

these cellular processes by loss of HSP40-farnesylation directly corresponds to HS-

sensitive pathways identified via both our forward-genetic screen and our gene-

expression analyses of the HS-Sensitive LRR5- and DHC4-mutant clones.

Few eukaryotes are known to be able to thrive in extreme-heat environments; most are

unable to complete their lifecycles above 40°C [40]. The survival mechanism of malaria

parasites could be attributed to the algal ancestral lineage of the apicoplast. Some

extant red algal-lineages (genus *Cyanidioschyzon*) are extremophilic inhabitants of

acidic hot-springs and are remarkably resistant to heat shock up to 63°C; green-algae

Chlamydamonas reinhardtii was also able to survive to 42°C [41]. Responsibility for this

extreme resistance to transient exposure to high temperatures was attributed to two

genes of the small heat shock protein (sHSP) family (CMJ100C and CMJ101C). The *P.*

falciparum ortholog for these genes (PF3D7_1304500) was upregulated in the wildtype

HS-response and dysregulated in both our HS-Sensitive mutants, indicating its

contribution to parasite survival in extreme temperatures. Mutations in this gene were

not represented in our pooled screens.

It is tempting to speculate that acquisition of the red-algal endosymbiont and its

associated plant stress-response mechanisms is what enabled the ancestral parasite to

survive host-fever, likely an important and early step leading to successful infection of

hominid hosts. Our findings of significant overlap between parasite-responses to three

disparate stressors (HS, artemisinin, oxidative stress) offers new insight into how *P.*

falciparum exhibited artemisinin-resistance even in the initial clinical trials [42], and then

further evolved resistance relatively quickly after mass-introduction of the drug by

"hijacking" and repurposing the parasite's in-built fever-response pathways.

Conclusion

Deeper knowledge of parasite biology is expected to enable more effective and likely

longer-lasting antimalarial interventions. Similarly, a better mechanistic understanding of

artemisinin MOA will lead to better combination therapies to combat emerging

resistance. With this first large-scale forward-genetic screen in *P. falciparum*, we

revealed the parasite's survival responses to malarial fever and artemisinin

chemotherapy share common underpinnings that heavily depend on metabolic

processes of plant origin.

ART-R ultimately hinges on highly efficient protein-degradation mechanisms. This

mechanistic knowledge allows for the application of intelligently considered counters to

ART-R, such as combinatorial therapy with proteasome-inhibitors, which has

experimentally shown great promise [43]. Our current study highlights the potential of

forward-genetic screens to elucidate unexpected processes and pathways, such as

DOXP and isoprenoid biosynthesis, that are associated with the artemisinin MOA which

may serve as synergistic druggable targets. Future studies can exploit a genome-wide

screening approach to iteratively ascribe function to every part of the malaria-parasite

genome to support targeted development of new, more-efficacious antimalarial

combination therapies to limit and potentially reverse artemisinin resistance.

Declarations

Ethics approval and consent to participate

Not applicable.

Consent for publication

Not applicable.

Availability of data and materials

The raw RNAseq dataset supporting the conclusions of this article are available in the

Mendeley Data repository *Malaria-parasite survival of host fever is linked to artemisinin*

resistance, http://dx.doi.org/10.17632/b8g3wbnd5v.1 [44]. Raw QIseq dataset accession

numbers are listed in Table S5.

Competing interests

The authors declare that they have no competing interests.

Funding

This work was supported by the National Institutes of Health grant R01 AI094973 and

R01 AI117017 (J.H.A.) and the Wellcome Trust grant 098051 (J.C.R.).

Authors' contributions

Conceptualization, M.Z., C.W., J.O., T.D.O., J.C.R., M.T.F., R.H.Y.J., and J.H.A.;

Methodology, M.Z., C.W., J.O., K.B.S., R.H.Y.J., and J.H.A.; Software, J.O. and C.W.;

Validation, M.Z., C.W., and J.O.; Formal Analysis, M.Z., C.W., J.O., S.R.A., and

R.H.Y.J.; Investigation, M.Z., C.W., J.O., P.T., D.C., S.B., S.X., M.M., and R.H.Y.J.; Data

Curation, M.Z., C.W., and J.O.; Writing—Original Draft, J.O., M.Z., and J.H.A.; Writing—

Review & Editing, J.O., M.Z., T.D.O., J.C.R. and J.H.A.; Visualization, C.W., M.Z., J.O.,

and R.H.Y.J.; Supervision, M.Z., J.O., T.D.O., J.C.R., M.T.F., R.H.Y.J., and J.H.A.;

Project Administration, M.Z.; Funding Acquisition, J.C.R. and J.H.A.

Acknowledgements

We appreciate the Wellcome Sanger Institute (United Kingdom) for performing QIseq

and Xiangyun Liao, Suzanne Li, and Kenneth Udenze for support of parasite cell culture.

We thank the USF Genomics Program Omics Hub for productive discussion.

FIGURE LEGENDS

Figure 1. Pooled screens of *P. falciparum piggyBac* mutants allow robust

identification of heat-shock phenotypes. A. Experimental design for pooled heat

shock (HS) phenotypic screens. The pilot library of *pB*-mutant clones (n=128) was

exposed to three rounds of temperature-cycling to simulate malarial fever (Methods). A

pilot-library control concurrently grown continuously at 37°C established inherent growth

of each *pB* mutant.

B. QIseq quantifies each *pB*-mutant in the pilot library from sequence-reads of the 5′ and

3′ ends of each *pB* insertion-site. Colored lines represent genes. Black boxes indicate

transposon location.

C. Pilot-library mutant growth-phenotypes at ideal temperatures, defined as fold change

in QIseq reads-count after three cycles growth at 37°C (FC-Growth; Methods) ranked

from Sensitive to Tolerant. Mutants with inherently slower or faster growth under ideal

conditions are shown in grey and blue, respectively.

D. Pilot-library mutant HS-phenotypes ordered from Sensitive (red) to Tolerant (green).

Mutant growth was defined as QIseq reads-count fold-change in response to HS (FC-

HS) vs. non heat-shocked control (Methods). HS-Sensitive mutants have lower FC-HS

(red, FC-HS < 1), while HS-Tolerant mutants have higher FC-HS (green, FC-HS > 1).

E-F. HS- and Growth-phenotypes of the pilot-library mutants. HS-phenotype of each

mutant (displayed as line-graph) is superimposed on its corresponding Growth-

phenotype (bar graph). Red = HS-Sensitive mutants (FC-HS < 0.5 and Phenotypic

Fitness Score in response to heat shock (PFS_{HS}) < 0.25, n = 28). Yellow = mutants

classified as both Growth-Sensitive and HS-Sensitive (FC-HS < 0.5, PFS_{HS} > 0.25, n =

14). Green = HS-Tolerant mutants (FC-HS > 1.5, n = 30). Mutants neither Sensitive nor

Tolerant to HS were classified as HS-Neutral (n = 49). *Known HS-Sensitive *pB*-mutant

clones validated by individual HS-assay [15]. **Known HS-Tolerant *pB*-mutant clones

validated by individual HS-assay [15].

G. Distributions of PFS_{HS} for mutant HS-phenotype classifications. HS-Sensitive mutants

are assigned the lowest PFS_{HS}, while HS-Tolerant mutations are assigned the highest

PFS_{HS} (**** Wilcoxon-test p-value < 1e-15) (Methods).

Figure 2. Large-scale pooled phenotypic screens enable identification of

processes driving the *P. falciparum* heat-shock response

A. HS-Sensitive mutations identified in pooled screens of 922 *pB*-mutants. Mutant pools

were screened in both ideal growth conditions and under HS and were assigned

phenotypes as per methods established in the pilot-library screens (Fig. 1). Mutants are

ranked by fold-change in response to HS (FC-HS; n = 922) from HS-Sensitive (red; n =

149) to HS-Tolerant (green; n = 139). Mean mutant fold-change in ideal growth (FC-

Growth) is superimposed as a bar plot (gray, FC-Growth < 1.0; blue, FC-Growth > 1.0).

Mutants performing poorly in both screens (yellow; n = 91) were classified as HS- and

Growth-Sensitive and were not considered further. Mutations neither HS-Sensitive nor

HS-Tolerant were classified as HS-Neutral. See Methods and Table S2.

B. Functional enrichment of GO terms for HS-Sensitive or Growth-Sensitive *pB*-mutants

vs all other mutants in the 1K-library. HS-Sensitive mutants were enriched in terms

associated with HS-response such as protein-folding, response to DNA-damage, DNA-

repair, and regulation of vesicle-mediated transport. Growth-Sensitive mutants tended to

be enriched for more general categories broadly important for survival in all conditions,

such as translation- or mRNA-metabolism-related terms. Circles represent GO category,

circle color represents ontology, and circle size represents number of significant genes

annotated to that category. Significant terms (Fisher/elim-hybrid test p. value <= 0.05)

fall within the light-green box.

Figure 3. Increased transcription of the unfolded protein response, organelle-

targeted stress-response pathways and host-cell remodeling characterize the

parasite HS-response. A. Genes were classified based on their NF54-expression with

and without HS-exposure across all three parasite lines (Methods). Genes were

classified as upregulated in response to HS (FC-HS > 1; ↑), down-regulated in response

to HS (FC-HS < -0.5; ↓), or not regulated by HS (-0.5 < FC-HS < 1; –), with upregulated

genes considered to be driving the HS-response. Genes were then assigned HS

expression-categories based on phenotype in NF54 vs. HS-Sensitive mutants *ΔLRR5*

and *ΔDHC*. HS-regulated genes shared between NF54 and both mutants are indicated

in red (↑↑↑) or blue (↓↓↓) for up- and down-regulated genes, respectively. Genes

dysregulated in one or both HS-Sensitive mutants fell into two main expression-profile

categories underlying mutant HS-Sensitivity phenotypes: those upregulated in NF54 that

failed to be regulated in the mutants (↑✕✕), and genes not regulated in response to HS

in NF54 that were inappropriately upregulated in the mutants (− ↑↑).

B. Functional enrichment analyses between wildtype/mutant HS-expression profiles as

defined in A. Red: Shared upregulated HS-responsive GO-terms between NF54 and the

two HS-Sensitive *pB*-mutants (↑↑↑). Blue: Shared down-regulated HS-responsive GO-

terms (↓↓↓). Ochre: GO-terms upregulated in NF54 but dysregulated in the two *pB*-

mutant (↑✕✕). Tan: GO-terms enriched in genes not regulated in the wildtype HS-

response but upregulated in the mutants (− ↑↑). Only enriched GO-terms are shown

(Fisher/elim-hybrid test p. value <= 0.05), with highest significance indicated in dark

green. Fraction of significant genes mapping to a GO-term in an HS expression-profile

category vs. genes mapping to that GO-term in the entire analysis is indicated by

distance to the center of the circle, with the outermost position on the circle indicating

100% of genes in that GO-term are significant. See Table S3D.

C. Several apicoplast and isoprenoid biosynthesis-related genes have a tendency to be

upregulated in the wildtype-response to HS and are dysregulated in one or both HS-

Sensitive *pB*-mutant clones (↑✕✕). * Isoprenoid biosynthesis-related genes upregulated

by HS confirmed in the pooled HS-Screen.

Figure 4. Apicoplast isoprenoid biosynthesis is critical for *P. falciparum* survival

of febrile temperatures. A. Apicoplast-targeted genes tend to be increased in response

to HS as compared to all non apicoplast-targeted genes detected above threshold in

RNAseq. Apicoplast-targeted genes are as defined in [45] (*** Fisher-test p-value < 1e-

5).

B. Apicoplast-targeted genes tend to be highly essential during blood-stage vs. all other

non-apicoplast-targeted genes detected above threshold in RNAseq. The median

Mutagenesis Index Score (MIS; [6]) for apicoplast-targeted genes is much lower than

median MIS for all other genes, indicating a lower tolerance for disruption and thus

higher likely essentiality during blood-stage development than non-apicoplast-targeted

genes (**** Wilcoxon-test p-value < 1e-15).

C. Apicoplast pathways regulated in response to HS. GO categories enriched in up- and

down-regulated apicoplast genes are shown on a scale from red to blue, respectively.

The horizontal direction indicates the log ratio between up- and down-regulated

apicoplast genes in each category. Circle-size represents gene-number per category.

D. All nine *pB*-mutants in genes related to apicoplast isoprenoid biosynthesis

represented in the 1K-library pooled screen were HS-Sensitive. Mutants are ranked by

phenotype from HS-Sensitive (red) to HS-Tolerant (green). Circles indicate each HS-

Sensitive mutant related to isoprenoid-biosynthesis. *The three isoprenoid biosynthesis-

genes we identified as directly upregulated in response to HS via RNAseq (DXS, tRNA

m(1)G methyltransferase and apicoplast RNA methyltransferase). See Table S3.

E. Key enzymes in the *P. falciparum* isoprenoid biosynthesis-pathway are up-regulated

in response to heat-shock (red circle), dysregulated in HS-Sensitive mutants (ochre) and

absent in malaria-parasites of hosts that do not present fever. Pathway diagram

modeled from [46]. Isoprenoid biosynthesis-genes upregulated in HS include DXS, 1-

deoxy-D-xylulose-5-phosphate synthase (DOXP), 2-C-methyl-D-erythritol 2,4-

cyclodiphosphate synthase (IspF), pyruvate kinase II (PyKII), triosephosphate isomerase

(TIM), triose phosphate transporter (TPT), and upstream-regulator of MEP-pathway

substrates HAD1-phosphotase [47]. All vesicular trafficking-proteins RAB GTP'ases,

direct downstream-targets prenylated by the MEP-pathway (zigzag) represented in

pooled screening were HS-Sensitive. The key thiamin-synthesis enzyme

hydroxyethylthiazole kinase (ThzK) is absent in *P. berghei* and *P. yoelii*, malaria-

parasites whose rodent-hosts do not present fever.

F. *P. falciparum* genes with plant orthologs (green circles) indicating potential

endosymbiont-ancestry tend to be increased in response to HS vs. genes that do not

have plant orthologs (grey circles). *P. falciparum* genes with potential endosymbiont-

ancestry were derived from 1919 ortholog-pairs between *Arabidopsis thaliana* and *P.*

falciparum (data from OrthoMCLv5.0). The listed processes are sorted based on the

ratio of "green" to "non-green" orthologs.

Figure 5. Increased sensitivity to fever is directly correlated with increased

sensitivity to artemisinin in the malaria parasite. A. HS-Sensitive *pB*-mutants (red)

are more sensitive to multiple concentrations of artemisinin derivatives Artesunate and

Dihydroartemisinin, proteasome-inhibitor Bortezomib (BTZ), and conditions of

heightened oxidative stress than HS-tolerant parasites (green) in all pooled screens of

the pilot library. *pB*-mutants were cultured continuously under oxidative stress-inducing

conditions for three to six cycles (T1 and T2, respectively). Apicoplast-mutants (n = 5)

have phenotypes similar to all HS-Sensitive mutants (n =28) in artemisinin-derivative

screens, but not to protein-inhibitors or oxidative stress (* Wilcoxon $p < 0.05$; ***

Wilcoxon $p < 1e-10$. See Methods).

B. Correlation between mutant phenotypes in all pooled screens of the pilot library.

Mutants performing in the bottom 25% or top 25% of each screen were classified as

having "Sensitive" and "Tolerant" phenotypes, respectively. Mutant classifications were

compared pair-wise between each screen, with mutants falling into the same category in

both screens considered to have correlating phenotypes.

C. HS-Sensitive *pB*-mutants (red) are positively correlated with genes linked to ART-R in

recent field isolates [10](* Wilcoxon test $p < 0.05$).

D. HS expression-categories as determined by comparative RNAseq between NF54 and

two HS-Sensitive mutant clones are positively correlated with ART-R in field-isolates.

E. Both K13-mediated mechanisms of artemisinin resistance (endocytosis, ubiquitin-

proteasome system) are similarly regulated in HS. The K13-defined endocytosis

pathway (shades of green) and key ubiquitinating-enzymes of the ubiquitin-proteasome

system, E2/E3 and K13, are downregulated in the wildtype NF54 HS-response, while

protein folding, stress, exported proteins, and proteasome genes are upregulated.

RNAseq data are plotted for each gene by average log2 fold-change in response to

HS and significance (–log2(p-value)). Circles in shades of blue and pink indicate genes

significantly down- or upregulated after exposure to HS, respectively.

F. Activation of pathways underlying DHA-mediated killing and febrile-temperature

survival is directly inverse. **Top**: Model of DHA-mediated killing in *P. falciparum* adapted

from [8]. Artemisinin (ART) damages and unfolds proteins, prevents folding of newly

synthesized proteins, and inhibits the proteasome while at the same time activating

E1/E2/E3 ubiquitin-machinery. Accumulation of toxic polyubiquitinated protein-substrates

(S) overwhelms the cell and leads to death. **Bottom**: Model of parasite fever-response.

Heat-stress causes globally damaged protein. The parasite increases the UPR as it

inhibits E2/E3 ubiquitination to prevent accumulation of toxic, polyubiquitinated (Ub)

protein-aggregates, while at the same time increasing its capacity for proteasome-

mediated degradation—ultimately enabling the parasite to resolve HS-instigated stress

and survive febrile-temperatures.

Methods

Pilot pB-mutant clone-library characteristics and validation

The single *piggyBac*-transposon insertion sites of each *pB*-mutant-clone in the pilot-

library were verified as previously described [13, 14]. Additionally, whole-genome

sequencing performed on 23% of 128 *pB*-mutant-clones in the pilot-library verified that

no major genomic changes occurred aside from the *piggyBac* insertion, ensuring any

detected phenotypes are attributable to the single disruption [15]. The pilot-library was

generated in a manner to ensure approximately equal representation of each of the 128

clones at thaw [13].

Generating the pilot-library of pB-*mutant parasite clones*

The pilot-library was built as described in our previous QIseq methods-development

study [13] and data are available in PlasmoDB (RRID:SCR_013331). Aliquots of the

pilot-library were generated by first growing each of the 128 extensively-characterized

mutant-clones individually in T25-flasks to 1-2% parasitemia. All clones were then

combined equally into one large flask and gently mixed. One-hundred equal-volume

aliquots of the pilot-library were then cryopreserved according to standard methods.

Pooled-screen assay-design

HS-screens

We exposed pools of *pB*-mutant parasites to three rounds of temperature-cycling to

simulate the cyclical pattern of fever characteristic of human malaria (Figure 1A).

Parasites under phenotypic selection (heat-shock) and ideal-growth controls originated

from the same thaw, grown at 37°C for one cycle then split equally into five flasks (three

flasks for exposure to heat-shock, two for the ideal-growth controls). Experimental and

control-flasks were maintained in parallel to minimize potential batch-effects. Parasites

were grown for one cycle at 37°C until they reached the ring-stage of development

(Time-point 0; T^0), at which point the experimental-group were exposed to febrile

temperatures (41°C) for 8 hours. Post-heat-shocked parasites were then returned to

37°C for the remainder of the 48-hour window until they again reached ring-stage.

Parasite-gDNA was harvested for QIseq after two more rounds of temperature-cycling in

successive growth cycles to ensure enough parasite-material was available for QIseq

(Time-point 1; T^1). Control-parasites were harvested for gDNA before and after three

cycles of pooled growth at 37°C (T^0 and T^1, respectively) for quantification via QIseq in

technical triplicate. We used QIseq-reads obtained for each mutant after the same

number of cycles of pooled growth at 37°C as our T^0 control as previously reported [13].

Drug-screens

As with the HS-screen, parasites were split from the same thaw of the pilot-library after

one cycle of growth into experimental flasks and control-flasks. Experimental flasks were

exposed to three cycles of continuous drug-pressure at two different concentrations

(IC10, IC25) of each artemisinin-compound (AS, DHA). Proteasome-inhibitor BTZ-

experiments were performed at IC10. Control-flasks were cultured continuously in

parallel at 37°C without drug. Parasites were harvested immediately at the conclusion of

three growth-cycles for gDNA-extraction and phenotype-analysis via QIseq.

Oxidative stress screens

Parasites were split after one cycle of growth from the same thaw of the pilot-library as

the HS-screen. Parasites were grown one more cycle, then split into four flasks: two

control-flasks to be cultured with standard, washed human red blood-cells (hRBC), and

two experimental flasks to be cultured with H_2O_2-treated hRBCs to mimic conditions of

oxidative stress. Experimental flasks (H_2O_2 treated-hRBC) and control-flasks (untreated-

hRBC) were cultured continuously in parallel at 37°C. Parasites were harvested

immediately after three growth-cycles (T1), then again after an additional three growth-

cycles (T2) for gDNA-extraction and phenotype-analysis by QIseq.

Methods for oxidative pre-treatment of hRBCs were as published previously [48]. Briefly,

O+ hRBCs (Interstate blood bank, packed, 100% hematocrit) were incubated with 1 mM

H_2O_2 (Sigma-aldrich, Cat. no. H1009-100ML) for one hour at room temperature. After

treatment, cells were washed three times with phosphate-buffered saline (PBS) before

dithiothreitol (DTT) was added to a final concentration of 1 mM to heal any reversible

oxidative damages. Cells were then treated with menadione sodium bisulphite for one

hour at room temperature (Sigma-aldrich Cat. no. M5750-100G) and washed five times.

A volume of 3–4 ml of AB medium (RPMI 1640 medium supplemented with 2 mM L-

glutamine, 25 mM HEPES, 100 µM hypoxanthine and 20 µg ml^{-1} gentamicin) was added

on top of the cell-pellet after discarding the final wash. Pre-treated erythrocytes were

stored at 4 °C before use in parasite culture.

All pooled screens (HS, AS, DHA, BTZ, oxidative stress, ideal growth) were performed in

biological duplicate.

QIseq

QIseq, which uses Illumina next-gen sequencing technology and custom library-

preparation to enable sequencing from both the 5' and 3' ends of the *piggyBac*

transposon out into the disrupted genome-sequence, allows quantitative identification of

each *pB*-mutant line by its unique insertion-site within mixed-population pools of *pB*-

mutants [13] (Figure 1B). The anatomy of the *piggyBac* transposon and its distinct 5' and

3' inverted terminal-repeat sequences (ITRs) allows double-verification of insertion-sites;

both 5' and 3' QIseq libraries were therefore generated and sequenced for each sample.

Counts per insertion-site were determined as described previously [13]. Measures

indicating reproducibility for QIseq-data for both the pilot-library and 1K-library screens

are shown in Fig. S5A-B and Fig. S6, respectively. We observed high correlation

between 5' and 3' QIseq-libraries from each pool (Fig. S5A, Fig. S6). Resulting Growth

and HS phenotype-assignments were highly reproducible across three technical and two

biological replicates (HS-Screen, R=0.94; Growth-Screen, R=0.89; Fig. S5B). We found

only weak correlation between Growth-Sensitive and HS-Sensitive phenotypes (R =

0.44; Fig. S5B, Fig. S6), suggesting that our heat-shock exposure-conditions were

sufficient to allow reproducible detection of mutants with specific selection response-

phenotypes from pooled screening.

Calculating mutant fold-change in pooled screening to assign HS- and Growth-

phenotypes

We defined FC-Growth by *pB*-mutant fold-change after three cycles of growth at ideal

temperatures (T^{1-37C}/ T^{0-37C}). FC-HS was defined as *pB*-mutant fold-change after

exposure to heat-shock vs. the non- heat-shocked control (T^{1-41C}/ T^{1-37C}). We used

changes in reads-number detected for each *pB*-mutant in the Growth-Screen and the

HS-Screen as compared to reads-number detected for that mutant in the respective

control-screen to calculate mutant Fold Change (FC) in both screens (Figure 1C-D;

Methods). We then ranked mutants from lowest to highest FC, with lowest FC indicating

highest sensitivity to the screened-condition.

We developed a scoring-system to distinguish mutants with phenotypes specifically in

the condition under selection (HS) vs. those with inherently compromised growth in ideal

conditions, called the Phenotypic Fitness-Score (PFS). PFS_{HS} is the mutant fold-change

in response to heat-shock (FC-HS, 41C/37C) multiplied by the ratio of FC-HS to mutant

fold-change under ideal growth-conditions (FC-HS/FC-Growth), with the smallest and

largest values indicating the largest mutant growth-differentials between the two screens

(smallest PFS_{HS} indicating worse mutant-fitness in the HS-Screen than the Growth-

Screen, and largest PFS $_{HS}$ indicating better mutant-fitness in the HS-Screen than the

Growth-screen; Fig. S6). Mutants exhibiting (1) poor growth in the HS-Screen (i.e., low

FC-HS of < 0.5 based on performance of known HS-Sensitive *pB*-mutant-clones), and

(2) comparatively much better growth in the Growth-Screen (i.e., low PFS $_{HS}$ of < 0.25)

were classified as HS-Sensitive in pooled phenotypic screens (indicated in red in Fig.

1E-F). Mutants exhibiting poor fitness in both the Growth- and HS-Screens (FC-HS < 0.5

and PFS$_{HS}$ > 0.25) are indicated in Fig. 1E-F in yellow (n = 14). These double-sensitive

mutants were not included in our "HS-Sensitive" classification to avoid overinterpretation

of possibly-confounding phenotypes. We classified mutants displaying a slight growth

advantage in response to heat shock (FC-HS > 1.5, n = 28, indicated in the green box,

Fig.1E-F) as "HS-Tolerant". Mutants exhibiting neither sensitivity nor tolerance to heat

shock were classified as HS-Neutral (n = 49).

Assigning drug- and oxidative stress-screen phenotypes

Mutant fold-change in response to the given condition was calculated against an ideal-

growth control as above. Mutants in the top 25% of reads recovered in QIseq in the

screened condition were classified as Tolerant, while mutants in the bottom 25% were

classified as Sensitive.

Comparative RNAseq between wild-type NF54 and two HS-Sensitive mutant parasite

lines in response to heat shock

RNAseq experimental design is outlined in Fig. S8A. Briefly, highly synchronized ring-

stage cultures of wildtype NF54 and HS-Sensitive mutants *LRR5* and *DHC* were split

equally into four T75 flasks each. All parasites were grown at the normal human body

temperature (37C) to early ring-stage. Two flasks of each parasite-line were then

exposed to febrile temperatures (41C) for 8 hours, while the remaining two flasks were

allowed to continue to grow at 37C for 8 hours without exposure to heat-stress. This

temperature-cycling was repeated three times, just as we allowed for the pooled screen.

After the third round of heat-shock (Time 1, T^1), RNA was harvested simultaneously from

both conditions for RNAseq as in [19]. Parasite fold-change in response to HS was

calculated at the time of sample-collection and verified mutant defects in response to HS

as compared to NF54 (Fig S8B). RNA-seq was performed in-house on an Illumina

MiSeq using a 300-cycle V2 MiSeq reagent kit.

RNA-seq data-analysis

RNA-seq reads from each sample were aligned to the P. falciparum reference genome

(PlasmoDB version 28, RRID:SCR_013331). A maximum of one mismatch per read was

allowed. The mapped reads from TopHat [49] were used to assemble known transcripts

from the reference and their abundances were estimated using Cufflinks [50]. The

expression level of each gene was normalized as FPKM (fragments per kilobase of exon

per million mapped reads). We defined expressed genes as those having FPKM > 20 for

at least one biological replicate at either 37°C or 41°C. The fold change of normalized

gene expression between 41°C and 37°C was calculated for every biological replicate.

Fold-change for genes not expressed in both temperatures was set equal to one. We

conservatively filtered out genes in the top and bottom 10% of fold-change to remove

outliers. We then fit a Gaussian model to the log2 fold change (*log2FC*) for every

biological replicate using maximum log likelihood estimation to normalize the fold-

change calculations and assess significance. For each gene *i* with positive fold-change

in response to HS, we calculated the p-value as the fraction of genes *g* with *log2FC$_g$* >

log2FC$_i$. For each gene *i* with negative fold-change in response to HS, p-value was

calculated as the fraction of genes *g* with *log2FC$_g$* > *log2FC$_i$*. The false discovery rate

(FDR) was calculated for each replicate. We defined genes for which FDR < 0.1 as

having significant fold-change in response to HS. Genes were assigned HS phenotype-

categories based on significance and direction of HS-response. We assigned HS

phenotype-categories for 2567 genes using these criteria (Table S3). Heat-shock

phenotypes as identified via pooled phenotypic screening and comparative RNAseq

were highly correlated (Fig S9A-B), supporting our methodology.

GO-term enrichment analyses

All GO-enrichment analyses were performed testing GO-terms mapped to genes in the

category of interest against a background of GO-terms mapped to all other genes in the

analysis. The GO-term database was created from the latest curated *P. falciparum*

ontology available at the time of analysis, downloaded from GeneDB (accessed May 2,

2019) [51]. For enrichment-analysis in the 1K-library screens: Mutants were divided into

HS-phenotype categories, and each category was tested for enrichment against a

background of GO-terms mapped to the genes represented by the remainder of the 922

mutants in the screen using the weighted Fisher/elim hybrid-method of the TopGO

package (v 1.0) available from Bioconductor [52] (Fig. 2B). For enrichment-analysis in

comparative RNAseq data: a database of all GO-terms mapped to the 1298 genes which

could be assigned a HS-phenotype in all three parasites was assembled. Genes were

divided into HS phenotype-categories based on direction of fold-change (Up, Down,

Unchanged) in response to HS in all three parasites, then evaluated for GO-term

enrichment against the background GO-term database of all other genes in the analysis

using the weighted-Fisher/elim hybrid-method of the TopGO package (Fig 3B, Table

S2). For enrichment of apicoplast-targeted genes by RNAseq HS-phenotype category:

enrichment for each investigated GO-term g was calculated as the number of non-HS-

regulated apicoplast genes C annotated to g (the background-distribution) vs. the

number of HS-regulated apicoplast genes C_r annotated to GO-term g. The fraction of

HS-regulated apicoplast-genes to non-HS-regulated apicoplast genes (C_r/C) was

assessed for significance using the Fisher exact test (Fig 4C).

SUPPLEMENTARY FIGURE and TABLE LEGENDS

Figure S1. Schematic overview of the phenotypic screening pipeline. *pB*-mutant

library resources from small (individual, well-characterized mutant-clones) to large (the

1K-Library, comprised of pools randomly selected from the Saturation-Library) were

used to design carefully validated pooled screens at increasingly large scale. High

correlation between mutant-phenotypes in HS-screens and artemisinin (ART)-screens

indicated mechanistic overlap in response to both stressors. Iterative rounds of pooled-

screening for various phenotypes over time enables higher-throughput functional-

annotation of the *P. falciparum* genome.

Figure S2. Extended screening data against the pilot-library and summary.

A. Full drug-screening data for artemisinin-compounds Artesunate (AS) and DHA, and

proteasome-inhibitor Bortezomib (BTZ) against the pilot-library. HS-Sensitive mutants

are significantly more sensitive to each drug than HS-Tolerant mutants. There is no

significant relationship between *pB*-mutant sensitivity to any drug and mutant sensitivity

in standard growth-conditions.

B. HS-Sensitive (red) and HS-Tolerant (green) mutants and their phenotypes across all

pooled phenotypic screens. Mutants are clustered by HS-phenotype.

Figure S3.

Mutants in members of the DV proteome, targets of ART alkylation, and putative

interacting partners of K13 tend to be sensitive to HS.

A. 1k HS-Screen mutants are ordered by FC-HS from HS-Sensitive to HS-Tolerant.

Mutants in digestive vacuole-associated proteins as defined by [26] are indicated in

lavender dots. Gene-symbols for mutants with HS-sensitivity are labeled with black text

(10 of 18 genes). Gene-symbols for HS-Neutral and HS-Tolerant mutants are labeled

with grey text.

B. All mutants in ART alkylation-targets as defined by [9] included in the 1K HS-Screen.

C. HS-screen phenotypes of mutants in putative K13-interacting proteins as defined by

[25].

Figure S4. Core proteasome-components are slightly but universally upregulated in

response to HS as compared to other aggregately upregulated processes which have

more heterogenous expression. Fold-change for most individual proteasome-

components did not meet our threshold to be designated "upregulated". ** Wilcoxon p-

value < 1e-5.

Figure S5. QIseq data-correlations within and between Pilot-Library Screens. A.

Pearson correlations between 5' and 3' QIseq data for 37°C_ideal-growth screen and

41°C_heat-shock screen indicate highly reproducible analyses across technical and

biological replicates in both screens.

B. Correlations within and between 37°C_ideal growth screen and 41°C_heat-shock

screen QIseq data. High correlations within both HS-screens and Growth-screens (HS-

Screen, R=0.94; Growth-Screen, R=0.89) and weak correlation between HS-screens

and Growth-screens (R = ~0.42) suggests heat-shock exposure-conditions were

sufficient to allow reproducible detection of mutants with specific selection response-

phenotypes from pooled screening.

Figure S6. QIseq data-correlations of 1K-Library screening for *P. falciparum* heat-

shock phenotypes. 1K-Library screens were as robust and reproducible as the pilot-

library screens. QIseq data correlations within and between HS-screens and Growth-

screens for a representative pool of the 1K-mutant library are shown.

Figure S7. Phenotypic Fitness-Score in HS (PFS$_{HS}$) distribution across mutant HS

phenotype-classifications in the 1K-Library screen. See Table S2 and Methods for

PFS$_{HS}$ calculation details. HS-Sensitive mutants (mutants displaying defective growth in

response to heat shock but not in response to ideal growth conditions) are assigned the

lowest PFS$_{HS}$, while HS-Tolerant mutations are assigned the highest PFS$_{HS}$.

Figure S8. Methods and validation for comparative RNAseq.

A. RNA sample-collection methods for wildtype malaria-parasite NF54 vs. two HS-

Sensitive *pB*-mutant clones Δ*DHC* **(PB4) and** Δ*LRR5* **(PB31)** in response to febrile

temperatures. Assays were performed in biological duplicate.

B. Validation of HS-Sensitive mutant-clones during RNA-Seq Sample preparation**.** Both

mutants grown individually had growth-defects in response to HS as compared to NF54.

Figure S9. Complementary methods (pooled phenotypic screening, phenotypic

transcriptional profiling of HS-Sensitive mutants vs. wildtype in response to heat

stress) indicate genes driving the parasite heat-stress response.

A. HS-Sensitive *pB* mutants tend to have mutations in genes that have significant

changes in expression in response to heat-stress, while mutants that are neutral to or

tolerant of heat-stress tend to have mutations in genes that are not regulated in

response to heat-stress.

B. *pB* mutants in genes normally up-regulated in response to heat-stress grow poorly in

response to heat-stress (i.e., have significantly lower phenotypic fitness-scores) than

mutants in genes that are neutral or down-regulated in response to heat-stress.

Supplemental Tables:

Table S1. Pooled HS-Screen results of the *P. falciparum pB*-mutant pilot-library (n

= 128).

Table S2. Pooled HS-Screen results of the 1K-Library (n = 922).

Table S3. Comparative RNAseq-results between NF54 and HS-Sensitive mutant-

clones *ΔLRR5* and *ΔDHC* in response to heat-shock.

S3A. All genes classified into HS response-categories in NF54 with or without exposure

to heat-shock using RNAseq data (n = 2567). HS-classifications for each gene in two

HS-Sensitive mutant-lines are indicated where available. Criteria for inclusion: NF54

expression above threshold (FPKM > or = 20 for at least one replicate in at least one

temperature-condition) and FC-HS supported by two biological replicates.

S3B. Genes included in functional enrichment-analyses. Criteria for inclusion: all genes

with expression above threshold AND agreement between replicates as to HS fold-

change classification for all three parasite lines (n = 1298).

S3C. Enriched GO-terms for specified HS-response-categories as included in Figure 3B.

"Annotated": the number of genes annotated to a given GO-term included in the analysis

for all HS response-categories. "Significant": the number of genes annotated to a given

GO-term in the HS response-category being tested for enrichment.

S3D. Full functional enrichment-results for all HS response-categories.

1004

Table S4. Drug- and oxidative stress-screen results of the pilot library (n = 128).

Table S5. Qlseq dataset accession numbers.

References

1.   WHO: **World Malaria Report.** *World Health Organization* 2018.

2.   Gardner MJ, Hall N, Fung E, White O, Berriman M, Hyman RW, Carlton JM, Pain A, Nelson KE, Bowman S, et al: **Genome sequence of the human malaria parasite *Plasmodium falciparum*.** *Nature* 2002, **419:**498-511.

3.   Aurrecoechea C, Brestelli J, Brunk BP, Dommer J, Fischer S, Gajria B, Gao X, Gingle A, Grant G, Harb OS, et al: **PlasmoDB: a functional genomic database for malaria parasites.** *Nucleic Acids Res* 2009, **37:**D539-543.

4.   Oakley MS, Gerald N, McCutchan TF, Aravind L, Kumar S: **Clinical and molecular aspects of malaria fever.** *Trends in parasitology* 2011, **27:**442-449.

5.   Oakley MSM, Kumar S, Anantharaman V, Zheng H, Mahajan B, Haynes JD, Moch JK, Fairhurst R, McCutchan TF, Aravind L: **Molecular factors and biochemical pathways induced by febrile temperature in intraerythrocytic *Plasmodium falciparum* parasites.** *Infection and immunity* 2007, **75:**2012-2025.

6.   Zhang M, Wang C, Otto TD, Oberstaller J, Liao X, Adapa SR, Udenze K, Bronner IF, Casandra D, Mayho M, et al: **Uncovering the essential genes of the human malaria parasite *Plasmodium falciparum* by saturation mutagenesis.** *Science* 2018, **360**.

7.   Rocamora F, Zhu L, Liong KY, Dondorp A, Miotto O, Mok S, Bozdech Z: **Oxidative stress and protein damage responses mediate artemisinin resistance in malaria parasites.** *PLoS pathogens* 2018, **14:**e1006930-e1006930.

8.   Bridgford JL, Xie SC, Cobbold SA, Pasaje CFA, Herrmann S, Yang T, Gillett DL, Dick LR, Ralph SA, Dogovski C, et al: **Artemisinin kills malaria parasites by damaging proteins and inhibiting the proteasome.** *Nature Communications* 2018, **9:**3801.

9.   Ismail HM, Barton V, Phanchana M, Charoensutthivarakul S, Wong MHL, Hemingway J, Biagini GA, O'Neill PM, Ward SA: **Artemisinin activity-based probes identify multiple molecular targets within the asexual stage of the malaria parasites *Plasmodium falciparum* 3D7.** *Proceedings of the National Academy of Sciences* 2016, **113:**2080-2085.

1042   10.   Mok S, Ashley EA, Ferreira PE, Zhu L, Lin Z, Yeo T, Chotivanich K, Imwong M,
1043   Pukrittayakamee S, Dhorda M, et al: **Drug resistance. Population**
1044   **transcriptomics of human malaria parasites reveals the mechanism of**
1045   **artemisinin resistance.** *Science* 2015, **347:**431-435.

1046   11.   Yang T, Yeoh LM, Tutor MV, Dixon MW, McMillan PJ, Xie SC, Bridgford JL,
1047   Gillett DL, Duffy MF, Ralph SA, et al: **Decreased K13 Abundance Reduces**
1048   **Hemoglobin Catabolism and Proteotoxic Stress, Underpinning Artemisinin**
1049   **Resistance.** *Cell Reports* 2019, **29:**2917-2928.e2915.

1050   12.   Birnbaum J, Scharf S, Schmidt S, Jonscher E, Hoeijmakers WAM, Flemming S,
1051   Toenhake CG, Schmitt M, Sabitzki R, Bergmann B, et al: **A Kelch13-defined**
1052   **endocytosis pathway mediates artemisinin resistance in malaria parasites.**
1053   *Science* 2020, **367:**51.

1054   13.   Bronner IFF, Ottoa TD, Zhang M, Udenze K, Wang CCQ, Quail MA, Jiang RHY,
1055   Adams JH, Rayner JC: **Quantitative Insertion-site Sequencing (QIseq) for**
1056   **high throughput phenotyping of transposon mutants.** *Genome Research*
1057   2016.

1058   14.   Balu B, Chauhan C, Maher SP, Shoue DA, Kissinger JC, Fraser MJ, Adams JH:
1059   **piggyBac is an effective tool for functional analysis of the *Plasmodium***
1060   ***falciparum* genome.** *BMC microbiology* 2009, **9:**83.

1061   15.   Thomas P, Sedillo J, Oberstaller J, Li S, Zhang M, Singh N, Wang CC, Udenze
1062   K, Jiang RH, Adams JH: **Phenotypic Screens Identify Parasite Genetic**
1063   **Factors Associated with Malarial Fever Response in *Plasmodium***
1064   ***falciparum piggyBac* Mutants.** *mSphere* 2016, **1**.

1065   16.   Bushell E, Gomes AR, Sanderson T, Anar B, Girling G, Herd C, Metcalf T,
1066   Modrzynska K, Schwach F, Martin RE, et al: **Functional Profiling of a**
1067   **Plasmodium Genome Reveals an Abundance of Essential Genes.** *Cell* 2017,
1068   **170:**260-272.e268.

1069   17.   Gisselberg JE, Zhang L, Elias JE, Yeh E: **The Prenylated Proteome of**
1070   ***Plasmodium falciparum* Reveals Pathogen-specific Prenylation Activity and**
1071   **Drug Mechanism-of-action.** *Molecular & Cellular Proteomics* 2017,
1072   **16:**S54-S64.

1073   18.   Suazo KF, Schaber C, Palsuledesai CC, Odom John AR, Distefano MD: **Global**
1074   **proteomic analysis of prenylated proteins in *Plasmodium falciparum* using**
1075   **an alkyne-modified isoprenoid analogue.** *Scientific Reports* 2016, **6:**38615.

1076   19.   Gibbons J, Button-Simons KA, Adapa SR, Li S, Pietsch M, Zhang M, Liao X,
1077   Adams JH, Ferdig MT, Jiang RHY: **Altered expression of K13 disrupts DNA**
1078   **replication and repair in *Plasmodium falciparum*.** *BMC Genomics* 2018,
1079   **19:**849.

1080   20.   Krishnan KM, Williamson KC: **The proteasome as a target to combat malaria:**
1081   **hits and misses.** *Translational Research* 2018, **198:**40-47.

21.   Ng CL, Fidock DA, Bogyo M: **Protein Degradation Systems as Antimalarial**
Therapeutic Targets. *Trends in Parasitology* 2017, **33:**731-743.

22.   Ariey F, Witkowski B, Amaratunga C, Beghain J, Langlois A-C, Khim N, Kim S,
Duru V, Bouchier C, Ma L, et al: **A molecular marker of artemisinin-resistant**
Plasmodium falciparum **malaria.** *Nature* 2014, **505:**50-55.

23.   Tilley L, Straimer J, Gnädig NF, Ralph SA, Fidock DA: **Artemisinin Action and**
Resistance in *Plasmodium falciparum*. *Trends in parasitology* 2016, **32:**682-
696.

24.   Bhattacharjee S, Coppens I, Mbengue A, Suresh N, Ghorbal M, Slouka Z,
Safeukui I, Tang H-Y, Speicher DW, Stahelin RV, et al: **Remodeling of the**
malaria parasite and host human red cell by vesicle amplification that
induces artemisinin resistance. *Blood* 2018, **131:**1234-1247.

25.   Gnädig NF, Stokes BH, Edwards RL, Kalantarov GF, Heimsch KC, Kuderjavy M,
Crane A, Lee MCS, Straimer J, Becker K, et al: **Insights into the intracellular**
localization, protein associations and artemisinin resistance properties of
Plasmodium falciparum **K13.** *PLOS Pathogens* 2020, **16:**e1008482.

26.   Lamarque M, Tastet C, Poncet J, Demettre E, Jouin P, Vial H, Dubremetz JF:
Food vacuole proteome of the malarial parasite *Plasmodium falciparum*.
Proteomics Clin Appl 2008, **2:**1361-1374.

27.   Dogovski C, Xie SC, Burgio G, Bridgford J, Mok S, McCaw JM, Chotivanich K,
Kenny S, Gnädig N, Straimer J, et al: **Targeting the Cell Stress Response of**
Plasmodium falciparum **to Overcome Artemisinin Resistance.** *PLOS Biology*
2015, **13:**e1002132.

28.   White JK, Handa S, Vankayala SL, Merkler DJ, Woodcock HL: **Thiamin**
Diphosphate Activation in 1-Deoxy-d-xylulose 5-Phosphate Synthase:
Insights into the Mechanism and Underlying Intermolecular Interactions.
The journal of physical chemistry B 2016, **120:**9922-9934.

29.   Imlay L, Odom AR: **Isoprenoid Metabolism in Apicomplexan Parasites.**
Current Clinical Microbiology Reports 2014, **1:**37-50.

30.   Kennedy K, Cobbold SA, Hanssen E, Birnbaum J, Spillman NJ, McHugh E,
Brown H, Tilley L, Spielmann T, McConville MJ, Ralph SA: **Delayed death in the**
malaria parasite *Plasmodium falciparum* **is caused by disruption of**
prenylation-dependent intracellular trafficking. *PLOS Biology* 2019,
17:e3000376.

31.   Howe R, Kelly M, Jimah J, Hodge D, Odom AR: **Isoprenoid biosynthesis**
inhibition disrupts Rab5 localization and food vacuolar integrity in
Plasmodium falciparum. *Eukaryotic cell* 2013, **12:**215-223.

32.   Pandey AV, Tekwani BL, Singh RL, Chauhan VS: **Artemisinin, an**
endoperoxide antimalarial, disrupts the hemoglobin catabolism and heme

detoxification systems in malarial parasite. *J Biol Chem* 1999, **274:**19383-
19388.

33.   del Pilar Crespo M, Avery TD, Hanssen E, Fox E, Robinson TV, Valente P,
Taylor DK, Tilley L: **Artemisinin and a series of novel endoperoxide**
antimalarials exert early effects on digestive vacuole morphology.
Antimicrob Agents Chemother 2008, **52:**98-109.

34.   Sussmann RAC, Fotoran WL, Kimura EA, Katzin AM: ***Plasmodium falciparum***
uses vitamin E to avoid oxidative stress. *Parasites & vectors* 2017, **10:**461-
461.

35.   Mène-Saffrané L: **Vitamin E Biosynthesis and Its Regulation in Plants.**
Antioxidants (Basel, Switzerland) 2017, **7:**2.

36.   Estévez JM, Cantero A, Reindl A, Reichler S, León P: **1-Deoxy-d-xylulose-5-**
phosphate Synthase, a Limiting Enzyme for Plastidic Isoprenoid
Biosynthesis in Plants. *Journal of Biological Chemistry* 2001, **276:**22901-
22909.

37.   Zhang F, Liu W, Xia J, Zeng J, Xiang L, Zhu S, Zheng Q, Xie H, Yang C, Chen
M, Liao Z: **Molecular Characterization of the 1-Deoxy-D-Xylulose 5-**
Phosphate Synthase Gene Family in *Artemisia annua*. *Frontiers in Plant*
Science 2018, **9**.

38.   Heuston S, Begley M, Gahan CGM, Hill C: **Isoprenoid biosynthesis in**
bacterial pathogens. *Microbiology* 2012, **158:**1389-1401.

39.   Brunetti C, Guidi L, Sebastiani F, Tattini M: **Isoprenoids and phenylpropanoids**
are key components of the antioxidant defense system of plants facing
severe excess light stress. *Environmental and Experimental Botany* 2015,
119:54-62.

40.   Clarke A: **The thermal limits to life on Earth.** *International Journal of*
Astrobiology 2014, **13:**141-154.

41.   Kobayashi Y, Harada N, Nishimura Y, Saito T, Nakamura M, Fujiwara T, Kuroiwa
T, Misumi O: **Algae sense exact temperatures: small heat shock proteins are**
expressed at the survival threshold temperature in *Cyanidioschyzon*
***merolae* and *Chlamydomonas reinhardtii*.** *Genome biology and evolution*
2014, **6:**2731-2740.

42.   Li GQ, Arnold K, Guo XB, Jian HX, Fu LC: **Randomised comparative study of**
mefloquine, qinghaosu, and pyrimethamine-sulfadoxine in patients with
falciparum malaria. *Lancet* 1984, **2:**1360-1361.

43.   Kirkman LA, Zhan W, Visone J, Dziedziech A, Singh PK, Fan H, Tong X, Bruzual
I, Hara R, Kawasaki M, et al: **Antimalarial proteasome inhibitor reveals**
collateral sensitivity from intersubunit interactions and fitness cost of
resistance. *Proceedings of the National Academy of Sciences* 2018, **115:**E6863.

44.   Oberstaller J, Zhang M, Wang C: **RNAseq dataset: Malaria-parasite survival**
1161   **of host fever is linked to artemisinin resistance** 2020.

1162   45.   Boucher MJ, Yeh E: **Evidence that disruption of apicoplast protein import in**
1163   **malaria parasites evades delayed-death growth inhibition.** *bioRxiv*
1164   2018**:**422618.

1165   46.   Ralph SA, van Dooren GG, Waller RF, Crawford MJ, Fraunholz MJ, Foth BJ,
1166   Tonkin CJ, Roos DS, McFadden GI: **Metabolic maps and functions of the**
1167   ***Plasmodium falciparum* apicoplast.** *Nature Reviews Microbiology* 2004, **2:**203.

1168   47.   Guggisberg AM, Park J, Edwards RL, Kelly ML, Hodge DM, Tolia NH, Odom AR:
1169   **A sugar phosphatase regulates the methylerythritol phosphate (MEP)**
1170   **pathway in malaria parasites.** *Nature Communications* 2014, **5:**4467.

1171   48.   Cyrklaff M, Srismith S, Nyboer B, Burda K, Hoffmann A, Lasitschka F, Adjalley S,
1172   Bisseye C, Simpore J, Mueller A-K, et al: **Oxidative insult can induce malaria-**
1173   **protective trait of sickle and fetal erythrocytes.** *Nature Communications* 2016,
1174   **7:**13401.

1175   49.   Trapnell C, Pachter L, Salzberg SL: **TopHat: discovering splice junctions with**
1176   **RNA-Seq.** *Bioinformatics* 2009, **25:**1105-1111.

1177   50.   Trapnell C, Williams BA, Pertea G, Mortazavi A, Kwan G, van Baren MJ,
1178   Salzberg SL, Wold BJ, Pachter L: **Transcript assembly and quantification by**
1179   **RNA-Seq reveals unannotated transcripts and isoform switching during**
1180   **cell differentiation.** *Nat Biotech* 2010, **28:**511-515.

1181   51.   Logan-Klumpler FJ, De Silva N, Boehme U, Rogers MB, Velarde G, McQuillan
1182   JA, Carver T, Aslett M, Olsen C, Subramanian S, et al: **GeneDB--an annotation**
1183   **database for pathogens.** *Nucleic acids research* 2012, **40:**D98-108.

1184   52.   Alexa A, Rahnenfuhrer J, Lengauer T: **Improved scoring of functional groups**
1185   **from gene expression data by decorrelating GO graph structure.**
1186   *Bioinformatics* 2006, **22:**1600-1607.
1187

Figure 1. Pooled screens of an extensively characterized *pB*-mutant clone-library allow robust identification of heat-shock phenotypes

A. Experimental design for pooled heat shock (HS) phenotypic screens. The pilot-library of *pB*-mutant clones (n=128) was exposed to conditions simulating malarial fever. A pilot-library control concurrently grown continuously at 37°C established inherent growth of each *pB*-mutant.

B. QIseq quantifies each *pB*-mutant in the pilot library from sequence-reads of the 5' and 3' ends of each *pB* insertion-site. Colored lines represent genes. Black boxes indicate transposon location.

C. Pilot-library mutant growth-phenotypes at ideal temperatures, ranked from Sensitive to Tolerant by FC-Growth. Mutants with inherently slower or faster growth under ideal conditions are shown in grey and blue, respectively.

D. Pilot-library mutant HS-phenotypes ordered from Sensitive (red) to Tolerant (green).

E-F. HS- and Growth-phenotypes of the pilot-library mutants. HS-phenotype of each mutant (displayed as line-graph) is superimposed on its corresponding Growth-phenotype (bar graph). Red = HS-Sensitive mutants (n = 28). Yellow = mutants classified as both Growth-Sensitive and HS-Sensitive (n = 14). Green = HS-Tolerant mutants (n = 30). Mutants neither Sensitive nor Tolerant to HS were classified as HS-Neutral (n = 49). *Known HS-Sensitive *pB*-mutant clones validated by individual HS-assay [15]. **Known HS-Tolerant *pB*-mutant clones validated by individual HS-assay [15].

G. Distributions of PFS_{HS} for mutant HS-phenotype classifications (**** Wilcoxon-test p-value < 1e-15).

Figure 2. Pooled phenotypic screens scaled up to a 1K *pB*-mutant library enable identification of processes driving the *P. falciparum* heat-shock response

A. HS-Sensitive mutations identified in pooled screens of 922 *pB*-mutants. Mutants are ranked by fold-change in response to HS (FC-HS; n = 922) from HS-Sensitive (red; n = 149) to HS-Tolerant (green; n = 139). Mean mutant fold-change in ideal growth (FC-Growth) is superimposed as a bar plot (gray, FC-Growth < 1.0; blue, FC-Growth > 1.0). Mutants performing poorly in both screens (yellow; n = 91) were classified as HS- and Growth-Sensitive. Mutations neither HS-Sensitive nor HS-Tolerant were classified as HS-Neutral. See Table S2.

B. Functional enrichment of GO terms for HS-Sensitive or Growth-Sensitive *pB*-mutants vs all other mutants in the 1K-library. HS-Sensitive mutants were enriched in terms associated with HS-response such as protein-folding, response to DNA-damage, DNA-repair, and regulation of vesicle-mediated transport. Growth-Sensitive mutants tended to be enriched for more general categories broadly important for survival in all conditions, such as translation- or mRNA-metabolism-related terms. Circles represent GO category, circle color represents ontology, and circle size represents number of significant genes annotated to that category. Significant terms (Fisher/elim-hybrid test p. value <= 0.05) fall within the light-green box.

Figure 3. **Increased transcription of the unfolded protein response, organelle-targeted stress-response pathways and host-cell remodeling characterize the parasite HS-response.**

A. (Top-left) HS expression-categories based on phenotype in NF54 vs. HS-Sensitive mutants *ΔLRR5* and *ΔDHC* with and without HS-exposure. HS-regulated genes shared between NF54 and both mutants are indicated in red (↑↑↑) or blue (↓↓↓) for up- and down-regulated genes, respectively. Genes dysregulated in one or both HS-Sensitive mutants fell into two main expression-profile categories underlying mutant HS-Sensitivity phenotypes: those upregulated in NF54 that failed to be regulated in the mutants (↑✗✗), and genes not regulated in response to HS in NF54 that were inappropriately upregulated in the mutants (− ↑↑).

B. (Top-right) Functional enrichment analyses between wildtype/mutant HS-expression profiles as defined in A. Red: Shared upregulated HS-responsive GO-terms between NF54 and the two HS-Sensitive *pB*-mutants (↑↑↑). Blue: Shared down-regulated HS-responsive GO-terms (↓↓↓). Ochre: GO-terms upregulated in NF54 but dysregulated in the two *pB*-mutant (↑✗✗). Tan: GO-terms enriched in genes not regulated in the wildtype HS-response but upregulated in the mutants (− ↑↑). Only enriched GO-terms are shown (Fisher/elim-hybrid test p. value <= 0.05), with highest significance indicated in dark green. Fraction of significant genes mapping to a GO-term in an HS expression-profile category vs. genes mapping to that GO-term in the entire analysis is indicated by distance to the center of the circle, with the outermost position on the circle indicating 100% of genes in that GO-term are significant. See Table S3D.

C. (Bottom-left) Several apicoplast and isoprenoid biosynthesis-related genes have a tendency to be upregulated in the wildtype-response to HS and are dysregulated in one or both HS-Sensitive *pB*-mutant clones (↑✗✗). * Isoprenoid biosynthesis-related genes upregulated by HS confirmed in the pooled HS-screen.

Figure 4. Apicoplast isoprenoid biosynthesis is critical for *P. falciparum* survival of febrile temperatures

Figure 4. Apicoplast isoprenoid biosynthesis is critical for *P. falciparum* survival of febrile temperatures.

A. Apicoplast-targeted genes tend to be increased in response to HS as compared to all non apicoplast-targeted genes detected above threshold in RNAseq. Apicoplast-targeted genes are as defined in [45] (*** Fisher-test p-value < 1e-5).

B. Apicoplast-targeted genes tend to be highly essential during blood-stage vs. all other non-apicoplast-targeted genes detected above threshold in RNAseq. The median Mutagenesis Index Score (MIS; [6]) for apicoplast-targeted genes is much lower than median MIS for all other genes, indicating a lower tolerance for disruption and thus higher likely essentiality during blood-stage development than non-apicoplast-targeted genes (**** Wilcoxon-test p-value < 1e-15).

C. Apicoplast pathways regulated in response to HS. GO categories enriched in up- and down-regulated apicoplast genes are shown on a scale from red to blue, respectively. The horizontal direction indicates the log ratio between up- and down-regulated apicoplast genes in each category. Circle-size represents gene-number per category.

D. All 8 *pB*-mutants in genes associated with the apicoplast isoprenoid biosynthesis pathway (including direct downstream prenylated targets) represented in the pooled screen were sensitive to heat-shock. Mutants are ranked from HS-Sensitive (red) to HS-Tolerant (green). Circles indicate each HS-Sensitive mutant related to isoprenoid-biosynthesis. See Table S3.

E. Key enzymes in the *P. falciparum* isoprenoid biosynthesis-pathway are up-regulated in response to HS (red circle), dysregulated in HS-Sensitive mutants (ochre) and absent in malaria-parasites of hosts that do not present fever. Pathway diagram modeled from [46]. Isoprenoid biosynthesis-genes upregulated in HS include DXS, 1-deoxy-D-xylulose-5-phosphate synthase (DOXP), 2-C-methyl-D-erythritol 2,4-cyclodiphosphate synthase (IspF), pyruvate kinase II (PyKII), triosephosphate isomerase (TIM), triose phosphate transporter (TPT), and upstream-regulator of MEP-pathway substrates HAD1-phosphotase [47]. All represented Rab GTPases, vesicular trafficking-proteins that are direct downstream-targets prenylated by the MEP-pathway (zigzag) were HS-Sensitive. The key thiamin-synthesis enzyme hydroxyethylthiazole kinase (ThzK) is absent in *P. berghei* and *P. yoelii*, malaria-parasites whose rodent-hosts do not present fever.

F. *P. falciparum* genes with plant orthologs (green circles; n = 1919) indicating potential endosymbiont-ancestry tend to be increased in response to HS vs. genes that do not have plant orthologs (grey circles). The listed processes are sorted based on the ratio of "green" to "non-green" orthologs.

Figure 5.

Increased sensitivity to fever is directly correlated with increased sensitivity to artemisinin in the malaria parasite

Figure 5. Increased sensitivity to fever is directly correlated with increased sensitivity to artemisinin in the malaria parasite.

A. HS-Sensitive *pB*-mutants (red) are more sensitive to multiple concentrations of artemisinin derivatives AS and DHA, proteasome-inhibitor BTZ, and conditions of heightened oxidative stress than HS-Tolerant parasites (green) in all pooled screens of the pilot library. Apicoplast-mutants (n = 5) have phenotypes similar to all HS-Sensitive mutants (n =28) in artemisinin-derivative screens, but not to proteasome-inhibitors or oxidative stress (* Wilcoxon p < 0.05; *** Wilcoxon p < 1e-10. See Methods).

B. Correlation between mutant phenotypes in all pooled screens of the pilot library.

C. HS-Sensitive *pB*-mutants (red) are positively correlated with genes linked to ART-R in recent field isolates [10](* Wilcoxon test p < 0.05).

D. HS expression-categories as determined by comparative RNAseq between NF54 and two HS-Sensitive mutant clones are positively correlated with ART-R in field-isolates.

E. Both K13-mediated mechanisms of artemisinin resistance (endocytosis, ubiquitin-proteasome system) are similarly regulated in HS. The K13-defined endocytosis pathway (shades of green) and key ubiquitinating-enzymes of the ubiquitin-proteasome system, E2/E3 and K13, are downregulated in the wildtype NF54 HS-response, while protein folding, stress, exported proteins, and proteasome genes are upregulated. RNAseq data are plotted for each gene by average log2 fold-change in response to HS and significance (−log2(p-value)). Circles in shades of blue and pink indicate genes significantly down- or upregulated after exposure to HS, respectively.

F. Activation of pathways underlying DHA-mediated killing and febrile-temperature survival is directly inverse. **Top**: Model of DHA-mediated killing in *P. falciparum* adapted from [8]. Artemisinin (ART) damages and unfolds proteins, prevents folding of newly synthesized proteins, and inhibits the proteasome while at the same time activating E1/E2/E3 ubiquitin-machinery. Accumulation of toxic polyubiquitinated protein-substrates (S) overwhelms the cell and leads to death. **Bottom**: Model of parasite fever-response. Heat-stress causes globally damaged protein. The parasite increases the UPR as it inhibits E2/E3 ubiquitination to prevent accumulation of toxic, polyubiquitinated (Ub) protein-aggregates, while at the same time increasing its capacity for proteasome-mediated degradation—ultimately enabling the parasite to resolve HS-instigated stress and survive febrile-temperatures.