Molecular evolutionary trends and biosynthesis pathways in the Oribatida revealed by the

genome of *Archegozetes longisetosus*

Adrian Brückner[1]*, Austen A. Barnett[2], Igor A. Antoshechkin[1] and Sheila A. Kitchen[1]

[1]Division of Biology and Biological Engineering, California Institute of Technology, 1200 East

California Boulevard, Pasadena, CA 91125, United States of America

[2]Department of Biology, DeSales University, 2755 Station Avenue, Center Valley, PA 18034,

United States of America

* corresponding author: adrian.brueckner@gmail.com

ABSTRACT

Background

Oribatid mites are a specious order of microarthropods within the subphylum Chelicerata,

compromising about 11,000 described species. They are ubiquitously distributed across different

microhabitats in all terrestrial ecosystems around the world and were among the first animals

colonizing terrestrial habitats as decomposers and scavengers. Noted for their biosynthesis

capacities and biochemical diversity, the majority of oribatid mites possess a pair of exocrine

opisthonotal oil-glands used for chemical defense and communication. Genomic resources are

lacking for oribatids despite their species richness and ecological importance.

Results

20   We used a comparative genomic approach to investigate the developmental, sensory and

21 glandular biosynthetic gene repertoire of the clonal, all-female oribatid mite species *Archegozetes*

longisetosus Aoki, a model species used by numerous laboratories for the past 30 years. Here, we present a 190-Mb genome assembly constructed from Nanopore MinION and Illumina sequencing platforms with 23,825 predicted protein-coding genes. Genomic and transcriptional analyses revealed patterns of reduced body segmentation and loss of segmental identity gene *abd-A* within Acariformes, and unexpected expression of key eye development genes in these eyeless mites across developmental stages. Consistent with the soil dwelling lifestyle, investigation of the sensory genes revealed a species-specific expansion of gustatory receptors, the largest chemoreceptor family in the genome used in olfaction, and evidence of horizontally transferred enzymes used in cell wall degradation of plant and fungal matter, both components of the *Archegozetes longisetosus* diet. Using biochemical and genomic data, we were able to delineate the backbone biosynthesis of monoterpenes, an important class of compounds found in the major exocrine gland system of Oribatida – the oil glands.

Conclusions

With the *Archegozetes longisetosus* genome, we now have the first high-quality, annotated genome of an oribatid mite genome. Given the mite's strength as an experimental model, the new sequence resources provided here will serve as the foundation for molecular research in Oribatida and will enable a broader understanding of chelicerate evolution.

Keywords

soil animal, terpenes, horizontal gene transfer, parthenogenesis, chemoreceptors, Hox genes, model organism, RNAseq, MinION long-read sequencing, Sarcoptiformes

Introduction

In the past couple of years, the number of sequenced animal genomes has increased dramatically, especially for arthropods about 500 genomes sequences are now available (1, 2). The majority of these genomes, however, belong to the insects (e.g. flies, beetles, wasp, butterflies and bugs (2)) which compromise the most diverse, yet evolutionarily young and more derived taxa of arthropods (3, 4). In strong contrast, genome assemblies, many of which are incomplete or not well annotated, exist for the Chelicerata (1) – the other major subphylum of arthropods (3, 4). Chelicerates include sea spiders, spiders, mites and scorpions among other organisms, as well as several extinct taxa (5, 6). Chelicerates originated as marine animals about 500 million years ago (5, 7). Molecular analyses suggest that one particular group, the omnivorous and detritivores acariform mites, may have been among the first arthropods that colonized terrestrial habitats and gave rise to ancient, simple terrestrial food webs (8-10).

So far, the well-annotated genomic data of chelicerates is limited to animal parasites (including human pathogens and ticks), plant parasites, and predatory mites used in pest control (11-17). Other than some lower-quality genome assemblies (18), there are no resources available for free-living soil and litter inhabiting species. Such data are, however, pivotal to understanding the evolution of parasitic lifestyles from a free-living condition and to bridge the gap between early aquatic chelicerates such as horseshoe crabs, and highly derived terrestrial pest species and parasites (19-21). Because the phylogeny of Chelicerata remains unresolved, additional chelicerate genomes are urgently needed for comparative analyses (6, 7, 22). To help address this deficit, we report here the genome assembly of the soil dwelling oribatid mite *Archegozetes longisetosus* (Aoki, 1965; **Figure 1**) (23) and a comprehensive analysis in the context of developmental genes, feeding biology, horizontal gene transfer and biochemical pathway evolution of chelicerates.

Archegozetes longisetosus (hereafter referred to as *Archegozetes*) is a member of the

Oribatida (Acariformes, Sarcoptiformes), an order of chelicerates well-known for their exceptional

biosynthesis capacities, biochemical diversity, unusual mode of reproduction, unusually

highpulling strength, mechanical resistances and pivotal ecological importance (24-32).

Archegozetes, like all members of its family Trhypochthoniidae (Figure 1a), reproduce via

thelytoky (33). That means the all-female lineages procreate via automictic parthenogenesis with

an inverted meiosis of the holokinetic chromosomes, resulting in clonal offspring (34-37). While

studying a parthenogenetic species is useful for the development of genetic tools as stable germ-

line modifications can be obtained from the clonal progeny without laboratory crosses, one is

confronted with the technical and philosophical problems of species delineation, cryptic diversity

and uncertain species distribution (33, 38). Reviewing all available data, Norton (39, 40) and

Heethoff et al. (33) concluded that *Archegozetes* is found widely on continents and islands

throughout the tropical and partly subtropical regions of the world and that it is a middle-derived

oribatid mite closely related to the suborder Astigmata.

One major feature of most oribatid mites is a pair of opisthonotal oil-glands and

Archegozetes is no exception (30, 41). These are a pair of large exocrine glands, each composed

of a single-cell layer invagination of the cuticle, which is the simplest possible paradigm of an

animal gland (42, 43). The biological role of these glands was rather speculative for a long time;

idea ranged from a lubricating and osmo- or thermoregulative function (44-46) to roles in chemical

communication (47-49). So far about 150 different gland components have been identified from

oribatid mites, including mono- and sesquiterpenes, aldehydes, esters, aromatics, short-chained

hydrocarbons, hydrogen cyanide (HCN) and alkaloids (26, 30, 50-52). While some chemicals

appear to be alarm pheromones (47, 49), most function as defensive allomones (48). Interestingly,

alkaloids produced by oribatids mites are the ultimate source of most toxins sequestered by poison-

frogs (51, 53).

Terrestrial chelicerates predominately ingest fluid food. While phloem-feeding plant pests

like spider mites and ecotoparasites likes ticks adapted a sucking feeding mode, scorpions, spiders

and others use external, pre-oral digestion before ingestion by morphologically diverse mouthparts

(9, 12, 54, 55). Exceptions from this are Opiliones and sarcoptiform mites, i.e. oribatid and

astigmatid mites,all of which ingest solid food (40, 56, 57). In general, oribatids feed on a wide

range of different resources and show a low degree of dietary specialization (58). The typical food

spectrum of Oribatida, includes leaf-litter, algae, fungi, lichens, nematodes, and small dead

arthropods such as collembolans (44, 59-62). In laboratory feeding trials, oribatid mites tend to

prefer dark pigmented fungi, but also fatty acid-rich plant-based food (58, 61). Additionally,

stable-isotope analyses of ^{15}N and ^{13}C suggested that Oribatida are primary- and secondary

decomposers feeding on dead plant material and fungi, respectively (59, 63, 64). The reasons for

these preferences are still unknown, but they raise the question of how oribatid mites are able to

enzymatically digest the cell walls of plants and fungi (58, 60, 64, 65).

Early studies on *Archegozetes* and other mites found evidence for cellulase, chitinase and

trehalase activity which was later attributed to symbiotic gut bacteria (65-71). While such bacterial

symbionts are a possible explanation, genomic data of other soil organisms and plant-feeding

arthropods suggest a high frequency of horizontal transfer of bacterial and fungal genes enabling

the digestion of cell walls (11, 72-76). For instance, an in-depth analysis of the spider mite

Tetranychus urticae revealed a massive incorporation of microbial genes into the mite's genome

(11, 75). Horizontal gene transfer appears to be a common mechanism for soil organisms,

including mites, to acquire novel metabolic enzymes (11, 17, 74, 76-78), and hence seems very

likely for *Archegozetes* and other oribatid mite species that feed on plant or fungal matter.

Archegozetes has been established as a laboratory model organism for three decades,

having been used in studies, ranging from ecology, morphology, development and eco-toxicology

to physiology and biochemistry (27, 33, 79-83). As such, *Archegozetes* is among the few

experimentally tractable soil organisms and by far the best-studied oribatid mite species (33, 81,

84). Since *the mite* meets the most desirable requirements for model organisms (84), that is a rapid

development under laboratory conditions, a dedicated laboratory strain was named *Archegozetes*

longisetosus **ran** in reference to its founder **R**oy **A**. **N**orton (33 , **Figure 1b-c**). Their large number

of offspring enables mass cultures of hundreds of thousands of individuals, and their cuticular

transparency during juvenile stages, and weak sclerotization as adults are general assets of an

amenable model system (33, 85-87). In the past 10 years, *Archegozetes* also received attention as

a model system for chemical ecology (27, 85, 86, 88-91). Some of these studies focusing on the

Archegozetes gland revealed basic insights into the chemical ecology and biochemical capabilities

of arthropods (27, 89, 91). Hence, *Archegozetes* is poised to become a genetically tractable model

to study the molecular basis of gland and metabolic biology.

The aim and focus of the current study were three-fold – to provide well-annotated, high-

quality genomic and transcriptomic resources for *Archegozetes longisetosus* (**Figure 1**), to reveal

possible horizontal gene transfers that could further explain the feeding biology of oribatids, and

to present *Archegozetes* as a research model for biochemical pathway evolution. Through a

combination of comparative genomic and detailed computational analyses, we were able to

generate a comprehensive genome of *Archegozetes* and provide it as an open resource for genomic,

developmental and evolutionary research. We further identified candidate horizontal gene transfer

events from bacteria and fungi that are mainly related to carbohydrate metabolism and cellulose

digestion, features correlated with the mite feeding biology. We also used the genomic data

together with stable-isotope labeling experiments and mass spectrometric investigation to

delineate the biosynthesis pathway of monoterpenes in oribatid mites.

Results and Discussion

Archegozetes longisetosus genus assembly

Archegozetes longisetosus (**Figure 1**) has a diploid chromosome number (2n) of 18 (36), most likely comprising 9 autosomal pairs, the typical number of nearly all studied oribatid mite species (92). There are no distinct sex chromosomes in *Archegozetes*; this appears to be ancestral in the Acariformes and persisted in the Oribatida (35, 36, 92). Even though some XX:XO and XX:XY genetic systems have been described in the closely related Astigmata, the sex determination mechanism in oribatids, including *Archegozetes,* remains unknown (33, 35, 36, 92, 93)*.* To provide genetic resources, we sequenced and assembled the genome using both Illumina short-read and Nanopore MinION long-read sequencing approaches (**Table 1**; see also "**Materials and Methods**"). Analyses of the *k-mer* frequency distribution of short reads (**Table 1; Supplementary Figure S1**) resulted in an estimated genome size range of 135-180 Mb, smaller than the final assembled size of 190 Mb (**Table 1**; see also "**Materials and Methods**"). This difference was suggestive of high repetitive content in the genome of *Archegozetes* and indeed, repeat content was predicted to be 32 % of the genome (**see below**) (94, 95). Compared to genome assemblies of other acariform mites, the assembled genome size of *Archegozetes* is on the large end, but is smaller than that of mesostigmatid mites, ticks and spiders (11-13, 15, 17, 18, 96). In the context of arthropods in general, *Archegozetes*'s genome (**Table 1**) is among the smaller ones and shares this feature with other arthropod model species like the spider mite, *Drosophila*, clonal raider ant and red flour beetle (11, 38, 97, 98). Even though we surface-washed the mites and only used specimens with empty alimentary tracts for sequencing, we removed 438 contigs with high bacterial or fugal homology making up approximately 8.5 Mb of contamination (**see**

supplementary Table S1). The final filtered genome assembly was composed of 1182 contigs

with an N_{50} contiguity of 994.5 kb (**Table 1**).

<u>The official gene set and annotation of *Archegozetes*</u>

We generated the official gene set (OGS) for *Archegozetes* by an automated, multi-stage

process combining *ab inito* and evidenced-based (RNAseq reads, transcriptomic data and curated

protein sequences) gene prediction approaches (see "**Materials and Methods**") yielding 23,825

gene models. In comparison to other mites and ticks as well as insects, this is well within the range

of the numbers discovered in other Chelicerata so far (**Figure 2a**). Chelicerates with a large OSG,

however, usually possess larger genomes (1-7 Gb), which suggests that *Archegozetes* may have a

relatively dense distribution of protein-coding genes in its genome. On the other hand, ticks can

have giga-base sized genomes, but only a rather small number of gene models, probably due to

high repetitive content (12, 99-101). Lacking more high-quality genomic resources of mites, it is

thus not clear whether the OGS of *Archegozetes* is the rule, or rather the exception within the

Oribatida.

To compare if *Archegozetes'* OSG is similar to predicted genes of other oribatid mites as

well as Prostigmata and Astigmata, we first clustered genes by ortholog inference (OrthoFinder;

(102)), removed species-specific genes and constructed a presence-absence matrix of orthogroups

to ordinate the data using non-metric multidimension scaling (NMDS, **Figure 2b**). Ordination

revealed that the OGS of *Archegozetes* is well nested with other oribatid mites and clearly

separated from their closest relative the astigmatid mites as well as prostigmatid mites (**Figure**

2b). As a first step in annotating the OSG, we ran KOALA (KEGG Orthology And Links

Annotation) to functionally characterize the genes (103). In total, 10,456 (43.9%) of all genes

received annotation and about two thirds of all genes were assigned either as metabolic genes

(36%) or genes related to genetic information processing (34%), while the remaining genes fell

into different KEGG categories (**Figure 2c**). To further annotate the genome, we followed the

general workflow of funannotate with some modifications (104, see **"Materials and Methods"**).

Overall, we found 15,236 genes (64%) of the OGS with homology to previously published

sequences (**Figure 2d**). For about half of all genes (51%), we were able to assign a full annotation,

4% of all genes only showed homology to bioinformatically predicted proteins of other species,

while 9% of all genes only showed homology to hypothetical proteins (**Figure 2d**). As only a few

high-quality, annotated mite genomes are available and the two-spotted spider mite is the sole

species with any experimentally confirmed gene models, it is not surprising that we were only able

to confidently annotate about 55% of all genes of the OGS (**Figure 2d**).

<u>Orthology and comparative genomics of chelicerates</u>

To further access the protein-coding genes of the mite, we compared the OGS to other

chelicerates. Both concatenated maximum likelihood and coalescent species-tree phylogenomic

approaches based on 1,121 orthologs placed *Archegozetes,* as expected, within the Nothrina (33,

105) with strong support and recovered previously found oribatid clade topologies (**Figure 3a**).

Our analysis placed the Astigmata as a sister group of Oribatida and not nested within oribatids as

suggested based on life-history, chemical defensive secretions, morphology and several molecular

studies (39, 41, 106-115). The relationship of Oribatida and Astigmata has been challenging to

resolve for the past decades and several studies using different set of genes, ultra-conserved

elements or transcriptomic data reconstructed discordant phylogenies, some of which are similar

to ours (22, 109, 110, 112-116). Overall, the Oribatid-Astigmatid relationship remains unresolved

and a broader taxon sampling, especially of more basal Astigmata, will be necessary (22, 39, 111,

114-116). We recovered Trombidiformes (Prostigmata and Sphaerolichida) as sister group of the

Sarcoptiformes (Oribatida and Astigmata) constituting the Acariformes (**Figure 3a**). Neither the

maximum likelihood phylogeny (**Figure 3a**), nor the coalescence-based phylogeny

(**Supplementary Figure S2**) reconstructed the Acari (i.e. Acariformes and Parasitiformes) as a

monophyletic taxon. Even though there is morphological, ultrastructural and molecular evidence

for a biphyletic Acari, aswe recovered here, this relationship and larger-scale chelicerate

relationships remain unclear (9, 22, 113, 116-120).

To further assess the quality and homology of both the genome assembly (**Table 1**) and

the OGS (**Figure 2**), we used the1066 arthropod Benchmarking Universal Single-Copy Ortholog

(BUSCO) genes data set (121). Nearly all BUSCO genes were present in the *Archegozetes*

assembly and OGS (96.2% and 97.3%, respectively; **Figure 3b**). Compared to other genomes

sequenced so far, the *Archegozetes* genome has the highest completeness among oribatid mites

and the OGS completeness is on par to the well curated genomes of other chelicerate species and

Drosophila melanogaster (**Figure 3b**). This result is not surprising because the *Archegozetes*

genome was assembled from long-read and short-read data, while all other oribatid mite genomes

were solely short reads sequenced on older Illumina platforms (18). The fraction of duplicated

BUSCO genes in *Archegozetes* (4%) was similar to that of the spider mite and deer tick (11, 12),

but very low compared to the house spider (**Figure 3c**), whose genome underwent an ancient

whole-genome duplication (96).

Overall, the high quality of both the genome assembly and OGS of *Archegozetes* compared

to those of other oribatid mites, strongly indicates the importance of this genomic resource. We

next categorized all protein models from the OGS by conversation level based on a global

clustering orthology analysis (OrthoFinder; 102) of 23 species (**Figure 3c; supplementary Figure**

S3) representing Acariformes, Parasitiformes, several other chelicerates and the fly *Drosophila*.

As for most other species (2, 122), about a third of all orthogroups was highly conserved (**Figure**

3c) across the arthropods, being either in all species (10%; **Figure 3d**) or is most (22%; **Figure**

3d). Only 1% of all *Archegozetes* orthogroups did not show homology and were species specific

(**Figure 3c and d**). Only a low proportion (**Figure 3c**) of orthogroups was conserved across the

higher taxonomic levels (all <1% in *Archegozetes;* **Figure 3d**), which is in line with previous

studies that included prostigmatid and mesostigmatid mites (13, 15, 17). Interestingly, there was a

large proportion of orthogroups conserved across all Oribatida (43% in *Archegozetes;* **Figure 3d**)

and also about 19% of orthogroups in *Archegozetes* were shared only with other Nothrina (**Figure**

3d). A fairly large percentage of these orthogroups may contain potentially novel genes that await

experimental verification and functional analyses (2, 102, 123). Especially the lack of homology

within the Sarcoptiformes (2-3%; **Figure 3c**) may explain the controversial placement of

Astigmata as a sistergroup of Oribatida that we recovered (**Figure 3a**). This grouping is likely

caused by a long-branch attraction artifact and the sister relationship was incorrectly inferred (109,

112, 114, 115, 120), because orthogroup clustering could not detect enough homology between

oribatids and the Astigmata so far sequenced , which are highly derived. Hence, a broad taxon

sampling of basal astigmatid mite genomes seems necessary to resolve Oribatida-Astigmata

relationship (39, 111-113, 116).

<u>Repeat content analysis and transposable elements (TEs)</u>

For clonal species like *Archegozetes,* reproducing in the absence of recombination, it has

been hypothesized that a reduced efficacy of selection could results in an accumulation of

deleterious mutations and repeats in the genome (124-129). There is, however, no evidence for

such an accumulation in oribatids or other arthropods (18). Generally, we found that most of the

repetitive content in *Archegozetes* could not be classified (57%; **Figure 4a**). The high proportion

of unknown repeats likely corresponds to novel predicted repetitive content, because of limited

repeat annotation of mites in common repeat databases such as RepBase (18). Regarding the two

major classes of repeat content, DNA transposons made up about 32% of total repeats, while only

5% represented retrotransposons (**Figure 4a**). About 6% of total repetitive content comprised

simple and low complexity repeats (**Figure 4a**). Overall, the total repetitive content (32%, **Figure**

4b) seems to be within a normal range for chelicerates and arthropods.

The repeat content found in other oribatid mites was lower (18), but recent studies suggest

that sequencing technology, read depth and assembly quality are paramount to the capacity of

identifying repeat content and TEs (130, 131). Hence, it is very likely the current genomic data for

other Oribatida underestimates the actual total repetitive content. More low-coverage, long-read

sequencing could reduce the assembly fragmentation and likely reveal a higher proportion of

repeats, closer to the actual repetitiveness of oribatid genomes (130).

Different classes of transposable elements (TEs) are characterized by the mechanism they

use to spread within genomes and are known to influence population dynamics differently (131-

133). We therefore analyzed the evolutionary history of TE activity in *Archegozetes* in more detail

(**Figure 4c**). The main TE superfamilies were DNA transposons (**Figure 4a and c**), which seems

to be a common pattern of oribatid mite genomes. For *Archegozetes,* they appear to have

accumulated in the genome for a long time (i.e. they are more divergent from the consensus; (134))

with Tc1/mariner – a superfamily of interspersed repeats DNA transposons (131) – being the most

abundant one (**Figure 4c**). Interestingly, we found an increase in TE activity with 0-3% sequence

divergence range, indicating a recent burst (**Figure 4c**). This burst contained an enrichment of

DNA Mavericks, which are the largest and most complex DNA transposons with homology to

viral proteins (131), but also several of retrotransposons. Among these, is the Long Terminal

Repeat (LTR) gypsy retroelement (**Figure 4c**), which is closely related to retroviruses (131). Like

retroviruses, it encodes genes equivalent to *gag*, *pol* and *env*, but relatively little is known about

how it inserts its DNA into the host genome (135, 136). So far, it is unknown what these TEs do

in *Archegozetes*, but the recent burst in TE abundance might suggest that some changes in the

genome might have happened since the became a laboratory model nearly 30 years (33).

The *Archegozetes* Hox cluster

The Hox genes are a group of highly conserved transcription factor-encoding genes that

are used to pattern the antero-posterior axis in bilaterian metazoans (137, 138). Ancestrally,

arthropods likely had ten Hox genes arranged in a cluster (139). During arthropod development,

the Hox genes specify the identities of the body segments, and mutations in Hox genes usually

result in the transformation of segmental identities (139). The importance of Hox genes in

development of metazoans makes knowledge of their duplication and disappearances important

for understanding their role in the evolution of body plans (139).

Mites largely lack overt, external signs of segmentation, other than the serially arranged

appendages of the prosoma (140). Signs of segmentation in the posterior body tagma, the

opisthosoma, do exist in adult members of Endeostigmata (141). However, these segmental

boundaries are largely present only in the dorsal opisthosoma, making it difficult to assess how

these correspond to the ventral somites (140, 141). Developmental genetic studies of the spider

mite and *Archegozetes* suggest that acariform mites only pattern two segments in the posterior

body region, the opisthosoma, during embryogenesis (11, 79-81). This stands in stark contrast to

other studied chelicerate embryos. For example, during embryogenesis the spider *Parasteatoda*

tepidariorum patterns twelve opisthosomal segments (142) and the opilionid *Phalangium opilio*

patterns seven (143). Furthermore, a member of Parasitiformes, the tick *Rhipicephalus microplus,*

appears to pattern eight opisthosomal segments during embryogenesis (144).

Parallel to the observation of segmental reduction in *T. urticae*, genomic evidence suggests

that this acariform mite has lost two of its Hox genes, *i.e., Hox3* and *abdominal-A (abd-A)* (11).

Interestingly, orthologs of *abd-A* in other studied arthropods pattern the posterior segments as well.

A genomic comparison of arthropod Hox clusters has also shown a correlation between

independent losses of *abd-A* and a reduction in posterior segmentation (145). To investigate

whether the loss of segmentation in *Archegozetes* is also due to an absence in *abd-A*, we annotated

its Hox cluster, paying close attention to the region between the Hox genes *Ultrabithorax* (*Ubx*)

and *Abdominal-B* (*Abd-B*), which is usually where this gene resides in other arthropods (139). Our

results suggest that the *Archegozetes* Hox genes are clustered in a contiguous sequence

(tig00005200_pilon, total size ~7.5 Mbp) in the same order as suggested for the ancestral arthropod

(89). Furthermore, we found no sequences suggestive of an *abd-A* ortholog in *Archegozetes*

(**Figure 5a**). These data also support the findings of a previous PCR survey that retrieved no *abd-*

A ortholog in *Archegozetes* (146). Genomic evidence from the Parasitiformes *Ixodes scapularis*

and *Metaseiulus occidentalis* reveal that these taxa maintain orthologs of all ten Hox genes,

however in *M. occidentalis* these genes are not clustered as they are in *I. scapularis* (12, 13).

Taken together, these observations suggest that the last common ancestor of acariform

mites likely lost its *abdominal-A* gene as well as experiencing a reduction in opisthosomal

segmentation (**Figure 5b**). Alternatively, these shared losses of *abd-A* may be due to convergence

due to similar selective pressures favoring a reduction in body size. The dorsal, external

segmentation of endeostigmatid mites does not necessarily contradict the hypothesis of a loss of

abd-A at the base of the acariform mites. As Hox genes are usually deployed after the genetic

establishment of segments in arthropods (139), the opisthosomal segments in endeostigmatid mites

may still develop in the absence of *abd-A*. However, this hypothesis needs further testing with

observations of segmental gene expression in endeostigmatids as well as additional acariform

species.

<u>Life-stage specific RNA expression patterns</u>

Developmental and gene expression data from *Archegozetes* embryos (**Figure 5 d** and **e**)

have elucidated many of the potential mechanisms driving the morphogenesis of many

developmental peculiarities. These peculiarities include the suppression of the fourth pair of

walking legs during embryogenesis as well as the reduction of opisthosomal segmentation (79-82,

84, 147). In typical acariform mites, embryogenesis ends with the first instar, the prelarva, which

usually remains within the egg chorion, as in *Archegozeetes*. Hatching releases the second instar,

the larva, which is followed by three nymphal instars (proto-, deutero- and tritonymph) and the

adult, for a total of six instars. (148). Thus far, methodological limitations have made it difficult

to examine how mite segmentation and limb development progress throughout these instars.

To this end, we used RNAseq to calculate the transcripts per million (tpm) values of genes

known to be, or suspected to be, involved in limb development and segmentation throughout the

six different instars of *Archegozetes*. Prior to comparing these tpm values, gene orthology was

confirmed *via* phylogenetic analyses (**supplementary Figures S4-S11**; see **Table S2** for

phylogenetic statistics and **Table S3** for tpm values). Regarding the total number of genes

expressed across the different life stages, we found that earlier instars generally expressed a higher

number of genes (**Figure 5c**). While most expressed genes were shared across all instars, more

transcripts were shared between the eggs and the larvae and among all five juvenile instars.

Additionally, we found that earlier instars expressed a larger number of stage-specific genes as

compared to later instars and adults (**Figure 5c**).

Gene expression, SEM and time-lapse data have revealed that the development of the

fourth pair of walking legs in *Archegozetes* is suppressed until after the larval instar (80, 81, 147).

The resulting larva is thus hexapodal (see also embryo in **Figure 5e**), which constitutes a putative

synapomorphy of Acari, if they are monophyletic (9). In arthropods, the development of the limbs

is generally accomplished via the activity of highly conserved regulatory genes, termed the "limb

gap genes." These genes are expressed along their proximo-distal axes to establish the specific

identities of the limb podomeres. The limb gap genes include *extradenticle* (*exd*) and *homothorax*

(*hth*), which act together to specify the proximal limb podomeres, *dachshund* (*dac*), which

specifies the medial podomeres, and *Distal-less* (*Dll*) which specifies the distal-most podomeres.

It was previously shown that the deployment of these genes in the anterior appendages of

Archegozetes, *i.e.*, the chelicerae, pedipalps and first three pairs of walking legs (**Figure 5d** and

e), is similar to that of other chelicerate taxa (82, 142, 149). However, in the anlagen of the fourth

pair of walking legs, only the proximal-specifying genes, *exd* and *hth*, are expressed (82).

Whether the limb gap genes are re-deployed during the transition from the prelarval to

larval instars in order to activate the development of the fourth pair of walking legs remains an

open question. We therefore compared the average tpm values of verified limb gap genes (i.e., Al-

Dll, Al-*Hth*, Al-*exd*, and Al-*dac* (82)) in embryos and at each instar stage (**Figure 5f**). We also

compared the tpm values of the *Archegozetes* orthologs of *Sp6-9* and *optomotor blind*, genes

shown to be involved in limb formation in spiders (150, 151). We hypothesized that limb

development genes would show high expression in the larval stage leading to the development of

the octopodal protonymph. We did observe an increase in the tpm averages of Al-*hth* as well as

Al-*optomotor-blind*, however the aforementioned limb gap gene expression levels were similar between these instars (**Figure 5f**). Taken together, these genes may not be up-regulated for the formation of the fourth pair of walking legs between these two instars.

Chelicerate embryos segment their bodies through a "short/intermediate germ" mechanism, whereby the anterior (prosomal) segments are specified asynchronously (142). This usually occurs well before the sequential addition of posterior segments from a posterior growth zone. Based on neontological and paleontological data, chelicerate arthropods may have ancestrally had an opisthosoma comprised of 12 or more segments (5, 7, 140). Embryonic expression data for the segment polarity genes, those genes that delineate the boundaries of the final body segments, have shown that in most studied chelicerate embryos opisthosomal segments are delineated during embryogenesis (140, 142). However, as discussed above, expression data in *Archegozetes* embryos suggest that only two opisthosomal segments are patterned during embryogenesis (80, 81); this indicates that mites have significantly reduced their number of opisthosomal segments either by loss or by fusion. Further complicating this is the observation that many mites add segments as they progress through the larval instars, a phenomenon known as anamorphic growth (140).

To determine by what genetic process *Archegozetes* may add segments during post-embryonic ontogeny, we assessed the expression of known chelicerate and arthropod segmentation genes in each instar transcriptome (**Figure 5f**) (142). We observed an up-regulation of the segmentation genes *hedgehog* and *engrailed* in the larvae, as well as the slight up-regulation of *patched* and *pax3/7*. Furthermore, the segmentation gene *wingless* was slightly up-regulated in the protonymph, as well as a slight up-regulation of *hedgehog* in the tritonymph. Lastly, we found that transcripts of the genes *pax3/7* and *runt* were up-regulated in adults. These results suggest that

Archegozetes does pattern body segments during the progression through the it's instars similar to other Chelicerata.

Another peculiarity of *Archegozetes* is that these mites lack eyes (see more details below). Eye loss has been documented in other arachnid clades, including independently in other members of Acari (10, 152), and it has been recently demonstrated that a species of whip spider has reduced its eyes by reducing the expression of retinal determination genes that are shared throughout arthropods (153). We sought to determine if eye loss in *Archegozetes* also is associated with the reduced expression of these genes (see also analysis of photoreceptor genes below). The genes, which have been shown to be expressed in the developing eyes of spiders and whip scorpions, include *Pax-6*, *six1/sine oculis* (*so*), *eyes absent* (*eya*), *Eyegone*, *Six3/Optix*, and *atonal* (153-155). We also followed the expression of Al-*orthodenticle*, a gene previously shown to be expressed in the ocular segment of *Archegozetes* (147). Surprisingly, all of these genes, excluding the *Pax-6* isoform A and *eyegone*, are indeed expressed during embryogenesis (**Figure 5f**). Aside from the larval expression of the *Pax-6* isoform A during the larval stage, these eye-development genes remain quiescent until the adult stage, where all but *Pax-6* isoform A, *six3* and *atonal* are up-regulated (**Figure 5f**). These results are exceedingly surprising, given the conserved role of genes in retinal patterning. They suggest a novel role for these genes, or alternatively, these expression patterns could be the result of early expression of a retinal determination pathway followed by negative regulation by other genes to suppress eye development.

Photoreceptor and chemosensory system of *Archegozetes longisetosus*

Unlike insects and crustaceans, chelicerates do not have compounds eyes – with horseshoe crab being an exception. Generally, mites are eyeless or possess one or two pairs of simple ocelli (156-160). Ocelli are common in Prostigmata and Endeostigmata, among Acariformes, as well

Opilioacarida – the most likely sister group to the Parasitiformes – but are absent in most Oribatida,

Astigmata, Mesostigmata and ticks (10, 161-163). This suggests that the presence of eyes might

be an ancestral condition for both Acariformes and Parasitiformes, while more derived mites rely

largely on chemical communication systems (156).

In oribatid mites, detailed morphological and ultrastructural investigations have suggested

that setiform sensilla are the most obvious sensory structures (**Figure 6a**) (10, 156, 164). The

trichobothria are very complex, highly modified (e.g. filiform, ciliate, pectinate, variously

thickened or clubbed) no-pore setae which are anchored in a cup-like base and likely serve as

mechanosensory structures. In contrast, the setal shafts of solenidia and eupathidia (**Figure 6a**)

both possess pores (10, 156, 164). Solenidia have transverse rows of small pores visible under a

light microscope and likely function in olfaction, while the eupathidia have one or several terminal

pores and likely are used as contact/gustatory sensilla (**Figure 6a**) (156, 164). Previous work

demonstrated that oribatid mites indeed use olfactory signals in the context of chemical

communication and food selection (47-49, 58, 64, 86).

Interestingly, detailed morphological and ultrastructural studies showed that light-sensitive

organs exist in some Palaeosomata and Enarthronota (probably true eyes) as well as in

Brachypylina (the secondary lenticulus), representing lower and highly derived oribatid mites,

respectively (156, 160-162). *Archegozetes* and most other oribatids, however, are eyeless, yet there

is scattered experimental and some anecdotal evidence that even these mites show some response

to light and seem to avoid it ('negative phototropism' or 'negative phototaxis') (10, 165-167).

Hence, we mined the genome of *Archegozetes* for potential photoreceptor genes and found two

genes of the *all-trans retinal perosin* class and one gene related to spider mite *rhodopsin-7-like*

gene (**Figure 6b**). *Perosin-like* genes are also present in other eyeless ticks. In jumping spiders it

encodes for nonvisual, photosensitive pigments, while *rhodopsin-7* may be involved in basic insect

circadian photoreception (168-173). Taken together, this might suggest that eyeless species like

Archegozetes use *perosin-* and *rhodopsin-7-like* genes for reproductive and diapause behaviors, or

to maintain their circadian rhythm, as well as negative phototaxis.

However, the main sensory modality soil mites use is chemical communication via

olfaction (10, 49, 58, 64, 156, 164, 173). In contrast to insects, but similar to crustaceans and

Myriapoda, mites do not have the full repertoire of chemosensory classes, they are missing odorant

receptors and odorant-binding proteins (**Table 2**) (13, 15, 17, 30, 31, 174-176). Although

chemosensory protein (CSP) encoding genes are absent in most mite genomes, we identified one

gene encoding for such a protein in *Archegozetes* and one CSP has been previously found in the

deer tick (Table 2). Hence, *Archegozetes* should primarily rely on gustatory receptors (GRs) and

ionotropic receptors (IRs). Both the number of GRs (68 genes; **Figure 6d**) and IRs (3 genes;

Figure 6c) was very well within the range of most mites and ticks and there was no evidence for

any massive chemoreceptor expansion like in the spider mite (**Table 2**) (177). This was surprising

because *Archegozetes*, like other acariform mites have many multiporous solenidia, present on all

legs and the palp, but appear to only have a limited number of chemoreceptors.

Canonical ionotropic glutamate receptors (iGluRs) are glutamate-gated ion channels with

no direct role in chemosensation, which come in two major subtypes: either NMDA iGluRs which

are sensitive to N-methyl-D-aspartic acid (NMDA) or non-NMDA iGluRs. The latter group – at

least in *Drosophila* – seems to have essential functions in synaptic transmission in the nervous

system and have been associated with sleep and vision (175-179). None of the IRs we found in the

Archegozetes genome belonged to the NMDA iGluRs and most were classified as non-NMDA

iGluRs (**Figure 6c**). Nothing is known about their functions in mites. It is, however, likely that

they perform similar tasks in synaptic transmission in the brain and musculature. In *Drosophila* a

specific set of chemosensory IRs, which do not bind glutamate, respond to acids and amines

(*IR25a*), but also to temperature (*IR21a*, *IR93a*). For *Archegozetes* we found 3 IRs, like *IR21a* and

IR93a of *Drosophila*, which fell into the antenna/1st leg IRs category (**Table 2**; **Figure 6c**) (180-

182). This is consistent with an assumed limited contribution of IRs to the perception of chemical

cues. Furthermore, it is so far unclear whether these IRs are expressed in the first pair of legs

(**Figure 6a** and **c**) in *Archegozetes*, but similar genes seem to be expressed in the legs of other mite

species, which could suggest a similar function as in the fruit fly.

GRs are multifunctional proteins and at least in insects they are responsible for the

perception of taste, heat or volatile molecules (183). In *Archegozetes* we found 68 GRs, over half

of which belonged to a species-specific expansion of the GR gene family (**Figure 6d**). Generally,

it is unclear if GRs in *Archegozetes* and other mites have similar functions as in insects, but the

GR gene family is heavily expanded in many acariform mites and also is present in ticks (**Table**

2), suggesting an important biological role (12, 13, 15, 17, 101, 177). This is supported by

experimental evidence which suggested that ticks and other mites, including *Archegozetes,* use

chemical cues to find their host, communicate or discriminate food (12, 49, 58, 64, 101, 184-186).

In general, not much is known about the nervous and sensory system of oribatid mites, or

about sensory integration or the neuronal bases of their behavior (40, 156, 164). Modern methods

like Synchrotron X-ray microtomography (SRμCT) recently made it possible to investigate the

organization and development of the nervous systems of oribatid mites (**Figure 6e;** (187)) and

here we provide the first genomic resource for the investigation of the photo- and chemosensory

systems of Oribatida (**Figure 6b-d**).

Horizontal gene transfer event sheds light on oribatid feeding biology

Horizontal gene transfer (HGT) is common among mites and other soil organisms (11, 17,

74-77). In some cases, genes that had been horizontally transferred now have pivotal biological

functions. For instance, terpene and carotenoid biosynthesis genes in trombidiid and tetranychid

mites, respectively, are found nowhere else in the animal kingdom (17, 188). Yet they show high

homology with bacterial (terpene synthase) or fungal (carotenoid cyclase/synthase/desaturase)

genes, suggesting horizontal gene transfer from microbial donors (17, 188). At least the carotenoid

biosynthesis genes in spider mites still code for functional enzymes and equip these phytophages

with the ability to *de novo* synthesize carotenoids, which can induce diapause in these animals

(188).

Soil microarthropods like collembolans show numbers of horizontally transferred genes

that are among the highest found in Metazoan genomes, exceeded only by nematodes living in

decaying organic matter (76, 77, 189). Interestingly, many HGT genes found in collembolans are

involved in carbohydrate metabolism and were especially enriched for enzyme families like

glycoside hydrolases, carbohydrate esterases or glycosyltransferases (76, 77). All three enzyme

families are involved in the degradation of plant and fungal cell walls (190, 191). Hence, it has

been hypothesized that cell-wall degrading enzymes acquired by HGT are beneficial for soil

organisms as it allowed such animals to access important food source in a habitat that is highly

biased towards polysaccharide-rich resources (76, 77, 192, 193).

To assess the degree of HGT in *Archegozetes* we first used blobtools (v1.0) (194) to

generate a GC proportion vs read coverage plot of our genome assembly, in order to remove

scaffolds of bacterial origin (**Figure 7a**; 438 contigs of ~ 8.5 Mb of contamination). Of the

remaining scaffolds, candidate HGTs were identified using the Alien Index (195, 196), where

HGTs are those genes with blast homology (bit score) closer to non-metazoan than metazoan

sequences (**supplementary Table S4**). We further filtered these HGT candidates to remove those

that overlapped predicted repeats by $\geq$ 50%, resulting in 617 genes. As HGT become integrated

into the host genome, they begin to mirror features of the host genome, including changes in GC

content and introduction of introns (197). Comparing the GC content of the HGT candidates

showed two distinct peaks, one at 54.3% and the other at 35.2%, slightly higher than the remaining

Archegozetes genes, GC content of 31.5% (**Figure 7b**). Of the 407 HGT genes that shared similar

GC content to the host genome, 73.5% had at least one intron (**Table S4**). In a final step, we used

the gene expression data (RNAseq) to filter the list of all putative HGT genes and only retained

candidates that were expressed in any life stage of *Archegozetes* (*n*= 298 HGT genes).

512   The majority of HGT candidates were of bacterial origin (75.2%), followed by genes likely

acquired from fungi (13.4%), while transfer from Archaea, plants, virus and other sources was

comparatively low (**Figure 7d**). This composition of HGT taxonomic origin is different from

genes found in collembolans, which appear to have acquired more genes of fungal and protist

origin (76, 77, 192). Subsequently, we performed an over-representation analysis of GO terms

associated with these genes. We found an over-representation of genes with GO terms related to

methyl transfer reactions and breaking glycosidic bonds (molecular function; **Figure 7c**) as well

as carbohydrate metabolism, among others (biological process; **Figure 7c**) providing a first line of

evidence that *Archegozetes* possess HGT related to plant- and fungal cell wall degradation similar

to springtails.

522   As mentioned previously, oribatid mites are among the few Chelicerata that ingest solid

food and are primary- and secondary decomposers feeding on dead plant material and fungi (9, 40,

55-57, 63). It was argued for decades that the enzymes necessary to break down these

polysaccharide-rich resources originate from the mite's gut microbes (45, 65-67, 71, 198, 199).

Microbes might be mixed with the food in the ventriculus and digest it while passing through the

alimentary tract as food boli enclosed in a peritrophic membrane (see **Figure 7f** for an example)

(198, 199). However, screening the HGT candidate list for potential cell-wall degrading enzymes

and mapping their overall and life-stage specific expression in *Archegozetes* using the RNAseq

reads, revealed at least seven HGT genes related to polysaccharide breakdown (**Figure 7g**). We

found that specifically members of the *glycoside hydrolases family 48* and *cellulose-binding*

domain genes showed high expression in most life stages - the egg being an obvious exception

(**Figure 7g**). Moreover, the majority of these genes were flanked by a predicted metazoan gene,

suggesting host transcriptional regulation (**Table S4**).

In a last step we blasted the highly expressed HGT candidates (**Figure 7g**) against the non-

redundant protein sequence database, aligned the sequences with the highest alignment score.

Eventually, we performed a phylogenetic maximum likelihood analysis. For the highest expressed

HGT related to cell-wall-degrading enzymes (*glycoside hydrolases family 48 gene II*), we

recovered that the *Archegozetes* sequences was well nested within a clade of *GH 48* sequences

from herbivores beetles (72), which appear to be related to similar genes from various

Streptomyces (**Figure 7e**) and we reconstructed similar phylogenies for other highly expressed

HGT candidates (**supplementary Figure S12**). All the sequences of beetle *glycoside hydrolases*

family 48 members (**Figure 7e**) were included in recent studies arguing for a convergent horizontal

transfer of bacterial and fungal genes that enabled the digestion of lignocellulose from plant cell

walls in herbivores beetles (72, 200). They showed that phytophagous beetles likely acquired all

genes of the *GH 48 family* from Actinobacteria (including *Streptomyces*) (72) and our phylogenetic

analysis (**Figure 7e**) revealed the same pattern as well as a highly similar tree topology (compare

to Fig 3B in (72)).

Overall, our findings indicate that genes encoding for enzymes in *Archegozetes* capable of

degrading plant and fungal cell walls were likely horizontally transferred from bacteria (likely

Streptomyces). Bacterial symbionts and commensal living in the mites' gut are still likely to

contribute to the breakdown of food (**Figure 7f**). Yet, the high expression of genes encoding cell-

wall degrading enzymes (**Figure 7g**) as well as the evolutionary analyses of such genes (**Figure**

7e) suggest that *Archegozetes* – and potentially many other oribatid mites – are able to exploit

polysaccharide-rich resources like dead plant material or chitinous fungi without microbial aid.

Enzymological and microscopical investigation of *Archegozetes* have suggested that certain

digestive enzymes (chitinase and cellulase) are only active when the mites consume a particular

type of food (e.g. algae, fungi or filter paper) (71). These results were interpreted as evidence that

these enzymes are directly derived from the consumed food source (71). By contrast, we argue

that this instead confirms our findings of HGT: upon consumption of food containing either chitin

or cellulose, gene expression of polysaccharide-degrading enzymes starts, and proteins can readily

be detected. Further enzymological studies have placed oribatid mites in feeding guilds based on

carbohydrase activity and also found highly similar enzyme activity between samples of mites

from different times and locations (67, 68, 201-203). Future functional studies can disentangle the

contribution of the host and microbes to cell wall digestion and novel metabolic roles of the HGTs

identified here.

<u>Biosynthesis of monoterpenes – a common chemical defense compound class across oribatid mite</u>

Oribatid and astigmatid mites are characterized by a highly diverse spectrum of natural

compounds that are produced by and stored in so-called oil glands (for an example see **Figure 8a**)

(30, 88, 204). These paired glands are located in the opisthosoma (i.e. the posterior part of

chelicerate arthropods, analogous to the abdomen of insects) and are composed of a single-cell

layer invagination of the cuticle (**Figure 8f**). As previously mentioned, mites use chemicals produced by these glands to protect themselves against environmental antagonists (predators or microbes) or use them as pheromones (30, 47-49, 52, 86, 89). The monoterpene aldehyde citral – a stereoisomeric mixture of geranial ((E)-3,7-dimethylocta-2,6-dienal) and neral ((Z)-3,7-dimethylocta-2,6-dienal) – and its derivatives are widely detected compounds in glandular secretions of oribatids and astigmatids (41, 107, 184, 205-209). These monoterpenes have been called "astigmatid compounds" (41) as they characterize the biochemical evolutionary lineage of major oribatid mite taxa (Mixonomata and Desmonomata) and almost all investigated astigmatid mites (30, 41, 118, 184, 208).

The chemical cocktail released by *Archegozetes* consists of a blend of 10 compounds (**Figure 8a**) including two terpenes (approx. 45%) – neral and neryl formate – six hydrocarbons (approx. 15%) and two aromatic compounds (approx. 40%) (206, 210). The hydrocarbons likely serve as solvents, while the terpenes and aromatics are bioactive compounds used in chemical alarm and defense (47-49, 206). Recently, it was shown that *Archegozetes* synthesizes the two aromatic compounds using a polyketide-like head-to-tail condensation of (poly)-β-carbonyls *via* a horizontally acquired putative polyketide synthetase (27). Studies in Astigmata found that the monoterpenes of these mites appeared to be made *de novo* from (poly)-β-carbonyls as well and one study identified a novel geraniol dehydrogenases (GeDH), unrelated to those of bacteria, in *Carpoglyphus lactis* (211-213). To learn about the biosynthesis of astigmatid compounds in *Archegozetes* and demonstrate the mite's applicability as research model for biochemical pathway evolution, we used the novel genomic resources presented in this study.

First, we delineated the basic biochemical reactions likely to happen in the *Archegozetes* gland through a stable-isotope labeling experiment. We supplemented the diet of the mite with

food containing 25% heavy $^{13}C_6$ D-glucose and 10% antibiotics (a combination of three different

antibiotics was fed, because this mixture is able to eliminate nearly all qPCR and FISH detectable

bacteria found on the food and in the alimentary tract (27)). To examine the incorporation of heavy

$^{13}C_6$ D-glucose and its metabolic products into neral (**Figure 8b**) and neryl formate (**Figure 8c**),

we compared selected fragment ions (M^+ and M^+-46, respectively) using single ion mass

spectrometry. Both neral and neryl formate showed consistent enrichment in their M^+ to $[M+4]^+$

and $[M-46]^+$ to $[M-46+4]^+$-ion series, indicating that *Archegozetes* used glycolysis breakdown

products of $^{13}C_6$ D-glucose for the biosynthesis of their monoterpenes. We then used the OGS

mapped to KEGG metabolic pathways (214) to reconstruct the backbone synthesis of terpenes in

Archegozetes (**Figure 8d**). We found mapped mite genes, which suggest that *Archegozetes*

synthesizes geranyl pyrophosphate (GPP) – the input substrate for further monoterpene synthesis

– via the mevalonate pathway using the Mevalonate-5P to Isopentenyl-PP route (**Figure 8d**). The

Mevalonate-5P pathway is used in most higher eukaryotes as compared to the Mevalonate-3P

pathway in Archaea and the MEP/DOXP pathway in bacteria, some plants and apicomplexan

prostists (215-220). This likely excludes any horizontal gene transfer of mevalonate pathway genes

as *Archegozetes* uses enzymes similar to those of other animals.

The biosynthesis of monoterpenes not only depends on very widespread enzymes, but also

requires more specific enzymes downstream of GPP (218-220). For instance, *Carpoglyphus lactis*

expresses a unique geraniol dehydrogenase (GeDH) – catalyzing the oxidation of geraniol to

geranial – different from all previously characterized geraniol-related and alcohol dehydrogenases

(ADHs) of animals and plants (212). We used the functionally validated Carpoglyphus-GeDH

(212), blasted its sequence against the *Archegozetes* OGS and found a homologous sequence. We

used both mite sequences in an alignment with plant, fungal and bacterial GeDHs and animal

ADHs and constructed a maximum likelihood phylogeny (**Figure 8e**). Similar to the previous analysis including only Carpoglyphus-GeDH, we found that the *Al-GeDH* represent a new class of geraniol dehydrogenases different from those in plants, fungi or bacteria and not nested within animal ADHs (**Figure 8e**). This is why we hypothesize that *Al-GeDH* is a novel expansion of the geraniol dehydrogenases gene family and has not been acquired by horizontal gene transfer, like other biosynthesis and digestive enzymes in *Archegozetes* (**Figure 7**; (27)).

Based on our mass spectrometry data of stable isotopes and genomic analysis, we propose that the following biochemical pathway leading to monoterpenes is of oribatid mites (**Figure 8f and g**): geraniol is likely to be synthesized from GPP – the universal precursor of all monoterpenes – either enzymatically by a geraniol synthase (GES) or a diphosphate phosphatase (DPP), but possibly also endogenously by dephosphorylation of GPP (221-224). For *Archegozetes*, we could not find any *GES* or specific *DPP* in the OGS, thus geraniol might be formed from GPP *via* endogenous dephosphorylation, but further research is required to verify or falsify this hypothesis. Subsequently, geraniol is oxidized to geranial by the pervious described *GeDH* (**Figure 8e**) and readily isomerized to neral. Trace amounts of geranial have been found in *Archegozetes* and it is common among other oribatid and astigmatid mites, supporting this idea (107, 184, 209, 225). Also, there is no evidence that geraniol is cconverted into nerol, or that neral is formed directly via oxidation of nerol (211-213). The most parsimonious explanation for neryl formate synthesis would be an esterification of the corresponding terpene alcohol nerol. There is, however, no evidence of nerol in the traces of any oribatid or astigmatid mite species (30, 88, 184). Aliphatic non-terpene formats in Astigmata are synthesized by dehomologation and generation of a one-carbon–shorter primary alcohol from an aldehyde via hydrolysis of formate in a biological Baeyer–Villiger oxidation catalyzed by a novel, uncharacterized enzyme (226). A similar reaction to

641   synthesize terpene formates is unlikely, as the terpenoid backbone would be shortened by one-

642   carbon and this does not happen in any possible scenario. The discovery of this Baeyer–Villiger

643   oxidation mechanism, however, highlights the probability that there are many very unusual

644   reactions that remain to be discovered in oribatid mites (27).

Conclusion

The integrated genomic and transcriptomic resources presented here for *Archegozetes longisetosus* allowed a number of insights into the molecular evolution and basic biology of decomposer soil mites. Our analysis of an oribatid mite genome also provides the foundation for experimental studies building on the long history of *Archegozetes'* as a chelicerate model organism, which now enters the molecular genetics era (23, 33, 34, 92). This includes the study of biochemical pathways, biochemistry, neuroethological bases of food searching behavior, and environmental impacts on genomes of complex, clonal organisms.

Our evolutionary comparisons across the Chelicerata revealed interesting patterns of genome evolution and how horizontal gene transfer might have shaped the feeding mode of soil mites. We also showed how oribatid glandular biology and chemical ecology are reflected in the genome. The community of researchers studying the fundamental biology of oribatid and other free-living, non-parasitic mites is growing. We think that providing these genomic and transcriptomic resources can foster a community effort to eventually transform basic research on these mites into a modern, molecular discipline.

Key priorities for a future community research effort include i) sequencing organ-specific transcriptomic data, ii) developing tools for genetic interrogation (RNAi or CRISPR/CAS9), iii) establishing reporter linages with germ-line stable modifications (e.g. GAL4/UAS misexpression systems), iv) constructing an whole-animal single-cell RNAseq expression atlas, and v) gathering more genomic data to improve the genome assembly. Please do not hesitate to contact the corresponding author, if you want to start your own culture of *Archegozetes*. He will be happy to provide you with starter specimens and share rearing protocols with you.

Materials and Methods

Mite husbandry

The lineage 'ran' (33) of the pantropical, parthenogenetic oribatid mite *Archegozetes longisetosus* was used in this study. Stock cultures were established in 2015 from an already existing line and fed with wheat grass (*Triticum* sp.) powder from Naturya. Cultures were maintained at 20-24°C and 90% relative humidity. Sterilized water and 3-5 mg wheat grass were provided three times each week.

DNA extraction and Illumina sequencing

For the short-read library, DNA was extracted from ~200 mites that were taken from the stock culture, starved for 24 h to avoid possible contamination from food in the gut, subsequently washed with 1% SDS for 10 sec. For extraction of living specimens, we used the Quick-DNA Miniprep Plus Kit (Zymo Research) according to the manufacturer's protocol. Amounts and quality of DNA were accessed with Qubit dsDNA HS Kit (ThermoFisher) and with NanoDrop One (ThermoFisher) with target OD 260/280 and OD 260/230 ratios of 1.8 and 2.0-2.2, respectively. Extracted DNA was shipped to Omega Bioservices (Norcross, GA, USA) on dry ice for library preparation and sequencing. DNA library preparation followed the KAPA HyperPrep Kit (Roche) protocol (150 bp insert size) and 200 million reads were sequenced as 150bp paired-end on a HighSeq4000 (Illumina) platform.

High-molecular weight DNA isolation and Nanopore sequencing

Genomic DNA was isolated from ~300-500 mites starved for 24 h using QIAGEN Blood & Cell Culture DNA Mini Kit. Briefly, mites were flash frozen in liquid nitrogen and homogenized with a pestle in 1 ml of buffer G2 supplemented with RNase A and Proteinase K at final

concentrations of 200 ng/µl and 1 µg/µl, respectively. Lysates were incubated at 50°C for 2 h,

cleared by centrifugation at 5 krpm for 5 min at room temperature and applied to Genomic tip

G/20 equilibrated with buffer QBT. Columns were washed with 4 ml of buffer QC and genomic

DNA was eluted with 2 ml of buffer QF. DNA was precipitated with isopropanol, washed with

70% EtOH and resuspended in 50 µl of buffer EB. DNA was quantified with Qubit dsDNA HS

Kit (ThermoFisher) and the absence of contaminants was confirmed with NanoDrop One

(ThermoFisher) with target OD 260/280 and OD 260/230 ratios of 1.8 and 2.0-2.2, respectively.

DNA integrity was assessed using Genomic DNA ScreenTape kit for TapeStation (Agilent

Technologies).

Libraries for nanopore sequencing were prepared from 1 µg of genomic DNA using 1D

Genomic DNA by Ligation Kit (Oxford Nanopore) following manufacturer's instructions. Briefly,

unfragmented DNA was repaired and dA tailed with a combination of NEBNext FFPE Repair Mix

(New England Biolabs) and NEBNext End repair/dA-tailing Module (New England Biolabs).

DNA fragments were purified with Agencourt AMPure XP beads (Beckman Coulter) and Oxford

Nanopore sequencing adapters were ligated using NEBNext Quick T4 DNA Ligase (New England

Biolabs). Following AMPure XP bead cleanup, ~500 ng of the library was combined with 37.5 µL

of SQB sequencing buffer and 25.5 µl of loading beads in the final volume of 75 µl and loaded on

a MinION Spot-ON Flow Cell version R9.4 (Oxford Nanopore). Two flow cells were run on

MinION device controlled by MinKNOW software version 3.1.13 for 48 hours each with local

basecalling turned off generating 9.7 and 5.1 GB of sequence data. Post run basecalling was

performed with Guppy Basecalling Software, version 3.4.5 (Oxford Nanopore). After filtering low

quality reads (Q<7), the combined output of the two runs was 13.69 GB and 4.7 million reads.

Genome assembly and contamination filtering

Read quality was assessed using FastQC v0.11.8 (227). Illumina adapters, low-quality

nucleotide bases (phred score below 15) from the 3' and 5' ends and reads shorter than 50 bp were

removed using cutadapt v1.18 (228). From the filtered reads, *in silico* genome size estimates were

calculated using *k-mer* based tools kmergenie v.1.7048 (229), GenomeScope v1.0 (230), and

findGSE v0.1.0 R package (231). The latter two required a *k-mer* histogram computed by jellyfish

v2.2.10 (232) with *k-mer* size of 21. The genome was assembled using 4.7 million long reads from

two MinION runs (60x coverage) using Canu v1.8 with default settings and setting the expected

genome size to 200 Mb (233). To improve assembly quality, paired end Illumina reads were

mapped to the genome with BWA aligner (234) using BWA-MEM algorithm and polished with

Pilon v. 1.23 with '—changes' and '--fix all' options (235). Assembled contigs identified as

bacterial and fungal contaminants based on divergent GC content from most *Archegozetes* contigs,

high coverage and blast homology to the nt database (downloaded February 2019, Evalue 1e^{-25})

were removed using Blobtools v1.0 (194).

Identification, classification and masking of repetitive element

Repetitive elements in the genome *Archegozetes* were identified using a species-specific

library generated with RepeatModeler v 1.0.11 (236, 237) and MITE tracker (132) and annotated

by RepeatClassifier, a utility of the RepeatModeler software that uses the RepBase database

(version Dfam_Consensus-20181026). Unclassified repeat families from both programs were run

through CENSOR v 4.2.29 (238) executable *censor.ncbi* against the invertebrate library v 19.03

to provide further annotation. Predicted repeats were removed if they had significant blast

homology (E-value 1e^{-5}) to genuine proteins in the NCBI nr database and/or a local database of

arthropod genomes (*Drosophila melanogaster, Tribolium castaneum, Tetranychus urticae,*

Leptotrombidium deliense, Dinothrombium tinctorium, Sarcoptes scabiei, Euroglyphus maynei,

Galendromus occidentalis, Dermatophagoides pteronyssinus). Unclassified repeats with blast

homology to known TEs were retained whereas those with no blast homology were removed (239).

The remaining repeat families were combined with the Arthropoda sequences in RepBase and

clustered using vsearch v 2.7.1 (--iddef 1 --id 0.8 --strand both; (240)). The filtered repeat library

was used to soft mask the *A. longisetosus* using RepeatMasker v 4.07 (241).A summary of the

masked repeat content was generated using the "buildSummary.pl" script, the Kimura sequence

divergence calculated using the "calcDivergenceFromAlign.pl" script and the repeat landscape

visualized using the "createRepeatLandscape.pl" script, all utilities of RepeatMasker.

Gene prediction and annotation

Both *ab inito* and reference-based tools were used for gene prediction using modified steps

of the funannotate pipeline (104). The *ab inito* tool GeneMark-ES v4.33 (242) was used along with

reference based tools BRAKER v2.1.2 (243) using RNAseq reads discussed below and PASA v

2.3.3 (244) using genome-guided transcriptome assembly from Trinity described below. Lastly,

Tetranychus urticae gene models from the NCBI database (GCF_000239435.1) were aligned to

the contigs using GeMoMa (245). All gene predictions were combined in EvidenceModeler (244)

with the following weights: GeMoMa =1, PASA = 10, other BRAKER = 1, and GeneMark = 1.

Predicted tRNAs using tRNAscan-SE v 2.0.3 (246) were combined with the gene predictions in

the final gene feature format (GFF) file and filtered for overlap using bedtools (247) *intersect* tool

(247).

The predicted genes were searched against the NCBI nr (February 2019) (248), SwissProt

(February 2019) (249), a custom-made Chelicerata database including genomes of *Tetranychus*

urticae, Leptotrombidium deliense, Dinothrombium tinctorium, Sarcoptes scabiei, Euroglyphus

maynei, Galendromus occidentalis, Metaseiulus occidentalis, Dermatophagoides pteronyssinus,

Trichonephila clavipes, Stegodyphus mimosarum, Centruroides sculpturatus, Ixodes scapularis

and *Parasteatoda tepidariorum* (all downloaded Feb 2019), PFAM (v 32, August 2018) (250),

merops (v 12, October 2017) (251) and CAZY (v 7, August 2018) (252) databases. The results of

the hmm-based (253) PFAM and CAZY searches were filtered using cath-tools v 0.16.2

(https://cath-tools.readthedocs.io/en/; E-value $1e^{-5}$) and the blast-based searches were filtered by

the top hit (E-value $1e^{-5}$ threshold). Predicted genes were also assigned to orthologous groups using

eggNOG-mapper (254). Gene annotation was prioritized by the SwissProt hit if the E-value < $1e^{-10}$

followed by NCBI annotation, the custom Chelicerata database and if no homology was

recovered, then the gene was annotated as, "hypothetical protein". Final annotation was added to

the GFF file using GAG (255).

<u>Analysis of the official gene set (OGS)</u>

To allow the OGS to be used as resources for functional studies, we assigned functional

categories based on Gene Ontology (GO) and the Kyoto Encyclopedia of Genes and Genomes

(KEGG) (256, 257). GO terms for the respective genes models of the OGS were assigned based

on the gene id with highest homology from the SwissProt database or NCBI nr database (248,

249). A custom database of GO terms was created with makeOrgPackage function in the R package

AnnotationForge v1.26.0 (258). Over-representation analysis of GO terms was tested using the

enrichGO function in the R package clusterProfiler v3.12.0 (259) with a hypergeometric

distribution and a Fisher's Exact test. P-values were adjusted for multiple comparisons using false

discovery rate correction (260). Resulting enriched GO terms were processed with GO slim (261)

and the final list of over represented GO terms was used to plot the number of genes in a respective

category.

KEGG orthology terms were assigned from single-directional best hit BLAST searches of

each gene model on the KEGG Automatic Annotation Server (262). Additionally, we ran

GhostKOALA (103) (GHOSTX searches for KEGG Orthology And Links Annotation) to obtain

KEGG orthology terms. Compared to conventional BLAST searches, GhostKOALA is about 100

times more efficient than BLAST to remote homologs by using suffix arrays (263).

<u>Orthology and Phylogenomic analyses</u>

Orthologs of *A. longisetosus*, other species within Acari, Chelicerata and the fruit fly

Drosophila were identified using OrthoFinder v 2.3.3 (-M msa –A mafft –T fasttree; (102)). Prior

to running OrthoFinder, isoform variants were removed from the gene predictions using CD-Hit

(264). Trees of orthogroups with at least 80% of taxa present (n= 4,553) were constructed using

fasttree v 2.1.10 (265), trimmed with TrimAl v 1.4.1 (-keepheader -fasta –gappyout; (266)) and

paralogs pruned using phylotreepruner v 1.0 (min_number_of_taxa =18, bootstrap_cutoff= 0.7,

longest sequence for a given orthogroup=u; (267)). Alignments shorter than 100 amino acids were

removed, leaving 1,121 orthogroups.

For the maximum likelihood analysis, the trimmed and pruned alignments were

concatenated into a supermatrix using FasConCat v1.04 (268) composed of 377,532 amino acids

and the best substitution models determined using PartitionFinder v 2.1.1 (269). The maximum

likelihood consensus phylogeny from the supermatrix and partition scheme was constructed using

IQ-tree and 1,000 ultrafast bootstrap replicates (270). For the coalescence species tree

reconstruction, gene trees were generated using IQ-tree v 1.6.12 on the trimmed alignments of the

1,121 filtered orthogroups and processed using ASTRAL v 5.6.3 (271). Branch lengths are

presented in coalescent units (differences in the 1,121 gene trees) and the node values reflect the

local posterior probabilities.

    RNA sequencing and transcriptome assembly

For RNA extraction, about 200 mites of all life stages were taken from stock culture and

subsequently washed with 1% SDS for 10 s. RNA was extracted from living specimens using the

Quick-RNA MiniPrep Kit (Zymo Research) according to the manufacturer's protocol. Quantity

and quality of RNA were accessed using a Qubit fluorometer and NanoDrop One (Thermo Fisher

Scientific), respectively.

Extracted RNA was shipped to Omega Bioservices (Norcross, GA, USA) on dry ice for

library preparation and sequencing. Whole animal RNA was used for poly-A selection, cDNA

synthesis and library preparation following the Illumina TruSeq mRNA Stranded Kit protocol. The

library was sequenced with 100 million 150 bp paired-end on a HighSeq4000 platform. For the

genome-guided assembly of the transcriptome a bam-file was created from the genome using

STAR (272). RNAseq reads were *in silico* normalized and subsequently used together with the

bam-file to assemble the transcripts using Trinity v2.8.4 (273, 274), yielding an assembly with a

total length of 162.8 Mbp, an N50= 2994 bp and a BUSCO score (121) of C:96.3%

[S:36.5%,D:59.8%], F:1.3%, M:2.4%.

Life-stage specific RNAseq

For life-stage specific RNAseq, we collected 15 specimens per life stage from the stock

culture that were split into three replicates of five individuals. Whole animals (for all stages but

eggs) were flash frozen in 50 µl TRIzol using a mixture of dry ice and ethanol (100%) and stored

at −80°. RNA was extracted using a combination of the TRIzol RNA isolation protocol (Life

Technologies) and RNeasy Mini Kit (Qiagen) (275). The TRIzol protocol was used for initial steps

up to and including the chloroform extraction. Following tissue homogenization, an additional

centrifugation step was performed at 12,000 × g for 10 min to remove tissue debris. After the

chloroform extraction, the aqueous layer was combined with an equal volume of ethanol and the

RNeasy Mini Kit was used to perform washes following the manufacturer's protocol. Eggs were

crushed using pipette tips and directly stored in a mixture of cell lysis buffer and murine RNase

Inhibitor (New England Biolab).

We used the NEBNext® Single Cell/Low Input RNA Library Prep Kit for Illumina®

together with NEBNext® Multiplex Oligos for Illumina® (New England Biolab) for library

preparation, including reverse transcription of poly(A) RNA, amplification full-length cDNA,

fragmentation, ligation and final library amplification according to the manufacturer's protocol.

We performed cDNA amplification for 16 (18 for egg samples) PCR cycles and final library

amplification 8 PCR cycles. In total, we constructed 18 libraries (three for each life stage). The

quality and concentration of the resulting libraries were assessed using the Qubit High Sensitivity

dsDNA kit (Thermo Scientific) and Agilent Bioanalyzer High Sensitivity DNA assay. Libraries

were sequenced on an Illumina HiSeq2500 platform (single-end with read lengths of 50 bp) with

~18 million reads per library.

Illumina sequencing reads were pseudoaligned to the bulk transcriptome and quantified

(100 bootstrap samples) with kallisto 0.46.0 (276) using default options for single-end reads.

Fragment length sizes were extracted from the Agilent Bioanalyzer runs. For life-stage specific

differential expression analysis, kallisto quantified RNAseq data was processes with sleuth 0.30.0

(277) using Likelihood Ratio tests in R 3.6.1 (278). The average transcripts per million (tpm)

values for each target transcript were extracted from the sleuth object (see R script) and used with

the Heatmapper tool (279) to produce an unclustered heatmap showing relative expression levels.

UpSetR (280) was used to compare the number of unique and shared expressed genes across life

stages.

<u>Horizontal gene transfer events identification</u>

To detect HGTs, we used the published tool "./Lateral_gene_transfer_predictor.py" (196)

to calculate the Alien Index described by (281) and (195). All predicted genes were compared to

the NCBI nr database as previously described (196). Results to Arthropoda (tax id 6656) were

ignored in the downstream calculations. The HGT candidates were filtered for contamination

identified by both Blobtools (194) and the Alien Index (AI > 30 and >70% percent identity to a

non-metazon sequence). The candidates were further filtered for > 50% overlap with predicted

repeats using the bedtools intersect tool with the RepeatMasker gff file and expression from any

developmental stage. Introns were scored manually from visualization in IGV genome browser

(282) and GC content for all predicted genes was calculated using the bedtools nuc tool.

<u>Analysis of chemosensory and photoreceptor gene families</u>

The search and analysis chemosensory genes largely followed the procedure outlined by

Dong et al. (17) with slight modifications. First, the *Archegozetes* official gene set (OGS) was

searched using BLASTP (E-value, $<1 \times 10^{-3}$) against the following queries for the different

chemosensory gene families. The OGS was queried against i) *D. melanogaster*, *D. mojavensis*,

Anopheles gambia, *Bombyx mori*, *T. castaneum*, *Apis mellifera*, *Pediculus humanus humanus*,

and *Acyrthosiphon pisum* odorant binding proteins (OBPs) (174); ii) *D. melanogaster*, *D.*

mojavensis, *A. gambiae*, *B. mori*, *T. castaneum*, *A. mellifera*, *P. humanus humanus*, *A. pisum, I.*

scapularis, and *Daphnia pulex* small chemosensory proteins (CSP) (174, 283, 284); iii) *D.*

melanogaster and *A. mellifera* odorant receptors (283, 284); iv) *D. melanogaster*, *A. mellifera*, *I.*

scapularis, *T. urticae*, *T. mercedesae*, and *M. occidentalis* gustatory receptors (GRs) (12, 13, 15,

177, 284, 285); v) a comprehensive list of iGluRs and IRs across vertebrates and invertebrates

(178), as well as those identified in the *T. mercedesae, D. tinctorium* and *L. deliense* genome

projects (15, 17). Second, all candidate *Archegozetes* sequences were reciprocally blasted

(BLASTP, E-value <1 × 10^{-3}) against the NCBI database (248) and all sequences that did not hit

one of the respective receptors or transmembrane proteins were removed from the list. Third, for

phylogenetic analysis of IRs and GRs from *Archegozetes* were aligned with IRs from *D.*

melanogaster, *T. urticae, D. tinctorium* and *L. deliense* and GRs from iv) *D. melanogaster, T.*

mercedesae, *I. scapularis*, and *M. occidentalis*, respectively, using MAFFT (v 7.012b) with default

settings (286). Poorly aligned and variable terminal regions, as well as several internal regions of

highly variable sequences were excluded from the phylogenetic analysis. Fourth, maximum

likelihood trees were constructed with the IQ-TREE pipeline (v 1.6.12) with automated model

selection using 1,000 ultrafast bootstrap runs (270).

Reference opsin genes and opsin-like sequences were obtained from Dong et al. (17) and

used to query the *Archegozetes* OGS using BLASTP (E-value, <1 × 10^{-5}). Subsequently,

candidates sequenced were reciprocally blasted against NCBI using the same settings and only

retained if they hit an opsin or opsin-like gene. The *Archegozetes* candidates were aligned with the

query sequence list using MAFFT (v 7.012b) with default settings (286). This opsin gene

alignment phylogenetically analyzed using the IQ-TREE pipeline (v 1.6.12) with automated model

selection and 1,000 ultrafast bootstrap runs (270).

Gene family phylogenies

We used the following workflow to analyses genes related to Figure 5 (hox and

developmental genes), Figure 7 (cell wall-degrading enzyme encoding genes) and Figure 8

(alcohol and geraniol dehydrogenases genes). Generally, protein orthologs were retrieved from

NCBI (248), and aligned using MUSCLE (287) or MAFFT (v 7.012b) (286) and ends were

manually inspected and trimmed. The resulting final protein sequence alignments used to construct

a maximum likelihood (ML) phylogenetic tree with either i) PhyML with Smart Model Selection

(288, 289) or ii) the IQ-TREE pipeline with automated model selection (270). The ML trees were

constructed using either 1,000 ultrafast bootstrap runs (IQ-TREE) or approximate-likelihood ratio

test (PhyML) was used to assess node support.

<u>Feeding experiments with labelled precursors and chemical analysis (GC/MS)</u>

Stable isotope incorporation experiments were carried out as previously described (27).

Briefly, mites were fed with wheat grass containing a 10% (w/w) mixture of three antibiotics

(amoxicillin, streptomycin and tetracycline) and additionally, we added 25% (w/w) of the stable

isotope-labelled precursors $[^{13}C_6]$ D-glucose (Cambridge Isotope Laboratories, Inc.) as well as a

control with untreated wheat grass. Cultures were maintained for one generation and glands of

adult specimens were extracted one week after eclosion by submersing groups of 15 individuals

in 50 µl hexane for 5 min, which is a well-established method to obtain oil gland compounds from

mites (26, 210, 225, 290).

Crude hexane extracts (2-5 µl) were analysed with a GCMS-QP2020 gas chromatography

– mass spectrometry (GCMS) system from Shimadzu equipped with a ZB-5MS capillary column

(0.25 mm x 30m, 0.25 µm film thickness) from Phenomenex. Helium was used a carrier gas with

a flow rate of 2.14 ml/min, with splitless injection and a temperature ramp was set to increase from

50°C (5 min) to 210°C at a rate of 6°C/min, followed by 35°C/min up to 320°C (for 5 min).

Electron ionization mass spectra were recorded at 70 eV and characteristic fragment ions were

monitored in single ion mode.

Acknowledgment

We thank Joe Parker for making his laboratory space and resources available to us. AB is Simons Fellow of the Life Sciences Research Foundation (LSRF). This project was supported by a grant from the Caltech Center for Environmental Microbial Interactions (CEMI) to AB. Michael Heethoff, Sebastian Schemlzle, Benjamin Weiss and Martin Kaltenpoth graciously allowed us to use some of their unpublished images. Roy A. Norton provided invaluable comments to the manuscript and collected the first specimens of *Archegozetes longisetosus* giving rise to the current laboratory strain.

Ethics statement

There are no legal restrictions on working with mites.

Authors contributions

All authors gave final approval for publication.

Authors contributions

AB had the initial idea for the study; AB, AAB and SAK design research; IAA performed long-read sequencing and assembled the genome; AB performed all other experimental work; AAB analyzed hox and life-stage specific expression data; AB analyzed chemical data; SAK and AB performed bioinformatic analyses; AB wrote the first draft of the manuscript with input from AAB and SAK; SAK revised the manuscript. All authors gave final approval for publication.

Data availability

Genomic and transcriptomic data generated for his project can be found on NCBI under the accession numbers PRJNA683935 and PRJNA683999.

Figure 1 The mite, *Archegozetes longisetosus*, in its phylogenetic and natural environment. **a**: Species tree of selected oribatid mites of the family Trhypochthoniidae based on phylogenetic analyses and divergence time estimates by (291). **b**: Two adults and one tritonymph of *Archegozetes* on a piece of leaf litter. The algae growing on the leaf serves as a food source for the mites. **c**: Habitus of an adult mite based on a surface rendering of a μCT-scan reconstruction. Image courtesy of Sebastian Schmelzle.

Figure 2 Comparisons and annotations of the official gene set (OGS) of *Archegozetes longisetosus.* **a**: Number of gene models of the mites compared to other mites, chelicerates and the fruit fly (11, 12, 96, 97, 292). **b**: Non-linear multidimensional scaling plot (NMDS) of clustered orthogroups based on the OGS or predicted proteins of several mite species. *Archegozetes longisetosus* is marked a square, nested within Oribatida. Prostigmata are depicted in blue, Astigmata in yellow and Oribatida in green. **c**: Pie chart showing the percentage composition of genes of the *Archegozetes* annotated to different broad biological categories by GhostKOALA. **d**: Pie chart describing the overall annotation of the OGS of the mite.

Figure 3 Orthology comparison and phylogenetic placement of *Archegozetes longisetosus* among other chelicerates. **a**: Maximum

likelihood phylogeny based on concatenation of 1,121 orthologs showing the mites phylogenetic environment within the Oribatida (all

nodes have 100% support; branch length unit is substitutions per site). For the fully expanded tree see supplementary Figure S2. **b**:

BUSCO-assessment of the *Archegozetes* genome assembly and protein set for both ortholog presence and copy number compared to

957   other oribatid mites and selected model species, respectively. **c**: Comparisons of protein-coding genes in seven oribatid mite species (for

958   a full comparison of all species see supplementary Figure S3) with *Archegozetes* highlighted in red. The bar charts show the proportion

959   of orthrogroup conservation with each species (see insert legend) based on OrthoFinder clustering. **d**: Detailed pie chart depicting the

960   conservation levels of orthogroup in *Archegozetes*.

Figure 4 Comparison of repeat content estimations and transposable element (TE) landscape of *Archegozetes longisetosus*. **a:** Repetitive element categories of *Archegozetes* based on the results from RepeatModeler and MITE Tracker. LINE= long interspersed nuclear element, LTR= long terminal repeat. **b:** Comparison of total repetitive content among *Archegozetes*, other model chelicerates and the fly. All values are from the respective genome paper of the species, except for the fly. **c:** Repeat divergence plot showing TE activity through time for the major TE superfamilies of *Archegozetes*. Transposable elements with a low divergence from the consensus were recently active, while TEs diverging from the consensus depicted older activities (x-axis).

Figure 5 The genomic organization of the Hox genes and life-stage specific expression patters of developmental genes in *Archegozetes longisetosus*. **a:** Schematic of the genomic region enclosing the *Archegozetes* Hox cluster. The genomic organization of the Hox cluster is collinear, as it is in many arthropod taxa, however an abdominal-A ortholog is absent. Arrowed boxes denote the direction of transcription. The scale bar represents 50,000 base pairs. **b:** A comparison of the Hox cluster organization of reported members of Acari with the fruit fly Drosophila melanogaster as

the outgroup. The last common ancestor of the parasitiform mites *M. occidentalis* and *I. scapularis*

likely had an intact Hox cluster (green branches and labels), whereas abdominal-A was likely lost

in the last common ancestor of acariform mites, as represented by *Archegozetes* and *T. urticae* (red

branches and labels). Boxes with white borders represent duplicated Hox genes. Lines through the

boxes indicate an intact Hox cluster. See text for further details. **c:** Number of transcripts shared

across the different life stages of *Archegozetes*. The barplot panel on the left shows the numbers

of transcripts in each stage. Exemplars of (**d**) early and (**e**) mid- germ-band embryos. Ch=

chelicera; L1-3= walking legs 1-3; Pp= pedipalp. Embryos are stained with the nuclear dye DAPI

and oriented with the anterior to the left of the page. **f:** Non-clustered heatmap showing the relative

expression (row z-score based on tpm) patterns of putative limb, eye, and body segmentation genes

throughout the embryonic, larval instars, and adult stages of *Archegozetes*. See **supplementary**

Table S3 for average tpm values. Life stages (for **c** and **f**): EGG= egg; LAR= larva; PRO=

protonymph; DEU= deutonymph; TRI= tritonymph; ADU= adult.

Figure 6 The sensory systems of *Archegozetes longisetosus* and phylogenetic analysis of selected photoreceptor and chemosensory genes. **a**: Scanning electron micrograph (SEM) showing the end of tarsus on *Archegozetes'* first leg. Images shows normal setae, but also modified chemosensory setae, namely eupathidia, both paired (p) and single (s), as well as an omega-3 solenidium. SEM picture

courtesy of Michael Heethoff. **b**: Phylogeny and classification of opsin genes across the Metazoa, including those of several Chelicerata.

The tree was constructed using a maximum likelihood approach and rooted with a jelly fish opsin. *Archegozetes* sequences are depicted

in red, Drosophila in turquoise; branch length unit is substitutions per site. **c**: Maximum likelihood phylogeny of ionotropic receptors

and ionotropic glutamate receptors of *Archegozetes* (Along), *Dinothrombium* (Dt), *Leptothrombidium* (Ld), *Tetranychus* (Tu) and

Drosophila (Dmel). IR25a/IR8a and antenna/1st leg IRs contain genes with known chemosensory function in *Drosophila*. The tree was

rooted to the middle point; *Archegozetes* sequences are depicted in red, *Drosophila* in turquoise; branch length unit is substitutions per

site. Bootstrap values can be found in the supplementary Figure S13. **d**: Maximum likelihood phylogenetic tree of gustatory receptors

of *Archegozetes* (Along), *Ixodes* (Is), *Tropilaelaps* (Tm), *Metaseiulus* (Mocc) and *Drosophila* (Dmel). The tree was rooted to the middle

point; *Archegozetes* sequences are depicted in red, *Drosophila* in turquoise; branch length unit is substitutions per site. Bootstrap values

can be found in the supplementary Figure S14. **e**: Combined image of volume rendering (grey) and reconstructed nervous system of

Archegozetes in dorsal view. Color-code corresponds to different parts of the nervous system, as depicted in the legend. The blue

structure in the middle of the synganglion is the part of the esophagus which penetrates the synganglion. Scale bar: 200 µm. Image

courtesy of Sebastian Schmelzle based on data in (187).

Figure 7 Horizontal gene transfer (HGT) and implications for the feeding biology of *Archegozetes*

longisetosus. **a:** Blob-plot of the long-read genome assembly contigs plotting the read coverage

against GC proportion [%]. Contigs are colored according to the taxonomic order of their best

Megablast hit to the NCBI nucleotide database. Size of circle corresponds to the nucleotides per

contigs. **b:** Comparison of the GC content of HGT and non-HGT genes. HGT genes shifted towards the host genome GC content indicate integration within the host genome while the higher GC content HGT genes might be the product of relatively recent HGT events. **c:** Enrichment of functional categories (GO terms) describing the molecular functions and biological processes related to the HGT candidate genes. **d:** Taxonomic origin of HGT. The category "others" includes mostly protozoan donor genes among other Eukaryotes. **e:** Unrooted maximum-likelihood tree of glycoside hydrolase family 48 members (GH_48) from Streptomyces bacteria and HGT genes from other arthropods as well as *Archegozetes* (GH_48 II). Bootstrap values and the full tree can be found in the supplementary Figure S15. The scale bar denotes substitutions per site. **f:** Fluorescence *in situ* hybridization (FISH) micrograph of a food bolus in the mites' alimentary tract. The food material (wheat grass power) is enclosed in a peritrophic membrane and there is a high bacterial prevalence in the food bolus. Image courtesy of Benjamin Weiss and Martin Kaltenpoth. **g:** RNAseq support of HGT candidates related to cell wall degrading enzymes. The first block (single column) shows the overall RNA expression (tpm) of the HGT in all life stages; red denotes high total expression, while blue depicts low total expression. The second block (six columns) shows the expression (row z-score based on tpm) of the same HGT candidates across the different life stages of *Archegozetes*. Abbreviations: GH_48= glycoside hydrolase family 48, GH_44= glycoside hydrolase family 44, cbd= cellulose-binding domain, cbp= cellulose-binding protein, chiA/celA = chitinase/cellulase, GH_75= glycoside hydrolase family 75.

Figure 8 Reconstruction of the biosynthetic pathway leading to monoterpenes in *Archegozetes longisetosus*. **a**: Representative gas chromatogram of the mite' gland content; in order of retention time: 2-hydroxy-6-methyl-benzaldehyde (2,6-HMBD), neral ((*Z*)-3,7-dimethylocta-2,6-dienal) neryl formate ((*Z*)-3,7-dimethyl-2,6-octadienyl formate), tridecane, 3-hydroxybenzene-1,2-

dicarbaldehyde (γ-acaridial). Further alkanes/alkenes (pentadec-7-ene, pentadecane, heptadeca-

6,9-diene, heptadec-8-ene, heptadecane) are not shown. Monoterpenes are marked in red. **b and**

c: Representative mass spectra of neral (**b**) and neryl formate (**c**) extracted from defensive glands

of mites fed with unlabeled wheatgrass powder (control), or wheatgrass infused with $^{13}C_6$-labelled

glucose recorded in single-ion mode. The mass spectra for neral (**b**) shows the M+-ion series, while

the spectra for neryl formate (**c**) show the diagnostic ion series at $[M-46]^+$. Mites fed with the $^{13}C_6$

glucose infused wheatgrass showed enriched ions. **d:** KEGG reference pathway map for terpenoid

backbone biosynthesis. Mapping genes from the *Archegozetes* genome encoding for pathway

enzymes (labeled in red) revealed that the mite can produce geranyl pyrophosphate (GPP) *via* the

mevalonate pathway from precursors provided by glycolysis. Enzymes names correspond to EC

numbers: 2.3.1.9= acetyl-CoA C-acetyltransferase; 2.3.3.10= hydroxymethylglutaryl-CoA

synthase; 1.1.1.34= hydroxymethylglutaryl-CoA reductase; 2.7.1.36= mevalonate kinase; 2.7.4.2=

phosphomevalonate kinase; 4.1.1.33= diphosphomevalonate decarboxylase; 5.3.3.2= isopentenyl-

diphosphate delta-isomerase; 2.5.1.1= farnesyl diphosphate synthase. **e:** Maximum-likelihood tree

based on an alignment of plant, fungal and bacterial geraniol dehydrogenases, animal alcohol

dehydrogenase and two mite (*Carpoglyphus lactis* and *Archegozetes*) geraniol dehydrogenases

(GeDH). Bootstrap values (based on 1000 replicates) are indicated along branches and the scale

bar denotes substitutions per site. The tree was rooted by the outgroup cinnamyl dehydrogenase

from sweet basil. **f:** Ultrastructure of the gland-tissue of *Archegozetes*, as observed by transmission

electron microscopy (TEM). Red error shows the border between the gland cell and the glandular

lumen. TEM picture courtesy of Michael Heethoff. **g**: Proposed biochemical pathway scenario

leading to neral and neryl formate in *Archegozetes* starting with GGP from the terpenoid backbone

biosynthesis.

Table 1 *Archegozetes longisetosus* genome metrics

1055

Feature	Value
Estimated genome size	135-180 Mb
Assembly size	190 Mb
Coverage based on assembly (short/long)	200x (short), ~60x (long)
# contigs	1182
N50 (contigs)	994.5 kb
Median contig length	50.3 kb
GC content	30.9%
# gene models	23,825

1056 **Table 2** Comparison of chemosensory receptor repertoires between *Archegozetes longisetosus* and

1057 other arthropods. GR= gustatory receptor, OR= odorant receptor, IR= ionotropic receptor, OBP=

1058 odorant binding protein, CSP= chemosensory protein.

	Chemosensory receptors				
	GR	OR	IR	OBP	CSP
A. longisetosus	68	0	3	0	1
spider mite	689	0	4	0	0
deer tick	60	0	22	0	1
house spider	634	0	108	4	0
fruit fly	73	62	66	51	4

1059

1060

 Supplementary Material

 Supplementary Figures

Figure S1 Results of *in silico* genome size estimations based on jellyfish *k-mer* counting using **a:**

GenomeScope v1.0 and **b** and **c**: the findGSE v0.1.0 R package (231).

Figure S2 Phylogenetic placement of *Archegozetes longisetosus* among other chelicerates. **a**:

Maximum likelihood phylogeny based on concatenation of 1,121 orthologs Branch lengths unit is

substitutions per site and the node values reflect bootstrap supports. **b**: Coalescence species tree

reconstruction of the 1,121 filtered orthogroups. Branch lengths are presented in coalescent units

(differences in the 1,121 gene trees) and the node values reflect the local posterior probabilities.

Figure S3 Comparisons of protein-coding genes of 23 arthropod species, including *Archegozetes*.

The bar charts show the proportion of orthrogroup conservation with each species (see insert

legend) based on OrthoFinder clustering.

Figure S4 Maximum likelihood phylogenetic analyses of the *A. longisetosus* Paired protein

orthologs. **(a)** Maximum likelihood tree showing the relationship of the Eyegone, Pax-3/7, and

Pax-6 clades as collapsed subtrees. **(b)** The un-collapsed clade in **A** showing the phylogenetic

relationships of selected Eyegone proteins and the putative *A. longisetosus* Eyegone ortholog. **(c)**

The un-collapsed clade in **A** showing the phylogenetic relationships of selected Pax-3/7 proteins

and the putative *A. longisetosus* Pax-3/7 orthologs. **(d)** The un-collapsed clade in **A** showing the

phylogenetic relationships of selected Pax-6 proteins and the putative *A. longisetosus* Pax-6

ortholog. All *A. longisetosus* orthologs are in red, and the *D. melanogaster* orthologs are in blue.

Node support was calculated using the approximate likelihood ratio (aLRT) method and is

represented by the color of each node. All taxa are represented by their species names, gene names

if given, and their NCBI accession numbers.

Figure S5 Maximum likelihood phylogenetic analyses of the *A. longisetosus* Eyes absent (Eya)

protein ortholog and selected metazoan Eya proteins. The *A. longisetosus* ortholog is in red, and

the *D. melanogaster* orthologs are in blue. Node support was calculated using the approximate

likelihood ratio (aLRT) method and is represented by the color of each node. All taxa are

represented by their species names, gene names if given, and their NCBI accession numbers.

Figure S6 Maximum likelihood phylogenetic analyses of the *A. longisetosus* Hairy protein

ortholog and selected metazoan Hairy proteins. Hairy/E(spl) proteins were used as an outgroup.

The *A. longisetosus* ortholog is in red, and the *D. melanogaster* orthologs are in blue. Node support

was calculated using the approximate likelihood ratio (aLRT) method and is represented by the

color of each node. All taxa are represented by their species names, gene names if given, and their

NCBI accession numbers.

Figure S7 Maximum likelihood phylogenetic analyses of the *A. longisetosus* Omb, T-box H15,

and TBX1 protein orthologs and selected metazoan T-box proteins. All *A. longisetosus* orthologs

are in red, and the *D. melanogaster* orthologs are in blue. Node support was calculated using the

approximate likelihood ratio (aLRT) method and is represented by the color of each node. All taxa

are represented by their species names, gene names if given, and their NCBI accession numbers.

Figure S8 Maximum likelihood phylogenetic analyses of the *A. longisetosus* Runt protein ortholog

and selected arthropod Runt proteins. The *A. longisetosus* ortholog is in red, and the *D.*

melanogaster orthologs are in blue. Node support was calculated using the approximate likelihood

ratio (aLRT) method and is represented by the color of each node. All taxa are represented by their

species names, gene names if given, and their NCBI accession numbers. See

Figure S9 Maximum likelihood phylogenetic analyses of the *A. longisetosus* Six family protein

orthologs and selected metazoan Six family proteins. All *A. longisetosus* orthologs are in red, and

the *D. melanogaster* orthologs are in blue. Node support was calculated using the approximate

likelihood ratio (aLRT) method and is represented by the color of each node. All taxa are

represented by their species names, gene names if given, and their NCBI accession numbers.

Figure S10 Maximum likelihood phylogenetic analyses of the *A. longisetosus* Sp-family protein

orthologs and selected metazoan Sp-family proteins. All *A. longisetosus* orthologs are in red, and

the *D. melanogaster* orthologs are in blue. Node support was calculated using the approximate

likelihood ratio (aLRT) method and is represented by the color of each node. All taxa are

represented by their species names, gene names if given, and their NCBI accession numbers.

Figure S11 Maximum likelihood phylogenetic analyses of the *A. longisetosus* Wnt-family protein

orthologs and selected metazoan Wnt proteins. The tree is organized as a cladogram for easier

viewing. All *A. longisetosus* orthologs are in red, and the *D. melanogaster* orthologs are in blue.

Node support was calculated using the approximate likelihood ratio (aLRT) method and is

represented by the color of each node. All taxa are represented by their species names, gene names

if given, and their NCBI accession numbers.

Figure S12 Unrooted maximum-likelihood phylogenetic trees of cell-wall degrading enzymes

based on the alignment of amino acid sequences. Branch lengths unit is substitutions per site and

the node values reflect bootstrap supports. *Archegozetes* sequences are highlighted in red.

Figure S13 Maximum likelihood phylogeny of ionotropic receptors and ionotropic glutamate

receptors of *Archegozetes* (Along), *Dinothrombium* (Dt), *Leptothrombidium* (Ld), *Tetranychus*

(Tu) and *Drosophila* (Dmel). The tree was rooted to the middle point. Branch lengths unit is

substitutions per site and the node values reflect bootstrap supports.

Figure S14 Maximum likelihood phylogenetic tree of gustatory receptors of *Archegozetes*

(Along), *Ixodes* (Is), *Tropilaelaps* (Tm), *Metaseiulus* (Mocc) and *Drosophila* (Dmel). The tree was

rooted to the middle point. Branch lengths unit is substitutions per site and the node values reflect

bootstrap supports.

Figure S15 Unrooted maximum-likelihood tree of glycoside hydrolase family 48 members

(GH_48) from Streptomyces bacteria and HGT genes from other arthropods as well as

Archegozetes (GH_48 II). Branch lengths unit is substitutions per site and the node values reflect

bootstrap supports.

1136

Supplementary Table

Table S1 Contamination contigs identified from Blobtools.

Table S2 Phylogenetic statistics of the PhyML constructed trees as well as the matrices selected

by the SMS model selection tool.

Table S3 The average transcript per million (tpm) values for the transcripts highlighted in the

heatmap (**Figure 5f**) for each instar stage.

Table S4 Candidate HGTs identified from the *Archegozetes* genome. The genes were filtered first

if they were predicted to be contamination from Blobtools and the Alien Index report, second if

they overlapped predicted repeats by $\geq$ 50%, and third if they were not expressed in any

developmental stage. Annotation is provided from similarity searches against the NCBI nr

database, other oribatid mite and eggNOG database. The taxonomy of the sequences upstream and

downstream of each candidate HGT was determine using the eggNOG predicted taxonomic group.

1149

References

1. Childers A (2020) Sequenced Arthropod Genomes. in http://i5k.github.io/arthropod_genomes_at_ncbi (i5k initiative).

2. Thomas GW, *et al.* (2020) Gene content evolution in the arthropods. *Genome biology* 21(1):1-14.

3. Giribet G & Edgecombe GD (2019) The phylogeny and evolutionary history of arthropods. *Current Biology* 29(12):R592-R602.

4. Regier JC*, et al.* (2010) Arthropod relationships revealed by phylogenomic analysis of nuclear protein-coding sequences. *Nature* 463(7284):1079-1083.

5. Dunlop J & Selden P (1998) The early history and phylogeny of the chelicerates. *Arthropod relationships*, (Springer), pp 221-235.

6. Ballesteros JA & Sharma PP (2019) A critical appraisal of the placement of Xiphosura (Chelicerata) with account of known sources of phylogenetic error. *Systematic Biology* 68(6):896-917.

7. Dunlop JA (2010) Geological history and phylogeny of Chelicerata. *Arthropod structure & development* 39(2-3):124-142.

8. Schaefer I, Norton RA, Scheu S, & Maraun M (2010) Arthropod colonization of land - Linking molecules and fossils in oribatid mites (Acari, Oribatida). *Molecular Phylogenetics and Evolution* 57:113-121.

9. Dunlop J & Alberti G (2008) The affinities of mites and ticks: a review. *Journal of Zoological Systematics and Evolutionary Research* 46(1):1-18.

10. Walter DE & Proctor HC (1999) Mites: ecology, evolution, and behaviour.

11.   Grbić M, *et al.* (2011) The genome of Tetranychus urticae reveals herbivorous pest

adaptations. *Nature* 479(7374):487-492.

12.   Gulia-Nuss M, *et al.* (2016) Genomic insights into the Ixodes scapularis tick vector of

Lyme disease. *Nature communications* 7(1):1-13.

13.   Hoy MA, *et al.* (2016) Genome sequencing of the phytoseiid predatory mite Metaseiulus

occidentalis reveals completely atomized Hox genes and superdynamic intron evolution.

Genome biology and evolution 8(6):1762-1775.

14.   Rider SD, Morgan MS, & Arlian LG (2015) Draft genome of the scabies mite. *Parasites*

& vectors 8(1):1-14.

15.   Dong X, *et al.* (2017) Draft genome of the honey bee ectoparasitic mite, Tropilaelaps

mercedesae, is shaped by the parasitic life history. *Gigascience* 6(3):gix008.

16.   Cornman RS, *et al.* (2010) Genomic survey of the ectoparasitic mite Varroa destructor, a

major pest of the honey bee Apis mellifera. *BMC genomics* 11(1):602.

17.   Dong X, *et al.* (2018) Genomes of trombidid mites reveal novel predicted allergens and

laterally transferred genes associated with secondary metabolism. *GigaScience*

7(12):giy127.

18.   Bast J, *et al.* (2016) No accumulation of transposable elements in asexual arthropods.

Molecular biology and evolution 33(3):697-706.

19.   Klimov PB & OConnor B (2013) Is permanent parasitism reversible?—Critical evidence

from early evolution of house dust mites. *Systematic biology* 62(3):411-423.

20.   Weinstein SB & Kuris AM (2016) Independent origins of parasitism in Animalia.

Biology Letters 12(7):20160324.

21. Shingate P*, et al.* (2020) Chromosome-level assembly of the horseshoe crab genome

provides insights into its genome evolution. *Nature communications* 11(1):1-13.

22. Lozano-Fernandez J*, et al.* (2019) Increasing species sampling in chelicerate genomic-

scale datasets provides support for monophyly of Acari and Arachnida. *Nature*

communications 10(1):1-8.

23. Aoki J (1965) Oribatiden (Acarina) Thailands. I. *Nature and Life in Southeast Asia*

4:129-193.

24. Schmelzle S & Blüthgen N (2019) Under pressure: force resistance measurements in box

mites (Actinotrichida, Oribatida). *Frontiers in zoology* 16(1):24.

25. Heethoff M & Koerner L (2007) Small but powerful: the oribatid mite Archegozetes

longisetosus Aoki (Acari, Oribatida) produces disproportionately high forces. *Journal of*

Experimental Biology 210(17):3036-3042.

26. Brückner A*, et al.* (2017) Storage and release of hydrogen cyanide in a chelicerate

(Oribatula tibialis). *Proc Natl Acad Sci U S A* 114(13):3469-3472.

27. Brückner A, Kaltenpoth M, & Heethoff M (2020) De novo biosynthesis of simple

aromatic compounds by an arthropod (Archegozetes longisetosus). *Proceedings of the*

Royal Society B 287(1934):20201429.

28. Norton RA & Palmer SC (1991) The distribution, mechanisms and evolutionary

significance of parthenogenesis in oribatid mites. *The Acari - Reproduction, Development*

and Life-History Strategies, eds Schuster R & Murphy PW (Chapman & Hall, London),

pp 107-136.

29. Heethoff M, Norton RA, Scheu S, & Maraun M (2009) Parthenogenesis in Oribatid Mites

(Acari, Oribatida): Evolution Without Sex. *Lost Sex: The Evolutionary Biology of*

Parthenogenesis, eds Schön I, Martens K, & van Dijk P (Springer, Dordrecht), pp 241-

257.

30.   Raspotnig G (2009) Oil gland secretions in Oribatida (Acari). *Trends in Acarology*, eds

Sabelis MW & Bruin J (Springer), pp 235-239.

31.   Maraun M, Schatz H, & Scheu S (2007) Awesome or ordinary? Global diversity patterns

of oribatid mites. *Ecography* 30(2):209-216.

32.   Maraun M & Scheu S (2000) The structure of oribatid mite communities (Acari,

Oribatida): Patterns, mechanisms and implications for future research. *Ecography*

23(3):374-383.

33.   Heethoff M, Bergmann P, Laumann M, & Norton RA (2013) The 20th anniversary of a

model mite: A review of current knowledge about Archegozetes longisetosus (Acari,

Oribatida). *Acarologia* 53(4):353-368.

34.   Palmer SC & Norton RA (1992) Genetic Diversity in Thelytokous Oribatid Mites (Acari;

Acariformes: Desmonomata). *Biochemical Systematics and Ecology* 20(3):219-231.

35.   Wrensch DL, Kethley JB, & Norton RA (1994) Cytogenetics of holokinetic

chromosomes and inverted meiosis: keys to the evolutionary success of mites, with

generalizations on eukaryotes. *Mites: Ecological and Evolutionary Analyses of Life-*

History Patterns, ed Houck MA (Chapman & Hall, New York), pp 282-343.

36.   Heethoff M, Bergmann P, & Norton RA (2006) Karyology and sex determination of

oribatid mites. *Acarologia* 46(1-2):127-131.

37.   Bergmann P, Laumann M, Norton RA, & Heethoff M (2018) Cytological evidence for

automictic thelytoky in parthenogenetic oribatid mites (Acari, Oribatida): Synaptonemal

complexes confirm meiosis in Archegozetes longisetosus. *Acarologia* in press.

38. Oxley PR*, et al.* (2014) The genome of the clonal raider ant Cerapachys biroi. *Current*

Biology 24(4):451-458.

39. Norton RA (1994) Evolutionary aspects of oribatid mite life histories and consequences

for the origin of the Astigmata. *Mites: Ecological and Evolutionary Analyses of Life-*

History Patterns, ed Houck MA (Chapman & Hall, New York), Chapman & Hall Ed, pp

99-135.

40. Norton RA (2007) Holistic acarology and ultimate causes: examples from the oribatid

mites. *Acarology XI: Proceedings of the International Congress.*, eds Morales-Malacara

JB, Behan-Pelletier V, Ueckermann E, Perez TM, Estrada-Venegas EG, & Badii M

(Sociedad Latinoamericana de Acarologia), pp 3-20.

41. Sakata T & Norton RA (2001) Opisthonotal gland chemistry of early-derivative oribatid

mites (Acari) and its relevance to systematic relationships of Astigmata. *International*

Journal of Acarology 27(4):281-292.

42. Heethoff M (2012) Regeneration of complex oil-gland secretions and its importance for

chemical defense in an oribatid mite. *J Chem Ecol* 38(9):1116-1123.

43. Brückner A & Parker J (2020) Molecular evolution of gland cell types and chemical

interactions in animals. *Journal of Experimental Biology* 223(Suppl 1).

44. Riha G (1951) Zur Ökologie der Oribatiden in Kalksteinböden. *Zoologische Jahrbücher*

80:407-450.

45. Smrž J (1992) Some adaptive features in the microanatomy of moss-dwelling oribatid

mites (Acari: Oribatida) with respect to their ontogenetical development. *Pedobiologia*

36(5):306-320.

46. Zachvatkin AA (1941) Fauna of the U.S.S.R., Vol. VI, No.1 Tyroglyphoidae (Acari).

Zoological Institute of the Acaemy of Science of the U.S.S.R., Moscow.

47. Shimano S, Sakata T, Mizutani Y, Kuwahara Y, & Aoki J-i (2002) Geranial: the alarm

pheromone in the nymphal stage of the oribatid mite, Nothrus palustris. *Journal of*

Chemical Ecology 28(9):1831-1837.

48. Heethoff M, Koerner L, Norton RA, & Raspotnig G (2011) Tasty but protected-first

evidence of chemical defense in oribatid mites. *Journal of Chemical Ecology* 37:1037-

1043.

49. Raspotnig G (2006) Chemical alarm and defence in the oribatid mite Collohmannia

gigantea (Acari: Oribatida). *Experimental and Applied Acarology* 39(3):177-194.

50. Heethoff M*, et al.* (2018) Life as a fortress–structure, function, and adaptive values of

morphological and chemical defense in the oribatid mite Euphthiracarus reticulatus

(Actinotrichida). *BMC Zoology* 3(1):7.

51. Saporito RA*, et al.* (2007) Oribatid mites as a major dietary source for alkaloids in poison

frogs. *Proceedings of the National Academy of Sciences of the USA* 104(21):8885-8890.

52. Brückner A, Stabentheiner E, Leis HJ, & Raspotnig G (2015) Chemical basis of

unwettability in Liacaridae (Acari, Oribatida): specific variations of a cuticular acid/ester-

based system. *Exp Appl Acarol* 66(3):313-335.

53. Saporito RA, Spande TF, Garraffo HM, & Donnelly MA (2009) Arthropod alkaloids in

poison frogs: A review of the dietary hypothesis. *Heterocycles* 79:277-297.

54. Bensoussan N*, et al.* (2016) Plant-herbivore interaction: dissection of the cellular pattern

of Tetranychus urticae feeding on the host plant. *Front Plant Sci* 7:1105.

55.   Cohen AC (1995) Extra-oral digestion in predaceous terrestrial Arthropoda. *Annual*

review of entomology 40(1):85-103.

56.   Shultz JW (2007) A phylogenetic analysis of the arachnid orders based on morphological

characters. *Zoological Journal of the Linnean Society* 150(2):221-265.

57.   Heethoff M & Norton RA (2009) A new use for synchrotron X-ray microtomography:

three-dimensional biomechanical modeling of chelicerate mouthparts and calculation of

theoretical bite forces. *Invertebrate Biology* 128(4):332-339.

58.   Brückner A*, et al.* (2017) Track the snack – Olfactory cues shape foraging behaviour of

decomposing soil mites (Oribatida). *Pedobiologia* in press - online early.

59.   Schneider K*, et al.* (2004) Trophic niche differentiation in soil microarthropods

(Oribatida, Acari): evidence from stable isotope ratios (N-15/N-14). *Soil Biology and*

Biochemistry 36(11):1769-1774.

60.   Schneider K, Renker C, Scheu S, & Maraun M (2004) Feeding biology of oribatid mites:

a minireview. *Phytophaga* 14:247-256.

61.   Schneider K & Maraun M (2005) Feeding preferences among dark pigmented fungal taxa

("Dematiacea") indicate limited trophic niche differentiation of oribatid mites (Oribatida,

Acari). *Pedobiologia* 49(1):61-67.

62.   Heidemann K, Scheu S, Ruess L, & Maraun M (2011) Molecular detection of nematode

predation and scavenging in oribatid mites: Laboratory and field experiments. *Soil*

Biology and Biochemistry 43:229-236.

63.   Maraun M*, et al.* (2011) Stable isotopes revisited: Their use and limits for oribatid mite

trophic ecology. *Soil Biology and Biochemistry* 43(5):877-882.

64.   Brückner A, Schuster R, Smit T, & Heethoff M (2018) Imprinted or innated food

preferences in the model mite Archegozetes longisetosus (Actinotrichida, Oribatida,

Trhypochthoniidae). *Soil Organisms* submitted.

65.   Smrž J & Čatská V (2010) Mycophagous mites and their internal associated bacteria

cooperate to digest chitin in soil. *Symbiosis* 52(1):33-40.

66.   Smrz J (2000) A modified test for chitinase and cellulase activity in soil mites.

Pedobiologia 44(2):186-189.

67.   Siepel H & de Ruiter-Dijkman EM (1993) Feeding guilds of oribatid mites based on their

carbohydrase activities. *Soil Biology and Biochemistry* 25(11):1491-1497.

68.   Luxton M (1979) Food and energy processing by oribatid mites. *Rev. Écol. Biol. Sol*

16:103-111.

69.   Haq MA (1993) Symbiotic association of mites and microbes in cellulose degradation.

Soil Organisms and Sustainability 4:81-85.

70.   Zinkler D (1971) Vergleichende Untersuchungen zum Wirkungsspektrum der

Carbohydrasen laubstreubewohnender Oribatiden. *Zool Ges Verh*.

71.   Smrž J & Norton RA (2004) Food selection and internal processing in Archegozetes

longisetosus (Acari : Oribatida). *Pedobiologia* 48(2):111-120.

72.   McKenna DD*, et al.* (2019) The evolution and genomic basis of beetle diversity.

Proceedings of the National Academy of Sciences 116(49):24729-24737.

73.   Wybouw N, Pauchet Y, Heckel DG, & Van Leeuwen T (2016) Horizontal gene transfer

contributes to the evolution of arthropod herbivory. *Genome biology and evolution*

8(6):1785-1801.

74. Mayer WE, Schuster LN, Bartelmes G, Dieterich C, & Sommer RJ (2011) Horizontal

gene transfer of microbial cellulases into nematode genomes is associated with functional

assimilation and gene turnover. *BMC evolutionary biology* 11(1):1-10.

75. Wybouw N, Van Leeuwen T, & Dermauw W (2018) A massive incorporation of

microbial genes into the genome of Tetranychus urticae, a polyphagous arthropod

herbivore. *Insect molecular biology* 27(3):333-351.

76. Wu C*, et al.* (2017) Analysis of the genome of the New Zealand giant collembolan

(Holacanthella duospinosa) sheds light on hexapod evolution. *BMC genomics* 18(1):795.

77. Faddeeva-Vakhrusheva A*, et al.* (2016) Gene family evolution reflects adaptation to soil

environmental stressors in the genome of the collembolan Orchesella cincta. *Genome*

biology and evolution 8(7):2106-2117.

78. Hoffmann A*, et al.* (1998) Intergeneric transfer of conjugative and mobilizable plasmids

harbored by Escherichia coli in the gut of the soil microarthropod Folsomia candida

(Collembola). *Applied and Environmental Microbiology* 64(7):2652-2659.

79. Barnett AA & Thomas RH (2013) Posterior Hox gene reduction in an arthropod:

Ultrabithorax and Abdominal-B are expressed in a single segment in the mite

Archegozetes longisetosus. *EvoDevo* 4(1):23.

80. Barnett AA & Thomas RH (2018) Early segmentation in the mite Archegozetes

longisetosus reveals conserved and derived aspects of chelicerate development.

Development genes and evolution 228(5):213-217.

81. Barnett AA & Thomas RH (2012) The delineation of the fourth walking leg segment is

temporally linked to posterior segmentation in the mite Archegozetes longisetosus (Acari:

Oribatida, Trhypochthoniidae). *Evol Dev* 14(4):383-392.

82. Barnett AA & Thomas RH (2013) The expression of limb gap genes in the mite

Archegozetes longisetosus reveals differential patterning mechanisms in chelicerates.

Evol Dev 15(4):280-292.

83. Brückner A, Hilpert A, & Heethoff M (2017) Biomarker function and nutritional

stoichiometry of neutral lipid fatty acids and amino acids in oribatid mites. *Soil Biology*

& Biochemistry 115:35-43.

84. Thomas RH (2002) Mites as models in development and genetics. *Acarid Phylogeny and*

Evolution: Adaptation in Mites and Ticks: Proceedings of the IV Symposium of the

European Association of Acarologists, eds Bernini F, Nannelli R, Nuzzaci G, & de Lillo

E (Kluwer Academic Publishers), pp 21-26.

85. Brückner A, Wehner K, Neis M, & Heethoff M (2016) Attack and defense in a gamasid-

oribatid mite predator-prey experiment - sclerotization outperforms chemical repellency.

Acarologia 56(4):451-461.

86. Heethoff M & Raspotnig G (2012) Expanding the 'enemy-free space' for oribatid mites:

evidence for chemical defense of juvenile Archegozetes longisetosus against the rove

beetle Stenus juno. *Experimental and Applied Acarology* 56:93-97.

87. Brückner A, Schuster R, Wehner K, & Heethoff M (2018) Adding to the reproductive

biology of Archegozetes longisetosus (Actinotrichida, Oribatida, Trhypochthoniidae)

again: the role of nutritional quality. *Soil Organisms* (submitted).

88. Raspotnig G, Norton RA, & Heethoff M (2011) Oribatid mites and skin alkaloids in

poison frogs. *Biology Letters* 7:555-556.

89. Heethoff M & Rall BC (2015) Reducible defence: chemical protection alters the

dynamics of predator–prey interactions. *Chemoecology* 25(2):53-61.

90.   Brückner A & Heethoff M (2018) Nutritional effects on chemical defense alter predator–

prey dynamics. *Chemoecology* 28(1):1-9.

91.   Thiel T, Brechtel A, Brückner A, Heethoff M, & Drossel B (2018) The effect of

reservoir-based chemical defense on predator-prey dynamics. *Theoretical Ecology*:1-14.

92.   Norton RA, Kethley JB, Johnston DE, & O'Connor BM (1993) Phylogenetic

perspectives on genetic systems and reproductive modes of mites. *Evolution and*

Diversity of Sex Ratio in Insects and Mites, eds Wrensch D & Ebbert M (Chapman &

Hall, London), pp 8-99.

93.   Oliver Jr JH (1983) Chromosomes, genetic variance and reproductive strategies among

mites and ticks. *Bulletin of the ESA* 29(2):8-17.

94.   Simpson JT (2014) Exploring genome characteristics and sequence quality without a

reference. *Bioinformatics* 30(9):1228-1235.

95.   Alfsnes K, Leinaas HP, & Hessen DO (2017) Genome size in arthropods; different roles

of phylogeny, habitat and life history in insects and crustaceans. *Ecology and Evolution*

7(15):5939-5947.

96.   Schwager EE*, et al.* (2017) The house spider genome reveals an ancient whole-genome

duplication during arachnid evolution. *BMC biology* 15(1):1-27.

97.   dos Santos G*, et al.* (2015) FlyBase: introduction of the Drosophila melanogaster Release

6 reference genome assembly and large-scale migration of genome annotations. *Nucleic*

acids research 43(D1):D690-D697.

98.   Consortium TGS (2008) The genome of the model beetle and pest Tribolium castaneum.

Nature 452(7190):949.

99.   Palmer M, Bantle J, Guo X, & Fargoxy1 WS (1994) Genome size and organization in the

1397   ixodid tick Amblyomma americanum (L.). *Insect molecular biology* 3(1):57-62.

1398   100.   Van Zee JP*, et al.* (2007) Tick genomics: the Ixodes genome project and beyond.

1399   *International journal for parasitology* 37(12):1297-1305.

1400   101.   Barrero RA*, et al.* (2017) Gene-enriched draft genome of the cattle tick Rhipicephalus

1401   microplus: assembly by the hybrid Pacific Biosciences/Illumina approach enabled

1402   analysis of the highly repetitive genome. *International journal for parasitology*

1403   47(9):569-583.

1404   102.   Emms DM & Kelly S (2015) OrthoFinder: solving fundamental biases in whole genome

1405   comparisons dramatically improves orthogroup inference accuracy. *Genome biology*

1406   16(1):157.

1407   103.   Kanehisa M, Sato Y, & Morishima K (2016) BlastKOALA and GhostKOALA: KEGG

1408   tools for functional characterization of genome and metagenome sequences. *Journal of*

1409   *molecular biology* 428(4):726-731.

1410   104.   Palmer J & Stajich J (2017) Funannotate: eukaryotic genome annotation pipeline.

1411   105.   Pachl P*, et al.* (2012) Convergent evolution of defense mechanisms in oribatid mites

1412   (Acari, Oribatida) shows no "ghosts of predation past". *Molecular Phylogenetics and*

1413   *Evolution* 65(2):412-420.

1414   106.   Alberti G & Michalik P (2004) *Feinstrukturelle Aspekte der Fortpflanzungssysteme von*

1415   *Spinnentieren (Arachnida)* (Denisia, Wien).

1416   107.   Koller LM, Wirth S, & Raspotnig G (2012) Geranial-rich oil gland secretions: a common

1417   phenomenon in the Histiostomatidae (Acari, Astigmata)? *International Journal of*

1418   *Acarology* 38(5):420-426.

1419   108.   Liana M & Witaliński W (2005) Sperm structure and phylogeny of Astigmata. *Journal of*

1420   *morphology* 265(3):318-324.

1421   109.   Dabert M, Witalinski W, Kazmierski A, Olszanowski Z, & Dabert J (2010) Molecular

1422   phylogeny of acariform mites (Acari, Arachnida): Strong conflict between phylogenetic

1423   signal and long-branch attraction artifacts. *Molecular Phylogenetics and Evolution*

1424   56:222-241.

1425   110.   Maraun M*, et al.* (2004) Molecular phylogeny of oribatid mites (Oribatida, Acari):

1426   evidence for multiple radiations of parthenogenetic lineages. *Experimental and Applied*

1427   *Acarology* 33(3):183-201.

1428   111.   Norton RA (1998) Morphological evidence for the evolutionary origin of Astigmata

1429   (Acari : Acariformes). *Experimental and Applied Acarology* 22(10):559-594.

1430   112.   Pepato A & Klimov P (2015) Origin and higher-level diversification of acariform mites–

1431   evidence from nuclear ribosomal genes, extensive taxon sampling, and secondary

1432   structure alignment. *BMC evolutionary biology* 15(1):178.

1433   113.   Li W-N & Xue X-F (2019) Mitochondrial genome reorganization provides insights into

1434   the relationship between oribatid mites and astigmatid mites (Acari: Sarcoptiformes:

1435   Oribatida). *Zoological Journal of the Linnean Society* 187(3):585-598.

1436   114.   Domes K, Althammer M, Norton RA, Scheu S, & Maraun M (2007) The phylogenetic

1437   relationship between Astigmata and Oribatida (Acari) as indicated by molecular markers.

1438   *Experimental and Applied Acarology* 42(3):159-171.

1439   115.   Klimov PB*, et al.* (2018) Comprehensive phylogeny of acariform mites (Acariformes)

1440   provides insights on the origin of the four-legged mites (Eriophyoidea), a long branch.

1441   *Molecular Phylogenetics and Evolution* 119:105-117.

1442   116.   Van Dam MH, Trautwein M, Spicer GS, & Esposito L (2019) Advancing mite

1443   phylogenomics: Designing ultraconserved elements for Acari phylogeny. *Molecular*

1444   *ecology resources* 19(2):465-475.

1445   117.   Jeyaprakash A & Hoy MA (2009) First divergence time estimate of spiders, scorpions,

1446   mites and ticks (subphylum: Chelicerata) inferred from mitochondrial phylogeny.

1447   *Experimental and Applied Acarology* 47(1):1-18.

1448   118.   Alberti G (1984) The contribution of comparative spermatology to problems of acarine

1449   systematics. *Acarology VI*, pp 479-490.

1450   119.   Alberti G (1991) Spermatology in the Acari: systematic and functional implications. *The*

1451   *Acari - Reproduction, Development and Life-History Strategies*, eds Schuster R &

1452   Murphy PW (Chapman & Hall, London), pp 77-105.

1453   120.   Dabert M (2006) DNA markers in the phylogenetics of the Acari. *Biological Letters*

1454   43(2):97-107.

1455   121.   Simão FA, Waterhouse RM, Ioannidis P, Kriventseva EV, & Zdobnov EM (2015)

1456   BUSCO: assessing genome assembly and annotation completeness with single-copy

1457   orthologs. *Bioinformatics* 31(19):3210-3212.

1458   122.   Siepel A*, et al.* (2005) Evolutionarily conserved elements in vertebrate, insect, worm, and

1459   yeast genomes. *Genome research* 15(8):1034-1050.

1460   123.   Nagy LG, Merényi Z, Hegedüs B, & Bálint B (2020) Novel phylogenetic methods are

1461   needed for understanding gene function in the era of mega-scale genome sequencing.

1462   *Nucleic Acids Research* 48(5):2209-2219.

1463   124.   Charlesworth B (2012) The effects of deleterious mutations on evolution at linked sites.

1464   *Genetics* 190(1):5-22.

1465 125. Barton NH (2010) Mutation and the evolution of recombination. *Philosophical*

1466   *Transactions of the Royal Society B: Biological Sciences* 365(1544):1281-1294.

1467 126. Schön I, Martens K, & van Dijk P (2009) *Lost Sex - The Evolutionary Biology of*

1468   *Parthenogenesis* (Springer, Dordrecht).

1469 127. Muller HJ (1964) The relation of recombination to mutaional advance. *Mutation*

1470   *Research* 106:2-9.

1471 128. Arkhipova I & Meselson M (2000) Transposable elements in sexual and ancient asexual

1472   taxa. *Proceedings of the National Academy of Sciences* 97(26):14473-14477.

1473 129. Nuzhdin SV & Petrov DA (2003) Transposable elements in clonal lineages: lethal

1474   hangover from sex. *Biological Journal of the Linnean Society* 79(1):33-41.

1475 130. Panfilio KA*, et al.* (2019) Molecular evolutionary trends and feeding ecology

1476   diversification in the Hemiptera, anchored by the milkweed bug genome. *Genome*

1477   *biology* 20(1):64.

1478 131. Bourque G*, et al.* (2018) Ten things you should know about transposable elements.

1479   *Genome biology* 19(1):1-12.

1480 132. Crescente JM, Zavallo D, Helguera M, & Vanzetti LS (2018) MITE Tracker: an accurate

1481   approach to identify miniature inverted-repeat transposable elements in large genomes.

1482   *BMC bioinformatics* 19(1):348.

1483 133. Finnegan DJ (1989) Eukaryotic transposable elements and genome evolution. *Trends in*

1484   *genetics* 5:103-107.

1485 134. Waterston RH*, et al.* (2002) Initial sequencing and comparative analysis of the mouse

1486   genome. *Nature* 420(6915):520-562.

1487   135.   Dej KJ, Gerasimova T, Corces VG, & Boeke JD (1998) A hotspot for the Drosophila

1488   gypsy retroelement in the ovo locus. *Nucleic acids research* 26(17):4019-4024.

1489   136.   Havecker ER, Gao X, & Voytas DF (2004) The diversity of LTR retrotransposons.

1490   *Genome biology* 5(6):1-6.

1491   137.   Hrycaj SM & Wellik DM (2016) Hox genes and evolution. *F1000Research* 5.

1492   138.   Holland P & Hogan B (1988) Expression of homeo box genes during mouse

1493   development: a review. *Genes & development* 2(7):773-782.

1494   139.   Hughes CL & Kaufman TC (2002) Hox genes and the evolution of the arthropod body

1495   plan 1. *Evolution & development* 4(6):459-499.

1496   140.   Dunlop JA & Lamsdell JC (2017) Segmentation and tagmosis in Chelicerata. *Arthropod

1497   structure & development* 46(3):395-418.

1498   141.   van der Hammen L (1970) La segmentation primitive des Acariens. *Acarologia* 12(1):3-

1499   10.

1500   142.   Schwager EE, Schönauer A, Leite DJ, Sharma PP, & McGregor AP (2015) Chelicerata.

1501   *Evolutionary Developmental Biology of Invertebrates 3*, (Springer), pp 99-139.

1502   143.   Sharma PP, Schwager EE, Extavour CG, & Giribet G (2012) Hox gene expression in the

1503   harvestman Phalangium opilio reveals divergent patterning of the chelicerate

1504   opisthosoma. *Evolution & development* 14(5):450-463.

1505   144.   Santos VT*, et al.* (2013) The embryogenesis of the tick Rhipicephalus (Boophilus)

1506   microplus: the establishment of a new chelicerate model system. *Genesis* 51(12):803-818.

1507   145.   Pace RM, Grbić M, & Nagy LM (2016) Composition and genomic organization of

1508   arthropod Hox clusters. *Evodevo* 7(1):11.

1509   146.   Cook CE, Smith ML, Telford MJ, Bastianello A, & Akam M (2001) Hox genes and the

1510   phylogeny of the arthropods. *Current Biology* 11(10):759-763.

1511   147.   Telford MJ & Thomas RH (1998) Expression of homeobox genes shows chelicerate

1512   arthropods retain their deutocerebral segment. *Proceedings of the National Academy of*

1513   *Sciences of the USA* 95(18):10671-10675.

1514   148.   Heethoff M, Laumann M, & Bergmann P (2007) Adding to the reproductive biology of

1515   the parthenogenetic oribatid mite, Archegozetes longisetosus (Acari, Oribatida,

1516   Trhypochthoniidae). *Turkish Journal of Zoology* 31:151-159.

1517   149.   Sharma PP*, et al.* (2015) A conserved genetic mechanism specifies deutocerebral

1518   appendage identity in insects and arachnids. *Proceedings of the Royal Society B:*

1519   *Biological Sciences* 282(1808):20150698.

1520   150.   Königsmann T, Turetzek N, Pechmann M, & Prpic N-M (2017) Expression and function

1521   of the zinc finger transcription factor Sp6–9 in the spider Parasteatoda tepidariorum.

1522   *Development genes and evolution* 227(6):389-400.

1523   151.   Heingård M, Turetzek N, Prpic N-M, & Janssen R (2019) FoxB, a new and highly

1524   conserved key factor in arthropod dorsal–ventral (DV) limb patterning. *EvoDevo* 10(1):1-

1525   16.

1526   152.   Evans GO (1992) *Principles of Acarology* (CAB International, Wallingford) p 563.

1527   153.   Gainett G*, et al.* (2020) Systemic paralogy and function of retinal determination network

1528   homologs in arachnids. *BMC Genomics* 21:811.

1529   154.   Samadi L, Schmid A, & Eriksson BJ (2015) Differential expression of retinal

1530   determination genes in the principal and secondary eyes of Cupiennius salei Keyserling

1531   (1877). *EvoDevo* 6(1):16.

155. Schomburg C*, et al.* (2015) Molecular characterization and embryonic origin of the eyes

in the common house spider Parasteatoda tepidariorum. *EvoDevo* 6(1):1-14.

156. Alberti G & Coons LB (1999) *Acari-Mites* (Wiley, New York) p 1265.

157. Patten W (1887) Eyes of molluscs and arthropods. *Journal of Morphology* 1(1):67-92.

158. Exner S (1989) *The physiology of the compound eyes of insects and crustaceans*

(Springer-Verlag GmbH & Co. KG).

159. Harzsch S*, et al.* (2006) Evolution of arthropod visual systems: development of the eyes

and central visual pathways in the horseshoe crab Limulus polyphemus Linnaeus, 1758

(Chelicerata, Xiphosura). *Developmental Dynamics* 235(10):2641-2655.

160. Alberti G & Moreno-Twose AI (2012) Fine structure of the primary eyes in

Heterochthonius gibbus (Oribatida, Heterochthoniidae) with some general remarks on

photosensitive structures in oribatid and other actinotrichid mites. *Soil Organisms*

84(2):391-408.

161. Norton RA & Fuangarworn M (2015) Nanohystricidae n. fam., an unusual,

plesiomorphic enarthronote mite family endemic to New Zealand (Acari, Oribatida).

Zootaxa 4027(2):151-204.

162. Norton RA & Franklin E (2018) Paraquanothrus n. gen. from freshwater rock pools in the

USA, with new diagnoses of Aquanothrus, Aquanothrinae, and Ameronothridae (Acari,

Oribatida). *Acarologia* 58(3):557-627.

163. Walter DE & Proctor HC (1998) Feeding behaviour and phylogeny: observations on

early derivative Acari. *Experimental & applied acarology* 22(1):39-50.

164.   Alberti G (1998) Fine structure of receptor organs in oribatid mites (Acari). *Arthropod*

1554   *biology: Contributions to morphology, ecology and systematics. Biosystematics and*

1555   *Ecology Series* 14:27-77.

1556   165.   Madge D (1965) Further studies on the behaviour of Belba geniculosa Oudms. in relation

1557   to various environmental stimuli. *Acarologia* 7(4):744-757.

1558   166.   Woodring J (1966) Color phototactic responses of an eyeless oribatid mite. *Acarologia*

1559   8(2):382-388.

1560   167.   Trägårdh I (1933) Methods of automatic collecting for studying the fauna of the soil.

1561   *Bulletin of Entomological Research* 24(2):203-214.

1562   168.   Nagata T, Koyanagi M, Tsukamoto H, & Terakita A (2010) Identification and

1563   characterization of a protostome homologue of peropsin from a jumping spider. *Journal*

1564   *of Comparative Physiology A* 196(1):51.

1565   169.   Koyanagi M, Nagata T, Katoh K, Yamashita S, & Tokunaga F (2008) Molecular

1566   evolution of arthropod color vision deduced from multiple opsin genes of jumping

1567   spiders. *Journal of Molecular Evolution* 66(2):130-137.

1568   170.   Eriksson BJ, Fredman D, Steiner G, & Schmid A (2013) Characterisation and localisation

1569   of the opsin protein repertoire in the brain and retinas of a spider and an onychophoran.

1570   *BMC evolutionary biology* 13(1):186.

1571   171.   Senthilan PR, Grebler R, Reinhard N, Rieger D, & Helfrich-Förster C (2019) Role of

1572   rhodopsins as circadian photoreceptors in the Drosophila melanogaster. *Biology* 8(1):6.

1573   172.   Senthilan PR & Helfrich-Förster C (2016) Rhodopsin 7–the unusual rhodopsin in

1574   Drosophila. *PeerJ* 4:e2427.

1575   173.   Shen WL*, et al.* (2011) Function of rhodopsin in temperature discrimination in

1576   Drosophila. *Science* 331(6022):1333-1336.

1577   174.   Vieira FG & Rozas J (2011) Comparative genomics of the odorant-binding and

1578   chemosensory protein gene families across the Arthropoda: origin and evolutionary

1579   history of the chemosensory system. *Genome Biology and Evolution* 3:476-490.

1580   175.   Sánchez-Gracia A, Vieira FG, Almeida FC, & Rozas J (2011) Comparative genomics of

1581   the major chemosensory gene families in Arthropods. *eLS*.

1582   176.   Sánchez-Gracia A, Vieira F, & Rozas J (2009) Molecular evolution of the major

1583   chemosensory gene families in insects. *Heredity* 103(3):208-216.

1584   177.   Ngoc PCT*, et al.* (2016) Complex evolutionary dynamics of massively expanded

1585   chemosensory receptor families in an extreme generalist chelicerate herbivore. *Genome*

1586   *biology and evolution* 8(11):3323-3339.

1587   178.   Croset V*, et al.* (2010) Ancient protostome origin of chemosensory ionotropic glutamate

1588   receptors and the evolution of insect taste and olfaction. *PLoS genetics* 6(8):e1001064.

1589   179.   Benton R, Vannice KS, Gomez-Diaz C, & Vosshall LB (2009) Variant ionotropic

1590   glutamate receptors as chemosensory receptors in Drosophila. *Cell* 136(1):149-162.

1591   180.   Knecht ZA*, et al.* (2016) Distinct combinations of variant ionotropic glutamate receptors

1592   mediate thermosensation and hygrosensation in Drosophila. *Elife* 5:e17879.

1593   181.   Budelli G*, et al.* (2019) Ionotropic receptors specify the morphogenesis of phasic sensors

1594   controlling rapid thermal preference in Drosophila. *Neuron* 101(4):738-747. e733.

1595   182.   Rytz R, Croset V, & Benton R (2013) Ionotropic receptors (IRs): chemosensory

1596   ionotropic glutamate receptors in Drosophila and beyond. *Insect biochemistry and*

1597   *molecular biology* 43(9):888-897.

1598   183.   Montell C (2009) A taste of the Drosophila gustatory receptors. *Current opinion in*

1599   *neurobiology* 19(4):345-353.

1600   184.   Kuwahara Y (2004) Chemical ecology of astigmatid mites. *Advances in Insect Chemical*

1601   *Ecology*, eds Cardé RT & Millar JG (Cambridge University Press, Cambridge), pp 76-

1602   109.

1603   185.   Bunnell T, Hanisch K, Hardege JD, & Breithaupt T (2011) The fecal odor of sick

1604   hedgehogs (Erinaceus europaeus) mediates olfactory attraction of the tick Ixodes

1605   hexagonus. *Journal of chemical ecology* 37(4):340.

1606   186.   Yunker C*, et al.* (1992) Olfactory responses of adultAmblyomma hebraeum andA.

1607   variegatum (Acari: Ixodiae) to attractant chemicals in laboratory tests. *Experimental &*

1608   *applied acarology* 13(4):295-301.

1609   187.   Hartmann K, Laumann M, Bergmann P, Heethoff M, & Schmelzle S (2016)

1610   Development of the synganglion and morphology of the adult nervous system in the mite

1611   Archegozetes longisetosus Aoki (Chelicerata, Actinotrichida, Oribatida). *J Morphol*

1612   277(4):537-548.

1613   188.   Altincicek B, Kovacs JL, & Gerardo NM (2012) Horizontally transferred fungal

1614   carotenoid genes in the two-spotted spider mite Tetranychus urticae. *Biology letters*

1615   8(2):253-257.

1616   189.   Crisp A, Boschetti C, Perry M, Tunnacliffe A, & Micklem G (2015) Expression of

1617   multiple horizontally acquired genes is a hallmark of both vertebrate and invertebrate

1618   genomes. *Genome biology* 16(1):1-13.

1619   190.   Gilbert HJ (2010) The biochemistry and structural biology of plant cell wall

1620   deconstruction. *Plant physiology* 153(2):444-455.

191. Latgé JP (2007) The cell wall: a carbohydrate armour for the fungal cell. *Molecular*

microbiology 66(2):279-290.

192. Faddeeva-Vakhrusheva A*, et al.* (2017) Coping with living in the soil: the genome of the

parthenogenetic springtail Folsomia candida. *BMC genomics* 18(1):493.

193. Mitreva M, Smant G, & Helder J (2009) Role of horizontal gene transfer in the evolution

of plant parasitism among nematodes. *Horizontal Gene Transfer*, (Springer), pp 517-

535.

194. Laetsch DR & Blaxter ML (2017) BlobTools: Interrogation of genome assemblies.

F1000Research 6(1287):1287.

195. Flot J-F*, et al.* (2013) Genomic evidence for ameiotic evolution in the bdelloid rotifer

Adineta vaga. *Nature* 500(7463):453-457.

196. Thorpe P, Escudero-Martinez CM, Cock PJ, Eves-van den Akker S, & Bos JI (2018)

Shared transcriptional control and disparate gain and loss of aphid parasitism genes.

Genome Biol. Evol. 10(10):2716-2733.

197. Lawrence JG (1997) Selfish operons and speciation by gene transfer. *Trends in*

microbiology 5(9):355-359.

198. Stefaniak O (1981) The effect of fungal diet on the development of Oppia nitens (Acari,

Oribatei) and on the microflora of its alimentary tract.

199. Stefaniak O (1976) The microflora of the alimentary canal of Achipteria coleoptrata

(Acarina, Oribatei).

200. McKenna DD*, et al.* (2016) Genome of the Asian longhorned beetle (Anoplophora

glabripennis), a globally significant invasive species, reveals key functional and

evolutionary innovations at the beetle–plant interface. *Genome biology* 17(1):1-18.

201.   Luxton M (1972) Studies on Oribatid Mites of a Danish Beech Wood Soil .1. Nutritional

Biology. *Pedobiologia* 12(6):434-463.

202.   Luxton M (1981) Studies on the Oribatid Mites of a Danish Beech Wood Soil .7. Energy

Budgets. *Pedobiologia* 22(2):77-111.

203.   Luxton M (1982) The Biology of Mites from Beech Woodland Soil. *Pedobiologia*

23(1):1-8.

204.   Heethoff M, Norton RA, & Raspotnig G (2016) Once Again: Oribatid Mites and Skin

Alkaloids in Poison Frogs. *J Chem Ecol* 42(8):841-844.

205.   Kuwahara Y*, et al.* (2001) Chemical ecology of astigmatid mites LIX. Neral, the alarm

pheromone of Schwiebea elongata (Banks)(Acari: Acaridae). *Journal of the Acarological*

Society of Japan 10:19-25.

206.   Sakata T & Norton RA (2003) Opisthonotal gland chemistry of a middle-derivative

oribatid mite, Archegozetes longisetosus (Acari : Trhypochthoniidae). *International*

Journal of Acarology 29(4):345-350.

207.   Sakata T, Tagami K, & Kuwahara Y (1995) Chemical ecology of oribatid mites. I. Oil

gland components of Hydronothrus crispus Aoki. *Journal of the Acarological Society of*

Japan 4(2):69-75.

208.   Sakata T (1997) Natural Chemistry of Mite Secretions. PhD-thesis (Kyoto University,

Kyoto, Japan).

209.   Raspotnig G, Schuster R, & Krisper G (2004) Citral in oil gland secretions of Oribatida

(Acari): a key component for phylogenetic analyses. *Abh Ber Naturkundemus Görlitz*

76:43-50.

210.   Brückner A & Heethoff M (2017) The ontogeny of oil gland chemistry in the oribatid

mite Archegozetes longisetosus Aoki (Oribatida, Trhypochthoniidae). *International*

Journal of Acarology 43(5):337-342.

211.   Noge K*, et al.* (2005) Biosynthesis of neral in Carpoglyphus lactis (Acari:

Carpoglyphidae) and detection of its key enzyme, geraniol dehydrogenase, by

electrophoresis. *Journal of the Acarological Society of Japan* 14:75-81.

212.   Noge K*, et al.* (2008) Geraniol dehydrogenase, the key enzyme in biosynthesis of the

alarm pheromone, from the astigmatid mite Carpoglyphus lactis (Acari: Carpoglyphidae).

The FEBS journal 275(11):2807-2817.

213.   Morita A, Mori N, Nishida R, Hirai N, & Kuwahara Y (2004) Neral (the Alarm

Pheromone) Biosynthesis via the Mevalonate Pathway, Evidenced by D-Glucose-1-13C

Feeding in Carpoglyphus lactis and 13C Incorporation into Other Opisthonotal Gland

Exudates. *Journal of Pesticide Science* 29:27-32.

214.   Kanehisa M*, et al.* (2007) KEGG for linking genomes to life and the environment.

Nucleic acids research 36(suppl_1):D480-D484.

215.   Eisenreich W, Bacher A, Arigoni D, & Rohdich F (2004) Biosynthesis of isoprenoids via

the non-mevalonate pathway. *Cellular and Molecular Life Sciences CMLS* 61(12):1401-

1426.

216.   Miziorko HM (2011) Enzymes of the mevalonate pathway of isoprenoid biosynthesis.

Arch Biochem Biophys 505(2):131-143.

217.   Oldfield E & Lin FY (2012) Terpene biosynthesis: modularity rules. *Angewandte Chemie*

International Edition 51(5):1124-1137.

218.   Breitmaier E (2006) *Terpenes: flavors, fragrances, pharmaca, pheromones* (John Wiley

1689   & Sons).

1690   219.   Degenhardt J, Köllner TG, & Gershenzon J (2009) Monoterpene and sesquiterpene

1691   synthases and the origin of terpene skeletal diversity in plants. *Phytochemistry* 70(15-

1692   16):1621-1637.

1693   220.   Trapp SC & Croteau RB (2001) Genomic organization of plant terpene synthases and

1694   molecular evolutionary implications. *Genetics* 158(2):811-832.

1695   221.   Liu W*, et al.* (2015) Utilization of alkaline phosphatase PhoA in the bioproduction of

1696   geraniol by metabolically engineered Escherichia coli. *Bioengineered* 6(5):288-293.

1697   222.   Oswald M, Fischer M, Dirninger N, & Karst F (2007) Monoterpenoid biosynthesis in

1698   Saccharomyces cerevisiae. *FEMS yeast research* 7(3):413-421.

1699   223.   Zhou J*, et al.* (2014) Engineering Escherichia coli for selective geraniol production with

1700   minimized endogenous dehydrogenation. *J Biotechnol* 169:42-50.

1701   224.   Beran F, Köllner TG, Gershenzon J, & Tholl D (2019) Chemical convergence between

1702   plants and insects: biosynthetic origins and functions of common secondary metabolites.

1703   *New Phytologist*.

1704   225.   Raspotnig G, Kaiser R, Stabentheiner E, & Leis HJ (2008) Chrysomelidial in the

1705   Opisthonotal Glands of the Oribatid Mite, Oribotritia berlesei. *Journal of Chemical*

1706   *Ecology* 34(8):1081-1088.

1707   226.   Shimizu N, Sakata D, Schmelz EA, Mori N, & Kuwahara Y (2017) Biosynthetic pathway

1708   of aliphatic formates via a Baeyer-Villiger oxidation in mechanism present in astigmatid

1709   mites. *P Natl Acad Sci USA* 114(10):2616-2621.

1710   227.   Andrews S (2010) FastQC: A quality control tool for high throughput sequence data.

1711   *Reference Source*:https://www.bioinformatics.babraham.ac.uk/projects/fastqc/.

1712   228.   Martin M (2011) Cutadapt removes adapter sequences from high-throughput sequencing

1713   reads. *EMBnet. journal* 17(1):10-12.

1714   229.   Chikhi R & Medvedev P (2014) Informed and automated k-mer size selection for

1715   genome assembly. *Bioinformatics* 30(1):31-37.

1716   230.   Vurture GW*, et al.* (2017) GenomeScope: fast reference-free genome profiling from short

1717   reads. *Bioinformatics* 33(14):2202-2204.

1718   231.   Sun H, Ding J, Piednoël M, & Schneeberger K (2018) findGSE: estimating genome size

1719   variation within human and Arabidopsis using k-mer frequencies. *Bioinformatics*

1720   34(4):550-557.

1721   232.   Marçais G & Kingsford C (2011) A fast, lock-free approach for efficient parallel

1722   counting of occurrences of k-mers. *Bioinformatics* 27(6):764-770.

1723   233.   Koren S*, et al.* (2017) Canu: scalable and accurate long-read assembly via adaptive k-mer

1724   weighting and repeat separation. *Genome research* 27(5):722-736.

1725   234.   Li H & Durbin R (2009) Fast and accurate short read alignment with Burrows–Wheeler

1726   transform. *bioinformatics* 25(14):1754-1760.

1727   235.   Walker BJ*, et al.* (2014) Pilon: an integrated tool for comprehensive microbial variant

1728   detection and genome assembly improvement. *PloS one* 9(11):e112963.

1729   236.   Smit AF & Hubley R (2008) RepeatModeler Open-1.0. *Available fom http://www.*

1730   *repeatmasker. org*.

1731   237.   Bao W, Kojima KK, & Kohany O (2015) Repbase Update, a database of repetitive

1732   elements in eukaryotic genomes. *Mobile Dna* 6(1):11.

1733   238.   Kohany O, Gentles AJ, Hankus L, & Jurka J (2006) Annotation, submission and

1734   screening of repetitive elements in Repbase: RepbaseSubmitter and Censor. *BMC*

1735   *Bioinformatics* 7(1):1-7.

1736   239.   Petersen M*, et al.* (2019) Diversity and evolution of the transposable element repertoire

1737   in arthropods with particular reference to insects. *BMC Evol. Biol.* 19(1):11.

1738   240.   Rognes T, Flouri T, Nichols B, Quince C, & Mahé F (2016) VSEARCH: a versatile open

1739   source tool for metagenomics. *PeerJ* 4:e2584.

1740   241.   Smit A, Hubley R, & Green P (1996-2010) RepeatMasker Open-3.0.

1741   242.   Ter-Hovhannisyan V, Lomsadze A, Chernoff YO, & Borodovsky M (2008) Gene

1742   prediction in novel fungal genomes using an ab initio algorithm with unsupervised

1743   training. *Genome Res.* 18(12):1979-1990.

1744   243.   Bruna T, Hoff K, Stanke M, Lomsadze A, & Borodovsky M (2020) BRAKER2:

1745   Automatic Eukaryotic Genome Annotation with GeneMark-EP+ and AUGUSTUS

1746   Supported by a Protein Database. *bioRxiv*.

1747   244.   Haas BJ*, et al.* (2008) Automated eukaryotic gene structure annotation using

1748   EVidenceModeler and the Program to Assemble Spliced Alignments. *Genome Biol.*

1749   9(1):R7.

1750   245.   Keilwagen J, Hartung F, & Grau J (2019) GeMoMa: Homology-based gene prediction

1751   utilizing intron position conservation and RNA-seq data. *Gene Prediction*, (Springer), pp

1752   161-177.

1753   246.   Chan PP & Lowe TM (2019) tRNAscan-SE: searching for tRNA genes in genomic

1754   sequences. *Gene Prediction*, (Springer), pp 1-14.

1755   247.   Quinlan AR & Hall IM (2010) BEDTools: a flexible suite of utilities for comparing

1756   genomic features. *Bioinformatics* 26(6):841-842.

1757   248.   Pruitt KD, Tatusova T, & Maglott DR (2005) NCBI Reference Sequence (RefSeq): a

1758   curated non-redundant sequence database of genomes, transcripts and proteins. *Nucleic*

1759   *acids research* 33(suppl_1):D501-D504.

1760   249.   Bairoch A & Apweiler R (2000) The SWISS-PROT protein sequence database and its

1761   supplement TrEMBL in 2000. *Nucleic acids research* 28(1):45-48.

1762   250.   Bateman A*, et al.* (2004) The Pfam protein families database. *Nucleic acids research*

1763   32(suppl_1):D138-D141.

1764   251.   Rawlings ND, Barrett AJ, & Bateman A (2010) MEROPS: the peptidase database.

1765   *Nucleic acids research* 38(suppl_1):D227-D233.

1766   252.   Cantarel BL*, et al.* (2009) The Carbohydrate-Active EnZymes database (CAZy): an

1767   expert resource for glycogenomics. *Nucleic acids research* 37(suppl_1):D233-D238.

1768   253.   Eddy SR (2011) Accelerated profile HMM searches. *PLoS computational biology*

1769   7(10):e1002195.

1770   254.   Huerta-Cepas J*, et al.* (2017) Fast genome-wide functional annotation through orthology

1771   assignment by eggNOG-mapper. *Mol. Biol. Evol.* 34(8):2115-2122.

1772   255.   Geib SM*, et al.* (2018) Genome Annotation Generator: a simple tool for generating and

1773   correcting WGS annotation tables for NCBI submission. *GigaScience* 7(4):giy018.

1774   256.   Kanehisa M & Goto S (2000) KEGG: kyoto encyclopedia of genes and genomes. *Nucleic*

1775   *acids research* 28(1):27-30.

1776   257.   Consortium GO (2004) The Gene Ontology (GO) database and informatics resource.

1777   *Nucleic acids research* 32(suppl_1):D258-D261.

1778   258.   Carlson M & Pagès H (2019) AnnotationForge: Tools for building SQLite-Based

1779   Annotation Data Packages. *R package version* 1(0).

1780   259.   Yu G, Wang L-G, Han Y, & He Q-Y (2012) clusterProfiler: an R package for comparing

1781   biological themes among gene clusters. *Omics: a journal of integrative biology*

1782   16(5):284-287.

1783   260.   Benjamini Y & Hochberg Y (1995) Controlling the False Discovery Rate - a Practical

1784   and Powerful Approach to Multiple Testing. *J Roy Stat Soc B Met* 57(1):289-300.

1785   261.   Consortium GO (2019) The gene ontology resource: 20 years and still GOing strong.

1786   *Nucleic acids research* 47(D1):D330-D338.

1787   262.   Moriya Y, Itoh M, Okuda S, Yoshizawa AC, & Kanehisa M (2007) KAAS: an automatic

1788   genome annotation and pathway reconstruction server. *Nucleic acids research*

1789   35(suppl_2):W182-W185.

1790   263.   Suzuki S, Kakuta M, Ishida T, & Akiyama Y (2014) GHOSTX: an improved sequence

1791   homology search algorithm using a query suffix array and a database suffix array. *PloS*

1792   *one* 9(8):e103833.

1793   264.   Fu L, Niu B, Zhu Z, Wu S, & Li W (2012) CD-HIT: accelerated for clustering the next-

1794   generation sequencing data. *Bioinformatics* 28(23):3150-3152.

1795   265.   Price MN, Dehal PS, & Arkin AP (2010) FastTree 2 – approximately Maximum-

1796   Likelihood trees for large alignments. *PLoS One* 5(3):e9490.

1797   266.   Capella-Gutiérrez S, Silla-Martínez JM, & Gabaldón T (2009) trimAl: a tool for

1798   automated alignment trimming in large-scale phylogenetic analyses. *Bioinformatics*

1799   25(15):1972-1973.

267. Kocot KM, Citarella MR, Moroz LL, & Halanych KM (2013) PhyloTreePruner: a

phylogenetic tree-based approach for selection of orthologous sequences for

phylogenomics. *Evol. Bioinform. Online* 9:429-435.

268. Kück P & Meusemann K (2010) FASconCAT: Convenient handling of data matrices.

Mol. Phylogenet. Evol. 56(3):1115-1118.

269. Lanfear R, Frandsen PB, Wright AM, Senfeld T, & Calcott B (2016) PartitionFinder 2:

new methods for selecting partitioned models of evolution for molecular and

morphological phylogenetic analyses. *Mol. Biol. Evol.* 34(3):772-773.

270. Nguyen L-T, Schmidt HA, Von Haeseler A, & Minh BQ (2015) IQ-TREE: a fast and

effective stochastic algorithm for estimating maximum-likelihood phylogenies.

Molecular biology and evolution 32(1):268-274.

271. Zhang C, Rabiee M, Sayyari E, & Mirarab S (2018) ASTRAL-III: polynomial time

species tree reconstruction from partially resolved gene trees. *BMC Bioinformatics*

19(6):153.

272. Dobin A*, et al.* (2013) STAR: ultrafast universal RNA-seq aligner. *Bioinformatics*

29(1):15-21.

273. Grabherr MG*, et al.* (2011) Trinity: reconstructing a full-length transcriptome without a

genome from RNA-Seq data. *Nature biotechnology* 29(7):644.

274. Haas BJ*, et al.* (2013) De novo transcript sequence reconstruction from RNA-seq using

the Trinity platform for reference generation and analysis. *Nat. Protoc.* 8(8):1494-1512.

275. Kitchen SA, Crowder CM, Poole AZ, Weis VM, & Meyer E (2015) De novo assembly

and characterization of four anthozoan (phylum Cnidaria) transcriptomes. *G3: Genes,*

genomes, genetics 5(11):2441-2452.

276.   Bray NL, Pimentel H, Melsted P, & Pachter L (2016) Near-optimal probabilistic RNA-

1824   seq quantification. *Nature biotechnology* 34(5):525-527.

1825   277.   Pimentel H, Bray NL, Puente S, Melsted P, & Pachter L (2017) Differential analysis of

1826   RNA-seq incorporating quantification uncertainty. *Nature Methods* 14(7):687.

1827   278.   R_Core_Team (2019) R: A language and environment for statistical computing. R

1828   Foundation for Statistical Computing, Vienna, Austria. 2019.

1829   279.   Babicki S*, et al.* (2016) Heatmapper: web-enabled heat mapping for all. *Nucleic acids*

1830   *research* 44(W1):W147-W153.

1831   280.   Conway JR, Lex A, & Gehlenborg N (2017) UpSetR: an R package for the visualization

1832   of intersecting sets and their properties. *Bioinformatics* 33(18):2938-2940.

1833   281.   Gladyshev EA, Meselson M, & Arkhipova IR (2008) Massive horizontal gene transfer in

1834   bdelloid rotifers. *Science* 320(5880):1210-1213.

1835   282.   Robinson JT*, et al.* (2011) Integrative genomics viewer. *Nat. Biotechnol.* 29(1):24-26.

1836   283.   Niimura Y & Nei M (2005) Evolutionary dynamics of olfactory receptor genes in fishes

1837   and tetrapods. *Proceedings of the National Academy of Sciences* 102(17):6039-6044.

1838   284.   Robertson HM & Wanner KW (2006) The chemoreceptor superfamily in the honey bee,

1839   Apis mellifera: expansion of the odorant, but not gustatory, receptor family. *Genome*

1840   *research* 16(11):1395-1403.

1841   285.   Robertson HM, Warr CG, & Carlson JR (2003) Molecular evolution of the insect

1842   chemoreceptor gene superfamily in Drosophila melanogaster. *Proceedings of the*

1843   *National Academy of Sciences* 100(suppl 2):14537-14542.

1844   286.   Katoh K & Standley DM (2013) MAFFT multiple sequence alignment software version

1845   7: improvements in performance and usability. *Molecular biology and evolution*

1846   30(4):772-780.

1847   287.   Edgar RC (2004) MUSCLE: multiple sequence alignment with high accuracy and high

1848   throughput. *Nucleic Acids Res.* 32(5):1792-1797.

1849   288.   Guindon S*, et al.* (2010) New algorithms and methods to estimate maximum-likelihood

1850   phylogenies: assessing the performance of PhyML 3.0. *Systematic biology* 59(3):307-

1851   321.

1852   289.   Lefort V, Longueville J-E, & Gascuel O (2017) SMS: smart model selection in PhyML.

1853   *Molecular biology and evolution* 34(9):2422-2424.

1854   290.   Brückner A & Heethoff M (2016) Scent of a mite: origin and chemical characterization

1855   of the lemon-like flavor of mite-ripened cheeses. *Exp Appl Acarol* 69(3):249-261.

1856   291.   Heethoff M, Laumann M, Weigmann G, & Raspotnig G (2011) Integrative taxonomy:

1857   Combining morphological, molecular and chemical data for species delineation in the

1858   parthenogenetic Trhypochthonius tectorum complex (Acari, Oribatida,

1859   Trhypochthoniidae). *Frontiers in Zoology* 8:2.

1860   292.   Cao Z*, et al.* (2013) The genome of Mesobuthus martensii reveals a unique adaptation

1861   model of arthropods. *Nature communications* 4(1):1-10.

1862