1  **Integrative genomics identifies lncRNA regulatory networks across 1,044 pediatric leukemias**
2  **and extra-cranial solid tumors**
3

Apexa Modi[1,2], Gonzalo Lopez[1], Karina L. Conkrite[1], Chun Su[3], Tsz Ching Leung[1], Sathvik Ramanan[1],

Elisabetta Manduchi[3], Matthew E. Johnson[3], Daphne Cheung[1], Samantha Gadd[4], Jinghui Zhang[5],

Malcolm A. Smith[6], Jaime M. Guidry Auvil[7], Daniela S. Gerhard[6], Soheil Meshinchi[8], Elizabeth J.

Perlman[4], Stephen P. Hunger[1,9], John M. Maris[1,9,11], Andrew D. Wells[3,11], Struan F.A. Grant[3,9,12,13],

Sharon J. Diskin[1,9,10*]

[1] Division of Oncology and Center for Childhood Cancer Research, Children's Hospital of Philadelphia,
Philadelphia, Pennsylvania 19104, USA.
12

13  [2] Genomics and Computational Biology Graduate Group, Biomedical Graduate Studies, Perelman
14  School of Medicine, University of Pennsylvania, Philadelphia, Pennsylvania 19104, USA.
15

16  [3] Center for Spatial and Functional Genomics, Children's Hospital of Philadelphia, Philadelphia,
17  Pennsylvania, USA.
18

19  [4] Department of Pathology and Laboratory Medicine, Ann & Robert H. Lurie Children's Hospital of
20  Chicago, Robert H. Lurie Cancer Center, Northwestern University, Chicago, Illinois 60208, USA.
21

22  [5] Department of Computational Biology, St Jude Children's Research Hospital, Memphis, Tennessee
23  38105, USA
24

25  [6] Cancer Therapy Evaluation Program, National Cancer Institute, Bethesda, Maryland 20892, USA.
26

27  [7] Office of Cancer Genomics, National Cancer Institute, Bethesda, Maryland 20892, USA.
28

29  [8] Clinical Research Division, Fred Hutchinson Cancer Research Center, Seattle, Washington 98109,
30  USA.
31

32  [9] Department of Pediatrics, Perelman School of Medicine at the University of Pennsylvania,
33  Philadelphia, Pennsylvania 19104, USA.
34

35  [10] Abramson Family Cancer Research Institute, Perelman School of Medicine at the University of
36  Pennsylvania, Philadelphia, Pennsylvania 19104, USA.
37

38  [11] Department of Pathology and Laboratory Medicine, Perelman School of Medicine at the University of
39  Pennsylvania, Philadelphia, Pennsylvania 19104, USA.
40

41  [12] Department of Genetics, Perelman School of Medicine at the University of Pennsylvania,
42  Philadelphia, Pennsylvania 19104, USA.
43

44  [13] Divisions of Human Genetics and Endocrinology & Diabetes, Children's Hospital of Philadelphia,
45  Philadelphia, Pennsylvania, 19104, USA.
46

48  * Corresponding Author: diskin@email.chop.edu

**Abstract**

Long non-coding RNAs (lncRNAs) play an important role in gene regulation and contribute to tumorigenesis. While pan-cancer studies of lncRNA expression have been performed for adult malignancies, the lncRNA landscape across pediatric cancers remains largely uncharted. Here, we curate RNA sequencing data for 1,044 pediatric leukemia and solid tumors and integrate paired tumor whole genome sequencing and epigenetic data in relevant cell line models to explore lncRNA expression, regulation, and association with cancer. We report a total of 2,657 robustly expressed lncRNAs across six pediatric cancers, including 1,142 exhibiting histotype-specific expression. DNA copy number alterations contributed to lncRNA dysregulation at a proportion comparable to protein coding genes. Application of a multi-dimensional framework to identify and prioritize lncRNAs impacting gene networks revealed that lncRNAs dysregulated in pediatric cancer are associated with proliferation, metabolism, and DNA damage hallmarks. Analysis of upstream regulation via cell-type specific transcription factors further implicated distinct histotype-specific and developmental lncRNAs. We integrated our analyses to prioritize lncRNAs for experimental validation and showed that silencing of *TBX2-AS1*, our top-prioritized neuroblastoma-specific lncRNA, resulted in significant growth inhibition of neuroblastoma cells, confirming our computational predictions. Taken together, these data provide a comprehensive characterization of lncRNA regulation and function in pediatric cancers and pave the way for future mechanistic studies.

Long non-coding RNAs (lncRNAs) are transcribed RNA molecules greater than 200 nucleotides in length

that do not code for proteins. These molecules account for 70% of the expressed human transcriptome

and provide a key aspect of gene regulation[1-4]. Compared to protein coding genes (PCGs), lncRNAs

typically have fewer exons, weaker conservation, and lower abundance[3]. Despite this, lncRNAs have

been shown to play significant roles in both transcriptional and post-transcriptional gene regulation[5].

LncRNAs perform these roles by physically interacting with a variety of substrates, including proteins

(transcription co-factors), RNAs (microRNA sponges), and DNA (chromatin interaction scaffolds)[1,2,6,7].

While the mechanisms and function for the majority of lncRNAs remain unknown[3,8], those that have been

experimentally characterized are involved in a variety of cellular processes[6] including gene silencing

(*ANRIL*)[9], modulation of chromatin architecture (*Xist*)[10], and pre-mRNA processing (*MALAT1*)[11].

LncRNAs are also important in development[12]. For example, the *H19* lncRNA is involved in imprinting[13],

while the well-conserved *TUNA* lncRNA controls stem cell pluripotency and lineage differentiation[14].

Dysregulation of lncRNA expression has been widely observed in cancer[3,15,16] and studies have

shown that lncRNAs play important roles in tumor initiation and progression[17]. LncRNAs can function as

tumor suppressors, such as the *PANDA* lncRNA which regulates DNA damage response in diffuse large

B-cell lymphoma[18]; however, many more lncRNAs appear to be oncogenes. Examples include the

*HOTAIR* and *PVT1* lncRNAs which promote proliferation in various cancers through tissue specific

mechanisms[19,20]. Pan-cancer analyses of lncRNA expression in adult malignancies have uncovered

many cancer-associated lncRNAs[3,15-17,21,22]. Identification of functional lncRNAs amongst the large set of

cancer-associated lncRNAs, however, remains challenging[15,23]. Current methods to identify putative

functional lncRNAs involve identifying lncRNA-specific genetic aberrations[15,16,24] or using lncRNA

expression to predict overall patient survival[16]. To more systematically address how lncRNAs drive the

pathogenesis of cancer, recent computational methods seek to assign function to these molecules based

on predicted target genes and regulatory network models. These methods have been applied to adult

malignancies and allow for more focused hypotheses to be tested[21,22].

LncRNA studies and evidence of related function in pediatric cancers have been primarily limited

to neuroblastoma (NBL)[25-30], T-lymphoblastic leukemia (T-ALL)[31,32], and more recently glioblastoma[33].

*CASC15* and *NBAT-1* are a sense-antisense lncRNA pair that map to a NBL susceptibility locus identified

by genome-wide association study[26,34]. Both lncRNAs are downregulated in high-risk NBL tumors and

have been shown to be involved in cell proliferation and differentiation[25,26]. In pediatric T-ALL, the

NOTCH-regulated lncRNA, *LUNAR1,* promotes T-ALL cell growth by sustaining IGF1 signaling[32]. To

date, it is unknown whether lncRNAs function as common drivers across multiple pediatric cancers, or if

instead, the majority of lncRNAs influence oncogenesis in a histotype-specific manner. Furthermore,

given that pediatric cancers typically arise from primitive embryonic and mesodermal cells, rather than

adult epithelial cells, it is unclear whether adult cancer lncRNA drivers will also be implicated in childhood

cancer.

Here, we perform a pan-pediatric cancer study of lncRNAs across 1,044 pediatric leukemias and

extra-cranial solid tumors[35,36]. We present the landscape of lncRNA expression across these childhood

cancers and perform integrative multi-omic analyses to assess tissue specificity, regulation, and putative

function. To validate our approach, we show that silencing of the top-prioritized NBL-specific

lncRNA, *TBX2-AS1*, impairs NBL cell growth in human-derived NBL cell line models.

**Results**

**The lncRNA landscape of pediatric cancers**

To define the repertoire of lncRNAs expressed in childhood cancers, we analyzed RNA-sequencing data

from six distinct pediatric cancer histotypes profiled through the Therapeutically Applicable Research to

Generate Effective Treatments (TARGET) project (https://ocg.cancer.gov/programs/target/data-matrix)

(**Online Methods; Supplementary Table 1).** This curated set of 1,044 leukemia and solid tumor samples

includes 280 acute myeloid leukemia (AML), 190 B-lymphoblastic leukemias (B-ALL), 244 T-

lymphoblastic leukemias (T-ALL), 121 Wilms tumors (WT), 48 extracranial rhabdoid tumors (RT), and

161 neuroblastomas (NBL) (**Fig. 1a**). Since one of our goals was to identify novel cancer-associated

lncRNAs, we performed guided *de novo* transcriptome assembly using StringTie v1.3.3[37] with the

GENCODE v19 database [38] as a gene annotation reference (**Supplementary Fig. 1**). Expressed gene

sequences that did not match exons and transcript structures of any known gene in the GENCODE v19

or RefSeq v74 databases were considered putative novel genes (**Supplementary Fig. 1, Online**

**Methods**). Of these novel genes, we identified candidate lncRNAs by using the PLEK v1 algorithm[39] to

assess non-coding potential, and then additionally filtered hits by transcript length, exon read coverage,

and genomic location (**Fig. 1a, Online Methods, Supplementary Fig. 1**). As validation of our lncRNA

discovery pipeline, we observed that 36% (87 of 242) of identified novel lncRNAs not annotated in

Gencode v19 (hg19) were indeed annotated in the more recent Gencode v29 (hg38) genome build

(**Supplementary Table 2**). To ensure that we selected robustly expressed genes in the setting of cancer

heterogeneity and sequencing variability, we applied a conservative expression cutoff of Fragments Per

Kilobase of transcript per Million mapped reads (FPKM) >1 in at least 20% of samples for each cancer.

Across all cancers there were 15,588 PCGs, 2,512 known lncRNAs, and 145 novel lncRNAs expressed,

though the total number of expressed genes varied per cancer (**Fig 1b, Supplementary Table 3**).

Principal component analysis (PCA) of lncRNA gene expression showed that blood (AML, B-ALL, T-ALL)

and solid (NBL, WT, RT) cancers form two distinct groups. Moreover, individual cancer histotypes

clustered more closely using lncRNA expression than PCG expression alone (**Supplementary Fig. 2a-**

**b**), consistent with the known tissue specific nature of lncRNA expression and function[3].

Overall, lncRNAs had lower average expression compared to PCGs resulting in fewer highly

expressed lncRNAs (**Supplementary Fig. 2c**). Between 10-100 (3.7%) lncRNAs accounted for 50% of

the total sum of lncRNA expression (**Fig. 1c**). In contrast, between 100-1000 (6.4%) PCGs accounted for

50% of the total sum of PCG expression (**Fig. 1d**). We examined the union of the top five most highly

expressed lncRNAs across pediatric cancers (total 11 lncRNAs). Some of these lncRNAs had higher

expression in the blood cancers (*MALAT1* and *RP11-386I14.4*), in the solid cancers (*H19)*, or in only one

cancer, such as *MEG3* and *RP11-386G11.10* in NBL (**Fig. 1e**). Five of these lncRNAs were among the

top 10 lncRNAs expressed across normal tissues in the Genotype-Tissue Expression (GTEx) project [40].

Specifically, *C17orf76-AS1 (LRRC75A-AS1), MALAT1, GAS5, SNHG6, SNHG8* were expressed

ubiquitously in 30 of the 49 GTEx tissues (**Supplementary Table 4**).

**Tissue specific lncRNA expression distinguishes pediatric cancers**

To evaluate more formally the tissue specific expression of lncRNAs, we annotated all genes with a tissue

specificity index (tau score)[41,42] (**Online Methods**). The established tau score ranges from 0 (ubiquitous

expression) to 1 (tissue-specific). As an example, the highly expressed lncRNA *C17orf76-AS1* yielded a

tau score of 0.296 in this study, indicating ubiquitous expression (**Supplementary Fig. 2d**). In contrast,

the highly expressed *MEG3* lncRNA, which is known to have tissue-specific expression in NBL[30,43],

yielded a tau score of 0.986 (**Supplementary Fig. 2e**). Overall, we observed that lncRNAs yielded a

higher tau score range and mean, and thus greater tissue specific expression than PCGs (t-test

$p=1.62 \times 10^{-42}$). Novel lncRNAs had the greatest tissue specific expression (t-test: vs proteins- $p=1.62 \times 10^{-42}$,

vs known lncRNAs- $p = 3.39 \times 10^{-13}$) (**Fig. 2a**). A tau score threshold of 0.8 has been suggested to

distinguish tissue specific genes[42], and using this cutoff we identified 1,142 (42%) tissue specific (TS)

lncRNAs (**Fig. 2b, Supplementary Table 5**). To assess how well TS lncRNAs distinguish cancers, we

performed clustering based on the top five highest expressed TS lncRNAs per cancer (30 total). The

expression of just these lncRNAs was sufficient to cluster samples of the same cancer type (**Fig. 2c**).

Furthermore, the blood and solid cancers separately clustered together with little expression overlap

observed between the two groups across the 30 genes (**Fig. 2c**). Finally, we identified a similar proportion

of TS lncRNAs (38%, n = 1624) across 12 adult cancers from The Cancer Genome Atlas (TCGA) (**Online

Methods**) and observed that adult cancer tissue types were also well distinguished based on the

expression of the top 5 most TS lncRNAs (**Supplementary Fig. 2f-g**).

Notably, NBL tumors expressed 2.5x more TS lncRNAs (n=522) than the cancer with the next

highest: WT (TS lncRNAs: n=211), and 10x more than AML, which had the least number of TS lncRNAs

(n=49) (**Fig. 2b**). To validate NBL's striking quantity of TS lncRNAs, we first assessed whether immune

and stromal cell infiltration[36] could be contributing to the variety of lncRNAs expressed. We ran the

ESTIMATE algorithm as previously described[36] (**Online Methods**) to determine levels of immune and

stromal cell presence in each tumor sample using expression data. Using these purity estimates, we re-

calculated each cancer's tau score and restricted our analysis to NBL samples with either 80% or 90%

purity. In both cases, we found that NBL still had the greatest number of TS lncRNAs (n =588 – NBL 90%

purity) compared to other cancers (**Supplementary Table 6**). Finally, given that the TARGET NBL RNA-

seq dataset is un-stranded, we validated our findings using stranded RNA-seq data in an independent

NBL cohort generated through the Gabriela Miller Kids First (GMKF) program (n=223). We observed that

48% of expressed lncRNAs were tissue specific in the GMKF cohort, an increase from the 31% observed

in the TARGET cohort (**Supplementary Table 6**). These results confirm lncRNA abundance in NBL and

demonstrate that the tau score robustly identifies TS lncRNAs across varying datasets.

**Somatic DNA copy number alterations impact lncRNA expression**

Many pediatric cancers are marked by a lower single nucleotide variant (SNV) and insertion-deletion

(indel) burden than observed in adult cancers[36]. Instead, large chromosomal events, such as somatic

copy number aberrations (SCNAs) and other structural variants (SVs) have been shown to dysregulate

protein coding driver genes[36,44]. However, the extent to which large chromosomal alterations impact

lncRNAs in pediatric cancers remains unknown. We thus sought to identify SCNAs and SVs using

whole genome sequencing (WGS) data from the TARGET project available for NBL (n=146), B-ALL

(n=302), AML (n=297), and WT (n=81) (**Online Methods**). We observed that NBL had the greatest

frequency of copy number events (**Supplementary Fig. 3a**). The GISTIC v2 algorithm[45] was applied

to detect regions of recurrent SCNA (q-value < 0.25). We identified 673 expressed lncRNAs

overlapping 176 significant SCNA regions across the cancers (**Supplementary Table 7**). WGS

samples with matched RNA-sequencing were then used to compare lncRNA expression in samples

with or without an SCNA event and determine significant differential expression (DE) (**Online**

**Methods**, **Supplementary Table 8**). Across all cancers, between 10-30% of expressed genes

overlapping SCNA regions showed significant differential expression based on SCNA, a proportion

that was similar for both PCGs and lncRNAs (**Fig 3a**). Altogether, there were 198 (29%) unique

lncRNAs with significant DE due to SCNA (**Supplementary Fig 3b**). The majority of the significantly

dysregulated lncRNAs were identified in the two cancers with the greatest overall number of

expressed lncRNAs, NBL and WT, and mapped to regions with highly recurrent SCNAs in those

cancers (chromosomes 1, 7, 11, and 17) (**Fig 3b**).

While SCNAs can cause the dysregulation of lncRNA expression based on gene dosage,

structural variant (SV) breakpoints within a lncRNA could cause loss or gain of function[36,44]. We utilized

WGS data to identify lncRNAs disrupted by SV breakpoints using a previously described combination

approach involving copy number read-depth and discordant junction approach[44] (**Online Methods**).

There were 650 unique expressed lncRNA genes disrupted by SVs, 89% of which were found in only

one sample (**Supplementary Fig. 4a)**. We observed 212 SV-impacted lncRNA genes located at

SCNA regions (**Fig. 3c**), and 65% of lncRNAs genes disrupted by SV breakpoints in at least five

samples were located at SCNA regions (**Supplementary Fig. 4b, Supplementary Table 9**). Indeed,

the top-ranked SV-impacted lncRNA in both NBL and WT, *MYCNOS,* associates with the disease-

driving chr2p24 amplification[46,47] (**Supplementary Fig. 4c-d**). In B-ALL, the SV-impacted lncRNAs:

*KIAA0125* and *CDKN2B-AS1 (ANRIL)* associate with the well-studied *IGH* translocation and

*CDKN2A*/*B* deletion locus (**Supplementary Fig. 4e**)[48]. The top-ranked SV-impacted lncRNA in AML,

*MIR181A1HG (MONC)*, associates with a recurrent SCNA deletion on 1q and is mildly up-regulated

in the AML dataset (p = 0.061, **Supplementary Fig. 4f**). *MIR181A1HG* (*MONC*) was described

previously as an oncogene in acute megakaryoblastic leukemia[49,50]. Finally, we observed 30 lncRNAs

with pan-cancer (n>3) expression and SV breakpoints(**Supplementary Fig. 4h**). The most number of

breakpoints across unique samples was observed in *LINC00910,* which was shown previously to be

essential for cell growth in the K562 cell line[51].

**Characterization of transcriptional network perturbation mediated by dysregulated lncRNAs**

To determine how lncRNAs may drive pediatric cancers, we examined the downstream impact of

lncRNAs on gene regulation. We focused on identifying lncRNAs that mediate transcriptional regulation

by modulating TF activity (lncRNA modulators)[52-55]. We wrote custom scripts implementing the lncMod

computational framework[56] (**Online Methods**) to first identify DE-lncRNAs and then to assess their

impact on correlated expression between a TF and its target genes[21,56] (**Fig. 4a, Online Methods**).

Across all cancers studied, we identified 313,370 unique, dysregulated lncMod triplets (lncRNA-TF-target

gene), representing 0.02-0.2% of possible triplets, which have significant correlation differences between

a TF and target gene upon lncRNA expression dysregulation (**Supplementary Table 10-11**). This

proportion was consistent with previous findings from the lncMap study in adult cancers[21], although more

triplets were identified in datasets with greater sample size (**Supplementary Table 10-11**). LncRNA

modulators were categorized into one of three categories based on their impact on TF-target gene

correlation; either the correlation was enhanced, attenuated, or inverted (**Fig 4a-b**). lncRNA modulators

have context specific function such that for different TF-target gene pairs they could exert different types

of regulation (**Supplementary Fig. 5b**). The majority of lncRNA modulators appeared to be active in only

one cancer, with only 15% (138 of 923 lncRNAs) having pan-cancer activity (n>3) (**Fig. 4c**).

To determine the biological impact of lncRNA modulators, we identified lncRNAs whose target

genes were enriched in MSigDB's Hallmark Gene Sets (HMS)[57] (Fisher's exact test, FDR < 0.1; **Online**

**Methods**). Across the majority of cancers, lncRNA modulator target genes had significant enrichment in

the proliferation, metabolism, and DNA damage hallmark categories (FDR range: 0.1 to $2.24 \times 10^{-36}$; **Fig.**

**4d**). Overall, the top-enriched hallmark pathways closely mirrored those found for lncRNA modulators in

adult cancers[22]. Consistent with its role in development and as an oncogene in certain cancers [23], the

top-enriched hallmarks for the *H19* lncRNA, dysregulated in NBL, were the EMT (development) and G2M-

checkpoint (proliferation) hallmarks (**Supplementary Fig. 5c**). The blood cancers exhibited strong

enrichment of lncRNA modulators regulating MYC targets, which has a well-established role in

leukemias[58]. Furthermore, in AML, we observed that gene targets of the myeloid-specific lncRNA,

*HOTAIRM1*, were most enriched for proliferation hallmarks (**Supplementary Fig. 5d**), consistent with

this lncRNA's known role in proliferation as an oncogene in adult AML[59].

Finally, we sought to determine potential lncRNA mechanism by identifying recurring patterns of

regulation amongst lncMod triplets. To this end, we nominated candidate lncRNA-TF associations by

ranking TF's based on the number of target genes regulated by each given TF (**Supplementary Table**

**12**). As proof-of-concept, we were able to detect known lncRNA-TF associations such as *GAS5* with

E2F4[60] (RNA-protein), and *SNHG1* with *TP53*[61] (RNA-RNA) amongst lncMod triplets in our study

(**Supplementary Fig. 5e-f**). A notable example from the hundreds of novel associations identified is

between the B-ALL specific lncRNA, *BLACE* (B-cell acute lymphoblastic leukemia expressed, tau score:

0.999) and its top associated TF, XBP1, which has known roles in pre-B-ALL cell proliferation and

tumorigenesis[62] (**Fig 4e-f**). These predictions of lncRNA transcriptional networks provide focused

avenues to elucidate the mechanisms through which lncRNAs can drive pediatric cancers.

**Defining the role of lncRNAs in childhood cancer development**

Pediatric cancers arise in the context of normal human development where cells do not differentiate as

they should, resulting in malignant cell transformation[63]. Some tumors are comprised of heterogenous

cells that resemble varying differentiation lineages with distinct transcriptomic states due to distinct super

enhancer transcription factor networks[64,65]. We sought to uncover lncRNAs associated with these varying

cell lineages as they may contribute to pediatric cancer etiology. We used NBL as a model given its

heterogeneity and two confirmed tumor cell states: the undifferentiated mesenchymal (MES) cells and

the committed adrenergic (ADRN) cells, which can interconvert[66]. Given that NBL precursor cells, the

neural crest cells, have been shown to have a more MES gene expression signature[65,66], we

hypothesized that lncRNAs correlated with an MES signature may play a role in NBL development. Using

the gene set variation analysis (GSVA) method[67] we assigned for each NBL Stage 4 sample, both a MES

and ADRN score (**Online Methods**). Using hierarchical clustering (**Supplementary Figure 6a**) we

categorized samples based on their primary gene expression phenotype as ADRN, MES, or mixed (**Fig**

**5a**). We next correlated the MES and ADRN score with lncRNA expression across NBL samples. We

observed 29 lncRNAs associated with MES samples and 21 lncRNAs associated with ADRN samples

(**Fig 5b**) (Spearman's |rho| >0.6, adj. pval < 0.01). We then performed a guilt-by-association analysis[68]

to determine the potential functional pathway for these lncRNAs based on the pathway of their correlated

protein coding genes (**Online Methods**). Gene set enrichment was performed using the gene ontology

(GO) biological processes gene set. Intriguingly, the ADRN group of lncRNAs showed enrichment for

DNA replication and cell cycle associated gene sets, whereas the MES lncRNAs were associated with

organ development and immune response (**Fig 5b**). We validated these same pathway results in an

independent analysis of the GMKF NBL cohort restricted to Stage 4 samples (n=67) (**Supplementary**

**Figure 6b**). Across both TARGET and GMKF cohorts we observed 13 lncRNAs strongly associated with

MES samples (**Supplementary Table 6c**), which warrant further study for their potential role in NBL

development.

**Identification of potential cancer driver lncRNAs via integration of epigenetic data**

To better identify ADRN lncRNAs we elucidated lncRNAs directly regulated by the known ADRN

transcription factors (TFs): MYCN, PHOX2B, HAND2, GATA3, ISL1, and TBX2[65,69]. This set of TFs, which

are co-bound and auto-regulated, are known as the core transcriptional circuitries (CRC) and drive the

ADRN cell lineage in NBL[65,69]. CRC gene regulation occurs both by direct promoter binding (**Fig. 5c-1**)

and by distal binding to either promoters (**Fig. 5c-2**) or enhancer regions (**Fig. 5c-3**) which then regulate

the gene of interest via long-range chromatin interactions[65,69-71]. CRC-bound regulatory loci were

identified from publicly available ChIP-seq data for all ADRN TFs across two MYCN-amplified NBL cell

lines: SKNBE(2)C and KELLY[69,72] (**Online Methods**). To comprehensively identify both short- and long-

range CRC gene regulation, we generated high-resolution (i.e. using 4-cutter restriction enzyme DpnII)

genome-wide promoter-focused Capture C[73] in the NBL cell line NB1643. After pinpointing gene

promoters interacting with CRC TF bound regulatory loci (promoters or enhancers) (**Fig. 5c, Online**

**Methods**), we identified 547 lncRNA genes associated with the NBL CRC (**Fig 5d, Supplementary**

**Table 13**), with only 249 of these lncRNA genes being bound by CRC TFs within their promoter regions.

We further distinguished 313 ADRN lncRNAs based on differential expression (DE) between ADRN and

MES samples (**Fig 5d, Supplementary Table 14**). The *TBX2-AS1* DE-lncRNA was highly correlated to

the CRC TF: *TBX2* (Pearson's r=0.77), and both are up-regulated in ADRN samples (**Fig 5e**). CRC

binding is observed at both the shared promoter region of *TBX2* and *TBX2-AS1* and at an interacting

distal enhancer (**Fig 5f**). TBX2 was recently shown to be involved in NBL cell proliferation,[74] but the role

of *TBX2-AS1* in NBL is unknown.

To further demonstrate the utility of this epigenetic based prioritization, we applied the same

method to T-ALL, which also has a well-established set of CRC TFs (TAL1, MYB, GATA3, and RUNX1)[71].

We used available ChIP-seq and ChIA-PET data for the TAL1 mutated T-ALL cell lines, Jurkat and

CCRF-CEM, to identify loci bound by the T-ALL CRC TF's[71] (**Online Methods**). We not only identified

the known leukemia associated lncRNA PVT1, but also 9 other T-ALL CRC lncRNAs prioritized based

on correlation with T-ALL PCGs and differential expression associated with a previously defined TAL1-

subgroup[75] (**Supplementary Figure 7, Supplementary Table 13-15**). Taken together, this novel data

integration method nominates multiple lncRNAs with previously unknown function for further study as

potential driver genes in pediatric cancer.

**Integrative multi-omic analysis prioritizes *TBX2-AS1* as a candidate functional lncRNA in NBL**

To obtain a comprehensive prioritization of candidate functional lncRNAs for each cancer histotype,

we integrated information for (1) tissue specific expression, (2) dysregulation due to DNA copy number

aberration, and (3) regulation by CRC TFs (**Supplementary Table 16**). Here, we focus on the NBL

cohort since this cancer has data available for all of the prioritization steps (**Supplementary Table**

**17**). The top ranked lncRNA in NBL was *MEG3*, which has a known role in both NBL and other

cancers[43]. The next notable lncRNA, *TBX2-AS1,* is up-regulated due to chromosome 17q gain (**Fig**

**6a**), has NBL-specific expression (tau score: *TBX2-* 0.807, *TBX2-AS1-* 0.86; **Supplementary Fig. 8a**),

and is co-regulated with *TBX2*. TBX2 has been shown to drive NBL proliferation via the *FOXM1*/*E2F1*

gene regulatory network[72] and we hypothesized that *TBX2-AS1* may play a similar role because

predictions from our lncMod analysis indicated that *TBX2-AS1* impacts E2F targets and G2M

checkpoint genes (**Fig. 6b**). Furthermore, the TFs primarily impacted by TBX2 knockdown[72], MYBL2

and E2F1, were found to have the most target genes predicted to be regulated by *TBX2-AS1* (**Fig 6c-**

**d**). Evidence for this association was further supported by the correlation (Spearman's rho > 0.4)

between *TBX2-AS1* and *TBX2*'s target TFs, including: *FOXM1*, *E2F1*, and *MYBL2* (**Supplementary**

**Fig. 8b**). While the strong correlation between *TBX2-AS1* and *TBX2* may confound our predictions, a

previous study showed positionally conserved lncRNAs[59], such as *TBX2-AS1*, often regulate their

neighboring developmental TFs (TBX2) and can play roles in genome organization and cancer[59].

Based on the promising *in silico* evidence, we prioritized *TBX2-AS1* for experimental study.

**Silencing of *TBX2-AS1* inhibits cell growth of neuroblastoma cells**

We assessed the role of *TBX2-AS1* using human-derived NBL cell line models. First, we evaluated

*TBX2-AS1* expression across 38 NBL cell lines using RNA-seq[76] (**Supplementary Fig 8c**). Expression

of *TBX2* and *TBX2-AS1* were subsequently validated in eight cell lines using RT-qPCR

(**Supplementary Fig. 8d**). We selected NLF and SKNSH models for further study based on their high

*TBX2-AS1* expression and differing expression levels of *TBX2*. Silencing of *TBX2-AS1* using small

interfering RNA (siRNA) achieved 92% and 63% reduction of *TBX2-AS1* expression in NLF and

SKNSH, respectively (**Fig. 6e**). We also observed down-regulation of TBX2 protein levels in the

siTBX2-AS1 treated cells for both cell lines (**Fig. 6f**). Given the known role of TBX2 in NBL cell

proliferation[74], we measured cell growth of siTBX2-AS1 treated NBL cells to determine if *TBX2-AS1*

has similar function. When the non-targeting control (siNTC) treated cells reached confluence, the

siTBX2-AS1 treated cell index was reduced by 42.6% and 36.8% (n=3, p < 0.01) in the NLF and

SKNSH cell line, respectively (**Fig. 6g-h, Supplementary Fig. 8e**). Live cell imaging using the

IncuCyte revealed changes in cell morphology for siTBX2-AS1 treated NLF cells, featuring an

appearance of disrupted cell to cell adhesion and elongated cell body (**Supplementary Fig. 8f**). To

identify pathways impacted by *TBX2-AS1* knockdown, we performed total RNA sequencing in triplicate

of NLF cells and compared gene expression in control (siNTC) vs siTBX2-AS1 treated cells

(**Supplementary Fig. 8g**). Gene set enrichment analysis (GSEA) of the 364 significantly up-regulated

genes (log-fold change > 1.5, adj pval < 0.1) revealed enrichment (FDR < 0.1) for hallmarks associated

with inflammation including: TNFA signaling and interferon gamma response (**Supplementary Table**

**18**). Across the 544 down-regulated genes, E2F target genes hallmark was most enriched. To

determine whether differentially expressed genes shared common regulation, we used the iRegulon

program[77], to search for TF motifs and ENCODE ChIP-seq tracks upstream of genes (**Online**

**Methods**). Using a normalized enrichment score (NES) of at least 3, we observed motif enrichment

for the neuronal differentiation repressor REST and the RFX family of transcription factors in 59% of

siTBX2-AS1 up-regulated genes (**Fig. 6i**). In 42% of downregulated genes, the top enriched TFs were

MYBL2 and E2F1, corroborating our GSEA results. Moreover, both the growth assays and gene

expression profiling confirmed our lncMod results, which showed that *TBX2-AS1* impacts NBL

proliferation by modulating target genes of E2F1 and MYBL2 (**Fig. 6b-d**). These data thus demonstrate

the utility of our integrative lncRNA characterization and prioritization approach for future validation

experiments across all cancers considered in this study. Furthermore, we uncovered a functional role for

*TBX2-AS1* in NBL proliferation likely mediated via the regulation of TBX2 and its known target genes:

E2F, MYBL2, and REST[72].

**Discussion**

LncRNAs have emerged as important regulators of gene expression and their dysregulation can impact

key cancer pathways and drive tumorigenesis[1-4]. Despite this, relatively few lncRNAs have been

experimentally characterized and the landscape of lncRNA expression across pediatric cancers has been

previously unknown. In this study, we explored lncRNA expression, cancer association, and regulatory

networks across 1,044 pediatric leukemias and solid tumors, representing six different cancer types. The

breadth of samples and cancer types included allowed for robust identification of novel, cancer-specific,

and developmental lncRNAs. Furthermore, we used systems modelling to identify expression patterns

for both up- and downstream lncRNA gene regulation. Altogether we provide multi-dimensional insight

into the predicted biological and functional relevance of lncRNAs by integrating WGS, ChIP-seq,

chromatin capture, and predictions of transcriptional networks.

Analysis of the lncRNA landscape across pediatric cancers revealed the histotype and context-

specific nature of lncRNAs. We report a total of 2,657 robustly expressed lncRNAs across the six cancer

types studied. This number is notably smaller than reports from pan-cancer studies of adult

malignancies[15,17], likely due to the smaller number of cancer types studied here and conservative

expression threshold applied. However, similar to our findings in adult cancers, 43% (1,142/ 2,657) of

expressed lncRNAs exhibited tissue-specific (TS) expression across pediatric cancers. Indeed, lncRNAs

had significantly greater tissue specificity than protein coding genes, making them more ideal candidates

as biomarkers. Currently there is one lncRNA, *PCA3*, that is FDA-approved as a biomarker for prostate

cancer[78] and multiple trials investigating ncRNAs in cancer prognostics are underway[79]. In this study, the

top five most TS lncRNAs per cancer were sufficient to differentiate each cancer histotype. Furthermore,

we identify lncRNAs specific to distinct cell lineages within NBL, suggesting there is potential for lncRNAs

to be used as highly sensitive markers to differentiate cancer subtypes more accurately.

Typically, investigation of lncRNA dysregulation involves comparing lncRNA expression between

cancer and normal control samples and is an analysis that amply yields adult-cancer associated

lncRNAs[15]. However, the lack of normal expression controls for the majority of pediatric cancers[36] is a

major complication in defining pediatric cancer-associated lncRNAs. To overcome this, we leveraged

information about how pediatric cancers are epigenetically regulated. In particular, NBL, is composed of

two cells lineages representing different development stages and each with distinct super-enhancer

transcription factor networks. Given the tie between organogenesis and tumorigenesis in pediatric

cancer[63], we hypothesized that lncRNAs associated with these cell states may also be involved in NBL

development. After correlation and pathway analysis, we discovered that lncRNAs associated with the

mesenchymal cell lineage had enrichment for organogenesis gene sets, while adrenergic-associated

lncRNAs were predicted to be involved in proliferation based on enrichment for DNA replication and cell

cycle gene sets. The majority of NBL samples have cells with an adrenergic gene expression signature,

which could suggest that ADRN lncRNAs are major drivers of disease and thus potential therapeutic

targets. To better identify these ADRN lncRNAs, we integrated ChIP-sequencing of core regulatory

(CRC) transcription factors for ADRN cells with our expression data to identify cancer driver lncRNAs.

CRC TFs bind to cell-type-specific enhancers and regulate the expression of cell-type-specific genes[80].

By taking advantage of this information we were able to prioritize lncRNAs likely to be important for cancer

cell identity based on CRC TF regulation. CRC TFs have been well defined for NBL and T-ALL[69,71];

however the fact that they largely bind enhancer regions necessitated that we also use chromatin

interaction data to accurately determine regulated genes. Incorporation of these datasets allowed us to

identify 2-fold more CRC regulated lncRNAs in NBL and 3-fold in T-ALL as compared to using just ChIP-

seq data alone, which restricts lncRNA identification to those with CRC TFs bound at their promoter.

Notably, there were ten common CRC-regulated lncRNAs between NBL and T-ALL, and an important

next step for further identification of pan-pediatric cancer associated lncRNAs is application of this novel

analysis to a broader set of pediatric cancers.

While upstream regulation can help nominate cancer-associated lncRNAs, determining the

mechanism through which dysregulated lncRNAs impact downstream target genes is also crucial.

However, prediction of lncRNA function is limited given that very few lncRNA mechanisms have been

fully established and lncRNAs lack conserved sequence and structure[81]. Many studies instead use

correlated protein coding gene expression as a proxy to define lncRNA pathways, but this approach often

results in many false positives and does not provide mechanistic insight[81]. To address this, we used the

lncMod method[21,56] to model the functional mechanism of dysregulated lncRNAs by examining correlated

changes in transcription factor to target gene regulation. We used motif presence and regression analysis

to identify TF-target gene relationships, though future studies will be strengthened by incorporating TF

ChIP-seq data, when it becomes more widely available for pediatric cancers. Nevertheless, we were able

to successfully associate lncRNAs to TFs with known interactions, such as SNHG1 with TP53, while also

providing a prioritized list of novel associations that serve as a starting point for future experimental

studies such as RIP/MS[82] and ChiRP-seq[83]. Finally, while our lncMod analysis was focused on

transcriptional regulation, the addition of microRNA binding and RNA-binding protein data, as utilized in

adult cancers[22], is an important next step in understanding how lncRNAs impact post-transcriptional

regulation in pediatric cancers.

Our study delineated high confidence lncRNA expression across pediatric cancers within the

restrictions set by the sequencing depth and RNA-seq type available per cancer dataset. We required

RNA-seq samples included in our study to have at least 10 million reads and read length of at least 75

bp; and with the exception of the T-ALL samples, all samples were poly-A selected. Future studies

involving total RNA-seq, greater sequencing depth, and longer read sizes could capture a larger diversity

and more accurate set of expressed lncRNAs by accounting for non-polyadenylated genes and

identifying scarcer or temporally expressed lncRNAs. Nevertheless, our high confidence set of lncRNAs

are very likely to be functional given that low or rare expression can be an indicator of transcriptional

noise[84]. In addition to having a limited number of RNA matched WGS samples, the Complete Genomics

short read technology limits the detection of structural variants based on size as previously described[36,44].

The use of long-read sequencing and greater sequencing depth in future studies will enable more

accurate copy number and structure variant detection in pediatric cancers.

Finally, multi-dimensional integration of our computational predictions resulted in the nomination

of functionally relevant lncRNAs in each pediatric cancer. We annotated tissue specificity, copy number,

pathway, and likely targets for these lncRNAs, providing a solid foundation for mechanistic studies. As

proof-of-principle, we demonstrate that the top-prioritized tissue-specific and copy number dysregulated

lncRNA, *TBX2-AS1,* impacts NBL cell growth, validating our approach, while transcriptomic profiling

corroborated our pathway predictions. Knockdown of *TBX2-AS1*, showed downregulation of genes

regulated by E2F1 and MYBL2, the same TFs impacted upon TBX2 knockdown[72]. Future studies could

reveal whether *TBX2-AS1* modulates TBX2 through direct binding or by impacting transcriptional

regulation at their shared locus. *TBX2-AS1* was previously shown to be among a group of lncRNAs which

are positionally conserved and near developmental associated TFs[59]. This group of lncRNAs and their

neighboring TFs,  typically have tissue specific expression, can be involved in cancer development, and

affect each other's expression[59], all of which we observed for TBX2 and *TBX2-AS1*. Together these

genes contribute to the proliferative state of NBL cells and could have potential as novel therapeutic

targets.

Altogether, this study provides a comprehensive characterization of lncRNAs across pediatric

cancers and serves as a rich resource for future mechanistic studies; these data may aid in the selection

of cancer biomarkers and candidate therapeutic lncRNA targets.

**Online Methods**

**RNA-seq data processing.** A comprehensive RNA-seq analysis pipeline was used on all samples

(Supplementary Table 1, Supplementary Fig 1). First FASTQC was run on all samples and any samples

that had a Phred score < 30 for more than 25% of read bases were removed. Samples were then aligned

using STAR_2.4.2a [85] with the following parameters: "STAR --runMode alignReads --runThreadN 10 --

twopassMode Basic --twopass1readsN -1 --chimSegmentMin 15 --chimOutType WithinBAM –genomeDir

X --genomeFastaFiles ucsc.hg19.fa --readFilesIn fasta1 fasta2 --readFilesCommand zcat --outSAMtype

BAM   SortedByCoordinate   --outFileNamePrefix   X   --outSAMstrandField   intronMotif   --quantMode

TranscriptomeSAM   GeneCounts   --sjdbGTFfile   gencode.v19.annotation.gtf   --sjdbOverhang   X."   To

assess the quality of the aligned RNA-seq data we ran MultiQC [86], and removed samples with < 70%

uniquely mapped reads and < 10 million mapped reads.

**Gene/transcript mapping and quantification.** To map reads to genes and quantify gene expression

we ran StringTie 1.3.3 [37]. StringTie involves three steps, first quantifying expression of both known and

novel gene transcripts using an annotation guided approach. We used the Gencode v19 gene annotation

to guide gene detection.1) "stringtie bamfile -G gencode.v19.annotation_stringtie.gtf -B --rf -o out.gtf -A

gene_abund.tab -C cov_refs.gtf -p 10. " In the second step, StringTie merges the gene annotation across

all samples such that there is a uniform annotation for known and novel gene transcripts in one

transcriptome   gtf   file.   2)   "stringtie   All_PanTARGET_PreMerge_StringTie_Files.txt   --merge   -G

gencode.v19.annotation_stringtie.gtf   -o   StringTie_PanCancer_AllMergedTranscripts.gtf."   Finally,

StringTie is run again to quantify expression using the PanTarget transcriptome gtf file and de novo gene

transcript detection is turned off. 3) "stringtie bamfile -G StringTie_PanCancer_AllMergedTranscripts.gtf

-B -e --rf -o out.gtf -A gene_abund.tab -C cov_refs.gtf -p 10"

**Comparison of pan-TARGET transcriptome with reference annotation.** Novel transcripts were

assigned as an isoform of a known gene based on exonic overlap (>50% by bp) with genes in either the

GENCODE v19 or RefSeq v74 databases using custom Python scripts. Any remaining novel transcripts

were assigned as novel genes (MSTRG_Merged.# or MSTRG.#) based on overlapping exon positions.

Novel genes were further filtered based on read coverage, in that we required that at least one transcript

for a novel gene have more than one exon with at least 5 reads in at least 20% of samples per cancer.

High confidence novel genes were required to have at least 3 exons. Finally, for all transcripts (known

and novel), to obtain gene level quantification, transcript FPKM and count values were summed to get a

gene level value.

**Prediction of novel gene coding potential and lncRNA gene annotation.** We predicted coding

potential of novel transcripts using the PLEK v1 algorithm tool [39]. PLEK uses a support vector machine

(SVM) for a binary classification model to distinguish a lncRNA versus a coding mRNA. The features

used as input for the SVM are calibrated k-mer usage frequencies of a transcript's sequence. PLEK has

previously been validated on RefSeq mRNAs and GENCODE lncRNAs (the main reference annotations

used in our study) and has achieved >90% accuracy in predicting gene coding potential [39]. To further

delineate lncRNAs, we removed any predicted novel non-coding transcripts that were < 200bp (sum of

total exon length). We updated the gene type of GENCODE v19 genes with the gene type of genes that

had matching gene names in GENCODE v29. Additionally we filtered out lncRNA genes that have been

deprecated in Gencode v29. Finally, some lncRNA genes in Gencode v19, have both a lncRNA and small

RNA transcript. For these 147 cases we did not include the small RNA transcript when summing gene

transcripts to obtain gene level expression.

**Tissue specific gene expression.** The tau score, a measure of the tissue specific expression of a gene

was calculated as described by Yanai et. al[41]. The formula for the score is listed below. $x_i$ is defined as

the mean expression of a gene in a particular cancer and n is the total number of cancers considered, in

this case n = 6.

$$\tau = \frac{\sum_{i=1}^{n}(1-\hat{x_i})}{n-1}; \hat{x_i} = \frac{x_i}{\max\limits_{1 \leq x \leq n}(x_i)}$$

**CNV detection, processing, and impact on gene expression.** Copy number calls were made by

Complete Genomics (CGI) from WGS for NBL, WT, AML, and B-ALL. We used CGI

files"somaticCnvDetailsDiploidBeta" containing ploidy estimates and tumor/blood coverage along 2kb

bins across the genome. To create segmentation files, we used custom scripts to reformat CGI coverage

data to meet requirements of the "copynumber" R bioconductor package as previously described[44]. We

used the winsorize function in this package, which performs data smoothing and segmentation via a

piecewise constant segmentation (pcf) algorithm (kmin =2 and gamma= 1000). Segmentation files were

visualized     using     the     R     package     svpluscnv     (https://github.com/ccbiolab/svpluscnv)

https://doi.org/10.1093/bioinformatics/btaa878. We then ran GISTIC2.0, using segmentation data as

inputs and parameters: "GISTIC2 -v 30 -refgene hg19 -genegistic 1 -smallmem 1 -broad 1 -twoside 1 -

brlen 0.98 -conf 0.90 -armpeel 1 -savegene 1 -gcm extreme -js 2 -rx 0". To determine which genes copy

number impacts, we intersected CNV regions listed in the "all_lesions.conf_90.txt" file from GISTIC output

with gene positions. We used section 1 from the "all_lesions.conf_90.txt" file to assign a binary descriptor

to each gene as either being not amplified or deleted (CNV-no) if the sample had actual copy gain 0 for

the region containing the gene. We assigned CNV-yes if the region containing the gene was amplified or

deleted, which included samples with actual copy gain 1 or 2, where 1 indicates low level copy number

aberration (exceeds low threshold of copy number: 1: 0.1<t< 0.9) and 2 indicates a high level of copy

number aberration, CNV exceeds high threshold (t>0.9) according to GISTIC. To determine CNV impact

on gene expression, we assessed differential expression of the gene in samples from the two groups

(CNV yes or no) using Wilcoxon rank sum test (p < 0.01). Genes were considered to have evidence of

differential expression due to copy number if the absolute value of the log2 fold change between the two

groups was > 0.58 and p < 0.05.

**Structural variant detection and filtering.** Structural variants were identified from WGS as previously

described[44]. Somatic sequence junctions that were completely absent in the normal genome are reported

by Complete Genomics (CGI) in the somaticAllJunctionsBeta file. To obtain a high confidence set of

junctions, where there is a likely true physical connection between the left and right sections of a junction,

the following filtering was applied by CGI to obtain the highConfidenceSomaticAllJunctionsBeta.

1) DiscordantMatePairAlignments ≥ 10 (10 or more discordant mate pairs in cluster
2) JunctionSequenceResolve = Y (local de novo assembly is successful)
3) Exclude interchromosomal junction if present in any genomes in baseline samples
(FrequencyInBaseline > 0)
4) Exclude the junction if overlap with known underrepresented repeats
(KnownUnderrepresentedRepeat = Y): ALR/Alpha, GAATGn, HSATII, LSU_rRNA_Hsa, and
RSU_rRNA_Hsa
5) Exclude the junction if the length of either of the side sections is less than 70 base pairs.

Further filtering of these high confidence structural variants included removing rare/common germline

variants that passed the CGI filters. We used the Database of Genomic Variants (DGV v. 2016-05-15,

GRCh37) in order to remove SVs that had at least 50% reciprocal overlap with DGV annotated common

events and were type matched.

**Structural variant analysis.** To obtain a comprehensive landscape of SVs we combined both the

sequence junction and copy number read depth approaches to identify SVs, with co-localizing break

points being orthogonally validated. Recurrence of SVs was considered based on overlap with genes

from our pan-pediatric cancer transcriptome. Genomic overlap between SVs and genes was determined

using the bedtools intersect tool (default parameters). Variants were assigned to genes based on if the

sequence junction (left/right position) + 100 bp overlapped gene coordinates +/- 2.5kb. Genes were then

ranked based on the number of unique samples per cancer with a SV breakpoint.

**Gene signature analysis.** We obtained a list of genes associated with the mesenchymal (MES) and

adrenergic (ADRN) NBL cell types from GEO (GSE90805). We then used the GSVA R package[67] with

the Poisson kernel (kcdf) parameter to assign a score per sample representing the total expression

enrichment of genes associated with either the MES or ADRN cell types. We performed hierarchical

clustering to divide NBL samples into three groups (MES, ADRN or mixed phenotype) based on

expression of MES and ADRN genes using the pheatmap R package and cutting the dendrogram at n=3.

We correlated the MES and ADRN score with lncRNA expression across Stage 4 NBL TARGET cohort

and GMKF cohort samples separately and identified lncRNAs as having significant correlation based on

absolute value Spearman's rho > 0.6. These lncRNAs were then labeled as MES or ADRN based on

significant correlation with either the MES or ADRN score. We next repeated score correlation with PCGs.

We performed a guilt-by-association analysis assigning MES/ADRN PCGs and by association their

correlated MES/ADRN  lncRNAs (Spearman rho > 0.5) to pathways using Fisher exact test, FDR < 0.1

for gene sets in the gene ontology (GO) biological processes collection.

**ChIP-seq data analysis.** To determine which lncRNAs are regulated by transcription factors involved in

the core regulatory circuitry (CRC) we utilized previously generated and analyzed histone and

transcription factor ChIP-sequencing data for NBL and T-ALL. For NBL, we used peak files for our

previously generated histone ChIP-seq data of: H3K27ac, H3K4me1, H3K4me3 for the BE(2)C cell line[87],

available on GEO: GSE138315. We downloaded raw sequencing files for CRC transcription factor ChIP-

seq data for MYCN, PHOX2B, HAND2, GATA3, TBX2, and ISL1 for the BE(2)C and KELLY cell lines

from GEO: GSE94822[69] and selected peaks with q-value < 0.001 for further analysis. We identified

regions in the genome where at least 4/6 of the transcription factors overlapped. This was obtained using

the homer mergePeaks tool: "mergePeaks -d 1000 -cobound 6 bed_file1… bed_file6" and the resulting

coBoundBy4 output file. For the T-ALL CRC we obtained overlapping CRC transcription factor loci for

TAL1, GATA3, and RUNX1 from the study by Sanda et. al[71], GEO: GSE29181 for both the Jurkat and

CCRF-CEM cell lines and integrated ChIP-seq data for the MYB transcription factor from GEO:

GSE59657[70], only available in the Jurkat line. We selected loci for further analysis if they were bound by

TAL1, GATA3, and RUNX1 as previously annotated by Sanda et. al.

**Identification of CRC transcription factor regulated genes.** To identify genes regulated by the NBL

or T-ALL CRC we considered CRC TF binding at both the gene's promoter and other regulatory region

interacting with the gene's promoter. We first overlapped CRC regions using bedtools intersect with gene

transcript promoter regions, which we defined as 3000bp upstream and downstream of the transcripts

first exon. For NBL, we then utilized the promoter-focused Capture C data, inclusive of all interactions

within 1Mb on the same chromosome, to identify genomic regions that were both bound by NBL CRC

TFs and interacting with a gene's promoter. To determine this, we used bedtools intersect to determine

overlap (minimum 1bp) between CRC bound loci with loci involved in chromatin interactions. From these

regions, we determined which interacting regions corresponded with a lncRNA promoter region. We

performed a similar analysis in T-ALL, however we utilized publicly available SMC1 (cohesin) ChIA-PET

data available on the ENCODE project to consider chromatin interactions.

**Promoter-focused Capture C data generation.** High resolution promoter-focused Capture C was

performed in the neuroblastoma cell line, NB1643, (untreated) in triplicate. Cell fixation, 3C library

generation, capture C, and sequencing was performed as described by Chesi et. al (2019) and Su et al

(2020). For each replicate, $10^7$ fixed cells were centrifuged to cell pellets and split to 6 tubes for a pre-

digestion incubation with 0.3%SDS, 1x NEB DpnII restriction buffer, and dH2O for 1hr at 37ºC shaking

at 1,000rpm. A 1.7% solution of Triton X-100 was added to each tube and shaking was continued for

another hour.10 ul of DpnII (NEB, 50 U/µL) was added to each sample tube and continued shaking for 2

days. 100uL Digestion reaction was then removed and set aside for digestion efficiency QC.The

remaining samples were heat inactivated incubated at 1000 rpm in a MultiTherm for 20 min, at 65°C to

inactivate the DpnII, and cooled on ice for 20 additional minutes. Digested samples were ligated with 8

uL of T4 DNA ligase (HC ThermoFisher, 30 U/µL) and 1X ligase buffer at 1,000 rpm overnight at 16°C

.The ligated samples were then de-crosslinked overnight at 65°C with Proteinase K (20 mg/mL, Denville

Scientific) along with pre-digestion and digestion control. Both controls and ligated samples were

incubated for 30 min at 37°C with RNase A (Millipore), followed by phenol/chloroform extraction, ethanol

precipitation at -20°C, then the 3C libraries were centrifuged at 3000 rpm for 45 min at 4°C to pellet the

samples. The pellets of 3C libraries and controls were resuspended in 300uL and 20µL dH2O,

respectively, and stored at −20°C. Sample concentrations were measured by Qubit. Digestion and

ligation efficiencies were assessed by gel electrophoresis on a 0.9% agarose gel and also by quantitative

PCR (SYBR green, Thermo Fisher).

Isolated DNA from 3C libraries was quantified using a Qubit fluorometer (Life technologies), and

10 µg of each library was sheared in dH2O using a QSonica Q800R to an average fragment size of

350bp.QSonica settings used were 60% amplitude, 30s on, 30s off, 2 min intervals, for a total of 5

intervals at 4 °C. After shearing, DNA was purified using AMPureXP beads (Agencourt). DNA size was

assessed on a Bioanalyzer 2100 using a DNA 1000 Chip (Agilent) and DNA concentration was checked

via Qubit. SureSelect XT library prep kits (Agilent) were used to repair DNA ends and for adaptor ligation

following the manufacturer protocol. Excess adaptors were removed using AMPureXP beads. Size and

concentration were checked again by Bioanalyzer 2100 using a DNA 1000 Chip and by Qubit fluorometer

before hybridization. One microgram of adaptor-ligated library was used as input for the SureSelect XT

capture kit using manufacturer protocol and custom-designed 41K promoter Capture-C probe set. The

quantity and quality of the captured libraries were assessed by Bioanalyzer using a high sensitivity DNA

Chip and by Qubit fluorometer. SureSelect XT libraries were then paired-end sequenced on Illumina

NovaSeq 6000 platform (51bp read length) at the Center for Spatial and Functional Genomics at CHOP.

**Promoter-focused Capture C data analysis.** Paired-end reads from each replicated were pre-

processed using the HICUP pipeline (v0.5.9), with bowtie2 as aligner and hg19 as the reference genome.

The unique ditags output from HiCUP were further processed by the chicagoTools bam2chicago.sh script

before significant promoter interaction calling. Significant promoter interactions at 1-DpnII fragment

resolution were called using CHiCAGO (v1.1.8) with default parameters except for binsize set to 2500.

Significant interactions at 4-DpnII fragment resolution were also called using CHiCAGO with artificial

baitmap and rmap files in which DpnII fragments were concatenated *in silico* into 4 consecutive fragments

using default parameters except for removeAdjacent set to False. Interactions with a CHiCAGO score >

5 in either 1-fragment or 4-fragment resolution were considered as significant interactions. The significant

interactions were finally converted to ibed format in which each line represents a physical interaction

between fragments.

**Differential gene expression analysis for T-ALL subtypes.** We identified differentially expressed

genes using the DESeq2 tool. We compared expression between the TAL1 subgroup and non-TAL1

subgroup, defined by Liu, et al[75]. We ran DESeq2 using default parameters and considered genes as

significantly differentially expressed if their absolute value of the log2 fold change was > 0.58 and their

Benjamini-Hoschberg adjusted-p value was < 0.01.

**lncMod implementation: transcription factor target gene regulation.** We developed custom Python

scripts to implement the general framework of the lncMod method. The first part of this framework

involved determining transcription factor target gene regulation specific to each cancer. Target genes

here are defined as any protein coding or lncRNA gene and excludes pseduogenes and small RNAs.

Given that ChIP-seq binding profiles for the majority of transcription factors were not available for tissues

associated with each of these cancers we instead used transcription factor motif analysis as a proxy. We

utilized motifs in the JASPAR database[88] and predictions of binding across the genome determined by

FIMO and available in the UCSC genome database:

http://expdata.cmmt.ubc.ca/JASPAR/downloads/UCSC_tracks/2018/hg19/tsv/.             For          each

transcript we determined potential regulatory transcription factors based on the presence of predicted

binding motifs in the gene promoter region. Promoter regions were defined as regions 3000 bp upstream

and downstream of the transcript's first exon. Next we selected transcription factors based on their

expression in each cancer and then performed linear regression considering the expression of the

transcription factor and target gene specific to each cancer. We adjusted the false discovery rate due to

multiple testing using the Benjamini-Hochberg method and selected TF-target gene pairs with

significantly associated expression (adjusted p-value < 1e-5).

**Identification of lncRNA modulators.** To identify transcriptional perturbation, we first delineated genes

(TF, target genes, or lncRNAs) that had high expression variance (IQR > 1.5). For each differentially

expressed lncRNA in each cancer we calculated the following, as has been done in previous

studies[21,22,56]. For a given cancer and given lncRNA we sorted samples in the cancer based on the given

lncRNAs expression (low to high). We then determined the correlation (Spearman's rho) between the

expression of all transcription factor and target gene pairs previously identified in the given cancer. This

correlation was calculated for the 25% of samples with the lowest lncRNA expression and separately for

the 25% of samples with the highest expression for the given lncRNA. To ensure that we observed TF-

target gene regulation we required that the correlation between the TF-target pair in either the low or high

lncRNA expressing group was at least R>0.4. We only further evaluated the lncRNA TF-target gene

triplet if the correlation difference between the low and high lncRNA expression group was R>0.45. To

formally compare the correlation difference we first normalized the correlation using the Fisher r to z

transformation. Then we calculated the rewiring score, z-statistic, as previously described [21], which is

used to describe the degree of regulation change between the TF and target gene.

$$F(R) = \frac{1}{2} \ln \frac{1+R}{1-R}$$

$$rewire_{TF-gene} = P\left(|X| \le \left|\frac{F(R_{high})-F(R_{low})}{\sqrt{\frac{1.06}{n_{high}-3}+\frac{1.06}{n_{low}-3}}}\right|\right), X \sim N(0,1)$$

As a departure from what is described by Li et. al (lncMod method)[56], we used permutation analysis to

robustly assess the significance of the rewire score in the context of multiple hypothesis testing as

described by Sham et. al[89,90]. We randomly shuffled target gene expression (TF-target gene pair labels)

and calculated the rewire score P value across all TF-target gene pairs per given lncRNA. We kept the

smallest observed P value and repeated the permutation 100 times. This empirical frequency distribution

of the smallest P values was then compared to the P value in our real data to calculate an empirical

adjusted P value (adj P value) as given by the formula below, where r is the number of permutations

where the smallest P value are less than our actual P value and n is the number of permutations.

$$adj\ Pvalue_j = \frac{1 + (\#\ permutations\ where\ q \le p_j)}{1 + (\#\ permutations)}$$

The lncRNA-TF-target gene triplets, with adjusted p < 0.1 were considered significant. Datasets with

smaller sample sizes had lower statistical power and thus fewer significant triplets. Triplets were then

classified into three patterns based on correlation changes between the low and high expressing lncRNA

group: increased correlation – enhanced, decreased correlation – attenuated, and inverted – positive to

negative correlation and vice versa. We annotated lncRNA target genes as cancer genes based on if

they were listed in the COSMIC database or a complied list from Chiu et. al[22].

**Cell lines and reagents.** NBL cell lines were obtained from the American Type Tissue Culture Collection

(ATCC) and grown in RPM1-1640 with HEPES, L-glutamine and phenol red, supplemented with 10%

FBS, 1% L-glutamine in an incubator at 37°C with 5% $CO_2$. Cell line identity was confirmed biennially

through genotyping and confirmation of STR (short tandem repeat) profiles, while routine testing for

Mycoplasma contamination was confirmed to be negative.

**siRNA and growth assays.** The NBL cell lines, NLF and SKNSH, were plated in a 96-well RTCES

microelectronic sensor array (ACEA Biosciences, San Diego, CA, USA). Cell density measurements

were made every hour and were normalized to 24 hours post-plating (at transfection time). We used

siRNAs to knockdown the expression of genes in NLF and SKNSH. The siRNAs utilized were either a

non-targeting negative control siRNA (Silencer™ Select Negative Control siRNA, cat #4390843), TBX2-

AS1 Silencer™ Select siRNA (cat # n514841), and SMARTpool: ON-TARGETplus PLK1 siRNA (cat # L-

003290-00-0010). Transfection of cells was done using the DharmaFECT 1 transfection reagent (cat #

T-2001-02). siRNA at a concentration of 50nM and 2% (NLF) and 4% (SKNSH) DharmaFECT was added

to RPMI medium without 10% FBS or any antibiotic separately and then incubated at room temperature

for 5 minutes. The siRNA medium was then added to the DharmaFECT and incubated for another 20

minutes to form a complex. This solution was then mixed with our normal growth media and applied to

cells 24 hours after they had been initially plated. All experiments were repeated in triplicate, with

technical replicates (n=3) being averaged per biological replicate.

**Real time quantitative PCR.** Total RNA was extracted from NBL cells using miRNeasy kit (Qiagen) and

the provided protocol for animal cells. The concentration of RNA was determined with the Nanodrop

(Thermo Scientific). cDNA synthesis was performed using the SuperScript™ First-Strand Synthesis

System for RT-PCR using the SuperScript™ reverse transcriptase (Invitrogen). 5-20ng of cDNA were

mixed with the TaqMan Universal PCR Master Mix (Thermo Fisher Scientific) and TaqMan

probes/primers for either TBX2-AS1 (Hs00417285_m1) or the house keeping gene, HPRT1

(Hs02800695_m1). Gene expression from these reactions were measured using RT-qPCR and TBX2-

AS1 expression was normalized to HPRT1 expression.

**NLF gene knockdown expression profiling.** Total RNA was isolated from the NLF cell line 48 hours

post treatment with siTBX2-AS1 and non-targeting control samples, siNTC, (three biological replicates

per condition) and 1000 ng/sample was used as input for library preparation with the TruSeq Stranded

mRNA Sample Prep Kit from Illumina (with Ribo-Zero treatment). RNA-seq libraries were sequenced on

the Nextseq 500 at depth 10 million reads per sample minimum. Library prep and sequencing was

performed by the Sidney Kimmel Cancer Center Genomics Facility of Thomas Jefferson University.

Sample and read quality was assessed using FastQC and reads were aligned and mapped using the

same methods as described above for TARGET cancer samples. Genes were retained if at least one

sample had expression greater than 0 FPKM. To identify differentially expressed genes between siNTC

and siTBX2-AS1 treated cells, we used the DESeq2 method with default parameters. Differentially

expressed genes were annotated based on absolute value log fold change > 1.5 and Benjamini-Hochberg

adjusted p-value < 0.1. Gene set enrichment analysis (GSEA) was performed across samples using the

MsigDB Hallmarks gene sets and significantly enriched gene sets with FDR q-val < 0.1 were retained.

Up-stream co-regulators of differentially expressed genes were identified using default parameters from

the iRegulon program part of the Cytoscape suite.

**Data Availability**

771
All TARGET RNA and DNA-sequencing data analyzed in this study are available through the database

of Genotypes and Phenotypes (dbGaP; https://www.ncbi.nlm.nih.gov/gap/) under study-id phs000218

and accession number phs000467. GMKF RNA-sequencing data are available through dbGAP study

accession phs001436.v1.p1. Neuroblastoma cell line RNA-sequencing data analyzed in this study are

available through GEO at accessions GSE89413. NBL histone ChIP-seq and transcription factor ChIP-

seq data used in this study are both available through GEO at accessions: GSE138315 and GSE94822,

respectively. T-ALL transcription factor ChIP-seq data and SMC1 ChIA-PET data are available through

GEO at accessions GSE29181, GSE59657, and GSE68977.

780
781
**Acknowledgements**

the Children's Oncology Group Chair's grant CA098543 and with federal funds from the National Cancer

Institute, National Institutes of Health, under Contract No. HHSN261200800001E to S.J.D and Complete

Genomics. Promoter Capture C studies were funded by the Center for Spatial and Functional Genomics

(A.D.W and S.FA.G) at CHOP. S.FA.G. was supported by the Daniel B. Burke Endowed and Chair for

Diabetes Research and R01 HG010067.

791
792
**Author Contributions**
794
A.M. and S.J.D. conceived and designed the study. M.A.S., J.M.G.A, D.S.G., E.J.P, S.M., S.P.H., S.J.D.

and J.M.M. generated the TARGET data. K.L.C., M.E.J., S.J.D., A.D.W. and S.F.A.G. generated

promoter-focused capture C data. E.M. C.S, and A.M. analyzed promoter-focused capture C data. A.M.,

G.L., S.R. analyzed TARGET data. A.M., K.L.C., T.C.L. and D.C. performed *TBX2-AS1* experiments.

S.J.D. supervised the study. A.M. and S.J.D drafted the manuscript. All authors edited and approved the

manuscript.

801
802
803

## References

1.  Rinn, J.L. & Chang, H.Y. Genome regulation by long noncoding RNAs. *Annu Rev Biochem* **81**, 145-66 (2012).
2.  Bonasio, R. & Shiekhattar, R. Regulation of transcription by long noncoding RNAs. *Annual review of genetics* **48**, 433-55 (2014).
3.  Iyer, M.K. *et al.* The landscape of long noncoding RNAs in the human transcriptome. *Nature genetics* **47**, 199-208 (2015).
4.  Gil, N. & Ulitsky, I. Regulation of gene expression by cis-acting long non-coding RNAs. *Nat Rev Genet* **21**, 102-117 (2020).
5.  Dykes, I.M. & Emanueli, C. Transcriptional and Post-transcriptional Gene Regulation by Long Non-coding RNA. *Genomics Proteomics Bioinformatics* **15**, 177-186 (2017).
6.  Marchese, F.P., Raimondi, I. & Huarte, M. The multidimensional mechanisms of long noncoding RNA function. *Genome Biol* **18**, 206 (2017).
7.  Villegas, V.E. & Zaphiropoulos, P.G. Neighboring gene regulation by antisense long non-coding RNAs. *International journal of molecular sciences* **16**, 3251-66 (2015).
8.  Kopp, F. & Mendell, J.T. Functional Classification and Experimental Dissection of Long Noncoding RNAs. *Cell* **172**, 393-407 (2018).
9.  Kotake, Y. *et al.* Long non-coding RNA ANRIL is required for the PRC2 recruitment to and silencing of p15(INK4B) tumor suppressor gene. *Oncogene* **30**, 1956-62 (2011).
10. Engreitz, J.M. *et al.* The Xist lncRNA exploits three-dimensional genome architecture to spread across the X chromosome. *Science* **341**, 1237973 (2013).
11. Tripathi, V. *et al.* The nuclear-retained noncoding RNA MALAT1 regulates alternative splicing by modulating SR splicing factor phosphorylation. *Mol Cell* **39**, 925-38 (2010).
12. Perry, R.B. & Ulitsky, I. The functions of long noncoding RNAs in development and stem cells. *Development* **143**, 3882-3894 (2016).
13. Monnier, P. *et al.* H19 lncRNA controls gene expression of the Imprinted Gene Network by recruiting MBD1. *Proc Natl Acad Sci U S A* **110**, 20693-8 (2013).
14. Lin, N. *et al.* An evolutionarily conserved long noncoding RNA TUNA controls pluripotency and neural lineage commitment. *Mol Cell* **53**, 1005-19 (2014).
15. Yan, X. *et al.* Comprehensive Genomic Characterization of Long Non-coding RNAs across Human Cancers. *Cancer Cell* **28**, 529-540 (2015).
16. Du, Z. *et al.* Integrative genomic analyses reveal clinically relevant long noncoding RNAs in human cancer. *Nat Struct Mol Biol* **20**, 908-13 (2013).
17. Lanzós, A. *et al.* Discovery of Cancer Driver Long Noncoding RNAs across 1112 Tumour Genomes: New Candidates and Distinguishing Features. *Scientific Reports* **7**, 1-16 (2017).
18. Wang, Y. *et al.* Discovery and validation of the tumor-suppressive function of long noncoding RNA PANDA in human diffuse large B-cell lymphoma through the inactivation of MAPK/ERK signaling pathway. *Oncotarget* **8**, 72182-72196 (2017).
19. Hajjari, M. & Salavaty, A. HOTAIR: an oncogenic long non-coding RNA in different cancers. *Cancer Biol Med* **12**, 1-9 (2015).
20. Onagoruwa, O.T., Pal, G., Ochu, C. & Ogunwobi, O.O. Oncogenic Role of PVT1 and Therapeutic Implications. *Front Oncol* **10**, 17 (2020).
21. Li, Y. *et al.* LncMAP: Pan-cancer Atlas of long noncoding RNA-mediated transcriptional network perturbations. *Nucleic Acids Research* **46**, 1113-1123 (2018).
22. Chiu, H.S. *et al.* Pan-Cancer Analysis of lncRNA Regulation Supports Their Targeting of Cancer Genes in Each Tumor Context. *Cell Rep* **23**, 297-312 e12 (2018).
23. Huarte, M. The emerging role of lncRNAs in cancer. *Nature medicine* **21**, 1253-61 (2015).
24. Xu, Y. *et al.* Identification and comprehensive characterization of lncRNAs with copy number variations and their driving transcriptional perturbed subpathways reveal functional significance for cancer. *Brief Bioinform* (2019).

25.  Mondal, T. *et al.* Sense-Antisense lncRNA Pair Encoded by Locus 6p22.3 Determines
Neuroblastoma Susceptibility via the USP36-CHD7-SOX9 Regulatory Axis. *Cancer Cell* **33**,
417-434.e7 (2018).
26.  Russell, M.R. *et al.* CASC15-S is a tumor suppressor lncRNA at the 6p22 neuroblastoma
susceptibility locus. *Cancer Res* **75**, 3155-3166 (2016).
27.  Pandey, G.K. *et al.* The Risk-Associated Long Noncoding RNA NBAT-1 Controls
Neuroblastoma Progression by Regulating Cell Proliferation and Neuronal Differentiation.
*Cancer Cell* **26**, 722-737 (2014).
28.  Sahu, D. *et al.* Co-expression analysis identifies long noncoding RNA SNHG1 as a novel
predictor for event-free survival in neuroblastoma. *Oncotarget* **7**, 58022-58037 (2016).
29.  Mazar, J. *et al.* The long non-coding RNA GAS5 differentially regulates cell cycle arrest and
apoptosis through activation of BRCA1 and p53 in human neuroblastoma. *Oncotarget* **5**, 6589-
6607 (2016).
30.  Rombaut, D. *et al.* Integrative analysis identifies lincRNAs up- and downstream of
neuroblastoma driver genes. *Sci Rep* **9**, 5685 (2019).
31.  Ngoc, P.C.T. *et al.* Identification of novel lncRNAs regulated by the TAL1 complex in T-cell
acute lymphoblastic leukemia. *Leukemia* **32**, 2138-2151 (2018).
32.  Trimarchi, T. *et al.* Genome-wide mapping and characterization of Notch-regulated long
noncoding RNAs in acute leukemia. *Cell* **158**, 593-606 (2014).
33.  Liu, Y., Liu, H. & Zhang, D. Identification of novel long non-coding RNA in diffuse intrinsic
pontine gliomas by expression profile analysis. *Oncol Lett* **16**, 6401-6406 (2018).
34.  McDaniel, L.D. *et al.* Common variants upstream of MLF1 at 3q25 and within CPZ at 4p16
associated with neuroblastoma. *PLoS Genet* **13**, e1006787 (2017).
35.  Downing, J.R. *et al.* The Pediatric Cancer Genome Project. *Nat Genet* **44**, 619-22 (2012).
36.  Ma, X. *et al.* Pan-cancer genome and transcriptome analyses of 1,699 paediatric leukaemias
and solid tumours. *Nature* **555**, 371-376 (2018).
37.  Pertea, M. *et al.* StringTie enables improved reconstruction of a transcriptome from RNA-seq
reads. *Nature biotechnology* **33**, 290-5 (2015).
38.  Frankish, A. *et al.* GENCODE reference annotation for the human and mouse genomes. *Nucleic*
*Acids Res* **47**, D766-D773 (2019).
39.  Li, A., Zhang, J. & Zhou, Z. PLEK: a tool for predicting long non-coding RNAs and messenger
RNAs based on an improved k-mer scheme. *BMC Bioinformatics* **15**, 311 (2014).
40.  Consortium, G.T. Human genomics. The Genotype-Tissue Expression (GTEx) pilot analysis:
multitissue gene regulation in humans. *Science* **348**, 648-60 (2015).
41.  Yanai, I. *et al.* Genome-wide midrange transcription profiles reveal expression level
relationships in human tissue specification. *Bioinformatics* **21**, 650-9 (2005).
42.  Kryuchkova-Mostacci, N. & Robinson-Rechavi, M. A benchmark of gene expression tissue-
specificity metrics. *Brief Bioinform* **18**, 205-214 (2017).
43.  Dong, K., Tang, W. & Dong, R. MEG3, HCN3 and linc01105 influence proliferation and
apoptosis of neuroblastoma cells via HIF-1 alpha and p53 pathway. *Pediatric Blood and Cancer*
**63**, S194 (2016).
44.  Lopez, G. *et al.* Somatic structural variation targets neurodevelopmental genes and identifies
SHANK2 as a tumor suppressor in neuroblastoma. *Genome Res* **30**, 1228-1242 (2020).
45.  Mermel, C.H. *et al.* GISTIC2.0 facilitates sensitive and confident localization of the targets of
focal somatic copy-number alteration in human cancers. *Genome Biol* **12**, R41 (2011).
46.  Pugh, T.J. *et al.* The genetic landscape of high-risk neuroblastoma. *Nat Genet* **45**, 279-84
(2013).
47.  Gadd, S. *et al.* A Children's Oncology Group and TARGET initiative exploring the genetic
landscape of Wilms tumor. *Nat Genet* **49**, 1487-1494 (2017).

48.    Harvey, R.C. *et al.* Identification of novel cluster groups in pediatric high-risk B-precursor acute
lymphoblastic leukemia with gene expression profiling: correlation with genome-wide DNA copy
number alterations, clinical characteristics, and outcome. *Blood* **116**, 4874-84 (2010).
49.    Emmrich, S. *et al.* LincRNAs MONC and MIR100HG act as oncogenes in acute
megakaryoblastic leukemia. *Mol Cancer* **13**, 171 (2014).
50.    Gruber, T.A. & Downing, J.R. The biology of pediatric acute megakaryoblastic leukemia. *Blood*
**126**, 943-9 (2015).
51.    Liu, Y. *et al.* Genome-wide screening for functional long noncoding RNAs in human cells by
Cas9 targeting of splice sites. *Nat Biotechnol* (2018).
52.    Zhao, X. *et al.* CTCF cooperates with noncoding RNA MYCNOS to promote neuroblastoma
progression through facilitating MYCN expression. *Oncogene*, 1-12 (2015).
53.    Ng, S.Y., Bogu, G.K., Soh, B. & Stanton, L.W. The long noncoding RNA RMST interacts with
SOX2 to regulate neurogenesis. *Molecular Cell* **51**, 349-359 (2013).
54.    Tseng, Y.Y. *et al.* PVT1 dependence in cancer with MYC copy-number increase. *Nature* **512**,
82-6 (2014).
55.    Jeon, Y. & Lee, J.T. YY1 tethers Xist RNA to the inactive X nucleation center. *Cell* **146**, 119-33
(2011).
56.    Li, Y. *et al.* Identification and characterization of lncRNA mediated transcriptional dysregulation
dictates lncRNA roles in glioblastoma. *Oncotarget* **7**, 45027-45041 (2016).
57.    Liberzon, A. *et al.* The Molecular Signatures Database (MSigDB) hallmark gene set collection.
*Cell Syst* **1**, 417-425 (2015).
58.    Delgado, M.D. & Leon, J. Myc roles in hematopoiesis and leukemia. *Genes Cancer* **1**, 605-16
(2010).
59.    Amaral, P.P. *et al.* Genomic positional conservation identifies topological anchor point RNAs
linked to developmental loci. *Genome Biol* **19**, 32 (2018).
60.    Wang, M. *et al.* Long noncoding RNA GAS5 promotes bladder cancer cells apoptosis through
inhibiting EZH2 transcription. *Cell Death Dis* **9**, 238 (2018).
61.    Zhao, Y. *et al.* Long non-coding RNA (lncRNA) small nucleolar RNA host gene 1 (SNHG1)
promote cell proliferation in colorectal cancer by affecting P53. *Eur Rev Med Pharmacol Sci* **22**,
976-984 (2018).
62.    Kharabi Masouleh, B. *et al.* Mechanistic rationale for targeting the unfolded protein response in
pre-B acute lymphoblastic leukemia. *Proc Natl Acad Sci U S A* **111**, E2219-28 (2014).
63.    Federico, S., Brennan, R. & Dyer, M.A. Childhood cancer and developmental biology a crucial
partnership. *Curr Top Dev Biol* **94**, 1-13 (2011).
64.    Cao, J. *et al.* A human cell atlas of fetal gene expression. *Science* **370**(2020).
65.    Boeva, V. *et al.* Heterogeneity of neuroblastoma cell identity defined by transcriptional
circuitries. *Nat Genet* **49**, 1408-1413 (2017).
66.    van Groningen, T. *et al.* Neuroblastoma is composed of two super-enhancer-associated
differentiation states. *Nat Genet* **49**, 1261-1266 (2017).
67.    Hanzelmann, S., Castelo, R. & Guinney, J. GSVA: gene set variation analysis for microarray
and RNA-seq data. *BMC Bioinformatics* **14**, 7 (2013).
68.    Signal, B., Gloss, B.S. & Dinger, M.E. Computational Approaches for Functional Prediction and
Characterisation of Long Noncoding RNAs. *Trends in Genetics* **32**, 620-637 (2016).
69.    Durbin, A.D. *et al.* Selective gene dependencies in MYCN-amplified neuroblastoma include the
core transcriptional regulatory circuitry. *Nat Genet* **50**, 1240-1246 (2018).
70.    Mansour, M.R. *et al.* An oncogenic super-enhancer formed through somatic mutation of a
noncoding intergenic element. *Science* **346**, 1373-1377 (2014).
71.    Sanda, T. *et al.* Core transcriptional regulatory circuit controlled by the TAL1 complex in human
T cell acute lymphoblastic leukemia. *Cancer Cell* **22**, 209-21 (2012).

72.   Verboom, K. *et al.* A comprehensive inventory of TLX1 controlled long non-coding RNAs in T-
cell acute lymphoblastic leukemia through polyA+ and total RNA sequencing. *Haematologica*
**103**, e585-e589 (2018).
73.   Chesi, A. *et al.* Genome-scale Capture C promoter interactions implicate effector genes at
GWAS loci for bone mineral density. *Nat Commun* **10**, 1260 (2019).
74.   Decaesteker, B. *et al.* TBX2 is a neuroblastoma core regulatory circuitry component enhancing
MYCN/FOXM1 reactivation of DREAM targets. *Nat Commun* **9**, 4866 (2018).
75.   Liu, Y. *et al.* The genomic landscape of pediatric and young adult T-lineage acute lymphoblastic
leukemia. *Nat Genet* **49**, 1211-1218 (2017).
76.   Harenza, J.L. *et al.* Transcriptomic profiling of 39 commonly-used neuroblastoma cell lines. *Sci*
*Data* **4**, 170033 (2017).
77.   Janky, R. *et al.* iRegulon: from a gene list to a gene regulatory network using large motif and
track collections. *PLoS Comput Biol* **10**, e1003731 (2014).
78.   Bourdoumis, A. *et al.* The novel prostate cancer antigen 3 (PCA3) biomarker. *Int Braz J Urol* **36**,
665-8; discussion 669 (2010).
79.   Slack, F.J. & Chinnaiyan, A.M. The Role of Non-coding RNAs in Oncology. *Cell* **179**, 1033-1055
(2019).
80.   Chen, Y., Xu, L., Lin, R.Y., Muschen, M. & Koeffler, H.P. Core transcriptional regulatory
circuitries in cancer. *Oncogene* (2020).
81.   Zhang, X. & Ho, T.T. Computational Analysis of lncRNA Function in Cancer. *Methods Mol Biol*
**1878**, 139-155 (2019).
82.   Scheibe, M., Butter, F., Hafner, M., Tuschl, T. & Mann, M. Quantitative mass spectrometry and
PAR-CLIP to identify RNA-protein interactions. *Nucleic Acids Res* **40**, 9897-902 (2012).
83.   Chu, C., Qu, K., Zhong, F.L., Artandi, S.E. & Chang, H.Y. Genomic maps of long noncoding
RNA occupancy reveal principles of RNA-chromatin interactions. *Mol Cell* **44**, 667-78 (2011).
84.   Hon, C.C. *et al.* An atlas of human long non-coding RNAs with accurate 5′ ends. *Nature* **543**,
199-204 (2017).
85.   Dobin, A. *et al.* STAR: ultrafast universal RNA-seq aligner. *Bioinformatics* **29**, 15-21 (2013).
86.   Ewels, P., Magnusson, M., Lundin, S. & Kaller, M. MultiQC: summarize analysis results for
multiple tools and samples in a single report. *Bioinformatics* **32**, 3047-8 (2016).
87.   Upton, K. *et al.* Epigenomic profiling of neuroblastoma cell lines. *Sci Data* **7**, 116 (2020).
88.   Khan, A. *et al.* JASPAR 2018: update of the open-access database of transcription factor
binding profiles and its web framework. *Nucleic Acids Res* **46**, D260-D266 (2018).
89.   Sham, P.C. & Purcell, S.M. Statistical power and significance testing in large-scale genetic
studies. *Nat Rev Genet* **15**, 335-46 (2014).
90.   Wagner, B.D., Zerbe, G.O., Mexal, S. & Leonard, S.S. Permutation-based adjustments for the
significance of partial regression coefficients in microarray data analysis. *Genet Epidemiol* **32**, 1-
8 (2008).
991

**FIGURE LEGENDS**

**Fig 1: Pan-pediatric transcriptome characterization.**

**a.** Overview of pan-pediatric cancer RNA-seq dataset and schematic of data processing and filtering.

Reads from RNA-seq fastq files were aligned using the STAR algorithm and then gene transcripts were

mapped in a guided *de novo* manner and quantified via the StringTie algorithm. Genes were considered

novel if they did not have transcript exon structures matching genes in the GENCODE v19 or RefSeq

v74 databases. Novel genes were assigned as lncRNAs based on length >200bp and non-coding

potential calculated using the PLEK algorithm. Transcripts with low expression (FPKM <1 in >80%

samples) were not considered for further analysis. **b.** Pie graph showing the quantity of expressed and

robustly expressed protein coding genes, GENCODE/RefSeq annotated lncRNAs, and novel lncRNAs.

High confidence expressed genes are distinguished from all expressed genes. Adjoining schematic gives

overview of additional data types that were integrated with transcriptome data: WGS, ChIP-seq, and

chromatin capture. Listed are the analyses used to elucidate lncRNAs with functional roles in pediatric

cancer. **c.** Cumulative expression plots comparing the number of lncRNAs and (**d**) protein coding genes,

respectively, that constitute the total sum of gene expression (FPKM) per pediatric cancer. **e.** Percentage

of total lncRNA expression (FPKM) accounted for by the union of top five expressed lncRNAs per cancer

(total 11 lncRNAs).

1011

1012

**Fig 2: lncRNAs exhibit tissue specific expression that can distinguish cancers.**

**a.** Tissue specificity index (tau score) which ranges from 0 (ubiquitously expressed) to 1 (tissue specific)

is plotted for genes across three gene types: protein coding genes, lncRNAs, and novel lncRNAs. Table

shows the tau score range and mean per gene type. **b.** Number of tissue specific known and novel

lncRNAs in each cancer as defined by tissue specific gene threshold: tau score > 0.8. **c.** Heatmap

showing the hierarchically clustered gene expression for the top five most tissue specific lncRNAs per

cancer, ranked by highest tau score. Samples from each cancer cluster together based on expression of

these genes alone.

1021

1022

**Fig 3: A similar proportion of lncRNAs and protein coding genes are dysregulated due to SCNA**

**a.** The proportion of protein coding and lncRNA genes that have significant differential expression due

SCNA, separated by copy number type (amplification or deletion). The number of genes found in SCNA

loci is shown per cancer. Genes were evaluated to have differential expression due to copy number using

the Wilcoxon rank sum test (p-value < 0.05) and  log |fold change| > 1.5), comparing samples with no

SCNA to samples with low/high SCNA as defined by GISTIC scores. **b.** The number of differentially

expressed lncRNAs per chromosome and per cancer, distinguished by color. Chromosome 1 and 17 had

the most dysregulated lncRNAs associating with the greater frequency of SCNA on these chromosomes

across cancers. **c.** Number of samples with structural variant breakpoints in or near (+/- 2.5kb) lncRNAs

and that are also located in copy number regions, stratified by amplification or deletion status of the locus.

1033

1034

**Fig 4: lncRNA modulators impact transcriptional networks involving proliferation.**

**a.** Schematic that shows the three ways (attenuate, enhance, or invert) in which differentially expressed

lncRNA modulators can impact transcription factor and target gene relationships. lncRNA modulators are

associated with a TF-target gene pair based on a significant difference between TF-target gene

expression correlation in samples with low lncRNA expression (lowest quartile) vs samples with high

lncRNA expression (highest quartile). **b.** The proportion of lncRNA modulator types associated with

significantly dysregulated lncRNA modulator- TF-target gene (lncMod) triplets. The number of

significantly dysregulated lncMod triplets is listed per cancer. **c.** Number of lncRNA modulators genes

that are common in lncMod triplets across cancers. Common lncRNA modulator genes tend to have a

lower tau score compared to lncRNA modulators only associated with one cancer. **d.** Gene set

enrichment using the MSigDB Hallmark gene set, of target genes associated with lncRNA modulators in

each cancer (Fisher's exact test, FDR < 0.1). **e.** Transcription factors associated with the B-ALL

expression specific lncRNA, *BLACE*, ranked based on number of regulated target genes. **f.** Expression

heatmap of *BLACE* and the target genes of the XBP1 transcription factor, grouped by associated hallmark

gene set, in samples within the bottom and top quartiles of *BLACE* expression in B-ALL.

1050

1051

**Fig 5: Identification of lncRNAs associated with distinct neuroblastoma cell states**

**a.** The MES and ADRN signature score for TARGET NBL samples, with each sample labeled with either

ADRN, Mixed, or MES phenotype based on clustering analysis. **b.** Heatmap of the expression of lncRNAs

that have significant correlation with either the MES or ADRN score (|r| >0.6, pval < 0.01). lncRNAs were

correlated with protein coding genes on the same chromosome and subsequent gene set enrichment

analysis was performed for MES and ADRN protein coding genes separately. **c.** Schematic of how ADRN

associated CRC regulated genes are identified using ChIP-seq and chromatin interaction data. We

identified lncRNAs based on three types of regulation. 1) CRC transcription factors binding directly at the

promoter of the lncRNA. 2) CRC TFs bind an enhancer region that interacts with a lncRNA promoter. 3)

CRC TFs bind the promoter of a different gene and this promoter interacts with a lncRNA promoter. CRC

TF binding was identified from ChIP-seq data, while enhancer-promoter and promoter-promoter

interactions were identified from chromatin capture data. **d.** Filtering of lncRNAs expressed in NBL based

on CRC TF regulation and differential expression based on sample phenotypes (ADRN or MES). **e.**

Expression of *TBX2* and *TBX2-AS1* stratified by NBL sample phenotype (ADRN or MES). **f.** ChIP-seq

tracks for histone marks and CRC transcription factors in the NBL cell line: BE(2)C, and promoter capture

C chromatin interactions in NBL cell line: NB1643, at the *TBX2*/*TBX2-AS1* locus.

1068

1069

**Fig 6: The *TBX2-AS1* lncRNA plays a role in neuroblastoma proliferation by modulating TBX2**

**a.** Expression of *TBX2* and *TBX2-AS1* in NBL tumor samples with and without 17q gain. **b.** The top

MSigDB Hallmarks enriched across targets genes (p-value < 0.01) regulated by *TBX2-AS1* as predicted

from lncMod analysis. **c.** The transcription factors with most target genes regulated by *TBX2-AS1* as

predicted from lncMod analysis. **d.** Expression of gene targets of the E2F1 transcription factor that are

enriched for proliferation hallmarks, in samples with low and high *TBX2* and *TBX2-AS1* expression. *TBX2*

expression is highly correlated with that of *TBX2-AS1* (Pearson's r=0.77). **e.** siRNA knockdown efficiency

of *TBX2-AS1* in the NBL cell line: NLF is 91% and in the SKNSH cell 63% knockdown was achieved. **f.**

Western blot analysis of TBX2 in siTBX2 and siTBX2-AS1 treated NLF and SKNSH cell lines. **g.**

Representative image of cell growth (as measured by RT-Ces assay) of the NBL cell lines: NLF and **h.**

SKNSH. Cell index is normalized to time point when siRNA reagent is added at 24 hours post cell plating.

**i.** Results from iRegulon analysis for genes that are up- or down-regulated upon siTBX2-AS1 treatment

in NLF. Number of genes shown in Venn diagram with evidence of motif or ChIP-seq binding of the listed

transcription factors.

# Figure 1

**Fig 1: Pan-pediatric transcriptome characterization.**
**a.** Overview of pan-pediatric cancer RNA-seq dataset and schematic of data processing and filtering. Reads from RNA-seq fastq files were aligned using the STAR algorithm and then gene transcripts were mapped in a guided *de novo* manner and quantified via the StringTie algorithm. Genes were considered novel if they did not have transcript exon structures matching genes in the GENCODE v19 or RefSeq v74 databases. Novel genes were assigned as lncRNAs based on length >200bp and non-coding potential calculated using the PLEK algorithm. Transcripts with low expression (FPKM <1 in >80% samples) were not considered for further analysis. **b.** Pie graph showing the quantity of expressed and robustly expressed protein coding genes, GENCODE/RefSeq annotated lncRNAs, and novel lncRNAs. High confidence expressed genes are distinguished from all expressed genes. Adjoining schematic gives overview of additional data types that were integrated with transcriptome data: WGS, ChIP-seq, and chromatin capture. Listed are the analyses used to elucidate lncRNAs with functional roles in pediatric cancer. **c.** Cumulative expression plots comparing the number of lncRNAs and (**d**) protein coding genes, respectively, that constitute the total sum of gene expression (FPKM) per pediatric cancer. **e.** Percentage of total lncRNA expression (FPKM) accounted for by the union of top five expressed lncRNAs per cancer (total 11 lncRNAs).

# Figure 2

**Fig 2: lncRNAs exhibit tissue specific expression that can distinguish cancers.**
**a.** Tissue specificity index (tau score) which ranges from 0 (ubiquitously expressed) to 1 (tissue specific) is plotted for genes across three gene types: protein coding genes, lncRNAs, and novel lncRNAs. Table shows the tau score range and mean per gene type. **b.** Number of tissue specific known and novel lncRNAs in each cancer as defined by tissue specific gene threshold: tau score > 0.8. **c.** Heatmap showing the hierarchically clustered gene expression for the top five most tissue specific lncRNAs per cancer, ranked by highest tau score. Samples from each cancer cluster together based on expression of these genes alone.

# Figure 3

**Fig 3: A similar proportion of lncRNAs and protein coding genes are dysregulated due to SCNA**

**a.** The proportion of protein coding and lncRNA genes that have significant differential expression due SCNA, separated by copy number type (amplification or deletion). The number of genes found in SCNA loci is shown per cancer. Genes were evaluated to have differential expression due to copy number using the Wilcoxon rank sum test (p-value < 0.05) and log |fold change| > 1.5), comparing samples with no SCNA to samples with low/high SCNA as defined by GISTIC scores. **b.** The number of differentially expressed lncRNAs per chromosome and per cancer, distinguished by color. Chromosome 1 and 17 had the most dysregulated lncRNAs associating with the greater frequency of SCNA on these chromosomes across cancers. **c.** Number of samples with structural variant breakpoints in or near (+/- 2.5kb) lncRNAs and that are also located in copy number regions, stratified by amplification or deletion status of the locus.

# Figure 4

**Fig 4: lncRNA modulators impact transcriptional networks involving proliferation.**
**a.** Schematic that shows the three ways (attenuate, enhance, or invert) in which differentially expressed lncRNA modulators can impact transcription factor and target gene relationships. lncRNA modulators are associated with a TF-target gene pair based on a significant difference between TF-target gene expression correlation in samples with low lncRNA expression (lowest quartile) vs samples with high lncRNA expression (highest quartile). **b.** The proportion of lncRNA modulator types associated with significantly dysregulated lncRNA modulator- TF-target gene (lncMod) triplets. The number of significantly dysregulated lncMod triplets is listed per cancer. **c.** Number of lncRNA modulators genes that are common in lncMod triplets across cancers. Common lncRNA modulator genes tend to have a lower tau score compared to lncRNA modulators only associated with one cancer. **d.** Gene set enrichment using the MSigDB Hallmark gene set, of target genes associated with lncRNA modulators in each cancer (Fisher's exact test, FDR < 0.1). **e.** Transcription factors associated with the B-ALL expression specific lncRNA, *BLACE*, ranked based on number of regulated target genes. **f.** Expression heatmap of *BLACE* and the target genes of the XBP1 transcription factor, grouped by associated hallmark gene set, in samples within the bottom and top quartiles of *BLACE* expression in B-ALL.

# Figure 5

**Fig 5: Identification of lncRNAs associated with distinct neuroblastoma cell states**
**a.** The MES and ADRN signature score for TARGET NBL samples, with each sample labeled with either ADRN, Mixed, or MES phenotype based on clustering analysis. **b.** Heatmap of the expression of lncRNAs that have significant correlation with either the MES or ADRN score (|r| >0.6, pval < 0.01). lncRNAs were correlated with protein coding genes on the same chromosome and subsequent gene set enrichment analysis was performed for MES and ADRN protein coding genes separately. **c.** Schematic of how ADRN associated CRC regulated genes are identified using ChIP-seq and chromatin interaction data. We identified lncRNAs based on three types of regulation. 1) CRC transcription factors binding directly at the promoter of the lncRNA. 2) CRC TFs bind an enhancer region that interacts with a lncRNA promoter. 3) CRC TFs bind the promoter of a different gene and this promoter interacts with a lncRNA promoter. CRC TF binding was identified from ChIP-seq data, while enhancer-promoter and promoter-promoter interactions were identified from chromatin capture data. **d.** Filtering of lncRNAs expressed in NBL based on CRC TF regulation and differential expression based on sample phenotypes (ADRN or MES). **e.** Expression of *TBX2* and *TBX2-AS1* stratified by NBL sample phenotype (ADRN or MES). **f.** ChIP-seq tracks for histone marks and CRC transcription factors in the NBL cell line: BE(2)C, and promoter capture C chromatin interactions in NBL cell line: NB1643, at the *TBX2/TBX2-AS1* locus.

# Figure 6

**Fig 6: The *TBX2-AS1* lncRNA plays a role in neuroblastoma proliferation by modulating TBX2**

**a.** Expression of *TBX2* and *TBX2-AS1* in NBL tumor samples with and without 17q gain. **b.** The top MSigDB Hallmarks enriched across targets genes (p-value < 0.01) regulated by *TBX2-AS1* as predicted from lncMod analysis. **c.** The transcription factors with most target genes regulated by *TBX2-AS1* as predicted from lncMod analysis. **d.** Expression of gene targets of the E2F1 transcription factor that are enriched for proliferation hallmarks, in samples with low and high *TBX2* and *TBX2-AS1* expression. *TBX2* expression is highly correlated with that of *TBX2-AS1* (Pearson's r=0.77). **e.** siRNA knockdown efficiency of *TBX2-AS1* in the NBL cell line: NLF is 91% and in the SKNSH cell 63% knockdown was achieved. **f.** Western blot analysis of TBX2 in siTBX2 and siTBX2-AS1 treated NLF and SKNSH cell lines. **g.** Representative image of cell growth (as measured by RT-Ces assay) of the NBL cell lines: NLF and **h**. SKNSH. Cell index is normalized to time point when siRNA reagent is added at 24 hours post cell plating. **i.** Results from iRegulon analysis for genes that are up- or down-regulated upon siTBX2-AS1 treatment in NLF. Number of genes shown in Venn diagram with evidence of motif or ChIP-seq binding of the listed transcription factors.