Accuracy responses in species identification varying DNA barcode lengths with a Naïve Bayes classifier: Efficacy of mini-barcode under a supervised machine learning approach

Mohimenul Karim[1◑], Md. Rashid Abid[1*◑]

1 Department of Computer Science and Engineering, Bangladesh University of Engineering and Technology, Dhaka, Bangladesh

◑These authors contributed equally to this work.
* 1018052031@grad.cse.buet.ac.bd

Abstract

Specific gene regions in DNA, like COI (Cytochrome c oxidase I) in case of animals, have been defined as DNA barcode and many studies proved that it can be used as an identifier to distinguish species. The standard length of a DNA barcode is approximately 650 bp. But because of the challenges in sequencing technologies or unavailability of high-quality genomic DNA, it is not always possible to get the full barcode sequence of an organism. As a result, recent studies suggest that mini-barcodes can provide a good contribution in the species identification process. Among various methods proposed for the identification task, supervised machine learning methods have been shown effective. In this study, we have analyzed the effect of different barcode lengths on species identification from the perspective of supervised machine learning and suggested a general approximation of required length of mini-barcode in this regard. We have implemented a Naïve Bayes classifier as our model and implied the effectiveness of mini-barcode by demonstrating the accuracy responses varying the length of DNA barcode sequences.

Introduction

Species identification is naturally an elementary part in biological researches. According to the Catalogue of Life (CoL), which is a hierarchically ranked system based on expert and coherent opinions provided by taxonomists, more than 1.6 million species have been identified throughout the world so far [1]. Still, the total number of species is estimated to be 8.7 millions and it is suggested that in spite of over 250 years of taxonomic classification, the existing records tend to cover only a small portion of species on earth (14%) and in the ocean (9%) while the rest are yet to be described [2]. From the taxonomic point of view, the process of species discovery can be said to be broadly comprised of three pivotal but onerous steps: specimen collection, species level sorting and lastly species identification/description [3]. In addition to being already exigent by virtue of their own nature of work and thereby creating taxonomic impediments, these steps bring morphological characteristics into account in the traditional 'evidence-based' taxonomic approach and thus inflict likewise challenges in the decision making process [4]. For instance, in the specimen collection phase due to adverse situations often organism-fragments, stomach contents, feces, saliva, skin-scraps, blood, pollen etc can be achievable as samples which lack substantial morphological characteristics.

Additionally, in case of archival museum specimens, distinguishing morphological details are often absent. Also in the next phase of species level sorting, it is observed that many congeneric species possess morphological differences so negligible that they are often misclassified as a single species. Furthermore, in cases of species-complex and cryptic species, morphological differences are almost indiscernible. It might also be the case that due to sexual dimorphism, two sexes of same species can exhibit significantly disparate morphological attributes misleadingly appearing to be of different species. Various organisms can also develop conspicuous morphological changes in different stages of their life-cycle.

For addressing all the aforementioned issues, a more advanced, accurate, fast and efficient alternative approach in taxonomy was long yearned for which would attenuate morphological subservience in such cases. It is deemed that microgenomic approach involving DNA sequences has been successfully established as an extremely promising technique in recent years for species identification in this regard [5]. In this technique, a short portion of DNA from a specific gene or genes of an organism is used to uniquely identify a species in comparison with a reference library [6]. That specific portion is referred to as *DNA barcode* of that species as the process is similar to the process of identifying retail products with the help of machine readable barcodes. Different gene regions are used as DNA barcodes for different groups, namely: cytochrome-c-oxidase-I (*COI/COX1*) gene found in mtDNA for animals [7], *rbcL* and *matK* for plants [8] and the Internal Transcribed Spacer (*ITS*) rRNA for fungi [9]. Species identification with DNA Barcode can be treated as a classification problem where an unknown DNA barcode sequence is matched against a reference library comprising of DNA barcode sequences of already known species [10]. Various methods have been proposed for addressing this problem [11] and supervised machine learning approach is a prominent one among them where the reference dataset of known species is used as *Train Set* for training corresponding machine learning models and barcode sequences requiring classifications are placed in *Test Set* for query [12].

Notwithstanding the prominence of DNA barcode in species identification, several shortcomings are however perceived in some specific cases. Although PCR amplification and sequencing is usually consistent in freshly collected and well-preserved specimens, obtaining a full-length (∼650 bp) barcode in case of archival museum specimens which are preserved under sub-optimal or DNA-unfriendly conditions for years is often difficult [13]. This is also the case for processed biological materials (e.g. food products, pharmaceuticals etc.) and decayed tissues due to their degraded quality of DNA [14]. In such instances where obtaining a full-length barcode is hard, shorter barcode sequences (∼100−300 bp), often referred to as *Mini-barcodes*, have been found efficacious for species-level identification owing to their comparatively better amplification gain [15]. For this reason, mini-barcode is also being popularly used in DNA metabarcoding strategies based on environmental DNA (eDNA) [16] and researching ecological biodiversity [17]. Mini-barcodes can also be advantageously applied in alternative next generation short-read sequencing platforms providing higher throughput but remaining more cost-effective at the same time [13].

Since DNA Barcode was exercised as an effective molecular approach in species diagnosis by Hebert *et al.* [6], it has been used as a robust viewpoint in a wide range of studies [18–20]. *iBOL* (International Barcode of Life) and *CBOL* (The Consortium for the Barcode of Life) has also patronized the development initiatives for establishing DNA Barcode as a global standard for species identification [21]. Methodologies perceived in the literature review of species identification approaches with DNA barcoding can be divided into few major categories [11] [22] [23]. In *Distance Based* approaches, the reference library sequences are ranked with respect to their distances with the query barcode sequence on basis of the impression that small distances suggest

two barcodes might be of specimens of same species while large distances evince their belonging to two different species. This distance, commonly referred to as *Barcode Gap* [24], might be calculated on the basis of pairwise p-distances [6] or K2P distance [22]. *TaxI* [25] uses pairwise distances as distance measures whereas *BOLD* [26] uses K2P distances by default. In *Similarity Based* approaches, the query sequences are usually assigned to species based on a *Similarity score* [27] indicating how much barcode characters they have in common. *BLAST* [28], *FASTA* [29], TaxonDNA [30], *NN* [22] are some of the prominent examples in this regard. *Phylogenetic Approaches* construct trees using hierarchical clustering [27] where distinct clusters represent distinct species [31]. It can adopt NJ [32], Parsimony (*PAR* [33]), Maximum Likelihood (*PhyML* [34]) or Bayesian Inference (*MRBAYES* [35]) as underlying methods. *Statistical Approaches* use population genetics assumptions [36] based on coalescent theory (*GMYC* [37]) which takes phylogenetic uncertainty (*SAP-NJ* [38]), likelihood ratio tests [39] etc. into account. *Character Based* approaches investigate the presence/absence of particular characters, known as *Diagnostic Sites*, instead of total sequence [23] imitating morphological approaches. *CAOS* [40], *BLOG* [10], *DNABar* [41], *BRONX* [27], *DOME-ID* [27] are some instances of this approach. There are also some *Alignment-Free tools* such as: *ATIM-TNT* (Tree-based) [27], *CVTreeAlpha1.0* (Component Vector based) [42], *Spectrum Kernel* [43], *Alfie* (python based) [44] as well as *Web Based Tools* such as: *Linker* [45], *iBarcode* [46], *BioBarcode* [47], *ConFind* [48] for DNA barcoding. Comparative analysis of these major classical barcoding approaches are also extensively studied [11] [27] but there is no general consensus about a single best method to analyze DNA barcode data in species identification [22] [23].

Machine learning methods have been introduced as state-of-the-art methods for their extensive applicability in species identification with DNA Barcoding [12]. Although supervised statistical classification methods (CART, RF and Kernel methods) were studied previously [22]; supervised machine learning algorithms e.g. SVM, Jrip, J48 (C4.5), Naïve Bayes were compared on WEKA platforms against those ad-hoc methods by Weitschek *et al.* and it was observed that SVM and Naïve Bayes outperform on average in both synthetic and empirical datasets [12]. Simple-logistic, IBK, PART, Attribute-selected Classifier, Bagging approaches were also implemented in another study [49] in this regard. SMO, BP-NN [50], RF [51], k-mer based approaches [52] [53] can also be perceived in recent studies. Naïve Bayes is also applied on COI [54] barcode database demonstrating misclassification rates and also on ribosomal databases [55] effectively.

With the exploration of the prospects of DNA minibarcode in species identification [56], the application of DNA barcode has also significantly broadened. It is observed that although full length barcodes perform best with 97% species resolution, 90% and 95% identification success can be obtained with 100bp and 250bp mini barcodes respectively [15]. In case of species diagnosis from damaged/degraded DNA samples, ∼100−300bp mini-barcodes have also shown promising results [14]. Tree-based methods, objective clustering, automatic barcode gap discovery (ABGD), Poisson Tree Process (PTP) were also examined for analyzing the utility of minibarcodes for species identification [3]. Although no simple straight-forward formula is available through which necessary sequence lengths for ensuring species identification can be easily predicted [7], the general assumption of achieving superior performance by using full barcode length can be questioned as a saturation point is often seen beyond which the additional length-data create only a little impact in comparison to the increased cost overheads [3]. In this work of ours, we have tried to shed some lights on this question from the machine learning perspective by demonstrating the accuracy responses of species identification at different partial barcode lengths after implementing a Naïve

Bayes classifier as our analyzing contrivance. 122

Materials and methods 123

Datasets 124

Empirical Data 125

Weitschek *et al.* stated that for species identification, a full reference set with all 126
possible nucleotide polymorphisms for the sequences of each species is required as well 127
as a sufficient reference set is important to avoid over-fitting or under-fitting [12]. For 128
this purpose, we included the datasets they used in their work of [12] in our analysis 129
which is basically a collection of published empirical datasets and simulated DNA 130
barcode datasets. The public empirical datasets of *Bats*, *Birds*, *Cypraeidae*, *Drosophila* 131
and *Fishes* were chosen from GenBank Nucleotide Database for high phylogenetic 132
diversity and were available for download at `dmb.iasi.cnr.it/supbarcodes.php` in 133
FASTA format [12]. The datasets referred here as *Bird2*, *Fish2* and *Butterfly2* were 134
extracted from BOLD and available in CSV format at 135
`http://dimacs.rutgers.edu/archive/Workshops/BarcodeDataSets/`. Another 136
dataset was considered while performing our analysis, which appears to be the first 137
large-scale genetic assessment of the butterflies of Southern South America (available 138
for download at `dx.doi.org/10.5883/DS-BUNEACAR`) highlighting several cases of both 139
deep intraspecific and shallow interspecific divergence [57]. COI gene is used as gene 140
region in DNA barcode sequences of all these datasets. The detailed overview of these 141
empirical datasets having COI gene as barcode sequence can be found in Table 1. 142

Table 1. Summary of The Empirical Datasets: COI Gene as Barcodes

Dataset	No of sequences	Sequence Length	No of Species	Gene Region	Ref
Cypraeidae	2,008	614	211	COI	[58]
Drosophila	615	663	19	COI	[59]
Bats	826	659	82	COI	[26]
Fishes	626	702	82	COI	[60]
Birds	1,700	691	150	COI	[61]
Bird 2	2589	990	656	COI	[62]
Fish 2	754	901	211	COI	[63]
Butterfly	2028	658	417	COI	[57]
Butterfly 2	4267	901	561	COI	[56]

With a view to inspecting the overall scenario of species identification using DNA 143
Barcoding and analyzing the accuracy responses with varying barcode lengths, we 144
incorporated datasets of plants and fungi as well. These empirical datasets use rbcL and 145
ITS gene regions respectively and are available for download at 146
`dmb.iasi.cnr.it/supbarcodes.php` in FASTA format [12]. The detailed overview of 147
these empirical datasets of plants and fungi can be found in Table 2. 148

Table 2. Summary of The Empirical Datasets of Fungi, Inga and Algae

Dataset	No of sequences	Sequence Length	No of Species	Gene Region	Ref
Fungi	50	930	8	ITS	[9]
Inga	913	1,838	56	ITS	[64]
Algae	26	1,128	5	rbcL	[8]

Simulated Data

We have also used simulated real DNA Barcode datasets in our work for further
analysis. These datasets were also available for downlaod at
`dmb.iasi.cnr.it/supbarcodes.php` in FASTA format [12]. As stated in [11], these
data were simulated considering time of species divergence and the effective population
size (N_e). $N_e = 1000, 10000, 50000$ were used respectively in simulating gene trees and
the dataset complexity was increased with population size. After that, DNA barcode
sequences were simulated on the additive gene trees with 650 base sequence-length
bearing similarity to the real-life size of a standard DNA barcode. Each dataset consists
of 50 species and 20 individuals per species. The detailed overview of simulated datasets
is given in Table 3.

Table 3. Summary of The Simulated Datasets

Dataset	N_e	No of Individual	Sequence Length	No of Species	Ref
Ne1000	1000	20	650	50	[11]
Ne10000	10000	20	650	50	[11]
Ne50000	50000	20	650	50	[11]

Methodology

Naïve Bayes

Naïve Bayes [65] is one of those methods which are often used when a large reference set
is available. Naive Bayes is based on Bayes' Theorem and it is assumed that the
predictors are independent of each other. If we have an instance S with n features which
is represented by a feature vector $x = \{x_1, x_2, \ldots, x_n\}$, this model assigns to S a
probability $P(C_m|x_1, x_2, \ldots, x_n)$. Here, C_m is the m^{th} class among all possible classes.
Using Bayes' theorem, we can calculate the posterior probability $P(C_m|x)$ from
likelihood $P(x|C_m)$, prior probability $P(C_m)$ and evidence $P(x)$.

$$P(C_m|x) = \frac{P(x|C_m)P(C_m)}{P(x)} \tag{1}$$

Under naïve conditional assumption,

$$P(x_i|x_{i+1}, \ldots, x_n, C_m) = P(x_i|C_m) \tag{2}$$

So the posterior probability can be calculated as follows.

$$\frac{P(C_m)\prod_{i=1}^{n} P(x_i|C_m)}{P(x)} \tag{3}$$

As P(x) is constant for given input, we can only consider the numerator.
Using the Naïve Bayes model, the classification rule can be implemented as follows,
where $\hat{y}$ is the class label.

$$\hat{y} = \underset{m}{\operatorname{argmax}} P(C_m) \prod_{i=1}^{n} P(x_i|C_m) \tag{4}$$

Species Identification Using Naïve Bayes

Because of the efficacy of Naïve Bayes Algorithm in supervised prediction problem [54],
we have used this model for species identification process. We can outline the
identification approach as follows:

Let, b_k is the k^{th} barcode sequence in the dataset where $k = 1, 2, \ldots, n_b$ and n_b= no
of barcode entries in the dataset. $x_{1k}, x_{2k}, \ldots, x_{nk}$ are the nucleotides in a barcode
sequence b_k of length n. If C_p is the p^{th} species among different classes of species, the
probability that the barcode sequence b_k belongs to the species C_p is:

$$P(C_p|b_k) = \frac{P(b_k|C_p)P(C_p)}{P(b_k)} \tag{5}$$

The prior probability in eq. 5 can be calculated as follows:

$$
\begin{aligned}
P(b_k|C_p) &= P(x_{1k}, x_{2k}, \ldots, x_{nk}|C_p) \\
&= P(x_{1k}|x_{2k}, \ldots, x_{nk}, C_p)P(x_{2k}|x_{3k}, \ldots, x_{nk}, C_p) \\
&\ldots P(x_{nk}|C_p) \\
&= P(x_{1k}|C_p)P(x_{2k}|C_p)\ldots P(x_{nk}|C_p)
\end{aligned}
\tag{6}
$$

In the above equation, x_{ik} refers to the nucleotide base at i^{th} position of k^{th} sequence in
the dataset and $P(x_{ik}|C_p)$ means the probability of occurring nucleotide base x_{ik} in
species C_p at position i where $i = 1, \ldots, n$. So, if N_{c_p} and $N_{x_{ik}}$ are the number of
samples of C_p and the number of occurrences of x_{ik} at position i respectively, then we
calculate this probability as:

$$P(x_{ik}|C_p) = \frac{N_{x_{ik}}}{N_{C_p}} \tag{7}$$

If the proportion is equal to 0, then the posterior probability will become 0. To avoid
this, we replaced zero with a very small fractional value α, where $0 < \alpha << 1$. This α
can be considered as the mutation rate of nucleotides in the COI region [54].
$\alpha = 9.7 \times 10^{-8}$ was proposed as the estimated value of this mutation rate in COI
genes [66] and hence is also used here. In case of ambiguous base (e.g., N) or missing
data (e.g., -) at any position, we have not considered those positions for identification
purpose. In some datasets, there is a significant number of missing data (i.e., -) in a
significant number of sequences because of sequence alignments. Ignoring these missing
data in such cases can affect the performance of our model in identification process. For
this reason, in these situations we replaced a missing value in a particular position with
the most frequently occurring nucleotide in that position within a species.

Data Preparation

The empirical and simulated DNA Barcode datasets discussed in "Datasets" segment
are used to perform our analysis. Most of these empirical datasets were already split
into 80% and 20% sequences per species in training and testing set respectively by
biologists maintaining necessary polymorphism and species divergences [11]. In cases
where only a single dataset file was provided, it was split into Train and Test set on the
basis of 80%-20% random split. But it was ensured that whenever possible, each dataset
consists 4 or more representing sequences per species in the training set as it was found
necessary for obtaining a reliable classification performance [12]. The simulated datasets
were also split into Train and Test set in 80%-20% ratio where for every species 16
sequences were placed into Train Set and 4 sequences were placed into Test set. In cases
where datasets were non-aligned, MAFFT [67] was used as multiple sequence alignemnt
tool for aligning the barcode sequences. Most of the datasets were in FASTA format and
a Python program was written in order to convert these FASTA files into corresponding
CSV files. In case of a large number of missing nucleotide in barcode sequences of a
species, the most frequently occurring certain nucleotide within that species was
considered as replacement for that missing value. The related Python scripts written by
us for converting necessary files to desired CSV format and for replacing missing values
are available at `https://github.com/MohimenulRafi/DNA_Barcode.git`.

Experimental Arrangement

For the purpose of species identification with DNA Barcode, we developed a Naïve
Bayes classifier from scratch in Java fitting our methodological necessities (available at
`https://github.com/MohimenulRafi/DNA_Barcode.git`). In the training phase, a
probability matrix for each known species is calculated. For accomplishing this task, a
Hashmap is used where the key is the species name and the value is the probability
matrix. The matrix basically stores the probability of each nucleotide bases at each
position in the Barcode sequence for a species. A worked-out example is demonstrated
at Table 4 where it is considered that there are four different species and each having
three different barcode sequences of length five. The resulting probability matrix of
species 1 is shown in Table 5.

Table 4. Sample Barcode Sequences of Four Species

Species 1	Species 2	Species 3	Species 4
ATTGC	TCCAG	GTATG	CGAAT
ATTCC	TCCGG	GCATG	CGGAT
ATTGC	TCCGC	GTATC	CGAAT

Table 5. Probability Matrix of Species 1

Bases	Position 1	Position 2	Position 3	Position 4	Position 5
A	3/3	0/3	0/3	0/3	0/3
T	0/3	3/3	3/3	0/3	0/3
G	0/3	0/3	0/3	2/3	0/3
C	0/3	0/3	0/3	1/3	3/3

In the test phase, for a query sequence, we calculate it's likelihood of being a
particular species. Our model calculates the maximum likelihood and classify the new
sequence to that corresponding species. For example, if we get a query sequence
TCCAC from test set, our model takes the probability of T, C, C, A, C in position 1, 2,
3, 4 and 5 respectively from probability matrix of species 1, 2 and so on. Finally if the
highest probability value considering the whole length of query sequence comes from the
matrix of species 2, the sequence will be classified as species 2. Fig 1 shows an overview
of this classification process.

To analyze the impact of length of DNA Barcode sequences on species identification,
we have run our model with different sequence lengths on all the empirical as well as
simulated datasets in a comprehensive manner.

Results and Discussions

After implementing our Naïve bayes classifier, we needed to test the effectiveness of our
method. For this reason, to ensure the efficacy of our model, we ran our Naïve bayes
classifier on first five of the COI gene based empirical datasets mentioned in Table 1
considering their full-length barcode sequences. After comparing the results with the
maximum accuracy mentioned in other related studies [12] [49], where comparative
performance analysis of different supervised machine learning approaches on those same
COI based empirical datasets are provided, we thereby find our Naïve Bayes classifier
competent with their performances (shown in Fig 2).

After testing our Naïve Bayes classifier on the COI based first five empirical
datasets, subsequently we ran our model on all of the COI based empirical datasets

Fig 1. Overview of Species Identification with Naïve Bayes Procedure.
Each test sequence is fetched from Test set and using the probability matrices the
posterior probabilities are calculated. Based on the maximum probability score, that
respective test sequence is classified to corresponding species label.

Fig 2. Performance Competency of our Naïve Bayes Classifier. The accuracy
comparison of our model and the maximum accuracy score giving models of the works
of [12] and [49] is depicted in this figure. Our model yields very competing performance
in each cases.

mentioned in Table 1 with a view to analyzing accuracy responses varying barcode   251
sequence lengths. As all of the datasets in Table 1 had barcode sequence lengths >600   252
bp commonly, we chose to set the upper bound of sequence length at 600 while varying   253
lengths. Then the sequence lengths were decreasingly varied from this upper bound of   254
600 bp with an interval of 50 bases and corresponding accuracy scores were calculated   255

in each case. As 50 bp falls too low as sequence length, we considered 70 base-length ₂₅₆ after 100 bp and tried to maintain this as the lower bound for varying sequence lengths ₂₅₇ throughout our study (with only a few exceptions). All these accuracy scores for ₂₅₈ varying DNA barcode lengths for COI based empirical datasets are represented in Table ₂₅₉ 6. These length-wise accuracy scores of Table 6 are also plotted in Fig 3 and ₂₆₀ demonstrated in the Heatmap of Fig 4. If we analyze the pattern of these length-wise ₂₆₁ changes of accuracy scores in these plottings and heatmap, it seems that in most cases ₂₆₂ accuracy scores remain almost persistent in the range of 200bp-600bp. Although there ₂₆₃ is a gradual but trifling fall of accuracy in this range with the decrease of barcode ₂₆₄ sequence length, it significantly falls below 200bp barcode length. ₂₆₅

Table 6. Accuracy Responses in COI based Empirical Dataset

Datasets	Lengths											
	70	100	150	200	250	300	350	400	450	500	550	600
Bats	93.06	97.22	99.31	99.31	99.31	99.31	99.31	99.31	99.31	99.31	99.31	99.31
Bird	90.54	97.48	98.42	98.42	98.42	97.48	97.16	97.48	97.16	97.16	97.16	97.16
Cypraeidae	89.77	91.19	93.75	94.60	94.03	94.60	95.17	95.17	95.17	95.45	95.45	95.45
Drosophila	96.55	98.28	98.28	98.28	98.28	99.14	99.14	99.14	99.14	99.14	99.14	99.14
Fish	88.29	93.69	99.1	100.0	100.0	100.0	100.0	100.0	100.0	100.0	100.0	100.0
Butterfly	89.8	93.47	95.92	95.92	95.92	95.92	95.92	95.92	95.92	95.92	96.33	96.33
Bird 2	79.03	90.32	95.97	96.37	97.18	97.18	97.58	97.58	98.79	98.79	98.79	98.79
Butterfly 2	35.19	86.36	95.96	97.14	97.14	99.16	99.49	99.66	99.66	99.66	99.66	99.66
Fish 2	45.92	84.69	89.8	91.84	91.84	91.84	96.94	96.94	97.96	97.96	98.98	98.98

Fig 3. Lengthwise Accuracy comparison for COI based Empirical Datasets. From the lengthwise accuracy variations of each dataset, it is apparent that the accuracy is relatively consistent in the 200bp-600bp range but considerably falls below 200bp.

This similar approach is applied for the simulated datasets mentioned in Table 3 ₂₆₆ with maintaing same upper bound of 600 bp and lower bound of 70bp. The length-wise ₂₆₇ accuracy responses for the simulated datasets are summarized in Table 7 and ₂₆₈ demonstrated in Fig 5. From this Fig 5 it is also evident that, the accuracy score drops ₂₆₉

Fig 4. Heatmap of Lengthwise Accuracy Comparison for COI based Empirical Datasets. The lengthwise accuracy variation of Fig 3 is depicted in this heatmap. It is observed from the lengthwise color variation at each row that, 200bp-600bp provides a relatively consistent accuracy.

slightly and gradually with the reduction of sequence length and falls significantly when   270
the sequence length is reduced to <200bp.   271

Table 7. Accuracy Responses in Simulated Datasets

Datasets	Lengths											
	70	**100**	**150**	**200**	**250**	**300**	**350**	**400**	**450**	**500**	**550**	**600**
Ne1000	75.91	82.63	87.52	90.34	91.88	93.21	94.06	94.84	95.26	95.66	96.02	96.32
Ne10000	81.05	86.85	91.28	93.21	94.43	95.14	95.54	95.84	96.08	96.26	96.46	96.60
Ne50000	81.62	85.80	88.65	90.14	91.09	91.56	91.99	92.26	92.55	92.73	92.92	93.09

Furthermore, in order to get a complete overview of the effect of length on the   272
species identification process, the datasets mentioned in Table 3, which are based on   273
ITS and rbcL genes, are analyzed. For the dataset *Fungi* and *Algae*, we set upper   274
bound and lower bound for varying sequence length as 850bp and 100bp respectively.   275
For the dataset *Inga*, we set 1800bp and 1050bp as respective upper and lower bounds   276
as the full lengths of barcode sequences in this dataset are >1800bp. In all three   277
datasets, we decrease the length with an interval of 50 bases. The complete length-wise   278
average accuracy analysis of these datasets is represented in Table 8. This analysis is   279
also depicted in Fig 6.   280

From there it can be seen that although the contexts are different from the previous   281
length wise accuracy patterns of COI based empirical datasets and simulated datasets,   282
still *Algae* and *Fungi* shows a sharp fall in accuracy scores around sequence lengths   283
below 200-250bp. But this corroboration cannot be extended in case of *Inga* as its   284
length-bounds had to be chosen differently; but it still shows a decline in accuracy score   285
like every other ones if the lengths are gradually decreased from full length. In order to   286
bring these three datasets on a common ground, average accuracy responses were   287
further calculated at various percentage levels of their full lengths. The results are   288
represented in Table 9 and depicted in Fig 7. From these representations, we can   289

Fig 5. Lengthwise Accuracy comparison for Simulated Datasets. This figure suggests a gradual but minor drop in the accuracy score when reducing the sequence length from 600 bp to 200 bp whereas a comparatively significant fall is noticed when the sequence length is reduced below 200 bp.

Table 8. Lengthwise Accuracy Responses in Fungi, Inga and Algae Dataset

		Lengths														
Fungi	**bp**	100	150	200	250	300	350	400	450	500	550	600	650	700	750	800	850
	%	80.56	88.89	88.89	88.89	88.89	88.89	88.89	88.89	88.89	88.89	88.89	88.89	88.89	88.89	88.89	88.89
Inga	**bp**	1050	1100	1150	1200	1250	1300	1350	1400	1450	1500	1550	1600	1650	1700	1750	1800
	%	78.79	78.79	78.79	78.91	85.09	86.63	87.53	87.40	87.40	87.40	87.14	89.71	90.48	90.10	95.11	95.11
Algae	**bp**	100	150	200	250	300	350	400	450	500	550	600	650	700	750	800	850
	%	90.0	90.0	90.0	90.0	100.0	100.0	100.0	100.0	100.0	100.0	100.0	100.0	100.0	100.0	100.0	100.0

Table Notes: In this table, accuracy responses are demonstrated at different lengths for each of the Fungi, Inga and Algae datasets where respective results for each dataset consist of two rows. For each dataset, the first row indicates the lengths of the sequence (denoted by bp) and the second row indicates the corresponding accuracies (denoted by %).

Fig 6. Lengthwise Accuracy comparison in Fungi, Inga and Algae Dataset. This figure demonstrates a sharp fall in accuracy scores for Algae and Fungi dataset around sequence lengths below 200bp-250bp. The length and subsequent length wise accuracy responses in Inga dataset are not similar to those of Fungi and Algae datasets, but still a decline in accuracy score can be observed with the gradual decrease of sequence length.

observe that although no persistent range like previous COI based empirical datasets is ²⁹⁰
evident from these ITS & rbCL based datasets, the length-wise accuracy of *Fungi* and ²⁹¹
Algae falls sharply at around 20%-30% of their full lengths whereas the accuracy in *Inga* ²⁹²
dataset considerably falls first at around 70% and later again at 40% of its full length. ²⁹³
It is worth mentioning in this regard that, in case of COI based datasets and simulated ²⁹⁴
datasets, the length (∼200bp) below which a sharp decline was observed happens to be ²⁹⁵
approx. 30% of the full length of barcode sequence. In both cases of Table 8 and Table ²⁹⁶
9, the first row denotes the length of the sequence and the second row denotes the ²⁹⁷
corresponding accuracy considering that length respectively for each entry. ²⁹⁸

Table 9. Accuracy responses in Fungi, Inga and Algae dataset at different length-percentage

		Percentages of Lengths								
		10%	**20%**	**30%**	**40%**	**50%**	**60%**	**70%**	**80%**	**90%**	**100%**
Fungi	**bp**	93	186	279	372	465	558	651	744	837	930
	%	79.44	93.33	93.33	93.33	93.33	93.33	93.33	93.33	93.33	93.33
Inga	**bp**	183	367	551	735	919	1102	1286	1470	1654	1838
	%	46.33	66.84	67.09	76.41	76.54	79.09	87.57	88.49	90.06	94.96
Algae	**bp**	112	225	338	451	564	676	789	902	1015	1128
	%	90.0	90.0	100.0	100.0	100.0	100.0	100.0	100.0	100.0	100.0

Table Notes: In this table, accuracy responses are demonstrated at different percentage levels of the full sequence lengths for each of the Fungi, Inga and Algae datasets where respective results for each dataset consist of two rows. For each dataset, the first row indicates the lengths of the sequence (denoted by bp) at the mentioned percentage levels and the second row indicates the corresponding accuracies (denoted by %).

In support of our findings we can allude to Fig 8, which is adapted and redrawn ²⁹⁹
from the work of [15]. In this figure, the proportion of species resolution with different ³⁰⁰
DNA Barcode size is depicted. The shapes in individual plottings of Fig 3 also comport ³⁰¹
with the shape in Fig 8 implying the consistency between these separate but similar ³⁰²
findings. In other related studies regarding mini-barcodes, it was also observed that full ³⁰³
length barcodes provide marginally better identification success rates than mini ³⁰⁴
barcodes having 200-400bp sequence length while the identification ability is ³⁰⁵
significantly compromised only when the barcode length falls too short below 200bp [3]. ³⁰⁶
In light of these above discussions, this proposition can be made that an approximation ³⁰⁷
of 200-350bp mini-barcode length can be suggested for using in the process of species ³⁰⁸
identification with DNA barcode approached with a supervised machine learning model ³⁰⁹
where it has a significant possibility to perform with a reasonable accuracy score. ³¹⁰

Conclusion ³¹¹

In this study, we tried to provide an approximation of required length of DNA ³¹²
Mini-barcode from the supervised learning perspective by analysing the length-wise ³¹³
accuracy in the species identification process. A Naïve Bayes classifier was implemented ³¹⁴
for the comprehensive analysis of the effect of varying DNA Barcode lengths on the ³¹⁵
identification accuracy. Our findings from this machine learning perspective bears ³¹⁶
consistency with the findings of other related biological works where DNA ³¹⁷

Fig 7. Accuracy responses in Fungi, Inga and Algae dataset at different length-percentage. As the sequence lengths of these three datasets are different, average accuracy responses were further calculated at various percentage levels of their full lengths to bring uniformity.

Fig 8. Change of Species Resolution in different DNA mini-barcode lengths (adapted and redrawn from [15]). In the demonstration of [15], it is asserted that although best performance can be achieved at full length barcode with 97% species resolution, 100 bp and 250 bp mini barcode provide 90% and 95% identification success respectively.

mini-barcodes were used and analyzed to facilitate the species identification process. 318
Although we could perceive the approximate effective length of mini barcodes in case of 319
COI gene based DNA sequences, further study is needed for plants and ITS gene 320
barcode based organisms. Improving the quality of the sequence alignment by using 321
better multiple sequence alignment tools/techniques might improve the identification 322
accuracy in this regard. Other machine learning algorithms are also encouraged to be 323
applied in this context to enrich the comprehensiveness of our finding. 324

Acknowledgments 325

We sincerely thank Md. Shamsuzzoha Bayzid for his valuable guidelines and continuous 326
supports throughout the overall process. 327

References

1. Ruggiero MA, Gordon DP, Orrell TM, Bailly N, Bourgoin T, Brusca RC, et al. A higher level classification of all living organisms. PloS one. 2015;10(4):e0119248.

2. Mora C, Tittensor DP, Adl S, Simpson AG, Worm B. How many species are there on Earth and in the ocean? PLoS Biol. 2011;9(8):e1001127.

3. Yeo D, Srivathsan A, Meier R. Longer is not always better: Optimizing barcode length for large-scale species discovery and identification. Systematic Biology. 2020;.

4. Dayrat B. Towards integrative taxonomy. Biological journal of the Linnean society. 2005;85(3):407–417.

5. Armstrong K, Ball S. DNA barcodes for biosecurity: invasive species identification. Philosophical Transactions of the Royal Society B: Biological Sciences. 2005;360(1462):1813–1823.

6. Hebert PD, Cywinska A, Ball SL, Dewaard JR. Biological identifications through DNA barcodes. Proceedings of the Royal Society of London Series B: Biological Sciences. 2003;270(1512):313–321.

7. Hebert PD, Ratnasingham S, De Waard JR. Barcoding animal life: cytochrome c oxidase subunit 1 divergences among closely related species. Proceedings of the Royal Society of London Series B: Biological Sciences. 2003;270(suppl_1):S96–S99.

8. Group CPW, Hollingsworth PM, Forrest LL, Spouge JL, Hajibabaei M, Ratnasingham S, et al. A DNA barcode for land plants. Proceedings of the National Academy of Sciences. 2009;106(31):12794–12797.

9. Schoch CL, Seifert KA, Huhndorf S, Robert V, Spouge JL, Levesque CA, et al. Nuclear ribosomal internal transcribed spacer (ITS) region as a universal DNA barcode marker for Fungi. Proceedings of the National Academy of Sciences. 2012;109(16):6241–6246.

10. Weitschek E, Van Velzen R, Felici G, Bertolazzi P. BLOG 2.0: a software system for character-based species classification with DNA Barcode sequences. What it does, how to use it. Molecular ecology resources. 2013;13(6):1043–1046.

11. van Velzen R, Weitschek E, Felici G, Bakker FT. DNA barcoding of recently diverged species: relative performance of matching methods. PloS one. 2012;7(1):e30490.

12. Weitschek E, Fiscon G, Felici G. Supervised DNA Barcodes species classification: analysis, comparisons and results. BioData mining. 2014;7(1):1–18.

13. Hajibabaei M, Smith MA, Janzen DH, Rodriguez JJ, Whitfield JB, Hebert PD. A minimalist barcode can identify a specimen whose DNA is degraded. Molecular Ecology Notes. 2006;6(4):959–964.

14. Hajibabaei M, McKenna C. DNA mini-barcodes. In: DNA barcodes. Springer; 2012. p. 339–353.

15. Meusnier I, Singer GA, Landry JF, Hickey DA, Hebert PD, Hajibabaei M. A universal DNA mini-barcode for biodiversity analysis. BMC genomics. 2008;9(1):1–4.

16. Staats M, Arulandhu AJ, Gravendeel B, Holst-Jensen A, Scholtens I, Peelen T, et al. Advances in DNA metabarcoding for food and wildlife forensic species identification. Analytical and Bioanalytical Chemistry. 2016;408(17):4615–4630.

17. Erickson DL, Reed E, Ramachandran P, Bourg NA, McShea WJ, Ottesen A. Reconstructing a herbivore's diet using a novel rbc L DNA mini-barcode for plants. AoB Plants. 2017;9(3):plx015.

18. Frézal L, Leblois R. Four years of DNA barcoding: current advances and prospects. Infection, Genetics and Evolution. 2008;8(5):727–736.

19. Goldstein PZ, DeSalle R. Review and interpretation of trends in DNA barcoding. Frontiers in Ecology and Evolution. 2019;7:302.

20. Adamowicz SJ, Boatwright JS, Chain F, Fisher BL, Hogg ID, Leese F, et al. Trends in DNA barcoding and metabarcoding. Genome. 2019;62(3):v–viii.

21. Schindel DE, Miller SE. DNA barcoding a useful tool for taxonomists. Nature. 2005;435(7038):17–18.

22. Austerlitz F, David O, Schaeffer B, Bleakley K, Olteanu M, Leblois R, et al. DNA barcode analysis: a comparison of phylogenetic and statistical classification methods. BMC bioinformatics. 2009;10(S14):S10.

23. Casiraghi M, Labra M, Ferri E, Galimberti A, De Mattia F. DNA barcoding: a six-question tour to improve users' awareness about the method. Briefings in bioinformatics. 2010;11(4):440–453.

24. Little DP. DNA barcode sequence identification incorporating taxonomic hierarchy and within taxon variability. PLoS one. 2011;6(8):e20552.

25. Steinke D, Vences M, Salzburger W, Meyer A. TaxI: a software tool for DNA barcoding using distance methods. Philosophical Transactions of the Royal Society B: Biological Sciences. 2005;360(1462):1975–1980.

26. Ratnasingham S, Hebert PD. BOLD: The Barcode of Life Data System (http://www. barcodinglife. org). Molecular ecology notes. 2007;7(3):355–364.

27. Little DP, Stevenson DW. A comparison of algorithms for the identification of specimens using DNA barcodes: examples from gymnosperms. Cladistics. 2007;23(1):1–21.

28. Altschul SF, Gish W, Miller W, Myers EW, Lipman DJ. Basic local alignment search tool. Journal of molecular biology. 1990;215(3):403–410.

29. Pearson W. Rapid and sensitive sequence comparison with FASTP and FASTA. Methods in enzymology. 1990;183:63–98.

30. Meier R, Shiyang K, Vaidya G, Ng PK. DNA barcoding and taxonomy in Diptera: a tale of high intraspecific variability and low identification success. Systematic biology. 2006;55(5):715–728.

31. Munch K, Boomsma W, Willerslev E, Nielsen R. Fast phylogenetic DNA barcoding. Philosophical Transactions of the Royal Society B: Biological Sciences. 2008;363(1512):3997–4002.

32. Saitou N, Nei M. The neighbor-joining method: a new method for reconstructing phylogenetic trees. Molecular biology and evolution. 1987;4(4):406–425.

33. Farris JS. Estimating phylogenetic trees from distance matrices. The American Naturalist. 1972;106(951):645–668.

34. Guindon S, Gascuel O. A simple, fast, and accurate algorithm to estimate large phylogenies by maximum likelihood. Systematic biology. 2003;52(5):696–704.

35. Huelsenbeck JP, Ronquist F. MRBAYES: Bayesian inference of phylogenetic trees. Bioinformatics. 2001;17(8):754–755.

36. Abdo Z, Golding GB. A step toward barcoding life: a model-based, decision-theoretic method to assign genes to preexisting species groups. Systematic biology. 2007;56(1):44–56.

37. Pons J, Barraclough TG, Gomez-Zurita J, Cardoso A, Duran DP, Hazell S, et al. Sequence-based species delimitation for the DNA taxonomy of undescribed insects. Systematic biology. 2006;55(4):595–609.

38. Munch K, Boomsma W, Huelsenbeck JP, Willerslev E, Nielsen R. Statistical assignment of DNA sequences using Bayesian phylogenetics. Systematic biology. 2008;57(5):750–757.

39. Matz MV, Nielsen R. A likelihood ratio test for species membership based on DNA sequence data. Philosophical Transactions of the Royal Society B: Biological Sciences. 2005;360(1462):1969–1974.

40. Sarkar IN, Planet PJ, Desalle R. CAOS software for use in character-based DNA barcoding. Molecular Ecology Resources. 2008;8(6):1256–1259.

41. DasGupta B, Konwar KM, Măndoiu I, Shvartsman AA. DNA-BAR: distinguisher selection for DNA barcoding. Bioinformatics. 2005;21(16):3424–3426.

42. Chu KH, Xu M, Li CP. Rapid DNA barcoding analysis of large datasets using the composition vector method. BMC bioinformatics. 2009;10(S14):S8.

43. Kuksa P, Pavlovic V. Efficient alignment-free DNA barcode analytics. BMC bioinformatics. 2009;10(S14):S9.

44. Nugent CM, Adamowicz SJ. Alignment-free classification of COI DNA barcode data with the Python package Alfie. Metabarcoding and Metagenomics. 2020;4:e55815.

45. Albu M, Nikbakht H, Hajibabaei M, Hickey DA. The DNA barcode linker. Molecular ecology resources. 2011;11(1):84–88.

46. Singer GA, Hajibabaei M. iBarcode. org: web-based molecular biodiversity analysis. In: BMC bioinformatics. vol. 10. Springer; 2009. p. S14.

47. Lim J, Kim SY, Kim S, Eo HS, Kim CB, Paek WK, et al. BioBarcode: a general DNA barcoding database and server platform for Asian biodiversity resources. In: BMC genomics. vol. 10. Springer; 2009. p. S8.

48. Smagala JA, Dawson ED, Mehlmann M, Townsend MB, Kuchta RD, Rowlen KL. ConFind: a robust tool for conserved sequence identification. Bioinformatics. 2005;21(24):4420–4422.

49. Kabir T, Shemonti AS, Rahman AH. Species Identification Using Partial DNA Sequence: A Machine Learning Approach. In: 2018 IEEE 18th International Conference on Bioinformatics and Bioengineering (BIBE). IEEE; 2018. p. 235–242.

50. He T, Jiao L, Wiedenhoeft AC, Yin Y. Machine learning approaches outperform distance-and tree-based methods for DNA barcoding of Pterocarpus wood. Planta. 2019;249(5):1617–1625.

51. Meher PK, Sahu TK, Gahoi S, Tomar R, Rao AR. funbarRF: DNA barcode-based fungal species prediction using multiclass Random Forest supervised learning model. BMC genetics. 2019;20(1):2.

52. Meher PK, Sahu TK, Rao A. Identification of species based on DNA barcode using k-mer feature vector and Random forest classifier. Gene. 2016;592(2):316–324.

53. Kusuma WA, Nurilmala M, et al. Identification of Tuna and Mackerel Based on DNA Barcodes using Support Vector Machine (SVM). Telkomnika. 2016;14(2):778.

54. Anderson MP, Dubnicka SR. A sequential naive Bayes classifier for DNA barcodes. Statistical applications in genetics and molecular biology. 2014;13(4):423–434.

55. Porter TM, Gibson JF, Shokralla S, Baird DJ, Golding GB, Hajibabaei M. Rapid and accurate taxonomic classification of insect (class Insecta) cytochrome c oxidase subunit 1 (COI) DNA barcode sequences using a naïve Bayesian classifier. Molecular Ecology Resources. 2014;14(5):929–942.

56. Hajibabaei M, Janzen DH, Burns JM, Hallwachs W, Hebert PD. DNA barcodes distinguish species of tropical Lepidoptera. Proceedings of the National Academy of Sciences. 2006;103(4):968–971.

57. Lavinia PD, Núñez Bustos EO, Kopuchian C, Lijtmaer DA, García NC, Hebert PD, et al. Barcoding the butterflies of southern South America: Species delimitation efficacy, cryptic diversity and geographic patterns of divergence. PloS one. 2017;12(10):e0186845.

58. Meyer CP, Paulay G. DNA barcoding: error rates based on comprehensive sampling. PLoS biol. 2005;3(12):e422.

59. Lou M, Golding GB. Assigning sequences to species in the absence of large interspecific differences. Molecular Phylogenetics and Evolution. 2010;56(1):187–194.

60. Bertolazzi P, Felici G, Weitschek E. Learning to classify species with barcodes. BMC bioinformatics. 2009;10(S14):S7.

61. Hebert PD, Stoeckle MY, Zemlak TS, Francis CM. Identification of birds through DNA barcodes. Plos biol. 2004;2(10):e312.

62. Kerr KC, Stoeckle MY, Dove CJ, Weigt LA, Francis CM, Hebert PD. Comprehensive DNA barcode coverage of North American birds. Molecular ecology notes. 2007;7(4):535–543.

63. Ward RD, Zemlak TS, Innes BH, Last PR, Hebert PD. DNA barcoding Australia's fish species. Philosophical Transactions of the Royal Society B: Biological Sciences. 2005;360(1462):1847–1857.

64. Dexter KG, Pennington TD, Cunningham CW. Using DNA to assess errors in tropical tree identifications: How often are ecologists wrong and when does it matter? Ecological Monographs. 2010;80(2):267–286.

65. Rish I, et al. An empirical study of the naive Bayes classifier. In: IJCAI 2001 workshop on empirical methods in artificial intelligence. vol. 3; 2001. p. 41–46.

66. Denver DR, Morris K, Lynch M, Vassilieva LL, Thomas WK. High direct estimate of the mutation rate in the mitochondrial genome of Caenorhabditis elegans. Science. 2000;289(5488):2342–2344.

67. Katoh K, Standley DM. MAFFT multiple sequence alignment software version 7: improvements in performance and usability. Molecular biology and evolution. 2013;30(4):772–780.