Efficacy and accuracy responses of DNA mini-barcodes in species identification under a supervised machine learning approach

Mohimenul Karim
Department of Computer Science and Engineering
Bangladesh University of Engineering and Technology
Dhaka, Bangladesh
1105097.mk@ugrad.cse.buet.ac.bd

Rashid Abid
Department of Computer Science and Engineering
Bangladesh University of Engineering and Technology
Dhaka, Bangladesh
1205016.mra@ugrad.cse.buet.ac.bd

Abstract—**Specific gene regions in DNA, such as cytochrome c oxidase I (COI) in animals, are defined as DNA barcodes and can be used as identifiers to distinguish species. The standard length of a DNA barcode is approximately 650 base pairs (bp). However, because of the challenges associated with sequencing technologies and the unavailability of high-quality genomic DNA, it is not always possible to obtain the full-length barcode sequence of an organism. Recent studies suggest that mini-barcodes, which are shorter (100-300 bp) barcode sequences, can contribute significantly to species identification. Among various methods proposed for the identification task, supervised machine learning methods are effective. However, any prior work indicating the efficacy of mini-barcodes in species identification under a machine learning approach is elusive to find. In this study, we analyzed the effect of different barcode lengths on species identification using supervised machine learning and proposed a general approximation of the required length of the mini-barcode. Since Naïve Bayes is seen to generally outperform other supervised methods in species identification in other studies, we implemented this classifier and showed the effectiveness of the mini-barcode by demonstrating the accuracy responses obtained after varying the length of the DNA barcode sequences.**

Index Terms—**DNA Barcoding, Species identification, Mini-barcode, Barcode length, Supervised learning, Naïve Bayes**

I. INTRODUCTION

Species identification is an elementary component in biological research. According to the Catalogue of Life, a hierarchically ranked system based on expert opinions provided by taxonomists, more than 1.6 million species have been identified to date [1]. However the total number of species is estimated to be 8.7 million [2]. Therefore, despite over 250 years of taxonomic classification, the existing records cover only a small portion of species on Earth (14%) and in the ocean (9%) with the remaining yet to be described [2]. From a taxonomic perspective, species discovery comprises three pivotal and onerous steps: specimen collection, species level sorting, species identification/description [3]. These steps consider morphological characteristics in the traditional 'evidence-based' taxonomic approach and cause challenges in the decision-making process [4]. Morphological characteristics can be significantly different for the same species in different stages of its life-cycle. In case of archival museum specimens, distinguishing morphological details are scarce. During the specimen collection phase organism fragments, stomach contents, feces, saliva, skin-scraps, blood, and pollens are collected, which often lack substantial morphological characteristics. Some congeneric, species-complex and cryptic species have only tiny morphological differences, and thus are often misclassified as a single species. In sexual dimorphism, different sexes of the same species can exhibit significantly disparate morphological attributes, leading to misidentification as different species.

To address all the aforementioned issues with classification using traditional morphological methods, a more advanced, accurate, fast and efficient alternative approach is required to attenuate morphological dependencies. A microgenomic approach involving DNA sequences has been established recently as a promising technique for species identification [5]. In this technique, a short portion of DNA from a specific gene of an organism is used to uniquely identify a species in comparison to a reference library [6]. This specific portion is called the DNA barcode of that species. Different gene regions are used as DNA barcodes for different groups, such as the cytochrome-c-oxidase-I (*COI/COX1*) gene found in the mtDNA of animals [7], *rbcL* and *matK* in plants [8], and the internal transcribed spacer (*ITS*) rRNA in fungi [9]. Species identification using DNA barcodes can be considered as a classification problem where an unknown DNA barcode sequence is matched against a reference library of already known species DNA barcode sequences [10]. Several methods have been proposed to assist with this classification matching [11] with the supervised machine learning approach being prominent. In this approach, the reference dataset of known species is used as the train set for training the corresponding machine learning models and the barcode sequences requiring classifications are placed in the test set for the query [12].

Notwithstanding the prominence of DNA barcode in species identification, several shortcomings arise. Although polymerase chain reaction amplification and sequencing are usually consistent in freshly collected and well-preserved specimens,

obtaining a full length (∼650 bp) barcode in archival museum specimens that are preserved under sub-optimal or DNA-unfriendly conditions is difficult [13]. This is also the case for processed biological materials and decayed tissues owing to the degraded quality of the DNA [14]. In cases where obtaining a full-length barcode is difficult, shorter barcode sequences (∼100−300 bp), often referred to as *mini-barcodes*, are useful for species-level identification owing to their comparatively better amplification gain [15]. Mini-barcodes are also commonly used in DNA metabarcoding strategies based on environmental DNA [16] and for studying ecological biodiversity [17]. Mini-barcodes can also be applied in alternative next generation short-read sequencing platforms providing higher throughput while remaining cost-effective [13].

II. Related Works

DNA Barcodes were proposed as an effective molecular approach in species diagnosis by Hebert *et al.* [6] and have since been used in a wide range of studies [18]–[20]. The International Barcode of Life and the Consortium for the Barcode of Life have supported development initiatives for establishing DNA barcodes as a global standard for species identification [21]. Species identification approaches using DNA barcoding can be partitioned into a few major categories: distance based approach, similarity based approach, phylogenetic approach, statistical approach, and character based approach [11], [22], [23].

In distance based approaches, the reference library sequences are ranked with respect to their distances with the query barcode sequence, as small distances suggest that two barcodes can be of specimens of the same species whereas large distances indicate that they belong to two different species. This distance, commonly referred to as barcode gap [24], can be calculated based on the pairwise p-distances [6] or K2P distance [22]. TaxI [25] uses pairwise distances as distance measures whereas BOLD [26] uses K2P distances by default. In similarity based approaches, the query sequences are assigned to species based on a similarity score [27] indicating similarities in barcode characters. BLAST [28], FASTA [29], TaxonDNA [30], NN [22] are some prominent examples in this regard. Phylogenetic approaches construct trees using hierarchical clustering [27] where distinct clusters represent distinct species [31]. Phylogenetic approaches can adopt NJ [32], parsimony (PAR [33]), maximum likelihood (PhyML [34]) or bayesian inference (MRBAYES [35]) as underlying methods. Statistical approaches use population genetics assumptions [36] based on coalescent theory (GMYC [37]) considering phylogenetic uncertainty (SAP-NJ [38]), likelihood ratio tests [39] etc. Character based approaches investigate the presence/absence of particular characters, known as diagnostic sites, instead of total sequence imitating morphological approaches [23]. CAOS [40], BLOG [10], DNABar [41], BRONX [27], DOME-ID [27] are some instances of this approach. There are alignment-free tools such as: ATIM-TNT (Tree-based) [27], CVTreeAlpha1.0 (Component Vector based) [42], Spectrum Kernel [43] and Alfie (python based)

[44]. Web Based Tools such as: Linker [45], iBarcode [46], BioBarcode [47] and ConFind [48] are also there for DNA barcoding. Extensive comparative analyses of these major classical barcoding approaches have been studied [11], [27]; however, there is no general consensus on a single optimal method to analyze DNA barcode data for species identification [22], [23].

Machine learning methods have been introduced as state-of-the-art techniques for their extensive applicability in species identification using DNA barcoding [12]. Although supervised statistical classification methods such as: Classification And Regression Tree, Random Forest, and Kernel methods have been studied previously [22], supervised machine learning algorithms e.g., SVM, Jrip, J48 (C4.5), and Naïve Bayes were compared on WEKA platforms to the ad-hoc methods by Weitschek *et al.* and SVM and Naïve Bayes outperformed the other methods for synthetic and empirical datasets [12]. Simple-logistic, IBK, PART, attribute-selected classifier, and bagging approaches were also implemented in [49]. SMO, BP-NN [50], RF [51], and k-mer based approaches [52], [53] can be seen in recent studies. Naïve Bayes was applied on the COI [54] barcode database to show misclassification rates and on ribosomal databases [55].

With the exploration of the prospects of DNA minibarcode in species identification [56], the application of DNA barcode significantly broadened. It is observed that although full length barcodes perform efficiently with 97 % species resolution, 90 % and 95 % identification success can be obtained with 100 bp and 250 bp mini-barcodes, respectively [15]. In case of species diagnosis from damaged/degraded DNA samples, ∼100−300 bp mini-barcodes have shown promising results [14]. Tree-based methods, objective clustering, automatic barcode gap discovery, and Poisson tree process were examined to analyze the use of mini-barcodes for species identification [3]. No simple formula exists to determine the necessary sequence lengths for species identification [7] but the expectation of achieving significantly better performance using full-length barcodes is questionable because a saturation point can be obtained beyond which the additional length data yield insignificant impact compared to the increased overhead costs [3]. In the present study, we examined this issue from the machine learning perspective by demonstrating the accuracy responses of species identification at different partial barcode lengths after using a Naïve Bayes classifier.

III. Datasets

A. Empirical data

For species identification, a full reference set with all possible nucleotide polymorphisms for the sequences of each species is required and an adequate reference set is important to prevent over-fitting and under-fitting [12]. Therefore, we included the datasets used in [12] in our analysis, which are a collection of published empirical datasets and simulated DNA barcode datasets. The public empirical datasets of *Bats*, *Birds*, *Cypraeidae*, *Drosophila*, and *Fishes* were selected from the GenBank Nucleotide Database for high phylogenetic diversity

and are available for download at [1] in FASTA format [12]. The datasets referred to here as *Bird2*, *Fish2*, and *Butterfly2* were extracted from BOLD and are available in CSV format at [2]. An additional dataset considered in our analysis is the first large-scale genetic assessment of the butterflies of Southern South America (available for download at [3]) showing several cases of both deep intraspecific and shallow interspecific divergence [57]. The *COI* gene was used as the gene region in the DNA barcode sequences for all these datasets. A detailed overview of these empirical datasets is presented in Table I.

TABLE I
SUMMARY OF THE EMPIRICAL DATASETS: COI GENE AS BARCODES.

Dataset	No of sequences	Sequence Length	No of Species	Gene Region	Ref
Cypraeidae	2,008	614	211	COI	[58]
Drosophila	615	663	19	COI	[59]
Bats	826	659	82	COI	[26]
Fishes	626	702	82	COI	[60]
Birds	1,700	691	150	COI	[61]
Bird2	2589	990	656	COI	[62]
Fish2	754	901	211	COI	[63]
Butterfly	2028	658	417	COI	[57]
Butterfly2	4267	901	561	COI	[56]

To investigate the overall scenario of species identification using DNA barcoding and analyze the accuracy responses with varying barcode lengths, we used datasets of plants and fungi as well. These empirical datasets use the *rbcL* and *ITS* gene regions, respectively and are available for download at [1] in FASTA format [12]. A detailed overview of these empirical datasets is presented in Table II.

TABLE II
SUMMARY OF THE EMPIRICAL DATASETS OF FUNGI, INGA, AND ALGAE.

Dataset	No of sequences	Sequence Length	No of Species	Gene Region	Ref
Fungi	50	930	8	ITS	[9]
Inga	913	1,838	56	ITS	[64]
Algae	26	1,128	5	rbcL	[8]

B. Simulated data

We used simulated DNA barcode datasets for additional analysis. These datasets are available for downlaod at [1] in FASTA format [12]. As stated in [11], these real DNA barcode datasets were simulated considering the time of species divergence and the effective population size (N_e). To simulate gene trees $N_e = 1000, 10000$, and 50000 was used and the dataset complexity was increased with population size. Thereafter, DNA barcode sequences were simulated on the additive gene trees with 650 base sequence-length with similarity to the real-life size of a standard DNA barcode. Each dataset consisted of

50 species and 20 individuals per species. A detailed overview of the simulated datasets is presented in Table III.

TABLE III
SUMMARY OF THE SIMULATED DATASETS.

Dataset	N_e	No of Individual	Sequence Length	No of Species	Ref
Ne1000	1000	20	650	50	[11]
Ne10000	10000	20	650	50	[11]
Ne50000	50000	20	650	50	[11]

IV. METHODOLOGY

A. Naïve Bayes

Naïve Bayes [65] is a method that is frequently used if a large reference set is available. Naïve Bayes is based on Bayes' theorem, and it is assumed that the predictors are independent of each other. If an instance S with n features is represented by a feature vector $x = \{x_1, x_2, \ldots, x_n\}$, the model assigns a probability $P(C_m|x_1, x_2, \ldots, x_n)$ to S. Here, C_m is the m^{th} class among all possible classes.

Using Bayes' theorem, we calculated the posterior probability $P(C_m|x)$ from the likelihood $P(x|C_m)$, prior probability $P(C_m)$, and evidence $P(x)$:

$$P(C_m|x) = \frac{P(x|C_m)P(C_m)}{P(x)} \tag{1}$$

Under naïve conditional assumption,

$$P(x_i|x_{i+1}, \ldots, x_n, C_m) = P(x_i|C_m) \tag{2}$$

Therefore, the posterior probability $P(C_m|x)$ was calculated as follows:

$$P(C_m|x) = \frac{P(C_m) \prod_{i=1}^{n} P(x_i|C_m)}{P(x)} \tag{3}$$

Because $P(x)$ is constant for a given input, we can only consider the numerator.

Using the Naïve Bayes model, the classification rule can be implemented as follows, where $\hat{y}$ is the class label.

$$\hat{y} = \underset{m}{\operatorname{argmax}} P(C_m) \prod_{i=1}^{n} P(x_i|C_m) \tag{4}$$

B. Species identification using Naïve Bayes

Because of the efficacy of Naïve Bayes algorithm in supervised prediction problems [54] and its better performance compared to other supervised machine learning methods in species identification [12], we have used this model for species identification process. The identification approach used was as follows:

Let b_k be the k^{th} barcode sequence of length n in the dataset where $k = 1, 2, \ldots, n_b$ and n_b= no of barcode entries in the dataset. $x_{1k}, x_{2k}, \ldots, x_{nk}$ are the nucleotides in the b_k barcode sequence. If C_p is the p^{th} species among the different classes of species, the probability that the barcode sequence b_k belongs to the species C_p is

$$P(C_p|b_k) = \frac{P(b_k|C_p)P(C_p)}{P(b_k)} \tag{5}$$

The prior probability in eq. 5 can be calculated as follows:

$$
\begin{aligned}
P(b_k|C_p) &= P(x_{1k}, x_{2k}, \ldots, x_{nk}|C_p) \\
&= P(x_{1k}|x_{2k}, \ldots, x_{nk}, C_p)P(x_{2k}|x_{3k}, \ldots, x_{nk}, C_p) \\
&\quad \ldots P(x_{nk}|C_p) \\
&= P(x_{1k}|C_p)P(x_{2k}|C_p) \ldots P(x_{nk}|C_p)
\end{aligned}
\tag{6}
$$

In the above equation, x_{ik} refers to the nucleotide base at the i^{th} position of k^{th} sequence in the dataset and $P(x_{ik}|C_p)$ means the probability of occurring nucleotide base x_{ik} in species C_p at position i where $i = 1, \ldots, n$. Therefore, if N_{c_p} and $N_{x_{ik}}$ are the number of samples of C_p and the number of occurrences of x_{ik} at position i, respectively, then we calculate this probability as

$$
P(x_{ik}|C_p) = \frac{N_{x_{ik}}}{N_{C_p}}
\tag{7}
$$

If the proportion is equal to 0, then the posterior probability becomes 0. To prevent this, we replaced zero with a significantly small fractional value α, where $0 < \alpha << 1$. This α can be considered as the mutation rate of nucleotides in the particular gene region [54]. $\alpha = 9.7 \times 10^{-8}$ was proposed as the estimated value of this mutation rate in COI genes [66] and hence it is also used here. For ambiguous bases (e.g., *N*) or missing data (e.g., **-**) at any position, we have not considered those positions for identification purposes. In some datasets, there is a significant number of missing data (i.e., **-**) in a significant number of sequences because of sequence alignment. Ignoring these missing data in such cases could affect the performance of our model during the identification process. Therefore, in these situations, we replaced a missing value in a particular position with the most frequently occurring nucleotide in that position within a species.

C. Data preparation

The empirical and simulated DNA barcode datasets discussed in the section III were used to perform our analysis. Most of these empirical datasets were partitioned by biologists into 80 % and 20 % of total sequences per species in training and testing sets, respectively, maintaining the necessary polymorphism and species divergences [11]. In cases where only a single dataset file was provided, it was partitioned into train and test sets based on an 80 %-20 % random split. However, each dataset consisted of four or more representing sequences per species in the train set where possible, since it was necessary for obtaining a reliable classification performance [12]. The simulated datasets were also split into train and test sets in a 80 %-20 % ratio where for every species 16 sequences were placed into the train set and 4 sequences were placed into the test set. In cases where the datasets were non-aligned, MAFFT [67] was used as a multiple sequence alignemnt tool for aligning the barcode sequences. Most of the datasets were in FASTA format and a Python program was written to convert these FASTA files into corresponding CSV files. For large numbers of missing nucleotides in the barcode sequences of a

species, the most frequently occurring nucleotide within that species was considered as the replacement for that missing value. The related Python scripts written for converting the files to the desired CSV format and for replacing missing values are available at [4].

D. Experimental arrangement

For species identification with DNA barcodes, we developed a Naïve Bayes classifier using Java (available at [4]). During the training phase, a probability matrix for each known species is calculated. To achieve the goal, a Hashmap was used where the key was the species name and the value was the probability matrix. The matrix stored the probability of each nucleotide base at each position in the barcode sequence for a species. A worked-out example is presented in Table IV where there are four different species and each has three different barcode sequences of length five. The resulting probability matrix of species 1 is presented in Table V.

TABLE IV
SAMPLE BARCODE SEQUENCES OF FOUR SPECIES.

Species 1	Species 2	Species 3	Species 4
ATTGC	TCCAG	GTATG	CGAAT
ATTCC	TCCGG	GCATG	CGGAT
ATTGC	TCCGC	GTATC	CGAAT

TABLE V
PROBABILITY MATRIX OF SPECIES 1.

Bases	Position 1	Position 2	Position 3	Position 4	Position 5
A	3/3	0/3	0/3	0/3	0/3
T	0/3	3/3	3/3	0/3	0/3
G	0/3	0/3	0/3	2/3	0/3
C	0/3	0/3	0/3	1/3	3/3

During the test phase, for a query sequence, we calculated the likelihood of this being a particular species. Our model calculated the maximum likelihood and classified the new sequence to that corresponding species. For example, if we obtained a query sequence TCCAC from the test set, our model took the probability of T, C, C, A, C in position 1, 2, 3, 4, and 5, respectively, from the probability matrix of species 1, 2 and so on. If the highest probability value considering the full length of query sequence came from the matrix of species 2, the sequence was classified as species 2. Figure 1 shows an overview of this classification process.

To analyze the influence of DNA barcode sequence length on species identification, we ran our model with different sequence lengths on all the empirical and simulated datasets.

V. RESULTS AND DISCUSSIONS

After implementing our Naïve Bayes classifier, we tested the efficacy of our method. To ensure the efficacy of our model, we ran our Naïve Bayes classifier on the first five of

[4]M. Karim, R. Abid. "DNA_Barcode." Github.com. Accessed: Aug. 2021. [Online]. Available: https://github.com/MohimenulRafi/DNA_Barcode

Fig. 1. **Overview of species identification with Naïve Bayes procedure.** Each test sequence is fetched from the test set and using the probability matrices the posterior probabilities are calculated. Based on the maximum probability score, that respective test sequence is classified to corresponding species label.

the COI gene-based empirical datasets presented in Table I considering their full length barcode sequences. By comparing the results to the maximum accuracy from related studies [12], [49], where comparative performance analysis of different supervised machine learning approaches on those same COI gene-based empirical datasets are provided, we found our Naïve Bayes classifier was competent in its performance (as shown in Figure 2).

Fig. 2. **Performance competency of our Naïve Bayes classifier.** The accuracy comparison of our model and the maximum accuracy score giving models of the works of [12] and [49] is depicted in this figure. Our model yields very competing performance in each cases.

After testing our Naïve Bayes classifier on the first five empirical COI-based datasets, we ran our model on all of the COI-based empirical datasets presented in Table I to analyze the accuracy responses while varying the barcode sequence lengths. As the datasets in Table I had barcode sequence lengths > 600 bp, we set the upper bound of sequence length at 600 while varying the lengths. The sequence lengths were then decreased from the upper bound 600 bp in steps of 50 bases and the corresponding accuracy scores were calculated in each case. As 50 bp was too low a sequence length, we

considered a base length of 70 after 100 bp and maintained this as the lower bound for varying sequence lengths in the present study (with a few exceptions). The accuracy scores for the different DNA barcode lengths for the COI-based empirical datasets are presented in Table VI. These lengthwise accuracy scores in Table VI are plotted in Figure 3 and demonstrated in the heatmap in Figure 4. From the pattern of these lengthwise changes of accuracy scores in these plottings and heatmap, it seems that in most cases the accuracy scores remained persistent in the range of 200-600 bp. Although there was a gradual but tiny decrease in accuracy in this range with the decrease of barcode sequence length, it was significantly reduced below 200 bp.

TABLE VI
ACCURACY RESPONSES IN THE COI-BASED EMPIRICAL DATASET.

Datasets	Lengths											
	70	100	150	200	250	300	350	400	450	500	550	600
Bats	93.06	97.22	99.31	99.31	99.31	99.31	99.31	99.31	99.31	99.31	99.31	99.31
Bird	90.54	97.48	98.42	98.42	98.42	97.48	97.16	97.48	97.16	97.16	97.16	97.16
Cypraeidae	89.77	91.19	93.75	94.60	94.03	94.60	95.17	95.17	95.17	95.45	95.45	95.45
Drosophila	96.55	98.28	98.28	98.28	98.28	99.14	99.14	99.14	99.14	99.14	99.14	99.14
Fish	88.29	93.69	99.1	100.0	100.0	100.0	100.0	100.0	100.0	100.0	100.0	100.0
Butterfly	89.8	93.47	95.92	95.92	95.92	95.92	95.92	95.92	95.92	95.92	96.33	96.33
Bird 2	79.03	90.32	95.97	96.37	97.18	97.18	97.58	97.58	98.79	98.79	98.79	98.79
Butterfly 2	35.19	86.36	95.96	97.14	97.14	99.16	99.49	99.66	99.66	99.66	99.66	99.66
Fish 2	45.92	84.69	89.8	91.84	91.84	91.84	96.94	96.94	97.96	97.96	98.98	98.98

This similar approach was applied for the simulated datasets presented in Table III maintaing the same upper bound of 600 bp and lower bound of 70 bp. The lengthwise accuracy responses for the simulated datasets are presented in Table VII and demonstrated in Figure 5. From Figure 5 it is evident that, the accuracy score dropped insignificantly and steadily with the reduction of sequence length; however, it dropped significantly when the sequence length was reduced to < 200 bp.

TABLE VII
ACCURACY RESPONSES IN THE SIMULATED DATASETS.

Datasets	Lengths											
	70	100	150	200	250	300	350	400	450	500	550	600
Ne1000	75.91	82.63	87.52	90.34	91.88	93.21	94.06	94.84	95.26	95.66	96.02	96.32
Ne10000	81.05	86.85	91.28	93.21	94.43	95.14	95.54	95.84	96.08	96.26	96.46	96.60
Ne50000	81.62	85.80	88.65	90.14	91.09	91.56	91.99	92.26	92.55	92.73	92.92	93.09

To obtain a complete overview of the effect of length on the species identification process, the datasets presented in Table III, based on the ITS and rbcL genes, were analyzed. For the datasets *Fungi* and *Algae*, we set the upper bound and lower bound for varying sequence lengths to 850 bp and 100 bp, respectively. For the dataset *Inga*, we set 1800 bp and 1050 bp as the respective upper and lower bounds because the full lengths of the barcode sequences in this dataset were > 1800 bp. In all three datasets, we decreased the length stepwise by 50 bases. The complete lengthwise average accuracy analysis of these datasets is presented in Table VIII. This analysis is also depicted in Figure 6.

Fig. 3. **Lengthwise accuracy comparison for COI-based empirical datasets.** From the lengthwise accuracy variations of each dataset, it is apparent that the accuracy is relatively consistent in the 200 bp-600 bp range but considerably falls below 200 bp.

Fig. 4. **Heatmap of lengthwise accuracy comparison for COI-based empirical datasets.** The lengthwise accuracy variation of Figure 3 is depicted in this heatmap. It is observed from the lengthwise color variation at each row that, 200 bp-600 bp provides a relatively consistent accuracy.

Fig. 5. **Lengthwise accuracy comparison for simulated datasets.** This figure suggests a gradual but minor drop in the accuracy score when reducing the sequence length from 600 bp to 200 bp whereas a comparatively significant fall is noticed when the sequence length is reduced below 200 bp.

TABLE VIII
LENGTHWISE ACCURACY RESPONSES IN FUNGI, INGA, AND ALGAE DATASETS.

		Lengths														
Fungi	bp	100	150	200	250	300	350	400	450	500	550	600	650	700	750	800	850
	%	80.56	88.89	88.89	88.89	88.89	88.89	88.89	88.89	88.89	88.89	88.89	88.89	88.89	88.89	88.89	88.89
Inga	bp	1050	1100	1150	1200	1250	1300	1350	1400	1450	1500	1550	1600	1650	1700	1750	1800
	%	78.79	78.79	78.79	78.91	85.09	86.63	87.53	87.40	87.40	87.40	87.14	89.71	90.48	90.10	95.11	95.11
Algae	bp	100	150	200	250	300	350	400	450	500	550	600	650	700	750	800	850
	%	90.0	90.0	90.0	90.0	100.0	100.0	100.0	100.0	100.0	100.0	100.0	100.0	100.0	100.0	100.0	100.0

Table Notes: In this table, accuracy responses are demonstrated at different lengths for each of the Fungi, Inga and Algae datasets where respective results for each dataset consist of two rows. For each dataset, the first row indicates the lengths of the sequence (denoted by bp) and the second row indicates the corresponding accuracies (denoted by %).

Figure 6 shows that although the contexts differed from the previous lengthwise accuracy patterns of COI-based empirical datasets and simulated datasets, *Algae* and *Fungi* showed a sharp fall in accuracy scores for sequence lengths below 200-250 bp. However, this corroboration cannot be extended to *Inga* because its length-bounds were selected differently. However, it still showed a decrease in accuracy similar to the others when the lengths were gradually decreased from full length. To compare these three datasets on a common ground, average accuracy responses were calculated at various percentage levels of their full lengths. The results are presented in Table IX and depicted in Figure 7. No persistent range similar to the previous COI-based empirical datasets were evident from these representations of ITS and rbCL-based

Fig. 6. **Lengthwise accuracy comparison in Fungi, Inga and Algae datasets.** This figure demonstrates a sharp fall in accuracy scores for Algae and Fungi dataset around sequence lengths below 200 bp-250 bp. The length and subsequent length wise accuracy responses in Inga dataset are not similar to those of Fungi and Algae datasets, but still a decline in accuracy score can be observed with the gradual decrease of sequence length.

datasets. However, the lengthwise accuracy of *Fungi* and *Algae* decreased sharply at approximately 20 %-30 % of their full lengths whereas the accuracy in the *Inga* dataset considerably decreased initially at around 70 % and later at 40 % of its full length. For COI-based and simulated datasets, the length (~200 bp) below which a sharp decline was observed was approximately 30 % of the full length of barcode sequence. In Table VIII and Table IX, for each dataset the first row presents the sequence lengths and the second row presents the corresponding accuracy scores.

TABLE IX
ACCURACY RESPONSES IN FUNGI, INGA, AND ALGAE DATASETS AT
DIFFERENT LENGTH PERCENTAGES

		Percentages of Lengths									
		10%	20%	30%	40%	50%	60%	70%	80%	90%	100%
Fungi	**bp**	93	186	279	372	465	558	651	744	837	930
	%	79.44	93.33	93.33	93.33	93.33	93.33	93.33	93.33	93.33	93.33
Inga	**bp**	183	367	551	735	919	1102	1286	1470	1654	1838
	%	46.33	66.84	67.09	76.41	76.54	79.09	87.57	88.49	90.06	94.96
Algae	**bp**	112	225	338	451	564	676	789	902	1015	1128
	%	90.0	90.0	100.0	100.0	100.0	100.0	100.0	100.0	100.0	100.0

Table Notes: In this table, accuracy responses are demonstrated at different percentage levels of the full sequence lengths for each of the Fungi, Inga and Algae datasets where respective results for each dataset consist of two rows. For each dataset, the first row indicates the lengths of the sequence (denoted by bp) at the mentioned percentage levels and the second row indicates the corresponding accuracies (denoted by %).

Fig. 7. **Accuracy responses in Fungi, Inga and Algae datasets at different length-percentages.** As the sequence lengths of these three datasets are different, average accuracy responses were further calculated at various percentage levels of their full lengths to bring uniformity.

To support our findings, Figure 8 was adapted and redrawn from the work of [15]. In this figure, the proportion of species resolution with different DNA barcode size is depicted. The shapes in the individual plots of Figure 3 match the shape in Figure 8 implying the consistency between these separate but similar findings. From other related studies regarding mini-barcodes, it was observed that full length barcodes provide marginally better identification success rates than mini-barcodes with 200-400 bp sequence length, whereas the identification ability is significantly compromised only if the barcode length falls below 200 bp [3]. From the present study findings, an approximation of 200-350 bp mini-barcode length is proposed for use in species identification using the DNA barcode approach with a supervised machine learning model because it has a significant possibility of performing with a reasonably accurate score.

VI. CONCLUSION

In the present study, we provide an approximation of the suitable length of DNA mini-barcodes from a supervised machine learning approach by analyzing the lengthwise accuracy during the species identification process. A Naïve Bayes classifier was used to determine the effect of varying DNA barcode lengths on identification accuracy. Our findings from this machine learning perspective are consistent with those of related biological works where DNA mini-barcodes were used and analyzed to aid the species identification process. Although we proposed an approximate effective length of mini-barcodes for *COI* gene based DNA sequences, further studies are required for plants and *ITS* gene barcode based organisms. Improving the quality of the sequence alignment using suitable multiple sequence alignment tools/techniques would improve the identification accuracy. Other machine learning algorithms should also be applied to enrich the comprehensiveness of the findings.

Fig. 8. **Change of species resolution in different DNA mini-barcode lengths (adapted and redrawn from [15]).** In the demonstration of [15], it is asserted that although best performance can be achieved at full length barcode with 97 % species resolution, 100 bp and 250 bp mini barcode provide 90 % and 95 % identification success respectively.

ACKNOWLEDGMENT

We sincerely thank Dr. Md. Shamsuzzoha Bayzid for his valuable guidelines and immense support during this study.

REFERENCES

[1] Ruggiero MA, Gordon DP, Orrell TM, Bailly N, Bourgoin T, Brusca RC, et al. A higher level classification of all living organisms. PloS one. 2015;10(4):e0119248.

[2] Mora C, Tittensor DP, Adl S, Simpson AG, Worm B. How many species are there on Earth and in the ocean? PLoS Biol. 2011;9(8):e1001127.

[3] Yeo D, Srivathsan A, Meier R. Longer is not always better: Optimizing barcode length for large-scale species discovery and identification. Systematic Biology. 2020;.

[4] Dayrat B. Towards integrative taxonomy. Biological journal of the Linnean society. 2005;85(3):407–417.

[5] Armstrong K, Ball S. DNA barcodes for biosecurity: invasive species identification. Philosophical Transactions of the Royal Society B: Biological Sciences. 2005;360(1462):1813–1823.

[6] Hebert PD, Cywinska A, Ball SL, Dewaard JR. Biological identifications through DNA barcodes. Proceedings of the Royal Society of London Series B: Biological Sciences. 2003;270(1512):313–321.

[7] Hebert PD, Ratnasingham S, De Waard JR. Barcoding animal life: cytochrome c oxidase subunit 1 divergences among closely related species. Proceedings of the Royal Society of London Series B: Biological Sciences. 2003;270(suppl_1):S96–S99.

[8] Group CPW, Hollingsworth PM, Forrest LL, Spouge JL, Hajibabaei M, Ratnasingham S, et al. A DNA barcode for land plants. Proceedings of the National Academy of Sciences. 2009;106(31):12794–12797.

[9] Schoch CL, Seifert KA, Huhndorf S, Robert V, Spouge JL, Levesque CA, et al. Nuclear ribosomal internal transcribed spacer (ITS) region as a universal DNA barcode marker for Fungi. Proceedings of the National Academy of Sciences. 2012;109(16):6241–6246.

[10] Weitschek E, Van Velzen R, Felici G, Bertolazzi P. BLOG 2.0: a software system for character-based species classification with DNA Barcode sequences. What it does, how to use it. Molecular ecology resources. 2013;13(6):1043–1046.

[11] van Velzen R, Weitschek E, Felici G, Bakker FT. DNA barcoding of recently diverged species: relative performance of matching methods. PloS one. 2012;7(1):e30490.

[12] Weitschek E, Fiscon G, Felici G. Supervised DNA Barcodes species classification: analysis, comparisons and results. BioData mining. 2014;7(1):1–18.

[13] Hajibabaei M, Smith MA, Janzen DH, Rodriguez JJ, Whitfield JB, Hebert PD. A minimalist barcode can identify a specimen whose DNA is degraded. Molecular Ecology Notes. 2006;6(4):959–964.

[14] Hajibabaei M, McKenna C. DNA mini-barcodes. In: DNA barcodes. Springer; 2012. p. 339–353.

[15] Meusnier I, Singer GA, Landry JF, Hickey DA, Hebert PD, Hajibabaei M. A universal DNA mini-barcode for biodiversity analysis. BMC genomics. 2008;9(1):1–4.

[16] Staats M, Arulandhu AJ, Gravendeel B, Holst-Jensen A, Scholtens I, Peelen T, et al. Advances in DNA metabarcoding for food and wildlife forensic species identification. Analytical and Bioanalytical Chemistry. 2016;408(17):4615–4630.

[17] Erickson DL, Reed E, Ramachandran P, Bourg NA, McShea WJ, Ottesen A. Reconstructing a herbivore's diet using a novel rbc L DNA mini-barcode for plants. AoB Plants. 2017;9(3):plx015.

[18] Frézal L, Leblois R. Four years of DNA barcoding: current advances and prospects. Infection, Genetics and Evolution. 2008;8(5):727–736.

[19] Goldstein PZ, DeSalle R. Review and interpretation of trends in DNA barcoding. Frontiers in Ecology and Evolution. 2019;7:302.

[20] Adamowicz SJ, Boatwright JS, Chain F, Fisher BL, Hogg ID, Leese F, et al. Trends in DNA barcoding and metabarcoding. Genome. 2019;62(3):v–viii.

[21] Schindel DE, Miller SE. DNA barcoding a useful tool for taxonomists. Nature. 2005;435(7038):17–18.

[22] Austerlitz F, David O, Schaeffer B, Bleakley K, Olteanu M, Leblois R, et al. DNA barcode analysis: a comparison of phylogenetic and statistical classification methods. BMC bioinformatics. 2009;10(S14):S10.

[23] Casiraghi M, Labra M, Ferri E, Galimberti A, De Mattia F. DNA barcoding: a six-question tour to improve users' awareness about the method. Briefings in bioinformatics. 2010;11(4):440–453.

[24] Little DP. DNA barcode sequence identification incorporating taxonomic hierarchy and within taxon variability. PLoS one. 2011;6(8):e20552.

[25] Steinke D, Vences M, Salzburger W, Meyer A. TaxI: a software tool for DNA barcoding using distance methods. Philosophical Transactions of the Royal Society B: Biological Sciences. 2005;360(1462):1975–1980.

[26] Ratnasingham S, Hebert PD. BOLD: The Barcode of Life Data System (http://www. barcodinglife. org). Molecular ecology notes. 2007;7(3):355–364.

[27] Little DP, Stevenson DW. A comparison of algorithms for the identification of specimens using DNA barcodes: examples from gymnosperms. Cladistics. 2007;23(1):1–21.

[28] Altschul SF, Gish W, Miller W, Myers EW, Lipman DJ. Basic local alignment search tool. Journal of molecular biology. 1990;215(3):403–410.

[29] Pearson W. Rapid and sensitive sequence comparison with FASTP and FASTA. Methods in enzymology. 1990;183:63–98.

[30] Meier R, Shiyang K, Vaidya G, Ng PK. DNA barcoding and taxonomy in Diptera: a tale of high intraspecific variability and low identification success. Systematic biology. 2006;55(5):715–728.

[31] Munch K, Boomsma W, Willerslev E, Nielsen R. Fast phylogenetic DNA barcoding. Philosophical Transactions of the Royal Society B: Biological Sciences. 2008;363(1512):3997–4002.

[32] Saitou N, Nei M. The neighbor-joining method: a new method for reconstructing phylogenetic trees. Molecular biology and evolution. 1987;4(4):406–425.

[33] Farris JS. Estimating phylogenetic trees from distance matrices. The American Naturalist. 1972;106(951):645–668.

[34] Guindon S, Gascuel O. A simple, fast, and accurate algorithm to estimate large phylogenies by maximum likelihood. Systematic biology. 2003;52(5):696–704.

[35] Huelsenbeck JP, Ronquist F. MRBAYES: Bayesian inference of phylogenetic trees. Bioinformatics. 2001;17(8):754–755.

[36] Abdo Z, Golding GB. A step toward barcoding life: a model-based, decision-theoretic method to assign genes to preexisting species groups. Systematic biology. 2007;56(1):44–56.

[37] Pons J, Barraclough TG, Gomez-Zurita J, Cardoso A, Duran DP, Hazell S, et al. Sequence-based species delimitation for the DNA taxonomy of undescribed insects. Systematic biology. 2006;55(4):595–609.

[38] Munch K, Boomsma W, Huelsenbeck JP, Willerslev E, Nielsen R. Statistical assignment of DNA sequences using Bayesian phylogenetics. Systematic biology. 2008;57(5):750–757.

[39] Matz MV, Nielsen R. A likelihood ratio test for species membership based on DNA sequence data. Philosophical Transactions of the Royal Society B: Biological Sciences. 2005;360(1462):1969–1974.

[40] Sarkar IN, Planet PJ, Desalle R. CAOS software for use in character-based DNA barcoding. Molecular Ecology Resources. 2008;8(6):1256–1259.

[41] DasGupta B, Konwar KM, Măndoiu I, Shvartsman AA. DNA-BAR: distinguisher selection for DNA barcoding. Bioinformatics. 2005;21(16):3424–3426.

[42] Chu KH, Xu M, Li CP. Rapid DNA barcoding analysis of large datasets using the composition vector method. BMC bioinformatics. 2009;10(S14):S8.

[43] Kuksa P, Pavlovic V. Efficient alignment-free DNA barcode analytics. BMC bioinformatics. 2009;10(S14):S9.

[44] Nugent CM, Adamowicz SJ. Alignment-free classification of COI DNA barcode data with the Python package Alfie. Metabarcoding and Metagenomics. 2020;4:e55815.

[45] Albu M, Nikbakht H, Hajibabaei M, Hickey DA. The DNA barcode linker. Molecular ecology resources. 2011;11(1):84–88.

[46] Singer GA, Hajibabaei M. iBarcode. org: web-based molecular biodiversity analysis. In: BMC bioinformatics. vol. 10. Springer; 2009. p. S14.

[47] Lim J, Kim SY, Kim S, Eo HS, Kim CB, Paek WK, et al. BioBarcode: a general DNA barcoding database and server platform for Asian biodiversity resources. In: BMC genomics. vol. 10. Springer; 2009. p. S8.

[48] Smagala JA, Dawson ED, Mehlmann M, Townsend MB, Kuchta RD, Rowlen KL. ConFind: a robust tool for conserved sequence identification. Bioinformatics. 2005;21(24):4420–4422.

[49] Kabir T, Shemonti AS, Rahman AH. Species Identification Using Partial DNA Sequence: A Machine Learning Approach. In: 2018 IEEE 18th International Conference on Bioinformatics and Bioengineering (BIBE). IEEE; 2018. p. 235–242.

[50] He T, Jiao L, Wiedenhoeft AC, Yin Y. Machine learning approaches outperform distance-and tree-based methods for DNA barcoding of Pterocarpus wood. Planta. 2019;249(5):1617–1625.

[51] Meher PK, Sahu TK, Gahoi S, Tomar R, Rao AR. funbarRF: DNA barcode-based fungal species prediction using multiclass Random Forest supervised learning model. BMC genetics. 2019;20(1):2.

[52] Meher PK, Sahu TK, Rao A. Identification of species based on DNA barcode using k-mer feature vector and Random forest classifier. Gene. 2016;592(2):316–324.

[53] Kusuma WA, Nurilmala M, et al. Identification of Tuna and Mackerel Based on DNA Barcodes using Support Vector Machine (SVM). Telkomnika. 2016;14(2):778.

[54] Anderson MP, Dubnicka SR. A sequential naive Bayes classifier for DNA barcodes. Statistical applications in genetics and molecular biology. 2014;13(4):423–434.

[55] Porter TM, Gibson JF, Shokralla S, Baird DJ, Golding GB, Hajibabaei M. Rapid and accurate taxonomic classification of insect (class Insecta) cytochrome c oxidase subunit 1 (COI) DNA barcode sequences using a naïve Bayesian classifier. Molecular Ecology Resources. 2014;14(5):929–942.

[56] Hajibabaei M, Janzen DH, Burns JM, Hallwachs W, Hebert PD. DNA barcodes distinguish species of tropical Lepidoptera. Proceedings of the National Academy of Sciences. 2006;103(4):968–971.

[57] Lavinia PD, Núñez Bustos EO, Kopuchian C, Lijtmaer DA, García NC, Hebert PD, et al. Barcoding the butterflies of southern South America: Species delimitation efficacy, cryptic diversity and geographic patterns of divergence. PloS one. 2017;12(10):e0186845.

[58] Meyer CP, Paulay G. DNA barcoding: error rates based on comprehensive sampling. PLoS biol. 2005;3(12):e422.

[59] Lou M, Golding GB. Assigning sequences to species in the absence of large interspecific differences. Molecular Phylogenetics and Evolution. 2010;56(1):187–194.

[60] Bertolazzi P, Felici G, Weitschek E. Learning to classify species with barcodes. BMC bioinformatics. 2009;10(S14):S7.

[61] Hebert PD, Stoeckle MY, Zemlak TS, Francis CM. Identification of birds through DNA barcodes. Plos biol. 2004;2(10):e312.

[62] Kerr KC, Stoeckle MY, Dove CJ, Weigt LA, Francis CM, Hebert PD. Comprehensive DNA barcode coverage of North American birds. Molecular ecology notes. 2007;7(4):535–543.

[63] Ward RD, Zemlak TS, Innes BH, Last PR, Hebert PD. DNA barcoding Australia's fish species. Philosophical Transactions of the Royal Society B: Biological Sciences. 2005;360(1462):1847–1857.

[64] Dexter KG, Pennington TD, Cunningham CW. Using DNA to assess errors in tropical tree identifications: How often are ecologists wrong and when does it matter? Ecological Monographs. 2010;80(2):267–286.

[65] Rish I, et al. An empirical study of the naive Bayes classifier. In: IJCAI 2001 workshop on empirical methods in artificial intelligence. vol. 3; 2001. p. 41–46.

[66] Denver DR, Morris K, Lynch M, Vassilieva LL, Thomas WK. High direct estimate of the mutation rate in the mitochondrial genome of Caenorhabditis elegans. Science. 2000;289(5488):2342–2344.

[67] Katoh K, Standley DM. MAFFT multiple sequence alignment software version 7: improvements in performance and usability. Molecular biology and evolution. 2013;30(4):772–780.