Temporal scaling of human scalp-recorded potentials

Cameron D. Hassall, Jack Harley, Nils Kolling, and Laurence T. Hunt*

Oxford Centre for Human Brain Activity, Wellcome Centre for Integrative

Neuroimaging, Department of Psychiatry, University of Oxford, Oxford OX3 7JX, United

Kingdom

*laurence.hunt@psych.ox.ac.uk

Abstract

Much of human behaviour is governed by common processes that unfold over varying timescales. Standard event-related potential analysis assumes *fixed-latency* responses relative to experimental events. However, recent single unit recordings in animals have revealed neural activity *scales* to span different durations during behaviours demanding flexible timing. Here, we employed a general linear modelling approach using a novel combination of fixed-duration and variable-duration regressors to unmix fixed-time and scaled-time components in human magneto/electroencephalography (M/EEG) data. We use this to reveal consistent temporal scaling of human scalp-recorded potentials across four independent EEG datasets, including interval perception, production, prediction and value-based decision making. Between-trial variation in the temporally scaled response predicts between-trial variation in subject reaction times, demonstrating the relevance of this temporally scaled signal for temporal variation in behaviour. Our results provide a general approach for studying flexibly timed behaviour in the human brain.

Action and perception in the real world require flexible timing. We can walk quickly

or slowly, recognize the same piece of music played at different tempos, and form temporal

expectations over long and short intervals. In many cognitive tasks, reaction time variability

is modelled in terms of internal evidence accumulation [1], whereby the same dynamical

process unfolds at different speeds on different trials.

Flexible timing is critical in our lives, yet despite several decades of research [2–5] its

neural correlates remain subject to extensive debate. Due to their high temporal resolution,

magnetoencephalography and electroencephalography (M/EEG) have played a particularly

prominent role in understanding the neural basis of timing [5–13], and the method typically

used to analyze such data has been the event-related potential (ERP), which averages event-

locked responses across multiple repetitions. For example, this approach has been used to

identify the presence of a slow negative-going signal during timed intervals. This signal,

called the contingent negative variation (CNV) [14], is thought to be timing related because

its slope depends inversely on the duration of the timed interval [7,8,12].

Crucially, the ERP analysis strategy implicitly assumes that neural activity occurs at

fixed-time latencies with respect to experimental events. However, it has recently been shown

that brain activity at the level of individual neurons can be best explained by a *temporal*

scaling model [15,16], in which activity is explained by a single response that is stretched or

compressed according to the length of the produced interval. When monkeys are cued to

produce intervals of different lengths, the temporal scaling model explains the majority of

variance in neural responses from medial frontal cortex (MFC) single units [15]. This

suggests that one mechanism by which flexible motor timing can be achieved is by adjusting

the speed of a common neural process, a perspective readily viewed through the lens of

dynamical systems theory [16]. Consistent with the broad role played by dynamical systems

in a range of neural computations [17,18], recent studies in neural populations have revealed

time-warping as a common property across many different population recordings and

behavioural tasks [19]. For example, temporal scaling is also implicit in the neural correlates

of evidence integration during sensory and value-based decision making [20] (which itself

has also been proposed as a mechanism for time estimation in previous work [21]).

Successfully characterising scaled-time components in humans could open the door to

studying the role of temporal scaling in more complex, hierarchical tasks such as music

production or language perception, as well as in patient populations in which timing is

impaired [22]. Yet it is currently unclear how temporal scaling of neural responses may

manifest at the scalp (if at all) using non-invasive recording in humans. This is because of the

fixed-time nature of the ERP analysis strategy. Again, one component of the ERP called the

CNV has been found to ramp at different speeds for different temporal intervals [7,8,12],

suggestive of temporal scaling. Crucially though, any scaled activity would appear mixed at

the scalp with fixed-time components due to the *superposition problem* [23].

We therefore developed an approach to *unmix* scaled-time and fixed-time components

in the EEG (Fig 1a). Our proposed method builds on recently developed least square

regression-based approaches [24–29] that have proven useful in unmixing fixed-time

components that overlap with one another, such as stimulus-related activity and response-

related activity. To overcome the superposition problem, these approaches use a

convolutional general linear model (GLM) to deconvolve neural responses that are

potentially overlapping. Following this work, we estimate the fixed-time ERPs using a GLM

in which the design matrix is filled with time-lagged 'stick functions' (a regressor which is

valued 1 around the timepoint of interest, and 0 otherwise). Importantly, the stick functions

can overlap in time to capture overlap in the underlying neural responses (Fig 1b), and the

degree of fit to neural data can be improved by adding a regularisation penalty to the model

estimation [29]. In situations without any overlap, the GLM would exactly return the

conventional ERP.

Fig 1. Regression based unmixing of simulated data successfully recovers scaled-time and fixed-time components. (a) EEG data were simulated by summing fixed-time components (cue and response), a scaled-time component with differing durations for different trials (short, medium, or long), and noise. (b) The simulated responses were unmixed via a GLM with stick basis functions: cue-locked, response-locked, and a single scaled-time basis spanning from cue to response (i.e., variable duration). (c) The GLM successfully recovered all three components, including the scaled-time component. (d) A conventional ERP analysis (cue-locked and response-locked averages) of the same data obscured the scaled-time component.

The key innovation that we introduce here is to allow for *variable-duration* regressors

in such models, in addition to fixed-duration regressors, to test for the presence of scaled-time

responses. In particular, we allow the duration of the stick function to vary depending upon

the interval between stimulus and response, meaning that the same neural response can span

different durations on different trials. Thus, rather than modelling the mean interval duration

of each condition (e.g., via traditional ERPs), the proposed method captures trial-to-trial

response variability. The returned scaled-time potential is no longer a function of real-world

('wall clock') time, but instead a function of the *percentage of time elapsed* between stimulus

and response.

As a proof of concept, we simulated data at a single EEG sensor for an interval timing

task, consisting of two *fixed-time components* (locked to cues and responses), and one *scaled-*

time component spanning between cues and responses (Fig 1a). Our proposed method was

successful in recovering all three components (Fig 1c), whereas a conventional ERP approach

obscured the scaled-time component (Fig 1d). Crucially, in real EEG data we repeated this

approach across all sensors, potentially revealing different scalp distributions (and hence

different neural sources) for fixed-time versus scaled-time components.

By unmixing fixed and scaled components, our method goes beyond previous

approaches for dealing with timing variability in EEG experiments. For example, the event-

related timing of EEG trials can be aligned by translating either the entire waveform [30] or

individual ERP components [31]. Trials can also be aligned by scaling EEG to a common

time frame, e.g., through upsampling/downsampling [32] or dynamic time warping [33].

Finally, the fixed-time effect of continuously varying stimuli (including fast/slow speech) on

EEG can be quantified using the method of temporal response functions (TRFs), another

regression-based approach [27,34–36]. Critically, unlike the proposed method, these methods

either fail to quantify a scaled signal (translation-based methods and TRFs) or involve scaling

but no unmixing.

We also note that a time-frequency decomposition might also readily separate the

responses at higher and lower-frequencies. Indeed, a wide range of neural oscillations have

been implicated in time perception [37]. One might reasonably expect stretched/compressed

signals to manifest differently in the time-frequency domain, e.g., as they correlate more

strongly with different stretched/compressed versions of the same wavelet function. Unlike

our proposed approach, however, a time-frequency decomposition is not readily designed to

look for temporal scaling of the scaled-time response, namely the *same* neural response

unfolding over *different* timescales on different trials. Nor will a time-frequency

decomposition separate fixed-time responses from scaled-time responses if the signals

occupy the same frequency band [38].

We used our approach to analyze EEG recorded across four independent datasets,

comprising three interval timing tasks and one decision-making task. In the first task,

participants produced a target interval (short, medium, or long) following a cue (Fig 2a).

Feedback was provided, and participants were able to closely match the target intervals. In

the second, participants evaluated a computer-produced interval (Fig 2b). The closer the

produced interval was to the target interval, the more likely participants were to judge the

response as 'on time'. In the third (previously analyzed [39,40]) task, participants made

temporal predictions about upcoming events based on rhythmic predictions (Fig 2c).

In the fourth task (also previously analyzed [41,42]) participant chose between pairs

of snack items (Fig 2d) – a process in which reaction time variability can be modelled as a

process of internal evidence accumulation across time [43]. Neural activity related to

evidence accumulation is measurable on the scalp as ramping activity that scales with

decision difficulty. EEG for fast, easy trials increases at a faster rate compared to EEG for

slow, difficult trials, indicating a higher rate of internal evidence accumulation [41]. Thus, we

Fig 2. Datasets from three time-estimation and one decision-making paradigm were analyzed. In the temporal production task (a) participants successfully produced one of three cued intervals. In the temporal perception task (b) participants were able to properly judge a computer-produced interval. In a previously analyzed temporal prediction task[11,12], participants responded quickly to targets following either a rhythmic or repeated (non-rhythmic) cue. In a previously analyzed decision-making task[13,14] participants were cued to choose one of two snack food items, resulting in a range of response times (mean shown as red line). Error bars represent 95% confidence intervals.

predicted that the EEG would contain an underlying scaled component associated with

different rates of evidence accumulation.

In all four tasks, we observed a scaled-time component that was distinct from the

preceding and following fixed-time components (Fig 3), which resembled conventional ERPs

(Supplementary Fig 3). Typically, ERP components are defined by their polarity and scalp

distribution [44]. The observed scaled-time components shared a common polarity (negative)

and scalp distribution (central). In each task, cluster-based permutation testing revealed that

Fig 3. Scaled-time components were consistently observed across all four paradigms, with distinct scalp topographies from fixed-time components. Each had distinct fixed-time components relative to task-relevant events (left/middle columns), and a common negative scaled-time component over central electrodes, reflecting interval time (right column). The scalp topographies represent the mean voltage across the intervals indicated by the grey bars. For the fixed-time components, the intervals were chosen to visualize prominent deflections in the average waveform. For the scaled-time components, the intervals represent regions of significance as determined by cluster-based permutation tests. The error bars represent 95% confidence intervals.

the scaled-time component differed significantly from zero. The differences were driven by

clusters spanning 13-91% in the production task (p = .01), 34-100% in the perception task (p

= .001), 11-56% in the prediction task (p = .004), and 4-68% in the decision-making task (p

= .01). In many cases, scaled-time components were reliably observed at the single-subject

level (Supplementary Fig 5-8). To further validate our method, we quantified temporal

scaling by computing a 'scaling index' [15] for each task and participant (Fig 4). To calculate

this, we stretched/compressed each epoch to match the longest interval in each task, averaged

by condition, then calculated the coefficient of determination for predicting the longer

interval using stretched versions of the shorter intervals. We did this first on the raw data

('Original'), then separately for the data containing only the fixed-time components ('Fixed-

only', i.e., scaled-time components regressed out) and the scaled-time components ('Scaled-

only', i.e., fixed-time components regressed out). In three out of four tasks, the scaling index

for the scaled component exceeded the scaling index for the fixed component (production:

$t(9) = 3.79$, $p = .004$, Cohen's $d = 1.20$; perception: $t(9) = 1.89$, $p = .09$, Cohen's $d = 0.60$;

prediction: $t(18) = 5.45$, $p < .001$, Cohen's $d = 1.27$; decision-making: $t(17) = 5.45$, $p < .001$,

Cohen's $d = 1.29$).

Fig 4. The unmixed signals differed quantitatively in their degree of scaling. The scaling index, a measure of similarity between epochs of differing duration, was greater for the scaled-time component compared to the fixed-time component. This result was replicated in three of the four tasks.

We then examined how the scaled-time component relates to behavioural variability:

does the latency of the scaled-time component predict participants' response time? We

focussed on the temporal production and decision-making tasks, in which the interval

duration was equal to the response time. As response time varied from trial to trial, so did the

modelled scaled component. To measure component latency, we applied an approach

developed in [45,46], using principal component analysis (PCA) to model delay activity over

central electrodes in the temporal production task.. The approach works by detecting latency

shifts in a common underlying component [45,46]. Unlike simple peak detection, PCA can

account for a range of waveform dynamics (e.g., multiple peaks). We first regressed out the

fixed-time component as identified by the GLM, resulting in a dataset that consisted only of

the residual scaled-time activity. We then computed the average scaled-time activity for each

of the three interval conditions (Fig 5a-c). PCA was applied separately to each interval. This

consistently revealed a first principal component that matched the shape of the scaled-time

component and a second principal component that matched its *temporal derivative*. This

analysis confirms the presence of the scaled-time component in our data, as it is the first

principal component of the residuals after removing fixed-time components. Crucially,

adding or subtracting the second principal component captures variation in the *latency* of the

scaled-time component (Supplementary Fig 4). Across response time quantiles, we found that

PC2 scores (Supplementary Table 6) were significantly related to response times (Fig 5d),

$F(2,18) = 8.31$, $p = .003$, $\eta_p^2 = 0.48$, $\eta_g^2 = 0.43$). This implies that the earlier in time that the

scaled-time component peaked, the faster the subject would respond on that trial. This result

was replicated in the decision-making task, $F(2,34) = 43.60$, $p < .001$, $\eta_p^2 = 0.72$, $\eta_g^2 = 0.61$

(Fig 5e).

Fig 5. Variation in scaled-time components predicts behavioural variation in time estimation. Cue-locked EEG, shown as ERPs in (a), was analyzed via GLM. To visualize the unmixing of scaled-time and fixed-time components, the residual (noise) was recombined with either the scaled-time component (b) or the fixed-time components (c). PCA was run on the 'scaled-time plus residual' EEG. The second principal component resembled the temporal derivative or 'rate' of the scaled component (Supplementary Fig 5). (d) PC2 scores depended on response time, implying the scaled-time component peaked earlier for fast responses and later for slow responses. (e) The effect replicated in a decision-making task. Error bars represent 95% confidence intervals.

149   **Discussion**

Our results provide a general method for recovering temporally scaled signals in

human M/EEG, where scaled-time components are mixed at the scalp with conventional

fixed-time ERPs. We focused here on tasks that have been widely used in the timing

literature, namely interval production, perception, and prediction, as well as an example of a

cognitive task that exhibits variable reaction times across trials (value-based decision

making). Distinct scaled-time components and scalp topographies were revealed in all four

tasks. These results suggest that flexible cognition relies on temporally scaled neural activity,

as seen in recent animal work [15,16].

The existence of temporally scaled signals at the scalp may not be surprising to those

familiar with the study of time perception. Because of its excellent temporal resolution, EEG

has long been used to study delay activity in interval timing tasks. As discussed, one signal of

interest has been a ramping frontal-central signal called the CNV, which we observed in our

conventional ERP analysis (Supplementary Fig 3). Notably, CNV slope has been interpreted

as an accumulation signal in pacemaker-accumulation models of timing [7,8,11,13]. Our

work differs from these previous studies in one important respect. In a conventional ERP

analysis, delay activity is assumed to occur over fixed latencies. The CNV is thus computed

by averaging over many cue-locked EEG epochs of the same duration. In contrast, we have

considered the possibility that scalped-recorded potentials reflect a mixture of both fixed-time

and scaled-time components. By modelling fixed-time and scaled-time components

separately, we revealed scaled activity that was common across all timed intervals. This, in

turn, is consistent with a recent class of models of timing that propose time estimation reflects

the variable speed over which an underlying dynamical system unfolds [16–18].

We also observed temporally scaled activity in a decision-making task, a somewhat

surprising result given that the task did not have an explicit timing component (participants

made simple binary decisions [41]). Nevertheless, time is the medium within which decisions

are made [47]. Computationally, the timing of binary decisions can be captured in a drift

diffusion model as the accumulation of evidence in favour of each alternative [1]. This

accumulation is thought to be indexed by an ERP component called the central parietal

positivity (CPP) [48]. There is evidence that the slope of the CPP – which can be either

stimulus-locked or response-locked – captures the rate of evidence accumulation [49]. For

faster/easier decisions the CPP climbs more rapidly compared to slower/harder decisions

[48,49]. Perhaps these effects can also be explained by stretching/compressing a common

scaled-time component while holding stimulus- and response-related activity constant.

Furthermore, variation in the scaled-time component is relevant to decision making according

to our results: it predicts *when* a decision will be made. However, we also note that the

topography observed in our scaled-time component was a negative-going potential rather

than positive (Fig 3d). This can potentially be explained by the standard CPP-like ERP [41]

being a mixture of our observed negative scaled-time topography with the positive fixed-time

topographies.

Although our approach makes no assumptions about the overall shape of the scaled-

time component, it does assume a consistent, linear scaling across intervals. This is an

assumption that could be relaxed in a more complex model, e.g., using spline regression [25].

We also note that although we made no a priori predictions about waveform shape, some

between-task similarities and differences were noted in the resulting scaled-time components.

For example, similar responses were seen in the tasks for which the interval of interest ended

with a motor response (temporal production and decision making – see Fig 3a,d). In both

cases, activity immediately preceding the response depended on a ramping, fixed-time,

motor-related component, with little contribution from a scaled component. A similar

observation was made in the temporal prediction task – activity just before the appearance of

the target depended on anticipatory fixed-time activity, not scaled activity (Fig 3c). In

contrast, pre-probe activity in the temporal perception task showed almost no fixed-time

activity, but a robust scaled-time component (Fig 3b). The reason for this difference cannot

be identified by the current experiments, however. First, the perception and prediction tasks

involved different tasks instructions ('listen for the probe' versus 'respond to the target').

Second, the probe/target distributions differed in the two tasks; the mean duration was 75%

likely in the prediction task, but only 20% likely in the perception task. We therefore

speculate that scaled activity may be somewhat task dependent.

Our approach is not only conceptually different from previous work that models

variability in timing using a regression framework [27,34–36], it is also a mechanistically

important finding. It indicates the brain may support flexible timing by adapting the duration

of an otherwise consistent neural response. This can be understood as varying the rate of a

dynamical system [17,18] during interval estimation. Although there is evidence for such

temporally scaled responses in the monkey neurophysiology literature (e.g., [15,16], which

inspired the current study), we are not aware of any direct evidence in support of this idea in

humans. Indeed, it goes against the dominant framework of fixed-latency responses that has

thus far dominated M/EEG analysis.

Although we have focused here on interval timing and decision-making tasks, we

anticipate other temporally-scaled EEG and MEG signals will be discovered for cognitive

processes known to unfold over varying timescales. For example, the neural basis of flexible

(fast/slow) speech production and perception is an active area of research [50–52], and may

involve a form of temporal scaling [32]. Flexible timing is also important across a vast array

of decision-making tasks, where evidence accumulation can proceed quickly or slowly

depending on the strength of the evidence [20]. Flexible timing helps facilitate a range of

adaptive behaviours via temporal attention [4], while disordered timing characterizes several

clinical disorders [53], underscoring the importance of characterising temporal scaling of

neural responses in human participants.

Acknowledgements

This research was funded by a Natural Sciences and Engineering Research Council of Canada (NSERC: https://www.nserc-crsng.gc.ca/) Postdoctoral Fellowship to C.D.H. (PDF 546078 – 2020), a BBSRC (https://bbsrc.ukri.org/) AFL Fellowship (BB/R010803/1) to N.K., a Sir Henry Dale Fellowship from the Royal Society (https://royalsociety.org) and Wellcome (https://wellcome.org/) to L.T.H. (208789/Z/17/Z), a NARSAD Young Investigator Award from the Brain and Behavior Research Foundation (https://www.bbrfoundation.org/) to L.T.H., and supported by the NIHR Oxford Health Biomedical Research Centre (https://oxfordhealthbrc.nihr.ac.uk/). The Wellcome Centre for Integrative Neuroimaging was supported by core funding from Wellcome Trust (203139/Z/16/Z). For the purpose of Open Access, the author has applied a CC BY public copyright licence to any Author Accepted Manuscript version arising from this submission.

Author Contributions

N.K. and L.T.H. conceived the experiments and methodology. C.D.H. and L.T.H. designed the experiments and developed the methodology. C.D.H. and J.H. performed the experiments (except for the prediction task, where data was downloaded from a previous publication [39,40]). C.D.H. and L.T.H. analyzed the data. C.D.H. and L.T.H. wrote the manuscript with input from the other authors.

Competing interests

The authors declare no competing interests.

Additional information

Supplementary information is available for this paper.

Correspondence and requests for materials should be addressed to L.T.H.

 Methods

 Simulations

228  We simulated cue-related and response-related EEG in a temporal production task

229 using MATLAB 2020a (Mathworks, Natick, USA). Cue and response were separated by

230 either a short, medium, or long interval. During the delay period, we simulated a scaled

231 response that stretched or compressed to fill the interval. All three responses (cue, response,

232 scaled) were summed together at appropriate lags (short, medium, or long), with noise – see

233 Fig 1a. In total, we simulated 50 trials of each condition (short, medium, long).

234  To unmix fixed-time and scaled-time components, we used a regression-based

235 approach [24,25,54] in which the continuous EEG at one sensor *Y* is modelled as a linear

236 combination of the underlying event-related responses β, which are unknown initially. The

237 model can be written in equation form as:

$$Y = X\beta + \varepsilon$$

239  where *X* is the design matrix and ε is the residual EEG not accounted for by the

240 model. *X* contains as many rows as EEG data points, and as many columns as predictors (that

241 is, the number of points in the estimated event-related responses). In our case, *X* was

242 populated by 'stick functions' – non-zero values around the time of the modelled events, and

243 zeros otherwise. We included in *X* two fixed-time components, the cue and the response, as

244 stick functions of fixed EEG duration (with variables set to 1). In other words, the height of

245 the fixed-time stick function was constant across events of the same type and equal to its

246 width. To model a temporally-scaled response, we used the MATLAB *imresize* function

247 (Image Processing Toolbox, R2020b) with 'box' interpolation to stretch/compress a stick

248 function so that it spanned the duration between cue and response (other interpolation

249 methods were tried – see Supplementary Fig 1 – but this choice had little effect on the

250 results). Thus, the duration of the scaled stick function varied from trial to trial (Fig 1b). The

goal here was to estimate a single scaled-time response to account for EEG activity across

multiple varying delay periods. For the fixed-time responses, each column of X represents a

latency in ms before/after an experimental event; by contrast, for the scaled-time responses,

each column of X represents the *percentage* of time that has elapsed between *two* events

(stimulus and response). Simulation code is available at

https://git.fmrib.ox.ac.uk/chassall/temporal-scaling.

Production and Perception Tasks

Participants

Participants completed both the production and perception tasks within the same

recording session. We tested ten university-aged participants, 5 male, 2 left-handed, $M_{age} =$

23.40, 95% CI [21.29, 25.51]. Participants had normal or corrected-to-normal vision and no

known neurological impairments. Participants provided informed consent approved by the

Medical Sciences Interdivisional Research Ethics Committee at the University of Oxford.

Following the experiment, participants were compensated £20 (£10 per hour of participation)

plus a mean performance bonus of £3.23, 95% CI [2.92, 3.55].

Apparatus and Procedure

Participants were seated approximately 64 cm from a 27-inch LCD display (144 Hz, 1

ms response rate, 1920 by 1080 pixels, Acer XB270H, New Taipei City, Taiwan). Visual

stimuli were presented using the Psychophysics Toolbox Extension [55,56] for MATLAB

2014b (Mathworks, Natick, USA). Participants were given written and verbal instructions to

minimize head and eye movements. The goal of the production task was to produce a target

interval and the goal of the perception task was to judge whether or not a computer-produced

interval was correct.

The experiment was blocked with ten trials per block. There were 18 production

blocks and 18 perception blocks, completed in random order. Prior to each block, participants

listened to five isochronic tones indicating the target interval. Beeps were 400 Hz sine waves

of duration 50 ms and an onset/offset ramping to a point 1/8 of the length of the wave (to

avoid abrupt transitions). The target interval was either short (0.8 s), medium (1.65 s), or

long (2.5 s).

In production trials, participants listened to a beep then waited the target time before

responding. Feedback appeared after a 400-600 ms delay (uniform distribution) and remained

on the display for 1000 ms. Feedback was a 'quarter-to' clockface to indicate 'too early', a

'quarter-after' clockface to indicate 'too late', or a checkmark to indicate an on-time

response. Feedback itself was determined by where the participant's response fell relative to a

window around the target duration. The response window was initialized to +/- 100 ms

around each target, then changed following each feedback via a staircase procedure:

increased on each side by 10 ms following a correct response and decreased by 10 ms

following an incorrect response (either too early or too late).

In perception trials, participants heard two beeps, then were asked to judge the

correctness of the interval, that is, whether or not the test interval matched the target interval.

Test intervals (very early, early, on time, late, very late) were set such that each subsequent

interval was 25% longer than the previous (see Supplementary Table 1). Participants were

then given feedback on their judgement – a checkmark for a correct judgement, or an 'x' for

an incorrect judgement.

For each task, participants gained 2 points for each correct response and lost 1 point

for each incorrect response. At the end of the experiment points were converted to a monetary

bonus at a rate of £0.01 per point.

Data Collection

In the perception task we recorded participant response time from cue, trial outcome

(early, late, on time), and staircase- response window. In the production task, we recorded

trial 'on time' judgements (yes/no), and trial outcome (correct/incorrect).

We recorded 36 channels of EEG, referenced to AFz. Data were recorded at 1000 Hz

using a Synamps amplifier and CURRY 8 software (Compumetrics Neuroscan, Charlotte,

USA). The electrodes were sintered Ag/AgCl (EasyCap, Herrsching, Germany). 31 of the

electrodes were laid out according to the 10-20 system. Additional electrodes were placed on

the left and right mastoids, on the outer canthi of the left and right eyes, and below the right

eye. The reference electrode was placed at location AFz, and the ground electrode at Fpz.

Prediction Task

In this previously published [39,40] experiment, 19 participants responded to the

onset of a visual target following a visual warning cue. The delay between cue and target was

either short (700 ms) or long (1300 ms) and, in some conditions, congruent with a preceding

stimulus stream. Only congruent trials were included in the current analysis (i.e., the 'valid'

trials in the 'rhythmic' and 'repeated' conditions). Each trial was preceded by a 500 ms

fixation cross subtending 0.6° of visual angle. During the pre-cue period, participants were

shown a flashing stimulus for 4-6 repetitions to indicate the target interval. The stimulus was

a centrally presented black disc (1.2°) that appeared on the display for 100 ms. In the

rhythmic condition the black disc appeared every 700 ms or 1300 ms ('short' or 'long'). In

the repeated condition, a red disc appeared either 700 ms or 1300 ms after the appearance of

the black disc, followed by a variable delay period of either 1500-1900 ms (short) or 1900-

2700 ms (long). Following the pre-cue period participants were then shown the warning cue,

a white disc (1.2°) that appeared for 100 ms. After either a short or long delay (700 ms or

1300 ms) the target appeared – a green 1.2° disc – for 100 ms, followed by the participant's

response. The experimental program recorded the response time (time since the onset of the

target). See Supplementary Fig 2c and [39,40] for more detail.

Decision-Making Task

In this experiment, also previously published [41,42], 18 participants were presented

with two snack foods and asked to pick one. This was not an interval timing task and on

average participants took 763 ms, 95% CI [713, 813], to respond. Trials began with the

appearance of a centrally presented fixation cross (0.6°) for 2-4 s followed by the

presentation of the snack items (3° across, in total). Participants were asked to indicate their

preference by making a left or right button press within a 1.25 s window. The experimental

program recorded the response time (time since the onset of the snack items). See

Supplementary Fig 2d and [41] for more detail.

Data Analysis

Behavioural data

For the production task, we computed the mean produced interval for each participant.

For the perception task, we computed mean likelihood of responding yes to the 'on time'

prompt, for each condition (short, medium, long) and interval (very early, early, on time, late,

very late). For the prediction task, we computed the mean reaction time for each analyzed

condition (rhythmic, repeated) and interval (short, long). For the decision-making task, we

computed the mean response (decision) time. See Fig 2 and Supplementary Tables 2-4 for

behavioural results.

EEG Preprocessing

For all three timing tasks, EEG was preprocessed in MATLAB 2020b (Mathworks,

Natick, USA) using EEGLAB [57]. We first down-sampled the EEG to 200 Hz, then applied

a 0.1-20 Hz bandpass filter and 50 Hz notch filter. The EEG was then re-referenced to the

average of the left and right mastoids (and AFz recovered in the production/perception tasks).

Ocular artifacts were removed using independent component analysis (ICA). The ICA was

trained on 3-second epochs of data following the appearance of the fixation cross at the

beginning of each trial. Ocular components were identified using the *iclabel* function and

then removed from the continuous data.

EEG for the decision-making task was already preprocessed prior to our analysis. This

was a simultaneous EEG-fMRI recording, and preprocessing included the removal of MR-

related artifacts via filtering and principal component analysis, as well as a 0.5-40 Hz

bandpass filter. In line with our other analyses, we re-referenced the EEG to the average of

TP7 and TP8 (located close to the mastoids) and applied an additional 20 Hz low-pass filter.

ERPs

To construct conventional event-related potentials (ERPs), we first created epochs of

EEG around cues (all tasks), responses (perception task), probes (production task), targets

(prediction task), and decisions (decision-making task). Cue-locked ERPs extended from 200

ms pre-cue to either 800, 1650, or 2500 ms post-cue (the short, medium, and long targets) in

the perception/production tasks, 700 or 1300 ms in the prediction task (the short and long

targets), and 600 ms in the decision-making task. Epochs were baseline-corrected using a 200

ms pre-cue window. We also constructed epochs from 800, 1650, or 2500 ms prior to the

response/probe in the production/perception tasks, 700 or 1300 ms prior to the target in the

prediction task, and 600 prior to the decision in the decision-making task to 200 ms after the

response/probe/target/decision. A baseline was defined around the event of interest (mean

EEG from -20 to 20 ms) and removed in all cases except for the decision-making task, in line

with the original analysis [41]. We then removed any trials in which the sample-to-sample

voltage differed by more than 50 μV or the voltage change across the entire epoch exceeded

150 μV. We then created conditional cue and response/probe/target/decision averages for

each participant and task: production/perception (short, medium, and long), prediction (short

and long), and decision-making (early and late, via a median split [41]). Finally, participant

averages in the timing tasks were combined into grand-average waveforms at electrode FCz,

a location where timing-related activity has been previously observed [5] and Pz in the

decision-making task, in line with the previously published analysis [41] (Supplementary Fig

3).

378    *rERPs*

379    To unmix fixed-time and scaled-time components in our EEG data, we estimated

regression-ERPs (rERPs) following the same GLM procedure we used with our simulated

data, but now applied to each sensor. We used a design matrix consisting of a regular stick

functions for cue and response/probe/target and a stretched/compressed stick function

spanning the interval from cue to response/probe/target/decision. In particular, we estimated

cue-locked responses that spanned from 200 ms pre-cue to 800 ms post-cue. The

response/probe/target/decision response interval spanned from -800 to 200 ms. Each fixed-

time response thus spanned 1000 ms, or 200 EEG sample points. The scaled-time component,

as described earlier, was modelled as a single underlying component (set width in X) that

spanned over multiple EEG durations (varying number of rows in X). Thus, our method

required choosing how many scaled-time sample points to estimate (the width in X). For the

production/perception tasks, we chose to estimate 330 scaled-time points, equivalent to the

duration of the 'medium' interval. For the prediction task, we chose to estimate 200 scaled-

time points, equivalent to the mean of the short and long conditions (700 ms, 1300 ms). For

the decision-making task, we estimated 153 scaled-time points (roughly equivalent to the

mean decision time). Unlike the conventional ERP approach, this analysis was conducted on

the continuous EEG. To identify artifacts in the continuous EEG, we used the *basicrap*

function from the ERPLAB [58] toolbox with a 150 µV threshold (2000 ms window, 1000

ms step size). A sample was flagged if it was 'bad' for any channel. Flagged samples were

excluded from the GLM (samples removed from the EEG and rows removed from the design

matrix). Additionally, we removed samples/rows associated with unusually fast or slow

responses in the production task (less than 0.2 s or more than 5 s). On average, we removed

2.17 % of samples in the production task (95% CI [1.49, 2.86]), 3.75 % of samples in the

perception task (95% CI [2.39, 5.10]), 1.03% of samples in the prediction task (95% CI [0.95,

1.10]), and 5.56% of samples in the decision-making task (95% CI [4.99, 6.12]).

To test for multicollinearity between the regressors we computed the variance

inflation factor (VIF) for each regressor, i.e., at each timepoint in the estimated waveforms.

This was done using the *uf_vif* function in the Unfold toolbox [25]. We were concerned about

multicollinearity because the fixed-time and scaled-time components occurred over the same

'real time' durations. For example, in the production task the early and later parts of the

scaled waveform always coincided with the start of the cue-locked and end of the response-

locked responses, respectively. The overlap was not consistent, however; alignment between

the fixed and scaled regressors was lessened due to distortions in the scaled stick function

(see Supplementary Fig 1). As a result, the VIF was low (< 10) at nearly all points other than

the start/end (Supplementary Fig 9). This was true in all tasks except for the temporal

prediction task (VIFs > 10), as these tasks incorporated greater temporal variability across

trials. We therefore expected the waveform estimates in the temporal prediction task to be

noisier relative to the other tasks. We note that future studies could use VIF to evaluate the

likelihood of successfully unmixing fixed-time and scaled-time components. Introducing

elements of experimental design (such as increased interval variability across trials) could

help to address concerns over multicollinearity.

To lessen the effect of multicollinearity and impose a smoothness constraint on our

estimates, we used a first-derivative form of Tikhonov regularization [29]. Tikhonov

regularization reframes the GLM solution as the minimization of:

$$423 \qquad \|X\beta - Y\|^2 + \lambda\|L\beta\|^2$$

424   where L is the regularization operator and λ is the regularization parameter. In other

425   words, we aimed to minimize a penalty term in addition to the usual residual. This has the

426   solution

$$427 \qquad (X^T X + \lambda L)^{-1} X^T Y$$

428   In our case, L approximated the first derivative as a scaled finite difference[59]:

$$429 \qquad L = \frac{1}{2}\begin{bmatrix} 1 & -1 & 0 & ... & 0 & 0 \\ 0 & 1 & -1 & ... & 0 & 0 \\ ... & ... & ... & ... & ... & ... \\ 0 & 0 & 0 & ... & 1 & -1 \end{bmatrix}$$

We then chose regularization parameters for each participant using 10-fold cross

validation. Our goal here was to minimize the mean-squared error of the residual EEG at

electrode FCz, our electrode of interest. The following λs were tested on each fold: 0.001,

0.01, 0, 1, 10, 100, 1000, 10000, 100000. An optimal λ was chosen for each participant

corresponding to the parameter with the lowest mean mean-squared error across all folds. See

Supplementary Table 7 for a summary.

Statistics

We quantified the amplitude of the scaled-time component in two ways. First, we

computed and plotted the 95% confidence interval at each 'timepoint' of the scaled-time

signal (right column of Fig 3). Next, we conducted a nonparametric statistical test of the

scaled-time component according to the procedure outlined in [60]. After computing a single-

sample t-statistic at each sample point, we identified clusters of points for with the t-value

exceeded a critical threshold corresponding to an alpha level of .05 (i.e., the 2.5th and 97.5th

percentiles, making this a two-sided test). For each cluster, we computed the 'cluster mass',

defined as the sum of the absolute values of the t-values within the cluster. To determine

whether the observed cluster masses exceeded what could occur by chance, we permuted the

scaled components by randomly flipping (multiplying by -1) the entire waveform. We then

computed the cluster masses for the permuted waveforms as before, and recorded the

maximum cluster mass (or zero if there were no clusters). Every possible permutation was

checked for the production and perception tasks ($2^{10} = 1024$ permutations). For the prediction

and decision-making tasks, 1000 random permutations were checked. We then labelled our

observed cluster masses as 'significant' if they exceeded 95% of the maximum cluster masses

of the permuted waveforms. These analyses were done at the scalp location where the mean

scaled-time signal reached its most negative value, i.e. electrode Cz in the

production/perception tasks, FCz in the prediction task, and C1 in the decision-making task.

Scaling Index

To validate the unmixing procedure, we regressed out either the scaled-time

component or the fixed-time components from the EEG in each task and participant to create

'fixed-only' or 'scaled-only' data sets. We then quantified the amount of temporal scaling

present in each task, participant, and data set (original, fixed-only, scaled only) using a

similar procedure as [15]. Specifically, we constructed epochs spanning the intervals of

interest (e.g., cue to response), then stretched or compressed each epoch to match a common

duration (the longest duration in the interval timing tasks; the mean of the 'late' responses in

the decision-making task, as defined above). For each task and participant, we averaged by

condition (e.g., short, medium, long) to create conditional ERPs with a common duration,

then computed a scaling index defined as the coefficient of determination. Specifically, we

asked how well the 'long' waveform could be predicted by the 'short' waveform. If there was

also a 'medium' waveform (the production/perception tasks) another coefficient of

determination was computed, and the two coefficients were averaged. A larger scaling index

can therefore be interpreted as a greater post-scaling similarity between conditions. Scaling

indices in the fixed-only and scaled-only data sets were compared via paired-samples *t*-tests.

For each *t*-test, we computed Cohen's *d* as:

$$Cohen's\ d = \frac{M_{diff}}{s_{diff}}$$

where M_{diff} is the mean difference between the scores being compared and s_{diff} is the

standard deviation of the difference of the scores being compared [61]. Interestingly, the

scaling index of the original signal appeared to be a mixture of the scaling indices of the

fixed-only and scaled-only signals in all tasks except for the temporal prediction task (Fig 4).

We interpreted this as further evidence that the unmixing procedure was less effective here

due to multicollinearity.

PCA

To explore the link between the scaled-time component and behaviour, we examined

the scaled-only data set described above – that is, the scaled-time regressors plus residuals.

Only mid-frontal electrodes were considered: FC1, FCz, FC2, Cz, CP1, CPz, and CP2. We

then constructed epochs starting at the cue and ending at the target interval (800 ms, 1650 ms,

or 2500 ms). Epochs within each condition (short, medium, long) were further grouped into

three equal-sized response-time bins (early, on time, late) and averaged for each electrode

and participant. We then conducted a PCA for each condition (short, medium, long) and

participant. See Supplementary Table 5 for amount of variance explained by PC1 and PC2.

To visualize the effect of PC2, we computed the mean PC2 across all participants. We then

added more or less of the mean PC2 to the mean PC1 projection and applied a 25-point

moving-mean window for visualization purposes (Supplementary Fig 4). In order to choose a

reasonable range of PC2 scores, we examined the average minimum and maximum PC2

score for each participant and condition (short, medium, long). The PC2 score ranges were -

21 to 15 (short), -41 to 38 (medium), and -40 to 55 (long). To assess the relationship between

PC2 score and behaviour, we binned PC2 scores according to our response time bins (early,

on time, late) and collapsed across conditions (short, medium, long). This gave us as single

mean PC2 score for each participant and response time bin (early, on time, late), which we

analyzed using a two-sided repeated-measures ANOVA (Fig 5d) after verifying the

assumption of normality using the Shapiro-Wilk test. Two different effect sizes, η_p^2 and η_g^2,

were computed, according to:

$$\eta_p^2 = \frac{SS_Q}{SS_Q + SS_{sQ}} \qquad \eta_g^2 = \frac{SS_Q}{SS_Q + SS_S + SS_{sQ}}$$

where SS_Q is the sum of squares of the quantile effect (early, on time, late), SS_{sQ} is the error

sum of squares of the quantile effect, and SS_S is the sum of squares between subjects [62].

We then replicated the PCA procedure for the decision-making task using an epoch

extending 800 ms from the cue at a central electrode cluster (FC3, FC1, FC2, FC4, C3, C1,

Cz, C2, C4, CP3, CP1, CP2, CP4, P3, P1, Pz, P2, and P4). Note that the assumption of

normality was violated for 'early' responses in the decision-making task. However, as

repeated-measures ANOVA is robust to violations of normality, no statistical correction was

made.

Data Availability

Raw and preprocessed EEG for the production and perception tasks will be made

publicly available at the time of publication. Raw data for the prediction task is available at

https://doi.org/10.5061/dryad.5vb8h. Raw data for the decision-making task is available at

https://openneuro.org/datasets/ds002734/versions/1.0.2.

Code Availability

Simulation and analysis scripts are available athttps://git.fmrib.ox.ac.uk/chassall/temporal-

scaling.

References

1. Bogacz R, Brown E, Moehlis J, Holmes P, Cohen JD. The Physics of Optimal Decision
Making: A Formal Analysis of Models of Performance in Two-Alternative Forced-Choice
Tasks. Psychological Review. 113: 700–765.

2. Merchant H, Harrington DL, Meck WH. Neural Basis of the Perception and Estimation of
Time. Annu Rev Neurosci. 2013;36: 313–336. doi:10.1146/annurev-neuro-062012-
170349

3. Wittmann M. The inner sense of time: how the brain creates a representation of
duration. Nature Reviews Neuroscience. 2013;14: 217–223. doi:10.1038/nrn3452

4. Nobre AC, van Ede F. Anticipated moments: temporal structure in attention. Nature
Reviews Neuroscience. 2018;19: 34–48. doi:10.1038/nrn.2017.141

5. Macar F, Vidal F. Event-Related Potentials as Indices of Time Processing: A Review.
Journal of Psychophysiology. 2004;18: 89–104. doi:10.1027/0269-8803.18.23.89

6. Kononowicz TW, van Rijn H. Decoupling Interval Timing and Climbing Neural Activity: A
Dissociation between CNV and N1P2 Amplitudes. J Neurosci. 2014;34: 2931–2939.
doi:10.1523/JNEUROSCI.2523-13.2014

7. Pfeuty M, Ragot R, Pouthas V. Relationship between CNV and timing of an upcoming
event. Neuroscience Letters. 2005;382: 106–111. doi:10.1016/j.neulet.2005.02.067

8. Macar F, Vidal F, Casini L. The supplementary motor area in motor and sensory timing:
evidence from slow brain potential changes. Exp Brain Res. 1999;125: 271–280.
doi:10.1007/s002210050683

9. Trillenberg P, Verleger R, Wascher E, Wauschkuhn B, Wessel K. CNV and temporal
uncertainty with 'ageing' and 'non-ageing' S1–S2 intervals. Clinical Neurophysiology.
2000;111: 1216–1226. doi:10.1016/S1388-2457(00)00274-1

10.   Miniussi C, Wilding EL, Coull JT, Nobre AC. Orienting attention in time: Modulation of
brain potentials. Brain. 1999;122: 1507–1518. doi:10.1093/brain/122.8.1507

11.   Macar F, Vidal F. The CNV peak: An index of decision making and temporal memory.
Psychophysiology. 2003;40: 950–954. doi:10.1111/1469-8986.00113

12.   Praamstra P, Kourtis D, Kwok HF, Oostenveld R. Neurophysiology of Implicit Timing in
Serial Choice Reaction-Time Performance. J Neurosci. 2006;26: 5448–5455.
doi:10.1523/JNEUROSCI.0440-06.2006

13.   Macar F, Vidal F. Time processing reflected by EEG surface Laplacians. Exp Brain Res.
2002;145: 403–406. doi:10.1007/s00221-002-1103-z

14.   Walter WG, Cooper R, Aldridge VJ, McCallum WC, Winter AL. Contingent Negative
Variation : An Electric Sign of Sensori-Motor Association and Expectancy in the Human
Brain. Nature. 1964;203: 380–384. doi:10.1038/203380a0

15.   Wang J, Narain D, Hosseini EA, Jazayeri M. Flexible timing by temporal scaling of
cortical responses. Nat Neurosci. 2018;21: 102–110. doi:10.1038/s41593-017-0028-6

16.   Remington ED, Egger SW, Narain D, Wang J, Jazayeri M. A Dynamical Systems
Perspective on Flexible Motor Timing. Trends in Cognitive Sciences. 2018;22: 938–952.
doi:10.1016/j.tics.2018.07.010

17.   Vyas S, Golub MD, Sussillo D, Shenoy KV. Computation Through Neural Population
Dynamics. Annual Review of Neuroscience. 2020;43: 249–275. doi:10.1146/annurev-
neuro-092619-094115

18.   Sussillo D. Neural circuits as computational dynamical systems. Current Opinion in
Neurobiology. 2014;25: 156–163. doi:10.1016/j.conb.2014.01.008

19.   Williams AH, Poole B, Maheswaranathan N, Dhawale AK, Fisher T, Wilson CD, et al.
Discovering Precise Temporal Patterns in Large-Scale Neural Recordings through
Robust and Interpretable Time Warping. Neuron. 2020;105: 246-259.e8.
doi:10.1016/j.neuron.2019.10.020

20.   O'Connell RG, Shadlen MN, Wong-Lin K, Kelly SP. Bridging Neural and Computational
Viewpoints on Perceptual Decision-Making. Trends in Neurosciences. 2018;41: 838–
852. doi:10.1016/j.tins.2018.06.005

21.   Boehm U, van Maanen L, Forstmann B, van Rijn H. Trial-by-trial fluctuations in CNV
amplitude reflect anticipatory adjustment of response caution. Neuroimage. 2014;96:
95–105. doi:10.1016/j.neuroimage.2014.03.063

22.   Meck WH. Neuropsychology of timing and time perception. Brain and Cognition.
2005;58: 1–8. doi:10.1016/j.bandc.2004.09.004

23.   Luck SJ. A Closer Look at ERPs and ERP Components. Second edition. An introduction to
the event-related potential technique. Second edition. Cambridge, Massachusetts: The
MIT Press; 2014. pp. 35–70.

24.   Smith NJ, Kutas M. Regression-based estimation of ERP waveforms: I. The rERP
framework. Psychophysiology. 2015;52: 157–168. doi:10.1111/psyp.12317

25.   Ehinger BV, Dimigen O. Unfold: an integrated toolbox for overlap correction, non-linear
modeling, and regression-based EEG analysis. PeerJ. 2019;7: e7838.
doi:10.7717/peerj.7838

26.   Litvak V, Jha A, Flandin G, Friston K. Convolution models for induced electromagnetic
responses. Neuroimage. 2013;64: 388–398. doi:10.1016/j.neuroimage.2012.09.014

27.   Lalor EC, Pearlmutter BA, Reilly RB, McDarby G, Foxe JJ. The VESPA: A method for the
rapid estimation of a visual evoked potential. NeuroImage. 2006;32: 1549–1561.
doi:10.1016/j.neuroimage.2006.05.054

28. Crosse MJ, Di Liberto GM, Bednar A, Lalor EC. The Multivariate Temporal Response
Function (mTRF) Toolbox: A MATLAB Toolbox for Relating Neural Signals to Continuous
Stimuli. Front Hum Neurosci. 2016;10. doi:10.3389/fnhum.2016.00604

29. Kristensen E, Guerin-Dugué A, Rivet B. Regularization and a general linear model for
event-related potential estimation. Behav Res. 2017;49: 2255–2274.
doi:10.3758/s13428-017-0856-z

30. Woody CD. Characterization of an adaptive filter for the analysis of variable latency
neuroelectric signals. Med & biol Engng. 1967;5: 539–554. doi:10.1007/BF02474247

31. Ouyang G, Herzmann G, Zhou C, Sommer W. Residue iteration decomposition (RIDE): A
new method to separate ERP components on the basis of latency variability in single
trials: RIDE: A new method to separate ERP components. Psychophysiology. 2011;48:
1631–1647. doi:10.1111/j.1469-8986.2011.01269.x

32. Lerner Y, Honey CJ, Katkov M, Hasson U. Temporal scaling of neural responses to
compressed and dilated natural speech. Journal of Neurophysiology. 2014;111: 2433–
2444. doi:10.1152/jn.00497.2013

33. Huang H-C, Jansen BH. EEG waveform analysis by means of dynamic time-warping.
International Journal of Bio-Medical Computing. 1985;17: 135–144. doi:10.1016/0020-
7101(85)90084-4

34. Herbst SK, Fiedler L, Obleser J. Tracking Temporal Hazard in the Human
Electroencephalogram Using a Forward Encoding Model. eNeuro. 2018;5:
ENEURO.0017-18.2018. doi:10.1523/ENEURO.0017-18.2018

35. Kulasingham JP, Joshi NH, Rezaeizadeh M, Simon JZ. Cortical Processing of Arithmetic
and Simple Sentences in an Auditory Attention Task. J Neurosci. 2021;41: 8023–8039.
doi:10.1523/JNEUROSCI.0269-21.2021

36. Gonçalves NR, Whelan R, Foxe JJ, Lalor EC. Towards obtaining spatiotemporally precise
responses to continuous sensory stimuli in humans: A general linear modeling
approach to EEG. NeuroImage. 2014;97: 196–205.
doi:10.1016/j.neuroimage.2014.04.012

37. Wiener M, Kanai R. Frequency tuning for temporal perception and prediction. Current
Opinion in Behavioral Sciences. 2016;8: 1–6. doi:10.1016/j.cobeha.2016.01.001

38. Herrmann CS, Rach S, Vosskuhl J, Strüber D. Time–Frequency Analysis of Event-Related
Potentials: A Brief Tutorial. Brain Topogr. 2014;27: 438–450. doi:10.1007/s10548-013-
0327-5

39. Breska A, Deouell LY. Neural mechanisms of rhythm-based temporal prediction: Delta
phase-locking reflects temporal predictability but not rhythmic entrainment. PLOS
Biology. 2017;15: e2001665. doi:10.1371/journal.pbio.2001665

40.   Breska A, Deouell LY. Data from: Neural mechanisms of rhythm-based temporal
prediction: delta phase-locking reflects temporal predictability but not rhythmic
entrainment. 2017. doi:doi:10.5061/dryad.5vb8h

41.   Pisauro MA, Fouragnan E, Retzler C, Philiastides MG. Neural correlates of evidence
accumulation during value-based decisions revealed via simultaneous EEG-fMRI. Nat
Commun. 2017;8: 1–9. doi:10.1038/ncomms15808

42.   Pisauro MA, Fouragnan E, Retzler C, Philiastides MG. Evidence Accumulation in Value-
Based decisions. 2020. doi:10.18112/openneuro.ds002734.v1.0.2

43.   Krajbich I, Armel C, Rangel A. Visual fixations and the computation and comparison of
value in simple choice. Nature Neuroscience. 2011;13: 10.

44.   Luck SJ. An introduction to the event-related potential technique. Second edition.
Cambridge, Massachusetts: The MIT Press; 2014.

45.   Möcks J. The Influence of Latency Jitter in Principal Component Analysis of Event-
Related Potentials. Psychophysiology. 1986;23: 480–484. doi:10.1111/j.1469-
8986.1986.tb00659.x

46.   Hunt LT, Behrens TE, Hosokawa T, Wallis JD, Kennerley SW. Capturing the temporal
evolution of choice across prefrontal cortex. Frank MJ, editor. eLife. 2015;4: e11945.
doi:10.7554/eLife.11945

47.   Ariely D, Zakay D. A timely account of the role of duration in decision making. Acta
Psychologica. 2001;108: 187–207. doi:10.1016/S0001-6918(01)00034-8

48.   O'Connell RG, Dockree PM, Kelly SP. A supramodal accumulation-to-bound signal that
determines perceptual decisions in humans. Nature Neuroscience. 2012;15: 1729–
1735. doi:10.1038/nn.3248

49.   Kelly SP, O'Connell RG. Internal and External Influences on the Rate of Sensory
Evidence Accumulation in the Human Brain. Journal of Neuroscience. 2013;33: 19434–
19441. doi:10.1523/JNEUROSCI.3355-13.2013

50.   Kösem A, Bosker HR, Takashima A, Meyer A, Jensen O, Hagoort P. Neural Entrainment
Determines the Words We Hear. Current Biology. 2018;28: 2867-2875.e3.
doi:10.1016/j.cub.2018.07.023

51.   Abrams DA, Nicol T, Zecker S, Kraus N. Abnormal Cortical Processing of the Syllable
Rate of Speech in Poor Readers. J Neurosci. 2009;29: 7686–7693.
doi:10.1523/JNEUROSCI.5242-08.2009

52.   Müller JA, Wendt D, Kollmeier B, Debener S, Brand T. Effect of Speech Rate on Neural
Tracking of Speech. Front Psychol. 2019;10. doi:10.3389/fpsyg.2019.00449

53.   Allman MJ, Meck WH. Pathophysiological distortions in time perception and timed
performance. Brain. 2012;135: 656–677. doi:10.1093/brain/awr210

54. Smith NJ, Kutas M. Regression-based estimation of ERP waveforms: II. Nonlinear
effects, overlap correction, and practical considerations. Psychophysiology. 2015;52:
169–181. doi:10.1111/psyp.12320

55. Brainard DH. The psychophysics toolbox. Spatial vision. 1997;10: 433–436.

56. Pelli DG. The VideoToolbox software for visual psychophysics: transforming numbers
into movies. Spatial Vision. 1997;10: 437–442. doi:10.1163/156856897X00366

57. Delorme A, Makeig S. EEGLAB: an open source toolbox for analysis of single-trial EEG
dynamics including independent component analysis. Journal of Neuroscience
Methods. 2004;134: 9–21. doi:10/bqr2f2

58. Lopez-Calderon J, Luck SJ. ERPLAB: an open-source toolbox for the analysis of event-
related potentials. Front Hum Neurosci. 2014;8. doi:10.3389/fnhum.2014.00213

59. Reichel L, Ye Q. Simple Square Smoothing Regularization Operators. Electronic
Transactions on Numerical Analysis. 2008;33: 63–83.

60. Maris E, Oostenveld R. Nonparametric statistical testing of EEG- and MEG-data. Journal
of Neuroscience Methods. 2007;164: 177–190. doi:10.1016/j.jneumeth.2007.03.024

61. Cumming G. The New Statistics: Why and How. Psychological Science. 2014;25: 7–29.
doi:10.1177/0956797613504966

62. Olejnik S, Algina J. Generalized eta and omega squared statistics: measures of effect
size for some common research designs. Psychol Methods. 2003;8: 434–447.
doi:10.1037/1082-989X.8.4.434