Myogenin controls via AKAP6 non-centrosomal microtubule

organizing center formation at the nuclear envelope

Robert Becker[1], Silvia Vergarajauregui[1], Florian Billing[1], Maria Sharkova[1],

Eleonora Lippolis[2], Kamel Mamchaoui[3], Fulvia Ferrazzi[2,4,5,6], Felix B. Engel[1,6,*]

[1]Experimental Renal and Cardiovascular Research, Department of Nephropathology,

Institute of Pathology, Friedrich-Alexander-Universität Erlangen-Nürnberg (FAU),

91054 Erlangen, Germany

[2]Institute of Human Genetics, Friedrich-Alexander-Universität Erlangen-Nürnberg,

91054 Erlangen, Germany

[3]Sorbonne Universités UPMC Université Paris 06, INSERM U974, CNRS FRE3617,

Center for Research in Myology, GH Pitié Salpêtrière, 47 Boulevard de l'Hôpital, 75013

Paris, France

[4]Department of Nephropathology, Institute of Pathology, Friedrich-Alexander-

Universität Erlangen-Nürnberg (FAU), 91054 Erlangen, Germany

[5]Institute of Pathology, Friedrich-Alexander-Universität Erlangen-Nürnberg (FAU),

91054 Erlangen, Germany

[6]Muscle Research Center Erlangen (MURCE), 91054 Erlangen, Germany

Short title: Myogenin controls ncMTOC formation

*Corresponding author:

Felix B. Engel, Universitätsklinikum Erlangen, Schwabachanlage 12 (TRC), 91054

Erlangen, Germany, felix.engel@uk-erlangen.de

Abstract

Non-centrosomal microtubule organizing centers (ncMTOC) are pivotal for the function of multiple cell types, but the processes initiating their formation are unknown. Here, we find that the transcription factor myogenin is required in myoblasts for recruiting centrosomal proteins. Moreover, myogenin is sufficient in fibroblasts for ncMTOC formation and centrosome attenuation. Bioinformatics combined with loss- and gain-of-function experiments identified induction of AKAP6 expression as one central mechanism for myogenin-mediated ncMTOC formation. Promoter studies indicate that myogenin preferentially induces the transcription of muscle- and ncMTOC-specific isoforms of *Akap6* and *Syne1*, which encodes nesprin-1α, the ncMTOC anchor protein in muscle cells. Overexpression of AKAP6β and nesprin-1α was sufficient to recruit endogenous centrosomal proteins to the nuclear envelope of myoblasts in the absence of myogenin. Taken together, our results illuminate how mammals transcriptionally control the switch from a centrosomal MTOC to an ncMTOC and identify AKAP6 as a novel ncMTOC component in muscle cells.

Keywords: myogenin; muscle differentiation; MTOC; non-centrosomal MTOC; AKAP6; centrosome; microtubules; terminal differentiation; nuclear envelope; transcription

Introduction

Correct organization of the microtubule cytoskeleton is essential for many cellular processes such as the establishment of cell shape, organelle positioning or intracellular transport (Akhmanova & Steinmetz, 2015; Conduit, Wainman, & Raff, 2015). In proliferating vertebrate cells, proteins that control microtubule nucleation and anchoring accumulate as pericentriolar material at the centrosome, which in turn functions as the dominant microtubule-organizing center (MTOC) (Prosser & Pelletier, 2015). The centrosomal MTOC is pivotal for cell cycle progression and correct chromosome segregation during mitosis (Hinchcliffe, Miller, Cham, Khodjakov, & Sluder, 2001; Khodjakov & Rieder, 2001; Sir et al., 2013). In contrast, MTOC function is assigned to non-centrosomal sites during differentiation of various cell types (Sanchez & Feldman, 2017). In epithelial cells, apically localized ncMTOCs participate in organelle positioning and help to establish apical-basal cell polarity (Brodu, Baffet, Le Droguen, Casanova, & Guichet, 2010; C. Lee, Scherr, & Wallingford, 2007; Meads & Schroer, 1995; Toya et al., 2016). In neurons, dendritic branch points, Golgi outposts, and pre-existing microtubules have been suggested to act as ncMTOC sites and precise control of microtubule array polarity helps to define the axonal and dendritic compartments (Luders, 2020; Nguyen et al., 2014; Ori-McKenney, Jan, & Jan, 2012; Sanchez-Huertas et al., 2016). In striated (i. e. heart and skeletal) muscle cells, ncMTOCs form at the nuclear envelope and contribute to correct nuclei positioning in skeletal myotubes (Elhanany-Tamir et al., 2012; Espigat-Georger, Dyachuk, Chemin, Emorine, & Merdes, 2016; Gimpel et al., 2017; Kronebusch & Singer, 1987; Tassin, Maro, & Bornens, 1985).

Despite progress in illuminating identity, composition, and function of ncMTOCs, it remains elusive which mechanisms initiate the switch from centrosomal to non-centrosomal MTOCs during differentiation of vertebrate cells. The only mechanistic

insight into ncMTOC induction has been gained by studying *Drosophila* tracheal cells. It was shown that the transcription factor trachealess, which specifies tracheal fate, is required for the spastin-mediated release of centrosomal components from the centrosome and their subsequent Piopio-mediated anchoring to the apical membrane (Brodu et al., 2010).

Here, we aimed to identify a transcription factor that controls ncMTOC formation in mammals utilizing skeletal muscle differentiation as an experimental system. Mammalian skeletal muscle differentiation is controlled by a family of transcription factors termed myogenic regulatory factors (MRFs) (Braun & Gautel, 2011; Buckingham & Rigby, 2014). Among those, myoblast determination protein (MyoD) regulates commitment to a myogenic fate and is thought to promote early differentiation of myoblasts (Comai & Tajbakhsh, 2014; Ishibashi, Perry, Asakura, & Rudnicki, 2005). The MRF myogenin acts as a unique regulator of terminal differentiation during myogenesis. In the absence of myogenin *in vivo*, embryonic myofiber formation is disturbed and the second wave of fetal myogenesis is largely abolished (Hasty et al., 1993; Nabeshima et al., 1993; Venuti, Morris, Vivian, Olson, & Klein, 1995). Notably, all MRFs are able to induce phenotypical markers of skeletal muscle in permissive non-muscle cells (Braun, Bober, Winter, Rosenthal, & Arnold, 1990; Braun, Buschhausen-Denker, Bober, Tannich, & Arnold, 1989; Comai & Tajbakhsh, 2014; Davis, Weintraub, & Lassar, 1987; Edmondson & Olson, 1989; Weintraub et al., 1989). Therefore, we examined whether MRFs regulate ncMTOC formation during skeletal muscle differentiation and whether they are sufficient for ncMTOC initiation in non-muscle cells.

Based on loss and gain of function experiments, quantitative analysis utilizing different cell types, and promoter studies, we show here that myogenin is required and sufficient for the formation of an ncMTOC at the nuclear envelope by controlling the

97   expression of muscle- and ncMTOC-specific isoforms of *Akap6* and *Syne1* which

98   encodes nesprin-1α.

Results

Myogenin is required for centrosomal protein recruitment to the nuclear envelope

To elucidate the regulation of ncMTOC establishment in skeletal muscle cells, we correlated the expression of MyoD and myogenin, transcription factors promoting early (MyoD) or late (myogenin) muscle differentiation, with the ncMTOC formation during mouse C2C12 myoblast differentiation. We analyzed two key steps during ncMTOC formation: i) Expression of nesprin-1α, the nuclear envelope anchor for centrosomal proteins (Espigat-Georger et al., 2016; Gimpel et al., 2017) and ii) recruitment of PCM-1, the first centrosomal protein to localize to the nuclear membrane (Srsen, Fant, Heald, Rabouille, & Merdes, 2009; Zebrowski et al., 2015). Immunofluorescence analyses of C2C12 cells one day after induction of differentiation revealed that 49.1 ± 7% of nuclei were positive for MyoD, 12 ± 0.9% were positive for myogenin, 7.8 ± 0.3% were positive for nesprin-1α, and 1.9 ± 0.3% were positive for PCM-1 (Figure 1A-C). Note, PCM-1 recruitment to the nuclear envelope is a gradual process (Figure 1 – figure supplement 1A). In our analysis intermediate stages were rare (<2% of total PCM-1-positive nuclei) and have therefore been included in the total percentage of PCM-1-positive nuclei.

All nuclei that were positive for PCM-1 were also positive for MyoD, myogenin, and nesprin-1α (Figure 1B,E). This suggests that MyoD and/or myogenin might be required for the recruitment of PCM-1 to the nuclear envelope. In contrast, only MyoD was present in all nesprin-1α-positive nuclei whereas only 42% of nesprin-1α-positive nuclei were also myogenin-positive (Figure 1A,D). Considering that nesprin-1α is anchoring centrosomal proteins at the nuclear envelope, this suggests that myogenin might be dispensable for ncMTOC formation.

To determine whether myogenin is dispensable for the establishment of the ncMTOC, we depleted myogenin in C2C12 cells via siRNA and analyzed nesprin-1α expression as well as nuclear envelope localization of the centrosomal proteins PCM-1, pericentrin, and AKAP9, which is essential for microtubule nucleation at the nuclear envelope, after two days of differentiation. The number of nesprin-1α-positive nuclei was not significantly affected by myogenin depletion (Figure 1E,F). By contrast, knockdown of myogenin reduced the number of PCM-1-positive nuclei from 11.6 ± 1.3% in control siRNA-treated cells to 2.3 ± 0.8% (Figure 1E,F). Similarly, pericentrin-positive nuclei were reduced from 6.9 ± 0.1% to 1.4 ± 0.1% and AKAP9-positive nuclei showed a reduction from 6.8 ± 0.5% to 1.4 ± 0.1% (Figure 1F and Figure 1 – figure supplement 1B). Contrary to our hypothesis, these data indicate that myogenin is required for centrosomal protein localization to the nuclear envelope during skeletal muscle differentiation.

Considering that PCM-1 is 1) important for recruitment of other proteins to the centrosome (Dammermann & Merdes, 2002; Prosser & Pelletier, 2020) and 2) the first centrosomal protein to localize to the nuclear envelope {Srsen, 2009 #412}{Zebrowski, 2015 #419}, we aimed to assess whether the loss of PCM-1affects the recruitment of other centrosomal proteins to the nuclear envelope. Depletion of PCM-1 reduced the percentage of C2C12 nuclei fully or partially positive for pericentrin after two days of differentiation from 5.8 ± 1% to 0.8 ± 0.2% and 2.7 ± 0.9% to 0.5 ± 0.1%, respectively (Figure 1G,H). In contrast, PCM-1 depletion did not affect recruitment of AKAP9 (Figure 1 – figure supplement 1C). Similar qualitative results have been described in myotubes (Espigat-Georger et al., 2016; Gimpel et al., 2017).

Collectively, our data suggest that the MyoD-driven early myogenic differentiation is sufficient to induce nesprin-1α expression whereas myogenin regulates the nuclear envelope targeting of AKAP9 and PCM-1, which in turn recruits pericentrin.

Figure 1. Myogenin is required for centrosomal protein recruitment to the nuclear envelope.
(A-B) C2C12 cells were differentiated for one day and immunostained for the myogenic regulatory factors (MRFs) MyoD or myogenin (Myog) and nesprin-1α (A) or PCM-1 (B). Orange asterisks: MRF+/PCM-1+ nuclei; yellow asterisks: MRF+/nesprin-1α+ nuclei; arrowheads: MRF-/nesprin-1α- nuclei; arrows: MRF-/nesprin-1α+ nuclei. Scale bars: 20 µm. **(C)** Quantification of (A) and (B). Data are represented as individual biological replicates, together with mean ± SD. **(D)** Quantification of MyoD and Myog in relation to nesprin-1α showing that not all nesprin-1α positive nuclei are myogenin positive. Data are represented as individual biological replicates, together with mean ± SD. **(E-F)** C2C12 myoblasts were transfected with negative control (si-ctrl) or myogenin siRNA (si-Myog) and differentiated for two days. Immunostaining (E) and subsequent quantification (F) shows that myogenin depletion affects nuclear recruitment of PCM-1, pericentrin (PCNT), and AKAP9 but not of nesprin-1α. 95% CI of differences si-Myog vs. si-ctrl = -3.11% to 0.98% (nesprin-1α+), -11.31% to -7.22% (PCM-1+), -7.53% to 3.43% (PCNT+), -7.52% to -3.42% (AKAP9+). Scale bars: 20 µm. **(G-H)** C2C12 myoblasts were transfected with si-ctrl or PCM-1 siRNA (si-PCM1) and differentiated for two days. PCNT was detected by immunostaining (G) and subsequent quantification (H) showed that PCM-1 depletion reduces PCNT+ nuclei. 95% CI of differences si-PCM1 vs. si-ctrl = -6.6% to -3.4% (full), -3.78% to -0.59% (partial). Scale bars: 20 µm. Data are represented as individual biological replicates, together with mean ± SD. ns: p > 0.05. *: p < 0.05 ***: p < 0.001. n = 3.

Ectopic myogenin expression is sufficient to induce a nuclear envelope MTOC

All MRFs are able to "transdifferentiate" permissive non-muscle cells with varying efficiency. They induce skeletal muscle markers such as the expression of contractile proteins or cell fusion into myotubes (Braun et al., 1990; Braun et al., 1989; Davis et al., 1987; Edmondson & Olson, 1989; Weintraub et al., 1989). However, ncMTOC formation has never been analyzed in "transdifferentiated" cells. To determine whether MRFs are sufficient to induce ncMTOC formation at the nuclear envelope in non-muscle cells, we ectopically expressed MyoD-GFP, myogenin-GFP or GFP alone in mouse NIH3T3 fibroblasts and analyzed the localization of PCM-1. NIH3T3 cells transfected with GFP exhibited centrosomal PCM-1 localization, typical for proliferating cells (Figure 2A). By contrast, expression of MyoD-GFP and myogenin-GFP induced nuclear envelope localization of PCM-1 in a subset of GFP-positive cells (Figure 2A,B), suggesting that both MRFs are sufficient individually to induce a switch from centrosomal MTOC to a nuclear envelope MTOC.

The results obtained upon MyoD-GFP expression appear surprising, as our C2C12 experiments suggested that myogenin is required for centrosomal protein recruitment. However, MyoD is known to activate myogenin transcription (Berkes & Tapscott, 2005). Analyzing endogenous myogenin expression in MyoD-GFP expressing cells, we observed high myogenin levels in cells that had recruited PCM-1 to the nuclear envelope compared to MyoD-GFP expressing cells where PCM-1 was absent from the nuclear envelope (Figure 2 – figure supplement 1A). In addition, depletion of myogenin in NIH3T3 cells abrogated the effect of MyoD-GFP expression on the recruitment of PCM-1 to the nuclear envelope (Figure 2 – figure supplement 1B). This further argues for myogenin to be required to recruit centrosomal proteins to the nuclear envelope.

In order to elucidate how myogenin induces nuclear envelope MTOC formation, we generated stable NIH3T3 cell lines that express either mScarlet or myogenin-2A-

mScarlet (MYOG-mScarlet) under control of a tetracycline-responsive promoter (Tet-ON). To induce myogenin expression and potentially centrosomal recruitment to the nuclear envelope, we treated MYOG-mScarlet cells with doxycycline (Dox) for three days. Immunofluorescence analysis revealed that Dox-treatment induced nuclear envelope localization of PCM-1, pericentrin, and AKAP9 (Figure 2C, 2E, and 2G). To better account for the dynamic of the recruitment process, we quantified the area of the nuclear envelope positive for each centrosomal protein in mScarlet and MYOG-mScarlet cells after Dox treatment (Figure 2D, 2F, and 2H). For this, we set an intensity threshold for the centrosomal protein signal and quantified the percentage of pixels above this threshold inside a 1 µm-wide band around the nucleus (identified by DAPI signal). The signal at the centrosome, which localizes in close proximity to the nucleus in most cells, accounts for the nuclear envelope coverage in mScarlet cells. Quantification revealed that median PCM-1 nuclear envelope coverage increases from 12.5% in mScarlet cells to 30.3% in MYOG-mScarlet cells (Figure 2D). Pericentrin was recruited with an increase of the median coverage from 8.2% in mScarlet cells to 23.1% in MYOG-mScarlet cells (Figure 2F). AKAP9, which is essential for microtubule nucleation at the nuclear envelope (Gimpel et al., 2017), coverage increased only moderately from 3.6% to 6.2%, suggesting that AKAP9 recruitment is less efficient in MYOG-mScarlet cells compared to PCM-1 and pericentrin (Figure 2H). However, ~13% of analyzed MYOG-mScarlet nuclei showed a higher coverage than the maximum observed in mScarlet-cells. Together, these data indicate that myogenin is sufficient to induce nuclear envelope recruitment of centrosomal proteins in non-muscle cells.

As nuclear envelope localization of centrosomal proteins requires nesprin-1α in C2C12 cells (Espigat-Georger et al., 2016; Gimpel et al., 2017), we examined whether myogenin is able to induce nesprin-1α expression in Dox-treated MYOG-mScarlet

cells. Dox-treatment resulted in nesprin-1-positive nuclei (Figure 2I) and RT-PCR

analysis confirmed that the α-isoform of nesprin-1 is upregulated upon myogenin

induction (Figure 2J). This indicate that while myogenin loss might be compensated by

another transcription factor in C2C12, myogenin can induce nesprin-1α expression.

Notably, chromatin immunoprecipitation (ChIP) sequencing data available through the

ENCODE consortium (Consortium et al., 2020; Yue et al., 2014) predict myogenin as

well as MyoD to bind a candidate regulatory element near the transcription start site of

nesprin-1α. To examine whether myogenin directly induces nesprin-1α transcription in

MYOG-mScarlet cells by binding to the α-isoform-specific promoter in the *Syne1* gene,

we performed ChIP, using an anti-myogenin antibody or an isotype control, followed

by PCR for the ENCODE-predicted site. PCR amplification was successful from

myogenin-precipitated DNA but not from IgG1 control (Figure 2K), indicating that

myogenin binds the nesprin-1α promoter and can induce expression of nesprin-1α in

permissive cells.

Finally, we examined if the myogenin-induced recruitment of centrosomal proteins

converts the nuclear envelope to a functional ncMTOC. For this we analyzed

microtubule regrowth after cold-induced depolymerization in Dox-treated mScarlet and

MYOG-mScarlet cells. In mScarlet-expressing cells, microtubule regrowth was

observed from the centrosome (Figure 2L). In contrast, MYOG-mScarlet cells exhibited

microtubule regrowth from the nuclear envelope to varying degrees. Quantification

revealed that median nuclear envelope coverage increased from 10.1% in mScarlet

cells to 18.4% in MYOG-mScarlet cells.

Collectively, these data demonstrate that myogenin is sufficient to induce ncMTOC

formation in NIH3T3 fibroblasts.

Figure 2. Myogenin expression is sufficient to induce ncMTOC formation in non-muscle cells.
(A) NIH3T3 fibroblasts were transfected with constructs encoding GFP, MyoD-GFP or myogenin-GFP (Myog-GFP). After three days, PCM-1 localization was assessed by immunostaining. Arrows indicate nuclei of transfected cells which have recruited PCM-1. Scale bars: 10 μm. **(B)** Quantification of (A) demonstrating that myogenin induces nuclear recruitment of PCM-1 more efficiently than MyoD. Data are represented as individual biological replicates, together with mean ± SD. ***: p < 0.001; 95% CI of difference Myog-GFP vs. MyoD-GFP = 30.99% to 49.22%. n = 3. **(C-H)** NIH3T3 Tet-ON mScarlet or MYOG-2A-mScarlet (MYOG-mScarlet) cells were treated with doxycycline (Dox) for three days. After immunostaining, nuclear envelope recruitment of PCM-1 (C-D), pericentrin (Pcnt) (E-F), and AKAP9 (G-H) was analyzed and quantified. Data are depicted as violin plots. Red line indicates the median, dotted lines indicate the 25% and 75% percentile. ***: p < 0.001. Scale bars = 20 μm **(I)** Immunostaining of MYOG-mScarlet cells treated with Dox for three days showing the presence of nesprin-1α-positive nuclei. Scale bars = 20 μm **(J)** RT-PCR analysis of MYOG-mScarlet cells in the absence of Dox (-Dox) or treated with Dox for the indicated time points demonstrating that *Syne1α* is upregulated upon myogenin expression. *Gapdh* was used as equal input control. **(K)** ChIP-PCR analysis of Dox-treated MYOG-mScarlet cells using an anti-myogenin antibody or an IgG1 control showing that myogenin binds an E-box in the *Syne1α* promoter region. **(L-M)** Immunostaining of α-tubulin and subsequent quantification of nuclear envelope coverage after 30s of microtubule regrowth following cold-induced microtubule depolymerization in mScarlet or MYOG-mScarlet cells treated with Dox for three days. Data are depicted as violin plots. Red line indicates the median, dotted lines indicate the 25% and 75% percentile. ***: p < 0.001. Scale bars = 20 μm.

Myogenin expression attenuates the centrosomal MTOC

In different cell types, it has been observed that ncMTOC formation is associated with attenuation of the centrosomal MTOC (Leask, Obrietan, & Stearns, 1997; Nguyen, Stone, & Rolls, 2011; Yang & Feldman, 2015; Zebrowski et al., 2015). Therefore, we examined if myogenin expression induces centrosome attenuation in MYOG-mScarlet fibroblasts. Dox stimulation resulted in a significant reduction of pericentrin levels at centrioles in MYOG-mScarlet cells when compared to Dox-treated mScarlet cells (Figure 3A,B). In contrast, levels of Cep135, a centriole-associated protein, which does not relocalize to the nuclear envelope in muscle cells, did not change significantly upon myogenin induction (Figure 3C,D) indicating that myogenin affects centrosomal localization of PCM proteins but not of centriole-associated proteins. To test if myogenin attenuates MTOC activity at the centrosome, we analyzed centrosomal levels of the microtubule nucleating factor γ-tubulin. Induced MYOG-mScarlet cells displayed a significant reduction in centrosomal γ-tubulin levels compared to mScarlet control cells (Figure 3E). Analyzing microtubule regrowth, we observed that centrosomes still nucleated microtubules in MYOG-mScarlet cells that exhibited microtubule nucleation at the nuclear envelope (Figure 3F). However, α-tubulin signal at centrioles was less intense in MYOG-mScarlet cells compared to mScarlet cells, indicating a reduced MTOC activity (Figure 3F). Taken together, these data indicate that myogenin expression – in parallel to inducing ncMTOC formation – attenuates the centrosomal MTOC and that centrosomal and nuclear envelope MTOC can be active at the same time.

Figure 3. Myogenin expression attenuates the centrosomal MTOC
(A-E) mScarlet or MYOG-mScarlet cells were stimulated with Dox for three days and pericentrin (Pcnt) (A), Cep135 (C), and γ-tubulin (A, C) were detected by immunostaining. Quantification shows that pericentrin (B) and γ-tubulin (E) intensities at the centrosome are reduced upon myogenin induction while Cep135 intensity (D) is not significantly affected. Single channel images of Pcnt, γ-tubulin, and Cep135 are false-colored to visualize different intensities. Data are shown as violin plots. The red line indicates the median and dotted lines indicate the 25% and 75% percentile. ns: p > 0.05. ***: p < 0.001. Scale bars = 5 μm **(F)** Immunostaining of α-tubulin and γ-tubulin in Dox-stimulated mScarlet or MYOG-mScarlet cells after 30s of microtubule regrowth. Intensity-based color-coding of α-tubulin shows that microtubule growth from centrioles is reduced after myogenin induction. Scale bars = 5 μm.

AKAP6 is a potential mediator of myogenin-induced ncMTOC formation

The myogenin depletion experiments suggested that the sole presence of nesprin-1α at the nuclear envelope does not allow efficient recruitment of MTOC components during muscle differentiation (Figure 1F). Thus, myogenin potentially contributes to the recruitment process by controlling the expression of proteins that are necessary for 1) inhibiting the localization of centrosomal proteins to the centrosome, 2) targeting centrosomal proteins to the nucleus, and/or 3) anchoring centrosomal proteins to the nuclear envelope via nesprin-1α.

In order to identify candidates that act downstream of myogenin and mediate ncMTOC formation, we integrated published myogenin ChIP-Seq data (Yue et al., 2014) with RNA-Seq data of C2C12 differentiation (Doynova, Markworth, Cameron-Smith, Vickers, & O'Sullivan, 2017) (Figure 4A). Myogenin ChIP-Seq data was obtained at four different time points (myoblasts, 24 h differentiation, 60 h differentiation, and 7 days differentiation; see Methods section for details) whereas the RNA-Seq data-set contained three time points (myoblasts as well as differentiating C2C12 cells at 3 days and 7 days of differentiation). PCM-1 nuclear envelope localization can already be observed 24 h after induction of differentiation (Figure 1A), but the number of cells which differentiate and form an ncMTOC increases over time. Additionally, we assumed genes required for the maintenance of the ncMTOC in differentiated cells have to be actively transcribed. Therefore, we considered only those myogenin binding sites in our analysis which were detected in the ChIP-Seq data-set at 24 h, 60 h, and 7 days of differentiation. The promoters of 2462 genes were bound by myogenin at these three time points (Figure 4B). We then intersected these 2462 genes with a list of 3800 genes, which were upregulated in the RNA-Seq data-set at both 3 days and 7 days of differentiation when compared to proliferating myoblasts (Figure 4B). This intersection yielded a list of 748 potential direct myogenin

target genes (Figure 4B). Considering that skeletal muscle cells and cardiomyocytes

(myogenin-negative) both express nesprin-1α and exhibit a nuclear envelope MTOC,

we hypothesized that ncMTOC formation in both cell types is controlled by similar

mechanisms. Thus, we assessed whether any of the 748 target genes are upregulated

during rat heart development from embryonic day 15 to postnatal day 3, the

developmental window in which ncMTOCs form in cardiomyocytes (Zebrowski et al.,

2015). For this purpose, we utilized a microarray-derived temporal expression data-set

spanning rat heart development (Patra et al., 2011). This strategy helped to further

reduce the number of candidate genes to 107 myogenin targets that potentially

mediate ncMTOC formation (Supplementary file 1). As our previous data suggested

that nesprin-1α alone does not allow efficient recruitment of centrosomal proteins to

the nuclear envelope, we first focused on candidates that potentially cooperate with

nesprin-1α in anchoring centrosomal proteins to the nuclear envelope. To this end, we

utilized Gene Ontology analysis to identify candidates that are annotated to localize at

the nuclear envelope (Figure 4A). Four genes matched the Gene Ontology cellular

component search terms "nuclear membrane" and "nuclear envelope": *Akap6*, *Dmpk*,

Rb1cc1, and *Tmem38a*. Previous studies indicated that myogenin directly binds and

activates the promoter of *Akap6* (S. W. Lee et al., 2015), which encodes the large

scaffold A-kinase anchoring protein (AKAP) 6 (also known as mAKAP). AKAP6 has

been described to localize to the nuclear envelope of cardiomyocytes through

interaction with the N-terminal spectrin domains of nesprin-1α and to act as a signaling

hub by assembling signaling proteins such as protein kinase A, ryanodine receptor,

phosphodiesterase 4D3 and phospholipase C (Kapiloff, Schillace, Westphal, & Scott,

1999; Pare, Easlick, Mislow, McNally, & Kapiloff, 2005; Passariello, Li, Dodge-Kafka,

& Kapiloff, 2015; Ruehr et al., 2003). Furthermore, proximity labeling indicated that

AKAP6 is an interactor of nesprin-1α in C2C12 myotubes (Gimpel et al., 2017). Taken

together, these data identify AKAP6 as a potential mediator of myogenin-induced ncMTOC formation.

AKAP6 occurs in two isoforms: The brain-specific α-isoform and the β-isoform, which is predominantly expressed in heart and skeletal muscle (Michel et al., 2005). We first determined if myogenin binds the β-isoform promoter of *Akap6* in MYOG-mScarlet fibroblasts and if AKAP6β expression is induced in these cells upon Dox treatment. We could amplify an E-Box containing region of the β-isoform promoter after ChIP using an anti-myogenin antibody (Figure 4C). RT-PCR analysis showed that AKAP6β is upregulated after Dox stimulation (Figure 4D). This indicates that, in fibroblasts, myogenin can bind the *Akap6*β-isoform promoter and induce AKAP6 expression.

To determine whether nesprin-1α is – similar to the situation in cardiomyocytes – involved in AKAP6β localization to the nuclear envelope of skeletal muscle cells (Pare et al., 2005), we analyzed the expression pattern of AKAP6 (refers to the β isoform if not specified) and nesprin-1α in C2C12 cells. Immunofluorescence analysis at two days differentiation showed that all AKAP6-positive nuclei were nesprin-1α-positive (Figure 4E). In addition, high-resolution microscopy suggested that AKAP6 mainly localizes at the cytoplasmic side of nesprin-1α (Figure 4F and Figure 4 – figure supplement 1). It has been reported that the C-terminus of nesprin-1α is inserted into the outer nuclear membrane, whereas the N-terminus extends into the cytoplasm (Apel, Lewis, Grady, & Sanes, 2000; Q. Zhang et al., 2001).

To test if nesprin-1α is required to anchor AKAP6 to the nuclear envelope, patient-derived myoblasts carrying a mutation in the *SYNE1* gene (23560 G>T causing a premature stop and loss nesprin-1α expression) and myoblasts of healthy donors were differentiated into myotubes and AKAP6 localization was compared (Figure 4G). Whereas nesprin-1α and AKAP6 localized to the nuclear envelope of myotubes from

381   healthy donors, expression of nesprin-1α and nuclear membrane localization of AKAP6

382   were abolished in myotubes carrying the *SYNE1* mutation. Taken together, these data

383   indicate that AKAP6 localizes to the nuclear envelope in differentiated skeletal muscle

384   cells via interaction with nesprin-1α.

385
Figure 4. The nesprin-1α interaction partner AKAP6 is a potential mediator of myogenin-induced
ncMTOC formation.
(A) Scheme illustrating the bioinformatics workflow used to identify potential myogenin downstream
candidates. **(B)** Venn diagram depicting the numbers of genes matching criteria for the individual data
sets and for intersection of data sets. Criteria for Myogenin ChIP-Seq data (red): Genes where myogenin
binding was detected at the promoter region; criteria for C2C12 RNA-Seq data (green) and for
microarray data of rat heart development (blue): up-regulated genes. **(C)** ChIP-PCR analysis of Dox-
treated MYOG-mScarlet cells using an anti-myogenin antibody or an IgG1 control showing that
myogenin binds an E-box in the *Akap6* β-promoter region. **(D)** RT-PCR analysis of MYOG-mScarlet
cells in the absence of Dox (-Dox) or treated with Dox for the indicated time points demonstrating that
Akap6β is upregulated upon myogenin expression. The two bands for *Akap6β* derive from alternative
splicing of the first exon of *Akap6β* which results in a ~200 bp insertion in the 5' untranslated region.
Gapdh was used as equal input control. **(E)** C2C12 cells were differentiated for two days and
immunostaining shows that all AKAP6+ nuclei are also nesprin-1α+. Scale bar: 20 μm. **(F)** High resolution
Airyscan image of (E). Arrowhead indicates AKAP6 localized at the cytoplasmic side of nesprin-1α
signal. Arrow marks nesprin-1α that is localized at the nuclear side of AKAP6 signal. Scale bar: 0.5 μm.
(G) Myoblasts from healthy donors (wt) and from patients carrying a mutation in the *SYNE1* gene
(SYNE1−/−) were differentiated for four days. Immunostaining analysis showed that loss of nesprin-1α is
associated with loss of AKAP6 from the nuclear envelope in differentiated myotubes (troponin I). Scale
bars: 10 μm.

AKAP6 is required for ncMTOC formation and maintenance

To examine the role of AKAP6 in ncMTOC formation we performed siRNA-mediated depletion experiments in differentiating C2C12 cells and MYOG-mScarlet fibroblasts. In C2C12 cultures differentiated for two days, 12.3 ± 0.6% of nuclei were positive for AKAP6 and 10.4 ± 1.2% of nuclei were positive for PCM-1 (Figure 5A,B). Importantly, AKAP6 was found at all nuclei that had recruited PCM-1. Transfection of differentiating C2C12 cultures with AKAP6 siRNA significantly reduced the number of AKAP6-positive nuclei from 12.3 ± 0.6% to 4.8 ± 0.1% and the number of PCM-1-positive nuclei from 10.4 ± 1.2% to 3.4 ± 0.3% (Figure 5B) but had no effect on nesprin-1α localization (Figure 5 - figure supplement 1). Correspondingly, treatment of Dox-induced MYOG-mScarlet with AKAP6 siRNA decreased median nuclear envelope coverage by PCM-1 and pericentrin from 22.6% to 9% and 18.7% to 5.5%, respectively (Figure 5C-E). Median coverage of AKAP9 was only moderately affected (7.2% to 6.5%) but nuclei showing more than ~18% AKAP9 coverage were completely lost after AKAP6 depletion (Figure 5F). Next, we examined if AKAP6 promotes nuclear envelope recruitment by directly interacting with centrosomal proteins. For this, we performed co-immunoprecipitation experiments using an anti-AKAP6 antibody. We could co-precipitate PCM-1 from MYOG-mScarlet lysates but not from mScarlet lysate (Figure 5G). These data indicate that AKAP6 is required for the localization of centrosomal proteins to the nuclear envelope, in part by interacting with PCM-1.

To determine if AKAP6 is required for maintaining MTOC protein localization at the nuclear envelope, we transfected C2C12 cultures enriched for myotubes with AKAP6 siRNA (Figure 5G,H). Depletion of AKAP6 resulted in the loss of PCM-1, AKAP9, and pericentrin from the nuclear envelope in myotubes.

Taken together, our data demonstrate that AKAP6 is required for recruiting centrosomal proteins to the nuclear envelope as well as maintaining nuclear envelope

432   localization of centrosomal proteins in myotubes, most likely by acting as an adaptor

433   between centrosomal proteins and the nuclear membrane anchor nesprin-1α.

434
Figure 5. AKAP6 is required for nuclear envelope localization of centrosomal proteins.
(A) C2C12 cells were differentiated for two days. Immunostaining shows that all PCM-1+ nuclei are also
AKAP6+. Scale bar: 20 μm. **(B)** Quantification of AKAP6+ and PCM-1+ nuclei in C2C12 cells treated with
negative control (si-ctrl) or Akap6 (si-Akap6) siRNA after two days of differentiation indicate that
AKAP6 is required for nuclear PCM-1 localization. Data are represented as individual biological
replicates, together with mean ± SD. **: p = 0.0012, 95% CI = 6.21% to 8.74%; ***: p < 0.0001, 95%
CI = 4.63% to 9.43%. n = 3. **(C)** MYOG-mScarlet cells were treated with si-ctrl or si-Akap6 and
subsequently treated with Dox for three days. Image analysis revealed that myogenin-induced
recruitment of PCM-1 to the nuclear envelope is AKAP6-dependent. Scale bars: 10 μm. **(D)**
Quantification of (C). Data are shown as violin plots. The red line indicates the median and dotted lines
indicate the 25% and 75% percentile. ***: p < 0.001. **(E-F)** Quantification of pericentrin (E) and AKAP9
(F) nuclear coverage in Dox-stimulated MYOG-mScarlet cells treated with si-ctrl or si-Akap6. Data are
shown as violin plots. The red line indicates the median and dotted lines indicate the 25% and 75%
percentile. *: p < 0.05. ***: p < 0.001. **(G)** Co-Immunoprecipitation (IP) of PCM-1 from MYOG-mScarlet
but not from mScarlet lysate (L) using an anti-AKAP6 antibody. **(H-I)** Enriched C2C12 myotubes
(troponin I) were transfected with negative control (si-ctrl) or Akap6 (siAkap6) siRNA and
immunostaining demonstrates that AKAP6 is required for maintaining nuclear envelope localization of
PCM-1 (H) as well as pericentrin and AKAP9 (I). Scale bars (H): 20 μm, (I): 10 μm.

AKAP6 is required for ncMTOC function

The nuclear envelope MTOC has been described to be required for correct positioning and distribution of nuclei in multinucleated myotubes via two different mechanisms: 1) PCM-1 enables the recruitment of the dynein regulator p150glued and other motor proteins to the nuclear envelope and promotes alignment of nuclei (Espigat-Georger et al., 2016), and 2) the Akap9-dependent nucleation of microtubules from the nuclear envelope contributes to the spreading of nuclei throughout the cell body (Gimpel et al., 2017) (Figure 6A). Our results indicate a potential role for AKAP6 in both aspects of nucleus positioning as it is required for PCM-1 and AKAP9 to localize to the nuclear envelope. Analyzing the positioning and distribution of nuclei in enriched C2C12 myotubes two days after siRNA-mediated depletion of AKAP6, we found that AKAP6 depletion reduced the number of myotubes with aligned nuclei (66.0 ± 7.5% to 34.7 ± 7.6%) and increased the number of myotubes with overlapping nuclei (26.0 ± 6.5% to 58.3 ± 9.6%), compared to control myotubes (Figure 6B,C). This indicates that AKAP6 is required for proper alignment and spreading of nuclei in myotubes.

Next, we aimed to confirm that the observed nuclei mispositioning in AKAP6-depleted cells is due to aberrant microtubule nucleation and motor protein recruitment at the nuclear envelope. Immunofluorescence analysis showed that two days post siRNA transfection the microtubule network organization was similar in AKAP6-depleted and control myotubes showing the typical organization of microtubules in longitudinal arrays (Figure 6 – figure supplement 1A). Similar results have been obtained previously when depleting nesprin-1α (Espigat-Georger et al., 2016). Yet, we observed that AKAP6 depletion resulted in a reduced intensity of detyrosinated (i.e. stable) microtubules compared to control myotubes (Figure 6 – figure supplement 1B). To test if AKAP6 depletion affects the nucleation of new microtubules, we assessed microtubule regrowth after nocodazole-induced depolymerization in C2C12 myotubes.

In myotubes treated with control siRNA, microtubules regrew from the nuclear

envelope and to a lesser extent from cytoplasmic loci (Figure 6D). In AKAP6-depleted

myotubes, microtubule regrowth from the nuclear envelope was abolished (Figure 6D).

This suggests that AKA6 depletion impairs spreading of myonuclei by preventing

microtubule growth from the nuclear envelope. Next, we analyzed the localization of

p150glued (also known as DCTN1). In control siRNA-treated myotubes, p150glued

localized at the nuclear envelope (Figure 6E). AKAP6 depletion resulted in a reduction

of p150glued at the nuclear envelope of C2C12 myotubes (Figure 6E) suggesting that

the reduced number of myotubes with aligned nuclei is due to impaired dynein

activation. Collectively, these data demonstrate that AKAP6 is required for the function

of the nuclear envelope MTOC in skeletal muscle cells.

491
492
Figure 6. AKAP6 is required for correct nuclear positioning in myotubes.
(A) Scheme illustrating the role of the ncMTOC in myonuclear positioning and the potential impact of
AKAP6 depletion. **(B)** Enriched C2C12 myotubes (troponin I) were transfected with negative control
(si-ctrl) or Akap6 (si-Akap6) siRNA. The upper si-Akap6 panel shows a representative image of a
myotube with misaligned nuclei, the lower si-Akap6 panel shows nuclei overlapping inside a myotube.
Scale bars: 20 μm. **(C)** Quantification of (B). Data are represented as individual biological replicates,
together with mean ± SD. *: p < 0.05, 95% CI of difference si-Akap6 vs. si-ctrl = 10.42% to 52.25% (left
graph); 95% CI = 9.65% to 55.02% (right graph). n = 3. **(D)** Enriched C2C12 myotubes (troponin I) were
transfected with si-ctrl or si-Akap6 and subsequently subjected to a nocodazole-based microtubule
(α-tubulin) regrowth assay. Image analysis showed that AKAP6 depletion abrogated microtubule
nucleation at the nuclear envelope. Scale bars: 10 μm. **(E)** Enriched C2C12 myotubes (troponin I)
transfected with si-ctrl or si-Akap6 were immunostained for the dynein regulator p150glued. Image
analysis showed that AKAP6 depletion reduces p150glued signal at the nuclear envelope. Scale bars:
5 μm.

Myogenin-induced isoforms of nesprin-1 and AKAP6 are sufficient for centrosomal protein recruitment

To examine if myogenin specifically induces expression of isoforms that are associated with the nuclear envelope MTOC in skeletal muscle, we performed ChIP on MYOG-mScarlet cell lysate using an anti-myogenin antibody and assessed the abundance of isoform-specific promoter regions of *Syne1* and *Akap6* in the precipitated DNA. For this, we performed semi-quantitative PCR (qPCR) using primer pairs targeting myogenin consensus binding sites (i.e. E-boxes) in regions predicted by ENCODE data (Consortium et al., 2020) to be associated with myogenin binding (Figure 7A,C). We found that the amount of template corresponding to the promoter region upstream of the nesprin-1α transcript (nesprin-1α2) was 4.5-fold higher than the promoter region of the long nesprin-1 isoform (nesprin-1-giant) (Figure 7B). Similarly, the promoter region upstream of the AKAP6β transcript was 3-fold enriched compared to the promoter region of the AKAP6α transcript (Figure 7D). This indicates that myogenin preferentially binds promoter regions of *Syne1* and *Akap6* isoforms that are involved in ncMTOC formation.

To test if the preferential binding is associated with an increased activation of transcription of specific isoforms, we constructed vectors with putative promoter regions of AKAP6α, AKAP6β, nesprin-1-giant, or nesprin-1α located directly upstream of a luciferase coding sequence and transfected them into human HEK293T cells together with GFP or myogenin-GFP. Co-transfection of myogenin with the AKAP6β promoter construct increased luciferase activity 10.9-fold, while co-transfection with the AKAP6α promoter construct did not show a significant increase compared to GFP-transfected control (Figure 7E). Similarly, we observed a 21.7-fold increase in activity when myogenin was co-transfected with the nesprin-1α promoter construct but only a mild 2.7-fold increase after co-transfection with the promoter construct of nesprin-1-

giant (Figure 7F). These results indicate that myogenin preferentially induces

transcription of nesprin-1α and AKAP6β.

Finally, we examined whether the myogenin-induced isoforms of nesprin-1 and

AKAP6 are sufficient to recruit centrosomal proteins in the absence of myogenin. For

this, we expressed nesprin-1α-mCherry alone or together with AKAP6β-GFP in

undifferentiated, myogenin-negative myoblasts. In nesprin-1α-mCherry-transfected

cells, PCM-1 did not localize to the nuclear envelope (Figure 7G). In contrast, co-

transfection of nesprin-1α-mCherry and AKAP6β-GFP was sufficient to recruit PCM-1

to the nuclear envelope (Figure 7G).

Taken together, our results demonstrate that myogenin specifically induces

transcription of isoforms which are 1) required for the ncMTOC in differentiated skeletal

muscle cells and 2) sufficient to recruit centrosomal proteins to the nuclear envelope

in cells with a centrosomal MTOC.

546
547

Figure 7. Myogenin preferentially induces MTOC-associated isoforms of *Syne1* and *Akap6*.
(A, C) Schematic representation of the murine *Syne1* (A) and *Akap6* (C) gene and derived transcripts.
Exons are indicated by grey rectangles and the first exon of each transcript is marked by color. E-boxes
(myogenin consensus sites) inside putative promoters are indicated as yellow boxes and small black
arrows mark the primers used for qPCR. **(B, D)** Myogenin ChIP from Dox-stimulated MYOG-mScarlet
cells followed by qPCR for the indicated E-boxes shows that myogenin preferentially binds the promoter
regions upstream of *nesprin-1α2* and *Akap6β* transcripts. Data are represented as individual biological
replicates, together with mean ± SD. ***: $p < 0.001$. **(E-F)** Luciferase assay testing the activity of the
indicated *Akap6* (E) or *Syne1* (F) promoters in the presence of GFP or myogenin-GFP (MYOG-GFP).
Data are represented as individual biological replicates, together with mean ± SD. ns: $p > 0.05$.
: $p < 0.01$. ***: $p < 0.001$. **(G) Overexpression of nesprin-1α-mCherry alone or together with AKAP6β-
GFP in undifferentiated (myogenin-negative) C2C12 myoblasts. Co-expression of nesprin-1α and
AKAP6β is sufficient for nuclear envelope recruitment of endogenous PCM-1. Scale bars: 20 µm.

Discussion

We conclude that the myogenic transcription factor myogenin controls ncMTOC formation at the nuclear envelope and that myogenin-induced AKAP6β expression is one of the central molecular components required for ncMTOC formation (Figure 8). This conclusion is supported by our findings that 1) myogenin is required to recruit the MTOC proteins PCM-1, pericentrin and AKAP9 to the nuclear envelope in differentiating muscle cells, 2) ectopic myogenin expression is sufficient to promote the formation of an ncMTOC at the nuclear envelope of fibroblasts, and 3) the myogenin-induced isoforms AKAP6β and nesprin-1α are required and sufficient for the recruitment of centrosomal proteins to the nuclear envelope.

Figure 8. Schematic overview of the role of myogenin and AKAP6 in ncMTOC formation.
Myogenin induces expression of AKAP6β which connects centrosomal proteins like pericentrin (Pcnt), AKAP9, and PCM-1 to the nuclear membrane protein nesprin-1α, whose transcription can be induced by myogenin as well. Depletion, overexpression, and co-immunoprecipitation experiments suggest that AKAP6β directly acts as an adapter between centrosomal proteins and nesprin-1α. Yet, other proteins might be involved and the here presented protein complex at the nuclear envelope is hypothetical. At the same time, myogenin is sufficient to attenuate centrosomal MTOC function. AKAP6-dependent anchoring of centrosomal proteins as well as microtubule nucleation from the nuclear envelope are required for correct positioning of nuclei inside differentiating myotubes.

Formation of ncMTOCs has been associated with cellular differentiation (Sanchez
& Feldman, 2017), but a direct regulation of ncMTOC formation by particular
differentiation pathways in vertebrate cells has remained elusive. Our results
demonstrate that myogenin, which is an essential regulator of terminal differentiation,
drives ncMTOC formation in mammalian cells. This shows that terminal differentiation
factors can control in vertebrates the switch of dominant MTOC localization from the
centrosome to non-centrosomal sites. Notably, MTOC formation at the nuclear
envelope occurs also in cells that lack myogenin or cell-type specific transcriptional
master regulators of terminal differentiation, such as cardiomyocytes or osteoclasts
(Kronebusch & Singer, 1987; Mulari et al., 2003; Zebrowski et al., 2015). A recent study
from our lab demonstrated that AKAP6β orchestrates the assembly of the ncMTOC in
cardiomyocytes and osteoclasts (Vergarajauregui et al., 2020), validating our findings
in skeletal muscle. Therefore, it would be important in future studies to identify
transcription factors that bind to AKAP6β and nesprin-1α promoters and regulate the
switch from centrosomal to ncMTOC in these cell types. For this purpose, promising
cardiomyocyte and osteoclast "transdifferentiation" tools using multiple transcription
factors are available (Chang, Cho, Kim, & Kim, 2019; Ieda et al., 2010; Klose, Gossen,
& Stamm, 2019; Yamamoto et al., 2015). Additionally, it appears important to
determine in future experiments the mechanisms underlying the preferential binding of
myogenin to the isoform-specific promoters, considering the abundance of myogenin
binding sites (E-boxes) throughout the genome. Notably, isoform upregulation or
switching might be a general mechanism that contributes to MTOC regulation during
differentiation. This assumption is supported by the recent identification of a spermatid-
specific isoform of centrosomin, the *Drosophila* orthologue of CDK5RAP2, which can
induce ncMTOC formation at mitochondria (Chen, Buchwalter, Kao, & Megraw, 2017).
Moreover, a non-centrosomal isoform of ninein contributes to neuronal differentiation

(X. Zhang et al., 2016) and a shorter isoform of pericentrin is upregulated in

differentiating cardiomyocytes (Zebrowski et al., 2015).

While site-specific anchor proteins for ncMTOCs, such as nesprin-1α, have been

identified (Espigat-Georger et al., 2016; Gimpel et al., 2017; Lechler & Fuchs, 2007;

Meng, Mushika, Ichii, & Takeichi, 2008), it remained unclear how centrosomal proteins

are connected to these site-specific anchors. Previously, it has been reported that

overexpression of nesprin-1α in undifferentiated myoblasts is sufficient to recruit an

ectopically expressed centrosomal targeting domain of pericentrin and AKAP9 (i.e. the

PACT domain) as well as minor amounts of endogenous PCM-1 to the nuclear

envelope in a subset of transfected cells (Espigat-Georger et al., 2016; Gimpel et al.,

2017). Here, we show that myogenin induced the expression of the large scaffold

protein AKAP6, which we prove to be essential for nuclear ncMTOC formation and

maintenance, most likely by connecting centrosomal proteins to nesprin-1α. Myogenin

preferentially binds and activates the putative promoters of AKAP6β and nesprin-1α

isoforms, which are known to be upregulated in differentiated muscle cells (Kapiloff et

al., 1999; Michel et al., 2005; Randles et al., 2010). Importantly, ectopic co-expression

of AKAP6β and nesprin-1α, but not nesprin-1α alone, was sufficient to recruit

endogenous centrosomal proteins in the absence of myogenin.

MTOC activity plays a pivotal role for cellular function and altered MTOC function

or mutations in genes encoding MTOC proteins are associated with disease such as

Emery-Dreifuss muscular dystrophy (*SYNE1*) (Q. Zhang et al., 2007) and

Microcephalic osteodysplastic primordial dwarfism, type II (*PCNT*) (Rauch et al., 2008)

as well as possibly Long QT syndrome 11 (*AKAP9*) (de Villiers et al., 2014) and Seckel

syndrome (*NIN*) (Dauber et al., 2012). In addition, amplified and/or hyperactive

centrosomes act as oncogene-like factors (Arnandis et al., 2018; Godinho & Pellman,

2014; Godinho et al., 2014; Levine et al., 2017; LoMastro & Holland, 2019). Therefore,

it is important to better understand mechanisms that control MTOC activity. Notably,

ectopic expression of myogenin in fibroblasts did not only induce ncMTOC formation

but also attenuated the centrosomal MTOC. In addition, myogenin expression in

fibroblast was only inducing in a subset of cells an ncMTOC. Thus, our cellular systems

combined with our bioinformatics approach provides new opportunities to tackle future

key questions of MTOC formation such as: What factors increase or decrease

efficiency of ncMTOC induction? What post-transcriptional processes contribute to

ncMTOC establishment? How are induction of ncMTOC and attenuation of

centrosomal MTOC linked?

In summary, our findings suggest that key differentiation factors can control the

switch from centrosomal MTOC to ncMTOC and cell-type specific adaptor proteins are

required to connect centrosomal proteins to anchor proteins at non-centrosomal sites.

Conclusively, our study 1) contributes to a better understanding of the striated muscle

ncMTOC, 2) presents a mechanistic framework that may be applicable to ncMTOC

formation in other cell types and tissues, and 3) provides a cellular system to elucidate

further molecular mechanisms inducing the switch from centrosomal to non-

centrosomal MTOCs.

Material and Methods

Further information and requests for resources and reagents should be directed to and

will be fulfilled by the lead contact, Felix Engel (felix.engel@uk-erlangen.de).

Cell lines, differentiation and doxycycline stimulation

The following cell lines have been used in the present study:

Cell Lines	Source	Catalogue number/ Identifier
Mouse: C2C12 myoblast cell line	ATCC	Cat# CRL-1772, RRID: CVCL_0188
Mouse: NIH3T3 fibroblast cell line	ATCC	Cat# CRL-1658, RRID:CVCL_0594
Human: HEK293T cell line	ATCC	Cat# PTA-4488, RRID:CVCL_0045
Human: myoblasts healthy donor	(Holt et al., 2016; Mamchaoui et al., 2011)	N/A
Human: myoblasts carrying SYNE1 mutation	(Holt et al., 2016; Mamchaoui et al., 2011)	N/A

Reagents used for cell culture are listed in the table below. All cells used in this

study were cultured at 37°C in a humidified atmosphere containing 5% CO_2. Growth

medium for C2C12, NIH3T3, Hela and Hct116 consisted of high glucose DMEM

supplemented with GlutaMAX containing 10% FBS, 1 mM sodium pyruvate, 100 U/ml

penicillin and 100 µg/ml streptomycin. C2C12 cells were maintained at 50% confluence

to preserve differentiation capacity. For differentiation, cells were cultured to 90%

confluence and subsequently changed to differentiation medium (high glucose DMEM

with GlutaMAX containing 0.5% FBS and Insulin, Transferrin, Selenium, Sodium

Pyruvate Solution (1:1000 of 100x ITS-A).

Human myoblasts from a healthy control or from a congenital muscular dystrophy

patient carrying a homozygous nonsense mutation within the *SYNE1* gene (nucleotide

23560 G>T) were immortalized by Kamel Mamchaoui and Vincent Mouly (Center for

Research in Myology, Paris, France) as previously described via transduction with

retrovirus vectors expressing hTERT and Cdk4 (Holt et al., 2016; Mamchaoui et al.,

2011). Growth medium consisted of DMEM supplemented with GlutaMAX and DMEM

199 in a 4:1 ratio containing 20% FBS, 25 µg/ml bovine fetuin, 5 ng/ml recombinant

human EGF, 0.5 mg/ml recombinant human FGF-basic, 5 µg/ml recombinant insulin,

0.2 µg/ml dexamethasone, and 50 µg/ml gentamicin (Gimpel et al., 2017). To induce

differentiation, immortalized myoblasts were grown to ~90% confluence and then

changed to Skeletal Muscle Differentiation Medium (PromoCell) containing 50 µg/ml

gentamicin. For immunofluorescence analysis of immortalized myoblasts, glass

coverslips were coated with Matrigel diluted 1:100 in DMEM.

Cell culture reagent	Source	Catalogue number/ Identifier
0.25% Trypsin/EDTA solution	Sigma Aldrich	Cat# T4049
Dulbecco's Modified Eagle Medium (DMEM), high glucose, GlutaMAX	Thermo Fisher Scientific	Cat# 61965059
DMEM 199 medium	Thermo Fisher Scientific	Cat# 41150
Skeletal Muscle Differentiation Medium	PromoCell	Cat# C-23061
Fetal Bovine Serum (FBS)	biowest	Cat# S1810
Fetal Bovine Serum (FBS), tetracycline free	PAN-Biotech	Cat# P30-3602
Doxycycline hydrochloride	Sigma-Aldrich	Cat# D3447
Horse Serum	Thermo Fisher Scientific	Cat# 16050122
Sodium pyruvate solution	Sigma-Aldrich	Cat# S8636
Penicillin-Streptomycin	Thermo Fisher Scientific	Cat# 15140122
Gentamicin	Thermo Fisher Scientific	Cat# 15750060

Insulin-Transferrin-Selenium-Sodium Pyruvate (ITS-A) (100X)	Thermo Fisher Scientific	Cat# 51300044
Bovine fetuin	Thermo Fisher Scientific	Cat# 10344026
EGF Recombinant Human Protein	Thermo Fisher Scientific	Cat# PHG0311
FGF-Basic (AA 10-155) Recombinant Human Protein	Thermo Fisher Scientific	Cat# PHG0026
Insulin, Human Recombinant	Sigma-Aldrich	Cat# 91077C
Dexamethasone	Sigma-Aldrich	Cat# D4902
Matrigel® hESC-Qualified Matrix, LDEV-free	Corning	Cat# 356277
Fibronectin bovine plasma	Sigma-Aldrich	Cat# F1141

Myotube enrichment

C2C12 cells were differentiated in 6-well plates or 10 cm dishes for 4-5 days as described above. To preferentially detach myotubes, cells were washed two times with PBS and treated with pre-warmed 0.0125% Trypsin/EDTA solution (0.25% Trypsin/EDTA stock diluted in PBS) for ~2 min at room temperature. Detachment of myotubes was constantly monitored by phase contrast microscopy. After sufficient myotube detachment was observed, Trypsin/EDTA solution was carefully aspirated and a myotube-enriched suspension was collected by rinsing the plates five times with normal growth medium. Enriched myotubes were then plated on glass coverslips coated with 25 µg/ml fibronectin in PBS for >45 min at 37°C. After 24 h incubation at 37°C myotube cultures were subjected to siRNA treatment and/or microtubule regrowth assays.

MRF plasmids construction

Primers used to generate fragments are listed in the table below. *Myod* and *Myog* cDNA were obtained by reverse transcription PCR using RNA from C2C12 cells differentiated for 2 days. The cDNAs were then cloned into the peGFP-N1 backbone

by Cold Fusion Cloning (System Biosciences, Cat# MC010B-1) following the manufacturer's instruction. Positive clones were identified by restriction digest and Sanger sequencing.

Oligonucleotides	Sequence
Primer: mouse MyoD cold fusion cloning forward	gggatccaccggtcgccaccatggagcttctatcgccgcc
Primer: mouse MyoD cold fusion cloning reverse	tcctcgcccttgctcaccataagcacctgataaatcgcat
Primer: mouse myogenin cold fusion cloning forward	gggatccaccggtcgccaccatggagctgtatgagacatc
Primer: mouse myogenin cold fusion cloning reverse	tcctcgcccttgctcaccatgttgggcatggtttcgtctg

Luciferase plasmids construction

Candidate regulatory elements associated with myogenin binding in *Syne1* and *Akap6* genes were identified from ENCODE data accessed through the SCREEN web interface (https://screen.wenglab.org/). Potential promoter regions were amplified from genomic DNA obtained from NIH3T3 cells using the primers listed in the table below. After amplification, promoter fragments were cloned in front of the *Renilla* luciferase ORF (*hRluc*) into the psiCHECK-2 vector (Promega, Cat# C8021; GenBank Accession Number AY535007) using NEBuilder HiFi DNA Assembly Master Mix (New England Biolabs, Cat# E2621L) according to the manufacturer's instructions.

Oligonucleotides	Sequence
Primer: psiCHECK™-2 assembly cloning forward	gctagccaccatggcttc
Primer: psiCHECK™-2 assembly cloning reverse	gtcgagccatgtgagcaaaag
Primer: AKAP6α promoter fragment assembly cloning forward	ttttgctcacatggctcgacgcaaaccatgattattatctaagtc
Primer: AKAP6α promoter fragment assembly cloning forward	tggaagccatggtggctagcaggagatgccttgtttgac
Primer: AKAP6β promoter fragment assembly cloning forward	ttttgctcacatggctcgactacagagaattaatgcttagaaac
Primer: AKAP6β promoter fragment assembly cloning reverse	tggaagccatggtggctagcgcttgcagtgaactgtaatg

Primer: Syne1-giant promoter fragment assembly cloning forward	ttttgctcacatggctcgacactctgtctctttcaacccaaacatc atgaacc
Primer: Syne1-giant promoter fragment assembly cloning reverse	tggaagccatggtggctagcggctgtgccctccctccc
Primer: Syne1-α promoter fragment assembly cloning forward	ttttgctcacatggctcgacgtgtcagattctgactcattg
Primer: Syne1-α promoter fragment assembly cloning reverse	tggaagccatggtggctagctggtcatgcttctaaataaaac

Plasmid transfections

Plasmid transfection into NIH3T3 cells was carried out with 500 ng DNA per well of a

24-well plate using 1 µl Lipofectamine LTX (Thermo Fisher Scientific, Cat# 15338100)

according to the manufacturer's instructions. Transfection complexes were formed by

incubating DNA with Lipofectamine LTX in Opti-MEM for 20 min at room temperature.

For transfection of luciferase constructs in HEK293T cells, 250 ng luciferase vector

and 250 ng myogenin-eGFP or eGFP control plasmid were used per well of a 24-well

plate. Transfection complexes were assembled by incubating DNA with PEI MAX

(Polysciences, Cat# 24765-1) in a 1:3 ratio in Opti-MEM for 20 min at room

temperature.

siRNA transfections

Cells were transfected using 2 µl Lipofectamine RNAiMAX reagent (Thermo Fisher

Scientific, Cat# 13778150) and 40 nM final siRNA concentration per well of a 24-well

plate. Transfection complexes were formed by incubating siRNA with RNAiMAX in

Opti-MEM for 20 min at room temperature. C2C12 cells were transfected 24 h prior to

induction of differentiation (~50% confluence) and enriched C2C12 myotubes were

transfected 24 h after re-plating. NIH3T3 were transfected with siRNA 48 h after

plasmid transfection. The following siRNAs have been used in this study:

Oligonucleotides and sequences	Source	Catalogue number
siRNA targeting sequence: mouse Myogenin AAUAAAGACUGGUUGCUAUCAAAAA	Integrated DNA technologies	Cat# mm.Ri.Myog.13.1
siRNA targeting sequence: mouse Akap6 GGACUACAUCAAGAACGAATT	Thermo Fischer Scientific	Cat# 4390771 s108732
Negative control (si-ctrl) siRNA	Integrated DNA technologies	Cat# 51-01-14-04

Luciferase assay

Luciferase activity was measured in Centro XS^3 LB 960 96-well plate reader

luminometer (BertholdTech, #50-6860) using the Dual-Luciferase® Reporter Assay

System (Promega, #E1910) according to the manufacturer's instructions. In brief,

HEK293T cells were harvested 48h after transfection in passive lysis buffer and stored

at -80°C until measurement. Activities of firefly luciferase (*hluc+*, internal control) and

Renilla luciferase (promoter activation) were measured sequentially for each sample.

Values of *Renilla* luciferase activity were normalized to those of firefly luciferase for

each measurement.

Production of lentiviral vectors

The lentiviral packaging plasmid psPAX2 and the VSV-G envelope plasmid pMD2.G

were gifts from Didier Trono (psPAX2: Addgene plasmid #12260;

http://n2t.net/addgene:12260; RRID: Addgene_12260; pMD2.G: Addgene plasmid

#12259; http://n2t.net/addgene:12259; RRID: Addgene_12259). The transfer plasmids

pLenti CMVtight Blast DEST (w762-1) and pLenti CMV rtTA3 Hygro (w785-1) used for

creating tetracycline-inducible cell lines were gifts from Eric Campeau (w762-1:

Addgene plasmid #26434; http://n2t.net/addgene:26434; RRID: Addgene_26434;

w785-1: Addgene plasmid #26730; http://n2t.net/addgene:26730; RRID:

Addgene_26730). The coding sequences of human myogenin (gift from Matthew

Alexander & Louis Kunkel (Addgene plasmid #78341 ; http://n2t.net/addgene:78341;

RRID: Addgene_78341) and mScarlet (Bindels et al., 2017) (gift from Dorus Gadella

(Addgene plasmid #85042; http://n2t.net/addgene:85042; RRID: Addgene_85042)

were cloned into w762-1 by NEBuilder® HiFi DNA Assembly according to the

manufacturer's instructions using the primers indicated in the table below.

To produce lentiviral vectors, psPAX2, pMD2.G, and the desired transfer plasmid

were transfected in a 1:1:2 ratio into HEK293T cells using PEI MAX. Supernatant

containing lentiviral vectors was harvested 72h after transfection, filtered through a

0.45 µm filter and aliquots were snap frozen.

Oligonucleotides	Sequence
Primer: human myogenin cloning into pLenti CMVtight Blast DEST forward	agccgccaccATGGAGCTGTATGAGACATC
Primer: human myogenin cloning into pLenti CMVtight Blast DEST reverse	ctgccctctccactgcccatGTTGGGCATGGTTTCATC
Primer: mScarlet cloning into pLenti CMVtight Blast DEST forward	tcctggcccaATGGTGAGCAAGGGCGAG
Primer: mScarlet cloning into pLenti CMVtight Blast DEST reverse	ggccgccactgtgctggataTCACTTGTACAGCTCGTCCATGC

Generation of stable cell lines

Lentiviral vector aliquots were rapidly thawed at 37°C and diluted in tetracycline-free

growth medium containing 10 µg/ml Polybrene (Sigma-Aldrich, Cat# 107689). Cells

were transduced overnight in 6-well plates using 1 ml of diluted lentiviral vector. The

following morning, medium was refreshed and cells were selected for transgene

integration 72h after transduction. We first generated cells expressing a reverse

tetracycline activator (rtTA3), which were subsequently transduced with lentiviral

vectors carrying the desired transgene to express mScarlet or MYOG-2A-mScarlet

under control of a tetracycline responsive element (TRE). After selection, Tet-ON cell lines were used for 20 passages.

Immunoprecipitation

Cells were harvested in lysis buffer containing 1% NP-40, 150 mM NaCl, 10% glycerol, 5 mM EDTA, 20 mM Tris-HCl (pH 7) and EDTA-free protease inhibitor cocktail (cOmplete™, Roche # 11873580001). After 30 min incubation on ice, samples were sonicated and lysates were cleared by centrifugation at 16000xg for 16 min at 4°C. For immunoprecipitation, 0.5 µg anti-AKAP6 antibody/mg of total protein were added to the lysate and incubated overnight at 4°C on a rotor. Subsequently, antibody complexes were purified by incubation of lysate with Protein A Sepharose® beads (Merck, GE17-5138-01) for 3h rotating at 4°C. Beads were washed three times with cold lysis buffer for 5 min and proteins were eluted from beads by incubation in 2x NuPAGE LDS sample buffer () at 95°C for 5 min. Lysates and immunoprecipitated samples were analyzed by SDS-PAGE (4-12% NuPAGE Novex Bis-Tris gels) under reducing conditions and transferred to a nitrocellulose membrane by wet transfer at 350 mA and <60 V for 1.5 h in 1x transfer buffer (25 mM Tris-HCl, pH 7.5, 192 mM glycine, 0.1% SDS, 10% methanol). The membrane was then blocked with 5% BSA in TBS-T (1×TBS, 0.05% Tween-20) and incubated with the following antibodies:

Antibodies and dilutions	Source	Catalogue number/ Identifier
rabbit polyclonal anti-PCM1 (H-262) dilution: 1:500	Santa Cruz	Cat# sc-67204, RRID:AB_2139591
rabbit polyclonal anti-Akap6 dilution: 1:2000	Sigma-Aldrich	Cat# HPA048741, RRID:AB_2680506
sheep polyclonal anti-rabbit IgG HRP-linked	GE Healthcare	NA934-1ML, RRID:AB_2722659

ChIP-qPCR

Protein and DNA were cross-linked by fixing cells for 10 min at 37°C using 1% formaldehyde in culture medium. Cross-linking was quenched by adding 125 mM glycine and gently agitating the cells for 5 min at room temperature. Then, cells were harvested in ice-cold PBS, centrifuged at 1000xG for 5 min at 4°C, and the resulting pellet was lysed in ChIP lysis buffer (1% SDS, 10 mM EDTA, 50 mM Tris-HCl pH 8.1) for 30 min on ice. After lysis, samples were sonicated for 30 cycles consisting of 20s sonication and 30s pause inside an ice bath. Sonicated samples were centrifuged at 4°C for 30 min at 10000xG. A small aliquot of the supernatant was saved as input control and the remaining supernatant was subjected to immunoprecipitation. Samples were diluted 1:5 in RIPA buffer (10x RIPA Millipore, # 20-188; 1:10 dilution in ddH_2O) and incubated overnight at 4°C with 1 µg/ml myogenin antibody (Santa Cruz Biotechnology, #sc-12732 X) or IgG1 isotype control (Thermo Fisher, # 16471482). Protein G agarose beads (Roche #11719416001) pre-blocked with salmon sperm (Thermo Fisher, # 15632011) were used to precipitate antibody complexes from diluted samples. Beads were sequentially washed at 4°C with low salt buffer (0.1% SDS, 1% Triton X-100, 2 mM EDTA, 20 mM Tris-HCl pH 8.0, 150 mM NaCl), high salt buffer (0.1% SDS, 1% Triton X-100, 2 mM EDTA, 20 mM Tris-HCl pH 8.0, 500 mM NaCl), LiCl buffer (0.25 M LiCl, 1% NP-40, 1% Sodium Deoxycholate, 1 mM EDTA, 10 mM Tris-HCl pH 8.0), and TE buffer (10 mM Tris-HCl pH 8.0, 1 mM EDTA). Antibody complexes were eluted by incubating beads for 15 min at 30°C in elution buffer (1% SDS, 100 mM $NaHCO_3$). Eluates were digested with proteinase K and RNAse A and DNA fragments were purified using a PCR purification kit (Macherey-Nagel, #740609).

Immunofluorescence and microscopy

Antibodies used in this study are listed in the table below. Note that the MANNES1E antibody detects different nesprin-1 isoforms (Holt et al., 2016; Randles et al., 2010). However, previous studies have shown that only the nesprin-1α isoform is up-regulated during muscle differentiation (Espigat-Georger et al., 2016; Gimpel et al., 2017; Holt et al., 2016). Prior to fixation, cells were rinsed once with PBS. Cells were fixed either with pre-chilled methanol at -20°C for 3 min or with 4% formaldehyde/PBS for 10 min at room temperature. Formaldehyde-fixed cells were permeabilized with 0.5% TritonX-100/PBS. Prior to antibody staining, samples were blocked for at least 20 min using 5% BSA in 0.2% Tween-20 in PBS. Primary antibodies were diluted in blocking reagent and incubated with the sample for 90 min at room temperature or overnight at 4°C. After removal of primary antibody solution and three 5 min washes with 0.1% NP40/PBS, samples were incubated for 60 min with fluorophore-coupled secondary antibodies. DNA was visualized with 0.5 µg/ml DAPI (4′,6′-diamidino-2-phenylindole) in 0.1% NP40/PBS. After DAPI staining cover slips were rinsed once with Millipore-filtered water and then mounted using Fluoromount-G mounting medium. Analysis, image acquisition and high-resolution microscopy were done using a LSM800 confocal laser scanning microscope equipped with an Airyscan detector and the ZEISS Blue software (Carl Zeiss AG, RRID: SCR_013672) with Airyscan image processing.

Antibodies and dilutions	Source	Catalogue number/ Identifier
rabbit polyclonal anti-PCM1 (H-262) dilution: 1:200	Santa Cruz	Cat# sc-67204, RRID:AB_2139591
mouse monoclonal anti-PCM1 (G-6) dilution: 1:200	Santa Cruz	Cat# sc-398365
rabbit polyclonal anti-Akap6 dilution: 1:1000	Sigma-Aldrich	Cat# HPA048741, RRID:AB_2680506
mouse monoclonal anti-nesprin-1 (MANNES1E) dilution: 1:50	G. Morris (Randles et al., 2010)	N/A

mouse monoclonal anti-Myogenin (F5D) dilution 1:500	Santa Cruz	Cat# sc-12732, RRID:AB_627980
mouse monoclonal anti-MyoD1 dilution: 1:500	Millipore	Cat# MAB3878, RRID:AB_2251119
rat anti-α-tubulin dilution: 1:500	Sigma-Aldrich	Cat# T9026, RRID:AB_477593
goat polyclonal anti-Troponin I dilution: 1:500	Abcam	Cat# ab56357, RRID:AB_880622
mouse monoclonal anti-γ-tubulin dilution: 1:100	Santa Cruz	Cat# sc-51715, RRID:AB_630410
rabbit polyclonal anti-Akap9 dilution: 1:200	Sigma-Aldrich	Cat# HPA026109, RRID:AB_1844688
rabbit polyclonal anti-Pericentrin dilution: 1:1000	BioLegend	Cat# PRB-432C, RRID:AB_291635
Donkey anti-Rabbit IgG (H+L) Highly Cross-Adsorbed Secondary Antibody, Alexa Fluor 488 dilution: 1:500	Thermo Fisher Scientific	Cat# A21206
Donkey anti-Rabbit IgG (H+L) Highly Cross-Adsorbed Secondary Antibody, Alexa Fluor 594 dilution: 1:500	Thermo Fisher Scientific	Cat# A21207
Donkey anti-Rabbit IgG (H+L) Highly Cross-Adsorbed Secondary Antibody, Alexa Fluor 647 dilution: 1:500	Thermo Fisher Scientific	Cat# A31573
Donkey anti-Mouse IgG (H+L) Highly Cross-Adsorbed Secondary Antibody, Alexa Fluor 488 dilution: 1:500	Thermo Fisher Scientific	Cat# A21202
Donkey anti-Mouse IgG (H+L) Highly Cross-Adsorbed Secondary Antibody, Alexa Fluor 594 dilution: 1:500	Thermo Fisher Scientific	Cat# A21203
Donkey anti-Mouse IgG (H+L) Highly Cross-Adsorbed Secondary Antibody, Alexa Fluor 647 dilution: 1:500	Thermo Fisher Scientific	Cat# A31571

Donkey anti-Goat IgG (H+L) Highly Cross-Adsorbed Secondary Antibody, Alexa Fluor 647 dilution: 1:500	Thermo Fisher Scientific	Cat# A21447
Donkey anti-Rat IgG (H+L) Highly Cross-Adsorbed Secondary Antibody, Alexa Fluor 594 dilution: 1:500	Thermo Fisher Scientific	Cat# A21209

Image analysis

All image analyses were carried out using Fiji (http://fiji.sc, RRID: SCR_002285). For

quantification of nuclear envelope coverage, confocal images were transformed into

binary images by setting a manual intensity threshold. Regions of interests (ROI) were

obtained by detecting nuclei outline via DAPI staining and subsequent transformation

of these outlines into 1 µm-wide bands. Coverage was quantified as the percentage of

positive pixels inside bands in the binary images. ROIs for measuring intensities at

centrosomes were generated by detecting local signal maxima in γ-tubulin channels

and subsequent generation of circular ROIs with 1 µm diameter using the maxima as

centers.

To measure nuclear envelope intensity profiles for nesprin-1α and AKAP6, we first

created ROIs by manually detected nuclear outlines using DAPI staining. We then

decreased the diameter of these ROIs by 1 µm and used the newly created ROIS as

starting points for linear intensity profiles perpendicular to the nuclear outlines.

Microtubule regrowth assay

Cells were treated with 5 µM nocodazole (Sigma-Aldrich, Cat# M1404) in culture media

for 3 h at 37°C to depolymerize microtubules. To observe microtubule regrowth,

nocodazole-containing medium was removed, cells were rinsed three times with cold

medium and either fixed (0 min time point) with 4% formaldehyde in PBS for 10 min or

immediately transferred to 37°C pre-warmed culture media for the desired length of

time followed by formaldehyde fixation. Myotubes were extracted with 1% Triton X-100

in PHEM buffer (60 mM PIPES, 25 mM HEPES, 10 mM EGTA, 2 mM MgCl2, pH 6.9)

for 30 s at room temperature prior to fixation.

Bioinformatics analysis

Data was analyzed with R (http://www.r-project.org/; RRID:SCR_001905) and

Bioconductor (http://www.bioconductor.org/, RRID: SCR_006442). Myogenin ChIP-

Seq data (GEO accession number: GSE36024) produced within the ENCODE project

(Consortium, 2012; Consortium et al., 2020; Yue et al., 2014) were obtained via the

UCSC Genome Browser at https://genome.ucsc.edu/index.html (Kent et al., 2002;

Rosenbloom et al., 2013). NarrowPeak tracks relative to ChIP-Seq data from four

different time points were considered: undifferentiated C2C12 myoblasts as well as

C2C12 cultures differentiated for 24 h, 60 h, or 7 days. To identify myogenin promoter

binding, peaks were annotated to the Ensembl release 67 mouse genome relying on

Bioconductor packages biomaRt v. 2.30.0 (Durinck, Spellman, Birney, & Huber, 2009)

and ChIPpeakAnno v. 3.8.9 (Zhu et al., 2010). Genes were considered as myogenin

targets if a peak (p-value < 10^{-5}) was localized at a maximum distance of 1 kb from the

annotated transcriptional start site.

Results of differential expression analysis for RNA-Seq data from C2C12

differentiation (GEO accession number: GSE84158) were obtained from the Gene

Expression Omnibus (GEO) repository (Doynova et al., 2017). Three sample types

were analyzed: C2C12 myoblasts (C1), C2C12 cultures differentiated for 3 days

containing myoblasts as well as myotubes (C2), and C2C12 cultures differentiated for

7 days and treated with AraC resulting in depletion of proliferating myoblasts (C3).

Genes were considered as up-regulated if they exhibited a positive fold change

(p-value < 0.05) from C1 to C2 as well as from C1 to C3.

Gene expression microarray data for rat heart development were obtained as

described previously using the Affymetrix GeneChip RAT 230 Expression Set (Patra

et al., 2011). Genes were considered up-regulated if: 1) the corresponding probe set

was identified as differentially expressed on the basis of a procedure that accounts for

the total area under the profile compared to a constant profile (Di Camillo, Toffolo, Nair,

Greenlund, & Cobelli, 2007), and 2) the difference between the maximum expression

value over time and the initial one was greater than the difference between the initial

value and the minimum value.

For Gene Ontology analysis, annotated cellular component terms for each of the

potential   myogenin   targets   were   retrieved   using   the   search   tool   at

http://geneontology.org/ (Ashburner et al., 2000; The Gene Ontology, 2017). Potential

targets annotated with the terms "nuclear membrane" and "nuclear envelope" were

considered.

QUANTIFICATION AND STATISTICAL ANALYSIS

Quantification of nesprin-1α positive nuclei

Low levels of nesprin-1 expression can be detected at nuclei of non-differentiated

muscle cells using MANNES1E antibody. Prior to scoring of nesprin-1α positive nuclei,

we therefore set a threshold for nesprin-1 signal in images of differentiated C2C12 cells

by measuring and subtracting maximal nesprin-1 signal intensity in undifferentiated

C2C12 cultures.

Statistical analysis

As preliminary experiments indicated a large effect size of siRNA treatments and

ectopic MRF expression, three biological replicates were performed per experiment

(i.e. n = 3). Biological replicate means that cells were freshly plated, treated, fixed and

stained, and then analyzed. For each biological replicate, two technical replicates were performed in the sense that two individual wells were processed at the same time. When analyzed, the two technical replicates were scored as one sample.

For quantification in C2C12 cells, >500 nuclei were analyzed per condition and biological replicate. For MRF-GFP experiments, >50 GFP-positive cells were analyzed per condition and biological replicate. For nuclear coverage and intensity quantifications of mScarlet and MYOG-mScarlet cells, >100 nuclei or centrosomes were analyzed per condition and pooled from three biological replicates to display distribution in violin plots.

Statistical analysis was carried out using GraphPad Prism 5.02 or Prism 8.2.1 (La Jolla, USA). Differences between groups were considered statistically significant when the p-value ≤ 0.05. The 95% confidence interval (CI) for the differences between compared groups are reported in the figure legends. Statistical significance of differences between groups was tested using the following:

Figures 1F, 1H, S1B, S1C, 5B, 7E: One-way ANOVA followed by Bonferroni's Post-Hoc test to compare selected pairs of groups.

Figures 2B, 6C, 7B, 7D, 7F: Student's t-test together with an F-test to assess equality of variances.

Figures 2D, 2F, 2H, 2M, 3B, 3D, 3E, 5D, 5E, 5F: Kolmogorov-Smirnov test to compare the cumulative distribution of groups.

Author Contributions

Conceptualization: R.B., S.V., and F.B.E.

Methodology: R.B., S.V., E.L., F.F., and F.B.E.

Formal Analysis: E.L., F.F.

Investigation: R.B., S.V., F.B., and M.S.

Resources: K.M.

Writing – Original Draft: R.B. and F.B.E.

Writing – Reviewing & Editing: R.B., S.V., F.F., and F.B.E.

Visualization: R.B. and F.B.E.

Supervision: F.F. and F.B.E.

Funding Acquisition: F.F. and F.B.E.

Acknowledgments

We acknowledge the Platform for Immortalization of Human Cells at the Institute de Myologie, Paris for the generation and distribution of immortalized human myoblasts. We thank Glenn E. Morris for providing us with nesprin-1 antibody (MANNES1E). We thank the ENCODE consortium, the laboratory of Barbara Wold and the Millard and Muriel Jacobs Genetics and Genomics Laboratory at the California Institute of Technology for providing myogenin ChIP-Seq data. We thank Christina Warnecke for support with ChIP experiments, Marc Stemmler, Eva Bauer, and Thomas Brabletz for their help with luciferase assays and Anna K. Großkopf and Alexander S. Hahn for experimental support. We acknowledge Les Laboratoires Servier for providing illustrations in the Servier Medical Art collection under a Creative Commons Attribution 3.0 Unported License (https://creativecommons.org/ licenses/by/3.0/). We thank Anna K. Großkopf, Manfred Frasch, Hanh Nguyen, Thomas U. Mayer, Rosa M. Puertollano,

Payel Das, Marina Leone, Gentian Musa, and Salvador Cazorla-Vazquez for critical

reading of the manuscript and all members of the Engel lab for critical discussions.

This work was supported by the Interdisciplinary Centre for Clinical Research

Erlangen (IZKF project J42 to F.F.), the Emerging Fields Initiative Cell "Cycle in

Disease and Regeneration" (CYDER to F.B.E.) and an ELAN Program Grant (ELAN-

16-01-04-1-Vergarajauregui to S.V.) from the Friedrich-Alexander-Universität

Erlangen-Nürnberg, by the German Research Foundation (DFG, INST 410/91-1

FUGG and EN 453/12-1 to F.B.E.), and by the Research Foundation Medicine at the

University Clinic Erlangen, Germany.

Declaration of Interests

The authors have no conflict of interest to declare.

References

Akhmanova, A., & Steinmetz, M. O. (2015). Control of microtubule organization and dynamics: two ends in the limelight. *Nat Rev Mol Cell Biol, 16*(12), 711-726. doi:10.1038/nrm4084

Apel, E. D., Lewis, R. M., Grady, R. M., & Sanes, J. R. (2000). Syne-1, a dystrophin- and Klarsicht-related protein associated with synaptic nuclei at the neuromuscular junction. *J Biol Chem, 275*(41), 31986-31995. doi:10.1074/jbc.M004775200

Arnandis, T., Monteiro, P., Adams, S. D., Bridgeman, V. L., Rajeeve, V., Gadaleta, E., . . . Godinho, S. A. (2018). Oxidative Stress in Cells with Extra Centrosomes Drives Non-Cell-Autonomous Invasion. *Dev Cell, 47*(4), 409-424 e409. doi:10.1016/j.devcel.2018.10.026

Ashburner, M., Ball, C. A., Blake, J. A., Botstein, D., Butler, H., Cherry, J. M., . . . Sherlock, G. (2000). Gene ontology: tool for the unification of biology. The Gene Ontology Consortium. *Nat Genet, 25*(1), 25-29. doi:10.1038/75556

Berkes, C. A., & Tapscott, S. J. (2005). MyoD and the transcriptional control of myogenesis. *Semin Cell Dev Biol, 16*(4-5), 585-595. doi:10.1016/j.semcdb.2005.07.006

Bindels, D. S., Haarbosch, L., van Weeren, L., Postma, M., Wiese, K. E., Mastop, M., . . . Gadella, T. W., Jr. (2017). mScarlet: a bright monomeric red fluorescent protein for cellular imaging. *Nat Methods, 14*(1), 53-56. doi:10.1038/nmeth.4074

Braun, T., Bober, E., Winter, B., Rosenthal, N., & Arnold, H. H. (1990). Myf-6, a new member of the human gene family of myogenic determination factors: evidence for a gene cluster on chromosome 12. *EMBO J, 9*(3), 821-831. Retrieved from https://www.ncbi.nlm.nih.gov/pubmed/2311584

Braun, T., Buschhausen-Denker, G., Bober, E., Tannich, E., & Arnold, H. H. (1989). A novel human muscle factor related to but distinct from MyoD1 induces myogenic conversion in 10T1/2 fibroblasts. *EMBO J, 8*(3), 701-709. Retrieved from https://www.ncbi.nlm.nih.gov/pubmed/2721498

Braun, T., & Gautel, M. (2011). Transcriptional mechanisms regulating skeletal muscle differentiation, growth and homeostasis. *Nat Rev Mol Cell Biol, 12*(6), 349-361. doi:10.1038/nrm3118

Brodu, V., Baffet, A. D., Le Droguen, P. M., Casanova, J., & Guichet, A. (2010). A developmentally regulated two-step process generates a noncentrosomal microtubule network in Drosophila tracheal cells. *Dev Cell, 18*(5), 790-801. doi:10.1016/j.devcel.2010.03.015

Buckingham, M., & Rigby, P. W. (2014). Gene regulatory networks and transcriptional mechanisms that control myogenesis. *Dev Cell, 28*(3), 225-238. doi:10.1016/j.devcel.2013.12.020

Chang, Y., Cho, B., Kim, S., & Kim, J. (2019). Direct conversion of fibroblasts to osteoblasts as a novel strategy for bone regeneration in elderly individuals. *Exp Mol Med, 51*(5), 1-8. doi:10.1038/s12276-019-0251-1

Chen, J. V., Buchwalter, R. A., Kao, L. R., & Megraw, T. L. (2017). A Splice Variant of Centrosomin Converts Mitochondria to Microtubule-Organizing Centers. *Curr Biol, 27*(13), 1928-1940 e1926. doi:10.1016/j.cub.2017.05.090

Comai, G., & Tajbakhsh, S. (2014). Molecular and cellular regulation of skeletal myogenesis.
Curr Top Dev Biol, 110, 1-73. doi:10.1016/B978-0-12-405943-6.00001-4

Conduit, P. T., Wainman, A., & Raff, J. W. (2015). Centrosome function and assembly in animal
cells. *Nat Rev Mol Cell Biol, 16*(10), 611-624. doi:10.1038/nrm4062

Consortium, E. P. (2012). An integrated encyclopedia of DNA elements in the human genome.
Nature, 489(7414), 57-74. doi:10.1038/nature11247

Consortium, E. P., Moore, J. E., Purcaro, M. J., Pratt, H. E., Epstein, C. B., Shoresh, N., . . .
Weng, Z. (2020). Expanded encyclopaedias of DNA elements in the human and mouse
genomes. *Nature, 583*(7818), 699-710. doi:10.1038/s41586-020-2493-4

Dammermann, A., & Merdes, A. (2002). Assembly of centrosomal proteins and microtubule
organization depends on PCM-1. *J Cell Biol, 159*(2), 255-266.
doi:10.1083/jcb.200204023

Dauber, A., Lafranchi, S. H., Maliga, Z., Lui, J. C., Moon, J. E., McDeed, C., . . . Hwa, V. (2012).
Novel microcephalic primordial dwarfism disorder associated with variants in the
centrosomal protein ninein. *J Clin Endocrinol Metab, 97*(11), E2140-2151.
doi:10.1210/jc.2012-2150

Davis, R. L., Weintraub, H., & Lassar, A. B. (1987). Expression of a single transfected cDNA
converts fibroblasts to myoblasts. *Cell, 51*(6), 987-1000. doi:10.1016/0092-
8674(87)90585-x

de Villiers, C. P., van der Merwe, L., Crotti, L., Goosen, A., George, A. L., Jr., Schwartz, P. J.,
. . . Corfield, V. A. (2014). AKAP9 is a genetic modifier of congenital long-QT syndrome
type 1. *Circ Cardiovasc Genet, 7*(5), 599-606.
doi:10.1161/CIRCGENETICS.113.000580

Di Camillo, B., Toffolo, G., Nair, S. K., Greenlund, L. J., & Cobelli, C. (2007). Significance
analysis of microarray transcript levels in time series experiments. *BMC Bioinformatics,*
8 Suppl 1, S10. doi:10.1186/1471-2105-8-S1-S10

Doynova, M. D., Markworth, J. F., Cameron-Smith, D., Vickers, M. H., & O'Sullivan, J. M.
(2017). Linkages between changes in the 3D organization of the genome and
transcription during myotube differentiation in vitro. *Skelet Muscle, 7*(1), 5.
doi:10.1186/s13395-017-0122-1

Durinck, S., Spellman, P. T., Birney, E., & Huber, W. (2009). Mapping identifiers for the
integration of genomic datasets with the R/Bioconductor package biomaRt. *Nat Protoc,*
4(8), 1184-1191. doi:10.1038/nprot.2009.97

Edmondson, D. G., & Olson, E. N. (1989). A gene with homology to the myc similarity region
of MyoD1 is expressed during myogenesis and is sufficient to activate the muscle
differentiation program. *Genes Dev, 3*(5), 628-640. doi:10.1101/gad.3.5.628

Elhanany-Tamir, H., Yu, Y. V., Shnayder, M., Jain, A., Welte, M., & Volk, T. (2012). Organelle
positioning in muscles requires cooperation between two KASH proteins and
microtubules. *J Cell Biol, 198*(5), 833-846. doi:10.1083/jcb.201204102

Espigat-Georger, A., Dyachuk, V., Chemin, C., Emorine, L., & Merdes, A. (2016). Nuclear
alignment in myotubes requires centrosome proteins recruited by nesprin-1. *J Cell Sci,*
129(22), 4227-4237. doi:10.1242/jcs.191767

Gimpel, P., Lee, Y. L., Sobota, R. M., Calvi, A., Koullourou, V., Patel, R., . . . Gomes, E. R. (2017). Nesprin-1alpha-Dependent Microtubule Nucleation from the Nuclear Envelope via Akap450 Is Necessary for Nuclear Positioning in Muscle Cells. *Curr Biol, 27*(19), 2999-3009 e2999. doi:10.1016/j.cub.2017.08.031

Godinho, S. A., & Pellman, D. (2014). Causes and consequences of centrosome abnormalities in cancer. *Philos Trans R Soc Lond B Biol Sci, 369*(1650). doi:10.1098/rstb.2013.0467

Godinho, S. A., Picone, R., Burute, M., Dagher, R., Su, Y., Leung, C. T., . . . Pellman, D. (2014). Oncogene-like induction of cellular invasion from centrosome amplification. *Nature, 510*(7503), 167-171. doi:10.1038/nature13277

Hasty, P., Bradley, A., Morris, J. H., Edmondson, D. G., Venuti, J. M., Olson, E. N., & Klein, W. H. (1993). Muscle deficiency and neonatal death in mice with a targeted mutation in the myogenin gene. *Nature, 364*(6437), 501-506. doi:10.1038/364501a0

Hinchcliffe, E. H., Miller, F. J., Cham, M., Khodjakov, A., & Sluder, G. (2001). Requirement of a centrosomal activity for cell cycle progression through G1 into S phase. *Science, 291*(5508), 1547-1550. doi:10.1126/science.1056866

Holt, I., Duong, N. T., Zhang, Q., Lam le, T., Sewry, C. A., Mamchaoui, K., . . . Morris, G. E. (2016). Specific localization of nesprin-1-alpha2, the short isoform of nesprin-1 with a KASH domain, in developing, fetal and regenerating muscle, using a new monoclonal antibody. *BMC Cell Biol, 17*(1), 26. doi:10.1186/s12860-016-0105-9

Ieda, M., Fu, J. D., Delgado-Olguin, P., Vedantham, V., Hayashi, Y., Bruneau, B. G., & Srivastava, D. (2010). Direct reprogramming of fibroblasts into functional cardiomyocytes by defined factors. *Cell, 142*(3), 375-386. doi:10.1016/j.cell.2010.07.002

Ishibashi, J., Perry, R. L., Asakura, A., & Rudnicki, M. A. (2005). MyoD induces myogenic differentiation through cooperation of its NH2- and COOH-terminal regions. *J Cell Biol, 171*(3), 471-482. doi:10.1083/jcb.200502101

Kapiloff, M. S., Schillace, R. V., Westphal, A. M., & Scott, J. D. (1999). mAKAP: an A-kinase anchoring protein targeted to the nuclear membrane of differentiated myocytes. *J Cell Sci, 112 (Pt 16)*, 2725-2736. Retrieved from http://www.ncbi.nlm.nih.gov/pubmed/10413680

Kent, W. J., Sugnet, C. W., Furey, T. S., Roskin, K. M., Pringle, T. H., Zahler, A. M., & Haussler, D. (2002). The human genome browser at UCSC. *Genome Res, 12*(6), 996-1006. doi:10.1101/gr.229102

Khodjakov, A., & Rieder, C. L. (2001). Centrosomes enhance the fidelity of cytokinesis in vertebrates and are required for cell cycle progression. *J Cell Biol, 153*(1), 237-242. Retrieved from http://www.ncbi.nlm.nih.gov/pubmed/11285289

Klose, K., Gossen, M., & Stamm, C. (2019). Turning fibroblasts into cardiomyocytes: technological review of cardiac transdifferentiation strategies. *FASEB J, 33*(1), 49-70. doi:10.1096/fj.201800712R

Kronebusch, P. J., & Singer, S. J. (1987). The microtubule-organizing complex and the Golgi apparatus are co-localized around the entire nuclear envelope of interphase cardiac myocytes. *J Cell Sci, 88 (Pt 1)*, 25-34. Retrieved from http://www.ncbi.nlm.nih.gov/pubmed/3327863

Leask, A., Obrietan, K., & Stearns, T. (1997). Synaptically coupled central nervous system
neurons lack centrosomal gamma-tubulin. *Neurosci Lett, 229*(1), 17-20.
doi:10.1016/s0304-3940(97)00412-6

Lechler, T., & Fuchs, E. (2007). Desmoplakin: an unexpected regulator of microtubule
organization in the epidermis. *J Cell Biol, 176*(2), 147-154. doi:10.1083/jcb.200609109

Lee, C., Scherr, H. M., & Wallingford, J. B. (2007). Shroom family proteins regulate gamma-
tubulin distribution and microtubule architecture during epithelial cell shape change.
Development, 134(7), 1431-1441. doi:10.1242/dev.02828

Lee, S. W., Won, J. Y., Yang, J., Lee, J., Kim, S. Y., Lee, E. J., & Kim, H. S. (2015). AKAP6
inhibition impairs myoblast differentiation and muscle regeneration: Positive loop
between AKAP6 and myogenin. *Sci Rep, 5*, 16523. doi:10.1038/srep16523

Levine, M. S., Bakker, B., Boeckx, B., Moyett, J., Lu, J., Vitre, B., . . . Holland, A. J. (2017).
Centrosome Amplification Is Sufficient to Promote Spontaneous Tumorigenesis in
Mammals. *Dev Cell, 40*(3), 313-322 e315. doi:10.1016/j.devcel.2016.12.022

LoMastro, G. M., & Holland, A. J. (2019). The Emerging Link between Centrosome Aberrations
and Metastasis. *Dev Cell, 49*(3), 325-331. doi:10.1016/j.devcel.2019.04.002

Luders, J. (2020). Nucleating microtubules in neurons: Challenges and solutions. *Dev
Neurobiol*. doi:10.1002/dneu.22751

Mamchaoui, K., Trollet, C., Bigot, A., Negroni, E., Chaouch, S., Wolff, A., . . . Mouly, V. (2011).
Immortalized pathological human myoblasts: towards a universal tool for the study of
neuromuscular disorders. *Skelet Muscle, 1*, 34. doi:10.1186/2044-5040-1-34

Meads, T., & Schroer, T. A. (1995). Polarity and nucleation of microtubules in polarized
epithelial cells. *Cell Motil Cytoskeleton, 32*(4), 273-288. doi:10.1002/cm.970320404

Meng, W., Mushika, Y., Ichii, T., & Takeichi, M. (2008). Anchorage of microtubule minus ends
to adherens junctions regulates epithelial cell-cell contacts. *Cell, 135*(5), 948-959.
doi:10.1016/j.cell.2008.09.040

Michel, J. J., Townley, I. K., Dodge-Kafka, K. L., Zhang, F., Kapiloff, M. S., & Scott, J. D. (2005).
Spatial restriction of PDK1 activation cascades by anchoring to mAKAPalpha. *Mol Cell,
20*(5), 661-672. doi:10.1016/j.molcel.2005.10.013

Mulari, M. T., Patrikainen, L., Kaisto, T., Metsikko, K., Salo, J. J., & Vaananen, H. K. (2003).
The architecture of microtubular network and Golgi orientation in osteoclasts--major
differences between avian and mammalian species. *Exp Cell Res, 285*(2), 221-235.
doi:10.1016/s0014-4827(03)00033-8

Nabeshima, Y., Hanaoka, K., Hayasaka, M., Esumi, E., Li, S., Nonaka, I., & Nabeshima, Y.
(1993). Myogenin gene disruption results in perinatal lethality because of severe
muscle defect. *Nature, 364*(6437), 532-535. doi:10.1038/364532a0

Nguyen, M. M., McCracken, C. J., Milner, E. S., Goetschius, D. J., Weiner, A. T., Long, M. K.,
. . . Rolls, M. M. (2014). Gamma-tubulin controls neuronal microtubule polarity
independently of Golgi outposts. *Mol Biol Cell, 25*(13), 2039-2050.
doi:10.1091/mbc.E13-09-0515

Nguyen, M. M., Stone, M. C., & Rolls, M. M. (2011). Microtubules are organized independently of the centrosome in Drosophila neurons. *Neural Dev, 6*, 38. doi:10.1186/1749-8104-6-38

Ori-McKenney, K. M., Jan, L. Y., & Jan, Y. N. (2012). Golgi outposts shape dendrite morphology by functioning as sites of acentrosomal microtubule nucleation in neurons. *Neuron, 76*(5), 921-930. doi:10.1016/j.neuron.2012.10.008

Pare, G. C., Easlick, J. L., Mislow, J. M., McNally, E. M., & Kapiloff, M. S. (2005). Nesprin-1alpha contributes to the targeting of mAKAP to the cardiac myocyte nuclear envelope. *Exp Cell Res, 303*(2), 388-399. doi:10.1016/j.yexcr.2004.10.009

Passariello, C. L., Li, J., Dodge-Kafka, K., & Kapiloff, M. S. (2015). mAKAP-a master scaffold for cardiac remodeling. *J Cardiovasc Pharmacol, 65*(3), 218-225. doi:10.1097/FJC.0000000000000206

Patra, C., Diehl, F., Ferrazzi, F., van Amerongen, M. J., Novoyatleva, T., Schaefer, L., . . . Engel, F. B. (2011). Nephronectin regulates atrioventricular canal differentiation via Bmp4-Has2 signaling in zebrafish. *Development, 138*(20), 4499-4509. doi:10.1242/dev.067454

Prosser, S. L., & Pelletier, L. (2015). Centrosome biology: the ins and outs of centrosome assembly. *Curr Biol, 25*(15), R656-659. doi:10.1016/j.cub.2015.06.038

Prosser, S. L., & Pelletier, L. (2020). Centriolar satellite biogenesis and function in vertebrate cells. *J Cell Sci, 133*(1). doi:10.1242/jcs.239566

Randles, K. N., Lam le, T., Sewry, C. A., Puckelwartz, M., Furling, D., Wehnert, M., . . . Morris, G. E. (2010). Nesprins, but not sun proteins, switch isoforms at the nuclear envelope during muscle development. *Dev Dyn, 239*(3), 998-1009. doi:10.1002/dvdy.22229

Rauch, A., Thiel, C. T., Schindler, D., Wick, U., Crow, Y. J., Ekici, A. B., . . . Reis, A. (2008). Mutations in the pericentrin (PCNT) gene cause primordial dwarfism. *Science, 319*(5864), 816-819. doi:10.1126/science.1151174

Rosenbloom, K. R., Sloan, C. A., Malladi, V. S., Dreszer, T. R., Learned, K., Kirkup, V. M., . . . Kent, W. J. (2013). ENCODE data in the UCSC Genome Browser: year 5 update. *Nucleic Acids Res, 41*(Database issue), D56-63. doi:10.1093/nar/gks1172

Ruehr, M. L., Russell, M. A., Ferguson, D. G., Bhat, M., Ma, J., Damron, D. S., . . . Bond, M. (2003). Targeting of protein kinase A by muscle A kinase-anchoring protein (mAKAP) regulates phosphorylation and function of the skeletal muscle ryanodine receptor. *J Biol Chem, 278*(27), 24831-24836. doi:10.1074/jbc.M213279200

Sanchez-Huertas, C., Freixo, F., Viais, R., Lacasa, C., Soriano, E., & Luders, J. (2016). Non-centrosomal nucleation mediated by augmin organizes microtubules in post-mitotic neurons and controls axonal microtubule polarity. *Nat Commun, 7*, 12187. doi:10.1038/ncomms12187

Sanchez, A. D., & Feldman, J. L. (2017). Microtubule-organizing centers: from the centrosome to non-centrosomal sites. *Curr Opin Cell Biol, 44*, 93-101. doi:10.1016/j.ceb.2016.09.003

Sir, J. H., Putz, M., Daly, O., Morrison, C. G., Dunning, M., Kilmartin, J. V., & Gergely, F. (2013). Loss of centrioles causes chromosomal instability in vertebrate somatic cells. *J Cell Biol, 203*(5), 747-756. doi:10.1083/jcb.201309038

Srsen, V., Fant, X., Heald, R., Rabouille, C., & Merdes, A. (2009). Centrosome proteins form
an insoluble perinuclear matrix during muscle cell differentiation. *BMC Cell Biol, 10*, 28.
doi:10.1186/1471-2121-10-28

Tassin, A. M., Maro, B., & Bornens, M. (1985). Fate of microtubule-organizing centers during
myogenesis in vitro. *J Cell Biol, 100*(1), 35-46. Retrieved from
http://www.ncbi.nlm.nih.gov/pubmed/3880758

The Gene Ontology, C. (2017). Expansion of the Gene Ontology knowledgebase and
resources. *Nucleic Acids Res, 45*(D1), D331-D338. doi:10.1093/nar/gkw1108

Toya, M., Kobayashi, S., Kawasaki, M., Shioi, G., Kaneko, M., Ishiuchi, T., . . . Takeichi, M.
(2016). CAMSAP3 orients the apical-to-basal polarity of microtubule arrays in epithelial
cells. *Proc Natl Acad Sci U S A, 113*(2), 332-337. doi:10.1073/pnas.1520638113

Venuti, J. M., Morris, J. H., Vivian, J. L., Olson, E. N., & Klein, W. H. (1995). Myogenin is
required for late but not early aspects of myogenesis during mouse development. *J Cell
Biol, 128*(4), 563-576. Retrieved from http://www.ncbi.nlm.nih.gov/pubmed/7532173

Vergarajauregui, S., Becker, R., Steffen, U., Sharkova, M., Esser, T., Petzold, J., . . . Engel, F.
B. (2020). AKAP6 orchestrates the nuclear envelope microtubule-organizing center by
linking golgi and nucleus via AKAP9. *Elife, 9*. doi:10.7554/eLife.61669

Weintraub, H., Tapscott, S. J., Davis, R. L., Thayer, M. J., Adam, M. A., Lassar, A. B., & Miller,
A. D. (1989). Activation of muscle-specific genes in pigment, nerve, fat, liver, and
fibroblast cell lines by forced expression of MyoD. *Proc Natl Acad Sci U S A, 86*(14),
5434-5438. doi:10.1073/pnas.86.14.5434

Yamamoto, K., Kishida, T., Sato, Y., Nishioka, K., Ejima, A., Fujiwara, H., . . . Mazda, O. (2015).
Direct conversion of human fibroblasts into functional osteoblasts by defined factors.
Proc Natl Acad Sci U S A, 112(19), 6152-6157. doi:10.1073/pnas.1420713112

Yang, R., & Feldman, J. L. (2015). SPD-2/CEP192 and CDK Are Limiting for Microtubule-
Organizing Center Function at the Centrosome. *Curr Biol, 25*(14), 1924-1931.
doi:10.1016/j.cub.2015.06.001

Yue, F., Cheng, Y., Breschi, A., Vierstra, J., Wu, W., Ryba, T., . . . Mouse, E. C. (2014). A
comparative encyclopedia of DNA elements in the mouse genome. *Nature, 515*(7527),
355-364. doi:10.1038/nature13992

Zebrowski, D. C., Vergarajauregui, S., Wu, C. C., Piatkowski, T., Becker, R., Leone, M., . . .
Engel, F. B. (2015). Developmental alterations in centrosome integrity contribute to the
post-mitotic state of mammalian cardiomyocytes. *Elife, 4*. doi:10.7554/eLife.05563

Zhang, Q., Bethmann, C., Worth, N. F., Davies, J. D., Wasner, C., Feuer, A., . . . Shanahan,
C. M. (2007). Nesprin-1 and -2 are involved in the pathogenesis of Emery Dreifuss
muscular dystrophy and are critical for nuclear envelope integrity. *Hum Mol Genet,
16*(23), 2816-2833. doi:10.1093/hmg/ddm238

Zhang, Q., Skepper, J. N., Yang, F., Davies, J. D., Hegyi, L., Roberts, R. G., . . . Shanahan,
C. M. (2001). Nesprins: a novel family of spectrin-repeat-containing proteins that
localize to the nuclear membrane in multiple tissues. *J Cell Sci, 114*(Pt 24), 4485-4498.
Retrieved from http://www.ncbi.nlm.nih.gov/pubmed/11792814

Zhang, X., Chen, M. H., Wu, X., Kodani, A., Fan, J., Doan, R., . . . Walsh, C. A. (2016). Cell-
Type-Specific Alternative Splicing Governs Cell Fate in the Developing Cerebral
Cortex. *Cell, 166*(5), 1147-1162 e1115. doi:10.1016/j.cell.2016.07.025

Zhu, L. J., Gazin, C., Lawson, N. D., Pages, H., Lin, S. M., Lapointe, D. S., & Green, M. R.
(2010). ChIPpeakAnno: a Bioconductor package to annotate ChIP-seq and ChIP-chip
data. *BMC Bioinformatics, 11*, 237. doi:10.1186/1471-2105-11-237

1226

1227

Figure Supplements

Figure 1 – figure supplement 1. Centrosomal protein recruitment to the nuclear envelope is gradual.
(A) C2C12 cells were differentiated for 24h and PCM-1 was detected by immunostaining. Different stages of PCM-1 recruitment are shown with increasing coverage of the nuclear envelope from left to right. Scale bar = 10 μm. **(B)** A more detailed quantification of pericentrin (PCNT) and AKAP9-positive nuclei in differentiating C2C12 (treated as described in Figure 1F) showing that myogenin depletion affects also partial recruitment of PCNT and AKAP9. Data are represented as individual biological replicates, together with mean ± SD. ***: $p < 0.001$. n = 3. **(C)** C2C12 myoblasts were transfected with si-ctrl or PCM-1 siRNA (si-PCM1) and differentiated for two days. AKAP9 was detected by immunostaining and subsequent quantification (H) showed that PCM-1 depletion does not affect AKAP) positive nuclei. Data are represented as individual biological replicates, together with mean ± SD. ns: $p > 0.05$. n = 3.

1243

Figure 2 – figure supplement 1. Myogenin is required for MyoD-induced nuclear envelope recruitment of PCM-1.
(A) Three days after ectopic expression of MyoD-GFP in NIH3T3 fibroblasts, immunostaining revealed that nuclear PCM-1 localization is associated with high myogenin (Myog) expression (yellow arrowhead). Scale bars: 10 µm. (B) NIH3T3 cells were treated with negative control (si-ctrl) or myogenin (si-Myog) siRNA and 24 h later transfected with a construct encoding MyoD-GFP. Immunofluorescence staining indicates that myogenin is required for MyoD-induced PCM-1 nuclear recruitment. Scale bars: 10 µm.

1253

Figure 4 – figure supplement 1. Quantification of nesprin-1α and AKAP6 signal at the nuclear envelope.

Differentiated C2C12 cells were immunostained for nesprin-1α and AKAP6. Signal intensity was quantified as line scan profiles across the nuclear envelope. Ten line scan profiles were obtained in regular intervals along the nuclear periphery and pooled from ten nuclei. Data are shown as mean ± SD.

1259

1260

Figure 5 – figure supplement 1. Akap6 depletion does not affect nesprin-1α.
Enriched C2C12 myotubes were transfected with scrambled or Akap6-siRNA and immunostained for
Akap6 and nesprin-1α. Scale bars: 10 μm.

1264

1265

Figure 6 – figure supplement 1. AKAP6 depletion does not affect steady state microtubule organization but reduces detyrosinated microtubules.

(A-B) Enriched C2C12 myotubes (troponin I) were transfected with negative control (si-ctrl) or Akap6 (si-Akap6) siRNA and subsequently analyzed for overall microtubule (α-tubulin) structure (A) or detyrosinated microtubules (detyr α-tubulin). Scale bars: 10 μm.