Physiological and transcriptomic response of *Medicago truncatula* to colonization with high and low benefit arbuscular mycorrhizal fungi

Kevin R. Cope[1†], Arjun Kafle[1,2,†], Jaya K. Yakha[1], Philip E. Pfeffer[3], Gary D. Strahan[3], Kevin Garcia[2], Senthil Subramanian[1,4], Heike Bücking[1,5*]

[1]South Dakota State University, Biology and Microbiology Department, Brookings, SD 57007 USA

[2]Department of Crop and Soil Sciences, North Carolina State University, Raleigh, NC 27695 USA

[3]USDA, Agricultural Research Service, Eastern Regional Research Center, Wyndmoor, PA, 19038 USA

[4]Department of Agronomy, Horticulture, and Plant Science, South Dakota State University, Brookings, SD, 57007 USA

[5]Division of Plant Science and Technology, University of Missouri, Columbia, MO 65211

[†]These Authors contributed equally to the work.

*Address correspondence to Heike Bücking (heike.buecking@missouri.edu)

Funding. This work was funded by the USDA (2017-67014-26530), the SD Soybean Research and Promotion Council, and the Agricultural Experiment Station at SDSU to HB and SS. KG acknowledges support of the North Carolina Agriculture Research Service

ABSTRACT

Arbuscular mycorrhizal (AM) fungi form a root endosymbiosis with many agronomically

important crop species and both enhance the ability of their host to obtain nutrients from

the soil and increase host tolerance to biotic and abiotic stressors. However, AM fungal

species differ in the benefits they provide to their host plants. Here, we examined the

putative molecular mechanisms involved in the regulation of the physiological response

of *Medicago truncatula* to either *Rhizophagus irregularis or Glomus aggregatum,* a high

or a low benefit AM fungus, respectively. Colonization with *R. irregularis* led to higher

growth and nutrient uptake benefits than the colonization with *G. aggregatum*. These

benefits were linked to an elevated expression in the roots of genes involved in

strigolactone biosynthesis (*NSP1*, *NSP2*, *CCD7*, and *MAX1a*), mycorrhiza-induced

phosphate (*PT8*), ammonium (*AMT2;3*), and nitrate (*NPF4.12*) transporters and the

putative ammonium transporter *NIP1;5*. *R. irregularis* also stimulated the expression of

photosynthesis-related genes in the shoot and the upregulation of the sugar transporters

SWEET1.2, SWEET3.3 and *SWEET 12* and the lipid biosynthesis gene *RAM2* in the

roots. In contrast, *G. aggregatum* induced the expression of biotic stress defense

response genes in the shoots and several genes associated with abiotic stress in the

roots. This suggests that either the host perceives colonization by *G. aggregatum* as a

pathogen attack or that *G. aggregatum* can prime host defense responses. Our findings

reveal novel insights into the molecular mechanisms that control the host plant response

to colonization with high- and low-benefit arbuscular mycorrhizal fungal symbionts.

KEYWORDS

Medicago truncatula, arbuscular mycorrhizal fungi, symbiosis, nutrient exchange,

phosphorus, nitrogen, carbon, transporters, and SWEET

INTRODUCTION

Nitrogen (N) and phosphorus (P) are the main nutrients that limit plant growth in

natural and agricultural ecosystems (LeBauer and Treseder 2008; Vitousek et al. 2010).

Conventional management practices typically address both N and P limitations through

the application of chemical fertilizers, but fertilizer production, transportation, and

application are both economically and environmentally expensive (Havlin et al. 2014).

Implementing more sustainable methods to mitigate nutrient deficiencies in crops is

essential to prevent continued environmental degradation and to still meet the nutritional

demands of a growing human population.

To overcome nutrient limitations, plants have evolved the capacity to form alliances

with mutualistic, root-associated microbes (Martin et al. 2017). The most widespread

mutualism is the arbuscular mycorrhizal (AM) symbiosis formed by 72% of all known plant

species and a relatively small number of fungal species in the Glomeromycotina

(Brundrett and Tedersoo 2018; Spatafora et al. 2016). AM fungi provide the host plant

with mineral nutrients, such as P and N, and improve the resistance of plants against

abiotic and biotic stressors (Bücking and Kafle 2015; Kafle et al. 2019). Under P

deficiency, plants increase the production of strigolactones, a class of phytohormones

that also serve as signaling molecules perceived by AM fungi (Yoneyama et al. 2007).

Strigolactones are exuded into the rhizosphere through the ATP-binding cassette

transporter PDR1 (Kretzschmar et al. 2012). AM fungal perception of strigolactones

triggers spore germination, hyphal branching, and the production of short-chain chitin

oligomers (Besserer et al. 2008; Genre et al. 2013). These, along with lipo-

chitooligosaccharides (Maillet et al. 2011; Rush et al. 2020), comprise the so-called "Myc

factors" that prime the host plant for colonization by activating the "common symbiosis

signaling pathway" (CSSP; reviewed in MacLean et al., 2017).

The key steps of the CSSP include: 1) the perception of Myc factors at the plasma

membrane of root epidermal cells by lysine-motif receptor-like kinases (*e.g.*, NFP and

LYK3; Feng et al., 2019); 2) the activation of the secondary messenger mevalonate that

likely activates multiple nuclear membrane-localized calcium channels (CASTOR,

DMI1/POLLUX, and CNGC15); 3) oscillations in calcium ion concentration in the

nucleoplasm that activate a calcium- and calmodulin-dependent protein kinase

(DMI3/CCaMK; Lévy et al., 2004; Singh and Parniske, 2012); 4) phosphorylation of IPD3

which, in concert with DELLA, induces the expression of RAM1; and 5) activation of

several other transcription factors required for the development and regulation of the AM

symbiosis (Pimprikar et al., 2016; Pimprikar and Gutjahr, 2018). Following root

penetration, AM hyphae proliferate both intra- and intercellularly and form highly branched

nutrient exchange structures called arbuscules in cortical cells (Park et al. 2015).

Surrounding the arbuscule is a plant-derived periarbuscular membrane (PAM) that is

enriched in AM-induced transporters that facilitate reciprocal nutrient exchange between

the fungus and the plant (Garcia et al. 2016). In *Medicago truncatula,* three AMT2 family

ammonium transporters (AMT2;3, AMT2;4, and AMT2;5; Breuillin-Sessoms et al., 2015)

and the phosphate transporters *PT4* (Javot et al. 2007) and *PT8* (Breuillin-Sessoms et al.

2015) are involved in the uptake of N and P from the periarbuscular space between the

fungal plasma membrane and the PAM. In exchange, host plants transfer about 20% of

their photosynthetically derived carbon to their fungal partners, and these high carbon

costs force host plants to strongly regulate the carbon flux to their fungal symbionts

(Wright et al. 1998). The host plant releases lipids and sugars into the mycorrhizal

interface likely via the putative lipid transporters *STR* and *STR2* (Zhang et al., 2010; Roth

and Paszkowski, 2017) and the bidirectional sugar transporter *SWEET1.2* (formerly

SWEET1b, An et al., 2019; Doidy et al., 2019).

AM fungi differ in the benefits that they provide for their host plant. The cooperative

AM fungus *Rhizophagus irregularis* 09 (Ri09) transfers more N and P to its host plant and

contributes to higher mycorrhizal growth responses than the less-cooperative fungus

Glomus aggregatum 165 (Ga165; Fellbaum et al., 2014). Symbiosis with Ga165 is also

more costly for the host plant (measured as carbon costs per P transferred) than with

Ri09, and *M. truncatula* preferentially allocates more carbon to Ri09 than to Ga165 (Kiers

et al. 2011). Collectively, these studies suggest that carbon to nutrient exchange

processes are driven by biological market dynamics, but the molecular mechanisms that

regulate these processes are largely unknown. We hypothesized that the host plant

regulates the expression of key genes in response to the colonization of AM fungi that

differ in the nutrient benefits that they provide for the host. To test this hypothesis, we

examined the physiological and transcriptomic responses of the host plant *M. trancatula*

to colonization by either high- (Ri09) or low-benefit (Ga165) AM fungal partners under N

and P limitation.

RESULTS

Colonization by *Rhizophagus irregularis* 09 leads to greater plant biomass and

phosphorus and nitrogen uptake than *Glomus aggregatum* 165

We used a dual-compartment pot system to compare the overall benefit that Ga165 and

Ri09 provided to *M. truncatula* (**Supplemental Fig. S1**). The percent root length

colonization by intraradical hyphae and arbuscules was comparable in roots colonized by

either fungus; however, the percent of roots containing vesicles was significantly higher

in Ga165-colonized roots (**Fig. 1A**). Arbuscular width was comparable for both species of

AM fungi (**Fig. 1B-D**). Compared to non-mycorrhizal (NM) plants, Ri09 colonization

resulted in greater shoot and root biomass, whereas Ga165 provided no growth benefits

(**Fig. 2A-B**). Both fungi led to increased P content (**Fig. 2C-D**) and concentration

(**Supplemental Fig. S2**) in the shoots and roots of their host, but Ri09 transferred more

P than Ga165. Nitrogen-15 (^{15}N) enrichment levels in the roots and shoots of NM plants

did not exceed natural abundance levels (**Fig. 2E-F**), which confirms that there was no

mass flow of ^{15}N from the HC into the RC. Both fungi transferred ^{15}N from the HC to their

host, but Ga165 only increased root ^{15}N enrichment, whereas Ri09 increased both root

and shoot ^{15}N enrichment. Overall, these data indicate that although Ga165 and Ri09

showed similar levels of root colonization, only Ri09 colonization led to increased growth,

most likely by providing the host plant with greater access to both P and N.

Figure 1. Arbuscular mycorrhizal colonization of *Medicago truncatula* roots and arbuscule size. **A,** percent root length colonization of roots inoculated with *Glomus aggregatum* 165 (Ga165; grey) or *Rhizophagus irregularis* 09 (Ri09; dark grey), measured for intraradical hyphae (IRH), arbuscules, and vesicles. Data points represent individual values for biological replicates (n = 5) and bars represent the mean ± SEM. **B-C**, representative confocal microscope images of Ga165 (**B**) and Ri09 (**C**) arbuscules stained with wheat germ agglutinin conjugated to Alexa Fluor 488. Scale bars = 50 µm. **D**, violin plot of arbuscule widths measured in roots colonized by Ga165 (grey) or Ri09 (dark grey) overlayed with box and whisker plots. In total, 488 and 508 arbuscules were measured in Ga165 and Ri09-colonized roots, respectively. All data were analyzed using a Welch's two-sample t-test and statistically significant differences (p ≤ 0.05) are indicated with an asterisk.

Figure 2. Biomass and nutrient content of *Medicago truncatula* plants colonized by different arbuscular mycorrhizal fungi. Shown are dry weight (**A**,**B**), phosphorus content (**C**,**D**), and % nitrogen-15 enrichment (**E**,**F**) of roots (left) and shoots (right). The plants were inoculated with either *Glomus aggregatum* 165 (Ga165) or *Rhizophagus irregularis* 09 (Ri09) and compared to non-mycorrhizal (NM) control plants. Data points represent individual values for biological replicates (n = 5) and bars represent the mean of each treatment ± SEM. Different letters on the bars indicate statistically significant (p ≤ 0.05) differences based on one-way ANOVA and LSD test. F statistics and p-values are shown in **Supplemental Table S1**.

Colonization by either fungus alters gene expression more dramatically in the roots than in the shoots of *Medicago truncatula*

To identify potential molecular mechanisms responsible for the observed physiological responses, we used RNA sequencing (RNA-seq) to evaluate gene expression patterns in roots and shoots of mycorrhizal and non-mycorrhizal plants (**Supplemental File 1**). Analysis of the root data revealed distinct, treatment-specific hierarchical clustering among biological replicates (n=3, Ga165 and NM; n=4, Ri09) using a dendrogram, multidimensional scaling, and principal component analyses (**Supplemental Fig. S3**). In contrast, analysis of the shoot data showed no distinct clustering among biological replicates (n=4, Ga165 and NM; n=3, Ri09) within each treatment (**Supplemental Fig. S4**), suggesting that particularly in the roots, gene expression is driven by the fungal partner. Consequently, the overall number of significant differentially expressed genes (DEGs) was lower in the shoots than in the roots (**Supplemental Fig. S5**).

Both shared and distinct sets of genes are differentially regulated in *Medicago truncatula* by colonization with either fungus

To identify key differences in the response of the host plant to both fungal species, we sorted all DEGs from each treatment and used an arbitrary threshold of $\log_2$(fold-change) > 2 and a q-value < 0.05 (**Fig. 3A**). In the roots, this threshold revealed that 364 genes were upregulated in roots colonized by either fungus, and 197 or 165 genes were upregulated by colonization with only Ga165 or Ri09, respectively (**Fig. 3B**). Ga165 and Ri09 colonization led to the downregulation of 41 shared genes, and 172 and 77 unique genes, respectively. The clustering of DEGs allowed us to identify specific groups of

Figure 3. Significantly differentially expressed genes in roots (**A**–**C**) and shoots (**D**–**F**) of *Medicago truncatula* colonized by different AM fungi. **A** and **D**, Number of significantly differentially expressed genes (DEGs; $\log_2$[fold-change] > 2 and a q-value < 0.05) in plants colonized by *Glomus aggregatum* 165 (Ga165) or *Rhizophagus irregularis* 09 (Ri09) compared either to non-mycorrhizal (NM) control plants or to each other. **B** and **E**, Venn diagrams of significantly upregulated (yellow) and downregulated (blue) DEGs unique to Ga165 and Ri09-colonized plants or shared by both. **C** and **F**, Heat maps of fragments per kilobase of transcript per million mapped reads (FPKM) values + 1 for all significant DEGs from each treatment. Gene clusters are assigned based on second-order and sub-clusters based on third- or fourth-order branching. Upregulated genes are shown in yellow and downregulated genes in blue with intensity based on row z-score. All DEGs in each plot for roots and shoots, respectively, are listed in **Supplemental Files 2 and 3**.

similarly expressed genes (**Fig. 3C**). Clusters A and B are comprised of genes that were

strongly downregulated in mycorrhizal roots, cluster C1 contains genes that were

upregulated in mycorrhizal roots, clusters C2 and D include genes primarily upregulated

by Ga165 colonization, and cluster C3 contains genes exclusively upregulated by Ri09

colonization (**Supplemental File 2)**.

Although fewer genes were differentially expressed in the shoots, many were

above our applied threshold (**Fig. 3D**). We found that 48 and 14 genes were up, and 13

and 27 genes were downregulated in the shoots of Ga165 and Ri09-colonized plants,

respectively (**Fig. 3E**). In contrast to the roots, clustering of DEGs based on expression

patterns in the shoots (**Fig. 3F)** revealed that only a small number of genes were

upregulated (cluster C1) or downregulated (cluster A) in mycorrhizal plants. In addition,

some genes were only downregulated by Ga165 (cluster B) or by Ri09 colonization

(cluster D), while others were primarily upregulated by Ga165 colonization (cluster C2;

Supplemental File 3). Collectively, these results demonstrate that compared to shoots,

gene expression patterns in roots respond more strongly to colonization with different AM

fungi, and that both fungi lead to distinct differences particularly in the root and to a lesser

degree also in the shoot transcriptome profiles.

Genes critical for the establishment of AM symbiosis are more strongly

upregulated by *Rhizophagus irregularis* 09 than *Glomus aggregatum* 165

We conducted a targeted analysis of the root RNA-seq data by first evaluating the

expression of many genes known to play a role in AM symbiosis (**Supplemental Table**

S2). Among six core strigolactone biosynthesis or exporter genes (**Fig. 4A**), we observed

Figure 4. Expression patterns of genes encoding enzymes involved in the strigolactone (**A**) and gibberellic acid (**B**) biosynthesis pathways in the roots of *Medicago truncatula* colonized by arbuscular mycorrhizal fungi. Each diagram portrays the relative expression levels of genes that encode components of the pathways shown as fragments per kilobase of transcript per million mapped reads (FPKM; blue=low, yellow=high) in roots from non-mycorrhizal (NM) plants or plants colonized by either *Glomus aggregatum* 165 (Ga165) or *Rhizophagus irregularis* 09 (Ri09). Letters within boxes indicate significant differences (q-value < 0.05) in FPKM values between treatments and were determined as part of the CuffDiff2 analysis (see **Supplemental Table S2**). Abbreviations: **A**, GAP, glyceraldehyde 3-phosphate; DXS2, 1-deoxyxylulose 5-phosphate synthase2; MEP, 2-methyl-erythritol-4-phosphate; GGDP geranylgeranyl pyrophosphate; D27, Dwarf27; CCD, carotenoid cleavage dioxygenase; and PDR1a, pleiotropic drug resistance 1a. **B**, CPS, *ent*-copalyl diphosphate synthase; KS, *ent*-kaurene synthase; KO, *ent*-kaurene oxidase; KAO, *ent*-kaurenoic acid oxidase; GA13ox, gibberellin 13-oxidase; GA20ox, gibberellin 20-oxidase; GA3ox, gibberellin 3-oxidase; and GA2ox, gibberellin 2-oxidase.

that two (*DXS2* and *D27*) were upregulated in AM roots compared to the NM control; one

(*MAX1a*) was only upregulated in Ri09-colonized roots, while three (*CCD7*, *CCD8-1* and

PDR1a) were not affected. Two GRAS transcription factors (*NSP1* and *NSP2*) that

regulate strigolactone biosynthesis and other symbiosis genes were also upregulated,

particularly in Ri09-colonized roots. The combined increased expression of *NSP1*, *NSP2*,

CCD7, and *MAX1a* in Ri09-colonized roots compared to Ga165-colonized roots suggests

that Ri09 more strongly induces strigolactone biosynthesis than Ga165.

Among the CSSP genes, the lipochitooligosaccharide receptor *NFP* was

downregulated and *PUB1*, a negative regulator of AM symbiosis (Vernié et al. 2016), was

upregulated in roots colonized by either fungus (**Supplemental Table S2**). None of the

other 13 analyzed CSSP genes were affected. However, gibberellic acid (GA) is known

to play an important role in regulating AM colonization (Floss et al. 2013; Nouri et al.

2021), and several genes regulating the biosynthesis and degradation of GA were

differentially expressed by mycorrhizal colonization (**Fig. 4B**). In Ri09-colonized roots, the

expression of the genes encoding *ent*-copalyl diphosphate synthase (*CPS*) and

GA3oxidase (*GA3ox*) were down-regulated compared to NM roots, but the gene encoding

ent-kaurene oxidase (*KO*) was upregulated. In Ga165-colonized roots, four GA

biosynthesis genes (*GA13ox* and three homologs of *GA20ox*) were significantly

upregulated compared to NM roots. In Ri09-colonized roots also two of the *GA20ox*

homologs were upregulated, but not as strongly as by GA165. Similarly, the GA

degradation gene *GA2ox1* was also upregulated in mycorrhizal roots, but more strongly

in Ga165-colonized roots (**Supplemental Table S2**).

Given the higher expression of more GA biosynthesis genes in Ga165 compared

to Ri09-colonized roots, we hypothesize that GA plays a stronger role in regulating the

symbiosis with GA165. Therefore, we evaluated the expression of genes downstream of

of the CSSP and DELLA1/2, which are proteins that are negatively regulated by GA

(Davière and Achard 2013). We evaluated the expression of 21 genes (**Supplemental**

Table S3), including: seven AM-induced transcription factors that regulate arbuscule

development, function, or degradation — *DIP1*, *RAD1*, *RAM1*, *MIG1/2/3*, and *MYB1*;

three cellular remodeling genes — *EXO70I*, *SYP132A*, and *VAPYRIN*; two phosphate

transporters (*PT4* and *PT8*); three ammonium transporters (*AMT2;3*, *AMT2;4*, and

AMT2;5); the putative lipid transporters *STR* and *STR2* and three lipid biosynthesis genes

— *RAM2*, *FatM*, and *WRI5a*; and the sugar transporter *SWEET1.2*. Almost all these

genes were upregulated by both AM fungal species; however, six were substantially more

upregulated by Ri09, including: *RAD1*, *MYB1*, *PT8*, *AMT2;3*, *RAM2*, and *SWEET1.2* (**Fig.**

5A-F). Together with the physiological results, the increased upregulation of *PT8*,

AMT2;3, and *SWEET1.2* in Ri09-colonized roots is suggestive of greater nutrient

exchange activity across the PAM than in Ga165-colonized roots.

Colonization with *Glomus aggregatum* 165 and *Rhizophagus irregularis* 09

differentially affect the expression of transporters

Gene ontology (GO) enrichment analysis revealed several cellular components,

molecular functions, and/or biological processes that were significantly enriched among

genes that are only upregulated (**Supplemental Fig. S6**) or downregulated in AM roots

(**Supplemental Fig. S7**). Predominant among these were GO terms related to

Figure 5. Relative expression of six genes in the roots of *Medicago truncatula* after colonization with different arbuscular mycorrhizal fungi. Expression of **A**, *Required for Arbuscule Development 1* (*RAD1*); **B**, *MYB-like Transcription Factor 1* (*MYB1*); **C**, *Phosphate Transporter 8* (*PT8*); **D**, *Ammonium Transporter 2;3* (*AMT2;3*); **E**, *Reduced Arbuscular Mycorrhization 2* (*RAM2*); and **F**, *Sugars Will Eventually Be Exported Transporter 1.2* (*SWEET1.2*) in the roots of non-mycorrhizal plants (NM, n=3) and plants inoculated with *Glomus aggregatum* 165 (Ga165, n=3) or *Rhizophagus irregularis* 09 (Ri09, n=4). Expression levels are shown as fragments per kilobase of transcript per million mapped reads (FPKM). Data points represent individual values for biological replicates (n = 5) and bars represent the mean of each treatment ± SEM. Significant differences (q-value < 0.05) in FPKM values between treatments were determined as part of the CuffDiff2 analysis and are indicated using different letters (see **Supplemental Table S3**).

transmembrane transport. We filtered all gene annotations containing the term "transport"

from the *M. truncatula* JCVI 4.0v2 annotation file (Tang et al. 2014) and found 259 genes

that were significantly differentially regulated in mycorrhizal roots (**Supplemental File 5**).

Based on their function, we further subdivided these genes into two groups: transport of

mineral nutrients (72 genes) and secondary metabolites (187 genes). Compared to NM

roots, 12 mineral transporters were up- and 17 were down-regulated by both AM fungi

(**Fig. 6A**). This suggests that a core set of mineral transporters in AM roots are similarly

regulated independent of fungal benefits. However, a unique set of transporters were

differentially regulated by each AM fungal species: 14 were up- and 13 were down-

regulated in Ga165-colonized roots, while only 7 were up- and 12 were down-regulated

in Ri09-colonized roots. From the identified 16 groups of mineral transporters (**Fig. 6B**),

we focused on ammonium, nitrate, phosphate, and sulfate transporters with FPKM ≥ 10,

due to the key role that AM fungi play in the transport of these nutrients (**Fig. 6C-F**).

Among the five differentially regulated ammonium transporters, one belongs to the AMT1

ammonium transporter family and four to the AMT2 family. Compared to AM roots,

AMT1;1 and two of the AMT2 family transporters (*AMT2;1* and *AMT2;6*) were highly

expressed in NM roots, while *AMT2;3* and *AMT2;5* were specifically induced in

mycorrhizal roots. *AMT2;3* was particularly up-regulated in Ri09-colonized roots (**Fig.**

6C).

We also identified seven differentially regulated nitrate transporters (**Fig. 6D**). Five

belong to the nitrate/peptide transporter (NPF) family, and two are putative nitrate

transporters. Interestingly, the most highly expressed nitrate transporter (*NPF4.12*) was

highly upregulated in AM roots, particularly in Ri09-colonized roots. In contrast, five nitrate

Figure 6. Expression patterns of significantly differentially expressed mineral transporter genes in the roots of *Medicago truncatula* after colonization with different arbuscular mycorrhizal fungi. **A**, Venn diagrams showing the number of mineral transporters that are significantly up- (yellow) or down-regulated (blue) in roots colonized by *Glomus aggregatum* 165 (Ga165) or *Rhizophagus irregularis* 09 (Ri09), or both. **B**, summary of the number of genes from specific gene annotation groups of mineral transporters that are up- or down-regulated in roots colonized by Ga165 or Ri09, or both (see also **Supplemental File 6**). Expression patterns of genes annotated as **C**, ammonium; **D**, nitrate; **E**, phosphate; and **F**, sulfate transporters in the roots of non-mycorrhizal plants (NM; n=3) and plants inoculated with Ga165 (n=3) or Ri09 (n=4). The bars in C–F represent the mean of each treatment ± SEM (data points for biological replicates are displayed in **Supplemental Figure S8**). Expression levels are shown as fragments per kilobase of transcript per million mapped reads (FPKM), only genes with FPKM ≥ 10 are shown, significant differences (q-value < 0.05) in FPKM values between treatments were determined as part of the CuffDiff2 analysis and are indicated using different letters on the bars.

transporters were more highly expressed in Ga165-colonized roots than in Ri09-colonized

roots (*NPF1.2*, *NPF1.5*, *NPF1.6*, *Medtr4g114340*, and *Medtr1g057460*), and two (*NPF1.2*

and *Medtr1g057460*) also differed from control roots. Only the nitrate transporter *NPF5.24*

and *Medtr1g057460* showed higher expression in NM roots than in AM-colonized roots.

Seven phosphate transporters were differentially regulated (**Fig. 6E**). Four belong

to the high-affinity inorganic phosphate transporter family PHT1 (*PT4*, *PT6*, *PT8*, and

PT9), two were annotated as high-affinity inorganic phosphate transporters

(*Medtr1g043290* and *Medtr1g074930*), and one as a phosphate transporter PHO1-like

protein. As expected, two of the PHT1 transporter genes, *PT4* and *PT8*, were only

expressed in AM roots, and *PT8* expression was higher in Ri09-colonized roots. By

contrast, the expression of *Medtr1g074930* was strongly upregulated in Ga165-colonized

roots compared to NM and Ri09-colonized roots. The PHO1-like transporter

(*Medtr1g041695*) showed higher expression levels in NM than in AM roots.

We identified six sulfate transporters that were differentially regulated among the

three treatments (**Fig. 6F**). Two were annotated as high affinity sulfate transporter 1

(SULTR1) family genes (*SULTR1.1* and *1.2*), three as sulfate/bicarbonate/oxalate

exchanger and transporter genes (*SULTR2.2*, *3.1*, and *4.1*), and one as a sulfate

transporter-like gene (*MOT1.1*). While the expression levels of *SULTR2.2* and *SULTR4.1*

were higher in NM roots than in AM roots, the remaining sulfate transporters were

upregulated by colonization with Ri09 (*SULTR1.1*), Ga165 (*SULTR3.1*), or by both fungi

(*SULTR1.2* and *MOT1.1*).

In total, we identified 20 groups of secondary metabolite transporters in our

dataset. Compared to NM control roots, 29 of these transporters were up- and 43 were

downregulated in AM roots (**Fig. 7A**). Several secondary metabolite transporters were

differentially regulated in Ga165 and Ri09-colonized roots. Among these, most belonged

to six gene annotation groups, including ATP-binding cassette (ABC), amino acid, major

facilitator superfamily (MFS), major intrinsic protein (MIP), peptide, and Sugars Will

Eventually be Exported Transporters (SWEETs) (**Fig. 7B**). Here, we will only discuss

those transporters that are known to play a role in AM symbiosis, or that were strongly

differentially regulated by mycorrhizal colonization, but have not yet been functionally

characterized.

Two white-brown-complex ABC transporter family proteins (*STR* and *STR2*; Zhang

et al., 2010), three ABC transporter B family proteins (*AMN2*, *AMN3* [Roy, 2015]*, and

Medtr4g124040), and one drug resistance transporter-like ABC domain protein

(*Medtr1g050525*) were strongly upregulated in AM roots, particularly in Ri09-colonized

roots (**Fig. 7C**). The expression levels in Ga165-colonized roots were higher but did not

always differ significantly from the NM roots.

Two amino acid transporters, *Medtr1g030660* and *Medtr2g101920*, and two

peptide transporters, *PTR3* and *OPT15*, were upregulated in mycorrhizal roots (**Fig. 7D**).

As expected, *NOPE1,* an N-acetylglucosamine transporter of the major facilitator

superfamily (MFS) that is required for AM symbiosis (Nadal et al. 2017), was also highly

upregulated in AM roots (**Fig. 7E**). Three major intrinsic protein (MIP) transporters,

NIP1;5, *NIP1;6*, and *NIP3;1*, which are often classified as aquaporins, were all

upregulated in mycorrhizal roots. *NIP1;5* was particularly highly expressed in Ri09-

colonized roots, while *NIP1;6*, *NIP3;1* and *NIP6;2* showed higher expression levels in

Ga165-colonized roots.

Figure 7. Expression patterns of significantly differentially expressed secondary metabolite transporters in the roots of *Medicago truncatula* after colonization with different AM fungi. **A**, Venn diagrams showing the number of secondary metabolite transporters that are significantly up- (yellow) or down-regulated (blue) in roots colonized by *Glomus aggregatum* 165 (Ga165) or *Rhizophagus irregularis* 09 (Ri09), or both. **B**, summary of the number of genes from specific gene annotation groups of secondary metabolite transports that are up- or down-regulated in roots colonized by Ga165 or Ri09, or both (see **Supplemental File 7**). Expression patterns of genes annotated as **C**, ATP-binding cassette (ABC); **D**, amino acid or peptide; **E**, major facilitator superfamily (MFS) or major intrinsic protein (MIP); and **F**, SWEET transporters in the roots of non-mycorrhizal plants (NM; n = 3) and plants inoculated with Ga165 (n =3) or Ri09 (n = 4). The bars in C-F represent the mean of each treatment ± SEM (data points for biological replicates are displayed in **Supplemental Figure S9**). Expression levels are shown as fragments per kilobase of transcript per million mapped reads (FPKM); only genes with FPKM ≥ 10 are shown; significant differences (q-value < 0.05) in FPKM values between treatments were determined as part of the CuffDiff2 analysis and are indicated using different letters on the bars. Abbreviations in B: G3P, glycerol-3-phosphate; NAT, nucleobase-ascorbate transporter; OST, organic solute transporter; TPT, triose-phosphate transporter; TRAPP, transport protein particle.

SWEETs are known to play a crucial role in AM symbiosis (Doidy et al. 2019), and

five were differentially expressed in AM roots. *SWEET1.2* was upregulated by

colonization with either fungus, but especially with Ri09. *SWEET3.3* and *SWEET12,* were

only upregulated in Ri09-colonized roots, whereas *SWEET7* and *SWEET13* were only

upregulated in Ga165-colonized roots (**Fig. 7F**). The variation in the expression patterns

of these five *SWEET* genes suggests that sugar transport is strongly regulated in

response to fungal benefit.

Photosynthesis-related genes in the shoots are upregulated in response to

colonization with *Rhizophagus irregularis* 09

GO enrichment analysis on several gene clusters shown in **Fig. 2F** revealed that cellular

components and biological processes associated with photosynthesis were enriched in

the shoots of Ri09-colonized plants (**Supplemental Fig. S10A**). The associated genes

were annotated as follows: *Medtr1g115410*, photosystem II reaction center family protein

(orthologous to PsbP in *Arabidopsis thaliana*); *Medtr2g082580*, oxygen-evolving

enhancer protein (orthologous to PsbQ-like protein 2 in *A. thaliana*); *Medtr5g018670*,

photosystem II oxygen-evolving enhancer protein; *Medtr1g015290*, ultraviolet-B-

repressible protein (orthologous to PsbX in *A. thaliana*); and *Medtr1588s0010*, ATP

synthase F1, gamma subunit. The expression levels of *PsbP*, *PsbQ-like protein 2*, *PsbX*,

and the ATP synthase gene were significantly higher in the shoots of Ri09-colonized

plants than in NM plants (**Fig. 8A**). Out of these genes, only *PsbP* was also significantly

upregulated in Ga165-colonized plants. By contrast, *Medtr5g018670* was more highly

expressed in the shoots of Ga165-colonized than in NM plants. Overall, these expression

Figure 8. Expression patterns of significantly differentially expressed genes in the shoots of *Medicago truncatula* colonized by different AM fungi. Expression levels of genes involved in **A**, photosynthesis; **B**, response to biotic stress; and **C**, response to abiotic stress in non-mycorrhizal plants (NM; n=4) or plants colonized by *Glomus aggregatum* 165 (Ga165, n=4) or *Rhizophagus irregularis* 09 (Ri09, n=3). Bars represent the mean of each treatment ± SEM (data points for biological replicates are displayed in **Supplemental Figure S11**). Expression levels are shown as fragments per kilobase of transcript per million mapped reads (FPKM), and significant differences (q-value < 0.05) in FPKM values between treatments were determined as part of the CuffDiff2 analysis and are indicated using different letters above the bars (see also **Supplemental Table S4**).

patterns imply that photosynthesis was more strongly induced in the shoots of Ri09-colonized plants, which could partially explain the increased biomass of Ri09-colonized plants.

Biotic and abiotic stress-related genes are differentially regulated in the shoots of *Glomus aggregatum* and *Rhizophagus irregularis* colonized plants

Many of the GO terms that were enriched in the shoots of Ga165-colonized plants encompassed biological processes related to biotic stress (**Supplemental Fig. S10B**). The genes linked to these GO terms were annotated as: *Medtr4g054920*, cytochrome P450 family 94 protein; *Medtr4g120760*, pathogenesis-related protein bet V I family protein; *Medtr8g096900,* pathogenesis-related thaumatin family protein; *Medtr4g092010*, (3S)-linalool/(E)-nerolidol/(E,E)-geranyl linalool synthase; and *Medtr5g053950*, allene oxide cyclase. Enrichment of these GO terms in the shoots of Ga165-colonized plants could indicate that the host plant is responding to Ga165 as it would to a pathogen by upregulating defense responses, or that Ga165 stimulates host plant defenses against pathogens (**Fig. 8B**).

In contrast, Ri09 colonization led to the downregulation of genes (**Fig. 3F**, cluster D), associated with biotic and abiotic stress (**Supplemental Fig. S10C**). The genes included in these GO terms were annotated as: *Medtr1g048990*, superoxide dismutase; *Medtr7g069980*, ferritin; *Medtr8g059170*, NAC transcription factor-like protein; *Medtr7g079180*, late embryogenesis abundant protein; and *Medtr3g087590*, myo-inositol 1-phosphate synthase. The expression of *Medtr7g079180* was upregulated by Ga165, but the remaining abiotic stress response genes were more highly expressed in NM or

Ga165-colonized plants than in Ri09-colonized plants (**Fig. 8C**). Collectively, the

expression patterns of these abiotic stress response genes could indicate that Ri09 is

more effective at conferring abiotic stress tolerance to *M. truncatula* than Ga165.

DISCUSSION

The growth and nutritional benefits of arbuscular mycorrhizal fungi are linked to

mycorrhiza-induced transporter expression

When compatible host plants are inoculated with AM fungi, colonization most often leads

to increased growth and nutrient uptake, especially in nutrient-deprived conditions

(Chandrasekaran 2020). We found that under N and P limitation, Ga165 and Ri09

colonized *M. truncatula* equally well (**Fig. 1**), but still, Ri09 promoted greater growth and

provided more N and P to the host than Ga165 (**Fig. 2**). This observation is consistent

with both the expression patterns that we observed for mycorrhiza-induced N and P

transporters and with previous findings where we demonstrated that Ri09 is more

cooperative than Ga165 (Kiers et al. 2011; Fellbaum et al. 2014; Wang et al. 2016). For

example, the known AM-induced ammonium transporters *AMT2;3* and *AMT2;5* were both

upregulated by Ga165 and Ri09 (**Fig. 6C**). Although the expression levels of *AMT2;3* in

AM roots were lower than that of *AMT2;5*, *AMT2;3* expression was more strongly

upregulated by Ri09 than by Ga165 colonization. Similarly, we observed the same

expression pattern for the putative ammonium transporter *NIP1;5* (**Fig. 7E**), which has

not been functionally characterized in *M. truncatula* during AM symbiosis, but the soybean

ortholog *Nod26* plays a crucial role in ammonium transport in the rhizobia-legume

symbiosis (Frare et al. 2018). Interestingly, although the AM fungi were provided with

$^{15}NH_4Cl$ as a N source, we also found that the nitrate transporter *NPF4.12* was strongly

induced in AM roots, especially in Ri09-colonized roots (**Fig. 6D**). Aloui et al. (2018) found

that *NPF4.12* is enriched in the proteome of mycorrhizal *M. truncatula* roots, and recently

the rice ortholog *OsNPF4.5* was functionally characterized as an AM-induced nitrate

transporter (Wang et al. 2020). Overall, the stronger expression of ammonium and nitrate

transporters in Ri09- than in Ga165-colonized roots aligns with the observed increase in

the ^{15}N labeling in the shoots of Ri09-colonized plants.

The AM-induced P transporters *PT4* and *PT8* were both highly expressed in AM

colonized roots, especially Ri09-colonized roots (**Fig. 6E**). Similarly, colonization with

both fungi, but particularly with Ri09, caused an increase in shoot P content. However,

Medtr1g074930, a putative high affinity inorganic phosphate transporter, was strongly

upregulated in Ga165-colonized roots (**Fig. 6E**). This P transporter is upregulated during

P deficiency (Wang et al. 2017), and the upregulation of *Medtr1g074930* in our study

could indicate that colonization by Ga165 resulted in P deficiency in the host plant.

However, this is inconsistent with the observation that *Medtr1g074930* was not

upregulated in the roots of NM plants (**Fig. 6E**) and Ga165-colonized roots had a higher

P concentration than NM roots (**Supplemental Fig. S2**). Regardless, others have

similarly observed that the expression of high affinity P transporters are differentially

regulated in the host by colonization with different species of AM fungi (Grunwald et al.

2009).

Like N and P, AM fungi also play a critical role in supplying sulfur to the host plant

(Sieh et al. 2013). We observed an upregulation of two sulfate transporters, *SULTR1.1*

and *SULTR1.2* (**Fig. 6F**). Both are upregulated in mycorrhizal *M. truncatula* roots,

although not consistently (Casieri et al. 2012; Wipf et al. 2014). The *SULTR1.2* ortholog

in *Lotus japonicus* appears to play an important role in both direct and mycorrhizal sulfur

uptake pathways, and its promoter is active in arbuscule-containing cells (Giovannetti et

al. 2014). We also observed that the putative sulfate transporter gene *MOT1.1* was

upregulated in Ga165 and Ri09-colonized roots (**Fig. 6F**). However, the involvement of

MOT1.1 in sulfur transport has not been experimentally demonstrated yet, but it is

phylogenetically related to *MOT1.3*, which is a molybdate transporter crucial for the

rhizobia-legume symbiosis (Tejada-Jiménez et al. 2017).

Patterns of photosynthesis, carbon transporter, and lipid biosynthesis gene

expression in the host plant are linked to the nutritional benefits provided by the

fungal symbiont

In exchange for the nutritional benefits that AM fungi provide, plants transfer between 4

and 20% of their net fixed carbon in the form of carbohydrates and lipids to these fungal

symbionts (Wright et al. 1998; Roth and Paszkowski 2017). Both the nutritional benefits

that AM fungi provide and the increased carbon sink they create are known to stimulate

an increase in photosynthesis (Kaschuk et al. 2009). In line with this, we observed that

the expression levels of photosynthesis-related genes were more strongly induced in

Ri09- than in Ga165-colonized plants (**Fig. 8A**); thus, the elevated growth response

provided by Ri09 (**Fig. 2A-B**) was likely not only due to an increased nutrient transfer, but

potentially also due to a stimulation in photosynthesis.

Our previous studies have shown that host plants preferentially allocate more

carbon to the more cooperative AM fungus Ri09 (Kiers et al. 2011). Yet, we found that

the putative lipid transporters *STR* and *STR2* were upregulated in AM roots, but did not

differ significantly between Ga165 and Ri09-colonized roots (**Fig. 7C**). Nevertheless, we

did observe that the expression levels of the lipid biosynthesis gene *RAM2* (**Fig. 5**) and

the sugar transporter *SWEET1.2* (**Fig. 5F**) were higher in Ri09-colonized roots, which is

suggestive of increased transport of carbon resources to the fungus in exchange for

elevated nutritional benefits (Bravo et al. 2017; An et al. 2019).

Although *SWEET1.2* is highly upregulated during AM colonization, particularly in

arbusculated cortical cells, it is not essential for the maintenance of the AM symbiosis,

most likely because of functional redundancy between SWEET1.2 and other SWEETs

(An et al. 2019). We found that *SWEET3.3* and *SWEET12* were more strongly

upregulated in Ri09-colonized roots (**Fig. 7F**); both are upregulated in arbusculated

cortical cells in *M. truncatula* (Sameeullah et al. 2016). Interestingly, *SWEET7* and

SWEET13 were more strongly induced in Ga165-colonized roots, but SWEETs perform

diverse physiological functions, and their upregulation does not necessarily increase

sugar transport to the fungus. For example, some SWEET transporters in *Arabidopsis*

localize to the tonoplast and restrict sugar efflux from the root (Chen et al. 2015). Given

the variable expression patterns of these five SWEETs in Ri09 and Ga165-colonized

roots and the varying roles they may play in sugar efflux and retention, functionally

characterizing them will improve our understanding of the mechanisms controlling carbon

allocation during AM symbioses.

Arbuscular mycorrhiza-induced changes in the expression of host abiotic and biotic stress response genes are species specific

Although the primary focus of this study was to evaluate both resource exchange and transporter expression, our gene expression data also provided some unique insights into biotic and abiotic stress responses. Mycorrhizal growth responses are context-dependent and fall along a mutualism to parasitism continuum (Johnson and Graham 2013). In natural environments, plants are colonized by communities of AM fungi, which supports the idea of functional complementarity, where a host plant experiences a broad range of benefits that are provided by specific members of the AM fungal community (*e.g.*, enhanced nutrient availability or increased disease resistance; Jansa et al., 2008). While Ga165 does not seem to provide sufficient nutritional benefits to elicit a positive growth response, it is possible that like other AM fungal species it may provide resistance to one or more abiotic stresses (*e.g.*, salinity or heavy metal accumulation), none of which were evaluated in this study.

We found that Ga165 more strongly up-regulated defense-response genes in the shoots (**Fig. 8B**), two of which (*Medtr4g120760* and *Medtr8g096900*) are upregulated in alfalfa when challenged with the foliar fungal pathogen *Phoma medicaginis* (Li et al. 2019a) and pea aphids (*Acyrthosiphon pisum*; Li et al., 2019b). There are two potential explanations for this observation. The plant could show similar responses to a low benefit AM fungus than to known fungal pathogens, or Ga165 could be more effective than Ri09 at priming host defense responses and thereby prevent infection from pathogens and mitigating arthropod attack as previously been described for other AM fungal species (Liu et al. 2007; Sharma et al. 2017).

Similarly, we also observed that genes associated with abiotic stress were

upregulated in Ga165-colonized roots (**Supplemental Fig. S6A)**. Three of these genes

— *Medtr5g063670*, *Medtr7g093170*, and *Medtr8g026960* — were very strongly

upregulated and are annotated as annexin D8, seed maturation protein, and homeobox

associated leucine zipper protein, respectively. Although the precise role of annexin D8

is not known, other annexins are differentially regulated during various abiotic and biotic

stresses, and some seem to play a role during AM symbiosis (Roux et al. 2012). The

seed maturation protein *Medtr7g093170* has been characterized as the late

embryogenesis abundant protein LEA1370, which is upregulated under various abiotic

stresses, including abscisic acid treatment, cold, salinity and dehydration (Zhang et al.

2020). Finally, the homeobox associated leucine zipper protein encoded by

Medtr8g026960 was characterized as a transcription factor that is upregulated during salt

stress in alfalfa (Postnikova et al. 2013). However, since these genes are not yet

functionally characterized, the question whether Ga165 colonization could contribute to

an alleviation of these abiotic stresses, or whether host plants respond similarly to abiotic

stress and to Ga165 colonization, requires further studies.

It is also possible that Ga165 truly is a low-benefit symbiont capable of persisting

in a community with high benefit AM fungi thereby avoiding both detection and

subsequent sanctioning by the host plant for poor performance (Hart et al. 2013).

Avoiding detection could be carried out using effectors, as has been described in the

model AM fungus *R. irregularis* DAOM197198 that counteracts the plant immune program

with the small, secreted peptide SP7 (Kloppholz et al. 2011). Indeed, in our root

expression data we observed that 11 defense response genes, including three defensins

(*Medtr8g012810, Medtr8g012815*, and *Medtr8g012845*) were strongly up-regulated in

Ri09, but not in Ga165-colonized roots (**Supplemental Fig. S6B**, **Supplemental Table**

S5). Defensins are known to inhibit pathogen growth (Maróti et al. 2015), so it is tempting

to speculate that Ga165 could temper host defense responses to avoid detection and

sanctioning. However, the defensin *DefMd1* when overexpressed upregulates the

expression of the GRAS transcription factor *RAM1*, which plays a critical role in

arbuscular branching (Park et al., 2015; Uhe et al., 2018). *RAM1* was highly upregulated

in both Ga165 and Ri09 colonized roots (**Supplemental Table S3**) suggesting that

defensins might play a role in AM symbiosis.

Transporter expression patterns in *Medicago truncatula* roots reveal interesting

candidates for future functional characterization

This study has provided novel insights into the physiological and transcriptomic response

of a host plant to colonization by both high and low benefit AM fungi and **Fig. 9** provides

an overview of these responses. The experimental approach allowed us to identify both

the molecular mechanisms controlling the observed physiological responses and

transporters that are both commonly and uniquely regulated by high (Ri09) and low

benefit (Ga165) AM fungal species. While the primary focus of our study was on N, P,

and carbon transporters in the roots of *M. truncatula* colonized by Ga165 or Ri09, our

analysis of differentially expressed transporters uncovered some interesting gene

candidates with unique expression patterns (**Figs. 6** and **7**). Although there was a core

set of transporters similarly induced by both fungi, a smaller set were expressed

differently, and many are prime candidates for future functional analysis.

Figure 9. Summary of the physiological and transcriptomic responses in roots and shoots of *Medicago truncatula* colonized by either *Glomus aggregatum* 165 or *Rhizophagus irregularis* 09. Physiological data include the analysis of AM colonization, plant biomass, and both phosphorus and nitrogen-15 contents in plant tissues. Transcriptomic data include the expression patterns of nutrient transporters and enrichment of gene ontology terms. Yellow up-arrows indicate an increase, equal signs (=) specify no change, and blue down-arrows designate a decrease in each response compared to non-mycorrhizal plants. Arrow size indicates the degree of the response.

Multiple ABC transporters were induced by AM colonization (**Fig. 7C**), and two of

these, *STR* and *STR2*, have already been studied extensively (Zhang et al., 2010). Two

other transporters, *AMN2* and *AMN3*, are also well studied, but the substrate they

transport remains elusive (Roy et al. 2021). The two remaining genes, *Medtr4g124040*

and *Medtr1g050525*, have not been functionally characterized and could represent

interesting targets for future studies. The amino acid transporters *Medtr1g030660* and

Medtr2g101920 and the peptide transporters *PTR3* and *OPT15* were all upregulated in

mycorrhizal roots. None of these have been functionally characterized, but the protein

product of *Medtr2g101920* was isolated from the plasma membrane of AM roots (Aloui et

al. 2018). Future characterization of these transporters will provide greater insight into the

extensive nutrient exchange mechanisms that support the AM symbiosis.

In conclusion, the comparison of the physiological and transcriptomic response of *M.*

truncatula to colonization with two AM fungal species that provide different nutritional

benefits to the host allowed us to identify some of the molecular mechanisms that control

reciprocal nutrient exchange. These include specific N and P transporters that have

already been functionally characterized (e.g., *AMT2;3*, *AMT2;5*, *NPF4.12*, *PT4,* and *PT8*)

and others that have not (e.g., NIP1;5). In addition, this study uncovered differential

expression of genes putatively involved in the allocation of carbon resources to AM fungi

depending on their nutritional benefit (e.g., *RAM2, SWEET1.2, SWEET3.3 and SWEET12*

were upregulated in Ri09-colonized roots, while *SWEET7* and *SWEET13* were

upregulated in Ga165-colonized roots). Root colonization with diverse fungal

communities that are composed of both low and high benefit fungi, allows the host plant

to choose among multiple trading partners, but also to take advantage of the diverse

benefits that fungi within this community can provide. The differential gene expression in

the roots and shoots after colonization with AM fungi that differ in the symbiotic benefit

that they provide, indicates that host plants are able to fine tune their response to different

fungi in their community.

MATERIALS AND METHODS

Biological materials and growth conditions

M. truncatula 'Jemalong A17' seeds were acid scarified in 36N H_2SO_4 (Fisher Scientific

Inc., Waltham, MA, USA), surface sterilized with 8.25% sodium hypochlorite (Clorox®

bleach, The Clorox Company, Oakland, CA, USA), rinsed with sterile Type 1 water and

imbibed at 4°C overnight. The seeds were then transferred onto moist autoclaved filter

paper in Petri dishes and kept in the dark for 3 d. Finally, the Petri dishes were placed in

ambient conditions for 7 d. Fully germinated seedlings were transferred into the root

compartment (RC) of a custom-made, two-compartment pot measuring 12 cm x 8 cm x 8

cm (L x W x H; **Supplemental Fig. S1**). The RC was separated from the hyphal

compartment (HC) by a 0.1 cm-thick plastic divider sealed on all sides by silicone

(Aqueon, Franklin, WI, USA). A hole in the middle of the divider (~3.12 cm in diameter)

was covered on both sides with a sheet of fine nylon mesh with 50 µm pores. Between

these sheets was a coarse nylon mesh with 1000 µm pores to create an air gap and inhibit

mass flow between the RC and the HC. This allowed fungal hyphae, but not roots, to

crossover from the RC to the HC (**Supplemental Fig. S1**). Both compartments were filled

with 250 ml of sterilized soil substrate containing 40% sand, 20% perlite, 20% vermiculite,

and 20% soil by volume. The soil substrate had an original concentration of 4.81 mg kg^{-1}

of Olsen's extractable phosphate, 10 mg kg^{-1} of NH_4^+, 34.4 mg kg^{-1} of NO_3^-, and a pH of

8.26. These nutrient levels were sufficiently low to induce host demand for N and P and

to stimulate AM colonization.

Fungal inoculum for *Rhizophagus irregularis* isolate 09 (Ri09; collected from

Southwest Spain by Mycovitro S.L. Biotechnología ecológica, Granada, Spain) and

Glomus aggregatum isolate 165 (Ga165; collected from the Long Term Mycorrhizal

Research Site, University of Guelph, Canada) were produced using axenic root organ

cultures of Ri T-DNA transformed carrot (*Daucus carota* clone DCI) grown on minimal

medium (St-Arnaud et al. 1996). After approximately eight weeks of growth, spores were

isolated by blending the medium in 10 mM citrate buffer (pH 6.0). Then, at transplanting,

each seedling was inoculated with ~0.4 g mycorrhizal roots and ~500 spores of either

Ri09 or Ga165. Non-mycorrhizal (NM) controls received a similar quantity of double

autoclaved roots and spores. After transplanting and fungal inoculation, the plants were

grown in a growth chamber (model TC30; Conviron, Winnipeg, MB, Canada) with a 25°C

day/20°C night cycle, 30% relative humidity, and a photosynthetic photon flux of ~225

μmol m^{-2} s^{-1}.

Experimental design

567 Throughout the experiment, the position of the plants within the growth chamber was

568 randomized three times. Five weeks post-inoculation, 4 mM $^{15}NH_4Cl$ (Sigma Aldrich, St.

569 Louis, USA) and 0.5 mM KH_2PO_4 were added to the HC of colonized and noncolonized

570 plants in a modified Ingestad solution (Ingestad 1960). The plants were harvested two

571 weeks later (seven weeks post-inoculation).

Biomass determination, mycorrhizal quantification, and analysis of nitrogen and phosphorus contents in plants

At plant harvest, shoots and roots were separated to determine their respective fresh weight. Subsamples of both tissues were flash frozen in liquid N_2 and stored at -80°C for transcriptome analysis. An additional root subsample was collected and stored in 50% ethanol at 4°C to assess AM colonization. The remaining plant tissue was dried at 70°C for 48 h and then shoot and root dry weights were recorded. To determine the level of AM colonization, the root samples were cleared with 10% KOH at 80°C for 30 min, rinsed with tap water, and stained with 5% Sheaffer ink-vinegar (v:v) at 80°C for 15 min (Vierheilig et al. 1998). Total AM colonization rates were assessed using the gridline intersection method (McGonigle et al. 1990). We did not observe any fungal structures in the roots of NC control plants. Subsamples of dried shoot and root tissues were pulverized using a tissue homogenizer (Precellys 24 Dual, Cayman Chemical Company, Ann Arbor, MI, USA) and digested with 2N HCl for 2 h at 95°C. The P concentration in the plant tissues was spectrophotometrically determined at 436 nm after adding ammonium molybdate vanadate solution (Fisher Scientific, Pittsburgh, USA). Measurements of nitrogen-15 in plant tissues were performed by quantitative NMR spectroscopy as described previously (Kafle et al. 2018).

Fluorescent staining and confocal analysis of arbuscules

Ink-stained roots that contained arbuscules were subsequently stained for confocal imaging as described in Cope et al. (2019). Briefly, they were incubated overnight at 4°C in 1X PBS containing 2 µg ml^{-1} Alexa Fluor 488^{TM} conjugated to wheat germ agglutinin

(WGA; Thermo Fischer Scientific, Waltham, MA, USA). The next day, stained roots were

rinsed with 1X PBS, mounted on a glass slide, and immediately observed on an Olympus

FV1200 confocal laser scanning microscope (Olympus, Shinjuku, Tokyo, Japan) with an

Olympus UPlanApo 40×/0.85 objective. The emission spectra of Alexa Fluor 488^{TM} WGA

(500 to 540 nm) was detected after excitation with a 488-nm laser. Other settings used in

the Fluoview v4.2c acquisition software (*e.g.*, laser intensity, gain, offset, magnification,

airy units) were similar for all sample observations. The width of 488 and 508 arbuscules

from 22 Ga165 and 21 Ri09-colonized root segments, respectively, were measured using

Fiji (Schindelin et al. 2012).

RNA sequencing and analysis

Root and shoot tissues from four biological replicates (n = 4) were homogenized in liquid

N2, and total RNA was extracted using the PureLinkTM RNA Mini Kit (Thermo Fisher

Scientific). Extracted RNA was treated with TURBO™ DNase (Thermo Fisher Scientific)

and quantified using a NanoDrop ND-2000 spectrophotometer (Thermo Fisher Scientific).

RNA quality was assessed using 2100 BioAnalyzer technology (Agilent Technologies).

Approximately 100 ng of total RNA was used to construct poly(A) selection libraries using

the Illumina TruSeq RNA Sample Preparation kit and 150-nucleotide single-end reads

were sequenced using an Illumina NextSeq500 with high output flow cells (Genomics

Sequencing Facility, South Dakota State University).

The RNA sequencing data were uploaded to Discovery Environment

(https://de.cyverse.org/de) and read quality of each fastq file was evaluated using FastQC

0.11.5 (https://www.bioinformatics.babraham.ac.uk/projects/fastqc/). One Ga165 and

one NC root sample and one Ri09 shoot sample were discarded due to poor quality. The

remaining root (n=3, Ga165 and NC; n=4, Ri09) and shoot (n=4, Ga165 and NC; n=3,

Ri09) samples were aligned to the reference genome of *M. truncatula* 'Jemalong A17'

(JCVI Mt4.0; Tang et al., 2014) using HISAT2.1 (Kim et al. 2019). Aligned reads were

then assembled and merged using StringTie-1.3.3 (Pertea et al. 2015) and the *M.*

truncatula JCVI 4.0v2 annotation file (Tang et al. 2014). Finally, Cuffdiff2.1.1 (Trapnell et

al. 2013) was used to quantify transcript abundance. The output files from CuffDiff2.1.1

were visualized in R v3.6.2 (The R Foundation 2020) using the CummeRbund package

(Goff et al. 2012). For further analyses, the differential gene expression output files from

CuffDiff2.1.1 were imported into Microsoft Excel (Microsoft 365®) and sorted to identify

significantly differentially expressed genes (DEGs) with a log2 fold-change > 2 and a false

discovery rate-adjusted q-value < 0.05. All DEGs outside this threshold were not

considered for further analysis. Venn diagrams of DEGs were generated using the

Bioinformatics & Evolutionary Genomics Web tool

(http://bioinformatics.psb.ugent.be/webtools/Venn/) and heatmaps of FPKM (Fragments

Per Kilobase of transcript per Million mapped reads) values from all DEGs across

inoculation regimes were generated using Cluster 3.0 and Java TreeView

(http://bonsai.hgc.jp/~mdehoon/software/cluster/software.htm). Gene ontology (GO)

enrichment analysis was conducted with the web-based tool PlantTFDB 4.0

(http://plantregmap.gao-lab.org/go.php).

Statistical analysis

For the physiological response variables, unless mentioned otherwise, mean and standard error of the mean were determined from five independent biological replicates for each colonization regime, including noncolonized and either Ga165 or Ri09-colonized plants. One-way ANOVA was used to identify statistically significant differences ($p \leq 0.05$) among all three colonization regimes for dry weight, P concentration, P content, and nitrogen-15 enrichment. The least significant difference (LSD) test was subsequently used for pairwise multiple comparisons to identify significant differences ($p \leq 0.05$) among treatments. These statistical analyses were conducted using Statistix 9 (analytical software, Tallahassee, FL, USA). For AM colonization and arbuscule size, statistically significant differences ($p \leq 0.05$) between the Ga165 and Ri09 mycorrhizal treatments were determined using a Welch's two-sample t-test in R v4.0.2 (The R Foundation 2020). Only results that are significant at the $p \leq 0.05$-level are primarily discussed. All F statistics and p-values are shown in **Supplemental Table S1**.

ACKNOWLEDGEMENTS

This work was funded by the USDA (2017-67014-26530), the SD Soybean Research and Promotion Council, and the Agricultural Experiment Station at SDSU to HK and SS. KG acknowledges support of the North Carolina Agriculture Research Service (NCARS), the North Carolina Soybean Producers Association (2019-1656), and the USDA (2020-67013-31800). We would like to thank Lindsay McKeever for performing the Kjeldahl degradations for the $^{15}N/^{14}N$ analyses.

LITERATURE CITED

Aloui, A., Recorbet, G., Lemaître-Guillier, C., Mounier, A., Balliau, T., Zivy, M., Wipf, D.,

and Dumas-Gaudot, E. 2018. The plasma membrane proteome of *Medicago*

truncatula roots as modified by arbuscular mycorrhizal symbiosis. Mycorrhiza.

28:1–16

An, J., Zeng, T., Ji, C., de Graaf, S., Zheng, Z., Xiao, T. T., Deng, X., Xiao, S., Bisseling,

T., Limpens, E., and Pan, Z. 2019. A *Medicago truncatula* SWEET transporter

implicated in arbuscule maintenance during arbuscular mycorrhizal symbiosis. New

Phytol. 224:396–408

Besserer, A., Becard, G., Jauneau, A., Roux, C., Sejalon-Delmas, N., Bécard, G.,

Jauneau, A., Roux, C., and Séjalon-Delmas, N. 2008. GR24, a synthetic analog of

strigolactones, stimulates the mitosis and growth of the abuscular mycorrhizal

fungus *Gigaspora rosea* by boosting its energy metabolism. Plant Physiol.

148:402–413

Bravo, A., Brands, M., Wewer, V., Dörmann, P., and Harrison, M. J. 2017. Arbuscular

mycorrhiza-specific enzymes FatM and RAM2 fine-tune lipid biosynthesis to

promote development of arbuscular mycorrhiza. New Phytol. 214:1631–1645

Breuillin-Sessoms, F., Floss, D. S., Karen Gomez, S., Pumplin, N., Ding, Y., Levesque-

Tremblay, V., Noar, R. D., Daniels, D. A., Bravo, A., Eaglesham, J. B., Benedito, V.

A., Udvardi, M. K., Harrison, M. J., Gomez, S. K., Pumplin, N., Ding, Y., Levesque-

Tremblay, V., Noar, R. D., Daniels, D. A., Bravo, A., Eaglesham, J. B., Benedito, V.

A., Udvardi, M. K., Harrison, M. J., Karen Gomez, S., Pumplin, N., Ding, Y.,

Levesque-Tremblay, V., Noar, R. D., Daniels, D. A., Bravo, A., Eaglesham, J. B.,

Benedito, V. A., Udvardi, M. K., and Harrison, M. J. 2015. Suppression of arbuscule

degeneration in *Medicago truncatula* phosphate transporter4 mutants is dependent

on the ammonium transporter 2 family protein AMT2;3. Plant Cell. 27:1352–1366

Brundrett, M. C., and Tedersoo, L. 2018. Evolutionary history of mycorrhizal symbioses

and global host plant diversity. New Phytol. 220:1108–1115

Bücking, H., and Kafle, A. 2015. Role of arbuscular mycorrhizal fungi in the nitrogen

uptake of plants: Current knowledge and research gaps. Agronomy. 5:587–612

Casieri, L., Gallardo, K., and Wipf, D. 2012. Transcriptional response of *Medicago*

truncatula sulphate transporters to arbuscular mycorrhizal symbiosis with and

without sulphur stress. Planta. 235:1431–1447

Chandrasekaran, M. 2020. A meta-analytical approach on arbuscular mycorrhizal fungi

inoculation efficiency on plant growth and nutrient uptake. Agriculture. 10:370

Chen, H. Y., Huh, J. H., Yu, Y. C., Ho, L. H., Chen, L. Q., Tholl, D., Frommer, W. B.,

and Guo, W. J. 2015. The Arabidopsis vacuolar sugar transporter SWEET2 limits

carbon sequestration from roots and restricts *Pythium* infection. Plant J. 83:1046–

1058

Cope, K. R., Bascaules, A., Irving, T. B., Venkateshwaran, M., Maeda, J., Garcia, K.,

Rush, T. A., Ma, C., Labbé, J., Jawdy, S., Steigerwald, E., Setzke, J., Fung, E.,

Schnell, K. G., Wang, Y., Schleif, N., Bücking, H., Strauss, S. H., Maillet, F.,

Jargeat, P., Bécard, G., Puech-Pagès, V., and Ané, J.-M. M. 2019. The

ectomycorrhizal fungus *Laccaria bicolor* produces lipochitooligosaccharides and

uses the common symbiosis pathway to colonize populus roots. Plant Cell.

31:2386–2410

Davière, J. M., and Achard, P. 2013. Gibberellin signaling in plants. Development.

140:1147–1151

Doidy, J., Vidal, U., and Lemoine, R. 2019. Sugar transporters in Fabaceae, featuring

SUT MST and SWEET families of the model plant *Medicago truncatula* and the

agricultural crop *Pisum sativum* J.J. Zwiazek, ed. PLoS One. 14:e0223173

Fellbaum, C. R., Mensah, J. A., Cloos, A. J., Strahan, G. E., Pfeffer, P. E., Kiers, E. T.,

and Bücking, H. 2014. Fungal nutrient allocation in common mycorrhizal networks

is regulated by the carbon source strength of individual host plants. New Phytol.

203:646–656

Floss, D. S., Levy, J. G., Lévesque-Tremblay, V., Pumplin, N., and Harrison, M. J. 2013.

DELLA proteins regulate arbuscule formation in arbuscular mycorrhizal symbiosis.

Proc. Natl. Acad. Sci. U. S. A. 110:E5025-34

Frare, R., Ayub, N., Alleva, K., and Soto, G. 2018. The ammonium channel NOD26 is

the evolutionary innovation that drives the emergence, consolidation, and

dissemination of nitrogen-fixing symbiosis in angiosperms. J. Mol. Evol. 86:554–

565

Garcia, K., Doidy, J., Zimmermann, S. D., Wipf, D., and Courty, P. E. 2016. Take a trip

through the plant and fungal transportome of mycorrhiza. Trends Plant Sci. 21:937–

950

Genre, A., Chabaud, M., Balzergue, C., Puech-Pagès, V., Novero, M., Rey, T.,

Fournier, J., Rochange, S., Bécard, G., Bonfante, P., and Barker, D. G. 2013.

Short-chain chitin oligomers from arbuscular mycorrhizal fungi trigger nuclear Ca^{2+}

spiking in *Medicago truncatula* roots and their production is enhanced by

732  strigolactone. New Phytol. 198:190–202

733 Giovannetti, M., Tolosano, M., Volpe, V., Kopriva, S., and Bonfante, P. 2014.

734  Identification and functional characterization of a sulfate transporter induced by

735  both sulfur starvation and mycorrhiza formation in *Lotus japonicus*. New Phytol.

736  204:609–619

737 Goff, L. A., Trapnell, C., and Kelley, D. 2012. CummeRbund: visualization and

738  exploration of Cufflinks high-throughput sequencing data. R Packag. version. 2:1–

739  45

740 Grunwald, U., Guo, W., Fischer, K., Isayenkov, S., Ludwig-Müller, J., Hause, B., Yan,

741  X., Küster, H., and Franken, P. 2009. Overlapping expression patterns and

742  differential transcript levels of phosphate transporter genes in arbuscular

743  mycorrhizal, Pi-fertilised and phytohormone-treated *Medicago truncatula* roots.

744  Planta. 229:1023–1034

745 Hart, M. M., Forsythe, J., Oshowski, B., Bücking, H., Jansa, J., and Kiers, E. T. 2013.

746  Hiding in a crowd - Does diversity facilitate persistence of a low-quality fungal

747  partner in the mycorrhizal symbiosis? Symbiosis. 59:47–56

748 Havlin, J. L., Beaton, J. D., Tisdale, S. L., and Nelson, W. L. 2014. *Soil Fertility and*

749  *Fertilizers: An Introduction to Nutrient Management*. Eighth. Pearson Education,

750  Inc.

751 Ingestad, T. 1960. Studies on the nutrition of forest tree seedlings. III mineral nutrition of

752  pine. Physiol. Plant. 13:513–533

753 Jansa, J., Smith, F. A., and Smith, S. E. 2008. Are there benefits of simultaneous root

754  colonization by different arbuscular mycorrhizal fungi? New Phytol. 177:779–789

Javot, H., Penmetsa, R. V., Terzaghi, N., Cook, D. R., and Harrison, M. J. 2007. A

Medicago truncatula phosphate transporter indispensable for the arbuscular

mycorrhizal symbiosis. Proc. Natl. Acad. Sci. 104:1720–1725

Johnson, N. C., and Graham, J. H. 2013. The continuum concept remains a useful

framework for studying mycorrhizal functioning. Plant Soil. 363:411–419

Kafle, A., Cope, K. R., Raths, R., Yakha, J. K., Subramanian, S., Bücking, H., and

Garcia, K. 2019. Harnessing soil microbes to improve plant phosphate efficiency in

cropping systems. Agronomy. 9:127

Kafle, A., Garcia, K., Wang, X., Pfeffer, P. E., Strahan, G. D., and Bücking, H. 2018.

Nutrient demand and fungal access to resources control the carbon allocation to

the symbiotic partners in tripartite interactions of *Medicago truncatula*. Plant. Cell

Environ. :1–15

Kaschuk, G., Kuyper, T. W., Leffelaar, P. A., Hungria, M., and Giller, K. E. 2009. Are the

rates of photosynthesis stimulated by the carbon sink strength of rhizobial and

arbuscular mycorrhizal symbioses? Soil Biol. Biochem. 41:1233–1244

Kiers, E. T., Duhamel, M., Beesetty, Y., Mensah, J. a, Franken, O., Verbruggen, E.,

Fellbaum, C. R., Kowalchuk, G. A., Hart, M. M., Bago, A., Palmer, T. M., West, S.

a, Vandenkoornhuyse, P., Jansa, J., Bücking, H., and Bucking, H. 2011. Reciprocal

rewards stabilize cooperation in the mycorrhizal symbiosis. Science (80-.).

333:880–882

Kim, D., Paggi, J. M., Park, C., Bennett, C., and Salzberg, S. L. 2019. Graph-based

genome alignment and genotyping with HISAT2 and HISAT-genotype. Nat.

Biotechnol. 37:907–915

Kloppholz, S., Kuhn, H., and Requena, N. 2011. A secreted fungal effector of *Glomus*

intraradices promotes symbiotic biotrophy. Curr. Biol. 21:1204–1209

Kretzschmar, T., Kohlen, W., Sasse, J., Borghi, L., Schlegel, M., Bachelier, J. B.,

Reinhardt, D., Bours, R., Bouwmeester, H. J., and Martinoia, E. 2012. A petunia

ABC protein controls strigolactone-dependent symbiotic signalling and branching.

Nature. 483:341–344

LeBauer, D. S., and Treseder, K. K. 2008. Nitrogen limitation of net primary productivity

in terrestrial ecosystems is globally distributed. Ecology. 89:371–379

Li, Y., Duan, T., Nan, Z., and Li, Y. 2019a. Arbuscular mycorrhizal fungus alleviates

alfalfa leaf spots caused by *Phoma medicaginis* revealed by RNA-seq analysis. J.

Appl. Microbiol. :jam.14387

Li, Y., Nan, Z., and Duan, T. 2019b. *Rhizophagus intraradices* promotes alfalfa

(*Medicago sativaM*) defense against pea aphids (*Acyrthosiphon pisum*) revealed by

RNA-Seq analysis. Mycorrhiza. 29:623–635

Liu, J., Maldonado-Mendoza, I., Lopez-Meyer, M., Cheung, F., Town, C. D., and

Harrison, M. J. 2007. Arbuscular mycorrhizal symbiosis is accompanied by local

and systemic alterations in gene expression and an increase in disease resistance

in the shoots. Plant J. 50:529–544

MacLean, A. M., Bravo, A., and Harrison, M. J. 2017. Plant signaling and metabolic

pathways enabling arbuscular mycorrhizal symbiosis. Plant Cell. 29:2319–2335

Maillet, F., Poinsot, V., André, O., Puech-Pagés, V., Haouy, A., Gueunier, M., Cromer,

L., Giraudet, D., Formey, D., Niebel, A., Martinez, E. A., Driguez, H., Bécard, G.,

and Dénarié, J. 2011. Fungal lipochitooligosaccharide symbiotic signals in

arbuscular mycorrhiza. Nature. 469:58–64

Maróti, G., Downie, J. A., and Kondorosi, É. 2015. Plant cysteine-rich peptides that

inhibit pathogen growth and control rhizobial differentiation in legume nodules.

Curr. Opin. Plant Biol. 26:57–63

Martin, F. M., Uroz, S., and Barker, D. G. 2017. Ancestral alliances: Plant mutualistic

symbioses with fungi and bacteria. Science (80-.). 356:eaad4501

McGonigle, T. P., Miller, M. H., Evans, D. G., Fairchild, G. L., and Swan, J. A. 1990. A

new method which gives an objective measure of colonization of roots by vesicular-

arbuscular mycorrhizal fungi. New Phytol. 115:495–501

Nadal, M., Sawers, R., Naseem, S., Bassin, B., Kulicke, C., Sharman, A., An, G., An, K.,

Ahern, K. R., Romag, A., Brutnell, T. P., Gutjahr, C., Geldner, N., Roux, C.,

Martinoia, E., Konopka, J. B., and Paszkowski, U. 2017. An N-acetylglucosamine

transporter required for arbuscular mycorrhizal symbioses in rice and maize. Nat.

Plants. 3:1–7

Nouri, E., Surve, R., Bapaume, L., Stumpe, M., Chen, M., Zhang, Y., Ruyter-Spira, C.,

Bouwmeester, H., Glauser, G., Bruisson, S., and Reinhardt, D. 2021. Phosphate

Suppression of Arbuscular Mycorrhizal Symbiosis Involves Gibberellic Acid

Signaling. Plant Cell Physiol.

Park, H.-J., Floss, D. S., Levesque-Tremblay, V., Bravo, A., and Harrison, M. J. 2015.

Hyphal branching during arbuscule development requires RAM1. Plant Physiol.

169:pp.01155.2015

Pertea, M., Pertea, G. M., Antonescu, C. M., Chang, T. C., Mendell, J. T., and Salzberg,

S. L. 2015. StringTie enables improved reconstruction of a transcriptome from

RNA-seq reads. Nat. Biotechnol. 33:290–295

Pimprikar, P., and Gutjahr, C. 2018. Transcriptional Regulation of Arbuscular

Mycorrhiza Development. Plant Cell Physiol. 59:673–690

Postnikova, O. A., Shao, J., and Nemchinov, L. G. 2013. Analysis of the alfalfa root

transcriptome in response to salinity stress. Plant Cell Physiol. 54:1041–1055

Roth, R., and Paszkowski, U. 2017. Plant carbon nourishment of arbuscular mycorrhizal

fungi. Curr. Opin. Plant Biol. 39:50–56

Roux, S. J., Clark, G. B., Morgan, R. O., Fernandez, M.-P. P., and Roux, S. J. 2012.

Evolutionary adaptation of plant annexins has diversified their molecular structures,

interactions and functional roles. New Phytol. 196:695–712

Roy, S. 2015. Analysis of gene expression, regulation, and function of three symbiotic

ABC subfamily-B transporters in *Medicago truncatula*.

Roy, S., Breakspear, A., Cousins, D., Torres-Jerez, I., Jackson, K. J., Kumar, A., Su, Y.,

Krom, N., Liu, C.-W., Udvardi, M., Xu, P., and Murray, J. D. 2021. Three common

symbiotic ABC-B transporters in Medicago truncatula are regulated by a NIN-

independent branch of the symbiosis signalling pathway. Mol. Plant-Microbe

Interact.

Rush, T. A., Puech-Pagès, V., Bascaules, A., Jargeat, P., Maillet, F., Haouy, A., Maës,

A. Q. M., Carriel, C. C., Khokhani, D., Keller-Pearson, M., Tannous, J., Cope, K. R.,

Garcia, K., Maeda, J., Johnson, C., Kleven, B., Choudhury, Q. J., Labbé, J., Swift,

C., O'Malley, M. A., Bok, J. W., Cottaz, S., Fort, S., Poinsot, V., Sussman, M. R.,

Lefort, C., Nett, J., Keller, N. P., Bécard, G., and Ané, J. M. 2020. Lipo-

chitooligosaccharides as regulatory signals of fungal growth and development. Nat.

Commun. 11:1–10

Sameeullah, M., Demiral, T., Aslam, N., Baloch, F. S., and Gurel, E. 2016. In silico

functional analyses of SWEETs reveal cues for their role in AMF symbiosis. Pages

45–58 in: Plant, Soil and Microbes: Volume 2: Mechanisms and Molecular

Interactions, Springer International Publishing.

Schindelin, J., Arganda-Carreras, I., Frise, E., Kaynig, V., Longair, M., Pietzsch, T.,

Preibisch, S., Rueden, C., Saalfeld, S., Schmid, B., Tinevez, J. Y., White, D. J.,

Hartenstein, V., Eliceiri, K., Tomancak, P., and Cardona, A. 2012. Fiji: An open-

source platform for biological-image analysis. Nat. Methods. 9:676–682

Sharma, E., Anand, G., and Kapoor, R. 2017. Terpenoids in plant and arbuscular

mycorrhiza-reinforced defence against herbivorous insects. Ann. Bot. 119:791–801

Sieh, D., Watanabe, M., Devers, E. A., Brueckner, F., Hoefgen, R., and Krajinski, F.

2013. The arbuscular mycorrhizal symbiosis influences sulfur starvation responses

of Medicago truncatula. New Phytol. 197:606–616

Spatafora, J. W., Chang, Y., Benny, G. L., Lazarus, K., Smith, M. E., Berbee, M. L.,

Bonito, G., Corradi, N., Grigoriev, I., Gryganskyi, A., James, T. Y., O'Donnell, K.,

Roberson, R. W., Taylor, T. N., Uehling, J., Vilgalys, R., White, M. M., and Stajich,

J. E. 2016. A phylum-level phylogenetic classification of zygomycete fungi based

on genome-scale data. Mycologia. 108:1028–1046

St-Arnaud, M., Hamel, C., Vimard, B., Caron, M., and Fortin, J. A. 1996. Enhanced

hyphal growth and spore production of the arbuscular mycorrhizal fungus Glomus

intraradices in an in vitro system in the absence of host roots. Mycol. Res.

100:328–332

Tang, H., Krishnakumar, V., Bidwell, S., Rosen, B., Chan, A., Zhou, S., Gentzbittel, L.,

Childs, K. L., Yandell, M., Gundlach, H., Mayer, K. F. X., Schwartz, D. C., and

Town, C. D. 2014. An improved genome release (version Mt4.0) for the model

legume Medicago truncatula. BMC Genomics. 15:312

Tejada-Jiménez, M., Gil-Díez, P., León-Mediavilla, J., Wen, J., Mysore, K. S., Imperial,

J., and González-Guerrero, M. 2017. *Medicago truncatula* Molybdate Transporter

type 1 (MtMOT1.3) is a plasma membrane molybdenum transporter required for

nitrogenase activity in root nodules under molybdenum deficiency. New Phytol.

216:1223–1235

The R Foundation. 2020. R: The R Project for Statistical Computing. Available at:

https://www.r-project.org/ [Accessed May 11, 2020].

Trapnell, C., Hendrickson, D. G., Sauvageau, M., Goff, L., Rinn, J. L., and Pachter, L.

2013. Differential analysis of gene regulation at transcript resolution with RNA-seq.

Nat. Biotechnol. 31:46–53

Uhe, M., Hogekamp, C., Hartmann, R. M., Hohnjec, N., and Küster, H. 2018. The

mycorrhiza-dependent defensin MtDefMd1 of *Medicago truncatula* acts during the

late restructuring stages of arbusculecontaining cells. PLoS One. 13:e0191841

Vernié, T., Camut, S., Camps, C., Rembliere, C., Carvalho-Niebel, F., Mbengue, M.,

Timmers, T., Gasciolli, V., Thompson, R., Le Signor, C., Lefebvre, B., Cullimore, J.,

and Hervé, C. 2016. PUB1 interacts with the receptor kinase DMI2 and negatively

regulates rhizobial and arbuscular mycorrhizal symbioses through its ubiquitination

activity in *Medicago truncatula*. Plant Physiol. 170:2312–2324

Vierheilig, H., Coughlan, A. P., Wyss, U., and Piché, Y. 1998. Ink and vinegar, a simple

staining technique for arbuscular-mycorrhizal fungi. Appl. Environ. Microbiol.

64:5004–5007

Vitousek, P. M., Porder, S., Houlton, B. Z., and Chadwick, O. A. 2010. Terrestrial

phosphorus limitation: mechanisms, implications, and nitrogen–phosphorus

interactions. Ecol. Appl. 20:5–15

Wang, S., Chen, A., Xie, K., Yang, X., Luo, Z., Chen, J., Zeng, D., Ren, Y., Yang, C.,

Wang, L., Feng, H., López-Arredondo, D. L., Herrera-Estrella, L. R., and Xu, G.

2020. Functional analysis of the OsNPF4.5 nitrate transporter reveals a conserved

mycorrhizal pathway of nitrogen acquisition in plants. Proc. Natl. Acad. Sci. U. S. A.

Wang, T., Zhao, M., Zhang, X., Liu, M., Yang, C., Chen, Y., Chen, R., Wen, J., Mysore,

K. S., and Zhang, W. H. 2017. Novel phosphate deficiency-responsive long non-

coding RNAs in the legume model plant *Medicago truncatula*. J. Exp. Bot. 68:5937–

5948

Wang, X., Zhao, S., and Bücking, H. 2016. Arbuscular mycorrhizal growth responses

are fungal specific but do not differ between soybean genotypes with different

phosphate efficiency. Ann. Bot. 118:11–21

Wipf, D., Mongelard, G., van Tuinen, D., Gutierrez, L., and Casieri, L. 2014.

Transcriptional responses of *Medicago truncatula* upon sulfur deficiency stress and

arbuscular mycorrhizal symbiosis. Front. Plant Sci. 5:680

Wright, D. P., Read, D. J., and Scholes, J. D. 1998. Mycorrhizal sink strength influences

whole plant carbon balance of Trifolium repens L. Plant, Cell Environ. 21:881–891

Yoneyama, K., Yoneyama, K., Takeuchi, Y., and Sekimoto, H. 2007. Phosphorus

deficiency in red clover promotes exudation of orobanchol, the signal for

mycorrhizal symbionts and germination stimulant for root parasites. Planta.

225:1031–1038

Zhang, Q., Blaylock, L. A., and Harrison, M. J. 2010. Two *Medicago truncatula* half-ABC

transporters are essential for arbuscule development in arbuscular mycorrhizal

symbiosis. Plant Cell. 22:1483–1497

Zhang, Y. M., Wang, H. Q., Liu, D. M., and Liu, R. J. 2020. Three tandemly aligned lea

genes from *Medicago truncatula* confer differential protection to *Escherichia coli*

against abiotic stresses. Biol. Plant. 64:95–103

FIGURE LEGENDS

Figure 1. Arbuscular mycorrhizal colonization of *Medicago truncatula* roots and arbuscule

size. **A,** percent root length colonization of roots inoculated with *Glomus aggregatum* 165

(Ga165; grey) or *Rhizophagus irregularis* 09 (Ri09; dark grey), measured for intraradical

hyphae (IRH), arbuscules, and vesicles. Data points represent individual values for

biological replicates (n = 5) and bars represent the mean ± SEM. **B-C**, representative

confocal microscope images of Ga165 (**B**) and Ri09 (**C**) arbuscules stained with wheat

germ agglutinin conjugated to Alexa Fluor 488. Scale bars = 50 μm. **D**, violin plot of

arbuscule widths measured in roots colonized by Ga165 (grey) or Ri09 (dark grey)

overlayed with box and whisker plots. In total, 488 and 508 arbuscules were measured in

Ga165 and Ri09-colonized roots, respectively. All data were analyzed using a Welch's

two-sample t-test and statistically significant differences (p ≤ 0.05) are indicated with an

asterisk.

Figure 2. Biomass and nutrient content of *Medicago truncatula* plants colonized by different arbuscular mycorrhizal fungi. Shown are dry weight (**A**,**B**), phosphorus content (**C**,**D**), and % nitrogen-15 enrichment (**E**,**F**) of roots (left) and shoots (right). The plants were inoculated with either *Glomus aggregatum* 165 (Ga165) or *Rhizophagus irregularis* 09 (Ri09) and compared to non-mycorrhizal (NM) control plants. Data points represent individual values for biological replicates (n = 5) and bars represent the mean of each treatment ± SEM. Different letters on the bars indicate statistically significant (p ≤ 0.05) differences based on one-way ANOVA and LSD test. F statistics and p-values are shown in **Supplemental Table S1**.

Figure 3. Significantly differentially expressed genes in roots (**A–C**) and shoots (**D–F**) of *Medicago truncatula* colonized by different AM fungi. **A** and **D**, Number of significantly differentially expressed genes (DEGs; $\log_2$[fold-change] > 2 and a q-value < 0.05) in plants colonized by *Glomus aggregatum* 165 (Ga165) or *Rhizophagus irregularis* 09 (Ri09) compared either to non-mycorrhizal (NM) control plants or to each other. **B** and **E**, Venn diagrams of significantly upregulated (yellow) and downregulated (blue) DEGs unique to Ga165 and Ri09-colonized plants or shared by both. **C** and **F**, Heat maps of fragments per kilobase of transcript per million mapped reads (FPKM) values + 1 for all significant DEGs from each treatment. Gene clusters are assigned based on second-order and sub-clusters based on third- or fourth-order branching. Upregulated genes are shown in yellow and downregulated genes in blue with intensity based on row z-score. All DEGs in each plot for roots and shoots, respectively, are listed in **Supplemental Files 2 and 3**.

Figure 4. Expression patterns of genes encoding enzymes involved in the strigolactone

(**A**) and gibberellic acid (**B**) biosynthesis pathways in the roots of *Medicago truncatula*

colonized by arbuscular mycorrhizal fungi. Each diagram portrays the relative expression

levels of genes that encode components of the pathways shown as fragments per

kilobase of transcript per million mapped reads (FPKM; blue=low, yellow=high) in roots

from non-mycorrhizal (NM) plants or plants colonized by either *Glomus aggregatum* 165

(Ga165) or *Rhizophagus irregularis* 09 (Ri09). Letters within boxes indicate significant

differences (q-value < 0.05) in FPKM values between treatments and were determined

as part of the CuffDiff2 analysis (see **Supplemental Table S2**). Abbreviations: **A**, GAP,

glyceraldehyde 3-phosphate; DXS2, 1-deoxyxylulose 5-phosphate synthase2; MEP, 2-

methyl-erythritol-4-phosphate; GGDP geranylgeranyl pyrophosphate; D27, Dwarf27;

CCD, carotenoid cleavage dioxygenase; and PDR1a, pleiotropic drug resistance 1a. **B**,

CPS, *ent*-copalyl diphosphate synthase; KS, *ent*-kaurene synthase; KO, *ent*-kaurene

oxidase; KAO, *ent*-kaurenoic acid oxidase; GA13ox, gibberellin 13-oxidase; GA20ox,

gibberellin 20-oxidase; GA3ox, gibberellin 3-oxidase; and GA2ox, gibberellin 2-oxidase.

Figure 5. Relative expression of six genes in the roots of *Medicago truncatula* after

colonization with different arbuscular mycorrhizal fungi. Expression of **A**, *Required for*

Arbuscule Development 1 (*RAD1*); **B**, *MYB-like Transcription Factor 1* (*MYB1*); **C**,

Phosphate Transporter 8 (*PT8*); **D**, *Ammonium Transporter 2;3* (*AMT2;3*); **E**, *Reduced*

Arbuscular Mycorrhization 2 (*RAM2*); and **F**, *Sugars Will Eventually Be Exported*

Transporter 1.2 (*SWEET1.2*) in the roots of non-mycorrhizal plants (NM, n=3) and plants

inoculated with *Glomus aggregatum* 165 (Ga165, n=3) or *Rhizophagus irregularis* 09

(Ri09, n=4). Expression levels are shown as fragments per kilobase of transcript per

million mapped reads (FPKM). Data points represent individual values for biological

replicates (n = 5) and bars represent the mean of each treatment ± SEM. Significant

differences (q-value < 0.05) in FPKM values between treatments were determined as part

of the CuffDiff2 analysis and are indicated using different letters (see **Supplemental**

Table S3).

Figure 6. Expression patterns of significantly differentially expressed mineral transporter

genes in the roots of *Medicago truncatula* after colonization with different arbuscular

mycorrhizal fungi. **A**, Venn diagrams showing the number of mineral transporters that are

significantly up- (yellow) or down-regulated (blue) in roots colonized by *Glomus*

aggregatum 165 (Ga165) or *Rhizophagus irregularis* 09 (Ri09), or both. **B**, summary of

the number of genes from specific gene annotation groups of mineral transporters that

are up- or down-regulated in roots colonized by Ga165 or Ri09, or both (see also

Supplemental File 6). Expression patterns of genes annotated as **C**, ammonium; **D**,

nitrate; **E**, phosphate; and **F**, sulfate transporters in the roots of non-mycorrhizal plants

(NM; n=3) and plants inoculated with Ga165 (n=3) or Ri09 (n=4). The bars in C–F

represent the mean of each treatment ± SEM (data points for biological replicates are

displayed in **Supplemental Figure S8**). Expression levels are shown as fragments per

kilobase of transcript per million mapped reads (FPKM), only genes with FPKM ≥ 10 are

shown, significant differences (q-value < 0.05) in FPKM values between treatments were

determined as part of the CuffDiff2 analysis and are indicated using different letters on

the bars.

1009

Figure 7. Expression patterns of significantly differentially expressed secondary

metabolite transporters in the roots of *Medicago truncatula* after colonization with different

AM fungi. **A**, Venn diagrams showing the number of secondary metabolite transporters

that are significantly up- (yellow) or down-regulated (blue) in roots colonized by *Glomus*

aggregatum 165 (Ga165) or *Rhizophagus irregularis* 09 (Ri09), or both. **B**, summary of

the number of genes from specific gene annotation groups of secondary metabolite

transports that are up- or down-regulated in roots colonized by Ga165 or Ri09, or both

(see **Supplemental File 7**). Expression patterns of genes annotated as **C**, ATP-binding

cassette (ABC); **D**, amino acid or peptide; **E**, major facilitator superfamily (MFS) or major

intrinsic protein (MIP); and **F**, SWEET transporters in the roots of non-mycorrhizal plants

(NM; n = 3) and plants inoculated with Ga165 (n =3) or Ri09 (n = 4). The bars in C-F

represent the mean of each treatment ± SEM (data points for biological replicates are

displayed in **Supplemental Figure S9**). Expression levels are shown as fragments per

kilobase of transcript per million mapped reads (FPKM); only genes with FPKM ≥ 10 are

shown; significant differences (q-value < 0.05) in FPKM values between treatments were

determined as part of the CuffDiff2 analysis and are indicated using different letters on

the bars. Abbreviations in B: G3P, glycerol-3-phosphate; NAT, nucleobase-ascorbate

transporter; OST, organic solute transporter; TPT, triose-phosphate transporter; TRAPP,

transport protein particle.

1029

Figure 8. Expression patterns of significantly differentially expressed genes in the shoots

of *Medicago truncatula* colonized by different AM fungi. Expression levels of genes

involved in **A**, photosynthesis; **B**, response to biotic stress; and **C**, response to abiotic

stress in non-mycorrhizal plants (NM; n=4) or plants colonized by *Glomus aggregatum*

165 (Ga165, n=4) or *Rhizophagus irregularis* 09 (Ri09, n=3). Bars represent the mean of

each treatment ± SEM (data points for biological replicates are displayed in

Supplemental Figure S12). Expression levels are shown as fragments per kilobase of

transcript per million mapped reads (FPKM), and significant differences (q-value < 0.05)

in FPKM values between treatments were determined as part of the CuffDiff2 analysis

and are indicated using different letters above the bars (see also **Supplemental Table**

S4).

1041

Figure 9. Summary of the physiological and transcriptomic responses in roots and shoots

of *Medicago truncatula* colonized by either *Glomus aggregatum* 165 or *Rhizophagus*

irregularis 09. Physiological data include the analysis of AM colonization, plant biomass,

and both phosphorus and nitrogen-15 contents in plant tissues. Transcriptomic data

include the expression patterns of nutrient transporters and enrichment of gene ontology

terms. Yellow up-arrows indicate an increase, equal signs (=) specify no change, and blue

down-arrows designate a decrease in each response compared to non-mycorrhizal

plants. Arrow size indicates the degree of the response.

1050

1051

Supplemental Table S1. F statistics and p-values from one-way ANOVA for each

experimental parameter.

Supplemental Table S2. Expression levels of genes associated with the strigolactone

biosynthesis pathway and the common symbiosis signaling pathway in roots of *Medicago*

truncatula.

Supplemental Table S3. Expression levels of genes downstream of the common

symbiosis signaling pathway in roots of *Medicago truncatula*.

Supplemental Table S4. List of differentially regulated genes from shoot gene ontology

enrichment analysis.

Supplemental Table S5. List of differentially regulated defense response genes from

root gene ontology enrichment analysis.

Supplemental File 1. Significantly differentially expressed genes in roots

Supplemental File 2. Significantly differentially expressed genes in shoots

Supplemental File 3. Significant gene ontology terms in roots

Supplemental File 4. Significantly differentially expressed transporters in roots

Supplemental File 5. List of mineral transporters by type

Supplemental File 6. List of secondary metabolite transporters by type

Supplemental File 7. Significant gene ontology terms in shoots

1093

Supplemental Figure S1. Schematic model of the dual-compartment pot system used for this study. Roots in the root compartment (RC) were mock-inoculated for the non-mycorrhizal control treatment and for the mycorrhizal treatments, roots were inoculated with either 500 spores of either *Glomus aggregatum* 165 or *Rhizophagus irregularis* 09. Five weeks post-inoculation, 0.5 mM KH_2PO_4 and 4 mM $^{15}NH_4Cl$ were added to the hyphal compartment (HC).

Supplemental Figure S2. Shoot and root phosphorous concentration of *Medicago truncatula* colonized by arbuscular mycorrhizal fungi. The plants were inoculated with either *Glomus aggregatum* 165 (Ga165) or *Rhizophagus irregularis* 09 (Ri09). Control plants were mock inoculated and served as non-mycorrhizal (NM) controls. Data points represent individual values for biological replicates (n = 5) and bars represent the mean of each treatment ± SEM. Different letters above the bars indicate statistically significant differences (p ≤ 0.05) based on one-way ANOVA and LSD test. F statistics and p-values are shown in **Supplemental Table S1**.

Supplemental Figure S3. Hierarchical clustering of biological replicates from root RNA sequencing data. Treatments include non-mycorrhizal (NM, n = 3), *Glomus aggregatum* 165 (Ga165, n = 3) and *Rhizophagus irregularis* 09 (Ri09, n = 4). The data were clustered using **A**, dendrogram analysis; **B**, multidimensional scaling; and principal component analyses (PCA), including PC1 vs. PC2 (**C**) and PC2 vs. PC3 (**D**). Note the distinct clustering of all biological replicates from each treatment.

Supplemental Figure S4. Hierarchical clustering of biological replicates from shoot RNA sequencing data. Treatments include non-mycorrhizal (NM, n = 4), *Glomus aggregatum* 165 (Ga165, n = 4) and *Rhizophagus irregularis* 09 (Ri09, n = 3). The data were clustered using **A**, dendrogram analysis; **B**, multidimensional scaling; and **C-D**, principal component analyses (PCA), including PC1 vs. PC2 (**C**) and PC2 vs. PC3 (**D**). Note that no distinct clustering of biological replicates was observed for any of the treatments.

Supplemental Figure S5. Volcano plots of differentially expressed genes (DEGs) between treatments. **A-C**, shoot data are shown on top and **D-F**, root data on bottom for the following comparisons: left, *Glomus aggregatum* 165 (Ga165) vs. non-mycorrhizal (NM); middle, *Rhizophagus irregularis* 09 (Ri09) vs. NM; and right, Ri09 vs. Ga165. Data points representing significant DEGs ($-\log_{10}$[p-value] > 2.3) are shown in red while non-significant DEGs ($-\log_{10}$[p-value] < 2.3) are shown in black. Note the greater abundance of significant DEGs among the root data compared to the shoot data.

Supplemental Figure S6. Summary of gene ontology enrichment analysis of genes upregulated in the roots of *Medicago truncatula*. All enriched gene ontology (GO) terms for **A**, *Glomus aggregatum* 165; **B**, *Rhizophagus irregularis* 09; or **C**, both species are shown for molecular function, cellular component, and biological process. The enrichment score (-log$_{10}$[p-value]) for each GO term is also shown ranging from low (light yellow) to high (red). A list of the genes from each GO term can be found in **Supplemental File 4**.

Supplemental Figure S7. Summary of gene ontology enrichment analysis of genes downregulated in the roots of *Medicago trunctaula*. All enriched gene ontology (GO) terms for **A**, *Glomus aggregatum* 165; **B**, *Rhizophagus irregularis* 09; or **C**, both species are shown for molecular function, cellular component, and biological process. The enrichment score ($-\log_{10}$[p-value]) for each GO term is also shown ranging from low (light yellow) to high (red). A list of the genes from each GO term can be found in **Supplemental File 4**.

Supplemental Figure S8. Alternative portrayal of expression patterns of genes annotated as **A**, ammonium; **B**, nitrate; **C**, phosphate; and **D**, sulfate transporters. Shown are individual data points for gene expression values from each biological replicate overlayed on the graphs from **Figure 6**, panels C through F, respectively.

Supplemental Figure S9. Alternative portrayal of expression patterns of genes annotated as **A**, ABC; **B**, amino acid or peptide; **C**, MIP or MFS; and **D**, SWEET transporters. Shown are individual data points for gene expression values from each biological replicate overlayed on the graphs from **Figure 7**, panels C through F, respectively.

Supplemental Figure S10. Summary of gene ontology enrichment analysis of differentially expressed genes in the shoots of *Medicago trunctaula*. Enriched gene ontology (GO) terms for molecular function, cellular component, and biological process among the following groups of genes based on the heatmap in **Figure 2F**: **A**, genes upregulated in the shoots of plants colonized by *Rhizophagus irregularis* 09 (Ri09); **B**, genes upregulated in the shoots of plants colonized by *Glomus aggregatum* 165 (Ga165); and **C**, genes downregulated in the shoots of plants colonized by Ri09. The enrichment score ($-\log_{10}$[p-value]) for each GO term is also shown ranging from low (light yellow) to high (red). A list of the genes from each GO term can be found in **Supplemental File 8**.

Supplemental Figure S11. Alternative portrayal of expression patterns of genes involved in **A**, photosynthesis; **B**, response to biotic stress; and **C**, response to abiotic stress. Shown are individual data points for gene expression values from each biological replicate overlayed on the graphs from **Figure 8**, panels A through B, respectively.

Supplemental Table S1. F statistics and p-values from one-way ANOVA for each experimental parameter.

Parameter	F statistics	p-value
Shoot dry wt. (g)	$F_{2,14} = 16.61$	0.0003
Root dry wt. (g)	$F_{2,14} = 13.14$	0.0009
Shoot P concentration (mg/g)	$F_{2,14} = 21.17$	<0.0001
Total shoot P content (mg)	$F_{2,14} = 26.79$	<0.0001
Root P concentration (mg/g)	$F_{2,14} = 127.01$	<0.0001
Total root P content (mg)	$F_{2,14} = 136.80$	<0.0001
Shoot ^{15}N enrichment (%)	$F_{2,14} = 12.01$	0.0014
Root ^{15}N enrichment (%)	$F_{2,14} = 49.75$	<0.0001

Supplemental Table S2. Expression levels of genes associated with the strigolactone biosynthesis pathway, the common symbiosis signaling pathway, and the gibberellic biosynthesis and degradation pathway in roots of *Medicago truncatula*. Mean FPKM (Fragments Per Kilobase of transcript per Million mapped reads) values for all listed genes are shown for all three treatments from this study, including nonmycorrhizal (NM, n=3) control plants and both *Glomus aggregatum* 165 (Ga165, n=3) and *Rhizophagus irregularis* 09 (Ri09, n=4) mycorrhizal plants. A green (low) to red (high) color scale is used to highlight differences in FPKM values. Significant differences (q-value < 0.05) in FPKM values between treatments were determined as part of the CuffDiff2 analysis and are indicated using different letters. References for the studies in which each gene was characterized are also included. Full citations can be found in the **Supplemental References**.

	Gene ID	Gene Name	NM	Ga165	Ri09	Reference
Strigolactone Biosynthesis & Transport	*Medtr8g068265*	*DXS2*	9.37 b	37.77 a	36.07 a	(Floß et al., 2008)
	Medtr1g471050	*D27*	37.83 b	64.74 a	80.61 a	(Müller et al., 2019)
	Medtr7g045370	*CCD7*	21.91 ab	15.87 b	28.47 a	(Gomez-Roldan et al., 2008)
	Medtr3g109610	*CCD8-1*	49.43 a	47.47 a	65.00 a	(Gomez-Roldan et al., 2008)
	Medtr3g104560	*MAX1a*	82.93 b	70.28 b	140.84 a	(Müller et al., 2019)
	Medtr8g020840	*NSP1*	6.02 c	13.98 b	22.41 a	(Liu et al., 2011)
	Medtr3g072710	*NSP2*	13.47 b	12.39 b	23.85 a	(Liu et al., 2011)
	Medtr3g107870	*PDR1a*	0.34 a	3.19 a	1.26 a	(Kretzschmar et al., 2012)
Common Symbiosis Signaling Pathway	*Medtr5g019040*	*NFP*	2.63 a	0.79 b	0.99 b	(Feng et al., 2019)
	Medtr3g080050	*LYK9*	0.88 a	2.37 a	0.27 a	(Gibelin-Viala et al., 2019)
	Medtr5g030920	*DMI2/NORK*	61.59 a	59.99 a	68.39 a	(Stracke et al., 2002)
	Medtr5g083030	*PUB1*	8.48 b	14.21 a	17.74 a	(Vernié et al., 2016)
	Medtr5g026500	*HMGR1*	29.53 a	35.08 a	22.69 a	(Kevei et al., 2007)
	Medtr2g005870	*DMI1*	3.42 a	2.91 a	3.95 a	(Ané et al., 2004)
	Medtr7g117580	*CASTOR*	20.94 a	25.02 a	26.92 a	(Charpentier et al., 2008)
	Medtr7g100110	*MCA8*	53.22 a	66.33 a	71.46 a	(Capoen et al., 2011)
	Medtr1g064240	*CNGC15a*	0.00 a	0.00 a	0.01 a	(Charpentier et al., 2016)
	Medtr4g058730	*CNGC15b*	0.29 a	3.98 a	6.61 a	(Charpentier et al., 2016)
	Medtr2g094860	*CNGC15c*	0.67 a	0.05 a	0.29 a	(Charpentier et al., 2016)
	Medtr8g043970	*DMI3/CCaMK*	19.29 a	19.19 a	16.05 a	(Lévy et al., 2004)
	Medtr5g026850	*IPD3/CYCLOPS*	25.28 a	33.95 a	37.35 a	(Horváth et al., 2011)
	Medtr3g065980	*DELLA1*	65.07 a	38.27 a	56.47 a	(Floss et al., 2013)
	Medtr4g122240	*DELLA2*	2.91 a	3.56 a	3.04 a	(Floss et al., 2013)
Gibberellic Acid	Medtr7g011663	CPS	17.09 a	11.06 ab	9.34 b	(Igielski et al., 2017)
	Medtr2g064295	KS	6.62 a	5.17 a	4.07 a	(Igielski et al., 2017)
	Medtr2g105360	KO	43.47 b	50.49 b	76.47 a	(Igielski et al., 2017)
	Medtr2g031930	KAO1	30.06 a	29.73 a	26.21 a	(Igielski et al., 2017)
	Medtr3g093530	CYP714_A1/GA13ox	6.71 b	13.26 a	10.26 ab	(Igielski et al., 2017)
	Medtr1g102070	GA20ox1	5.36 b	12.48 a	8.23 ab	(Igielski et al., 2017)
	Medtr1g086550	GA20ox6	7.66 c	41.71 a	16.02 b	(Nouri et al., 2021)
	Medtr8g033380	GA20ox8	0.16 c	8.21 a	2.42 b	(Igielski et al., 2017)
	Medtr2g102570	GA3ox1	10.68 a	7.96 ab	5.24 b	(Igielski et al., 2017)
	Medtr2g019370	GA2ox1	0.96 c	16.73 a	9.99 b	(Nouri et al., 2021)

Supplemental Table S3. Expression levels of genes downstream of the common symbiosis signaling pathway in roots of *Medicago truncatula*. Mean FPKM (Fragments Per Kilobase of transcript per Million mapped reads) values for all listed genes are shown for all three treatments from this study, including nonmycorrhizal (NM, n=3) control plants and both *Glomus aggregatum* 165 (Ga165, n=3) and *Rhizophagus irregularis* 09 (Ri09, n=4) mycorrhizal plants. A green (low) to red (high) color scale is used to highlight differences in FPKM values. Significant differences (q-value < 0.05) in FPKM values between treatments were determined as part of the CuffDiff2 analysis and are indicated using different letters. References for the studies in which each gene was identified as playing a role in the arbuscular mycorrhizal symbiosis are also included. Full citations can be found in the **Supplemental References**.

	Gene ID	Gene Name	NM	Ga165	Ri09	Reference
Transcription Factors	Medtr7g027190	RAM1	0.01 b	45.05 a	52.08 a	(Park et al., 2015)
	Medtr4g104020	RAD1	0.09 c	84.69 b	139.93 a	(Park et al., 2015)
	Medtr7g088990	DIP1	111.61 a	145.81 a	137.94 a	(Yu et al., 2014)
	Medtr2g034280	MIG1	0.56 a	0.97 a	1.15 a	(Heck et al., 2016)
	Medtr2g034260	MIG2	9.07 b	20.31 a	18.69 a	(Heck et al., 2016)
	Medtr2g034250	MIG3	7.73 b	18.70 a	25.75 a	(Heck et al., 2016)
	Medtr7g068600	MYB1	0.06 c	594.35 b	1262.52 a	(Floss et al., 2017)
Cellular Remodeling	Medtr6g027840	VAPYRIN/VPY	3.20 b	34.54 a	33.33 a	(Pumplin et al., 2010)
	Medtr1g017910	EXO70I	2.53 b	24.18 a	18.44 a	(Zhang et al., 2015)
	Medtr2g088700	SYP132A	56.60 b	92.13 a	79.90 ab	(Pan et al., 2016)
Lipid Biosynthesis & Transport	Medtr8g468920	WRI5a	0.00 b	2.26 a	3.15 a	(Jiang et al., 2018)
	Medtr8g107450	STR	0.68 b	51.53 b	71.44 a	(Zhang et al., 2010)
	Medtr5g030910	STR2	0.22 b	55.77 a	69.28 a	(Zhang et al., 2010)
	Medtr1g040500	RAM2	0.23 c	192.67 b	353.74 a	(Luginbuehl et al., 2017)
	Medtr1g109110	FatM	0.07 a	12.57 a	17.47 a	(Bravo et al., 2017)
Mineral Transport	Medtr1g028600	PT4	0.09 b	277.24 a	381.65 a	(Javot et al., 2007)
	Medtr5g068140	PT8	0.05 c	81.31 b	133.25 a	(Breuillin-Sessoms et al., 2015)
	Medtr8g074750	AMT2;3	0.00 c	7.08 b	13.07 a	(Breuillin-Sessoms et al., 2015)
	Medtr7g115050	AMT2;4	0.15 b	6.53 a	3.59 a	(Breuillin-Sessoms et al., 2015)
	Medtr1g036410	AMT2;5	0.11 b	192.75 a	213.55 a	(Breuillin-Sessoms et al., 2015)
Sugar Transport	Medtr3g089125	SWEET1.2	0.04 c	6.81 b	32.93 a	(An et al., 2019)

Supplemental Table S4. List of differentially regulated genes from shoot gene ontology enrichment analysis. Mean FPKM (Fragments Per Kilobase of transcript per Million mapped reads) values for all listed genes are shown for all three treatments from this study, including nonmycorrhizal (NM, n=3) control plants and both *Glomus aggregatum* 165 (Ga165, n=3) and *Rhizophagus irregularis* 09 (Ri09, n=4) mycorrhizal plants. A green (low) to red (high) color scale is used to highlight differences in FPKM values. Significant differences (q-value < 0.05) in FPKM values between treatments were determined as part of the CuffDiff2 analysis and are indicated using different letters.

	Gene ID	Gene Annotation	NM	Ga165	Ri09
Photosynthesis	*Medtr1g015290*	ultraviolet-B-repressible protein	268.2 b	662.9 ab	992.1 a
	Medtr2g082580	oxygen-evolving enhancer protein	15.9 b	28.6 ab	54.4 a
	Medtr1g115410	photosystem II reaction center PsbP family protein	17.3 b	44.6 a	52.8 a
	Medtr1588s0010	ATP synthase F1, gamma subunit	605.3 b	1556.0 ab	1993.1 a
	Medtr6g084320	ABC transporter-like family-protein	12.8 b	21.8 ab	47.0 a
Biotic Stress	*Medtr4g054920*	cytochrome P450 family 94 protein	8.26 b	62.56 a	15.45 b
	Medtr4g120760	pathogenesis-related protein bet V I family protein	16.02 ab	51.03 a	13.83 b
	Medtr5g053950	allene oxide cyclase	77.55 b	391.10 a	162.61 ab
	Medtr8g096900	pathogenesis-related thaumatin family protein	17.47 b	45.09 a	36.91 ab
	Medtr4g092010	(3S)-linalool/(E)-nerolidol/(E,E)-geranyl linalool synthase	11.90 b	50.43 a	33.24 ab
Abiotic Stress	*Medtr1g048990*	superoxide dismutase	105.08 a	63.93 ab	30.32 b
	Medtr3g087590	myo-inositol 1-phosphate synthase	322.70 ab	388.70 a	84.82 b
	Medtr7g069980	ferritin	60.16 a	33.23 a	6.36 b
	Medtr8g059170	NAC transcription factor-like protein	49.87 a	41.35 a	6.64 b
	Medtr5g018670	photosystem II oxygen-evolving enhancer protein	35.97 b	103.56 a	87.44 ab
	Medtr7g079180	late embryogenesis abundant protein	29.47 b	103.91 a	78.75 ab

Supplemental Table S5. List of differentially regulated defense response genes from root gene ontology enrichment analysis. Mean FPKM (Fragments Per Kilobase of transcript per Million mapped reads) values for all listed genes are shown for all three treatments from this study, including nonmycorrhizal (NM, n=3) control plants and both *Glomus aggregatum* 165 (Ga165, n=3) and *Rhizophagus irregularis* 09 (Ri09, n=4) mycorrhizal plants. A green (low) to red (high) color scale is used to highlight differences in FPKM values. Significant differences (q-value < 0.05) in FPKM values between treatments were determined as part of the CuffDiff2 analysis and are indicated using different letters.

	Gene ID	Gene Annotation	NM	Ga165	Ri09
	Medtr0012s0290	disease resistance protein (TIR-NBS-LRR class), putative	0.91 a	3.53 b	5.78 c
	Medtr1g015140	WRKY family transcription factor	0.67 a	1.06 ab	2.68 b
	Medtr2g008430	LL-diaminopimelate aminotransferase	2.08 a	7.54 b	14.17 c
	Medtr3g028040	LRR and NB-ARC domain disease resistance protein	0.08 a	0.53 ab	1.10 b
Defense Response	*Medtr3g072140*	disease resistance protein (TIR-NBS-LRR class), putative	0.15 a	0.56 ab	0.82 b
	Medtr6g006990	carbonic anhydrase family protein	0.56 a	3.53 ab	7.55 b
	Medtr8g012810	Defensin related	0.00 a	0.67 ab	45.59 b
	Medtr8g012815	Defensin related	0.00 a	1.00 ab	34.68 b
	Medtr8g012845	Defensin related	0.00 a	0.36 ab	51.54 b
	Medtr8g012870	Defensin related	0.00 a	0.49 ab	5.59 b
	Medtr8g059405	Defensin	0.97 a	0.73 a	8.89 b