Conformational Ensembles of Non-Coding Elements in the SARS-CoV-2 Genome from Molecular Dynamics Simulations

Sandro Bottaro[1,*]**, Giovanni Bussi**[2]**, Kresten Lindorff-Larsen**[1,*]

*For correspondence:
sandro.bottaro@bio.ku.dk (SB);
lindorff@bio.ku.dk (KLL)

[1]Structural Biology and NMR Laboratory & Linderstrøm-Lang Centre for Protein Science
Department of Biology, University of Copenhagen. Ole Maaløes Vej 5, DK-2200
Copenhagen N, Denmark; [2]Scuola Internazionale Superiore di Studi Avanzati (SISSA),
Via Bonomea 265, 34136, Trieste, Italy

Abstract The 5' untranslated region (UTR) of the SARS-CoV-2 genome is a conserved, functional and structured genomic region consisting of several RNA stem-loop elements. While the secondary structure of such elements has been determined experimentally, their three-dimensional structure is not known yet. Here, we predict structure and dynamics of five RNA stem loops in the 5'-UTR of SARS-CoV-2 by extensive atomistic molecular dynamics simulations, more than 0.5ms of aggregate simulation time, in combination with enhanced sampling techniques. We compare simulations with available experimental data, describe the resulting conformational ensembles, and identify the presence of specific structural rearrangements in apical and internal loops that may be functionally relevant. Our atomic-detailed structural predictions reveal a rich dynamics in these RNA molecules, could help the experimental characterisation of these systems, and provide putative three-dimensional models for structure-based drug design studies.

Introduction

The genome of the SARS-CoV-2 virus is a single, positive stranded RNA consisting of approximately 30 thousand nucleotides. The 5' and 3' untranslated regions (UTR) contain cis-acting sequences required for viral transcription and replication (*Lal, 2010*; *Yang and Leibowitz, 2015*). Recent studies suggest the 5'-UTR to enhance viral transcription (*Schubert et al., 2020*), and undergoes liquid-liquid phase separation with the SARS-CoV-2 nucleocapsid protein (*Iserman et al., 2020*; *Carlson et al., 2020*).

Chemical probing experiments, NMR chemical shift measurements, and computational predictions have shown that the 5'-UTR is highly structured (*Wacker et al., 2020*; *Andrews et al., 2020*; *Manfredonia et al., 2020*; *Rangan et al., 2020b*; *Miao et al., 2020*), and consists of several stem-loop elements (Fig 1). The secondary structures of the isolated elements are in close agreement with the full-length construct (*Wacker et al., 2020*), suggesting that they can fold independently. The three-dimensional structure of stem loop 2 (SL2) from SARS-CoV-1, which is completely conserved in SARS-CoV-2, has been determined by NMR spectroscopy (*Lee et al., 2011*), while no experimental structures exist for all the other elements in the 5'-UTR. In a recent computational study, *Rangan et al.* (*2020a*) predicted the three-dimensional fold of all stem-loop elements in both 5' and

3'-UTR using Rosetta's FARFAR2 algorithm (*Watkins et al., 2020*). The predicted SL2 model in the

5'-UTR was later refined using MD simulations, (*Bergonzo and Szakal, 2020*) leading to an improved

agreement with available NMR data (*Lee et al., 2011*).

In this work we have used atomistic molecular dynamics simulations to predict the structure

and dynamics of five 5'-UTR stem-loops. We have generated three-dimensional configurations

compatible with the secondary structure recently determined by chemical probing and NMR ex-

periments (*Wacker et al., 2020*). While several fragment-based tools exist to perform such task

(*Parisien and Major, 2008*; *Cheng et al., 2015*), we here adapt an MD-based procedure (*Poblete*

et al., 2018). These initial structures are subject to extensive MD sampling, amounting to more

than 0.5ms of aggregate simulation time. More precisely, we enhance the sampling of apical and

internal loop regions (*Wang et al., 2011*; *Bussi, 2014*), where we currently lack detailed structural

information and where significant conformational changes are more likely to occur. Our simula-

tions are thus a Boltzmann-weighted ensemble of three-dimensional conformations constituting

our prediction of structure and conformational heterogeneity of these elements.

First, we apply the computational protocol to stem loop 2 (SL2), and validate the accuracy of

our prediction against available experimental NMR measurements on SL2 from SARS-CoV-1 (*Lee*

et al., 2011). We then proceed by describing our predictions on systems of increasing complexity,

stem loop 5a (SL5a), SL1, SL4, and SL2+SL3 (Fig.1), for which no detailed structural information is

available.

Figure 1. Reference sequence and secondary structure of the 5'-UTR in SARS-CoV-2 genome. The numbering follows the convention in *Wacker et al.* (*2020*). The five structured elements considered in this work are labeled: SL1, SL2, SL2+SL3, SL4, and SL5a. The conserved sequence of the transcription regulatory sequence and the AUG start codon are highlighted in orange and purple, respectively. The centroids of the most populated clusters from MD simulations corresponding to each element are shown.

Our predictions could help the atomic-detailed experimental characterisation of such systems,
as simulations can be easily integrated a posteriori with NMR or SAXS data (*Cesari et al., 2018*;
Bottaro and Lindorff-Larsen, 2018; *Bottaro et al., 2020*). The predicted models provide a starting
point for structure-based drug design studies (*Stelzer et al., 2011*; *Costales et al., 2020*), and could
serve as additional benchmark systems to evaluate the accuracy of atomistic RNA force-field (*Dans*
et al., 2019).

Results

66 In this section we describe the conformational ensembles obtained from MD simulations. We first
67 show the results obtained on SL2, the simplest system considered in this study. Crucially, in this
68 case it is possible to validate the procedure by comparing with available structural data and NMR
69 measurements.

Stem Loop 2

We run a plain MD simulation of a single-stranded, extended RNA with sequence GGAUCUCUUGUAGAUCU.
After 10ns, we apply a biasing force to promote the formation of an ideal A-form RNA structure in
the region where the secondary structure is known (blue nucleotides in Fig. 2a). By construction,
the eRMSD distance (*Bottaro et al., 2014*) from the template decreases during pulling (Fig. 2b, top
panel). After 40 ns the external bias is removed, and the system fluctuates around a stable struc-
ture in which all five Watson-Crick (WC) base-pairs are formed. (Fig. 2b, bottom panel).

While pulling simulations make it possible quickly to generate conformations with a desired
stem structure, these runs are not sufficiently long to allow rearrangements in the loop region.
Previous simulation studies have shown these motions to occur on long timescales (μs to ms) that
require extensive simulations (*Šponer et al., 2018*). We thus enhance the sampling in the loop
region (red in Fig. 2a) by running extensive partial tempering simulations (20 replicas, each 2.6
μs-long) initialised from a successful pulling run (see Methods).

Analyses of the enhanced simulations show that all five base-pairs in the stem are stable, and
a comparison of two independent simulations shows that our simulations are well converged
(Fig. 2c). In addition, we observe the formation of a stable terminal GU wobble base-pair, as well
as of a C50-G53 Watson-Crick interaction in the loop, resulting in a structure resembling a CUUG
tetraloop (*Jucker and Pardi, 1995*) with a bulged U at position 54. A three-dimensional cluster analy-
sis (*Bottaro et al., 2019*) reveals that the largest cluster ($\approx 90\%$) is remarkably similar to the available
NMR loop structure from SARS-CoV-1 (*Lee et al., 2011*), whose sequence differs from our construct
in the two terminal base-pairs. For example, the centroid of the most populated cluster from MD
simulations superimposes extremely well to the first NMR model in PDB 2L6I (Fig. 2d). In the loop,
C50 forms a canonical base-pair with G53, U52 stacks on top of C50, and both U51/U54 are solvent
exposed. The median distance between the experimental structure and the MD simulation is 1.6
Å heavy-atom root mean square deviation and 0.5 eRMSD (SI4).

The sugar puckers of U51, U52, and U54 are preferably in C2'-endo conformation, in full agree-
ment with NMR measurements (*Lee et al., 2011*) (Fig. 2e). The great majority (169 out of 179) of
upper-bound NOE-derived distances are satisfied in MD simulations (Fig. 2f). The most significant
discrepancies consist of short proton-proton distances between G53-U54 that are not sampled in
our MD runs (SI5).

Stem loop 5a

101 Stem loop 5a (SL5a) consists of two helices (blue nucleotides in Fig. 3a) connected by an internal
102 loop and capped by an hexaloop with sequence UUUCGU. We use the same procedure as described
103 above for SL2 to generate initial configurations and to sample the ensemble of SL5a by enhancing
104 regions where secondary structure is not known (red regions in Fig. 3a). Helix 1 is highly stable
105 throughout the 20 replicas $\times$ 3.6 μs partial tempering simulations (Fig. 3b).

Figure 2. Description of the computational procedure and validation on SL2. **a** Sequence and secondary structure of SL2. The genomic numbering follows the convention in *Wacker et al.* (*2020*). The region used for constructing the template A-form three-dimensional structure is shown in blue. The sampling in the region shown in red is enhanced by partial tempering. **b** Behaviour of a successful pulling simulation. eRMSD (top panel) and distance between ring centres of the five WC base-pairs (bottom panel) during the three stages of the procedure: initial plain MD, pulling stage, and final plain MD stage, as labelled. The eRMSD threshold of 0.75 is shown as a dashed horizontal line. **c** Population of base-pairs from 20 replicas × 2.6 μs partial tempering simulations. Following the Leontis-Westhof classification (*Leontis and Westhof, 2001*), base-pairs are annotated as cis Watson-Crick/Watson-Crick (WWc), GU wobble cis (GUc) or non-Watson-Crick (nWC). For each interaction, the two bars show the statistics from independent simulations. **d** Centroid of the most-populated cluster from MD simulations (blue/red) superposed onto the SL2 loop from SARS-CoV-1 structure (white) (*Lee et al., 2011*). **e** Sugar pucker population in the loop region. **f** Calculated and experimental NOE-derived upper bound distances. The 10 of the 179 calculated NOEs that are larger than the experimental values are located in the white upper-triangular region.

Figure 3. SL5a results. **a** Sequence and secondary structure of SL5a. The genomic numbering follows the convention in *Wacker et al.* (*2020*). The region used for constructing the A-form templates is shown in blue. The sampling in the region shown in red is enhanced by partial tempering. **b** Population of base-pairs from 20 replicas × 3.6 μs partial tempering simulations. Centroids of the most populated internal and apical loop structures discussed in the text are shown. **c** Representative straight conformation. The axes defined by helix 1 and 2, together with the angle θ between them are shown in green. The helical axes are calculated as the vector normal to the base-pair plane, averaged over all base-pairs within the helix. **d** Representative kinked conformation. **e** Time series of the angle between helices, θ, (top panel) and of the distance between U212 and G213 (bottom panel) in the neutral replica. We note that due to the replica-exchange setup used in our simulations, these simulations do not reflect the timescales needed to switch between the two conformations.

In the internal loop we observe the formation of a U194-U212 base-pair with a population of
$\approx 60\%$ (Fig. 3b). The wobble GU base-pair between U195 and G210 in helix 2 is in equilibrium with
an alternative WC/WC base-pair between U195 and U211. The most stable conformation of the
apical loop is characterised by a wobble interaction between U200 and G204, with U205 completely
solvent exposed. In one of the two simulations we observe the transient formation of a U200-U205
base-pair capped by a canonical UUCG tetraloop fold (*Cheong et al., 1990*), as also suggested by
NMR measurements (*Wacker et al., 2020*). The low population of this state is not surprising, as this
fold has been shown not to be stabilised to a sufficient degree within this force-field (*Mráziková*
et al., 2020).

In addition to the analysis of individual base-pairs, we identify in simulations frequent transi-
tions from straight to kinked conformations (Figs. 3c-d). The latter structure is reminiscent of a
kink-turn motif, but lacks its signature interactions and sequence propensities (*Klein et al., 2001*).
This large-scale motion does not affect the stem structures, but only their relative orientation. We
also note that kinked conformations are observed only when U212 is solvent exposed, and there-
fore not engaging base-pairs with U194 nor U195 (Fig. 3e), and suggest that the relative populations
could be modulated by binding of proteins or small molecules to this motif.

Stem loop 1

Similarly to SL5a, stem loop 1 (SL1) consists of two helices connected by an asymmetrical internal
loop (Fig. 4a). We use the pulling procedure to initialise our runs from the previously determined
secondary structure (*Wacker et al., 2020*) where U13-A26 are base-paired and thus represent the
first WC interaction of helix 2. During partial tempering simulations we observe a rich dynamics
involving the bulge region, together with its neighbourhood, as well as the apical loop.

While helix 1 is stable, the terminal base-pair U11-A29 either interacts through the Watson-
Crick/Watson-Crick or Hoogsteen-sugar edges (Fig. 4b). The presence of this non-canonical inter-
action allows the formation of a A12-C28 sugar-Hoogsteen interaction as well as of a U13-A27 base-
pair. A26 is not completely solvent-exposed, but forms a base triple together with A14 and U25.
In our simulations we additionally observe two alternative, lowly populated secondary structure
arrangements (Fig. 4c-d) where A12-A27/U13-A26 are base-paired, and either C28 or A29 are par-
tially excluded from the stack. Qualitatively, the nucleobase arrangements shown in Fig. 4b-d all
preserve the coaxial stacking of the two helices. Taken together, our predictions are compatible
with the NMR-derived secondary structure that does not show the presence of the U13-A26 WC
base-pair. Our models suggest the central role of A27 in mediating alternative bulge conformations
that might be required for viral viability (*Li et al., 2008*).

The apical loop is highly dynamic and lacks a major, stable conformation with specific base-base
interactions. U17-A22 base-pair interconverts from Watson-Crick/Watson-Crick to sugar/Hoogsteen,
and U18 can engage a sugar-Hogsteen interaction with C21, albeit with a small population (SI6).

Stem loop 4

Stem loop 4 (SL4) consists of four A-form helices of different lengths linked by short pyrimidine
internal loops and capped by a UGCAU pentaloop (Fig. 5a). The experimentally-derived secondary
structure is generally preserved in the parallel tempering simulations (Fig. 5b). The A-form geom-
etry between helix 1 and helix 2 is stabilised by a WC/WC C92-C119 base-pair (Fig. 5b). U95 is
solvent-exposed and transiently engages in non-canonical interactions with G94 (Fig. 5b).

In the third internal loop the two simulations behave differently suggesting slow motions that
are not fully sampled in our simulations. In the first run C100 preferentially base-pairs with U112,
while in the second run additional configurations are sampled, where either C100 or U112 are not
in the stack. The partial overlap between the two runs, that we ascribe to insufficient sampling, is
further confirmed by a principal component analysis of this internal loop region (SI7). Experimen-
tal NMR measurements on SL4 agree with A-form helix compatible arrangement of the opposing
residues C100 and U112 (*Wacker et al., 2020*), thus supporting the presence of pairing between

Figure 4. SL1 results. **a** Population of base-pairs from partial tempering simulations. **b** Three-dimensional structure of the most populated internal loop configuration showing the U11-A29 Hogsteen-Sugar, A12-C28 Sugar-Hoogsteen base-pairs, as well as the base A14-U25-U26 base triple. The corresponding secondary structure is shown on the right. The genomic numbering follows the convention in *Wacker et al.* (*2020*). The base-paired regions selected for creating initial configurations is shown in blue, while partial tempering acts in regions shown in red. **c-d** Secondary structures of two alternative internal loop arrangements.

Figure 5. SL4 results. **a** Sequence and secondary structure of SL4. The genomic numbering follows the convention in *Wacker et al.* (*2020*). The region used for constructing the A-form templates is shown in blue. The sampling in the region shown in red is enhanced by partial tempering. **b** Population of base-pairs from 20 replicas × 1.6 μs partial tempering simulations. Centroid of the most populated internal and apical loops discussed in the text are shown.

155 the two bases. The apical loop structure resembles a tetraloop with U108 solvent-exposed and
156 U104:A107 forming either a Watson-Crick/Watson-Crick or a sugar/Hogsteen non-canonical inter-
157 action, as shown in Fig. 5b.

SL2+SL3

We here report the results on the SL2+SL3 construct, consisting of two consecutive stem-loop struc-
tures: SL2 (described above) and SL3, containing the transcription-regulating sequence required
for subgenomic viral RNA synthesis (*Dufour et al., 2011*). In our simulations we sample structures
where the two helices are stacked coaxially tail-to-tail, thereby forming a unique A-form-like stem
(Fig. 6).
The SL2 loop conformations are similar to those observed for SL2 alone, with a stable C50-
G53 base-pair (see above). The UCUAAAC heptaloop in SL3 is highly flexible, and adopts several
conformations that are partially structured in terms of base-pairing (Fig. 6). Helix 2, containing the
core sequence ACGAAC is highly stable in simulations, at variance with NMR measurements (*Wacker*
et al., 2020). Since SL3 is expected to be only partially stable (*Yang and Leibowitz, 2015*), it is pos-
sible that the force-field overstabilises A-form structures, making it difficult to sample alternative
conformations.

Discussion

172 We report the prediction of the structure and conformational heterogeneity of five stem-loop ele-
173 ments, ranging in size between 17 and 44 nucleotides, located at the 5'-UTR in SARS-CoV-2 genome.

Figure 6. SL2+SL3 results. Top: Sequence and secondary structure of SL2+SL3. The genomic numbering follows the convention in *Wacker et al. (2020)*. The region used for constructing the A-form templates is shown in blue. The sampling in the region shown in red is enhanced by partial tempering. Middle panel: Population of base-pairs from partial tempering simulations. Bottom panel: Centroid of the most populated cluster.

As a starting point for our study we use the secondary structure determined by NMR experiments (*Wacker et al., 2020*).

Structures of several of these elements have also been predicted using FARFAR2 (*Rangan et al., 2020a*). A principal component analysis of the structures generated for SL1 and SL2 by FARFAR2 and MD, suggest the two methods to be complementary (SI8). In the case of SL2, the MD predictions mostly agree with the available NMR data, albeit small discrepancies remain, and for this structure it appears that the ensemble generated by MD provides a more accurate description compared to FARFAR2 (SI4). We note, however, that our simulations require significant computational resources, amounting to more than 0.5ms of aggregate simulation time.

As in most MD studies, the results presented in this work depend on the amount of sampling. We took care in identifying and discussing states that are visited multiple times during one simulation and in both replicates, indicating these conformations to correspond to *bona fide*, highly populated metastable states. Moreover, the overall high agreement between the two independent simulations suggests that the conformational ensembles are relatively well converged.

Despite the recent improvement to RNA force-fields (*Vangaveti et al., 2017*; *Šponer et al., 2018*; *Dans et al., 2019*) simulation results also depend on the accuracy of the physical models used. While we have validated our simulations against available data, we note also that it should be straightforward to integrate our extensive conformational sampling with data from future solution measurements (e.g. NMR or SAXS) using reweighting approaches (*Cesari et al., 2018*; *Bottaro and Lindorff-Larsen, 2018*; *Bottaro et al., 2020*) to refine the atomic-level description of SARS-CoV-2 RNA structure and dynamics.

Methods

MD simulations were conducted using the Amber ff99SB force field with parmbsc0 (*Pérez et al., 2007*), OL3 (*Zgarbová et al., 2011*), and corrections to the Van der Waals parameters for oxygen (*Steinbrecher et al., 2012*) in conjunction with OPC water model (*Izadi et al., 2014*). This combination has been shown to improve the description of RNA structures compared to other force-fields (*Bergonzo and Cheatham III, 2015*), and its accuracy has been assessed by independent groups (*Bottaro et al., 2018*; *Kuhrova et al., 2019*). The initial conformations were minimised in vacuum and then solvated in a dodecahedral box with minimum distance from box edge of 1.2nm. Potassium ions were added to neutralise the box (*Joung and Cheatham III, 2008*). The systems were equilibrated in the NVT and NPT ensembles at 1 bar for 1 ns. Production runs were performed in the canonical ensemble using a stochastic velocity rescaling thermostat (*Bussi et al., 2007*). All bonds were constrained using the LINCS (*Hess et al., 1997*) algorithm and equations of motion were integrated with a time step of 2 fs. Simulations were performed using GROMACS 2019.4 (*Pronk et al., 2013*) patched with PLUMED 2.5.3 (*Tribello et al., 2014*). Force-field parameters are available at http://github.com/srnas/ff/tree/opc-water-model/amber_na.ff.

Pulling simulations

The aim of the pulling simulations is quickly to generate three-dimensional RNA conformations with a specific input secondary structure. We adapt a procedure originally introduced for reconstructing atomistic structures from coarse-grained models (*Poblete et al., 2018*). Our protocol consists of three stages, as illustrated in Fig. 2a-b for SL2:

1. Plain MD simulation initialised from extended, single stranded, A-form RNA structures.
2. Pulling. We construct a reference structure with an ideal A-form double stranded RNA template for regions with known secondary structure (shown in blue in Fig. 2a) and perform pulling simulations to minimise the eRMSD (*Bottaro et al., 2014*) distance between the RNA molecule and the reference. We here use adiabatic bias MD simulations (*Marchi and Ballone, 1999*), that introduce a biasing potential damping the fluctuations only when the system moves further away from the template. In this way the pulling procedure follows the thermal

System	Lenght (nt)[a]	Simulation time [μs][b]	Acceptance rate [%] [c]	Round trips [d]
SL1	29	3.6	15-41	44-191
SL2	17	2.6	26-51	220-435
SL2+SL3	32	1.6	4-23	0-10
SL4	44	1.6	7-30	0-20
SL5a	33	3.6	12-34	6-33

Table 1. Simulated systems and related statistics. (a) number of nucleotides (nt). (b) Per-replica simulation time. Each simulation consists of 20 replicas and two independent runs for each system are carried out. The first 100ns of each run are discarded during analysis. (c) Minimum and maximum acceptance rate across all replicas and runs (d) Minimum and maximum number of round trips in temperature space across all replicas and runs.

<p>222   fluctuations of the system, and does not require a pre-defined scheduling for the moving re-</p>
<p>223   straint (*Poblete et al., 2018*). When more than one stem is present, the bias acts on multiple</p>
<p>224   templates simultaneously. The constant of the harmonic damping bias is set to 2 kJ/mol.</p>
<p>225   3. Plain MD. After 40ns of pulling simulation, the external bias is removed and the system freely</p>
<p>226   fluctuates without external forces.</p>

At the end of the procedure we extract samples from the final plain MD stage satisfying the
secondary structure constraints, i.e. with eRMSD from template < 0.75. This criterion guarantees
that all base-pairs corresponding to those in the target structure are correctly formed (Fig. 2b,
bottom panel). Note that individual pulling simulations may fail: stems are sometimes not formed
correctly and unfold during the final plain MD stage (Supporting Information 1). The success rate
of this procedure ranges from $\approx 80\%$ for SL2 to $\approx 10\%$ for SL4, which consists of 4 stem structures,
consistent with previously reported benchmarks (*Poblete et al., 2018*). During preliminary tests we
have observed that the formation of the correct secondary structure is hampered by early-forming
non-native contacts. For this reason we conduct pulling simulations at 340K.

Partial tempering

We resolvate and minimise the RNA structures obtained from pulling as described above. We use
partial tempering to enhance sampling in regions with unknown secondary structure (*Bussi, 2014*;
Ebrahimi et al., 2019). One reference replica exchanges via the Metropolis-Hastings algorithm with
parallel simulations that are conducted with a modified Hamiltonian. The modification is such that
interactions in the cold region (blue and grey in Fig. 2a, together with ions and water molecules)
are kept at the reference temperature T=300K. Interactions in the hot region (shown in red in
Fig. 2a) are rescaled by a factor $1/\lambda$, while the interactions between cold and hot regions are at an
intermediate effective temperature of $T/\sqrt{\lambda}$, with $1 \leq \lambda < 0$. In this study we use 20 geometrically
distributed replicas in the effective temperature range of 300 to 1000 K, and attempting exchanges
every 240 steps. Hot regions are shown in red in Figs. 2-6, and include the phosphate group at the
3'-end of the selected region. We have found this small addition to be beneficial in preliminary test
runs. System length, simulation time per replica, average acceptance rate, and number of round
trips in temperature space are reported in Table 1, and additional statistics are available in SI2. To
help the identification of slow degrees of freedom and assess convergence we conduct two inde-
pendent runs with different initial conformations obtained from two separate pulling simulations
(SI3).

Analysis and visualisation

254 Ideal A-form structures were constructed using the nucleic acids builder (NAB) software in Amber-
255 tools (*Macke and Case, 1998*). Trajectories were analysed using PLUMED (*Tribello et al., 2014*) and
256 MDtraj (*McGibbon et al., 2015*). Base-pair annotation and puckering angles were calculated with
257 Barnaba (*Bottaro et al., 2019*).

258 Acknowledgements

259 We acknowledge Drs. Harald Schwalbe, Anna Wacker, Julia E. Weigand and other members of
260 the Covid19-NMR consortium for discussions and insights. SB and KLL acknowledge funding from
261 the Lundbeck Foundation BRAINSTRUC structural biology initiative (R155-2015-2666). We acknowl-
262 edge access to computational resources from PRACE for the COVID-RNA project (COVID19-72).

263 Data Availability

264 Analysis scripts, MD trajectories, and input files are available at https://github.com/KULL-Centre/
265 papers/tree/master/2020/COVID_5UTR_MD PLUMED input files are available in the PLUMED NEST
266 (*Bonomi et al., 2019*) under the accession code 20.034 https://www.plumed-nest.org/eggs/20/034/.

267 References

Andrews RJ, Peterson JM, Haniff HS, Chen J, Williams C, Grefe M, Disney MD, Moss WN. An in silico map of the
SARS-CoV-2 RNA Structurome. BioRxiv. 2020; .

Bergonzo C, Cheatham III TE. Improved force field parameters lead to a better description of RNA structure.
Journal of Chemical Theory and Computation. 2015; 11(9):3969–3972.

Bergonzo C, Szakal AL. Using All-Atom Potentials to Refine RNA Structure Predictions of SARS-CoV-2 Stem
Loops. International Journal of Molecular Sciences. 2020; 21(17):6188.

Bonomi M, Bussi G, Camilloni C, Tribello GA, Banáš P, Barducci A, Bernetti M, Bolhuis PG, Bottaro S, Branduardi
D, et al. Promoting transparency and reproducibility in enhanced molecular simulations. Nature Methods.
2019; 16(8):670–673.

Bottaro S, Bengtsen T, Lindorff-Larsen K. Integrating molecular simulation and experimental data: a
Bayesian/maximum entropy reweighting approach. In: *Structural Bioinformatics* Springer; 2020.p. 219–240.

Bottaro S, Bussi G, Kennedy SD, Turner DH, Lindorff-Larsen K. Conformational ensembles of RNA oligonu-
cleotides from integrating NMR and molecular simulations. Science Advances. 2018; 4(5):eaar8521.

Bottaro S, Bussi G, Pinamonti G, Reißer S, Boomsma W, Lindorff-Larsen K. Barnaba: software for analysis of
nucleic acid structures and trajectories. RNA. 2019; 25(2):219–231.

Bottaro S, Di Palma F, Bussi G. The role of nucleobase interactions in RNA structure and dynamics. Nucleic
Acids Research. 2014; 42(21):13306–13314.

Bottaro S, Lindorff-Larsen K. Biophysical experiments and biomolecular simulations: A perfect match? Science.
2018; 361(6400):355–360.

Bussi G. Hamiltonian replica exchange in GROMACS: a flexible implementation. Molecular Physics. 2014; 112(3-
4):379–384.

Bussi G, Donadio D, Parrinello M. Canonical sampling through velocity rescaling. Journal of Chemical Physics.
2007; 126(1):014101.

Carlson CR, Asfaha JB, Ghent CM, Howard CJ, Hartooni N, Safari M, Frankel AD, Morgan DO. Phosphoregulation
of phase separation by the SARS-CoV-2 N protein suggests a biophysical basis for its dual functions. Molecular
Cell. 2020; .

Cesari A, Reißer S, Bussi G. Using the maximum entropy principle to combine simulations and solution exper-
iments. Computation. 2018; 6(1):15.

Cheng CY, Chou FC, Das R. Modeling complex RNA tertiary folds with Rosetta. In: *Methods in Enzymology*, vol.
553 Elsevier; 2015.p. 35–64.

Cheong C, Varani G, Tinoco I. Solution structure of an unusually stable RNA hairpin, 5GGAC (UUCG) GUCC.
Nature. 1990; 346(6285):680–682.

Costales MG, Childs-Disney JL, Haniff HS, Disney MD. How we think about targeting RNA with small molecules.
Journal of Medicinal Chemistry. 2020; .

302 **Dans PD**, Gallego D, Balaceanu A, Darré L, Gómez H, Orozco M. Modeling, simulations, and bioinformatics at
303 the service of RNA structure. Chem. 2019; 5(1):51–73.

304 **Dufour D**, Mateos-Gomez PA, Enjuanes L, Gallego J, Sola I. Structure and functional relevance of a transcription-
305 regulating sequence involved in coronavirus discontinuous RNA synthesis. Journal of Virology. 2011;
306 85(10):4963–4973.

307 **Ebrahimi P**, Kaur S, Baronti L, Petzold K, Chen AA. A two-dimensional replica-exchange molecular dynamics
308 method for simulating RNA folding using sparse experimental restraints. Methods. 2019; 162:96–107.

309 **Hess B**, Bekker H, Berendsen HJ, Fraaije JG. LINCS: a linear constraint solver for molecular simulations. Journal
310 of Computational Chemistry. 1997; 18(12):1463–1472.

311 **Iserman C**, Roden CA, Boerneke MA, Sealfon RS, McLaughlin GA, Jungreis I, Fritch EJ, Hou YJ, Ekena J, Weidmann
312 CA, et al. Genomic RNA elements drive phase separation of the SARS-CoV-2 nucleocapsid. Molecular Cell.
313 2020; .

314 **Izadi S**, Anandakrishnan R, Onufriev AV. Building water models: a different approach. The Journal of Physical
315 Chemistry Letters. 2014; 5(21):3863–3871.

316 **Joung IS**, Cheatham III TE. Determination of alkali and halide monovalent ion parameters for use in explicitly
317 solvated biomolecular simulations. Journal of Physical Chemistry B. 2008; 112(30):9020–9041.

318 **Jucker FM**, Pardi A. Solution structure of the CUUG hairpin loop: a novel RNA tetraloop motif. Biochemistry.
319 1995; 34(44):14416–14427.

320 **Klein D**, Schmeing T, Moore P, Steitz T. The kink-turn: a new RNA secondary structure motif. The EMBO journal.
321 2001; 20(15):4214–4221.

322 **Kuhrova P**, Mlynsky V, Zgarbová M, Krepl M, Bussi G, Best RB, Otyepka M, Sponer J, Banas P. Improving the
323 performance of the amber RNA force field by tuning the hydrogen-bonding interactions. Journal of Chemical
324 Theory and Computation. 2019; 15(5):3288–3305.

325 **Lal SK**. Molecular Biology of the SARS-Coronavirus. Springer Science & Business Media; 2010.

326 **Lee CW**, Li L, Giedroc DP. The solution structure of coronaviral stem-loop 2 (SL2) reveals a canonical CUYG
327 tetraloop fold. FEBS Letters. 2011; 585(7):1049–1053.

328 **Leontis NB**, Westhof E. Geometric nomenclature and classification of RNA base pairs. RNA. 2001; 7(4):499–512.

329 **Li L**, Kang H, Liu P, Makkinje N, Williamson ST, Leibowitz JL, Giedroc DP. Structural lability in stem–loop 1 drives
330 a 5 UTR–3 UTR interaction in coronavirus replication. Journal of Molecular Biology. 2008; 377(3):790–803.

331 **Macke TJ**, Case DA. Modeling unusual nucleic acid structures. In: *Molecular Modeling of Nucleic Acids* ACS
332 Publications; 1998.p. 379–393.

333 **Manfredonia I**, Nithin C, Ponce-Salvatierra A, Ghosh P, Wirecki TK, Marinus T, Ogando NS, Snijder EJ, van
334 Hemert MJ, Bujnicki JM, et al. Genome-wide mapping of SARS-CoV-2 RNA structures identifies therapeutically-
335 relevant elements. Nucleic acids research. 2020; .

336 **Marchi M**, Ballone P. Adiabatic bias molecular dynamics: a method to navigate the conformational space of
337 complex molecular systems. The Journal of Chemical Physics. 1999; 110(8):3697–3702.

338 **McGibbon RT**, Beauchamp KA, Harrigan MP, Klein C, Swails JM, Hernández CX, Schwantes CR, Wang LP, Lane
339 TJ, Pande VS. MDTraj: a modern open library for the analysis of molecular dynamics trajectories. Biophysical
340 Journal. 2015; 109(8):1528–1532.

341 **Miao Z**, Tidu A, Eriani G, Martin F. Secondary structure of the SARS-CoV-2 5'-UTR. RNA Biology. 2020; p. 1–10.

342 **Mráziková K**, Mlynsky V, Kuhrova P, Pokorná P, Kruse H, Krepl M, Otyepka M, Banas P, Sponer J. UNCG RNA
343 tetraloop as a formidable force-field challenge for MD simulations. bioRxiv. 2020; .

344 **Parisien M**, Major F. The MC-Fold and MC-Sym pipeline infers RNA structure from sequence data. Nature.
345 2008; 452(7183):51–55.

346 **Pérez A**, Marchán I, Svozil D, Sponer J, Cheatham III TE, Laughton CA, Orozco M. Refinement of the AMBER
347 Force Field for Nucleic Acids: Improving the Description of alpha gamma Conformers. Biophysical Journal.
348 2007; 92(11):3817–3829.

Poblete S, Bottaro S, Bussi G. Effects and limitations of a nucleobase-driven backmapping procedure for nucleic
acids using steered molecular dynamics. Biochemical and Biophysical Research Communications. 2018;
498(2):352–358.

Pronk S, Páll S, Schulz R, Larsson P, Bjelkmar P, Apostolov R, Shirts MR, Smith JC, Kasson PM, van der Spoel D,
Hess B, Lindhal E. GROMACS 4.5: a high-throughput and highly parallel open source molecular simulation
toolkit. Bioinformatics. 2013; 29(7):845–854.

Rangan R, Watkins AM, Kladwang W, Das R. De novo 3D models of SARS-CoV-2 RNA elements and small-
molecule-binding RNAs to guide drug discovery. BioRxiv. 2020; .

Rangan R, Zheludev IN, Hagey RJ, Pham EA, Wayment-Steele HK, Glenn JS, Das R. RNA genome conservation
and secondary structure in SARS-CoV-2 and SARS-related viruses: a first look. RNA. 2020; 26(8):937–959.

Schubert K, Karousis ED, Jomaa A, Scaiola A, Echeverria B, Gurzeler LA, Leibundgut M, Thiel V, Mühlemann
O, Ban N. SARS-CoV-2 Nsp1 binds the ribosomal mRNA channel to inhibit translation. Nature Structural &
Molecular Biology. 2020; 27(10):959–966.

Šponer J, Bussi G, Krepl M, Banáš P, Bottaro S, Cunha RA, Gil-Ley A, Pinamonti G, Poblete S, Jurečka P, Walter
NG, Otyepka M. RNA structural dynamics as captured by molecular simulations: a comprehensive overview.
Chemical Reviews. 2018; 118(8):4177–4338.

Steinbrecher T, Latzer J, Case D. Revised AMBER Parameters for Bioorganic Phosphates. Journal of Chemical
Theory and Computation. 2012; 8(11):4405–4412.

Stelzer AC, Frank AT, Kratz JD, Swanson MD, Gonzalez-Hernandez MJ, Lee J, Andricioaei I, Markovitz DM, Al-
Hashimi HM. Discovery of selective bioactive small molecules by targeting an RNA dynamic ensemble. Nature
Chemical Biology. 2011; 7(8):553.

Tribello GA, Bonomi M, Branduardi D, Camilloni C, Bussi G. PLUMED 2: New feathers for an old bird. Computer
Physics Communications. 2014; 185(2):604–613.

Vangaveti S, Ranganathan SV, Chen AA. Advances in RNA molecular dynamics: a simulator's guide to RNA
force fields. Wiley Interdisciplinary Reviews: RNA. 2017; 8(2):e1396.

Wacker A, Weigand JE, Akabayov SR, Altincekic N, Bains JK, Banijamali E, Binas O, Castillo-Martinez J, Cetiner E,
Ceylan B, et al. Secondary structure determination of conserved SARS-CoV-2 RNA elements by NMR spec-
troscopy. Nucleic Acids Research. 2020; .

Wang L, Friesner RA, Berne B. Replica exchange with solute scaling: a more efficient version of replica exchange
with solute tempering (REST2). The Journal of Physical Chemistry B. 2011; 115(30):9431–9438.

Watkins AM, Rangan R, Das R. FARFAR2: Improved de novo Rosetta prediction of complex global RNA folds.
Structure. 2020; .

Yang D, Leibowitz JL. The structure and functions of coronavirus genomic 3 and 5 ends. Virus Research. 2015;
206:120–133.

Zgarbová M, Otyepka M, Sponer J, Mlaadek A, Banas P, Cheatham III TE, Jurecka P. Refinement of the Cornell et
al. nucleic acids force field based on reference quantum chemical calculations of glycosidic torsion profiles.
Journal of Chemical Theory and Computation. 2011; 7(9):2886–2902.