Benzoquinones in the defensive secretion of a bug (*Pamillia behrensii*): a common chemical

trait retrieved in the Heteroptera

Julian M. Wagner[#], Thomas H. Naragon[#] and Adrian Brückner*

Division of Biology and Biological Engineering, California Institute of Technology, 1200 East

California Boulevard, Pasadena, CA 91125, United States of America

[#] both authors contributed equally

* corresponding author: adrian.brueckner@gmail.com

Abstract

Benzoquinones are a phylogenetically widespread compound class within arthropods,

appearing in harvestman, millipedes and insects. Whereas the function of benzoquinones as

defensive compounds against potential predators and microbes has been well established, the full

extent of benzoquinone usage across arthropods, and especially within Insecta, has yet to be

established. Adding to the growing list of unique evolutionary origins of benzoquinone

employment, we describe in this paper the metathoracic scent gland secretion of the mirid bug

Pamillia behrensii, which is composed of heptan-2-one, 2-heptyl acetate, 2,3-dimethyl-1-4-

benzoquinone, 2,3-dimethyl-1-4-hydroquinone as well as one unknown compound. Similarly, to

many other arthropods that use benzoquinones, *Pamillia* releases the contents of its gland as a

defensive mechanism in response to harassment by other arthropod predators. Morphological

investigation of the gland showed that the benzoquinone-producing gland complex of *P. behrensii*

follows a similar blueprint to metathoracic scent glands described in other Heteropterans. Overall,

our data further underpins the widespread convergent evolution and use of benzoquinones for

defense across the Arthropoda, now including the order Hemiptera.

Keywords

Quinones, chemical defense, exocrine glands, convergence, evolutionary biology

Introduction

Chemical convergence is the evolution and widespread use of certain compound classes by different taxa for similar purposes (Beran et al. 2019; Brückner and Parker 2020). Evolutionary patterns of this nature suggest potential genetic biases at the biosynthetic level or the efficiency of certain compound classes at eliciting a given response in other organisms within certain ecological settings as drivers of selection (Brückner and Parker 2020; Chevrette et al. 2020). '

One striking example of chemical convergence is the frequent evolution of aromatic benzoquinones (BQs) as defensive compounds in arthropods (**Fig 1A**). The past 60 years of research has uncovered a smorgasbord of quinone containing glands in insects, harvestment, and millipedes (e.g., Blum 1981; Meinwald et al. 1966; Raspotnig et al. 2017; Roth and Stay 1958; Shear 2015). In insects, BQs have a broad taxonomic distribution, spanning both hemi- and holometabolous insects, with representatives in the earwigs (Dermaptera), lubber grasshoppers (Orthoptera), cockroaches (Blattodea), caddisflies (Trichoptera) and most prominently beetles (Coleoptera) (Blum 1981; Eisner et al. 2005; Francke and Dettner 2005).

Despite the similarity of the chemical phenotype, it is yet to be determined if the biosynthetic pathways for the BQs are also convergent (Blum 1981; Morgan 2010). Some have argued that BQs may be produced by enzymes related to tyrosine dependent cuticle tanning (Duffey 1974; He et al. 2018; Roth and Stay 1958), while others have demonstrated the possibility of *de novo* BQ synthesis from poly-β-carbonyls via head-to-tail condensation (Meinwald et al. 1966; Morgan 2010; Rocha et al. 2013). In contrast, the function of BQs in chemical defense

against predators and as antimicrobial agents is non-controversial and widely documented (e.g.,

Eisner et al. 2005; Gasch et al. 2013; Li et al. 2013; Ruther et al. 2001; Shear 2015).

Here we provide another example for the widespread evolution and chemical convergence

of benzoquinones, and report on the first evidence of this compound class in the Heteroptera

(Aldrich 1988). We chemically analyzed metathoracic scent gland (MTG) secretion of the mirid

bug *Pamillia behrensii,* Uhler (**Fig 2D**), investigated the internal and external morphology of the

gland via confocal microscopy and investigated the biological function of MTG compounds.

Materials and Methods

Collection

Individual *Pamillia behrensii* were collected in close proximity to nests of the velvety tree

ant *(Liometopum occidentale)* on October 23rd 2019 and November 17th 2020 at Chaney Canyon,

Altadena, CA (34°13'01.7"N 118°09'15.1"W). Insects were carefully transferred to the laboratory

in tubes lined with moist tissue paper and directly used for chemical analysis or imaging.

Chemical analysis

To identify compounds stored in the metathoracic scent gland (MTG) of *Pamillia*

behrensii we used whole body extraction in 50 µl hexane for 10 min, which is a well-established

method in Heteropterans (e.g., Leal et al. 1994; Zarbin et al. 2000). Crude hexane sample aliquots

(1 µl) were analyzed on a GCMS-QP2020 gas chromatography/mass-spectrometry system

(Shimadzu, Kyōto, Japan) equipped with a ZB-5MS fused silica capillary column (30 m x 0.25

mm ID, df= 0.25 µm) from Phenomenex (Torrance, CA, USA). Samples were injected using an

AOC-20i autosampler system from Shimadzu into a split/splitless-injector operating in splitless-

mode at a temperature of 310°C. Helium was used as the carrier-gas with a constant flow rate of

2.13 ml/min. The chromatographic conditions were as follows: The column temperature at the start

was 40°C with a 1-minute hold after which the temperature was initially increased 30°C/min to

250°C and further increased 50°C/min to a final temperature of 320°C and held for 5 minutes.

Electron impact ionization spectra were recorded at 70 eV ion source voltage, with a scan rate of

0.2 scans/sec from *m/z* 40 to 450. The ion source of the mass spectrometer and the transfer line

were kept at 230°C and 320°C, respectively.

We integrated chromatograms manually using LabSolutions Postrun Analysis (Shimadzu,

Kyōto, Japan), quantified the ion abundance and calculated the relative composition of individual

compounds compared to the total ion abundance. Compounds were identified based on the *m/z*

fragmentation patterns and retention indices (RIs) calculated after Van den Dool and Kratz (1963)

using a standard alkane mixture. Synthetic heptan-2-one was purchased from Sigma-Aldrich (St.

Louis, MO, USA) and natural 2-ethyl-1-4-benzoquinone and 2-ethyl-1-4-hydroquinone from the

secretion of *Tribolium castaneum* (Li et al. 2013) were used as reference compounds. 2-Heptyl

acetate was synthesized by mixing acetic acid with an excess of 2-heptanol and H_2SO_4 (all Sigma-

Aldrich) as catalyst and subsequently heating the mixture for 48 h at 45 °C. The crude ester was

extracted with hexane and the organic layer was washed with water, saturated Na_2CO_3 solution

(2x) and saturated NaCl solution and finally dried over anhydrous Na_2SO_4.

Histochemistry and imaging

Adult bugs were immersed in PBS and legs were removed with forceps. The dorsal parts

of the thorax were removed by cutting around the abdominal margin with dissection scissors and

the ventral thorax was fixed in 4% paraformaldehyde (25 minutes, room temperature), washed in

PBS+0.02% Triton X100, and subsequently stained with Alexa-647-Phalloidin (Thermo Fisher,

Waltham, MA, USA) to label muscles and Alexa-546-WGA (Thermo Fisher, Waltham, MA,

USA) to stain membranes. Metathoracic scent glands from both sides were imaged as whole

mounts of ventral thoraxes in ProLong Gold Antifade Mountant (Thermo Fisher, Waltham, MA,

USA), using a Zeiss LSM 880 (Carl Zeiss, Jena, Germany) with airyscan.

Behavioral assay

To confirm that P. behrensii deploys gland contents in response to threat we performed a

behavioral trial in tandem with solid phase microextraction (SPME). A 4-cm circular behavioral

arena was constructed from 1/8th inch infrared transmitting acrylic (Plexiglass IR acrylic 3143)

with a port on the side for a 65 µm polydimethylsiloxane/divinylbenzene fiber from Supelco

(Sigma-Aldrich; St. Louis, MO) to sample released volatiles. For compound desorption the fiber

was placed in the injector port for 1 min operating at 230°C and the GC/MS run was carried out

as outlined below. After each trial the fiber was baked for 30 min at 230°C. SPME-GC/MS runs

were used to identify the fraction of volatile metathoracic scent gland (MTG) compounds (see e.g.

Krajicek et al. 2016).

Behavior was recorded for thirty minutes at 25 fps with a machine vision camera (Flir

BFS-U3-16S2M-CS). Bugs were anesthetized on ice and placed in an arena either alone, with five

L. occidentale, or with five *L. occidentale* with superglued mandibles to reduce ant aggressiveness.

To assess behavior, bugs were labeled with DeepLabCut (DLC) to quantify movement over time

(Mathis et al. 2018). Median filtered, labeled bug locations were used to sum to total distance

traversed during the trial. SPME-GC/MS runs were used to identify the fraction of volatile

metathoracic scent gland (MTG) compounds (see e.g. Krajicek et al. 2016). A linear regression

was performed on 10,000 bootstrap samples of the data in python to calculate confidence intervals

for the regression slope. A Jupyter notebook is available which reproduces the analysis at

https://github.com/julianmwagner/bq_bug.

Results and Discussion

In total, we found five compounds in the MTG extracts of *Pamillia behrensii* of which we were able to identify four (**Fig 2A; Tab 1**). Compound I elicited a prominent fragment ion at $m/z=$ 58 arising from McLafferty rearrangement, which is characteristic for saturated methyl-ketones, together with a molecular ion at $m/z=$ 114 we identified I as heptan-2-one and eventually confirmed our identification with a synthetic standard.

Compound II showed a molecular ion at $m/z=$ 158 and a base ion at $m/z=$ 43 indicating an acetate ester [additional R'COOH$_2$ at $m/z=$ 61; Urbanová et al. (2012)] with a heptanol alcohol moiety (ion at m/z= 115). Both diagnostic ions at $m/z=$87 and the pair at $m/z=$69/70 indicate a possible heptan-2-ol moiety (**Tab 1**). Hence, we tentatively identified II as 2-heptyl acetate (1-methylhexyl acetate) and eventually confirmed the compound's identity by synthesis.

Compound III showed molecular ions at $m/z=$ 136 as well as diagnostic ions at $m/z=$ 108, $m/z=$ 82, $m/z=$ 79 and $m/z=$ 54, indicating either a 2,3-dimethyl- or 2-ethyl- substituted 1-4-benzoquinone. An ion at $m/z=$ 107 (i.e. 5-methyl-4-methylidenecyclopentenone ion) and III's retention index (**Tab 1**) were, however, more indicative of 2,3-dimethyl-1-4-benzoquinone. Compound V showed a molecular ions at $m/z=$ 138 (also the base ion) and diagnostic ions at $m/z=$ 137 $m/z=$ 95, as well as $m/z=$ 91 which together with V's retention index (**Tab 1**), were all indicative of 2,3-dimethyl-1-4-hydroquinone. We used the secretion from defensive stink gland of *Tribolium castaneum* (Li et al. 2013) which contains 2-ethyl-1,4-benzoquinone and 2-ethyl-1,4-hydroquinone as a natural standard and found no correspondence to III and V. Hence, we eventually identified III as 2,3-dimethyl-1,4-benzoquinone and V as 2,3-dimethyl-1,4-hydroquinone.

For compound IV we did not find any close hit in any compound library used (NIST14,

Wiley-NIST 2009, FCSN2) and the identity of this compound remains to be elucidated (see **Tab**

1 for *m/z* fragmentation). The MTG extracts of *P. behrensii* (**Fig 2A**), overall showed a very

consistent chemical composition, with heptan-2-one always appearing in the greatest abundance,

2,3-dimethyl-1,4-benzoquinone at an intermediate abundance and 2-heptyl acetate, unknown IV

as well 2,3-dimethyl-1-4-hydroquinone being minor compounds (**Fig 2B; Tab 1**).

Based on synthetic heptan-2-one and 2-methyl-1,4-benzoquinone, as external standards,

we estimated 14.1±5.8 µg of heptan-2-one and 4.9±2.2 µg of 2,3-dimethyl-1,4-benzoquinone per

individual, respectively. Based on the composition and amounts, we interpreted heptan-2-one as a

solvent for the more noxious benzoquinone and hydroquinone, while the ester 2-heptyl acetate

may be an additional repellent or act as surfactant as previously described for glandular secretions

of rove beetles (Dettner 1984; Steidle and Dettner 1993).

Detecting 2,3-dimethyl-1,4-benzoquinone in the MTG extracts of a heteropteran (**Fig 2A**),

adds another order of arthropods to the list of BQ-producers and underpins the remarkable pattern

of chemical convergent evolution of BQs. Even though other aromatics compounds (e.g. vanillin,

2-phenylethanol, benzyl alcohol, or p-hydroxbenzaldehyde) have been described in other bugs,

benzoquinones add another compound class to the chemical skillset of the Heteroptera. Focusing

on aliphatic compounds, MTG secretions of bugs are typically composed of n-alkanes, alkenals,

alkenyl alkanoates and occasionally monoterpenoids. While heptan-2-one appears to be a rather

unusual compound for bugs as well, methyl ketones are common in arthropods. For instance, in

ants, other hymenopterans, trichopterans as well as cyphophthalmid and dyspnoan harvestman

they serve as alarm and sex pheromones or as defensive compounds (Blum 1981; Cheng et al.

2017; Löfstedt et al. 2008; Raspotnig et al. 2005; Schaider et al. 2018).

2,3-Dimethyl-1-4-hydroquinone in the MTG secretions of *P. behrensii* (**Fig 2A**) can be

interpreted as a precursor molecule that is oxidized to the final benzoquinone *via* a laccase

enzymes, as a recent transcriptomic study in termite soldiers suggested (He et al. 2018). Whether

the hydroquinone is derived from tyrosine or was produced utilizing a polyketide-like mechanism

(Brückner and Parker 2020) is, however, unknown.

As is typical for chemical release from the metathoracic scent gland of other bugs, we

found that the yellow, quinone secretion is also expelled from this gland complex as small droplets

(**Fig 2C**). The MTG was located between thorax segment II and III (**supplement Fig S1A**) and

generally showed an outer morphology comparable to other Heteropteran glands (Aldrich 1988;

Gonzaga-Segura et al. 2013; Hepburn and Yonke 1971): two adjacent cuticular folds form

flaps/lips on the anterior and posterior edges of the gland opening (**Fig 2C**). The edges of the

cuticular folds constitute the ostiolar peritreme, which likely aids in carrying the glandular

secretions away from the opening and fosters increased compound dispersion (**Fig 2C**).

Additionally, the roughly sculpted cuticle around the gland opening serves as an evaporation

surface (**Fig 2C**), increasing the volatility of expelled MTG compounds (Aldrich 1988; Hepburn

and Yonke 1971).

The internal part of the MTG complex (**Fig S1B and C**) consists of a lateral extension

which connects the gland opening to the median reservoir where the secretions produced by

primary gland are stored (Aldrich 1988; Hepburn and Yonke 1971). Despite the capability of the

MTG of *P. behrensii* to synthesize quite unusual Heteropteran compounds, the general outer and

inner morphology of the MTG complex appears to be similar to that found in other MTG

containing species and we did not find any specialized adaptions to house these compounds.

To confirm *P. behrensii* uses its MTG secretion (**Fig 2A**) as a defensive compound,

we performed behavioral experiments coupled with chemical headspace profiling with solid phase

microextraction. We expected the chemical secretion expelled from the gland should function as

an allomone in chemical defense, as quinones are known to be noxious and serve as predator

repellents in many other arthropods (e.g., Eisner et al. 2005; Gasch et al. 2013; Li et al. 2013;

Ruther et al. 2001). Like other chemicals, benzoquinones activate the transient receptor potential

ankyrin 1 (TRPA1), a nonselective cation channel, which has a conserved function as a noxious

chemical receptor in animals (Arenas et al. 2017; Ibarra and Blair 2013). TRPA1 might thus be a

major target of many convergently evolved defensive chemicals like benzoquinones and initiate

protective behavior after a chemical attack (Blair et al. 2016; Ibarra and Blair 2013).

We found direct evidence of active usage of the MTG gland for defense in our behavioral

recordings. In cases where ants successfully grabbed the bug, deployed gland content could be

seen smeared on the behavioral arena (**Fig 3A**). The compounds quickly evaporated off the arena

surface and were readily measurable *via* SPME-GCMS. We additionally profiled the correlation

of amount of benzoquinone deployed with bug movement in the behavioral arena (**Fig 3B**). We

found a positive correlation (95% confidence interval of slope: 0.154-1.75) of BQ amount with the

distance the bug traveled during the behavioral trial (a proxy bug agitation/escape response). This

behavioral readout supports the idea that *P. behrensii* deploys its gland to confuse or repel its

predators before fleeing. We thus demonstrated that *P. behrensii* uses the compounds stored in its

MTG as a repellent against predators, which again underpins the chemical convergence of

benzoquinones.

We found first evidence of a benzoquinone in the Heteroptera (Aldrich 1988; Morgan

2010). More specifically we detected 2-ethyl-1-4-benzoquinone produced by and stored in the

metathoracic scent gland complex of the mirid bug *Pamillia behrensii* and showed that it uses this

BQ in combination with three other compounds for chemical defense against predators. Our study

therefore highlights the remarkable convergent evolution of benzoquinones as defensive

compounds across the Arthropoda, now including true bugs.

Acknowledgment

208 We thank Joe Parker for making his laboratory available to us. JW was supported by NN.

209 THN received a NN. AB is Simons Fellow of the Life Sciences Research Foundation (LSRF).

Ethics statement

211 There are no legal restrictions on working with bugs. Field collection permissions were

212 issues by California Department of Fish and Wildlife and the Angeles National Forest (US Forest

213 Service; USDA).

Authors contributions

215 JMW, THN and AB design research; JMW and THN collected specimens; THN performed

216 microscopy; JMW performed and analyzed behavioral assays; JMW collected chemical data; AB

217 analyzed chemical data and performed organic synthesis; AB supervised the project; AB wrote the

218 paper with input from THN and JMW. JMW and THN contributed equally. All authors gave final

219 approval for publication.

References

Aldrich J (1988) Chemical ecology of the Heteroptera Annual Review of Entomology 33:211-238

Arenas OM, Zaharieva EE, Para A, Vásquez-Doorman C, Petersen CP, Gallio M (2017) Activation of planarian TRPA1 by reactive oxygen species reveals a conserved mechanism for animal nociception Nature neuroscience 20:1686-1693

Beran F, Köllner TG, Gershenzon J, Tholl D (2019) Chemical convergence between plants and insects: biosynthetic origins and functions of common secondary metabolites New Phytologist

Bertoli A, Leonardi M, Krzyzanowska J, Oleszek W, Pistelli L (2011) Mentha longifolia in vitro cultures as safe source of flavouring ingredients Acta Biochimica Polonica 58

Blair NT, Philipson BI, Richards PM, Doerner JF, Segura A, Silver WL, Clapham DE (2016) Naturally produced defensive alkenal compounds activate TRPA1 Chemical senses 41:281-292

Blum MS (1981) Chemical Defenses of Arthropods Academic Press, Inc., New York

Brückner A, Parker J (2020) Molecular evolution of gland cell types and chemical interactions in animals Journal of Experimental Biology 223

Cheng Y-n, Wen P, Dong S-h, Tan K, Nieh JC (2017) Poison and alarm: the Asian hornet Vespa velutina uses sting venom volatiles as an alarm pheromone Journal of experimental biology 220:645-651

Chevrette MG et al. (2020) Evolutionary dynamics of natural product biosynthesis in bacteria Nat Prod Rep 37:566-599

Dettner K (1984) Isopropylesters as wetting agents from the defensive secretion of the rove beetle Coprophilus striatulus F.(Coleoptera, Staphylinidae) Insect biochemistry 14:383-390

243   Duffey SS (1974) The biosynthesis of defensive chemicals by millipedes: parrallelism with plant

244   biosynthetic pathways. University of British Columbia

245   Eisner T, Eisner M, Siegler M (2005) Secret Weapons. Secret Weapons. Harvard University Press,

246   Cambridge

247   Francke W, Dettner K (2005) Chemical signalling in beetles. In: The Chemistry of Pheromones

248   and Other Semiochemicals II. Springer, pp 85-166

249   Gasch T, Schott M, Wehrenfennig C, Düring R-A, Vilcinskas A (2013) Multifunctional weaponry:

250   the chemical defenses of earwigs Journal of insect physiology 59:1186-1193

251   Gonzaga-Segura J, Valdez-Carrasco J, Castrejon-Gomez VR (2013) The metathoracic scent gland

252   of the leaf-footed bug, Leptoglossus zonatus Journal of Insect Science 13:149

253   He S et al. (2018) Termite soldiers contribute to social immunity by synthesizing potent oral

254   secretions Insect molecular biology 27:564-576

255   Hepburn H, Yonke T (1971) The metathoracic scent glands of coreoid Heteroptera Journal of the

256   Kansas Entomological Society:187-210

257   Ibarra Y, Blair NT (2013) Benzoquinone reveals a cysteine-dependent desensitization mechanism

258   of TRPA1 Molecular pharmacology 83:1120-1132

259   Krajicek J et al. (2016) Comparative analysis of volatile defensive secretions of three species of

260   Pyrrhocoridae (Insecta: Heteroptera) by gas chromatography-mass spectrometric method

261   PloS one 11:e0168827

262   Leal WS, Panizzi AR, Niva CC (1994) Alarm pheromone system of leaf-footed bugLeptoglossus

263   zonatus (Heteroptera: Coreidae) Journal of Chemical Ecology 20:1209-1216

Li J, Lehmann S, Weißbecker B, Naharros IO, Schütz S, Joop G, Wimmer EA (2013) Odoriferous

defensive stink gland transcriptome to identify novel genes necessary for quinone synthesis

in the red flour beetle, Tribolium castaneum PLoS genetics 9:e1003596

Löfstedt C, Bergmann J, Francke W, Jirle E, Hansson BS, Ivanov VD (2008) Identification of a

sex pheromone produced by sternal glands in females of the caddisfly Molanna angustata

Curtis Journal of chemical Ecology 34:220

Mathis A, Mamidanna P, Cury KM, Abe T, Murthy VN, Mathis MW, Bethge M (2018)

DeepLabCut: markerless pose estimation of user-defined body parts with deep learning

Nature neuroscience 21:1281-1289

Meinwald J, Koch KF, Rogers Jr JE, Eisner T (1966) Biosynthesis of arthropod secretions. III.

Synthesis of simple p-benzoquinones in a beetle (*Eleodes longicollis*) Journal of the

American Chemical Society 88:1590-1592

Misof B et al. (2014) Phylogenomics resolves the timing and pattern of insect evolution Science

346:763-767

Morgan ED (2010) Biosynthesis in Insects. RSC Publishing, Camebridge

Raspotnig G, Fauler G, Leis M, Leis HJ (2005) Chemical profiles of scent gland secretions in the

cyphophthalmid opilionid harvestmen, *Siro duricorius* and *S. exilis* Journal of Chemical

Ecology 31:1353-1368 doi:10.1007/s10886-005-5291-4

Raspotnig G, Schaider M, Föttinger P, Schönhofer A (2017) A model for phylogenetic

chemosystematics: evolutionary history of quinones in the scent gland secretions of

harvestmen Frontiers in ecology and evolution 5:139

Rocha D, Wouters F, Zampieri D, Brocksom T, Machado G, Marsaioli A (2013) Harvestman

phenols and benzoquinones: characterisation and biosynthetic pathway Molecules

18:11429-11451

Roth LM, Stay B (1958) The occurrence of para-quinones in some arthropods, with emphasis on

the quinone-secreting tracheal glands of *Diploptera punctata* (Blattaria) Journal of Insect

Physiology 1:305-318

Ruther J, Podsiadlowski L, Hilker M (2001) Quinones in cockchafers: additional function of a sex

attractant as an antimicrobial agent Chemoecology 11:225-229

Schaider M, Novak T, Komposch C, Leis H-J, Raspotnig G (2018) Methyl-ketones in the scent

glands of Opiliones: a chemical trait of Cyphophthalmi retrieved in the dyspnoan

Nemastoma triste Chemoecology 28:61-67

Shear WA (2015) The chemical defenses of millipedes (diplopoda): Biochemistry, physiology and

ecology Biochemical Systematics and Ecology 61:78-117 doi:10.1016/j.bse.2015.04.033

Steidle JLM, Dettner K (1993) Chemistry and morphology of the tergal gland of freeliving adult

Aleocharinae (Coleoptera: Staphylinidae) and its phylogenetic significance Systematic

Entomology 18:149-168

Stork NE (2018) How many species of insects and other terrestrial arthropods are there on Earth?

Annual review of entomology 63:31-45

Urbanová K, Vrkoslav V, Valterová I, Háková M, Cvacka J (2012) Structural characterization of

wax esters by electron ionization mass spectrometry Journal of lipid research 53:204-213

Zarbin PH, Borges M, Santos AAd, de Oliveira AR, Simonelli F, Marques FdA (2000) Alarm

pheromone system of stink bug Piezodorus guildinii (Heteroptera: Pentatomidae) Journal

of the Brazilian Chemical Society 11:424-428

Table 1 Gas chromatographic and mass spectrometric data of the metathoracic scent gland secretion released by the mirid bug *Pamillia*

behrensii. Molecular ions (M+) are marked in bold. Retention indices (RI) were calculated according to van den Dool and Kratz (1963).

RI	RI literature	Identified as	Mass spectrometric fragmentation m/z (relative intensity)	Secretion profile [%]
890	892[a]	heptan-2-one	114 (M^+, 4), 99 (3), 85 (3), 71 (15), 58 (55), 43 (100)	64.1±5.2
1041	1043[b]	2-heptyl acetate	158 (M^+, 1), 115 (3), 102 (2), 98 (7), 87 (22), 70 (8), 69 (7), 61 (3), 56 (15), 43 (100)	6.5±6.5
1118	1119[c]	2,3-dimethyl-1-4-benzoquinone	136 (M^+, 85), 108 (71), 107 (59), 82 (59), 80 (31), 79 (76), 54 (100), 53 (44)	26.5±4.5
1223	-	unknown	138 (M^+, 52), 110 (14), 95 (16), 82 (29), 67 (40), 54 (100)	2.1±1.6
1432	1433[c]	2,3-dimethyl-1-4-hydroquinone	138 (M^+, 100), 137 (25), 123 (53), 95 (29), 91 (25)	0.8±0.7

[a]RI as reported by Schaider et al. (2018)

[b]RI as reported by Bertoli et al. (2011)

[c]RI as reported by Rocha et al. (2013)

Figure 1 Convergent evolution of benzoquinones across the arthropods. Conceptual

phylogenetic tree of arthropods - focusing on insects - depicting taxa producing benzoquinones

based on previous research (in red) and results presented in this study (in orange). The tree

topology is based on Misof et al. (2014) and the group size is proportional to the number of species

based on Stork (2018).

Figure 2 Metathoracic scent gland (MTG) chemistry of *Pamillia behrensii*. A: Example GC

traces of *P. behrensii* gland exudates. Compounds are labeled in the trace and mass spectrometric

data is detailed in Table 1. **B:** Matrix plot shows the composition of eleven individuals. The gland

compounds in the order of their retention indices are heptan-2-one, 2-heptyl acetate, 2-ethyl-1,4-

benzoquinone, unknown IV and 2-ethyl-1,4-hydroquinone. The legend indicates the percentage

composition for each compound per specimen. **C:** Outer morphology of MTG of *P. behrensii.* The

top image shows a droplet of the glandular secretion shortly after gland use. The yellow color

indicates the presence of benzoquinones. The bottom image details the surface morphology around

the gland opening. op= gland opening; evp= evaporation surface; per= peritreme; can= canal from

ostiole to coxal pit. **D:** Habitus image of *P. behrensii* with scale bar.

Figure 3 Behavioral evidence active defense gland use against predators. A: When placed in

a behavioral arena ants attack *Pamillia behrensii* (left panel of A). In response to aggression, *P.*

behrensii deploys defensive gland secretion, which can be seen by eye (A central panel, indicated

with arrow). The gland secretion volatilizes, diffusing to nearby attackers (A right panel); volatile

compound identity was confirmed *via* SPME-GCMS. **B:** Quantitative analysis of behavior

indicated a significant positive correlation of bug movement (a proxy for bug agitation/attempts to

flee predators) with benzoquinone quantity deployed. Grey intervals represent 95th, 75th and 10th

percentile for regression confidence intervals from 10,000 bootstrap replicates. Together, these

suggest that *P. behrensii* employs a joint strategy of gland deployment with an escape response to

evade aggressive predators.

Supplementary Figure S1 Overview of the morphology of the *P. behrensii* gland. **A:** Location

of the metathoracic scent gland (MTG) opening located between thorax segment II and II. **B** + **C:**

Inner morphology of the MTG showing the median reservoir where the secretions produced by

primary gland are stored.

CHELICERATA
MYRIAPODA
PROTURA
COLLEMBOLA
DIPLURA
ARCHAEOGNATHA
ODONATA
EPHEMEROPTERA
ZYGENTOMA
PLECOPTERA
ORTHOPTERA
EMBIOPTERA
GRYLLOBLATTODEA
ZORAPTERA
PHASMATODEA
MANTODEA
BLATTODEA
THYSANOPTERA
HEMIPTERA
PSOCODEA
HYMENOPTERA
RAPHIDIOPTERA
MEGALOPTERA
NEUROPTERA
STREPSIPTERA
COLEOPTERA
TRICHOPTERA
LEPIDOPTERA
SIPHONAPTERA
MECOPTERA
DIPTERA

(A)

relative intensity [%]

I — heptan-2-one

II

III

IV

V

2x

retention time

*

(B)

%

I - heptan-2-one

II - 2-heptyl acetate

III - 2-ethyl-1,4-BQ

IV - unknown

V - 2-ethyl-1,4-HQ

(C)

evp

op

can

per

(D)

1 mm

(A) ant attack   gland used   volatiles diffuse

(B)

- bug alone
- bug + 5 ants glued
- bug + 5 ants

log(BQ counts) vs log(cm traveled by bug during trial)