Evolution of the recombination regulator PRDM9 in minke whales

Elena Damm*§, Kristian K Ullrich*§, William B Amos†, Linda Odenthal-Hesse*#

Affiliations:

*Department Evolutionary Genetics, Research Group Meiotic Recombination and

Genome Instability, Max Planck Institute for Evolutionary Biology, August-Thienemann

Str. 2, Plön, D-24306, Germany

†Department of Zoology, University of Cambridge, Cambridge, United Kingdom

#corresponding author: email: odenthalhesse@evolbio.mpg.de

§These authors contributed equally

ORCID IDs: 0000-0002-6261-6076 (E.D.), 0000-0003-4308-9626 (K.K.U.), 0000-

0002-5519-2375 (L.O.-H.)

Running title

PRDM9 diversity in minke whales

Keywords

Abstract

We explored the structure and variability of the *Prdm9* gene, which codes for the

PRDM9 protein, in samples of the minke whales from the Atlantic, Pacific and Southern

Oceans. The PRDM9 protein controls the reshuffling of parental genomes in most

metazoans and we show that minke whale possess all the features characteristic of

PRDM9-directed recombination initiation, including complete KRAB, SSXRD and SET

domains and a rapidly evolving array of C2H2-type-Zincfingers (ZnF). We uncovered

eighteen novel PRDM9 variants and evidence of rapid evolution, particularly at DNA-

recognizing positions that evolve under positive selection. At different geographical

scales, we observed extensive *Prdm9* allelic diversity in Antarctic minke whales

(*Balaenoptera bonarensis*) that, conversely, lack observable population differentiation

in mitochondrial DNA and microsatellites. In contrast, a single PRDM9 variant is shared

between all Common Minke whales and even across subspecies boundaries of North

Atlantic (*B. a. acutorostrata*) and North Pacific (*B. a. scammoni*) minke whale, which

do show clear population differentiation. PRDM9 variation of whales predicts distinct

recombination initiation landscapes genome-wide, which has possible consequences

for speciation.

Introduction

The gene *Prdm9* encodes "PR-domain-containing 9" (PRDM9), a meiosis-specific

four-domain protein that regulates meiotic recombination in mammalian genomes. The

four functional domains of the PRDM9 protein are essential for the placement of

double-stranded DNA breaks (DSBs) at sequence-specific target sites. Three of the

domains are highly conserved: *i)* the N-terminal Kruppel-associated box- domain

(KRAB) that promotes protein-protein binding, for example with EWSR, CXXC1, CDYL

and EHMT2 [1, 2]; *ii)* the SSX-repression-domain (SSXRD) of yet unknown function;

iii) the central PR/SET domain, a subclass of the SET domain, with methyltransferase

activity at H3K4me3 and H3K36me3. The fourth, C-terminal, domain comprises an

array of type $Cystin_2Histidin_2$ zinc-fingers (C_2H_2-ZnF), encoded by a minisatellite-like

sequence of 84 bp tandem repeats. This coding minisatellite reveals evidence of

positive selection and concerted evolution, with many functional variants having been

found in humans [3, 4], mice [5, 6], non-human primates [7] and other mammals [8, 9].

Even highly domesticated species like equids [8], bovids [10] and ruminants [11, 12]

show high diversity and rapid evolution. In light of this extreme variability between

minisatellite-like repeat units in most other mammals, the allele found in the genome

of North Pacific minke whale (*Balaenoptera acutorostrata scammoni)* is puzzling, as it

comprises of an array of only a single type of C_2H_2-ZnF type repeat unit.

Minke whales are marine mammals of the genus *Balaenoptera*, in the order of baleen

whales (*Cetaceans*), that are of particular interest not only because little is known

about their population biology, seasonal migration routes and breeding behavior, but

also to support future conservation efforts. Minke whales were long considered to be

a single species, but have been classified as two distinct species, the Common minke

whale (*Balaenoptera acutorostrata)* and the Antarctic minke whale (*Balaenoptera*

bonaerensis Burmeister, 1867). The Common minke whale (*B. acutorostrata)* is

cosmopolitan in the waters of the Northern Hemisphere. And can be separated into

two subspecies, the Atlantic minke whale *(B. acutorostrata acutorostrata)* and the

North Pacific minke whale (*B. acutorostrata scammoni*), separated from each other by

landmasses and the polar ice cap*.* Antarctic minke whal*e*s inhabit the waters of the

Antarctic ocean in the Southern Hemisphere during feeding season but seasonally

migrate to the temperate waters near the Equator during the breeding season [13].

Although overlapping habitats exist near the Equator, seasonal differences in migration

and breeding behavior largely prevent inter-breeding between Common and Antarctic

minke whales [14]. Despite this, occasional migration across the Equator has been

observed [15] and recent studies have uncovered two instances of hybrid individuals,

both females and one with a calf most likely sired by an Antarctic minke whale [16, 17].

Thus, hybrids may be both viable and fertile, allowing backcrossing [17]. However, it

is unclear whether occasional hybridization events have always occurred or whether

they are a recent phenomenon driven by anthropogenic changes, including climate

change [17]. More importantly, since both hybrids were female, current data do not

exclude postzygotic reproductive isolation through males, as expected under

Haldane's rule. Hybrid sterility is a universal phenomenon observed in many eukaryotic

inter-species hybrids, including yeast, plants, insects, birds, and mammals [18, 19].

Minke whales offer an unusual opportunity to study *Prdm9*, the only hybrid sterility

gene that is known in vertebrates thus far, sometimes called a speciation gene [20].

Material and Methods

PRDM9 occurrence and protein domain prediction in diverse taxa

To infer PRDM9 occurrence and to subsequent investigate protein domain architecture

in diverse taxa, the genome sequences were downloaded as given in

Supplementary Table 2. To infer PRDM9 occurrence in the given species, the

annotated protein XP_028019884.1 from *Balaenoptera acutorostrata scammoni* was

used as the query protein with exonerate [21] (v2.2.0) and the --protein2genome model

to extract the best hit. Subsequently, for each species, the extracted coding sequences

(CDS) were translated and investigated with InterProScan [22] (v5.46-81.0) and

HMMER3 [23] (v3.3) using KRAB, SSXRD, SET and C_2H_2-ZnF as bait to obtain the

protein domain architecture.

Phylogenetic Analyses

For the phylogenetic reconstruction across Cetartiodactyla, we used only KRAB,

SSXRD and SET domains (Supplementary Figure 1). These protein domains were

concatenated and used as input for the software BAli-phy [24] (v3.5.0). BAli-phy was

run twice with 10,000 iterations each and the default settings for amino acid input.

Subsequently, the majority consensus tree was obtained by skipping the first 10% of

trees as burn-in, rooted on the branch outside the Cetacea and visualized in Figtree

(http://tree.bio.ed.ac.uk/software/figtree/; v1.4.4).

The phylogenetic tree from the mitochondrial D-loop region was reconstructed,

including several whale species as given in Supplementary Table 3. The

corresponding D-loop was aligned with MAFFT version 5 [25](v7.471) with the L-INS-

i algorithm and manually curated. The maximum-likelihood tree was calculated under

the TN+F+I+G4 model using IQ-TREE [26] (v1.6.12) mid-point rooted and visualised

in Figtree (http://tree.bio.ed.ac.uk/software/figtree/; v1.4.4).

Long-range sequencing of the Prdm9 gene

We first extracted the full-length DNA sequence of the PRDM9 Protein from the minke

whale genome from UCSC genome browser and designed primers to flank the entire

protein-coding portion, encompassing the KRAB, SSXDR, PR/SET and C_2H_2-ZnF

binding domains using Primer-BLAST [27](Supplementary Table 4). We amplified the

Balaenoptera acutorostrata Prdm9 gene, including all introns and exons across a

~11kb interval by long-range PCR. We then sequenced the entire interval using

phased long-range Nanopore Sequencing with MinION (Oxford Nanopore). Whole-

length consensus sequences generated from 639 sequencing reads that cover the

entire 10582 bp, with an average per base pair coverage of 269x. We used this

consensus sequence as reference for *in-silico* predictions of functional domains and

mapped human PRDM9 Exons 3 – 11 from ENSEMBL (ENST00000296682.3). By

manually splicing all unaligned sequence fragments, we generated an in-silico

predicted mRNA of *Balaenoptera acutorostrata* PRDM9. This sequence was then

submitted as '.fasta' to the Entrez Conserved Domains Database (CDD) home page,

(https://www.ncbi.nlm.nih.gov/Structure/cdd/wrpsb.cgi) with the following parameters;

searching against Database CDD v3.18 – 52910 PSSMs Expect Value threshold: 0,01,

without low-complexity filter, composition-based statistics adjustment, rescuing

borderline hits (ON) a maximum number of 500 hits and concise result mode).

Wild minke whale samples

Performed investigations are not considered to be animal trials in accordance with the

German animal welfare act, since samples were obtained from commercial whaling

between 1980 and 1984. A total of n=143 DNA samples of minke whale with vague

information about the subspecies, from four different defined commercial whaling

areas. These include North Atlantic (NA; n=17) and North Pacific (NP; n=4) individuals

captured during migration season, as well as Antarctic Ocean Areas IV (AN IV; n=65)

and V (AN V; n=57) were obtained during feeding season. A subset of these samples

had been used in van Pijlen et al., 1995 [28]. Samples from the Antarctic areas

included numerous duplicate samples that were used as internal controls, analyzed for

all parameters and excluded from the sample set once identical results had been

confirmed (318, 325, 327, 1661, 1663).

For the detailed DNA-extraction protocol, precise catch data, as well as providers of

the samples of most individuals used in this project, see van Pijlen, 1994. Briefly,

genomic DNA was extracted from the skin (NA) and muscle biopsies (NP, AN) with

Phenol-Chlorophorm and stored in TE at - 80°C in the 1980s. If the DNA was dried out

it was first eluted with TE. All DNA concentrations were determined with fluorescent

Nanodrop-1000 (Thermo Fisher Scientific). For all performed analysis, 20 ng/µl DNA

working stocks were prepared and stored at -20°C.

STR genotyping

Nine autosomal, as well as X/Y microsatellite loci with di- or tetramer repeat motifs,

were analysed for all samples: EV001, EV037 [29], GATA028, GATA098, GATA417

[30], GT023, GT310, GT509, GT575 [31] and sex loci X and Y [32]. Four separate

multiplexing reactions were performed for each individual, and each contained 40 ng

of DNA, 0.2 µM of each primer, 5 mM Multiplex-Kit (Qiagen) and HPLC water to a total

volume of 10 µL per sample. Primers (**Supplementary Table 4**) were purchased from

Sigma Aldrich; the reverse Primers were tagged at their 5' end with fluorescent tags

(HEX, FAM or JOE). The first multiplexing reaction contained primers GT023 (HEX),

EV037 (HEX) (FAM) in forward and reverse direction. The multiplexing conditions were

denaturation at 95°C for 15:30 minutes, annealing 1:30 minutes at 59°C, elongation at

72°C for 11:30 minutes and held at 12°C. The second multiplexing reaction contained

primers GT575 (HEX), GATA028 (FAM) in forward and reverse direction, the

multiplexing conditions were denaturation at 95°C for 15:30 minutes, annealing 1:30

minutes at 54°C, elongation at 72°C for 11:30 minutes and a final temperature of 12°C.

The third multiplexing reaction was set up with primers GATA098 (FAM), GT509 (FAM)

and GATA417 (JOE) in forward and reverse direction. The multiplexing conditions

were denaturation at 95°C for 15:30 minutes, annealing 1:30 minutes at 54°C,

elongation at 72°C for 11:30 minutes and a final temperature of 12°C. The fourth

multiplexing reaction contained primers GT310 (HEX) and EV001 (JOE) in forward and

reverse direction. The multiplexing conditions were denaturation at 95°C for 15:30

minutes, annealing 1:30 minutes at 60°C, elongation at 72°C for 11:30 minutes and a

final temperature of 12°C. After amplification, the multiplexing reactions were diluted

with 100 µl water (HPLC grade). Genescan ROX[500] dye size standard (Thermo Fisher

Scientific) and HiDi Formamide (Thermo Fisher Scientific) were mixed 1:100 and used

for sequencing in a mix of 1 µl of diluted multiplexing product and 10 µl HiDi-ROX[500].

STR Multiplex fragment analysis was carried out on the 16-capillary electrophoresis

system ABI 3300 Genetic Analyser (Applied Biosystems).

STR analysis:

We first performed MSA analysis [33] using standard parameters, which calculated

Weinberg expectation (Fis), Shannon Index (Hs), allele numbers (A) and allele sizes.

To detect both weak and strong population structure, we ran simulations with

LOCPRIOR and USEPOPINFO, respectively. For both simulations, the more

conservative "correlated allele frequencies" -model was used, which assumes a level

of non-independence. To ensure that a sufficient number of steps and runs have been

performed we used a burn-in period of 1.000 and runs of 100.000 MCMC repeats for

both types of simulations, each for 50 iterations for successive K values from 1 - 10.

[34]. We used the *a-priori* location to be able to detect even weak population

differentiation. In both datasets, we used the web-based STRUCTURE Harvester

software [35] to determine the rate of change in the log probability between successive

K values, via the *ad-hoc* statistic ΔK from [36]. Figures were rendered using

STRUCTURE PLOT V2.0. [37]

Sequencing of the Mitochondrial Hypervariable Region on the D-loop

The noncoding mtDNA-D-Loop region of 143 individuals was amplified in two

overlapping PCR-reactions. We PCR amplified two different lengths fragments for

each individual: 1066 bp and 331 bp and sequenced the longer product in forward and

the shorter in the reverse direction. The used primers (found in **Supplementary Table**

4) were MT4 (M13F) and MT3 (M13R) for the longer product, and BP15851 (M13F)

and MN312 (M13R) for the shorter PCR product from [38]. The 10 µL reaction was

carried out with 40 ng genomic DNA and 0.2 µM of each primer and 5 mM Multiplex

PCR Kit (Qiagen). Cycling conditions were identical for both directions with 95°C 15:30

min, 53°C 1:30 min, 72°C 13:30 min and hold at 4°C in a Veriti Thermal Cycler (Applied

Biosystems). The PCR products were purified with 3 µl Exo/SAP and then cycle-

sequenced with the BigDye^{TM} Terminator v3.1 Cycle Sequencing Kit (Thermo Fisher

Scientific) according to manufacturer instructions with BP15851 (M13F) for the forward

PCR product and MN312 (M13R) for the reverse PCR product, respectively. The mixes

were then purified with BigDye X-Terminator^{TM} Purification Kit (Thermo Fisher

Scientific) and sequenced by capillary electrophoresis on an ABI 16-capillary 3130xl

Analyzer (Applied Biosystems). The sequences were *de-novo* assembled, and

consensus sequences were generated with Geneious Software 10.2.3 [39].

Prdm9 coding minisatellite array PCR and Sequencing

To characterise the minisatellite-coding for the C_2H_2-ZnF array in more detail, we designed primers nested between the first two conserved ZnFs, and the reverse primer identical to the long-range amplification primer distal to the coding sequence for the C_2H_2-ZnF array. The minisatellite coding for the C_2H_2-ZnF array of PRDM9 of 143 *Balaenoptera* individuals was amplified from 20 ng genomic DNA, in a 20 µl PCR reaction, optimised for the amplification of minisatellites. With 0.5 mM primers that were designed for this study using the *Balaenoptera. acutorostrata scammoni (*XM_007172595) reference and included the PRDM9 minisatellite-like C_2H_2-ZnF array and additional 100bp flanking regions at 5' and 3'. Primers Prdm9ZnFA_Bal_R: and Prdm9ZnFA_Bal_F (**Supplementary Table 4**) and 1x AJ-PCR Buffer described in [40] and 0.025 U/µl Taq-Polymerase and 0.0033 U/µl Pfu-Polymerase. Cycling conditions were: initial denaturation at 95°C for 1:30 min followed by 33 cycles including 96°C 15 s, 61°C 20 s and 70 °C 2:00 min and finally 70°C 5 min, and hold at 4°C in a Veriti Thermal Cycler (Applied Biosystems). Agarose gel electrophoresis (1.5% Top Vision Low Melting Point Agarose gel (Thermo Fisher Scientific) with SYBR Safe (Thermo Fisher Scientific) was used to visualise allele sizes as well as zygosity. All bands were excised from the gel (Molecular Imager® Gel Doc™ XR System with Xcita-Blue™ Conversion Screen (Biorad), and PCR products were recovered from the gel fragments with 2 U/100 mg Agarase. If the individuals were homozygous, the extracted DNA was directly Sanger sequenced from in 5' and 3' directions for each sample with the BigDye™ Terminator v3.1 Cycle Sequencing Kit (Thermo Fisher Scientific) according to the manufacturer's protocol with the same primers as for the amplification (Prdm9ZnFA_Bal_R/Prdm9ZnFA_Bal_F). The sequencing-reaction was carried out in

the ABI 16-capillary 3130xl Analyzer (Applied Biosystems). Heterozygous samples

were subcloned before sequencing.

Subcloning of heterozygous alleles of identical lengths

Subcloning was performed for a subset of samples when Sanger sequencing revealed

heterozygous alleles of the same length, that could not be distinguished by

electrophoresis, but revealed heterozygous nucleotides in the chromatogram. Instead

of inferring haplotypes, we wanted to experimentally ascertain both alleles. Thus, we

cloned the remaining PCR product into the TOPO TA (Invitrogen) vector and

transferred the vectors into OneShotTop10 chemically competent cells (Invitrogen). All

steps were carried out according to the manufacturer's manuals. Eight positive clones

were picked for each sample, and the DNA was extracted by boiling in HPLC-grade

water at 96°C for 10 Minutes. A centrifugation step then removed the cell debris, and

the supernatant was directly used for PCR, gel-purification and Sanger-sequencing as

described in the section above.

Sequence Analysis of the PRDM9 C_2H_2-ZnF array

Sequences determined different alleles and numbers of repeat units per array, which

were *de novo* annotated and the resulting consensus sequences were translated into

the corresponding protein variants of each C_2H_2-ZnF by Geneious Software 10.2.3

[39]. All PCR products contained the complete minisatellite-like coding sequence for

the C_2H_2 domain and additional 100 base pairs flanking regions at the 5' and 3' ends.

PRDM9 C_2H_2-ZnF array coding sequence dN/dS analysis

The internal C_2H_2-ZnFs were obtained as described for the phylogenetic

reconstruction. Subsequently, all repeat units were stacked using only non-unique

alleles. The resulting codon alignment was used to determine basic population

parameters for each population (population theta using segregating sites and

nucleotide diversity) by either including or excluding hypervariable amino acid positions

(-1, +3, +6) with the R package "pegas" [41]. Episodic diversifying selection among

Zinc-finger sites using was determined by a mixed-effects model of evolution (MEME)

at https://www.datamonkey.org as described earlier [42].

Phylogenetic analyses of the PRDM9 hypervariable domain

The gene tree of PRDM9 was inferred either using only the repeats coding for the

internal C_2H_2-ZnFs or the concatenated KRAB, SSXRD and SET domain. For the

internal C_2H_2-ZnFs based phylogenetic reconstruction (**Figure 1**), internal C_2H_2-ZnFs

were extracted, and pairwise distances were calculated as in Vara et al. 2019 [43] with

the developmental R package repeatR (https://gitlab.gwdg.de/mpievolbio-it/repeatr). In

brief, only species were used with one leading C_2H_2-ZnF and at least eight internal

C_2H_2-ZnFs (see supplementary Figure 1). For each possible repeat combination (r, r')

the hamming distances of the corresponding repeat units $r = (r_1; r_2; r_3; …)$ and $r' = (r'_1;$

$r'_2; r'_3; …)$ were used to derive the edit distance between r and r'. Before calculating

the edit distance, the codons coding for the hypervariable amino acid positions (-1, +3,

+6) were removed for each repeat unit and weighting cost of $w_{mut} = 1$, $w_{indel} = 3.5$ and

$w_{slippage} = 1.75$ as given in [43]. The resulting pairwise edit distance matrix was used to

calculate a neighbour-joining tree with the *bionj* function of the R package ape [44],

rooted on the branch leading to the bottlenose dolphin (*Tursiops truncatus)* and

visualized in Figtree (http://tree.bio.ed.ac.uk/software/figtree/ v1.4.4).

Prediction of DNA-binding motifs of different PRDM9 variants

DNA-binding Specificities of the different Cys_2His_2 Zinc Finger Protein variants were

predict *in-silico* using the SVM polynomial kernel method within "Princeton ZnF"

(http://zf.princeton.edu/) [45].

Results

The evolutionary context of PRDM9 in *Cetartiodactyla*

For a broad view on the evolution of PRDM9 in *Cetartiodactyla*, we identified PRDM9

orthologs from all publicly available genomic resources (

Supplementary Table 2). To first establish an evolutionary context based on the

protein-coding regions for the proximal conserved domains, we used the *Balaenoptera*

acutorostrata scammoni protein as a query and extracted coding sequences of the N-

terminal (conserved) domains. Previous studies of PRDM9 in vertebrates found rapid

evolution of DNA binding sites only when the KRAB and SSXRD domains were present

[46]. Consequently, we used InterProScan to create a curated dataset of

Cetartiodactyla PRDM9 orthologues which contained the KRAB, SSXRD, and SET

domains, including a PRDM9 orthologue in Antarctic minke whale. Our initial analyses

revealed that minke whales, and *Cetartiodactyla* in general, possessed all the features

characteristic of organisms with PRDM9-directed recombination initiation, including

complete KRAB, SSXRD and SET domains. However, organisms with PRDM9

directed recombination, also display a rapidly-evolving PRDM9 C_2H_2-ZnF array [46]

With the sole exception of Hippopotamus, all *Cetartiodactyla* also contained the C_2H_2-

ZnF array, with large variation in the number of C_2H_2-ZnF between species.

Characterizing the *Prdm9* gene in minke whales

To characterize the sequence and structure of the *Prdm9* gene, we began by amplifying and sequencing it from a pooled sample, containing DNA from six individuals, five Antarctic minke whales and one common minke whale. The consensus sequence was subjected to *in-silico* prediction that successfully recovered all relevant PRDM9 protein domains with high-confidence: the KRAB domain (E-value: $1.84e^{-11}$); the SSXRD motif (E-value: $4.46e^{-10}$); the PR/SET domain (E-value: $9.69e^{-05}$); and several Zinc-Fingers (**Figure 1**). The complete C_2H_2-domain comprises an array of C_2H_2-ZnFs (the C_2H_2-ZnF-array), as well as a single C_2H_2-ZnF (ZnF#1) that is located proximally (E-value: $6.52e^{-04}$). This PRDM9-ZnF#1 is identical to that of mice [47] as well as rats, elephants, human, chimpanzee, macaque and orangutan [48], and thus appears to be conserved across large evolutionary timescales. The second C_2H_2-ZnF (E-value: $1.65e^{-11}$) appears conserved across *Balaenoptera*. The same second C_2H_2-ZnF is seen in our pooled sample, as well as the genome reference of the North Pacific minke whale (*Balaenoptera acutorostrata scammoni)* and Antarctic minke whale (*Balaenoptera bonarensis)* as well as the blue whale (*Balaenoptera musculus).* However, from the third ZnF onwards, sequencing across pooled samples exhibited high variability, which we could not resolve.

Variation of the PRDM9 coding minisatellite in minke whales

We assessed the variability of the C_2H_2-ZnF-array by analyzing the coding minisatellite in a large number of minke whales including Antarctic minke whale (*Balaenoptera bonaerensis* Burmeister, 1867) and two Common minke whale subspecies, Atlantic minke whale and North Pacific minke whale. Amplification and subsequent electrophoresis on agarose gels resolved six different allele sizes of the last exon of

the *Prdm9* gene. A single size was observed in all Common minke whales from

sampled populations in the North Atlantic as well as North Pacific. In contrast, five

different sizes were identified across Antarctic minke whale populations sampled in the

Southern Hemisphere. All common minke whales have a size consistent with eleven

84 bp repeats while the most prevalent genotype in the Antarctic minke whales

corresponds to nine repeats, with variation between six and ten repeats. Therefore,

Prdm9 shows size variation length of the coding minisatellite, that result from variation

in the number of repeat units.

Sequencing of the DNA purified from gel fragments, as well as cloning of apparently

homozygous bands, revealed a remarkable diversity of minisatellite alleles. Over 80%

of animals possessed repeat-arrays of nine repeats; however, these comprised six

different alleles. Because in addition to variation in repeat-number, there is nucleotide

variation between 84bp-minisatellite repeats. And finally, not only which repeat-types

are present, but also their order varies between individuals.

Diversity and diversifying selection on C_2H_2-zinc fingers of PRDM9

Each 84bp repeat unit translates into 28 amino-acids coding for a single C_2H_2-ZnF. To

explore diversity between the ZnF, we extracted all 84bp repeat units, which we used

to predict C_2H_2-ZnFs through an HMMER algorithm [45]. This yielded 26 different

versions with HMMER bit scores >17 (**Figure 1** and **Supplementary Figure 4**). Based

on amino acid variation within each predicted C_2H_2-ZnFs, we identified 14 "common"

variants found in multiple individuals as well as twelve "rare" variants found in only a

single individual (**Supplementary Figure 4**). The most variable amino acids are 13,

16 and 19, which correspond to positions -1, +3 and +6 of the alpha-helices that are

responsible for DNA binding specificity. However, five of the fourteen "common"

variants have the same amino sequence LNG at these positions, differing instead at

three positions in beta-helices flanking the cysteine residues that bind the zinc-ion

(positions 3, 10 and 12). Three of the five LNG variants are also found in the reference

sequences of both the Antarctic minke whale and the North Atlantic minke whale. We

next used the fourteen common C_2H_2-ZnFs to test for signals of selection, finding

Evidence that amino acid 16, is under episodic diversifying selection (**Supplementary**

Figure 5). This amino acid is located at position +3 of the alpha helix, one of the three

positions responsible for DNA-binding specificity.

Evolutionary turnover of PRDM9 in minke whales

Using all minisatellite-coding sequences for the array, we determined C_2H_2-ZNF

domain variants that each individual possessed. Minisatellite size homoplasy equates

to an identical PRDM9 C_2H_2-ZnF array in Common minke whales, but not Antarctic

minke whales. The only C_2H_2 domain variant found in all samples from the Northern

Hemisphere NH12_A consists of eleven C_2H_2-ZnF in the array and the C_2H_2-ZnF, that

is located proximally and appears conserved across *Balaenoptera*. In contrast,

samples from the Southern Hemisphere displayed seventeen different C_2H_2-Domains

of PRDM9 (**Figure 1**).

We want to understand the evolutionary relationships of the different alleles that we

observed in minke whale species. However, phylogenetic analysis of minisatellite

repeat-structures is challenging due to their rapid rate of evolution. Minisatellites,

including that of *Prdm9*, evolve mainly by unequal crossing-over and gene conversion

in meiosis [49]. Stepwise mutation models that are commonly assumed are based on

microsatellites, and thus give only a poor fit to minisatellite evolution [50]. So, instead

of comparing full-length-*Prdm9* allelic variants, we separated each array into 84bp

repeat types. We then calculated pairwise Hamming (i.e. minimum edit) distances

between all repeats, after removing the nucleotides coding for the hypervariable amino

acid positions (-1, +3, +6), and applying specific weighting costs as given in [43]. In

addition to the repeats present in our minke whale samples, we also included repeats

coding for the internal C_2H_2-ZnFs of all *Cetartiodactyla* species into our phylogenetic

analyses (see **Supplementary Figure 1**) Here, we included only those species that

possessed the complete N-terminal domain architecture, the proximal C_2H_2-ZnF and

at least eight successive C_2H_2-ZnF in the C-terminal DNA-binding domain. We

restricted our phylogenetic analyses on the internal minisatellite repeats since our

analyses revealed that the DNA-binding positions of the most proximal ZnF in the array

(ZnF#2) are not only conserved in minke whales but also generally within *Cetacea*

(**Supplementary Figure 2**). In fact, we observed only a single DNA-binding amino-

acid change between C_2H_2-ZnF #2 of all *Cetacea* compared to all *Ruminantia*

(**Supplementary Figure 3**).

The PRDM9 phylogenetic analyses in **Figure 2** show that all species cluster into

distinct phylogenetic groups. Reference alleles cluster with their own subspecies, even

including the unusual reference allele of *Balaenoptera acutorostrata scammoni*, which

nevertheless clusters with our common minke whale samples. Similarly, the genome

reference allele of *Balaenoptera bonarensis* (Antarctic minke whale) clusters within the

group of SH_8A and SH-8B alleles found in our Antarctic minke whale samples, which

appear to be more ancestral than other minke whale alleles. The phylogenetic

reconstruction also suggests that common minke whale alleles have appeared more

recently and evolved mainly by an increase in repeat-copy after splitting from the

Antarctic minke whale. Our phylogenetic analysis thus fits well with an evolutionary

history where common minke whales split from Antarctic minke whales approximately

5 million years ago.

Diversity of PRDM9 at different geographical scales

To explore the allelic diversity of *Prdm9* at different geographical scales, we partitioned

the data by sampling location wherever accurate catch-locations were available

(**Figure 2**). Allelic diversity varied between sampling locations, even within the

Antarctic sampling sites. The most common alleles are SH_10A, with SH_10B, and

these occur at frequencies of 10-20% in all sampling locations. However, all other

alleles differ between different sampling populations, each of which carries at least one

allele found exclusively in a single population. Sample size does not predict allele

number, with the highest allele number (ten) being associated with an average-sized

sample (n=34).

Population structure of minke whales

We are interested in the possible evolutionary consequences of PRDM9 for minke

whale speciation in the face of what may be recent secondary mixing, and this requires

a context of current levels of population isolation. We thus explored variability at the

84bp *Prdm9* minisatellite repeat across the four sample regions for population genetic

analyses (**Table 1**). Taking each repeat type as an individual allele, we computed the

mutation rate per base pair per generation (population θ), as well as the average

pairwise nucleotide diversity. The latter was calculated in two ways: (i) using the entire

minisatellite-like repeat sequence; (ii) after removing nucleotides coding for the

hypervariable sites that translate into the DNA binding positions of individual C_2H_2-

ZnF, which are known to be under positive selection. We observed the highest

population θ in Antarctic Area IV, compared to all other sampling locations, as seen in

Table 1. Across the microsatellite-repeat types, the Average Pairwise Nucleotide

Diversity (APND, including nucleotides coding for the hypervariable sites) is similar

across all sampled populations and between common minke whale and Antarctic

minke whales. After excluding the nucleotides coding for hypervariable sites (-1, +3,

+6), the APND is decreased roughly 2.5-fold, across minke whales.

We used two approaches to quantify genetic differentiation of PRDM9 between the

different minke whale populations: the per-site distance for multiple alleles [51] and

Jost's D (D_J), the fraction of allelic variation among populations [52]. Both measures

indicate weak population differentiation, the highest differentiation being between the

North Atlantic and North Pacific ($G_{ST} = -0.0039$, $D_J = 0.0077$). The second highest

population differentiation was seen between the North Pacific and both Antarctic

regions, $G_{ST} = -0.0035$, $D_J = 0.0070$, with slightly lower differentiation between Atlantic

and Antarctic, $G_{ST} = -0.0011$, $D_J = 0.0022$. The lowest population differentiation was

between Antarctic Areas IV and V (ANV vs ANIV, $G_{ST} = -0.0004$, $D_J = 0.0009$).

To explore population structure differentiation further, we also analysed polymorphic

microsatellites and the hypervariable region of the mitochondrial D-Loop (mtDNA HVR)

(**Figure 3**). Despite the low *Prdm9* diversity in the Northern Hemisphere, genetic

structuring was revealed by the mtDNA HVR, with a clear distinction between the North

Atlantic and North Pacific whales. We also examined nine unlinked autosomal

microsatellite loci, chosen for their high information content in minke whales [17, 29]

and sex-chromosome markers ZFX/ZFY, from [38]. Microsatellite summary statistics

reveal high heterozygosity of effectively zero F_{IS} (Supplementary Table 1), so we

continued with a Bayesian analysis of population structure. We used *STRUCTURE*

with a recessive allele model because any null alleles were due to polymorphisms,

rather than missing data [53] and the "admixture" model. However, given that minke

whales occur in geographically isolated groups, are separated by hemispheres, and

have asynchronous breeding seasons, we nevertheless assumed a low probability for

incomplete lineage sorting, which could otherwise confound structure inference,

particularly for weak population differentiation. The most likely number of clusters

returned by ΔK is *k* = 2, which distinguishes the two hemispheres. Increasing to *k = 3*

and *k = 4*, further separates the common minke whale subspecies (as shown in **Figure**

3), a result that holds even without *a priori* location data (NOLOCS) where only strong

population structure should be detected (**Supplementary Figure 6**).

Putative minke whale Recombination initiation motifs

Different C_2H_2-ZnF domains target different recombination hotspots [3, 4, 54, 55].

Target hotspots in humans [4, 48] and mice [55] can be predicted from C_2H_2 ZnF

sequences using the SVM polynomial kernel model for *de-novo* binding prediction [45].

To understand the situation in minke whales, we identified a number of "core motifs",

based on ZnFs #3-#6, that appear particularly important for DNA binding [56, 57]

(**Figure 4**). As must be true, the single PRDM9 variant found in common minke whales

results in a single "core motif". However, a large majority of Antarctic minke whales

also share a single "core motif", despite a much larger diversity of PRDM9 (**Figure 4**).

Core motifs show no overlap between common minke whales and Antarctic minke

whales, a trend that is particularly evident at C_2H_2-ZnF#3 where it creates distinctive

DNA binding motifs. Thus, it appears as if core motifs are shared between the majority

of minke whale PRDM9 variants within a given species, but not between species

inhabiting different Hemispheres.

Discussion:

PRDM9 shows high diversity, especially in Antarctic minke whales

We characterized the PRDM9 C_2H_2-ZnF domain diversity of 134 individuals from four

natural populations of minke whales and discovered a total of eighteen variants. The

minisatellite encoding the DNA-binding C_2H_2-ZnF array apparently evolves rapidly in

minke whales, with at least one of the DNA-recognizing position likely under positive

selection. This high diversity is similar to that observed at the *Prdm9* gene in other

vertebrate species, such as humans, mice, bovines and primates [4-6, 12, 58].

However, PRDM9 diversity is not equally abundant in both hemispheres. In contrast to

the extensive PRDM9 diversity we observe in Antarctic minke whales (*Balaenoptera*

bonarensis), only a single PRDM9 variant is found in Common minke whales. This

variant is shared between two subspecies, the North Atlantic minke whale *(B. a.*

acutorostrata) and North Pacific minke whales *(B. a. scammoni).* In most mammals,

extensive PRDM9 variation typically is seen between, and within, a given subspecies.

This holds true also for Antarctic minke whales, thus the observation of an identical

PRDM9 variant shared between two subspecies of common minke whales is

unexpected. The PRDM9 protein level diversity observed within each Hemisphere

contrasted with the levels of population differentiation seen in microsatellite as well as

mitochondrial haplotypes. However, diversity between repeat types of the *Prdm9*

minisatellite can reveal even weak population structure and is consistent with data on

mitochondrial haplotypes and microsatellites. This contrasting pattern between

protein-level conservation and nucleotide differentiation is fascinating and points to

functional constraints operating on different levels of PRDM9 evolution.

Population genetic analyses of minke whales and potential speciation

Taxonomists have separated the common minke whale into two subspecies, the North

Atlantic minke whale *(B. a. acutorostrata)* and the North Pacific minke whale (*B. a.*

scammoni), which diverged approximately 1.5 million years ago. variability between

84bp *Prdm9* minisatellite repeat units, as well as microsatellite markers and mtDNA all

reveal weak population structure and differentiation between minke whales in North

Atlantic and North Pacific. Our population genetic data support this classification and

suggest that these populations may be in the process of speciating. Yet fascinatingly,

all individuals of both Northern Hemisphere populations had identical C_2H_2 domains,

which suggests that these subspecies should still be able to interbreed if given a

chance. As geographical isolation of the two subspecies is currently still upheld by the

permanent polar ice, allopatric speciation may be promoted. However, due to global

warming, the two subspecies might come into secondary contact again in the future.

The removal of the current geographical barrier could therefore disrupt speciation,

especially in the light of identical C_2H_2 domains encoded by the mammalian hybrid

sterility gene *Prdm9.*

Hybrids between Antarctic minke whale and Common minke whale

Analysis of the hypervariable region (HVR) on the mitochondrial D-Loop suggests that

Antarctic minke whale and Common minke whale evolved from a common ancestor in

the Southern Hemisphere during a period of global warming approximately 5 million

years ago [59]. Common minke whales and Antarctic minke whales are now separated

by both geography and seasonality. However, while their winter habitats and breeding

grounds remain unknown, it is assumed that *B. a. acutorostrata* migrates south

between November and March to give birth in warmer waters and occasional

individuals are seen as far south as the Gulf of Mexico [60]. Interbreeding between the

populations of the Northern and Southern Hemisphere appears unlikely at the time our

samples were collected (the 1980s) because our study confirms previous findings [28]

that common minke whales and Antarctic minke whales exhibit extensive genetic

differences in their mtDNA and microsatellites. However, recent reports have

uncovered rare hybridization events between Common minke whale and Antarctic

minke whales [16], which shows that interbreeding is unlikely but possible. The

question remains whether known reproductive isolation mechanism exists in hybrids

between the two minke whale species.

The *Prdm9* gene remains the only hybrid sterility gene known in vertebrates. Large

evolutionary turnover of the coding minisatellite is seen in all mammals characterised

to date- including minke whales (*this study*). One consequence is genetic variation in

the DNA-binding C_2H_2-ZnF-domain that in turn introduces variation in species'

recombination landscapes, that differ even within populations of the same species [4,

61]. Based on computed PRMD9 C_2H_2-ZnF binding motifs found in Common minke

whale and Antarctic minke whale, we would speculate that the recombination initiation

landscapes of the two variants would not overlap. The associated variation in

recombination landscapes is implicated in generating reproductive isolation in

mammals, which has been well characterized in mice. Here, historical erosion of

genomic binding sites of PRDM9 ZnF domains has been shown to lead to the

introduction of asymmetric sets of DSBs in evolutionary divergent homologous genomic

sequences [61, 62]. This asymmetry is likely caused because PRDM9 initiation can

erode its own binding via biased gene conversion over long evolutionary timescales.

As a result, in hybrid genomes, the variant of one species preferentially binds the

ancestral, binding sites on the homologue of the other species, that have not been

eroded, and vice versa. The resulting asymmetry of recombination initiation sites is

believed to be responsible for the inefficient DSB repair, defective pairing, asynapsis

and segregation of the chromosomes that are observed in intersubspecific mouse

hybrids [63-65]. The low PRDM9 diversity in the Northern Hemisphere would predict

extensive historical erosion of PRDM9 binding sites in the genomes of Common Minke

Whales. A degree of erosion may also be true across the Southern Hemisphere, where

most animals share identical core motifs. Thus, the predicted high level of PRDM9

binding site erosion, coupled with non-overlapping minke whale PRDM9 binding motifs

would predict asymmetric PRDM9 recombination imitation in minke whale hybrids

(**Supplementary Figure 7**), which is implicated in hybrid male sterility.

Despite this potential incompatibility, two viable female minke whale hybrids have been

observed, one of which showed signs of successful backcrossing [15, 16]. This

observation alone is not sufficient to clarify whether postzygotic isolation mechanisms

do or do not generally operate. After all, Haldane's rule predicts that the sterile sex will

be the heterogametic males, rather than the females on which observations were

made, an expectation that holds in PRDM9-mediated hybrid sterility in mice [20, 66,

67]. Therefore, it still remains to be understood whether sporadically occurring

hybridization events will generate infertile males or, instead, signify that start of a

breakdown of genetic isolation. Further studies on minke whale hybrids are necessary

to elucidate this matter.

Conclusion

We uncovered that Minke whales, and related species, possess coding sequences for

a complete set of KRAB, SSXRD and SET domains, a necessary feature of organisms

with PRDM9-regulated recombination [46]. Sequencing revealed evolutionary turnover

of the PRDM9 coding minisatellite. Our phylogenetic analysis revealed a good

association of groups with their geographical origin, a high level of size homoplasy in

common minke whales and a lower level of size homoplasy in Antarctic minke whales.

In Antarctic minke whales (*Balaenoptera bonarensis*), we observe extensive diversity

within the C_2H_2-ZNF domain of PRDM9, even at fine geographical scale. In direct

opposition, we observe no genetic diversity within the C_2H_2-ZNF in common minke

whales. In each Hemisphere, the PRDM9 protein diversity is contrasting strongly with

the underlying population structure we observe between and within subspecies.

However, on the nucleotide level, variation between *Prdm9* minisatellite repeat types

mirrored the variation in mitochondrial DNA and microsatellites in respective animals.

This apparent contradiction points to maintenance of conserved protein sequence,

even across minke whale subspecies boundaries.

Acknowledgements

We thank the Max Planck Society for funding this study.

We thank Nicole Thomsen and Olga Eitel for technical support. We would also like to

thank Kevin Glover and Bjørghild B. Seliussen for sharing primer sequences and

genotyping conditions for microsatellites markers and the hypervariable segments of

the mtDNA D-loop.

References:
1.   Imai Y, Baudat F, Taillepierre M, Stanzione M, Toth A, de Massy B: **The PRDM9**
KRAB domain is required for meiosis and involved in protein interactions.
Chromosoma 2017, **126**(6):681-695.
2.   Parvanov ED, Tian H, Billings T, Saxl RL, Spruce C, Aithal R, Krejci L, Paigen
K, Petkov PM: **PRDM9 interactions with other proteins provide a link**
between recombination hotspots and the chromosomal axis in meiosis.
Mol Biol Cell 2017, **28**(3):488-499.
3.   Berg IL, Neumann R, Lam KW, Sarbajna S, Odenthal-Hesse L, May CA,
Jeffreys AJ: **PRDM9 variation strongly influences recombination hot-spot**
activity and meiotic instability in humans. *Nature genetics* 2010, **42**(10):859-
863.
4.   Berg IL, Neumann R, Sarbajna S, Odenthal-Hesse L, Butler NJ, Jeffreys AJ:
Variants of the protein PRDM9 differentially regulate a set of human
meiotic recombination hotspots highly active in African populations.
Proceedings of the National Academy of Sciences of the United States of
America 2011, **108**(30):12378-12383.
5.   Kono H, Tamura M, Osada N, Suzuki H, Abe K, Moriwaki K, Ohta K, Shiroishi
T: **Prdm9 polymorphism unveils mouse evolutionary tracks**. *DNA research*
: an international journal for rapid publication of reports on genes and genomes
2014, **21**(3):315-326.
6.   Buard J, Rivals E, Dunoyer de Segonzac D, Garres C, Caminade P, de Massy
B, Boursot P: **Diversity of Prdm9 zinc finger array in wild mice unravels**
new facets of the evolutionary turnover of this coding minisatellite. *PloS*
one 2014, **9**(1):e85021.
7.   Schwartz JJ, Roach DJ, Thomas JH, Shendure J: **Primate evolution of the**
recombination regulator PRDM9. *Nat Commun* 2014, **5**:4370.
8.   Steiner CC, Ryder OA: **Characterization of Prdm9 in equids and sterility in**
mules. *PloS one* 2013, **8**(4):e61746.
9.   Ahlawat S, Sharma P, Sharma R, Arora R, Verma NK: **Evidence of positive**
selection and concerted evolution in the rapidly evolving PRDM9 zinc
finger domain in goats and sheep. 2016, **47**(6):740-751.
10.   Ahlawat S, De S, Sharma P, Sharma R, Arora R, Kataria RS, Datta TK, Singh
RK: **Evolutionary dynamics of meiotic recombination hotspots regulator**
PRDM9 in bovids. *Mol Genet Genomics* 2017, **292**(1):117-131.
11.   Ahlawat S, Sharma P, Sharma R, Arora R, De S: **Zinc Finger Domain of the**
PRDM9 Gene on Chromosome 1 Exhibits High Diversity in Ruminants but
Its Paralog PRDM7 Contains Multiple Disruptive Mutations. *PloS one* 2016,
11(5):e0156159.
12.   Ahlawat S, Sharma P, Sharma R, Arora R, Verma NK, Brahma B, Mishra P, De
S: **Evidence of positive selection and concerted evolution in the rapidly**
evolving PRDM9 zinc finger domain in goats and sheep. *Anim Genet* 2016,
47(6):740-751.

13. Bakke I, Johansen S, Bakke Ø, El-Gewely MR: **Lack of population**
subdivision among the minke whales (Balaenoptera acutorostrata) from
Icelandic and Norwegian waters based on mitochondrial DNA sequences.
Marine Biology 1996, **125**(1):1-9.
14. Perrin WF, Jr. RLB: **Minke Whales: Balaenoptera acutorostrata and B.**
bonaerensis. In: *Encyclopedia of Marine Mammals (Second Edition).* Edited by
Perrin WF, Würsig B, Thewissen JGM, Second edn. Academic Press; 2009:
733-735.
15. Glover KA, Kanda N, Haug T, Pastene LA, Oien N, Goto M, Seliussen BB,
Skaug HJ: **Migration of Antarctic minke whales to the Arctic**. *PloS one* 2010,
5(12):e15197.
16. Glover KA, Kanda N, Haug T, Pastene LA, Oien N, Seliussen BB, Sorvik AG,
Skaug HJ: **Hybrids between common and Antarctic minke whales are**
fertile and can back-cross. *BMC Genet* 2013, **14**:25.
17. Malde K, Seliussen BB, Quintela M, Dahle G, Besnier F, Skaug HJ, Oien N,
Solvang HK, Haug T, Skern-Mauritzen R *et al*: **Whole genome resequencing**
reveals diagnostic markers for investigating global migration and
hybridization between minke whale species. *BMC genomics* 2017, **18**(1):76.
18. Maheshwari S, Barbash DA: **The genetics of hybrid incompatibilities**. *Annual*
review of genetics 2011, **45**:331-355.
19. Coyne JA, Orr HA: **Speciation**: Sinauer Associates, Inc.; 2004.
20. Mihola O, Trachtulec Z, Vlcek C, Schimenti JC, Forejt J: **A mouse speciation**
gene encodes a meiotic histone H3 methyltransferase. *Science* 2009,
323(5912):373-375.
21. Slater GS, Birney E: **Automated generation of heuristics for biological**
sequence comparison. *BMC Bioinformatics* 2005, **6**:31.
22. Quevillon E, Silventoinen V, Pillai S, Harte N, Mulder N, Apweiler R, Lopez R:
InterProScan: protein domains identifier. *Nucleic Acids Res* 2005, **33**(Web
Server issue):W116-120.
23. Mistry J, Finn RD, Eddy SR, Bateman A, Punta M: **Challenges in homology**
search: HMMER3 and convergent evolution of coiled-coil regions. *Nucleic*
Acids Res 2013, **41**(12):e121.
24. Suchard MA, Redelings BD: **BAli-Phy: simultaneous Bayesian inference of**
alignment and phylogeny. *Bioinformatics* 2006, **22**(16):2047-2048.
25. Katoh K, Kuma K, Toh H, Miyata T: **MAFFT version 5: improvement in**
accuracy of multiple sequence alignment. *Nucleic Acids Res* 2005,
33(2):511-518.
26. Nguyen LT, Schmidt HA, von Haeseler A, Minh BQ: **IQ-TREE: a fast and**
effective stochastic algorithm for estimating maximum-likelihood
phylogenies. *Molecular biology and evolution* 2015, **32**(1):268-274.
27. Ye J, Coulouris G, Zaretskaya I, Cutcutache I, Rozen S, Madden TL: **Primer-**
BLAST: A tool to design target-specific primers for polymerase chain
reaction. *BMC Bioinformatics* 2012, **13:134**.
28. van Pijlen IA, Amos B, Burke T: **Patterns of genetic variability at individual**
minisatellite loci in minke whale Balaenoptera acutorostrata populations
from three different oceans. *Molecular biology and evolution* 1995, **12**(3):459-
472.
29. Valsecchi E, Amos W: **Microsatellite markers for the study of cetacean**
populations. *Mol Ecol* 1996, **5**(1):151-156.

30.   Palsboll PJ, Berube M, Larsen AH, Jorgensen H: **Primers for the amplification
of tri- and tetramer microsatellite loci in baleen whales**. *Mol Ecol* 1997,
6(9):893-895.
31.   Berube M, Jorgensen H, McEwing R, Palsboll PJ: **Polymorphic di-nucleotide
microsatellite loci isolated from the humpback whale, Megaptera
novaeangliae**. *Mol Ecol* 2000, **9**(12):2181-2183.
32.   Berube M, Palsboll P: **Identification of sex in cetaceans by multiplexing with
three ZFX and ZFY specific primers**. *Mol Ecol* 1996, **5**(2):283-287.
33.   Mukaj A, Pialek J, Fotopulosova V, Morgan AP, Odenthal-Hesse L, Parvanov
ED, Forejt J: **Prdm9 inter-subspecific interactions in hybrid male sterility
of house mouse**. *Molecular biology and evolution* 2020.
34.   Orozco-terWengel P, Corander J, Schlotterer C: **Genealogical lineage sorting
leads to significant, but incorrect Bayesian multilocus inference of
population structure**. *Mol Ecol* 2011, **20**(6):1108-1121.
35.   Earl DA, vonHoldt BM: **STRUCTURE HARVESTER: a website and program
for visualizing STRUCTURE output and implementing the Evanno method**.
Conservation Genetics Resources 2011, **4**(2):359-361.
36.   Evanno G, Regnaut S, Goudet J: **Detecting the number of clusters of
individuals using the software STRUCTURE: a simulation study**. *Mol Ecol*
2005, **14**(8):2611-2620.
37.   Ramasamy RK, Ramasamy S, Bindroo BB, Naik VG: **STRUCTURE PLOT: a
program for drawing elegant STRUCTURE bar plots in user friendly
interface**. *Springerplus* 2014, **3**:431.
38.   Glover KA, Haug T, Øien N, Walløe L, Lindblom L, Seliussen BB, Skaug HJ:
**The Norwegian minke whale DNA register: a data base monitoring
commercial harvest and trade of whale products**. *Fish and Fisheries* 2012,
13(3):313-332.
39.   Kearse M, Moir R, Wilson A, Stones-Havas S, Cheung M, Sturrock S, Buxton
S, Cooper A, Markowitz S, Duran C *et al*: **Geneious Basic: an integrated and
extendable desktop software platform for the organization and analysis of
sequence data**. *Bioinformatics* 2012, **28**(12):1647-1649.
40.   Jeffreys AJ, Neumann R, Wilson V: **Repeat unit sequence variation in
minisatellites: a novel source of DNA polymorphism for studying variation
and mutation by single molecule analysis**. *Cell* 1990, **60**(3):473-485.
41.   Paradis E: **pegas: an R package for population genetics with an integrated-
modular approach**. *Bioinformatics* 2010, **26**(3):419-420.
42.   Murrell B, Wertheim JO, Moola S, Weighill T, Scheffler K, Kosakovsky Pond SL:
Detecting individual sites subject to episodic diversifying selection. *PLoS
genetics* 2012, **8**(7):e1002764.
43.   Vara C, Capilla L, Ferretti L, Ledda A, Sanchez-Guillen RA, Gabriel SI, Albert-
Lizandra G, Florit-Sabater B, Bello-Rodriguez J, Ventura J *et al*: **PRDM9
diversity at fine geographical scale reveals contrasting evolutionary
patterns and functional constraints in natural populations of house mice**.
Molecular biology and evolution 2019.
44.   Paradis E, Claude J, Strimmer K: **APE: Analyses of Phylogenetics and
Evolution in R language**. *Bioinformatics* 2004, **20**(2):289-290.
45.   Persikov AV, Singh M: **De novo prediction of DNA-binding specificities for
Cys2His2 zinc finger proteins**. *Nucleic Acids Res* 2014, **42**(1):97-108.

46. Baker Z, Schumer M, Haba Y, Bashkirova L, Holland C, Rosenthal GG, Przeworski M: **Repeated losses of PRDM9-directed recombination despite the conservation of PRDM9 across vertebrates**. *Elife* 2017, **6**.

47. Paigen K, Petkov PM: **PRDM9 and Its Role in Genetic Recombination**. *Trends Genet* 2018, **34**(4):291-300.

48. Myers S, Bowden R, Tumian A, Bontrop RE, Freeman C, MacFie TS, McVean G, Donnelly P: **Drive against hotspot motifs in primates implicates the PRDM9 gene in meiotic recombination**. *Science* 2010, **327**(5967):876-879.

49. Jeffreys AJ, Cotton VE, Neumann R, Lam KW: **Recombination regulator PRDM9 influences the instability of its own coding sequence in humans**. *Proceedings of the National Academy of Sciences of the United States of America* 2013, **110**(2):600-605.

50. Jin L, Chakraborty R: **Population structure, stepwise mutations, heterozygote deficiency and thei implications in DNA forensics**. *Heredity* 1995, **74**:274-285.

51. Nei M: **Analysis of Gene Diversity in Subdivided Populations**. *Proceedings of the National Academy of Sciences of the United States of America* 1973, **70**(12):pp. 3321-3323.

52. Jost L: **G(ST) and its relatives do not measure differentiation**. *Mol Ecol* 2008, **17**(18):4015-4026.

53. Porras-Hurtado L, Ruiz Y, Santos C, Phillips C, Carracedo A, Lareu MV: **An overview of STRUCTURE: applications, parameter settings, and supporting software**. *Front Genet* 2013, **4**:98.

54. Parvanov ED, Petkov PM, Paigen K: **Prdm9 controls activation of mammalian recombination hotspots**. *Science* 2010, **327**(5967):835.

55. Cole F, Baudat F, Grey C, Keeney S, de Massy B, Jasin M: **Mouse tetrad analysis provides insights into recombination mechanisms and hotspot evolutionary dynamics**. *Nature genetics* 2014, **46**(10):1072-1080.

56. Billings T, Parvanov ED, Baker CL, Walker M, Paigen K, Petkov PM: **DNA binding specificities of the long zinc-finger recombination protein PRDM9**. *Genome biology* 2013, **14**(4):R35.

57. Striedner Y, Schwarz T, Welte T, Futschik A, Rant U, Tiemann-Boege I: **The long zinc finger domain of PRDM9 forms a highly stable and long-lived complex with its DNA recognition sequence**. *Chromosome Res* 2017, **25**(2):155-172.

58. Groeneveld LF, Atencia R, Garriga RM, Vigilant L: **High diversity at PRDM9 in chimpanzees and bonobos**. *PloS one* 2012, **7**(7):e39064.

59. Pastene LA, Goto M, Kanda N, Zerbini AN, Kerem D, Watanabe K, Bessho Y, Hasegawa M, Nielsen R, Larsen F *et al*: **Radiation and speciation of pelagic organisms during periods of global warming: the case of the common minke whale, Balaenoptera acutorostrata**. *Mol Ecol* 2007, **16**(7):1481-1495.

60. Rosel PE, Wilcox LA, Monteiro C, Tumlin MC: **First record of Antarctic minke whale, Balaenoptera bonaerensis, in the northern Gulf of Mexico**. *Marine Biodiversity Records* 2016, **9**(1).

61. Pratto F, Brick K, Khil P, Smagulova F, Petukhova GV, Camerini-Otero RD: **DNA recombination. Recombination initiation maps of individual human genomes**. *Science* 2014, **346**(6211):1256442.

62. Smagulova F, Brick K, Pu Y, Camerini-Otero RD, Petukhova GV: **The**
evolutionary turnover of recombination hot spots contributes to
speciation in mice. *Genes Dev* 2016, **30**(3):266-280.
63. Davies B, Hatton E, Altemose N, Hussin JG, Pratto F, Zhang G, Hinch AG,
Moralli D, Biggs D, Diaz R *et al*: **Re-engineering the zinc fingers of PRDM9**
reverses hybrid sterility in mice. *Nature* 2016, **530**(7589):171-176.
64. Forejt J: **Genetics: Asymmetric breaks in DNA cause sterility**. *Nature* 2016,
530(7589):167-168.
65. Zelazowski MJ, Cole F: **X marks the spot: PRDM9 rescues hybrid sterility**
by finding hidden treasure in the genome. *Nat Struct Mol Biol* 2016,
23(4):267-269.
66. Dzur-Gejdosova M, Simecek P, Gregorova S, Bhattacharyya T, Forejt J:
Dissecting the genetic architecture of F1 hybrid sterility in house mice.
Evolution 2012, **66**(11):3321-3335.
67. Bhattacharyya T, Reifova R, Gregorova S, Simecek P, Gergelits V, Mistrik M,
Martincova I, Pialek J, Forejt J: **X chromosome control of meiotic**
chromosome synapsis in mouse inter-subspecific hybrids. *PLoS genetics*
2014, **10**(2):e1004088.
68. Tamura K, Nei M: **Estimation of the number of nucleotide substitutions in**
the control region of mitochondrial DNA in humans and chimpanzees.
Molecular biology and evolution 1993, **10**(3):512-526.
830

Availability of data and materials

Most data generated or analyzed during this study are included in this published article

and its supplementary information files, Genetic data generated and analyzed during

the current study are available in the Zenodo repository,

https://doi.org/10.5281/zenodo.4309437, and R Scripts are available from

https://gitlab.gwdg.de/mpievolbio-it/repeatr.

Author Contributions

ED designed the experiments, analyzed data and wrote parts of the manuscript. KKU

designed experiments and analyzed the data, WA provided the samples and wrote

parts of the manuscript and L.O.-H. conceptualized the study, designed the

experiments, analyzed data and wrote the manuscript. All authors read and approved

the final version of the manuscript.

Figures and Tables

846
Figure 1 Diversity in the Cys2His2-ZnF domain in minke whales. (A) *Prdm9* gene
annotation in *Balaenoptera acutorostrata scammoni* genome reference, with primer
site annotations. We predicted the PRDM9 protein (B) Representative PCR products
of DNAs from individuals from AN IV (294,271), AN V (1653), NP (K8030) and NA
(MN2) showing variation in the number of minisatellite repeats units (number of ZnFs
in brackets) (C) Stylized structure of C2H2-ZnF binding to DNA, with nucleotide-
specificity conferred by amino-acids in positions -1, +2, +3, and +6 of alpha-helices.
(D) PRDM9 C2H2-ZnF arrays identified in minke whales, named with broad-scale
sampling location, Northern Hemisphere (NH) and Southern Hemisphere (SH) and
total number of ZnFs in the C2H2-ZnFs domain. (E) Types of PRDM9 C2H2-ZnFs in
minke whales. We generate three-letter-codes using the IUPAC nomenclature of
amino-acids involved in DNA-binding. All variable amino acids are colored, and
asterisks label C2H2-ZnFs also present in genome references of *Balaenoptera*
acutorostrata scammoni (*Ba) or *Balaenoptera bonarensis* (*Bb), see also
Supplementary Figure 4.
862

A *Prdm9* Phylogenetic Analyses

B *Balaenoptera acutorostrata*

C *Balaenoptera bonarensis*

D *Balaenoptera bonarensis sampling locations*

Figure 2 *Prdm9* Phylogenetic analyses and allele frequencies at different geographical scales. (A) PRDM9 phylogenetic analyses across all geographical regions, including several outgroups, from top to bottom: **-1, +3, +6**: PRDM9 variants as in Figure 1. **# ZnFs:** number of ZnFs **pop:** assigned population of individuals (dark blue): North Pacific, (light blue) North Atlantic, (light green) Antarctic Area IV, (dark green) Antarctic Area V, as well as Genome Reference alleles: (blue*) Balaenoptera acutorostrata scammoni*, (red) *Balaenoptera bonarensis* and outgroups (pink) *Tursiops Truncatus* **allele:** PRDM9 coding minisatellite allele colored as the translated variant from Figure 1. **Tree:** Nucleotide Phylogeny of PRDM9 alleles. (B) allelic diversity in Common minke whales (C) allelic diversity in Antarctic minke whales (D) fine-scale diversity by Sampling locations of Antarctic minke whales on a map of the Southern Ocean (yellow delineations show protected areas).

Table 1 Nucleotide diversity [68] of the minisatellite coding for the PRDM9-C$_2$H$_2$-ZnF array, analyzed per sample region. Average pairwise nucleotide diversity was analyzed for the all sites, and also excluding the nucleotides coding for amino-acids at hypervariable sites (-1, +3, +6) in the alpha-helix of C$_2$H$_2$-ZnFs.

Sample region	Area V	Area IV	North Atlantic	North Pacific

Number of analyzed 84bp repeat units per population	554	577	170	40
Population theta (segregating sites)	2,031	1,731	2,102	2,116
Average pairwise nucleotide diversity (excluding hypervariable sites)	0.0148±0.0001	0.0143±0.0001	0.0184±0.0002	0.0183± 0.0002
Average pairwise nucleotide diversity (including hypervariable sites)	0.0382±0.0005	0.0374±0.0005	0.0434±0.0006	0.0437±0.0006

882
Figure 3 Population structure of minke whales (NP) North Pacific, (NA) North Atlantic (ANIV) Antarctic Area IV (ANV) Antarctic Area V (A) Phylogeny of minke whale mitochondrial D-loop region, containing the hypervariable segments (HVS). (B) Population STRUCTURE analyses of minisatellites with *a priori* location information. (C) Log probability of data L(K) as a function of K (mean ± SD) (D) Magnitude of ΔK as a function of K (mean ± SD over 50 replicates).

889
890

891
Figure 4 PRDM9 Array binding predictions and population frequency based on
ZnFs#3-#6 "core motif" combinations across all individuals
894

896
897 **Supplementary Figure 1: PRDM9 Phylogenetic analysis across *Cetartiodactyla*,**
898 using N-terminal KRAB, SSXRD and SET domains, with data from public genomic
899 databases collected in (
900 **Supplementary Table 2**)
901

Supplementary Figure 2: **Alignment of proximal ZnFs of *Cetartiodactyla***, in which

this ZnF (outside the ZnF array) was present in the public genomic databases (

Supplementary Table **2**). The proximal ZnF is fully conserved across *Balaenoptera*. A

single amino-acid change at position 22 in bowhead whale (*Balaena mysticetus)* is

observed, compared to *Balaenoptera*, with the DNA contacting residues remaining

conserved.

Supplementary Figure 3: Alignment of second ZnFs of all *Cetartiodactyla*, where

this ZnF was present in public genomic databases (

Supplementary Table **2**). Across *Cetacea*, ZnF#2 are conserved at the amino-acids

13, 16 and 19, which code for the DNA-binding positions of the alpha-helix of each

ZnF. Amino-acids 19 is changed between *Cetacea* and *Ruminantia*, and additional

changes are observed at amino-acids 5, 6, 18, and 24.

Supplementary Figure 4: All PRDM9 C$_2$H$_2$-ZnFs types in minke whales. We generate three-letter-codes for each C$_2$H$_2$-ZnF using the IUPAC nomenclature of amino-acids involved in DNA-binding. All variable amino acids are colored, and asterisks label C$_2$H$_2$-ZnFs also present in genome references of *Balaenoptera acutorostrata scammoni* (*Ba) or *Balaenoptera bonarensis* (*Bb).

926
927 **Supplementary Figure 5: Signals of selection on 84bp nucleotide repeats coding**
928 **for each ZnF identified in this study.** "Site" shows zinc-fingers amino-acid positions,
929 from top to bottom: a site is classified based on whether α>β (negative selection) or
930 α<β (positive selection). (A) p-value is significant when <0,05 (B) fel.α instantaneous
931 synonymous site rate (C) fel.β instantaneous non-synonymous site rate.
932

Supplementary Table 1: Descriptive statistics of microsatellites used in this study
analyzed using (Dieringer and Schlötterer 2003). The richness of allelic variants
(number of observed alleles) (**A**) Expected Heterozygosity (**H$_E$**) observed
Heterozygosity (**H$_O$**)**,** excess or deficiency from the average number of heterozygotes
(**F$_{IS}$**) and (**H$_S$**), the probability that two random alleles from the same, randomly
chosen subpopulation are different by state.

Locus	Origin	A	Allele sizes	H$_E$	H$_O$	FIS	H$_s$
GATA028	AN IV	20	148-272	0,93	0,74	0,21	2,73
	AN V	20	148-274	0,92	0,88	0,05	2,67
	NA	-	-	-	-	-	-
	NP	-	-	-	-	-	-
GT575	AN IV	26	190-262	0,95	0,95	0	3,00
	AN V	29	185-262	0,95	0,97	-0,02	3,06
	NA	8	212-240	0,85	0,89	-0,06	1,89
	NP	5	206-216	0,89	1,00	-0,22	1,56
GT310	AN IV	18	81-135	0,88	0,78	0,11	2,41
	AN V	18	81-137	0,90	0,82	0,09	2,46
	NA	5	113-127	0,59	0,33	0,43	1,01
	NP	6	107-121	0,93	0,75	0,14	1,73
EV001	AN IV	13	103-147	0,83	0,95	-0,14	1,97
	AN V	14	105-141	0,82	0,78	0,04	1,97
	NA	5	139-153	0,72	0,56	0,22	1,35
	NP	4	131-151	0,86	0	1,00	1,39
GATA098	AN IV	9	76-108	0,83	0,89	-0,08	1,85
	AN V	10	70-108	0,84	0,97	-0,16	1,94
	NA	5	80-96	0,73	0,72	-0,01	1,35
	NP	3	84-92	0,68	1,00	-0,65	0,97
GT509	AN IV	19	179-219	0,92	0,89	0,03	2,66
	AN V	20	179-217	0,93	0,92	0,00	2,70
	NA	10	191-213	0,85	1,00	-0,19	1,97
	NP	5	195-211	0,89	1,00	-0,22	1,56
GATA417	AN IV	17	188-264	0,92	0,93	-0,02	2,58
	AN V	19	184-260	0,92	0,92	-0,01	2,60
	NA	8	212-240	0,86	0,94	-0,11	1,91
	NP	5	204-242	0,89	0,75	0,11	1,56
GT023	AN IV	14	90-118	0,87	0,81	0,07	2,24
	AN V	15	86-118	0,88	0,95	-0,08	2,36
	NA	8	92-108	0,77	0,78	-0,02	1,64
	NP	5	94-110	0,86	1,00	-0,27	1,49
EV037	AN IV	18	184-222	0,91	0,49	0,46	2,53
	AN V	16	184-222	0,93	0,52	0,44	2,59
	NA	7	192-208	0,72	0,72	-0,02	1,52
	NP	5	176-196	0,86	1,00	-0,27	1,49
	Origin	A		H$_e$	H$_o$	FIS	H$_s$

Average	AN IV	17,1		0,89	0,82	0,07	2,44		
	AN V	17,9		0,90	0,86	0,04	2,48		
	NA	7		0,76	0,74	0,03	1,58		
	NP	4,75		0,86	0,81	-0,05	1,47		

Supplementary Table 2: Public genome resources for PRDM9 occurrence and
subsequent protein domain prediction with InterProScan. Note, that a 'normal'
PRDM9 domain structure consists of a single C_2H_2 ZnF domain followed by an C_2H_2-
ZnF-array. C_2H_2-ZnF prediction is based on http://zf.princeton.edu/fingerSelect.php.
According to HMMER, most confident ZnF domains should have scores > 17.7, C_2H_2
ZnF domains with lower scores are indicated in brackets and were excluded from all
downstream analyses.

SPECIES	COMMON	ALIAS	TAXID	KRAB	SSXRD	SET	PROXIMAL C_2H_2 ZNF	C_2H_2-ZNF ARRAY	URL-GENOME
BALAENOPTERA ACUTOROSTRATA	minke whale	BALACU	9767	x	x	x	x	13(1)	https://ftp.ncbi.nlm.nih.gov/genomes/all/GCF/00(/GCF_000493695.1_BalAcu1.0/GCF_0004936 cu1.0_genomic.fna.gz
BALAENOPTERA BONARENSIS	Antarctic minke whale	BALBON	33556	x	x	x	x	4(2)	https://ftp.ncbi.nlm.nih.gov/genomes/all/GCA/000/97 A_000978805.1_ASM97880v1/GCA_000978805.1_ 0v1_genomic.fna.gz
BALAENOPTERA MUSCULUS	blue whale	BALMUS	9771	-	x	x	x	5(2)	https://ftp.ncbi.nlm.nih.gov/genomes/all/GCA/009/87 A_009873245.2_mBalMus1.v2/GCA_009873245.2_ .v2_genomic.fna.gz
BALAENA MYSTICETUS	bowhead whale	BALMYS	27602	x	x	x	x	3(2)	http://alfred.liv.ac.uk/downloads/bowhead_whale/bov ale_scaffolds.zip
BOS TAURUS	cattle	BOSTAU	9913	x	x	x	x	13(1)	https://ftp.ncbi.nlm.nih.gov/genomes/all/GCF/002/26 _002263795.1_ARS-UCD1.2/GCF_002263795.1_AI UCD1.2_genomic.fna.gz
CAMELUS DROMEDARIUS	arabian camel	CAMDRO	9838	x	x	x	x	4(1)	https://ftp.ncbi.nlm.nih.gov/genomes/all/GCF/000/80 _000803125.2_CamDro3/GCF_000803125.2_CamD mic.fna.gz
CATAGONUS WAGNERI	chacoan peccary	CATWAG	51154	x	x	x	x	7	https://ftp.ncbi.nlm.nih.gov/genomes/all/GCA/004/02 A_004024745.2_CatWag_v2_BIUU_UCD/GCA_004 CatWag_v2_BIUU_UCD_genomic.fna.gz

Species	Common name	Code	Number					Ratio	URL
DELPHINAPTERUS LEUCAS	beluga whale	D E L L E U	9749	x	x	x	x	2(1)	ftp://ftp.ensembl.org/pub/release-100/fasta/delphinapterus_leucas/dna/Delphinapterus_ASM228892v3.dna.toplevel.fa.gz
ESCHRICHTIUS ROBUSTUS	grey wahle	E S C R O B	9764	x	x	x	x	2(3)	https://ftp.ncbi.nlm.nih.gov/genomes/all/GCA/002/18...A_002189225.1_ASM218922v1/GCA_002189225.1...922v1_genomic.fna.gz
EUBALAENA JAPONICA	north pacific right whale	E U B J A P	302098	x	x	x	x	2(2)	https://ftp.ncbi.nlm.nih.gov/genomes/all/GCA/004/36...A_004363455.1_EubJap_v1_BIUU/GCA_00436345...p_v1_BIUU_genomic.fna.gz
GLOBICEPHALA MELAS	long-finned pilot whale	G L O M E L	9731	x	x	x	x	3(1)	https://ftp.ncbi.nlm.nih.gov/genomes/all/GCF/006/54..._006547405.1_ASM654740v1/GCF_006547405.1_A...0v1_genomic.fna.gz
HIPPOPOTAMUS AMPHIBIUS	hippopotamus	H I P A M P	9833	x	x	x	-	-	https://ftp.ncbi.nlm.nih.gov/genomes/all/GCA/004/02...A_004027065.2_HipAmp_v2_BIUU_UCD/GCA_004...HipAmp_v2_BIUU_UCD_genomic.fna.gz
INIA GEOFFRENSIS	boutu	I N I G E O	9725	x	x	x	x	3(2)	https://ftp.ncbi.nlm.nih.gov/genomes/all/GCA/004/36...A_004363515.1_IniGeo_v1_BIUU/GCA_004363515..._v1_BIUU_genomic.fna.gz
LAGENORHYNCHUS OBLIQUIDENS	pacific white-sided dolphin	L A G O B L	90247	x	x	x	x	1(2)	https://ftp.ncbi.nlm.nih.gov/genomes/all/GCF/003/67..._003676395.1_ASM367639v1/GCF_003676395.1_A...9v1_genomic.fna.gz
LIPOTES VEXILLIFER	yangtze river dolphin	L I P V E X	118797	x	x	x	x	3(2)	https://ftp.ncbi.nlm.nih.gov/genomes/all/GCF/000/44..._000442215.1_Lipotes_vexillifer_v1/GCF_0004422...es_vexillifer_v1_genomic.fna.gz
MEGAPTERA NOVAEANGLIAE	humpback whale	M E G N O V	9773	x	x	x	x	4(2)	https://ftp.ncbi.nlm.nih.gov/genomes/all/GCA/004/32...A_004329385.1_megNov1/GCA_004329385.1_meg...omic.fna.gz
MESOPLODON BIDENS	sowerby's beaked whale	M E S B I D	48745	x	x	x	-	4(1)	https://ftp.ncbi.nlm.nih.gov/genomes/all/GCA/004/02...A_004027085.1_MesBid_v1_BIUU/GCA_00402708...d_v1_BIUU_genomic.fna.gz
MONODON MONOCEROS	narwhal	M O N M O N	40151	x	x	x	x	1(1)	https://ftp.ncbi.nlm.nih.gov/genomes/all/GCF/005/19..._005190385.1_NGI_Narwhal_1/GCF_005190385.1_...whal_1_genomic.fna.gz

NEOPHOCAENA ASIAEORIENTALIS ASIAEORIENTALIS	yangtze finless porpoise	N E O A S I	170 633 7	x	x	x	x	6	https://ftp.ncbi.nlm.nih.gov/genomes/all/GCF/003/03 _003031525.1_Neophocaena_asiaeorientalis_V1/G 1525.1_Neophocaena_asiaeorientalis_V1_genomic.
ORCINUS ORCA	killer whale	O R C O R C	973 3	x	x	x	x	3(1)	https://ftp.ncbi.nlm.nih.gov/genomes/all/GCF/000/33 _000331955.2_Oorc_1.1/GCF_000331955.2_Oorc_ mic.fna.gz
OVIS ARIES	sheep	O V I A R I	994 0	x	x	x	x	13(1)	https://ftp.ncbi.nlm.nih.gov/genomes/all/GCF/002/74 _002742125.1_Oar_rambouillet_v1.0/GCF_0027421 _rambouillet_v1.0_genomic.fna.gz
PHOCOENA PHOCOENA	harbor porpoise	P H O P H O	974 2	x	x	x	x	2	https://ftp.ncbi.nlm.nih.gov/genomes/all/GCA/003/07 A_003071005.1_ASM307100v1/GCA_003071005.1 100v1_genomic.fna.gz
PHYSETER CATODON	sperm whale	P H Y C A T	975 5	x	x	x	x	4(1)	https://ftp.ncbi.nlm.nih.gov/genomes/all/GCF/002/83 _002837175.2_ASM283717v2/GCF_002837175.2_A 7v2_genomic.fna.gz
SUS SCROFA	pig	S U S S C R	982 3	x	x	x	x	9	https://ftp.ncbi.nlm.nih.gov/genomes/all/GCF/000/00 _000003025.6_Sscrofa11.1/GCF_000003025.6_Ssc genomic.fna.gz
TRAGULUS KANCHIL	lesser mouse-deer	T R A K A N	108 813 1	x	x	x	x	5(1)	https://ftp.ncbi.nlm.nih.gov/genomes/all/GCA/006/40 A_006408655.1_LMD/GCA_006408655.1_LMD_ger gz
TURSIOPS ADUNCUS	indopacific bottlenose dolphin	T U R A D U	797 84	x	x	x	x	1(1)	https://ftp.ncbi.nlm.nih.gov/genomes/all/GCA/003/22 A_003227395.1_ASM322739v1/GCA_003227395.1 739v1_genomic.fna.gz
TURSIOPS TRUNCATUS	common bottlenose dolphin	T U R T R U	973 9	x	x	x	x	8(1)	https://ftp.ncbi.nlm.nih.gov/genomes/all/GCF/011/76 _011762595.1_mTurTru1.mat.Y/GCF_011762595.1 1.mat.Y_genomic.fna.gz
VICUGNA PACOS	alpaca	V I C P A C	305 38	x	x	x	x	4(1)	https://ftp.ncbi.nlm.nih.gov/genomes/all/GCF/000/16 _000164845.3_VicPac3.1/GCF_000164845.3_VicPa omic.fna.gz
ZIPHIUS CAVIROSTRIS	cuvier's beaked whale	Z I P C A V	976 0	x	x	x	-	4(1)	https://ftp.ncbi.nlm.nih.gov/genomes/all/GCA/004/36 A_004364475.1_ZipCav_v1_BIUU/GCA_00436447 _v1_BIUU_genomic.fna.gz

Supplementary Table 3: Public mitochondrial resources.

SPECIES	COMMON	ALIAS	TAXID	URL-SEQUENCE
BALAENOPTERA ACUTOROSTRATA	minke whale	BALACU	9767	https://www.ncbi.nlm.nih.gov/nuccore/X7200
BALAENOPTERA ACUTOROSTRATA	minke whale	BALACU	9767	https://www.ncbi.nlm.nih.gov/nuccore/AY230
BALAENOPTERA BONAERENSIS	Antarctic minke whale	BALBON	33556	https://www.ncbi.nlm.nih.gov/nuccore/M6040
BALAENOPTERA BOREALIS	sei whale	BALBOR	9768	https://www.ncbi.nlm.nih.gov/nuccore/NC_00
BALAENOPTERA BOREALIS	sei whale	BALBOR	9768	https://www.ncbi.nlm.nih.gov/nuccore/AP006
BALAENOPTERA BRYDEI	bryde whale	BALBRY	255365	https://www.ncbi.nlm.nih.gov/nuccore/AP006
BALAENOPTERA MUSCULUS	blue whale	BALMUS	9771	https://www.ncbi.nlm.nih.gov/nuccore/NC_00
BALAENOPTERA OMURAI	omura whale	BALOMU	255217	https://www.ncbi.nlm.nih.gov/nuccore/AB201
BALAENOPTERA PHYSALUS	fin whale	BALPHY	9770	https://www.ncbi.nlm.nih.gov/nuccore/NC_00
BALAENOPTERA PHYSALUS	fin whale	BALPHY	9770	https://www.ncbi.nlm.nih.gov/nuccore/KC572
BALAENA MYSTICETUS	bowhead whale	BALMYS	27602	https://www.ncbi.nlm.nih.gov/nuccore/NC_00
ESCHRICHTIUS ROBUSTUS	grey whale	ESCROB	9764	https://www.ncbi.nlm.nih.gov/nuccore/NC_00
ESCHRICHTIUS ROBUSTUS	grey whale	ESCROB	9764	https://www.ncbi.nlm.nih.gov/nuccore/X7220
EUBALAENA GLACIALIS	north atlantic right whale	EUBGLA	27606	https://www.ncbi.nlm.nih.gov/nuccore/NC_03
LIPOTES VEXILLIFER	yangtze river dolphin	LIPVEX	118797	https://www.ncbi.nlm.nih.gov/nuccore/NC_00
MEGAPTERA NOVAEANGLIAE	humpback whale	MEGNOV	9773	https://www.ncbi.nlm.nih.gov/nuccore/NC_00
ORCINUS ORCA	killer whale	ORCORC	9733	https://www.ncbi.nlm.nih.gov/nuccore/NC_02
PHYSETER CATODON	sperm whale	PHYCAT	9755	https://www.ncbi.nlm.nih.gov/nuccore/NC_00
TURSIOPS TRUNCATUS	common bottlenose dolphin	TURTRU	9739	https://www.ncbi.nlm.nih.gov/nuccore/NC_01

Supplementary Table 4: Primers used in this study

Method	Name	Sequence
Whole length Prdm9 amplification	Prdm9Bal_full_F	CCA CGG CCA AAG ATG CTT TC
PRDM9-Zinc-Finger Typing	Prdm9Bal_ZnFA_F	GAG CTG AGG TAT GTG GTC GC
	Prdm9Bal_full+ZnFA_R	ACT CAT GGG GGC AGG AGT AT
STR genotyping	EV001_F	[JOE] CCC TGC TCC CCA TTC TC
	EV001_R	ATA AAC TCT AAT ACA CTT CCT CCA AC
	EV037_F	[HEX] AGC TTG ATT TGG AAG TCA TGA
	EV037_R	TAG TAG AGC CGT GAT AAA GTG C
	GATA028_F	[6FAM] AAA GAC TGA GAT CTA TAG TTA
	GATA028_R	CGC TGA TAG ATT AGT CTA GG
	GATA098_F	[6FAM] TGT ACC CTG GAT GGA TAG ATT
	GATA098_R	TCA CCT TAT TTT GTC TGT CTG
	GATA417_F	[JOE] CTG AGA TAG CAG TTA CAT GGG
	GATA417_R	TCTGCTCAGGAAATTTTCAAG
	GT023_F	[HEX] GTT CCC AGG CTC TGCACT CTG
	GT023_R	CATTTCCTACCCACCTGTCAT
	GT211_F	[HEX] CTG CTC TAT TCT ATG AAA GCA
	GT211_R	CTCCAGTATACCTATCTTGTC
	GT310_F	[6FAM] GAA TAC TCC CAG TAG TTT CTC
	GT310_R	TAA CTT GTG GAA GAT GCC AAC
	GT509_F	[6FAM] CAG CTG CAA AAC CTT GAC ATT
	GT509_R	GTA AAA TGT TTC CAG TGC ATC
	GT575_F	[HEX] TAT AAG TGA ATA CAA AGA CCC
	GT575_R	ACC ATC AAC TGG AAG TCT TTC
Sex-Determination	ZFX_R	CAC TTA TGG GGG TAG TCC TTT
	ZFY_R	ATT ACA TGT CGT TTC AAA TCA
	ZFYX_F	[6FAM] ATA GGT CTG CAG ACT CTT CTA
Mitochondrial D-loop Sequencing	BP15851(M13F)	GTAAAACGACGGCCAGTGAAGAAGTATTACACTCCACCAT
	MN312(M13R)	CAGGAAACAGCTATGACCCGTGATCTAATGGAGCGGCCA
	MT3(M13Rev)	CAGGAAACAGCTATGACCCATCTAGACATTTTCAGTG
	MT4(M13F)	GTAAAACGACGGCCAGTCCTCCCTAAGACTCAAGGAAG

Ancestry

954 **Supplementary Figure 6 Population structure analyses on a set of ten**
955 **hypervariable microsatellite loci** using STRUCTURE without *a priori* location
956 information (NOLOCS)

958
Supplementary Figure 7 Principle of PRDM9 binding symmetry and asymmetry (A)
Identical "core motif" combinations are predicted to result in fully symmetric binding
and efficient DSB formation (Berg, et al. 2010; Berg, et al. 2011) (B) variants with
somewhat dissimilar "core motifs", are predicted to result in some degree of
asymmetry. In humans a similar degree of motif-match still allowed DSBs necessary
for successful recombination (Berg, et al. 2010; Berg, et al. 2011; Odenthal-Hesse, et
al. 2014) (C) hypothetical combination of the most common variant of minke whale
species, of two hemispheres, generating a putative interspecies hybrid combination.
Here we would predict asymmetric positioning of recombination initiation sites, which
is implicated in F_1-hybrid male sterility in mammals (Forejt 2016).
969
970