Title: Measuring Compound Eye Optics with Microscope and MicroCT Images

Author Names and Affiliations

John Paul Currea[1], Yash Sondhi[2], Akito Y. Kawahara[3], and Jamie Theobald[2]

[1]Department of Psychology, Florida International University, Miami, FL 33199, U.S.A.

[2]Department of Biological Sciences, Florida International University, Miami, FL 33199, U.S.A.

[3]Florida Museum of Natural History, University of Florida, Gainesville, FL, 32611, U.S.A.

Article Type: Methods

Abstract (≤150 words):

With a great variety of shapes and sizes, compound eye morphologies give insight into visual ecology,

development, and evolution, and inspire novel engineering. In contrast to our own *camera-type* eyes,

compound eyes reveal their resolution, sensitivity, and field of view externally, provided they have

spherical curvature and orthogonal ommatidia. Non-spherical compound eyes with skewed ommatidia

require measuring internal structures, such as with MicroCT (μCT). Thus far, there is no efficient tool

to characterize compound eye optics, from either 2D or 3D data, automatically. Here we present two

open-source programs: (1) the ommatidia detecting algorithm (ODA), which measures ommatidia

count and diameter in 2D images, and (2) a μCT pipeline (ODA-3D), which calculates anatomical

acuity, sensitivity, and field of view across the eye by applying the ODA to 3D data. We validate these

algorithms on images, images of replicas, and μCT eye scans from ants, fruit flies, moths, and a bee.

Arthropods, with about 1.3 million described species, represent roughly 80% of all known animal

species (Zhang 2013). They range vastly in size, with body lengths from the 85 µm ectoparasitic

crustacean *Tantulacus dieteri* (Mohrbeck, Arbizu, and Glatzel 2010) to the over 50 cm green rock

lobster, *Sagmariasus verreauxi* (Holthuis 1991; Kensler 1967), and lifestyle, with activity in nearly

every ecological niche. Likewise, arthropods wield an array of eye architectures, most commonly

compound eyes (Cronin et al. 2014; Land and Nilsson 2012a). The arthropod compound eye has been a

model for understanding cellular fate and neural development (Callier and Nijhout 2013; Perry et al.

2016; Ready, Hanson, and Benzer 1976) and comparing eyes across species can reveal underlying

selective pressures driving eye evolution (Casares and McGregor 2020; Currea, Smith, and Theobald

2018; Friedrich 2003; Gaspar et al. 2019; Harzsch and Hafner 2006). It has further sparked

improvements in artificial eyes and computer vision, and innovations including anti-reflective coatings

that imitate the graded refractive indices of some insect eyes, and three-dimensional eye ultrastructure

designs that enhance solar panel light absorption (Yang et al. 2013; Yano, Sugawara, and Taniguchi

2020; Zheng et al. 2019).

Eye morphology is fundamental to how animals see because it sets physical limitations on the capacity

to form images (Land and Nilsson 2012a). Depending on light intensity, spectral characteristics, and

image motion, some optimal eye architecture will maximize the ability to gather image information

(Land 1997; Snyder, Laughlin, and Stavenga 1977; Snyder, Stavenga, and Laughlin 1977). Because of

the critical role of eye morphology in understanding visual ecology, development, and evolution, we

offer a program to accurately and automatically characterize compound eye optics.

In contrast to the camera-type eyes we possess, compound eyes are made up of multiple, repeated

optical elements that are externally visible. These *ommatidia* individually direct light onto

photoreceptors. Contrary to popular belief, they generally do not produce a myriad of tiny images on

the retina, but average into the functional pixels of the transduced image. The number of ommatidia

therefore determines the total number of images an eye can form, or its spatial information capacity.

Ommatidia can be counted in micrographs, ranging from about 20 in the fairyfly *Kikiki huna* (body

length=158 µm; Huber and Beardsley 2000; Huber and Noyes 2013) to over 30,000 in large

dragonflies (Cronin et al. 2014). Compound eyes further divide into two structural groups: apposition

eyes, in which pigment cells between ommatidia restrict incoming light to a single rhabdom, such that

lens size limits optical sensitivity (Figure 1A), and superposition eyes, in which light travels through a

clear zone that allows many facets to contribute to each point (Figure 1B), thereby multiplying the final

sensitivity.

Previous studies have relied on painstaking manual counts and estimates to describe compound eye structure. Fortunately, ommatidia count and diameter estimation from 2D and 3D eye images can be automated. Although several algorithms and software plugins have been proposed, they currently require user input for each image, and frequently underestimate ommatidia counts (Woodman, Todd, and Staveley 2011), overestimate ommatidial diameter (Schramm et al. 2015), or were not validated against manual measurements or measurements in the literature (Diez-Hermano et al. 2015; Iyer et al. 2016; Vanhoutte, Michielsen, and Stavenga 2003). They do work in limited cases with a few hundred clearly separated ommatidia, but have not been tested on multiple species, over a substantial range of eye sizes, or with different media. Since the pre-print of this manuscript, a method has been proposed for processing CT data but it relies on access to proprietary MATLAB software (Tichit et al. 2022).

Here, we offer two open-source programs written in Python to characterize compound eyes: (1) the ommatidia detecting algorithm (ODA), which identifies individual facets, (the outward visible portion of the ommatidia), in 2D images, and (2) a multi-stage µCT pipeline (ODA-3D) which applies the ODA to segment ommatidia components and characterize the visual field. We test the reliability and validity of this technique on single images of 6 eye molds of 5 different ant species ranging in size, light micrographs of 29 fruit flies (*Drosophila melanogaster*), scanning electron micrographs (SEMs) of fruit flies (*D. melanogaster* and *D. mauritania*) and processed images of MicroCT (µCT) scans of one fruit fly (*D. mauritania*), 2 moths (*Manduca sexta* and *Deilephila elpenor*) and one bee (*Apis mellifera*).

For spherical eyes, the lens diameter measurements provided by the ODA can be divided by measurements of eye radius (using the luminous pseudopupil technique, for instance) to measure the angular separation of ommatidia, called the interommatidial (IO) angle (Figure 1 $\Delta\varphi$). The inverse of this angle limits spatial acuity (Land 1997; Snyder, Laughlin, et al. 1977; Snyder, Stavenga, et al. 1977). High spatial acuity affords many behaviors, such as prey, predator, and mate detection, and perceiving small changes in self-motion (Land 1997; Land and Nilsson 2012a). For spherical eyes, the IO angle is approximately: $\Delta\varphi = D/R$, where D is the ommatidial lens diameter and R is the radius of curvature, assuming the axes of all ommatidia converge to a central point. Fortunately, many compound eyes closely approximate the spherical model. Smaller compound eyes are often spherical and homogenous because photon noise and diffraction constrain the range of viable IO angles and ommatidial sizes (Snyder, Stavenga, et al. 1977). Likewise, superposition eyes are often roughly spherical because they must optically combine light from many ommatidia, which constrains their heterogeneity (Figure 1B; Land and Nilsson 2012a; Warrant, Bartsch, and Günther 1999).

Eyes with the longitudinal axes of ommatidia askew to the eye surface are not well approximated by a

spherical model. Skewed ommatidia can improve acuity at the expense of field of view (FOV) by

pointing more ommatidia onto a small visual field (Figure 1C), or increase FOV at the expense of

acuity by spreading a few ommatidia over a large visual field (Figure 1D), but in both cases sacrifice

sensitivity by reducing the effective aperture as a function of the skewness angle and refraction

(Stavenga 1979). For more information on the optical consequences of ommatidial skewness, see

Stavenga (1979:371–77). MicroCT (μCT) allows calculating visual parameters for non-spherical

compound eyes, measuring anatomical IO angles at very high spatial resolution, and segmenting

different tissues, such as visual neuropils, within the same dataset. It is a quickly growing technique for

comparative morphology (Baird and Taylor 2017; Buser et al. 2020), used on arthropods to study

muscles (Walker et al. 2014), brains (Smith et al. 2016), ocelli (Taylor et al. 2016), and eyes (Brodrick

et al. 2020; Gaspar et al. 2019; Taylor et al. 2018, 2020). Our second proposed method, ODA-3D,

segments individual corneal lenses or crystalline cones from a μCT image stack (using the ODA) to

measure the size, orientation, and spatial distribution of ommatidia. We validate it on μCT scans of

spherical fruit fly eyes, bees, and nonspherical honeybee eyes, and further demonstrate how ODA-3D

can detect oval eye features, measure regional changes in skewness, spatial acuity, and sensitivity, and

project onto world-referenced coordinates to accurately measure FOV.

Because there are so many arthropod species and compound eye morphologies, it is challenging but

valuable to characterize them in a meaningful, fast manner. Here we demonstrate the operational range

of two programs to automate this task as a function of image resolution and contrast and benchmark its

performance against estimates of the time needed to take comparable measurements by hand. Overall,

our proposed methods minimize the substantial labor that is typically required in characterizing optical

performance in compound eyes and therefore facilitate understanding their role in vision.

Methods

Specimens and eye imaging

Micrographs of glue eye molds or replicas were taken previously on 5 ant specimens from four ant

species: two *Notoncus ectatommoides* of the Formicinae subfamily (from Palavalli-Nettimi and

Narendra, 2018), a jumper ant (*Myrmecia nigrocincta*) and a bull ant (*M. tarsata*) of the Myrmeciinae

subfamily, and *Rhytidoponera inornata* of the Ectatomminae subfamily (from Palavalli-Nettimi et al.,

2019). Micrographs of 29 fruit fly eyes (*D. melanogaster*) were also drawn from Currea et al. (2018)

and SEMs of two fruit fly species (*D. melanogaster* and *D. mauritiana*) collected by Maike

Kittelmann. For the SEMs, fly heads were removed from the body and placed into Bouin's solution

(Sigma Aldrich) overnight at room temperature. Heads were then dehydrated in an ethanol series of

50%, 70% and 3x 100%. Heads were then critical point dried, mounted onto sticky carbon tabs on

12mm SEM stubs, sputter coated with 15 nm gold and imaged at 5kV in a Hitachi S-3400N with

secondary electrons.

We obtained micro-computed tomographs (μCTs) of a fruit fly (*D. mauritania*), tobacco hornworm

(*Manduca sexta*), elephant hawkmoth (*Deilephila elpenor*), and honeybee (*Apis mellifera*). The fruit fly

μCT was collected by Maike Kittelman and used with her permission. The head was fixed and

dehydrated among other heads in the same way as SEM samples. Once in 100% ethanol, they were

stained with 1% Iodine in ethanol before scanning at the TOMCAT beamline of the Swiss Light Source

(Paul Scherrer Institute, Switzerland). For more information on the scanning procedure, see Torres-

Oliva et al. (2021). Vouchered moth specimens from the Florida Natural History Museum were stored

at -20°C in 95% ethanol, then heads were sliced, with antennae removed, and soaked in staining

solution (I_2+KI, equal proportions 1.25% I_2 and 2.5% KI solutions) in Eppendorf vials or falcon tubes

for 36–48 hours. *M. sexta* was scanned with a Phoenix V|Tome|X M system with: a 180kv x-ray tube, a

diamond-tungsten target, 80 kV tube voltage, 110 μA current, 17.8 mm source object distance, 793

mm object-detector distance, and capture time adjusted to maximize absorption range for each scan.

The acquisition consisted of 2300 projections, 8 s each. GE's datos|x r software version 2.3 processed

raw x-ray data, producing voxel size of 4.50074 μm. Volume files were imported into VG StudioMax

version 3.3.3 (Volume Graphics, Heidelberg, Germany), eyes isolated with the segmentation tools, then

exported as Tiff stacks. *D. elpenor* was scanned with a Zeiss Xradia 520 Versa (Carl Zeiss Microscopy

GmbH, Jena, Germany), with: 80 kV tube voltage, 88 μA current, low energy filtering, 22.5 mm source

object distance, 210 mm object-detector distance, an indirect detector comprising a scintillator, a

0.392x optical lens, and a camera provided to us by Deborah Glass. The acquisition consisted of 3201

projections, 8 s each, with the adaptive motion correction option in Scout-and-Scan software (Carl

Zeiss Microscopy GmbH). The tomographic reconstruction automatically generated a 32-bit txrm set of

tomograms with an isotropic voxel size of 3.3250 μm. The XRM controller software (Carl Zeiss

Microscopy GmbH) converted data to a stack of 16-bit tiff file. A previously scanned honeybee eye

(Taylor et al. 2018; Tichit et al. 2022) was downloaded from *Morphosource* at:

https://www.morphosource.org/concern/parent/000396179/media/000396182. Voxel size: 5.0 μm. This

sample was scanned with a synchrotron x-ray source, (Baird and Taylor 2017; see Taylor et al., 2018

for the details).

Data Validation and Manual counts

For the microscope images, ommatidial counts and diameters were measured manually by the

researchers providing the datasets (Ravi Palavalli-Nettimi measured the ant eye replicas, Maike

Kittelman the SEMs, and John P. Currea the *D. melanogaster* micrographs). Image processing software

like ImageJ allowed us to manually annotate each ommatidium and estimate their diameter as the

average of several diameters taken across the eye. Our approach was different for the various CT

datasets. For the *D. mauritania* and *A. mellifera* datasets, we compared our results to previous

measurements taken on the same scan. Note that for *D. mauritania* this was a direct count but was a

density-based estimate for *A. mellifera*. Density-based estimates approximate the ommatidial density

across the surface of the eye by taking local measurements and averaging across the whole eye surface.

Our other CT stacks, *D. elpenor* and *M. sexta*, are new and don't have previous measurements in the

literature, so we compared our results to density-based estimates of other conspecifics.

Ommatidia Detecting Algorithm

The fourier transform is a mathematical transformation that decomposes arbitrary functions into

component sinusoids, which can highlight periodic or repeating patterns in a signal. For digital images,

the sinusoidal elements of a 2D Fast Fourier Transform (FFT) are plane waves (gratings), characterized

by contrast, frequency, phase, and orientation. Operations applied to the frequency representation

(reciprocal space) can be inverse-transformed to generate a filtered image. The hexagonal arrangement

of typical ommatidia has 3 major axes (Figure 2 B), each approximated by a grating, and filtering

frequencies higher than these generates a smooth image, with maxima near ommatidia centers. The

inverse of these frequencies, approximating the ommatidial diameter, also provides useful bounds for

easily applying local maxima detection algorithms to the smoothed image. In particular, our program

searches for maxima within 25% of the FFT-derived ommatidial diameter, which we found to be robust

even for less regular ommatidial lattices.

We developed a Python language module, the ommatidia detecting algorithm (ODA), which: (1)

generates a 2D FFT, (2) finds the three fundamental gratings as the local maxima closest to the

reciprocal image center, using autocorrelation to amplify periodic elements, (3) filters higher image

frequencies, (4) inverts the filtered 2D FFT, and (5) finds local maxima in the smoothed image (Figure

2 B). There are several options when running this that are described in depth in the documentation.

Importantly, the ODA can check for just the first 2 instead of the 3 fundamental frequencies, in

principle allowing the program to work on ommatidia arranged in a square lattice such as that found in

the reflecting superposition eyes of decapod crustaceans (Land and Nilsson 2012b). This option also

helps for noisy images where the highest fundamental frequency is sometimes mistaken as a harmonic

of one of the other two. For instance, although we used the default settings for all other results, we used

this option on the dataset of 29 *D. melanogaster* micrographs, resulting in much more accurate results

than those found without selecting this option. Also, users can check the results in the reciprocal image

with maxima superimposed using a graphical user interface we developed (Supplemental Figure 1 A).

The program stores ommatidia coordinates and calculates ommatidial diameter. An optional imported

mask (a white silhouette on a black background) can help avoid false positives outside of the eye.

Measuring Ommatidia using μCT

We have further used the ODA to process 3D μCT data. This pipeline: (1) imports a stack of images

representing a 3D dataset of points in the crystalline cones (Figure 3 A) or corneal lenses. Images can

be edited to delete irrelevant pixels, and our program can pre-filter data by choosing a density range in

a graphical interface, to further isolate crystalline cones, and can preview the whole dataset (Figure 4

A). Supplemental Figure 1 A shows a frame of the user interface offered by our program.

(2) projects coordinates onto 2D images processed by the ODA (Figure 3 B). The layer of crystalline

cones curves with little variation normal to its surface, allowing a sphere fit with least squares

regression. The algorithm transforms points to spherical coordinates and interpolates a continuous

surface, modeling the points' radii as a function of elevation and azimuth. Finally, it selects the cross-

sectional surface containing 50% of residuals (Figure 4 B). Supplemental Figure 1 B shows the residual

distance of each point from this surface, which is optionally displayed at the end of this stage of the

program. This allows you to check that the surface fit is not biased to any particular region of the eye.

(3) forms images of this cross-sectional sheet by taking 2D histograms of elevation and azimuth (as in

Figure 3 C but ignoring residual distance) in 90°X90° segments. Processing the surface in smaller

segments and recentering before forming each image avoids extreme spherical warping and improves

the accuracy of the algorithm. The ODA approximates lens centers within each segment, then

recombines them into the original coordinate system. Finally, each point is labeled by its nearest center,

effectively segmenting points into clusters corresponding to separate ommatidia (Figure 4 C). Although

this works well for spherical eyes, when ommatidia are substantially skewed the overlap of projected

clusters prevents the algorithm from segmenting correctly, often splitting up proper clusters among

their neighbors. Optionally, our program can identify these problematic clusters and attempt to correct

them using a custom clustering algorithm. Problematic clusters are identified as those with a large

proportion ($\geq$10%) of elements whose nearest neighbors were elements of a different cluster. For each

problematic cluster, the algorithm finds the immediate neighborhood of 6 clusters and applies a

nonlinear minimization algorithm (Scipy's differential evolution) to minimize both 1) the mean

projected distance between each point and its cluster center and 2) the aspect ratio of the cluster's

longest side to its shortest in 3D. Because the clusters are approximately parallel on a small scale and

elongated along the ommatidial axis, this process often converges on the plane orientation orthogonal

to the clusters' longitudinal axes. The program only incorporates these cluster corrections if they result

in substantially less problematic elements (<5%). This dramatically improves cluster segmentation for

non-spherical eyes like the honeybee scan and adds a trivial amount of time to the program's total

duration. Optionally, our program also allows the user to manually edit cluster centers (Supplemental

Figure 1 C–D) and then displays the outcome of this segmentation (Supplemental Figure 1 E.)

(4) approximates the ommatidia count and lens diameters with clusters. The distance between adjacent

centroids approximates lens diameter. The ideal ommatidial axis is derived from planes formed by

triplets of centers near the cluster, and we approximate the surface normal by averaging the normal

vector for each plane. Singular value decomposition finds the semi-axes of the ellipsoid of a cluster and

our program selects the one closest to the ideal axis to estimate the anatomical ommatidial axis. The

angular difference between ideal and anatomical axes estimates anatomical skewness.

Anatomical axes of neighboring ommatidia should yield reliable IO angles, but the raw axes are highly

variable. While greater resolution, or using other structures to extend approximations closer to the

intersection, could improve accuracy, our program reduces variability by replacing each anatomical

axis with the average axis of a local neighborhood of clusters (akin to the rosette averaging procedure

used in Wardill et al., 2017). The program allows the user to determine the radius of this neighborhood

in terms of ommatidial diameter: a radius of 1 specifies immediate neighbors within one diameter, 2

specifies immediate neighbors and immediate neighbors of those, and so on. Here we used a radius of

5. In general, this has the effect of reducing local variability in IO angles while maintaining global

patterns. However, it also skews axes close to the boundary of the eye towards the center due to the

asymmetrical participation of neighboring points, thus reducing FOV estimates. As a compromise, in

addition to allowing the user to determine the neighborhood radius, our program uses the

neighborhood-averaged angles for IO angle measurements but the raw axes for measuring the projected

FOV.

To calculate anatomical IO angles, our program partitions coordinates into evenly spaced vertical and

horizontal sections (Figure 4 D), in which we projected clusters onto a parallel plane. For instance, for

vertical sections, all clusters within a range of x values are considered and the 2D clusters formed by y

and z values determine the vertical angle component. Angles are then calculated along the 2D plane

using the locally averaged axes described above. This repeats independently for each vertical and

horizontal section. The process approximates a horizontal and vertical subtended angle for each cluster

pair and calculates the total angle as their hypotenuse. We call this hypotenuse the anatomical IO angle.

This method allows independent approximations for horizontal and vertical IO angles across the eye,

and by keeping track of cluster pair orientation along the eye surface, we can calculate horizontal and

vertical IO angles using the two-dimensional lattice proposed by Stavenga (1979; illustrated in Figure

6C).

Finally, the program generates two spreadsheets: (1) for each crystalline cone cluster, the Cartesian and

spherical coordinates of the centroid, the number and location of the points within its cluster, the

approximate lens diameter, and the ideal and anatomical axis direction vectors are saved per row; and

(2) for each pair of adjacent crystalline cones, the cones' indexes from spreadsheet 1, the center

coordinates for the two cones, the resulting orientation in polar coordinates, and the anatomical IO

angle between the two are saved per row. These allow approximations of how spatial acuity, optical

sensitivity, and the eye parameter (Snyder 1979) vary across the eye (Taylor et al. 2018). The full code

for measuring ommatidia with μCT is available at GitHub where you can download the Python

package and basic examples on how to use it (see Data Availability section).

ODA testing

To validate performance and speed of the ODA software, we applied the ODA to each image after

programmatically lowering image resolution (by bin-averaging the images into larger and larger bins)

and contrast in order to determine the performance constraints of the ODA (by reducing the 8-bit range

of brightness values to narrower and narrower distributions). We recorded the output ommatidial count

and diameter for each image and deterioration level and compared them to manual estimates of the

same. Using manual measurements of ommatidial diameter, we converted the image resolution into

units of pixels per diameter. Runtime was measured using the cProfile module in Python and the

degradation experiments were run on a Microsoft Surface Pro laptop with an 11th Gen Intel(R)

Core(TM) i7-1185G7 @ 3.00GHz processor and 15.8 GB of usable RAM.

Statistics and Reproducibility

269 For the most part, we report descriptive statistics to show the accuracy of our program. As in Table 1,

270 variables that follow a generally symmetrical distribution are reported as a mean ± the standard

271 deviation and those with skewed distributions, like IO angle, are reported as the median (interquartile

272 range). When reporting correlations, such as between manual and automatic measurements, we report

273 the Pearson correlation and the corresponding p-value of a two-tailed F-test.

Results

Microscope Images

We tested the ODA on 4 sets of images: 1) light micrographs of the flattened eye molds of five ants of

4 different species (Figure 5A), 2) light micrograph focus stacks of 5 *D. melanogaster* specimens, 3)

SEMs of 5 different *D. melanogaster* specimens, and 4) SEMs of 5 *D. mauritiana* specimens. To assess

the performance limitations of the ODA, we applied it to each image after programmatic degradation of

image resolution and contrast (see methods for more detail). We report the runtime, output number, and

diameter of ommatidia for each image and degradation level to compare with manual measurements.

Images at full resolution and contrast produced the most accurate automated measurements of

ommatidial count (automated/manual = 94 ± 13%; mean ± standard deviation) and diameter (86 ± 7%).

Among media types, the ant eye replicas were closest to manual measurements (count: 99 ± 3%;

diameter: 93 ± 3%), followed by the *D. melanogaster* micrographs (c: 105 ± 12%; d: 82 ± 9%), the *D.*

mauritiana SEMs (c: 88 ± 2%; d: 82 ± 5%), and the *D. melanogaster* SEMs (c: 81 ± 6%; d: 86 ± 4%).

The ant eye replicas likely performed best because they physically unwrap the eye surface, reducing

the distortion due to the eye curvature, and have a sharp, high contrast spot at the center of each

ommatidium unlike the smooth, low contrast SEMs.

Reducing spatial resolution had a predictable effect on ODA output (Figure 5B, left column). At

degraded resolutions, measurements of ommatidia count and diameter did not change substantially as

long as the pixel resolution was sufficiently above the Nyquist limit set by the ommatidial diameter.

Based on the Nyquist criterion (Shannon 1948), the image must have at least two pixels for every

ommatidial diameter to properly resolve the ommatidial lattice. To characterize this threshold

resolution, we measured the lowest image resolution resulting in a measurement greater than 50% of

the maximum relative ommatidial count and less than 50% of the maximum relative lens diameter per

subject. Across all media, the threshold resolution for ommatidial count was 4.2 ± 1.9 pixels per

diameter (px/D) and for lens diameter was 4.5 ± 2.3 px/D, close to the theoretical Nyquist limit of 2

px/D. The *D. melanogaster* SEMs performed better than the other media (2.9 ± 0.1 px/D), followed

closely by the ant eye replicas (3.2 ± 1.0 px/D). In terms of lens diameter, the ant eye replicas

performed just above the theoretical limit (2.1 ± 0.7 px/D). The runtime also followed a predictable

trend where higher resolutions resulted in longer runtimes. However, not even the highest resolutions

took longer than 8 seconds, which is a substantial improvement over the time needed to count the

ommatidia by hand (which is on the order of an hour depending on the number of ommatidia).

At lower contrasts, measurements of ommatidia count and diameter changed substantially for the

SEMs and 2 of the ant eye replicas at the lower contrasts (Figure 5B, right column). We characterized

the threshold contrasts in the same way as the threshold resolutions above. Across all media, the

threshold contrast for ommatidial count was 0.01 ± 0.01 root mean square (RMS) and for lens diameter

was 0.01 ± 0.003 RMS. Although the threshold contrast was roughly the same for all media, the SEMs

were particularly susceptible to reduced contrast, dropping from a relative mean ommatidial count of

81% to 2% for the *D. melanogaster* and 88% to 0% for the *D. mauritiana* SEMs at the lowest contrast

as opposed to reductions from 105% to 99% for the *D. melanogaster* micrographs and 99% to 69% for

the ant eye replicas. As for mean ommatidial diameter, the SEMs again showed the most sensitivity to

reduced contrast, increasing from 86% to 234% for the *D. melanogaster* and 82% to 93% for the *D.*

mauritiana SEMs at the lowest contrast as opposed to increases from 81.7% to 82.3% for the *D.*

melanogaster micrographs and 92% to 93% for the ant eye replicas. This may be due to the already low

contrast of SEMs, whereas the other images have high contrast reflections of a light source near the

center of each lens. Even so, the ODA performed well over a nearly tenfold range of contrast

reductions for all media. The ODA was successful on images that were such low contrast that we

struggled to see individual ommatidia ourselves. Further, contrast had no clear effect on runtime, which

again never exceeded 8 seconds. The runtime was roughly the same across all ant eye replicas—

ranging from 153 to 2626 ommatidia—suggesting that runtime is not substantially affected by

ommatidia count. The ODA should therefore work quickly on images with a resolution exceeding the

Nyquist limit and a reasonable contrast. A resolution of about 10 px/D would provide equivalent results

to higher resolutions and significantly reduce the program's runtime.

We also tested microscope images of the eyes of 29 vinegar flies (*D. melanogaster*; Figure 5C and D)

to determine ODA performance on comparisons within a species. Visual inspection of preliminary

results found that using just the first 2 fundamental frequencies resulted in substantially less false

positives than checking for all 3. These false positives—likely responsible for the overestimates

above—may be due to noise induced by the curvature of the eye and the quality of the image.

Automated counts and diameters shared strong and significant correlations with manual measurements

(counts: r = .81, df = 27, p ≪ .001; diameters: r = .76, df = 27, p ≪ .001), and automated counts were

$100 \pm 4\%$ and diameters $95 \pm 2\%$ of those taken by hand offering precise and relatively accurate

estimations.

μCT

We tested ODA-3D on eye scans of a fruit fly (*D. mauritiana*) two moth species (*M. sexta* and *D.*

elpenor) that we collected, and one honeybee (*A. mellifera*) used in Taylor et al. (2018; Figure 6 A).

The measurements of lens diameter, skewness, skewness-adjusted lens diameter, and spherical,

anatomical, and modeled IO angles for each specimen can be found in Table 1. We additionally tested

methods for non-spherical eyes on the honeybee scan, including features of an oval eye and projecting

onto world-referenced coordinates.

Automated ommatidia counts were 100.4% of the manual count on the same *D. mauritiana* scan, 85%

of the density-based count of the same *A. mellifera* scan (Taylor et al. 2018; Tichit et al. 2022), 112%

of measurements of a different *D. elpenor* specimen (Stöckl, O'Carroll, and Warrant 2017), and 98%

and 104% of measurements of different *M. sexta* specimens (Stöckl et al. 2017; White 2003). These

ommatidia had diameters (Figure 6 B) that were consistent with measurements in the literature: 95 –

115% for *D. mauritiana* (Posnien et al. 2012; Torres-Oliva et al. 2021), 96% for *A. mellifera* despite

missing ~15% of ommatidia (Taylor et al. 2018), 99 – 103% for *D. elpenor* (Stöckl et al. 2017;

Theobald, Warrant, and O'Carroll 2010), and 96 – 108 % for *M. sexta* (Stöckl et al. 2017; Theobald et

al. 2010; White 2003). We also profiled the runtime of the ODA-3D on the 4 specimens in comparison

to estimates of how long it would take to extract the clusters manually. Estimates of manual time are

based on Tichit et al. (2022), which recorded about 1.8 minutes to manually segment each of 100

crystalline cones in the *A. mellifera* stack. Assuming this same rate for all of our specimens, our

program offered comparable results 883 to 2425 times faster than manual segmentation (note the 1000-

fold difference in the y-axes). Moreover, these estimates only account for the segmentation stages of

the procedure, omitting the time needed for taking ommatidial and interommatidial measurements

which represented the majority of the ODA runtime for the two moth scans.

The fly eye and both moth eyes showed ommatidial axes with minor skew from spherical alignment.

The bee, however, showed axes with substantial skew, consistent with other anatomical measurements

(Baumgärtner 1928). Skew geometrically reduces effective lens diameter by the cosine of skew angle,

and further still by refraction of the image space (Stavenga 1979). The adjustment without accounting

for refraction produced marginal reductions for the spherical eyes: 0.3% in *D. mauritiana,* 0.5% in *D.*

elpenor, and 0.5% in *M.sexta,* but more substantial reduction for the oval eye: 2.5% in *A. mellifera*.

The spherical approximations for IO angle were closer to anatomical IO angle in moth eyes than the

honeybee eye: 86% in *D. mauritiana*, 82% in *D.elpenor*, and 80% for *M. sexta* versus 65% in *A.*

mellifera. Aside from the spherical approximation for *A. mellifera*, IO angles were consistent with

previous measurements in the literature: 54% for the spherical and 82% for the anatomical

approximation in *A. mellifera* (Taylor et al. 2018), 89 – 97% for the spherical and 108 – 127% for the

anatomical approximation in *D. elpenor* (Stöckl et al. 2017; Theobald et al. 2010), and 86 – 91% for

the spherical and 108 – 114% for the anatomical approximation in *M. sexta*, (Stöckl et al. 2017;

Theobald et al. 2010). To our knowledge, previous measurements of *D. mauritiana* IO angles are

unavailable in the literature, but our measurement of 3.9° is in the range of *Drosophila melanogaster*,

which is on average 4.5° and as low 3.4° (Gonzalez-Bellido, Wardill, and Juusola 2011).

Non-Spherical Properties

In bees, spherical coordinates largely accounted for vertical axis curvature, but vastly underestimated

horizontal curvature. To better characterize their visual field, we projected the ommatidial axes onto a

sphere outside of the eye, like the world-referenced projection of Taylor et al. (2018). We used the

center from step B of ODA-3D, which is near the center of the head and chose a radius of 10 cm based

on visual fixation behavior (Wehner and Flatt 1977). As opposed to *D. elpenor*, which had very similar

spherical and world-referenced projections, the *A. mellifera* spherical projection largely underestimated

the vertical FOV as 54° and horizontal FOV as 21°. The world-referenced visual field, subtending

about 110° horizontally and 126° vertically, was closer to previous measurements of 140° horizontally

and 162° vertically based on vectors orthogonal to the eye surface (Taylor et al. 2018). This

discrepancy was likely due to ODA-3D errors during the segmentation of highly skewed ommatidia in

the periphery (for example, see the horizontal slice insets in Figure 4D). Glial pigment cell interactions

between the crystalline cones or deformation of the soft cells could also have caused the error. For

precisely measuring the FOV of eyes with highly skewed ommatidia, we recommend a process that

uses subvolume unfolding like Tichit et al. (2022).

Nonetheless, by accounting for some ommatidial skewness, ODA-3D allows us to compare the

structural properties of spherical and non-spherical eyes. The bee eye is an oval eye, with ommatidial

axes intersecting at different points for horizontal and vertical IO pairs ($I_h \neq I_v$ in Figure 6C).

Anatomical IO angles are therefore separable into independent horizontal and vertical components ($\Delta\varphi_h$

and $\Delta\varphi_v$ in Figure 6C). For ommatidia arranged in a regular hexagonal lattice, the orientations of IO

pairs should fall into 3 modes separated by 60°; 2) the horizontal angle for horizontal IO pairs is $2\Delta\varphi_h$

while the horizontal angle for diagonal IO pairs is $\Delta\varphi_h$; 3) the vertical angle is 0 for horizontal pairs and

$\Delta\varphi_v$ for diagonal pairs; and 4) the proportion $\Delta\varphi_v / \Delta\varphi_h$ is approximately $1/\sqrt{3}$ (Stavenga 1979). Finally,

for an oval eye to follow a regular hexagonal lattice, 5) the vertical radius of curvature, R_v, must be 3

times the horizontal radius, R_h, whereas a spherical eye requires $R_v = R_h$.

Both eyes are consistent with the criteria of a regular hexagonal lattice. 1) The IO orientations of both

eyes follow trimodal distributions with modes separated by about 60° (Figure 6E). IO pairs within 15°

of the three modes were selected to measure horizontal and vertical angles and calculate the horizontal

and vertical IO angle components as in Figure 6C. For the moth eye, 2) the horizontal angle for

horizontal IO pairs, 1.42°, is nearly twice the horizontal angles for diagonal pairs, 0.71° + 0.60° =

1.31°; 3) the vertical angles 0.31°, are close to 0° for horizontal pairs and are nearly equal for diagonal

pairs, 1.14° and 1.21°; and 4) the proportion $\Delta\varphi_v / \Delta\varphi_h$= 0.40 is close to $1/\sqrt{3}$=.58. For the bee eye, 2)

the horizontal angle for horizontal IO pairs, 1.59°, is nearly twice the horizontal angle for diagonal

pairs, 0.96° + 0.74° = 1.70°; 3) the vertical angles for horizontal pairs, 0.54°, are close to 0° and are

nearly equal for diagonal pairs, 0.76° and .92°; and 4) the proportion $\Delta\varphi_v / \Delta\varphi_h$= 0.73 is close to .58.

For the oval eye criterion, we calculated the radius for each mode based on the equation for the IO

angle introduced above: $R = D/\Delta\varphi$, where R is the radius of curvature or intersection, D the ommatidial

diameter, and $\Delta\varphi$ the IO angle. If the radius of curvature follows an elliptical function of IO orientation,

then diagonal pairs should have a radius of curvature ~2.23 times the horizontal radius. In the moth

eye, both diagonal and horizontal pairs had roughly the same radius of intersection (R_d = 1188μm; R_h =

1079μm; R_d/R_h = 1.10). In the bee eye, however, the mean diagonal radius (R_d = 923μm) is 1.44 times

the mean radius for horizontal pairs (R_h = 640 μm), closer to an oval eye.

By combining the horizontal angle of diagonal pairs and half the horizontal angle of horizontal pairs,

we approximated the horizontal anatomical IO angle component, $\Delta\varphi_h = 0.82 \pm .81°$ (N = 11,312).

Again, this is almost certainly an underestimate due to poor horizontal segmentation of crystalline

cones in the lateral eye. Previous measurements in the literature which found $\Delta\varphi_h$ to be 1.1–1.9° across

different regions of the eye (Horridge 2005). By combining the vertical angles of diagonal pairs we

approximate $\Delta\varphi_v = .84 \pm 0.83°$ (N = 7,111) and the total anatomical IO angle ($\sqrt{\Delta\varphi_v^2 + \Delta\varphi_h^2}$) as $\Delta\varphi = $

1.17°. This is close to the estimate based on the anatomical angle of all IO pairs, 1.33 ± 1.07°. Despite

underestimated horizontal IO angle components, both estimates are consistent with measurements on

the same scan assuming ommatidial axes orthogonal to the eye surface, which found IO angles between

0.9° and 1.7°(Taylor et al. 2018).

426   **Discussion**

Our methods successfully automate the estimation of multiple visual parameters of compound eyes.

We tested compound eye images with the ODA, which filters spatial frequencies based on the

hexagonal arrangement of most ommatidia and applies a local maximum detector to identify their

centers. The ODA calculated ommatidial count and lens diameter from different media (eye molds,

microscope images, and µCT scans), taxa (ants, flies, moths, and a bee), sizes (hundreds to tens of

thousands of ommatidia), and eye types (apposition, neural superposition, and optical superposition). In

all cases, measurements provided by the program matched with manual measurements on the same

data, previous measurements in the literature, or both. Ommatidial counts were accurate when

compared to previous measurements on the same dataset or in the literature: 95 ± 15% for the datasets

in the ODA benchmark, with ant eye replicas performing the best at 99 ± 3%; 100 ± 4% for the *D.*

melanogaster micrographs; and 99.6% (range=85–112%) for the 4 CT scans. Ommatidial diameters

were also accurate: 86 ± 6% for the datasets in the ODA benchmark, with ant eye replicas performing

the best at 93 ± 3%; 95 ± 2% for the *D. melanogaster* micrographs; and 101% (96–105%) for the 4 CT

scans.

The ODA-3D, which integrated the ODA into a µCT pipeline, proved successful on scans of one

spherical fruit fly eye (*D. mauritania*), two spherical moth eyes (*D. elpenor* and *M. sexta*) and one oval

bee eye (*A. mellifera*). In addition to counts and diameters, ODA-3D estimated anatomical IO angles,

FOV, and skewness. Skewness angles were insignificant in moth eyes, which generally require

approximate sphericity for proper optical superposition. However the oval honeybee eye showed

significant skewness angles, implying reduced optical sensitivity, lower resolution and a greater FOV

horizontally. Again, estimates were consistent with manual measurements on the same data, previous

measurements in the literature, or both. High resolution 3D data additionally offered world-referenced

coordinates and measurements of resolution at different angles along the eye to better characterize the

visual field.

The great eye size range between and among invertebrate species (Casares and McGregor 2020; Currea

et al. 2018; Gaspar et al. 2019; Land 1997; Land and Nilsson 2012a; Shingleton et al. 2007), makes

compound eyes ideal for studying environmental reaction norms and allometry. Little allometry

research deals with compound eyes, and instead favors organs easily measured in one or two

dimensions (McDonald et al. 2018; Shingleton et al. 2007; Shingleton, Mirth, and Bates 2008). By

automating the more tedious tasks of characterizing compound eyes, our programs should help with

this challenge. For instance, ODA counts and diameters allow total cell count approximations and

correspond to the independent effects of cellular proliferation and growth during eye development.

Further, our program facilitates measuring allometry of visual performance, addressing the

environmental reaction norms of the anatomical determinants of vision. Further, progress in

understanding fruit fly eye development (*D. melanogaster*; Callier and Nijhout 2013; Casares and

McGregor 2020; Gaspar et al. 2019; Ready et al. 1976), makes compound eyes ideal for assessing

principles of eye development across different taxa (Casares and McGregor 2020; Friedrich 2003;

Harzsch and Hafner 2006). And because optics are the first limit to incoming visual information (Land

1997; Stavenga 1979), they inform electrophysiological and behavioral data to infer intermediate

neural processing (Currea et al. 2022, 2018; Gonzalez-Bellido et al. 2011; Juusola et al. 2017; Land

and Nilsson 2012a; Theobald et al. 2010; Wardill et al. 2017; Warrant 1999).

Our programs have some known limitations. Images for the ODA require sufficient resolution to

properly detect ommatidia. For regular images, pixel length must be at most half the smallest

ommatidial diameter according to the theoretical limit and is closer to .25 or lower in practice. For

μCT, if individual crystalline cones cannot be resolved at each layer, they are likely indiscriminable to

the ODA. Further, some species, preparations, and scanning procedures, capture better contrast

between crystalline cones and other structures while avoiding structural damage to the specimen. For

example, *M. sexta* crystalline cones contrasted sharply with the background scan when prefiltered with

just the threshold function. *D. elpenor*, however, had additional noise outside of the eye, and *A.*

mellifera had uneven exposure, altering density measurements across locations in the scan and

ultimately forcing omission of some data. This may be an unintentional consequence of the contrast

enhancing property of a synchrotron light source (Baird and Taylor 2017). Most importantly, ODA-3D

erroneously segmented highly skewed ommatidia in the *A. mellifera* scan, resulting in inaccuracies

downstream. This could likely be improved by incorporating nonspherical subvolume unfolding, like in

Tichit et al. (2022). Preservation techniques can cause small deformations in the eye and while ODA-

3D still works on minor deformations, scans of highly deformed eyes break certain assumptions about

the uniformity of the lattice and cannot be analyzed using this method. We also make certain

assumptions in calculating parameters like ommatidial diameter, interommatidial angle that may

introduce some inaccuracies at the edges, however we allow the user to omit data points that are

significant outliers.

Ultimately, anatomical measurements cannot replace optical techniques in measuring compound eye

optics (Stavenga 1979; Taylor et al. 2018). Light passing through an eye refracts depending on the

incident angle and index of refraction (Stavenga 1979). But our approximations used only incident

angle, so our measurements of the aperture-diminishing effect of skewness represent lower bounds, and

our measurements of IO angles are *anatomical*, not functional IO angles. Though skewness can be

somewhat corrected for, nothing can match optical techniques (Stavenga 1979; Taylor et al. 2018).

Future work will be needed to understand the limitations of both the ODA and ODA-3D. Because the

ODA depends on spatial frequencies corresponding inversely to the ommatidial diameter, an eye with a

wide range of diameters or high curvature may not work, and the ODA should be tested on eyes

containing acute or sensitive zones, such as the robberfly (Wardill et al. 2017) and lattices with

transitioning arrangements, such as hexagon to square in houseflies (Stavenga 1975) and male

blowflies (Smith et al. 2015). Likewise, the ODA-3D should be tested on non-spherical non-oval eyes.

While our program appropriately measured anatomical IO angles across the honeybee's oval eye and

actually corroborated its oval eye properties, it may not work when IO angles change dramatically like

in the robberfly (Wardill et al. 2017). Finally, the ODA-3D should be tested on non-insect arthropods.

We also plan to improve the program and are working on adding more features like 1) GUI tools to

manually detect and fixing ommatidial coordinates; 2) machine learning tools to to generate a mask for

the eye and to clean the 3D datasets; 3) use large, clean datasets to train a machine learning algorithm

to find ommatidia faster and more reliably.

Conclusion

Compound eyes are the most common eye type on Earth, found in nearly every ecological habitat and

visual environment, and varying widely in size, shape, and architecture. Because they are diverse,

ubiquitous, and subject to heavy selection pressure, they are crucial to understanding the evolution of

vision. Our programs contribute to this effort and are open source, easy to install, easy to incorporate

into custom pipelines, and downloadable as Python modules. By successfully measuring parameters

from a wide range of eye shapes and sizes, they should facilitate the study of the development,

evolution, and ecology of visual systems in non-model organisms.

References

Baird, Emily, and Gavin Taylor. 2017. "X-Ray Micro Computed-Tomography." *Current Biology*
27(8):R289–91. doi: 10.1016/j.cub.2017.01.066.
Baumgärtner, Herbert. 1928. "Der Formensinn Und Die Sehschärfe Der Bienen." *Zeitschrift Für*
Vergleichende Physiologie 7(1):56–143.
Brodrick, Emelie A., Nicholas W. Roberts, Lauren Sumner-Rooney, Christian M. Schlepütz, and
Martin J. How. 2020. "Light Adaptation Mechanisms in the Eye of the Fiddler Crab *Afruca*
Tangeri." *Journal of Comparative Neurology* cne.24973. doi: 10.1002/cne.24973.
Buser, T. J., O. F. Boyd, Á. Cortés, C. M. Donatelli, M. A. Kolmann, J. L. Luparell, J. A.
Pfeiffenberger, B. L. Sidlauskas, and A. P. Summers. 2020. "The Natural Historian's Guide to
the CT Galaxy: Step-by-Step Instructions for Preparing and Analyzing Computed Tomographic
(CT) Data Using Cross-Platform, Open Access Software." *Integrative Organismal Biology*
2(obaa009). doi: 10.1093/iob/obaa009.
Callier, Viviane, and H. Frederik Nijhout. 2013. "Body Size Determination in Insects: A Review and
Synthesis of Size- and Brain-Dependent and Independent Mechanisms." *Biological Reviews*
88(4):944–54. doi: 10.1111/brv.12033.
Casares, Fernando, and Alistair P. McGregor. 2020. "The Evolution and Development of Eye Size in
Flies." *WIREs Developmental Biology* n/a(n/a):e380. doi: 10.1002/wdev.380.
Cronin, Thomas W., Sönke Johnsen, N. Justin Marshall, and Eric J. Warrant. 2014. *Visual Ecology*.
Princeton University Press.
Currea, J. P., Joshua L. Smith, and J. C. Theobald. 2018. "Small Fruit Flies Sacrifice Temporal Acuity
to Maintain Contrast Sensitivity." *Vision Research* 149:1–8. doi: 10.1016/j.visres.2018.05.007.
Currea, John P., Rachel Frazer, Sara M. Wasserman, and Jamie Theobald. 2022. "Acuity and
Summation Strategies Differ in Vinegar and Desert Fruit Flies." *Iscience* 25(1):103637.
Diez-Hermano, Sergio, Jorge Valero, Cristina Rueda, Maria D. Ganfornina, and Diego Sanchez. 2015.
"An Automated Image Analysis Method to Measure Regularity in Biological Patterns: A Case
Study in a Drosophila Neurodegenerative Model." *Molecular Neurodegeneration* 10(1):9. doi:
10.1186/s13024-015-0005-z.
Friedrich, Markus. 2003. "Evolution of Insect Eye Development: First Insights from Fruit Fly,
Grasshopper and Flour Beetle." *Integrative and Comparative Biology* 43(4):508–21. doi:
10.1093/icb/43.4.508.
Gaspar, Pedro, Saad Arif, Lauren Sumner-Rooney, Maike Kittelmann, David L. Stern, Maria D. S.
Nunes, and Alistair P. McGregor. 2019. "Characterisation of the Genetic Architecture
Underlying Eye Size Variation within Drosophila Melanogaster and Drosophila Simulans."
BioRxiv 555698. doi: 10.1101/555698.
Gonzalez-Bellido, Paloma T., Trevor J. Wardill, and Mikko Juusola. 2011. "Compound Eyes and
Retinal Information Processing in Miniature Dipteran Species Match Their Specific Ecological
Demands." *Proceedings of the National Academy of Sciences* 108(10):4224–29. doi:
10.1073/pnas.1014438108.
Harzsch, Steffen, and Gary Hafner. 2006. "Evolution of Eye Development in Arthropods: Phylogenetic
Aspects." *Arthropod Structure & Development* 35(4):319–40. doi: 10.1016/j.asd.2006.08.009.
Holthuis, Lipke B. 1991. "Marine Species Identification Portal : Green Rock Lobster - Jasus
Verreauxi." *Food and Agriculture Organization of the United Nations* 13(125).
Horridge, Adrian. 2005. "The Spatial Resolutions of the Apposition Compound Eye and Its Neuro-
Sensory Feature Detectors: Observation versus Theory." *Journal of Insect Physiology*
51(3):243–66. doi: 10.1016/j.jinsphys.2004.11.018.
Huber, John, and John Noyes. 2013. "A New Genus and Species of Fairyfly, Tinkerbella Nana

(Hymenoptera, Mymaridae), with Comments on Its Sister Genus Kikiki, and Discussion on
Small Size Limits in Arthropods." *Journal of Hymenoptera Research* 32:17–44. doi:
10.3897/jhr.32.4663.
Huber, John T., and John W. Beardsley. 2000. "A New Genus of Fairyfly, Kikiki, from the Hawaiian
Islands (Hymenoptera: Mymaridae)." *Hawaiian Entomological Society*.
Iyer, Janani, Qingyu Wang, Thanh Le, Lucilla Pizzo, Sebastian Grönke, Surendra S. Ambegaokar,
Yuzuru Imai, Ashutosh Srivastava, Beatriz Llamusí Troisí, and Graeme Mardon. 2016.
"Quantitative Assessment of Eye Phenotypes for Functional Genetic Studies Using Drosophila
Melanogaster." *G3: Genes, Genomes, Genetics* 6(5):1427–37.
Juusola, Mikko, An Dau, Zhuoyi Song, Narendra Solanki, Diana Rien, David Jaciuch, Sidhartha Anil
Dongre, Florence Blanchard, Gonzalo G. de Polavieja, Roger C. Hardie, and Jouni Takalo.
2017. "Microsaccadic Sampling of Moving Image Information Provides Drosophila Hyperacute
Vision." *ELife* 6. doi: 10.7554/eLife.26117.
Kensler, Craig B. 1967. "The Distribution of Spiny Lobsters in New Zealand Waters (Crustacea:
Decapoda: Palinuridae)." *New Zealand Journal of Marine and Freshwater Research* 1(4):412–
20. doi: 10.1080/00288330.1967.9515216.
Land, Michael F. 1997. "Visual Acuity in Insects." *Annual Review of Entomology* 42(1):147–77.
Land, Michael F., and Dan-Eric Nilsson. 2012a. *Animal Eyes*. OUP Oxford.
Land, Michael F., and Dan-Eric Nilsson. 2012b. *Animal Eyes*. OUP Oxford.
McDonald, Jeanne M. C., Shampa M. Ghosh, Samuel J. L. Gascoigne, and Alexander W. Shingleton.
2018. "Plasticity Through Canalization: The Contrasting Effect of Temperature on Trait Size
and Growth in Drosophila." *Frontiers in Cell and Developmental Biology* 6:156. doi:
10.3389/fcell.2018.00156.
Menzel, J. G., H. Wunderer, and D. G. Stavenga. 1991. "Functional Morphology of the Divided
Compound Eye of the Honeybee Drone (Apis Mellifera)." *Tissue and Cell* 23(4):525–35. doi:
10.1016/0040-8166(91)90010-Q.
Mohrbeck, Inga, Pedro Martínez Arbizu, and Thomas Glatzel. 2010. "Tantulocarida (Crustacea) from
the Southern Ocean Deep Sea, and the Description of Three New Species of Tantulacus Huys,
Andersen & Kristensen, 1992." *Systematic Parasitology* 77(2):131–51. doi: 10.1007/s11230-
010-9260-0.
Palavalli-Nettimi, Ravindra, and Ajay Narendra. 2018. "Miniaturisation Decreases Visual Navigational
Competence in Ants." *Journal of Experimental Biology* 221(7):jeb177238.
Palavalli-Nettimi, Ravindra, Yuri Ogawa, Laura A. Ryan, Nathan S. Hart, and Ajay Narendra. 2019.
"Miniaturisation Reduces Contrast Sensitivity and Spatial Resolving Power in Ants." *Journal of
Experimental Biology* 222(12). doi: 10.1242/jeb.203018.
Perry, Michael, Michiyo Kinoshita, Giuseppe Saldi, Lucy Huo, Kentaro Arikawa, and Claude Desplan.
2016. "Molecular Logic behind the Three-Way Stochastic Choices That Expand Butterfly
Colour Vision." *Nature* 535(7611):280–84. doi: 10.1038/nature18616.
Posnien, Nico, Corinna Hopfen, Maarten Hilbrant, Margarita Ramos-Womack, Sophie Murat, Anna
Schönauer, Samantha L. Herbert, Maria D. S. Nunes, Saad Arif, Casper J. Breuker, Christian
Schlötterer, Philipp Mitteroecker, and Alistair P. McGregor. 2012. "Evolution of Eye
Morphology and Rhodopsin Expression in the Drosophila Melanogaster Species Subgroup."
PLOS ONE 7(5):e37346. doi: 10.1371/journal.pone.0037346.
Ready, Donald F., Thomas E. Hanson, and Seymour Benzer. 1976. "Development of the Drosophila
Retina, a Neurocrystalline Lattice." *Developmental Biology* 53(2):217–40. doi: 10.1016/0012-
1606(76)90225-6.
Schramm, Bartosz W., Agnieszka Gudowska, Filip Kapustka, Anna Maria Labecka, Marcin
Czarnoleski, and Jan Kozłowski. 2015. "Automated Measurement of Ommatidia in the
Compound Eyes of Beetles." *BioTechniques* 59(2):99–101. doi: 10.2144/000114316.

Shannon, C. E. 1948. "A Mathematical Theory of Communication." 53.

Shingleton, Alexander W., W. Anthony Frankino, Thomas Flatt, H. Frederik Nijhout, and Douglas. J.
Emlen. 2007. "Size and Shape: The Developmental Regulation of Static Allometry in Insects."
BioEssays 29(6):536–48. doi: 10.1002/bies.20584.

Shingleton, Alexander W., Christen K. Mirth, and Peter W. Bates. 2008. "Developmental Model of
Static Allometry in Holometabolous Insects." *Proceedings of the Royal Society of London B:*
Biological Sciences 275(1645):1875–85. doi: 10.1098/rspb.2008.0227.

Smith, Dylan B., Galina Bernhardt, Nigel E. Raine, Richard L. Abel, Dan Sykes, Farah Ahmed, Inti
Pedroso, and Richard J. Gill. 2016. "Exploring Miniature Insect Brains Using Micro-CT
Scanning Techniques." *Scientific Reports* 6(1). doi: 10.1038/srep21768.

Smith, J. L., N. A. Palermo, J. C. Theobald, and J. D. Wells. 2015. "Body Size, Rather Than Male Eye
Allometry, Explains Chrysomya Megacephala (Diptera: Calliphoridae) Activity in Low Light."
Journal of Insect Science 15(1):133. doi: 10.1093/jisesa/iev114.

Snyder, Allan W. 1979. "Physics of Vision in Compound Eyes." Pp. 225–313 in *Comparative*
Physiology and Evolution of Vision in Invertebrates, *Handbook of Sensory Physiology*, edited
by H. Autrum. Springer Berlin Heidelberg.

Snyder, Allan W., Simon B. Laughlin, and Doekele G. Stavenga. 1977. "Information Capacity of
Eyes." *Vision Research* 17(10):1163–75. doi: 10.1016/0042-6989(77)90151-1.

Snyder, Allan W., Doekele G. Stavenga, and Simon B. Laughlin. 1977. "Spatial Information Capacity
of Compound Eyes." *Journal of Comparative Physiology* 116(2):183–207.

Stavenga, Doekele G. 1975. "The Neural Superposition Eye and Its Optical Demands." *Journal of*
Comparative Physiology ? A 102(4):297–304. doi: 10.1007/BF01464342.

Stavenga, Doekele G. 1979. "Pseudopupils of Compound Eyes." Pp. 357–439 in *Comparative*
Physiology and Evolution of Vision in Invertebrates. Vol. 7 / 6 / 6 A, edited by H. Autrum.
Berlin, Heidelberg: Springer Berlin Heidelberg.

Stöckl, A. L., D. O'Carroll, and E. J. Warrant. 2017. "Higher-Order Neural Processing Tunes Motion
Neurons to Visual Ecology in Three Species of Hawkmoths." *Proceedings of the Royal Society*
B: Biological Sciences 284(1857):20170880. doi: 10.1098/rspb.2017.0880.

Taylor, Gavin J., Stephen A. Hall, Johan A. Gren, and Emily Baird. 2020. "Exploring the Visual World
of Fossilized and Modern Fungus Gnat Eyes (Diptera: Keroplatidae) with X-Ray
Microtomography." *Journal of The Royal Society Interface* 17(163):20190750. doi:
10.1098/rsif.2019.0750.

Taylor, Gavin J., Willi Ribi, Martin Bech, Andrew J. Bodey, Christoph Rau, Axel Steuwer, Eric J.
Warrant, and Emily Baird. 2016. "The Dual Function of Orchid Bee Ocelli as Revealed by X-
Ray Microtomography." *Current Biology* 26(10):1319–24. doi: 10.1016/j.cub.2016.03.038.

Taylor, Gavin J., Pierre Tichit, Marie D. Schmidt, Andrew J. Bodey, Christoph Rau, and Emily Baird.
2018. "Bumblebee Visual Allometry Results in Locally Improved Resolution and Globally
Improved Sensitivity." *BioRxiv*. doi: 10.1101/380527.

Theobald, J. C., Eric J. Warrant, and David C. O'Carroll. 2010. "Wide-Field Motion Tuning in
Nocturnal Hawkmoths." *Proceedings of the Royal Society of London B: Biological Sciences*
277(1683):853–60. doi: 10.1098/rspb.2009.1677.

Tichit, Pierre, Tunhe Zhou, Hans Martin Kjer, Vedrana Andersen Dahl, Anders Bjorholm Dahl, and
Emily Baird. 2022. "InSegtCone: Interactive Segmentation of Crystalline Cones in Compound
Eyes." *BMC Zoology* 7(1):10. doi: 10.1186/s40850-021-00101-w.

Torres-Oliva, Montserrat, Elisa Buchberger, Alexandra D. Buffry, Maike Kittelmann, Lauren Sumner-
Rooney, Pedro Gaspar, Georg C. Bullinger, Genoveva Guerrero, Fernando Casares, Saad Arif,
Nico Posnien, Maria D. S. Nunes, Alistair P. McGregor, and Isabel Almudi. 2021. "Differences
in Orthodenticle Expression Promote Ommatidial Size Variation between Drosophila Species."
2021.03.17.435774.

Vanhoutte, Kurt J. A., Kristel F. L. Michielsen, and Doekele G. Stavenga. 2003. "Analyzing the
Reflections from Single Ommatidia in the Butterfly Compound Eye with Voronoi Diagrams."
Journal of Neuroscience Methods 131(1):195–203. doi: 10.1016/j.jneumeth.2003.08.011.
Walker, Simon M., Daniel A. Schwyn, Rajmund Mokso, Martina Wicklein, Tonya Müller, Michael
Doube, Marco Stampanoni, Holger G. Krapp, and Graham K. Taylor. 2014. "In Vivo Time-
Resolved Microtomography Reveals the Mechanics of the Blowfly Flight Motor." *PLOS*
Biology 12(3):e1001823. doi: 10.1371/journal.pbio.1001823.
Wardill, Trevor J., Samuel T. Fabian, Ann C. Pettigrew, Doekele G. Stavenga, Karin Nordström, and
Paloma T. Gonzalez-Bellido. 2017. "A Novel Interception Strategy in a Miniature Robber Fly
with Extreme Visual Acuity." *Current Biology* 27(6):854–59. doi: 10.1016/j.cub.2017.01.050.
Warrant, ERIC, Klaus Bartsch, and CLAUDIA Günther. 1999. "Physiological Optics in the
Hummingbird Hawkmoth: A Compound Eye without Ommatidia." *Journal of Experimental*
Biology 202(5):497–511.
Warrant, Eric J. 1999. "Seeing Better at Night: Life Style, Eye Design and the Optimum Strategy of
Spatial and Temporal Summation." *Vision Research* 39(9):1611–30. doi: 10.1016/S0042-
6989(98)00262-4.
Wehner, R., and I. Flatt. 1977. "Visual Fixation in Freely Flying Bees." *Zeitschrift Für Naturforschung*
C 32(5–6):469–72.
White, R. H. 2003. "The Retina of Manduca Sexta: Rhodopsin Expression, the Mosaic of Green-, Blue-
and UV-Sensitive Photoreceptors, and Regional Specialization." *Journal of Experimental*
Biology 206(19):3337–48. doi: 10.1242/jeb.00571.
Woodman, Peter N., Amy M. Todd, and Brian E. Staveley. 2011. "Eyer: Automated Counting of
Ommatidia Using Image Processing Techniques." *Drosophila Information Service* 94:142.
Yang, Qiaoyin, Xu A. Zhang, Abhijeet Bagal, Wei Guo, and Chih-Hao Chang. 2013. "Antireflection
Effects at Nanostructured Material Interfaces and the Suppression of Thin-Film Interference."
Nanotechnology 24(23):235202. doi: 10.1088/0957-4484/24/23/235202.
Yano, Tomoya, Hiroyuki Sugawara, and Jun Taniguchi. 2020. "Moth-Eye Structured Mold Using
Sputtered Glassy Carbon Layer for Large-Scale Applications." *Micro and Nano Engineering*
9:100077. doi: 10.1016/j.mne.2020.100077.
Zhang, Zhi-Qiang. 2013. "Animal Biodiversity: An Update of Classification and Diversity in 2013. In :
Zhang, Z.-Q. (Ed.) Animal Biodiversity: An Outline of Higher-Level Classification and Survey
of Taxonomic Richness (Addenda 2013)." *Zootaxa* 3703(1):5–11. doi:
10.11646/zootaxa.3703.1.3.
Zheng, Yelong, Le Song, Jingxiong Huang, Haoyang Zhang, and Fengzhou Fang. 2019. "Detection of
the Three-Dimensional Trajectory of an Object Based on a Curved Bionic Compound Eye."
Optics Letters 44(17):4143–46. doi: 10.1364/OL.44.004143.
696
697

Author Contributions

JC: Conceptualization, Methodology, Software, Visualization, Writing- Original Draft.
YS: Conceptualization, Methodology, Writing- Review and Editing, Data Collection and Curation.
JT: Conceptualization, Methodology, Writing- Review and Editing, Supervision, Funding acquisition.
AK: Conceptualization, Methodology, Writing- Review and Editing, Funding acquisition.

Funding

This research was supported by grants from the National Science Foundation: IOS-1750833 to JT, BCS-1525371 to JC, DEB-1557007 to AYK and JT and IOS-1920895 to AYK. The content is solely the responsibility of the authors and does not necessarily represent the official views of the National Institutes of Health or the National Science Foundation.

Acknowledgements

We thank members of the Kawahara Lab at the Florida Museum of Natural History, McGuire Center for Lepidoptera and Biodiversity, for raising and vouchering moth specimens. We thank the University of Florida CT facility for helping acquire the *M. sexta* CT scan, and Kelly Dexter for help with sample staining. We thank Deborah Glass and Ian Kitching from the Natural History Museum of London for helping with the CT procedure and sharing the *D. elpenor* CT scan, which study was funded by NERC grant number NE/P003915/1 to IJK. We thank Maike Kittelman for sharing her SEMs and µCT of two fruit fly species (*D. melanogaster* and *D. mauritiana*) and for advice in using available 3D software. Gavin Taylor and Emily Baird shared their CT scans of *A. mellifera*. Michael Reisser and Arthur Zhao provided valuable discussions regarding the µCT pipeline.

Code Availability

The ODA module, written in Python, is available for download: www.github.com/jpcurrea/ODA/. The version of the package used in this paper is available in our online dataset at https://doi.org/10.6084/m9.figshare.21521142.

Data Availability

All datasets and code are freely available at https://doi.org/10.6084/m9.figshare.21521142. The image stack of the µCT data for the *A. mellifera* scan were drawn from Taylor et al. (2018) and are available at https://www.morphosource.org/Detail/ProjectDetail/Show/project_id/646.

Figure 1: Diagrams of apposition and superposition eyes demonstrating the geometric tradeoffs between lens diameter (D),
interommatidial angle (Δφ), and field of view (FOV) for spherical (A. and B.) and non-spherical eyes (C. and D.). **A.** In
spherical apposition eyes, D directly determines sensitivity while Δφ inversely determines acuity. **B.** In superposition eyes,
migrating pigment (indicated by the arrows) allows the ommatidia to share light, increasing the eye's sensitivity. As a result,
these eyes must adhere to a spherical design. **C-D.** In non-spherical eyes, the intersection of ommatidial axes differs from the
center of curvature, with ommatidial axes askew from the surface of the eye. Consequently, FOV and Δφ are not externally
measurable and the effect of D on sensitivity is reduced by greater angles of skewness. **C.** When the distance to the intersection
is greater than the radius of curvature, FOV and Δφ decrease, increasing average spatial acuity by directing more ommatidia
over a smaller total angle. **D.** Inversely, when the distance to the intersection is less than the radius of curvature, FOV and Δφ
increase, decreasing average spatial acuity by directing fewer ommatidia over a smaller total angle. In both cases, optical
sensitivity is lost because skewness reduces the effective aperture of the ommatidia.

Figure 2: The ommatidia detecting algorithm (ODA) extracts periodic signals in a 2D image using the FFT. **A**. In the frequency domain of a 2D FFT, called the reciprocal space, gratings are represented by an x- and y-frequency. The polar coordinates represent visual properties of the corresponding grating. The radial distance is a grating's spatial frequency, with high frequencies farther from the origin. The polar angle is the grating's orientation , which is perpendicular to the grating's wavefront. Notice that the reciprocal space has local maxima (in red) approximately equal to the input grating parameters (polar angle=45° and radial distance=.047±.005). **B**. In a hexagonal lattice, there are three major axes (here in blue, green, and red). Each axis corresponds to a 2D spatial frequency (and it's negative), visible in the image's reciprocal space. The periodic nature of the axes results in harmonic frequencies. A low-pass filter returns a version of our original image primarily representing these three axes. The center of each ommatidium is found at the local maxima of the filtered image.

Figure 3: **A.** ODA-3D starts with an image stack, which may be pre-filtered and cleaned of unrelated structures. **B.** Then we
filter the relevant density values and fit a surface to the coordinates, allowing us to split the points into two sets or cross
sections, inside and outside the fitted surface. **C.** For each cross section, we generate spherically projected, rasterized images
that are processed by the FFT method for locating ommatidia, yielding approximate centers for the ommatidia. With these
centers, we can find the coordinate clusters corresponding to independent ommatidia **D.** These can then be used to
automatically measure eye parameters.

Figure 4: Checkpoints along ODA-3D for the honeybee scan. **A. Import Stack**: The pipeline started with an image stack that we pre-filtered and manually cleaned of unrelated structures. **B. Get Cross-Section:** A surface was fitted to the coordinates identifying a cross-section of points within 50% of the residuals. We generated a spherically projected, 2D histogram of the crystalline cone coordinates colored by their residual distance from the cross-section. **C. Find Ommatidial Clusters**: We apply the ODA to label individual crystalline cones. The ODA locates the cluster centers, allowing us to partition the coordinates based on their distance to those centers (boundaries indicated by color filled polygons). Then we cluster points based on their nearest center and apply our custom clustering algorithm to improve segmentation of more skewed ommatidia. **D. Measure Visual Parameters:** Ommatidial diameter corresponds to the average distance between clusters and their adjacent neighbors. Ommatidial count corresponds to the number of clusters. For non-spherical eyes, we can partition the 3D coordinates into vertical (*left*) and horizontal (*right*) cross-sections in order to calculate independent vertical and horizontal anatomical IO angle components accounting for skewness in the ommatidial axes. The insets zoom into regions around the 25th, 50th, and 75th percentile ommatidia along the y-axis (*left*) and x-axis (*right*). Note: in **C** and **D** the different colors signify different crystalline cone clusters. In **D**, the black lines follow the ommatidial axes and the black dots indicate the cluster centroids. Also in **D**, note how some ommatidia in the periphery (for example, horizontal insets row 2, columns 2 and 3) were erroneously segmented and thus underestimate their skewness and IO angles.

Figure 5: The ODA successfully approximated ommatidial counts and diameters when compared to manual measurements.
A. When applied to 6 ant eye molds of 5 species ranging in overall size and lattice regularity, the automated counts were 99%

and the diameters were 93% of those measured by hand. For each image, the program missed relatively few ommatidia and
the lens diameter measurements were successful even when they varied substantially within an image (as in #5 from the left).
Species from left to right: *Notoncus ectatommoides*, *Notoncus ectatommoides*, *Rhytidoponera inornata*, *Myrmecia*
nigrocincta, and *Myrmecia tarsata*. Scale bars are 50μm. **B.** Benchmark performance of ODA on micrographs of diminished
spatial resolution (left) and contrast (right). We present 3 performance metrics as a function of resolution and contrast: relative
lens count equal to the ratio of automatic to manual ommatidia counts (top); relative lens diameter equal to the ratio of
automatic to manually measured lens diameters (middle); and the total duration or runtime of the ODA (bottom). **C.** An
example fruit fly eye (*D. melanogaster)* micrograph with the automated ommatidia centers superimposed as points colored
according to the measured diameter using the same colormap as in A. **D.** When applied to 29 microscope images of fruit fly
eyes from the same species (*D. melanogaster*), the automated counts were 100% and the diameters were 95% of those
measured by hand, with correlations of .81 and .76. Again, there were relatively low rates of false positives and negatives.

Figure 6: Comparing visual fields based on the spherical approximation. **A.** ODA-3D allows us to map the visual fields of our four specimens *(Drosophila mauritania, Apis mellifera, Deilephila elpenor, and Manduca sexta)*. Using their spherical projection, we map the azimuth (along the lateral-medial axis) and elevation (along the ventral-dorsal axis) angles of ommatidia and represent lens diameter with color. Note that the bee eye was flipped horizontally from the images in Figure 4. The colorbar to the right indicates lens diameter and shows diameter histograms for each species in white and gray. Insets zoom in on a 20°x20° region in the center of each eye, showing ommatidial lattices in more detail. Note that the spherical projection of the bee eye underestimates its visual field as explained in the text. **B.** The estimated time to segment each stack, based on the 2 minute per cone estimate from Tichit et al. (2022), is plotted above the ODA-3D runtime. Note the 1000-fold difference in the y-axes illustrating that the ODA-3D runs at about 1000 times faster than the estimated time to segment the crystalline cones manually. **C.** Adapted from Stavenga, 1979. Bee eyes are approximately oval, and, unlike spherical eyes, have different intersection points for horizontal (I_h) and vertical (I_v) ommatidial pairs corresponding to different IO angle components horizontally ($\Delta\varphi_h$) and vertically ($\Delta\varphi_v$). For oval eyes, the horizontal component of the IO angle of horizontal pairs (orientation=0°) is $2\Delta\varphi_h$ while the vertical component is ~0° and the horizontal component of diagonal pairs (orientation=±60°) is $\Delta\varphi_h$ while the vertical component is $\Delta\varphi_v$. **D.** To demonstrate the difference between spherical and oval eye visual fields, we plot the spherical and world referenced projections of the ommatidial axes for the moth (*D. elpenor*) and bee (*A. mellifera*) eyes. Notice that there is little difference for the spherical moth eye but a substantial difference horizontally for the bee eye. **E.** The vertical and horizontal subtended angles of ommatidial pairs with respect to their orientation also demonstrate differences between the spherical moth (*D. elpenor)* eye and the oval bee (*A. mellifera)* eye. The pair orientations form trimodal distributions with means close to ±60° (diagonal pairs) and 0° (horizontal pairs) for both species, but the distributions are more uniform for the moth eye. Instead, the bee eye has significantly larger horizontal than vertical pairs and each pair group is more variable, resulting in a larger combined distribution of IO angles. Grayscale heatmaps present 2D histograms and medians (red point), IQRs (half-opacity red line), and 99% C.I.s (full opacity error bars) are plotted for each orientation ±15°.

Supplemental Figure 1: Graphical user interfaces for checking the progress of ODA-3D. **A.** Layers of the stack can be
displayed using the slider on the bottom and then prefiltered by selecting the lower and upper bounds of density measurements
using the vertical sliders on the right. **B.** Once the thin surface is fit to the filtered voxels, the goodness of fit across the eye
surface can be displayed. This allows the user to see if the surface was biased to any particular region. **C–D.** Once the
ommatidial centers are approximated for a given segment of the eye, the user can add, delete, or clear all estimated centers.
Each center point is colored based on its estimated diameter, so outliers are often highlighted by their extreme color. For
instance, the user can identify and delete the two largest points (white arrows) in the top right of C. producing the frame in D.
with an updated colormap. **E.** Then, the user can check the success of applying those centers to segment individual crystalline
cones, each colored with a random color to assist in comparisons. **F.** In addition, the user can view the outcome of each stage
in an interactive 3D interface at the end of the procedure.

	D. mauritiana	*A. mellifera*	*D. elpenor*	*M. sexta*
Count	979	4725	13004	26552
	975 *, 957 ± 22.89 [a], 1018 [b]	5440 *[c]	11508 [d]	27000 [e], 25641 [d]
Lens Diameter (µm)	17.69 ± 0.92	21.79 ± 1.75	28.47 ± 1.86	32.32 ± 1.53
	15.37 ± 1.89 [a], 18.55 ± 1.07 [b]	22.61 ± 1.96 *[c]	28 [d], 29 [g]	30 [e], 31 [d], 34 [g]
Skewness (°)	4.55 (7.02)	12.81 (11.73)	5.92 (4.28)	5.79 (4.51)
		≤ 50 [h]		
Adjusted Lens Diameter (µm)	17.40 ± 1.15	20.71 ± 2.33	28.18 ± 1.93	32.04 ± 1.65
Spherical IO Angle (°)	3.31 (0.26)	0.87 (0.11)	1.17 (0.10)	0.83 (0.05)
Anatomical IO Angle (°)	3.87 (1.64)	1.33 (1.07)	1.42 (0.69)	1.04 (0.52)
Modeled IO Angle (°)	3.85	1.05	1.34	0.93
		1.62 (0.61) *[c]	1.12 [d], 1.31 [g]	0.91 [d], 0.96 [g]

Table 1: Optical parameters for the four species are presented as a table. Values are mean ± s.d except for angular measurements, which show median (IQR). Comparable measurements from the literature or measured manually are in gray below corresponding values, with an asterisk indicating manual measurements of the same dataset. References are signified in superscript letters: [a] Posnien et al. (2012), [b] Torres-Oliva et al. (2021), [c] (Taylor et al. 2018), [d] Stöckl et al. (2017), [e] White (2003), [f] Menzel, Wunderer, and Stavenga (1991), [g] Theobald et al. (2010), [h] Baumgärtner (1928). Asterisks signify measurements taken on the same datasets.