Fetal estrogens are not involved in sex determination but critical for early ovarian differentiation in rabbits.

Geneviève Jolivet[1][*], Nathalie Daniel-Carlier[1], Erwana Harscoët[1], Eloïse Airaud[1], Aurélie Dewaele[1], Cloé Pierson[1], Frank Giton[2], Laurent Boulanger[1], Nathalie Daniel[1], Béatrice Mandon-Pépin[1], Maëlle Pannetier[1], Eric Pailhoux[1]

[1] Université Paris-Saclay, INRAE, ENVA, UVSQ, BREED, 78350, Jouy-en-Josas, France

[2] AP-HP, Pôle biologie-Pathologie Henri Mondor, Créteil, France; INSERM IMRB U955, Créteil, France

*Corresponding author

Key words: CYP26A1, AROMATASE, sex determination, ovary, rabbit, TALEN

Abstract

AROMATASE, encoded by the *CYP19A1* gene, is the cytochrome enzyme responsible for the

synthesis of estrogens in vertebrates. In most mammals a peak of expression of the *CYP19A1*

gene occurs in the fetal XX gonad when sexual differentiation starts up. To elucidate the role

of this peak, we produced 3 lines of TALEN genetically edited *CYP19A1* KO rabbits, that were

void of any production of estradiol. All KO XX rabbits developed as females, with aberrantly

small sized ovaries at adulthood, an almost empty reserve of primordial follicles and very few

large antrum follicles. Ovulation never occurred. Our histological, immunohistological and

transcriptomic analyses showed that the surge of estradiol in the XX fetal rabbit gonad is

dispensable for its determination as an ovary, or for meiosis. However, it is mandatory for the

high proliferation and differentiation of both somatic and germ cells, and consequently for

the establishment of the ovarian reserve.

In Vertebrates, AROMATASE is the unique enzyme responsible for the synthesis of estrogens.

It converts specifically three androgenic substrates (androstenedione, testosterone and

17alpha-testosterone) to estrone, estradiol, and estriol respectively. It is the product of the

CYP19A1 gene, which is the unique member of its gene family. Although the *CYP19A1* gene is

commonly known to be mainly expressed by gonads in adults, a physiologically significant

expression has already been reported in numerous tissues or organs such as bones, adipose

tissue, brain or placenta (the expression in the latter depending on the species, see for review

in (Conley and Hinshelwood 2001) and references herein).

In mammals, the expression of the *CYP19A1* gene begins early at around implantation in the

embryo in rabbits (Dickmann *et al.* 1975; George and Wilson 1978b), pigs and ruminants

(Gadsby *et al.* 1980). A second burst of expression was reported later in fetal life in the ovaries

shortly after the sex determination process. This was described in rabbits (George *et al.* 1978;

George and Wilson 1978b; Gondos *et al.* 1983; Daniel-Carlier *et al.* 2013), sheep (Mauleon *et*

al. 1977; Payen *et al.* 1996; Torley *et al.* 2011), goats (Pannetier *et al.* 2006), bovines (Shemesh

1980; Ross *et al.* 2009; Garverick *et al.* 2010) and humans (George and Wilson 1978a) by RNA

measurements or by estrogen assays in gonads or gonadal culture media. This peak is

transitory and spans until meiosis initiation. Later on, *CYP19A1* gene expression increases

progressively with follicle differentiation. It culminates in antrum follicles, the mural granulosa

cells being the site of expression. By the past, the physiological significance of this fetal peak

of expression has been searched in the rabbit species. Namely, George et al (George *et al.*

1978) have supposed that the local production of estrogens in the XX gonad was a key event

for the initiation of the differentiation of the female fetal gonad but this hypothesis has still

not been proven. In birds (Elbrecht and Smith 1992; Vaillant *et al.* 2001) and fishes (Guiguen

et al. 2010), it was shown that estrogens are major determinants of the ovary orientation of

the undifferentiated gonad in fetal or larval life. The question was thus to investigate the role

of the fetal peak of estrogen in the mammalian gonad and more specifically to search whether

in mammals it is mandatory for ovary differentiation.

To reach this aim, we decided to suppress the expression of the *CYP19A1* gene in a mammal

species other than the mice. Indeed, rodents represent some of the few mammalian species

without detectable peak of expression at this developmental stage in the fetal XX gonad i.e.

before meiosis. Thus, thanks to the TALEN methodology already shown to be efficient for gene

targeting in the rabbit species (Lee *et al.* 2018; Peyny *et al.* 2020), we produced *CYP19A1*

knocked-out (KO) rabbits completely devoid of estrogens secretion. Here we report the

phenotype of the XX *CYP19A1* KO (*ARO^-/-^*) female rabbits, which are completely infertile at

adulthood due to ovarian developmental defects detectable as soon as the first stages of their

differentiation.

Results

Three aromatase mutant rabbit lines which do not produce estradiol were generated.

The TALEN subunits were designed to generate *InDel* mutations through non-homologous end

joining near the initiation site of translation of the *CYP19A1* gene as described in Figure 1.

After transferring 67 injected embryos in 3 recipient female rabbits, 6 founders (2 males and

4 females) harboring mutant alleles were characterized from 11 neonates. Each founder F0

was mated with a wild type counterpart in order to transmit a mutant allele to the

descendants and generate mutant lines carrying one mutated allele per line.

Three mutant lines harboring each one a specific mutant allele were generated and studied.

Homozygous (*ARO^-/-^*) rabbits were generated by crossing heterozygous (*ARO^+/-^*) rabbits.

Mating was performed between rabbits within each line in order to avoid mixing the various

mutant alleles.

Mutations are shown in Figure 1. A 339, 498 or 829 nucleotide-long (nt) deletion was observed

in each of the alleles respectively named Δ339, Δ498 and Δ829. Insertion of a small number

of nucleotides was also detected in Δ339 and Δ498 alleles. Briefly, mutations Δ339 and Δ498

eliminated the 3' end of exon II including the translation initiation site (ATG codon) plus a large

part of the second intron; the mutation Δ829 eliminated a larger fragment encompassing the

proximal promoter upstream of exon II, the whole exon II and a large part of the second intron.

Thus, in the absence of the translation initiation site, no protein was expected to be translated

from any of these mutant alleles. Moreover, the deletion of the 3' part of the exon II

theoretically should lead to the elimination of 48 amino acids in the N-terminus of the

AROMATASE enzyme. As previously published (Kaur and Bose 2014), these amino acids

encompass the signal anchor responsible for the translocation of the protein to the

endoplasmic reticulum. It is thus expected that if a protein is translated from the mutant

mRNA, it has a good chance to be not correctly embedded in the endoplasmic reticulum

bilayer, and consequently not functional.

Figure 1: TALEN induced indel mutations in the rabbit *CYP19A1* gene. (A): structure of the rabbit aromatase gene; **(B):** sequences of the mutant alleles Δ339, Δ498 and Δ829

Figure 1: TALEN induced indel mutations in the rabbit *CYP19A1* gene.

(A): structure of the rabbit aromatase gene, with the various tissue specific promoters i.e. the I.1 placental promoter at around 80kb upstream of the transcription start point (tsp, marked by squared arrows) and the two proximal ovarian promoters I.r and II as previously described (Bouraïma *et al.* 2001). The enlargement shows the sequence of the exon II with the initiation site of translation (ATG in green letters). The two subunits (left and right arm) of the TALEN are shown as green symbols and the targeted sequence is written in red. The horizontal black arrows point the primers used for PCR detection of mutants in founders (F0/R0 set) and for routine qPCR genotyping (F1/R1 set).

(B): sequences of the mutant alleles Δ339, Δ498 and Δ829. Capital letters report to exon sequence, lowercase letters to intron sequence. The deletion spans over 339 nucleotides (nt) in the Δ339 mutant allele (149 nt in the 3' end of exon II and 190 nt of intron 2) and over 498 nt in the Δ498 mutant allele (148 nt in the 3' end of exon II and 350 nt of intron 2), both with insertion of few nt at the repair position. In the Δ829 allele, the mutation consisted in the elimination of 829 nt encompassing 250 nt upstream of the beginning of exon II (transcription

start site of the gene as regard to the ovarian promoter), the totality of the second exon (263 nt) and 316 nt in the second intron. All mutations suppressed the ATG codon and the splice donor site at the 3' extremity of exon II.

Finally, to confirm the absence of any active AROMATASE enzyme, we searched for estradiol by Gas Chromatography / Mass Spectrometry (GC/MS) in rabbit serum samples collected after birth and in fetal gonads. The concentrations of estradiol in sera from $ARO^{-/-}$ XX females after birth were under the limit of quantification (LOQ = 0,2pg/ml) whatever the age of the animals, when in wild type ($ARO^{+/+}$) and heterozygous ($ARO^{+/-}$) females the estradiol concentration increased progressively with the age of the animal (supplementary Figure 1). The concentration of testosterone was measured in the same samples. In $ARO^{-/-}$ sera, the values ranged in similar way than in WT or $ARO^{+/-}$ XX rabbits.

We measured also by GC/MS the steroid content of the gonads in fetuses aged of 28 days *post coitum* (d*pc*). The estradiol content was undetectable in gonads from all $ARO^{-/-}$ XX fetuses and in all XY gonads, when it was always detected, even sometimes in very low amounts, in XX gonads from $ARO^{+/-}$ and WT genotypes (supplementary figure 2). Testosterone was not detected in any XX gonads whatever their genotype at the *CYP19A1* locus. This shows that androgens were not over-synthesized nor accumulated in XX $ARO^{-/-}$ gonads. Finally, these data also show that neither estradiol nor testosterone from maternal or placental origin was detected in the 28d*pc* fetal gonads.

XY and XX $ARO^{+/-}$ rabbits were fertile, with normal genital tracts and normal gonads morphology

In the three mutant lines, all $ARO^{+/-}$ and XY $ARO^{-/-}$ rabbits were viable until adults and did not suffer apparently from any metabolic or postural disease at least until 2 years after birth. No apparent modification of external genitalia was visible. All were fertile and gave birth to progeny. The sexual behavior was not studied. The number of neonates per litter from heterozygous mating (♀ $ARO^{+/-}$ x ♂ $ARO^{+/-}$) was similar to that of wild type mating (♀ WT x ♂ WT) (supplementary Figure 3). The percentage of mating without pregnancy was not different: 37,9% (11 null per 29 mating) in WT x WT mating and 35,3% (23 null per 65 mating) in heterozygous x heterozygous mating. All this suggests that the fertility of heterozygous animals was not affected.

At puberty (5-6 months after birth), genital tracts and gonads were not different in

heterozygous and wild type XX rabbits (supplementary Figure 4). Since all measured

parameters did not differ in heterozygous and wild type females (similar estradiol and

testosterone serum levels, similar prolificacy, similar overall structure of genital tracts and

ovaries), we used ovary samples indifferently from wild type or heterozygous females as

reference ones. Nevertheless, the genotype has always been distinctly reported in the legends

of all figures.

152　XX $ARO^{-/-}$ rabbits developed small female genital tracts, ovaries with almost no follicular

153　reserve and very few antral follicles.

All XX $ARO^{-/-}$ rabbits were viable without any metabolic or postural disease at least until 1.5

years after birth. No apparent modification of external genitalia was visible. The sexual

behavior of XX $ARO^{-/-}$ rabbits was not studied in detail.

Obvious modifications of the genital tracts and ovaries were observed in XX $ARO^{-/-}$ rabbits

from the three lines with similar characteristics. After puberty, a typical phenotype was

observed, with under-developed uterus horns, oviducts and pavilions resembling those of

neonatal females. Their size was clearly smaller than in wild type or heterozygous females

(supplementary Figure 4). In all three lines, XX $ARO^{-/-}$ gonads developed like ovaries (Figure 2,

supplementary Figure 5) but with a very small size. As the phenotype was identical in Δ339,

Δ498, and Δ829 lines, it is hardly attributable to any off-target alteration of the genome.

Ovaries contained almost no primordial follicles, scarce primary and secondary follicles, and

were filled with numerous remnants of atretic follicles. The abundant follicle reserve

composed of primordial follicles observed at the cortex of wild type ovaries (Figure 2A and B)

was absent in $ARO^{-/-}$ gonad mutants (Figure 2C and D). Nevertheless, some follicles developed

to reach the large antrum stage in $ARO^{-/-}$ ovaries, showing granulosa and theca cell layers,

oocyte and cumulus. The number of large antrum follicles was high in some females (see the

ovary of the Δ829 female in supplementary Figure 5) possibly reflecting the surge of

gonadropins that occurs at early puberty (around 5-6 months in the rabbit). Numerous Call-

Exner bodies (spherical deposits of basal lamina matrix surrounded by granulosa cells,

observed in rabbits and some mammals) were observed in the granulosa of growing follicles

in $ARO^{-/-}$ ovaries as classically observed in wild type ones (Lee et al. 1996). The number of

pyknotic cells or of cells with dense colored nuclei was low.

Figure 2: ovaries of adult *ARO⁻ᐟ⁻* rabbits were small with few follicles and filled with collagen rich connective tissue. Ovaries from wild type female rabbits (*ARO⁺ᐟ⁺*, A and B) and from the mutant *ARO⁻ᐟ⁻* line Δ339 (C and D) were fixed in PAF then HES colored. Females were 5,5 months old.

Figure 2: ovaries of adult *ARO⁻ᐟ⁻* rabbits were small with few follicles and filled with collagen rich connective tissue.

Ovaries from wild type female rabbits (*ARO⁺ᐟ⁺*, A and B) and from the mutant *ARO⁻ᐟ⁻* line Δ339 (C and D) were fixed in PAF then HES colored. Females were 5,5 months old. Two distinct sections of the same ovary are presented for the mutant *ARO⁻ᐟ⁻* to show the follicles that were otherwise not visible. Enlarged zones point the

different types of follicles. Black arrows = Call-Exner bodies. Black stars = accumulation of fibrous tissue. Red stars = remnants of degenerated follicles. "p" = primordial follicle; I = primary follicle; II = secondary follicle; IC = interstitial cells. The phenotype of the $ARO^{-/-}$ ovary was similar in the other lines Δ498 and Δ829 (supplementary Figure 5).

The expression of the marker genes *RSPO2* and *FOXL2* specific respectively for the oocyte and for the granulosa cells in differentiating follicles was studied by *in situ* hybridization (ISH, Figure 3). In the $ARO^{-/-}$ ovary, the rare detected oocytes were positive for the *RSPO2* labelling (Figure 3A), and granulosa cells were positive for *FOXL2* (Figure 3B). In wild type ovaries, AROMATASE was immuno-detected in a small number of follicles with antrum, thus characterizing the pre-ovulatory ones (Figure 3C). The Anti Müllerian Hormone (AMH) was immuno-detected in granulosa cells of many follicles in wild type ovaries except those positive for AROMATASE. In $ARO^{-/-}$ ovaries, granulosa cells of all follicles were AROMATASE negative and AMH positive (Figure 3D). However, the AMH serum level was lower in $ARO^{-/-}$ females, reflecting the small number of follicles and/or their poor synthesis capacity.

Figure 3: *RSPO2, FOXL2* and AMH were detected in wild type and ARO$^{-/-}$ adult ovaries. In Situ Hybridization to localize mRNA of *RSPO2* and *FOXL2* genes and immunodetection of AROMATASE and AMH in adult ovaries. Ovaries were collected in wild type (WT) or *ARO*$^{-/-}$ Δ329 females aged 5-6 months.

Figure 3: *RSPO2, FOXL2* and AMH were detected in wild type and ARO$^{-/-}$ adult ovaries. In Situ Hybridization to localize mRNA of *RSPO2* and *FOXL2* genes and immunodetection of AROMATASE and AMH in adult ovaries. Ovaries were collected in wild type (WT) or *ARO*$^{-/-}$ Δ329 females aged 5-6 months. The expression of *RSPO2* (A) and *FOXL2* (B) genes was detected by ISH probes. AROMATASE (C) and AMH (D) were detected by immunohistochemistry on two adjacent sections. The *RSPO2* and *FOXL2* probes labelled respectively the

cytoplasm of oocytes (brown dots) and of granulosa cells (red dots) from all follicles, from primordial to antrum stages in wild type and *ARO*$^{-/-}$ rabbits as well. Some thecal cells were also positive for the *FOXL2* labelling in both genotypes. Mural granulosa cells from pre-ovulatory follicles were positively labelled by the anti-AROMATASE antibody in wild type ovaries (brown colored); no cells were positive in *ARO*$^{-/-}$ ovaries. The anti-AMH antibody labelled most granulosa cells of growing follicles in wild type ovaries (brown colored), excepted those positive for AROMATASE. In *ARO*$^{-/-}$ ovaries, all follicles with antrum were positively labelled. Similar staining were observed in females from the two other strains. The graph represents the AMH serum level in *WT* (black points), *ARO*$^{+/-}$ (green points) and *ARO*$^{-/-}$ (red points) females.

Intense modifications affected the medulla of the ovary. In wild type rabbits, the medulla consisted mostly in interstitial tissue composed of cells with large clear cytoplasm and round regular shape nuclei (Figure 2B); besides, in *ARO*$^{-/-}$ ovary, the interstitial tissue appeared as a mixture of cells with heterogeneous shape and angular nuclei. More, several areas filled with fibrous material rich in collagen and devoid of cells were detected (Figure 2C and D; supplementary Figure 5).

Some of the heterozygous (*ARO*$^{+/-}$) and homozygous (*ARO*$^{-/-}$) 10 months old XX rabbits were hormonally treated to induce ovulation, mated and sacrificed to collect gonads for histological analysis. Ovulation rupture points were observed in ovaries from *ARO*$^{+/-}$ rabbits, but not in *ARO*$^{-/-}$ ovaries (supplementary Figure 6). While several primordial, primary, pre-antral and antral follicles were detected at the cortical periphery of ovaries in *ARO*$^{+/-}$ rabbits after super-ovulation treatment, no primordial or primary follicles were seen in the treated *ARO*$^{-/-}$ ovarian cortex. Nevertheless, some large antrum type follicles were observed in *ARO*$^{-/-}$ ovaries, with developed granulosa, theca cells, and oocyte.

The mutation impacted the XX gonad from the first stages of their differentiation

Our goal was to search when and how ovaries were impacted by the lack of functional AROMATASE enzyme and of estradiol during fetal life. In the rabbit species, pregnancy spans for 31 days. The present study analyzed the fetal gonads from 18 d*pc* onwards. At this stage in rabbits, the gonadic sex is already determined, the multiplication of germ cells and somatic cells is intense (Chretien 1966) and the peak of *CYP19A1* gene expression starts in the ovary, concomitantly with the increase of *FOXL2* gene expression (Daniel-Carlier *et al.* 2013). At 20 d*pc*, we detected a specific labelling with a *CYP19A1* ISH probe in cells of the surface epithelium and cells forming invaginated structures issued from this epithelium

(supplementary Figure 7). Simultaneously, we observed that these same cells were labelled

by an *ESR1* specific ISH probe and we detected a surge of the expression of the estrogen

receptor alpha gene (*ESR1*) by RT-qPCR (supplementary Figure 7). The *FOXL2* gene was mainly

expressed in somatic cells underneath the surface epithelium and those surrounding

invaginated structures (ISH labelling, supplementary Figure 7). Only few cells of the surface

epithelium were clearly positive for both the *CYP19A1* and *FOXL2* ISH probes. The expression

of the *ESR2* gene coding for the estrogen receptor beta was low at these stages compared

with the levels determined later at birth (34 d*pc*). In agreement, a faint labelling was observed

by ISH labelling (supplementary Figure 7).

Therefore, the morphology of the gonads and the location of specific markers were

investigated at various fetal stages and compared between wild type or heterozygous gonads

versus ARO$^{-/-}$ ones from the stage of gonadal determination onwards. At 18 d*pc*, the gonads

from all XX fetuses, *ARO*$^{-/-}$ and *ARO*$^{+/-}$, were differentiated as ovaries with the same

morphology except a thinner coelomic epithelium in *ARO*$^{-/-}$ gonads (Figure 4A). Regardless of

the developmental stages, XX *ARO*$^{-/-}$ gonads showed no evidence of testicular appearance (see

for comparison the 18 d*pc* testis in the supplementary Figure 8). A similar density of OCT4

positive germ cells was detected in *ARO*$^{-/-}$ and *ARO*$^{+/-}$ gonads, as shown by the labelling with

specific antibodies directed against OCT4 (Figure 4B). Mitotic activity was detected by KI67

immunostaining (Figure 4C). KI67 positive cells were mainly present in the ovarian surface

facing the abdominal cavity and in invaginated structures below in both *ARO*$^{+/-}$ and *ARO*$^{-/-}$

ovaries. Interestingly, the KI67-positive ovarian surface appeared thinner in KO gonads than

in the *ARO*$^{+/-}$ controls (Figure 4C).

Figure 4: morphology and detection of OCT4 and KI67 in ovaries from *ARO⁺ᐟ⁻* and *ARO⁻ᐟ⁻* 18 d*pc* old fetuses.

Figure 4: morphology and detection of OCT4 and KI67 in ovaries from *ARO⁺ᐟ⁻* and *ARO⁻ᐟ⁻* 18 d*pc* old fetuses.

Gonads were collected at 18 d*pc* in *ARO⁺ᐟ⁻* and *ARO⁻ᐟ⁻* fetuses, fixed in Bouin's then HES colored (A) or fixed in

PAF then treated for antibody labelling (B and C). **(A): HES coloration**. The surface epithelium appeared as a

continuous layer of epithelial cells. Red stars point dense nuclei that could correspond to pyknotic nuclei. Arrows

point figures of mitoses. **(B): detection of OCT4 positive cells.** The immune-labelling was visualized using a

fluorescent secondary antibody. Nuclei of positive cells (large round green-labelled) were dispersed through the
gonad in both *ARO$^{+/-}$* and *ARO$^{-/-}$* ovaries. **(C): detection of KI67 positive cells.** The immune-labelling was visualized
using a peroxidase coupled secondary antibody. Positive cells with brown colored nuclei were found at the
surface epithelium, and in rows of cells delimiting cell clusters. Green stars point brown light-colored large nuclei,
possibly corresponding to germ cells.

From 20 d*pc* onwards, more important alterations progressively appeared in *ARO$^{-/-}$* ovaries.
The cortex was thinner, composed of a single layer of epithelial cells, connected to small size
nascent ovigerous cords, by opposition to the thick cortex observed in *ARO$^{+/-}$* ovaries,
connected to numerous large size ovigerous cords (Figure 5A and 5D). A discontinuous layer
of cells with small and dense colored nuclei was observed underneath the surface epithelium,
forming at 22 d*pc* (Figure 5D) and later on (34 d*pc*, Figure 9) a loose connective tissue between
the thin cortex and the inside of the ovary.
The cells of the surface epithelium expressed the *ESR1* gene in both genotypes (Figures 5B and
5E). The intensity of the ISH *ESR1* labelling varied greatly at the surface of the *ARO$^{+/-}$* ovaries
as shown in figure 5 (compare 5B, E, C and F, which are different sections from the same
gonad). Nevertheless, in all treated sections, clusters of labelled *ESR1* cells connected to the
surface epithelium were easily observed. Besides, in *ARO$^{-/-}$* ovaries, labelled *ESR1* cells were
mainly dispersed in the inner of the gonad. Similarly, the *FOXL2* ISH probe labelled evenly
dispersed cells in the inner of the *ARO$^{-/-}$* gonad, which differed from the strong *FOXL2* labelled
cell cords of the *ARO$^{+/-}$* ovary. *OCT4* positive germ cells were detected in both genotypes but
the hybridization signals were lighter in *ARO$^{-/-}$* germ cells (Figures 5C and 5F-H). Numerous
were found in the surface epithelium, interestingly in close contact with somatic cells
expressing the *ESR1* gene.

Figure 5: gradual modifications of *ARO⁻ᐟ⁻* ovaries at 20 and 22 d*pc*. *ESR1*, *FOXL2* and *OCT4* gene expressions were labelled by dual HIS.

Figure 5: gradual modifications of *ARO⁻ᐟ⁻* ovaries at 20 and 22 d*pc*.

Gonads were collected from *ARO⁺ᐟ⁻* and *ARO⁻ᐟ⁻* fetuses, fixed in Bouin's then HES colored **(A, D).** Ovigerous nests

and cords are underlined by a dotted white line in **A-F**. The red double arrow points the thickness of the nascent

cortex. Blue stars point the connective tissue. In situ hybridization was performed to visualize simultaneously

two RNA targets to show the *ESR1* (blue points) and *FOXL2* (fast red points) RNAs **(B, E)**, and the *ESR1* (blue) and

OCT4 (fast red) RNAs **(C, F)**. The lower panel is the fluorescence observation of the fast red *OCT4* ISH labelling to

show the overall density of the *OCT4* positive cells (**G, H**).

KI67 positive cells were overall detected in both types of ovaries at 22 d*pc* (Figure 6A) but

there was still a clear difference of mitotic activity in the coelomic epithelium between *ARO⁺ᐟ⁻*

and *ARO⁻ᐟ⁻* ovaries. Indeed, while many KI67 positive cell layers were detected in the coelomic

epithelium of *ARO⁺ᐟ⁻* gonads, only one layer was visible in the *ARO⁻ᐟ⁻* one. The γH2AX-antibody,

marker of double strand breaks of the DNA, labelled mostly germ cells in both types of ovaries

as suggested by their large round nuclei (Figure 6B). The density of γH2AX positive cells (probably apoptotic cells) was apparently not different in *ARO*^{+/-} and *ARO*^{-/-} ovaries.

Figure 6: mitotic activity and DNA fragmentation at 22 d*pc*.
The marker of mitotic activity KI67 **(A)**, and that of double strand breaks gH2AX **(B)** were localized by immunohistochemistry in sections from PAF fixed ovaries of *ARO*^{+/-} and *ARO*^{-/-} fetuses at 22 d*pc*.

Figure 6: mitotic activity and DNA fragmentation at 22 d*pc*.

The marker of mitotic activity KI67 **(A)**, and that of double strand breaks γH2AX **(B)** were localized by immunohistochemistry in sections from PAF fixed ovaries of *ARO*^{+/-} and *ARO*^{-/-} fetuses at 22 d*pc*. The KI67 antibody labelled nuclei of most cells of the coelomic epithelium, and numbers of cells inside the ovary in both genotypes. Cells with flat nuclei KI67 positive (red arrowheads) surrounded from place to place the ovigerous cords. Cells with large round nuclei positive for KI67 labelling (probably germ cells) were found mainly at the inside of the cords (green stars). The γH2AX antibody labelled large round shape nuclei corresponding probably to germ cells. Note the disconnection between the coelomic epithelium and the new forming ovigerous cords in *ARO*^{-/-} ovaries.

The relative expression levels of key marker genes were determined at these developmental stages on total RNAs extracted from the whole gonads (Figure 7). The expression levels of the *OCT4* and *FOXL2* genes were reduced in *ARO*^{-/-} ovaries, reflecting a reduction of the number of *OCT4* positive germ cells and *FOXL2* positive somatic cells respectively. Interestingly, the expression level of the *DDX4* gene, known to be its lowest at this ovarian developmental stage (Daniel-Carlier *et al.* 2013), did not differ between WT and *ARO*^{-/-}. It increased between 20 and 22 d*pc*, which reflects the early differentiation of germ cells from the pluripotent stage. The

SOX9 gene expression level was 5- to 10 times lower in the ovaries from all genotypes than in

testes at the same age. The *SOX9* gene was not detected by immunohistology either in wild

type or in *ARO^-/-* ovaries, while a clear labeling was observed in the fetal testis at 22 d*pc*

(supplementary Figure 8). This reinforces the demonstration that *ARO^-/-* ovaries did not show

any sex-reversal phenotype. The expression levels of the *ESR1* and *ESR2* genes did not change

significantly.

Figure 7: gene expression in 20 and 22 d*pc* ovaries from WT and *ARO^-/-* fetuses.
Ovaries were collected from wild type and mutant fetuses. The diagrams show the relative mRNA expression levels.

Figure 7: gene expression in 20 and 22 d*pc* ovaries from WT and *ARO^-/-* fetuses.

Ovaries were collected from wild type and mutant fetuses. The diagrams show the relative mRNA expression

levels. Each point represents one RNA sample extracted from both gonads of one animal at the indicated

developmental stage. The same RNA samples were analyzed in the 6 diagrams. Red triangles refer to mutant

from Δ339 line, and red point from Δ829 line. No samples from line Δ498 were analyzed. The blue point in the

SOX9 diagram shows the *SOX9* mRNA level measured in the two testes of one 20 d*pc* WT fetus. Horizontal bars

represent the medians. At these stages *DDX4* and *ESR2* gene expression levels were very low, but significantly

measured.

Thus, from 20 d*pc* onwards, the differentiation of the *ARO*$^{-/-}$ gonad was profoundly modified,

with a decrease of proliferation of the somatic cells of the coelomic epithelium and a reduction

of the number of germ cells. Finally this leads to a clearly underdeveloped small organ with

small ovigerous nests, an abnormal distribution and differentiation of the *FOXL2* positive cells

and the spread of connective tissue underneath the surface epithelium.

Impact of estrogens deprivation on germ cell meiosis and follicular differentiation

In the rabbit ovary, the first signs of meiosis are histologically detected at 30-31 d*pc,* i.e.

around birth. Later on, typical meiotic germ cells (prophase I) are observed for less than 2

weeks. After meiosis arrest at diplotene stage, follicles begin to differentiate. The peak of

expression of the *STRA8* gene that starts at 24-28 d*pc* (Daniel-Carlier *et al.* 2013) marks the

commitment of germ cells to meiosis. Thus, we sought to analyze at 28 d*pc* (meiosis

commitment), 34 d*pc* (beginning of meiosis) and few days or weeks after birth

(folliculogenesis) whether initiation and arrest of meiosis I and early follicular formation were

impacted in *ARO*$^{-/-}$ ovaries.

In wild type and *ARO*$^{-/-}$ 28 d*pc* ovaries, numerous germ cells were positively labelled by the

OCT4, *DDX4* and *STRA8* ISH specific probes (Figures 8B, 8C, 8D). The *STRA8* labelling proves

the overall commitment of germ cells in meiosis in WT and *ARO*$^{-/-}$ ovaries. However, in these

later, the number of positive germ cells for the three probes was low. More, they were

clustered in small ovigerous-like structures, while they were all observed in large ovigerous

cords in wild type ovaries. Positive *FOXL2* labelled cells were also much less numerous in *ARO*$^{-/-}$

$^{/-}$ ovaries, as shown by the few numbers of hybridization labels. Nevertheless, some of these

FOXL2 positive cells harbored a flattened shape and surrounded the small ovigerous-like

structures embedding germ cells, as in wild type ovaries (Figure 8E).

Figure 8: germ cells were committed to meiosis in WT and *ARO*^(-/-) ovaries from 28 d*pc* old fetuses.

365 **Figure 8: germ cells were committed to meiosis in WT and *ARO*^(-/-) ovaries from 28 d*pc* old fetuses.**

366 Ovaries collected at 28 d*pc* were fixed in PAF, then HES colored or analyzed by ISH. **(A): morphology of the**

367 **ovaries**. The pictures show the difference of the overall size of the gonad and the different thickness of the

cortex. The *OCT4* **(B)** and *DDX4* **(C)** ISH probes labelled respectively pluripotent germ cells and differentiating ones. The *STRA8* ISH probe **(D)** labelled germ cells committed to meiosis. The *FOXL2* ISH probe **(E)** labelled the differentiating somatic cells. Note that the latter were mainly distributed around and inside the ovigerous cords in the WT ovaries, and in contrast were dispersed throughout the gonad in the *ARO⁻/⁻* rabbit. As in all ISH pictures, positive labeling appears as colored dots.

At 34 d*pc*, in wild type ovaries, most ovigerous cords were filled with numerous zygotene germ cells, thus showing that meiosis had begun in a majority of germ cells (Figure 9A and a1). In contrast, in the *ARO⁻/⁻* ovaries, germ cells at zygotene were sparse (Figure 9B and b1, b2). The differentiation of the *ARO⁻/⁻* ovaries was further altered as shown by the low staining observed with *DDX4*, *STRA8, SPO11* and *FOXL2* probes and by the qPCR low expression of these genes (figure 9 C-F). The ratios of the *STRA8/DDX4* and *SPO11/DDX4* mRNA levels were lower in *ARO⁻/⁻* ovaries, suggesting that germ cells expressing *STRA8* and *SPO11* gene expression, and thus undergoing meiosis, were less abundant in *ARO⁻/⁻* ovaries.

Figure 9: meiosis in WT and *ARO⁻/⁻* ovaries. Ovaries were collected at 34 d*pc* (3-4 days after birth)

Figure 9: meiosis in WT and *ARO⁻/⁻* ovaries. Ovaries collected at 34 d*pc* (3-4 days after birth) were fixed in Bouin's then HES colored **(A-B)** or PAF then ISH treated **(C-F)**. The stages of meiosis were identified as previously published (Peters *et al.* 1965). **(A):** In wild type ovaries, large ovigerous cords were filled with numerous germ

cells in zygotene (a1, red arrows). Few pachytene stages were observed at the inner part of the cords (a1, blue arrows). **(B):** In *ARO$^{-/-}$* ovaries, the rare ovigerous cords were filled with a few numbers of germ cells in zygotene (b1). As in wild type ovaries, some germ cells in pachytene were found at the inner part of the cords (b2). In both ovaries, aberrant pictures of cell nuclei were characterized by punctiform or condensed and dark colored chromatin (red stars). Blue stars show the connective tissue already observed mostly in *ARO$^{-/-}$* ovaries. **(C-F):** ISH was performed using *DDX4* (**C**), *FOXL2* (**D**), *STRA8* (**E**) and *SPO11* (**F**) genes specific probes. In wild type ovaries, numerous germ cells positive for *DDX4*, *STRA8* and *SPO11* filled the ovigerous cords; pre-granulosa cells *FOXL2* positive were found around and inside the cords. In *ARO$^{-/-}$* ovaries, germ cells positive for these same ISH probes were also found but much less numerous; the *FOXL2* labelling was very faint. The diagrams show the results of a quantitative analysis of gene expression. Values are relative expressions measured using total RNA extracted from the two gonads of each animal from wild type (WT), *ARO$^{+/-}$* (+/-) *and ARO$^{-/-}$* (-/-) rabbits; the same RNA samples were analyzed in the four diagrams. Horizontal bars represent the medians. The STRA8/DDX4 and SPO11/DDX4 diagrams represent the ratio between the expression of the genes for each sample. Stars represent the statistical significance (p<0.01).

These differences intensified further (supplementary Figure 9). As meiosis progressed (10 days *post partum*, d*pp*), zygotene, pachytene and diplotene stages were simultaneously observed in wild type ovaries. At the interface between the cortex and the medulla, the large ovigerous cords broke down, forming numerous early primordial follicles encompassing diplotene stage germ cells surrounded by a ring of flat pre-granulosa cells. This was far to be observed in *ARO$^{-/-}$* ovaries. Indeed, at 7 days after birth, almost no germ cells in zygotene or pachytene stages were detected; a small number of diplotene-like stage cells were nonetheless detected, some surrounded by a discontinuous ring of somatic cells that could be assumed to be pre-granulosa cells.

Later on at 16-19 d*pp*, the gonad size was approximately twice as small in *ARO$^{-/-}$* rabbits, harboring a thin cortex with a small number of primordial follicles (Figure 10A). All ovigerous cords broke down both in wild type and *ARO$^{-/-}$* ovaries and follicles were isolated. However, in *ARO$^{-/-}$* ovaries, the ring of pre-granulosa cells surrounding the oocytes was frequently discontinuous and rare well-developed primordial follicles were detected. The *FOXL2* ISH labelling was very weak reflecting a low level of well-differentiated granulosa cells (Figure 10B). Nevertheless, oocytes of primordial follicles were positive for the *RSPO2* ISH specific probe in *ARO$^{-/-}$* follicles, as in wild type ones (Figure 10C).

Figure 10: early follicle differentiation. Ovaries were collected at 16 or 19 d*pp* then treated with Bouin's or PAF

fixative and processed for HES staining or ISH labelling respectively. **(A): HES staining of 19 d*pp* sections**. Blue

stars point the abnormal persistence of connective tissue, void of differentiating oocytes or follicles, at the outer

part of the cortex and in the medulla of *ARO*−/− ovaries. The green arrows point some of the primordial follicles in

formation with a discontinuous wall of granulosa cells. Early formed primordial follicles surrounded by a wall of

regular shape granulosa cells are found in wild type and *ARO*−/− ovaries (white arrows). **(B) and (C): ISH labelling**

using *FOXL2* and *RSPO2* gene specific probes. The *FOXL2* and *RSPO2* probes labelled granulosa cells and oocytes

respectively in both wild type and *ARO*−/− ovaries. Note the huge decrease of *FOXL2* positive cells in *ARO*−/− ovaries

compared to the WT control ones.

To conclude, in the *ARO*−/− ovaries, a small number of oocytes had differentiated after birth;

these oocytes were surrounded by a ring of granulosa cells. Despite the probably abnormal

differentiation of the latter (shown by the weak expression of the *FOXL2* gene), a small

population of primordial follicles could differentiate. At the same time, the ovary was invaded

by the connective tissue filled with fibers rich in collagen as previously observed from 20 d*pc*

onwards; this latter was not replaced by the ovigerous nests as normally occurs in non-mutant

ovaries. The *ARO*−/− ovaries may have failed to fill-in this connective tissue due to the disturbed

formation of ovigerous nests.

Discussion

The already long-standing demonstration of a peak of *CYP19A1* gene expression, AROMATASE activity and estradiol production by the fetal ovary in several mammal species between the stage of sex determination and the initiation of meiosis has raised questions on the role of locally produced estrogens on the fetal differentiation of this organ. The mouse model was for a long time the sole available model to study the function of genes *in vivo*. So, strains of *Cyp19a1* gene or *estrogen-receptor* genes KO mice have been created and a series of papers have reported their phenotype (Fisher *et al.* 1998; Couse *et al.* 1999; Britt *et al.* 2000; Britt *et al.* 2001; Findlay *et al.* 2001; Hamilton *et al.* 2014). All XX mice developed ovaries, but follicle differentiation was abnormal after birth. However, this model could not answer to our question since the mouse is one species where *Cyp19a1* gene expression is detected after meiosis initiation in the "late" fetal ovary (Dutta *et al.* 2014) and not before (Greco and Payne 1994). Thanks to the recent possibility to target genome modifications in numerous other species, we created a rabbit model harboring a mutation in the *CYP19A1* gene (*ARO$^{-/-}$*), a species with a peak of *CYP19A1* gene expression in fetal ovary before meiosis. The *ARO$^{-/-}$* rabbits were totally void of any estrogen production. Particularly, the amount of estradiol in gonads of all *ARO$^{-/-}$* fetuses was undetectable, which eliminates the possibility of any compensatory mechanism through estradiol from maternal or placental origins. More, no testosterone was detected in the *ARO$^{-/-}$* fetal gonads, thus eliminating any possibility of interference due to the local accumulation of androgens in response to the lack of androgen aromatization. The key results of this study regarding XX *ARO$^{-/-}$* ovary phenotype were: 1) no XX sex reversal in *ARO$^{-/-}$* rabbits, 2) an abnormal differentiation of *FOXL2* positive cells, leading to the poor differentiation of ovigerous nests and the reduction of the number of meiotic germ cells and 3) a quasi-absence of follicular reserve, also linked to early folliculogenesis failure.

AROMATASE is not an ovarian determining enzyme in mammals

The question of the role of estrogens in sex determination in mammals was prompted by the finding that in goats, the *FOXL2* gene was a main actor for ovarian determination, since the *FOXL2* gene loss of function caused an ovary-to-testis sex reversal of the fetal gonad (Boulanger *et al.* 2014). Previously, FOXL2 was shown to be a crucial transcriptional activator

of the *CYP19A1* gene in goats (Pannetier *et al.* 2006) and fishes (Govoroun *et al.* 2004;
Pannetier *et al.* 2006; Guiguen *et al.* 2010), and possibly in chicken (Govoroun *et al.* 2004).
Thus, the question was raised to know whether FOXL2 could act on sex determination through
CYP19A1 gene transcriptional activation. This question was all the more important that in non-
mammalian vertebrate species, anti-aromatase drugs induced female to male sex-reversal
(see (Vaillant *et al.* 2001) in the chicken and (Yin *et al.* 2017) in fish). The rabbit was possibly
an interesting model since, as in goats, the surge of expression of the *FOXL2* gene in the fetal
ovary occurs when starts the peak of *CYP19A1* gene expression at the gonadal sex
determination (Diaz-Hernandez *et al.* 2008; Daniel-Carlier *et al.* 2013). The results of the
present study demonstrate that estrogens play no role on the female/male determinism of
the rabbit gonad. A similar finding had already been described in humans, where several cases
of total *CYP19A1* gene inactivation were reported (Akcurin *et al.* 2016; Zhu *et al.* 2016; Mazen
et al. 2017; Unal *et al.* 2018; Praveen *et al.* 2020). In spite of numerous symptoms affecting
the development of gonads, of external genitalia and of breast in the girl, with multiple
disorders in energy and bone metabolisms, there was no sex reversal of the human ovary as
regard to the genetic sex (Zhu *et al.* 2016). In contrast, opposite data were reported in the
mouse species, describing an increase in the serum testosterone levels in *Cyp19a1*[-/-] females
(Fisher *et al.* 1998) and the presence of seminiferous-like structures in the gonads of adult
Esr1/Esr2 double knock out females (Couse *et al.* 1999; Amano *et al.* 2017). However, this has
been controversial and not found by all authors.
490
Estrogens are involved in early ovarian development
The fetal ovarian differentiation is known to be intimately linked to mechanisms that involve
cellular interactions and cell migration, depending on autocrine and paracrine signaling
pathways. These processes involve genes coding for the components of the extracellular
matrix and cell-cell adhesion molecules. Estrogens have been reported as potent actors in
these mechanisms (Juengel *et al.* 2002; Gentilini *et al.* 2007; Wilhelm *et al.* 2007; Zalewski *et*
al. 2012; Hummitzsch *et al.* 2013). The reduced size of the gonads together with the reduced
number of somatic and germ cells is one of the major characteristics of the *ARO*[-/-] rabbit
ovaries. It was detected shortly after gonadal sex determination (18 d*pc*), and still clearly
visible around birth with a dramatic decrease of ovarian nests. We have also shown that the
estrogen receptor α gene (*ESR1*) was strongly expressed by the ovarian cortical cells

specifically by cells located within the coelomic epithelium (Figure 5B). This is worth to be pointed out because it resembles to what has been described in chicken. Indeed, in this species the female to male sex reversal induced by the AROMATASE inhibitor fadrozole induced a reduction of *FOXL2* expression (Hudson *et al.* 2005). More, the experimental down-regulation of the epithelial *Esrα* gene was sufficient to severely affect the ovarian cortex differentiation (Guioli *et al.* 2020), thus suggesting that the natural absence of *Esrα* expression in the cortex of the right ovary of the hen was at least partially responsible for the asymmetric degeneration of the right ovary (Guioli *et al.* 2014). We can therefore hypothesize that in rabbit *ARO$^{-/-}$* ovaries, the absence of estrogens can lead to a dramatic decrease of proliferative activity in the coelomic epithelium, evidenced by the KI67 staining (Figure 7A) and by the decrease of *FOXL2*-expressing somatic cells from 20-22 d*pc* (Figure 6).

Anyway, without estrogens, the proliferation of ovarian cells seemed to be affected but not abolished since some nests were clearly visible in *ARO$^{-/-}$* gonads at birth although under-developed. So, given that estrogens are involved in ovigerous nests formation, other factors are likely to play a role in this process as well. It has been shown in mice that the *Rspo1/Wnt4/β-catenin* pathway is involved in cell proliferation of the coelomic epithelium of both sexes during the gonadal switch (Chassot *et al.* 2012). This pathway could thus be responsible at least in part of the formation of the few ovigerous nests in *ARO$^{-/-}$* rabbit ovaries. Transcriptomics (even at the single cell level) and epigenetics studies are underway on WT rabbit ovaries in our team to search molecular targets of estrogens in early ovarian development by comparison with *ARO$^{-/-}$* gonads.

The number of germ cells was affected by the lack of estrogens.

The reduced number of germ cells is one of the major characteristics of the *ARO$^{-/-}$* rabbit ovaries. One of the first sign was the low expression level of the pluripotency marker gene *OCT4* and its reduced ISH labelling (see Figures 5 and 6). In mammals, germ cells colonize early the genital crests (Juengel *et al.* 2002; Wilhelm *et al.* 2007; Hummitzsch *et al.* 2013). In the rabbit species, it starts at 9 d*pc* and is over at around 18 d*pc* (Chretien 1966). Since the *CYP19A1* gene expression starts at 18 d*pc*, it is likely that the germ cell colonization occurs independently of *CYP19A1* expression. Thus, in the *ARO$^{-/-}$* ovary, in spite of the absence of estradiol, the initial stock of germ cells was probably not altered. In mammalian ovaries, once germ cells enter the genital ridges, they multiply intensely by mitosis until meiosis. Then, the

number of germ cells does not increase any more. Numerous articles have shown that in mammals estrogens stimulate the mitotic activity in the fetal ovary (see review in (Auersperg *et al.* 2001; Chou and Chen 2018)). During fetal life in rabbits, the mitotic activity of germ cells, which is maximum around 16-18 d*pc*, remains high up to 26 d*pc* (Chretien 1966), when the *CYP19A1* and *ESR1* genes are strongly expressed (Supplementary Figure 7). It can be suggested that the estradiol produced by AROMATASE in the fetal ovary could regulate the mitotic activity of germ cells. This hypothesis is supported by the fact that the germ cells, labelled at this stage by the *OCT4* HIS probe, were co-localized in the surface epithelium with cells expressing the *ESR1* gene. Germ cell death has previously been reported to occur continuously in the ovary from early fetal stages until puberty and later on (Hussein 2005; Myers *et al.* 2014). In *ARO*$^{-/-}$ ovaries, the number of γH2AX positive germ cells (highlighting DNA fragmentation, and at this stage dying germ cells) was similar to that in wild type ones (Figure 7). Thus, our data suggest that this is probably not germ cell death but rather the lack of germ cell divisions that was responsible for the low number of germ cells in the early *ARO*$^{-/-}$ ovary before meiosis.

Germ cells were committed in meiosis in spite of the lack of estrogens.

The commitment of germ cells in meiosis marks the first step of their differentiation as oocytes. In the rabbit species, it occurs at the third part of pregnancy (at around 28 d*pc*), and is characterized by the surge of the *STRA8* gene expression in germ cells specifically. Interestingly, in the *ARO*$^{-/-}$ ovaries, numerous germ cells were committed to meiosis as shown by the significant surge of *STRA8* gene expression; more, the various steps of the meiotic prophase were distinguishable, and the meiosis specific maker gene *SPO11* was significantly expressed as well. Thus, it suggests that nor the estradiol neither the AROMATASE enzyme were necessary for meiotic induction and progression. As discussed previously, the number of germ cells possible to be committed in meiosis was reduced in *ARO*$^{-/-}$ ovaries. More, it should be pointed out that the ratio of gene expression *STRA8/DDX4* and *SPO11/DDX4* was low in *ARO*$^{-/-}$ ovaries, showing that an abnormal loss of germ cells occurred at commitment or during meiosis. Thus, although estrogens are not mandatory for meiosis commitment, the fact remains that they are important for limiting meiotic germ cell losses.

Estrogens deprivation impacts follicle formation

Another key feature of the phenotype of the *ARO*[-/-] females concerns the abnormal genital
tract and ovarian follicle differentiation. At puberty, the genital tract of the XX *ARO*[-/-] rabbit
was profoundly altered, with massive atrophy, as already reported in *Cyp19a1*[-/-] mice (Fisher
et al. 1998; Britt *et al.* 2000), and in agreement with the fact already demonstrated that the
development of the female genital tract is highly dependent on estrogens. The follicle
differentiation that starts 12-15 days after birth in rabbit at the arrest of meiosis prophase is
a continuous process that involves the surrounding of oocytes with flat pre-granulosa cells
and the breakdown of ovigerous nests thus giving rise to primordial follicles. Then, the
differentiation of pre-granulosa cells from flat- to cuboidal-form cells signs the differentiation
of primary follicles. Of course, the size of the population of primordial follicles is linked to the
number of oocytes, the latter being the consequence of all the events which occurred during
fetal and neonatal life as discussed previously. In the *ARO*[-/-] rabbit ovary, this is of particular
importance since the pool of oocytes and that of *FOXL2* expressing pre-granulosa cells are
extremely reduced. More, it has been reported that the nests breakdown is an event
specifically sensitive to apoptotic losses (Myers *et al.* 2014). This could explain why the pool
of primordial follicles was very low in *ARO*[-/-] rabbit ovary few days after birth (Figure 10) and
almost null at puberty (Figure 2). The poor expression of the *FOXL2* marker gene by the
granulosa cells 16 days after birth signs the abnormal differentiation of the primary follicles,
the number of which was extremely low (Figure 2). Surprisingly, controversial results were
reported in *Cyp19a1* KO mice, where the number of primary follicles was significantly higher
than in wild type mice (Britt *et al.* 2000). To date, we have no explanation for such differences.
Furthermore, we observed that despite the absence of estradiol, large antral follicles with
oocytes developed in the *ARO*[-/-] ovary, however these follicles were very few. This proves that
estrogens are not the only actors in follicular growth and differentiation at puberty and
onwards. Many previous studies have shown that these events depend on the interactions
between several signaling pathways, among which are estrogens but also the anti-Müllerian
hormone (AMH) and gonadotropins (Grynberg *et al.* 2012; Dewailly *et al.* 2016; Chou and Chen
2018). Additional studies are underway to further analyze the mechanisms of differentiation
of these follicles in the absence of estradiol.
595
To conclude, the present work shows that in the rabbit species, the *CYP19A1* gene mutation
and the absence of estrogens from early embryo life was at the source of a series of main

disorders of the ovarian differentiation process. It did not disturb the sex determination of the

early fetal gonad but impacted profoundly the number of germ cells and the early

differentiation of somatic cells. It then had repercussions on the subsequent follicular growth

and differentiation and finally led to female infertility. This paper brings major and original

data on the role of estrogens in species where the *CYP19A1* gene is expressed in the fetal

ovary before meiosis. Deciphering the gene pathways sensitive to estrogens is currently

underway in our team in order to better understand specifically the role of endocrine

disruptors such as xeno-estrogens on female fertility.

Materials and methods

Animals

New Zealand rabbits (NZ1777, Hypharm, Rousssay, France) were bred at the SAAJ rabbit

facility (Jouy-en-Josas, France). All experiments were performed with the approval of the

French Ministry MENESR (accreditation number APAFIS#6775-2016091911407980 vI) following

the recommendation given by the local committee for ethic in animal experimentation

(COMETHEA, Jouy-en-Josas). All researchers working directly with the animals possessed an

animal experimentation license delivered by the French veterinary services. Hormonal

treatments for superovulation and surgery for embryo transfer were performed as previously

described (Peyny *et al.* 2020).

Design of TALEN sequences and plasmid constructions

Exon II of rabbit AROMATASE gene was targeted to create InDel mutations near the site of

initiation of translation (ATG site). The target sequences (left arm: 5'-TGCTTCATCTGAAGCCA-

3' (sense); right arm: 5'-TGGGTTCAGTATTTCCA-3' (antisense); 16 bases spaced) were chosen

with the ZiFiT Targeter sofware (http://zifit.partners.org). No homology was identified at any

other location in the rabbit genome that could represent a potential off-target site

(supplementary table 1). The TALEN were constructed as described (Sander *et al.* 2011), see

in supplementary methods). Each TALEN RNA was diluted (100 ng/µl) in injection buffer

(Millipore, France) and stored at -80°C until used.

Generation of mutant rabbits

Embryos produced from superovulated females were injected at single-cell stage with an

equimolar mixture of the left and right arm TALEN mRNAs (50ng/µl each). Injected embryos

were implanted 3-4 hours after injection into the oviducts of anesthetized recipient rabbits

via laparotomy. For all details concerning handling of females and embryos, see in (Peyny *et*

al. 2020).

Offspring were screened for the presence of InDel mutations using genomic DNA extracted

from ear clips (Jolivet *et al.* 2014). First detection of founders and characterization of mutation

were performed using one set of primers located far upstream and downstream the targeted

ATG (F0/R0, Figure 1 and supplementary table 2 for all primer sequences). The amplified

fragments were sequenced (Eurofins Genomics, Courtaboeuf, France) and the extent of the

mutation was deduced by comparing with the sequence of a wild type rabbit.

For further routine screening of mutants, a quantitative PCR (Fast SYBR Green Master Mix,

Applied Biosystems, ThermoFisher, France) was performed using two sets of primers, the

CYP19A1 specific one (*CYP19A1* gene specific primers) flanking the position of the

characterized InDel mutations (F1/R1, supplementary table 2) and another set to amplify a 2

copies reference gene (the rabbit *beta GLOBIN* gene, ENSOCUG00000000568). By using the

$\Delta(\Delta Ct)$ method with a wild type rabbit DNA as reference DNA, we deduced the copy number

of the *CYP19A1* allele in each DNA sample (no amplification = the two alleles are mutant (*ARO^{-}*

$^{/-}$ genotype); one copy = one mutant allele (*ARO$^{-/+}$* genotype); two copies = wild type (WT or

ARO$^{+/+}$ genotype).

The presence/absence of the Y chromosome was deduced from the amplification of the *SRY*

gene through PCR analyses with the rabbit *beta GLOBIN* gene as positive control of the PCR

and *SRY* specific primers (supplementary table 2). All rabbits were tested to determine the

genomic sex in parallel to the external and histological observations. In the present paper,

mentions of the XY or XX genotype always refer to the PCR determination.

Histological and immunohistological analyses

Immediately after sampling, gonads were immersed in Bouin's fixative or paraformaldehyde

(4% PAF in PBS 1x), fixed for 24 to 72 hours then paraffin embedded. Microtome sections of

5µm thickness were processed. Hematoxylin-eosin-saffron (HES) colorations were performed

by the @Bridge platform (INRAE, Jouy-en-Josas) using an automatic Varistain Slide Stainer

(Thermo Fisher Scientific). Sirius Red-Fast Green colorations were performed manually.

In situ hybridization (ISH) was performed using the RNAscope ISH methodology (ACD, Bio-

Techne SAS, Rennes, France). Briefly, probes at around 1000nt long were designed and

produced by the manufacturer in order to match to the full-length cDNA of interest, taking

care to reduce any cross-hybridization with non-specific targets. The list of all synthesized

probes is given in supplementary table 3. Hybridization was performed on 5μm sections from

PAF fixed tissue using labelling kits (RNAscope 2.5HD assay-brown or -red, or RNAscope 2.5HD

duplex chromogenic assay blue (conjugated to horse radish peroxidase) and red (conjugated

to alkaline phosphatase)) as recommended by the manufacturer. The red labelling (Fast Red)

was observed as visible or fluorescent signals. Hybridization was considered as positive when

at least one dot was observed in one cell.

Immunohistological analyses were performed on 5μm sections from PAF fixed tissue. The

supplementary table 4 recapitulates the antibodies used.

All colored sections (visible, or fluorescent ones) were scanned using a 3DHISTECH panoramic

scanner at the @Bridge platform (INRAE, Jouy-en-Josas).

RNA extraction and RT-qPCR analyses

Gonads from rabbit fetuses were collected and immediately frozen at -80°C. Total RNA from

each gonad was extracted using the RNeasy® MicroKit (Qiagen, France). Quantitative PCR was

performed on reverse transcribed RNAs (High Capacity Reverse cDNA Transcription kit with

the included set of random primers, Applied Biosystems, ThermoFisher, France). Several PCR

sets of primers and analyses were those published previously (Daniel-Carlier *et al.* 2013).

Other sets of primers are given in the supplementary table 2. Data from significant and specific

amplifications only were reported, thus excluding all amplifications with too high or variable

C_t values, and non-specific or multiple amplifications.

Measurement of estradiol, testosterone and anti Mülllerian hormone levels in serum

samples and fetal gonads

Estradiol and testosterone were assayed by GC/MS according to the protocol described by

Giton et al. (Giton *et al.* 2015) with modifications (Devillers *et al.* 2019). Female rabbit serum

from $ARO^{-/-}$ genotype was used as the matrix for calibrators and quality control (QC) standards

after twice charcoal dextran treatments. Sample extraction and purification, derivatization

and determination of estradiol and testosterone levels in serum samples and fetal gonads are

described in details as supplementary information (see also supplementary Table 5) or can be

provided upon request. Anti Müllerian hormone levels were determined in 50 µl aliquots of

serum samples by using an ELISA kit (AMH GenII ELISA, with AMH Gen II calibrators and

controls, Beckman Coulter, Villepinte, France) as previously described (Bourdon *et al.* 2018).

Statistics

The statistical analyses were performed using the GraphPad Prism 7 Software (GraphPad

Software Inc., La Jolla, CA, USA). Because of the small number of samples in groups,

comparisons between values were made by the Mann-Whitney test for non-parametric

values. A probability lower than 0.05 was required for significance.

Acknowledgements

The authors would like to thank the staff of the facility (SAAJ, INRAE, Jouy-en-Josas) for the

care of the rabbits and Julie Rivière and Marthe Vilotte (UMR GABI, INRAE, Jouy-en-Josas) for

their assistance on the histological plateau (@Bridge platform) and for the access to the virtual

slide scanner (MIMA2 platform). Special thanks go to Danielle Monniaux (UMR PRC, INRAE,

Nouzilly) for the time she spent in enriching discussions and highly valuable comments. This

study was supported by ANR grants (GENIDOV:ANR-09-GENM-009; ARGONADS: ANR-13-

BSV2-0017; ARDIGERM: ANR-2020-CE14). The authors declare that they have no competing

interests.

714   References
715
716   Akcurin, S., Turkkahraman, D., Kim, W.Y., Durmaz, E., Shin, J.G., and Lee, S.J. (2016) A
717   Novel Null Mutation in P450 Aromatase Gene (CYP19A1) Associated with Development of
718   Hypoplastic Ovaries in Humans. *J Clin Res Pediatr Endocrinol* **8**(2), 205-10
719
720   Amano, A., Kondo, Y., Noda, Y., Ohta, M., Kawanishi, N., Machida, S., Mitsuhashi, K.,
721   Senmaru, T., Fukui, M., Takaoka, O., Mori, T., Kitawaki, J., Ono, M., Saibara, T., Obayashi,
722   H., and Ishigami, A. (2017) Abnormal lipid/lipoprotein metabolism and high plasma
723   testosterone levels in male but not female aromatase-knockout mice. *Arch Biochem Biophys*
724   **622**, 47-58
725
726   Auersperg, N., Wong, A.S.T., Choi, K.-C., Kang, S.K., and Leung, P.C.K. (2001) Ovarian
727   Surface Epithelium: Biology, Endocrinology, and Pathology*. *Endocr Rev* **22**(2), 255-288
728
729   Boulanger, L., Pannetier, M., Gall, L., Allais-Bonnet, A., Elzaiat, M., Le Bourhis, D., Daniel,
730   N., Richard, C., Cotinot, C., Ghyselinck, N.B., and Pailhoux, E. (2014) FOXL2 is a female sex-
731   determining gene in the goat. *Curr Biol* **24**(4), 404-8
732
733   Bouraïma, H., Hanoux, V., Mittre, H., Féral, C., Benhaïm, C., and Leymarie, P. (2001)
734   Expression of tthe rabbit cytochroma P450 aromatase encoding gene uses alternative tissue-
735   specific promoters. *Eur. J. Biochem.* **268**, 4506-4512
736
737   Bourdon, M., Torres-Rovira, L., Monniaux, D., Faure, C., Levy, R., Tarrade, A., Rousseau-
738   Ralliard, D., Chavatte-Palmer, P., and Jolivet, G. (2018) Impact of a gestational exposure to
739   diesel exhaust on offspring gonadal development: experimental study in the rabbit. *J Dev Orig*
740   *Health Dis* **9**(5), 519-529
741
742   Britt, K.L., Drummond, A.E., Cox, V.A., Dyson, M., Wreford, N.G., Jones, M.E., Simpson,
743   E.R., and Findlay, J.K. (2000) An age-related ovarian phenotype in mice with targeted
744   disruption of the Cyp 19 (aromatase) gene. *Endocrinology* **141**(7), 2614-23
745
746   Britt, K.L., Drummond, A.E., Dyson, M., Wreford, N.G., Jones, M.E., Simpson, E.R., and
747   Findlay, J.K. (2001) The ovarian phenotype of the aromatase knockout (ArKO) mouse. *J*
748   *Steroid Biochem Mol Biol* **79**(1-5), 181-5
749
750   Chassot, A.A., Bradford, S.T., Auguste, A., Gregoire, E.P., Pailhoux, E., de Rooij, D.G.,
751   Schedl, A., and Chaboissier, M.C. (2012) WNT4 and RSPO1 together are required for cell
752   proliferation in the early mouse gonad. *Development* **139**(23), 4461-72
753
754   Chou, C.H., and Chen, M.J. (2018) The Effect of Steroid Hormones on Ovarian Follicle
755   Development. *Vitam Horm* **107**, 155-175
756
757   Chretien, F.C. (1966) [A study of the origin, migration and multiplication of the germ-cells of
758   the rabbit embryo]. *J Embryol Exp Morphol* **16**(3), 591-607
759
760   Conley, A., and Hinshelwood, M. (2001) Mammalian aromatases. *Reproduction* **121**(5), 685-
761   95
762

763   Couse, J.F., Hewitt, S.C., Bunch, D.O., Sar, M., Walker, V.R., Davis, B.J., and Korach, K.S.
764   (1999) Postnatal sex reversal of the ovaries in mice lacking estrogen receptors alpha and beta.
765   *Science* **286**(5448), 2328-31
766

767   Daniel-Carlier, N., Harscoet, E., Thepot, D., Auguste, A., Pailhoux, E., and Jolivet, G. (2013)
768   Gonad differentiation in the rabbit: evidence of species-specific features. *PLoS One* **8**(4),
769   e60451
770

771   Devillers, M.M., Petit, F., Cluzet, V., Francois, C.M., Giton, F., Garrel, G., Cohen-Tannoudji,
772   J., and Guigon, C.J. (2019) FSH inhibits AMH to support ovarian estradiol synthesis in infantile
773   mice. *J Endocrinol* **240**(2), 215-228
774

775   Dewailly, D., Robin, G., Peigne, M., Decanter, C., Pigny, P., and Catteau-Jonard, S. (2016)
776   Interactions between androgens, FSH, anti-Müllerian hormone and estradiol during
777   folliculogenesis in the human normal and polycystic ovary. *Hum Reprod Update* **22**(6), 709-
778   724
779

780   Diaz-Hernandez, V., Leon del Rio, A., Zamora, M., and Merchant-Larios, H. (2008) Expression
781   profiles of SRY and SOX9 in rabbit gonads: the classical model of mammalian sex
782   differentiation. *Sex Dev* **2**(3), 152-66
783

784   Dickmann, Z., Dey, S.K., and Gupta, J.S. (1975) Steroidogenesis in rabbit preimplantation
785   embryos. *Proc Natl Acad Sci U S A* **72**(1), 298-300
786

787   Dutta, S., Mark-Kappeler, C.J., Hoyer, P.B., and Pepling, M.E. (2014) The steroid hormone
788   environment during primordial follicle formation in perinatal mouse ovaries. *Biol Reprod*
789   **91**(3), 68
790

791   Elbrecht, A., and Smith, R.G. (1992) Aromatase enzyme activity and sex determination in
792   chickens. *Science* **255**(5043), 467-70
793

794   Findlay, J.K., Britt, K., Kerr, J.B., O'Donnell, L., Jones, M.E., Drummond, A.E., and Simpson,
795   E.R. (2001) The road to ovulation: the role of oestrogens. *Reprod Fertil Dev* **13**(7-8), 543-7
796

797   Fisher, C.R., Graves, K.H., Parlow, A.F., and Simpson, E.R. (1998) Characterization of mice
798   deficient in aromatase (ArKO) because of targeted disruption of the cyp19 gene. *Proc Natl*
799   *Acad Sci U S A* **95**(12), 6965-70
800

801   Gadsby, J.E., Heap, R.B., and Burton, R.D. (1980) Oestrogen production by blastocyst and
802   early embryonic tissue of various species. *J Reprod Fertil* **60**(2), 409-17
803

804   Garverick, H.A., Juengel, J.L., Smith, P., Heath, D.A., Burkhart, M.N., Perry, G.A., Smith,
805   M.F., and McNatty, K.P. (2010) Development of the ovary and ontongeny of mRNA and
806   protein for P450 aromatase (arom) and estrogen receptors (ER) alpha and beta during early fetal
807   life in cattle. *Anim Reprod Sci* **117**(1-2), 24-33
808

809   Gentilini, D., Busacca, M., Di Francesco, S., Vignali, M., Vigano, P., and Di Blasio, A.M.
810   (2007) PI3K/Akt and ERK1/2 signalling pathways are involved in endometrial cell migration
811   induced by 17beta-estradiol and growth factors. *Mol Hum Reprod* **13**(5), 317-22
812

813   George, F.W., Milewitch, L., and Wilson, J.D. (1978) Oestrogen content of the embryonic
814   rabbit ovary. *Nature* **274**, 172-173
815
816   George, F.W., and Wilson, J.D. (1978a) Conversion of Androgen to Estrogen by the Human
817   Fetal Ovary. *J Clin Endocr Metab* **47**, 550-555
818
819   George, F.W., and Wilson, J.D. (1978b) Estrogen formation in the early rabbit embryo. *Science*
820   **199**, 200-201
821
822   Giton, F., Sirab, N., Franck, G., Gervais, M., Schmidlin, F., Ali, T., Allory, Y., de la Taille, A.,
823   Vacherot, F., Loric, S., and Fiet, J. (2015) Evidence of estrone-sulfate uptake modification in
824   young and middle-aged rat prostate. *J Steroid Biochem Mol Biol* **152**, 89-100
825
826   Gondos, B., George, F., and Wilson, J. (1983) Granulosa cell differentiation and estrogen
827   synthesis in the fetal rabbit ovary. *Biol Reprod* **29**(3), 791-798
828
829   Govoroun, M.S., Pannetier, M., Pailhoux, E., Cocquet, J., Brillard, J.P., Couty, I., Batellier, F.,
830   and Cotinot, C. (2004) Isolation of chicken homolog of the FOXL2 gene and comparison of its
831   expression patterns with those of aromatase during ovarian development. *Dev Dyn* **231**(4), 859-
832   70
833
834   Greco, T.L., and Payne, A.H. (1994) Ontogeny of expression of the genes for steroidogenic
835   enzymes P450 side-chain cleavage, 3 beta-hydroxysteroid dehydrogenase, P450 17 alpha-
836   hydroxylase/C17-20 lyase, and P450 aromatase in fetal mouse gonads. *Endocrinology* **135**(1),
837   262-8
838
839   Grynberg, M., Pierre, A., Rey, R., Leclerc, A., Arouche, N., Hesters, L., Catteau-Jonard, S.,
840   Frydman, R., Picard, J.Y., Fanchin, R., Veitia, R., di Clemente, N., and Taieb, J. (2012)
841   Differential regulation of ovarian anti-mullerian hormone (AMH) by estradiol through alpha-
842   and beta-estrogen receptors. *J Clin Endocr Metab* **97**(9), E1649-57
843
844   Guiguen, Y., Fostier, A., Piferrer, F., and Chang, C.F. (2010) Ovarian aromatase and estrogens:
845   a pivotal role for gonadal sex differentiation and sex change in fish. *Gen Comp Endocrinol*
846   **165**(3), 352-66
847
848   Guioli, S., Nandi, S., Zhao, D., Burgess-Shannon, J., Lovell-Badge, R., and Clinton, M. (2014)
849   Gonadal asymmetry and sex determination in birds. *Sex Dev* **8**(5), 227-42
850
851   Guioli, S., Zhao, D., Nandi, S., Clinton, M., and Lovell-Badge, R. (2020) Oestrogen in the chick
852   embryo can induce chromosomally male ZZ left gonad epithelial cells to form an ovarian cortex
853   that can support oogenesis. *Development* **147**(4)
854
855   Hamilton, K.J., Arao, Y., and Korach, K.S. (2014) Estrogen hormone physiology: reproductive
856   findings from estrogen receptor mutant mice. *Reprod Biol* **14**(1), 3-8
857
858   Hudson, Q.J., Smith, C.A., and Sinclair, A.H. (2005) Aromatase inhibition reduces expression
859   of FOXL2 in the embryonic chicken ovary. *Dev Dyn* **233**(3), 1052-5
860

861 Hummitzsch, K., Irving-Rodgers, H.F., Hatzirodos, N., Bonner, W., Sabatier, L., Reinhardt,
862 D.P., Sado, Y., Ninomiya, Y., Wilhelm, D., and Rodgers, R.J. (2013) A new model of
863 development of the mammalian ovary and follicles. *PLoS One* **8**(2), e55578
864
865 Hussein, M.R. (2005) Apoptosis in the ovary: molecular mechanisms. *Hum Reprod Update*
866 **11**(2), 162-77
867
868 Jolivet, G., Braud, S., DaSilva, B., Passet, B., Harscoet, E., Viglietta, C., Gautier, T., Lagrost,
869 L., Daniel-Carlier, N., Houdebine, L.M., and Harosh, I. (2014) Induction of body weight loss
870 through RNAi-knockdown of APOBEC1 gene expression in transgenic rabbits. *PLoS One* **9**(9),
871 e106655
872
873 Juengel, J.L., Sawyer, H.R., Smith, P.R., Quirke, L.D., Heath, D.A., Lun, S., Wakefield, S.J.,
874 and McNatty, K.P. (2002) Origins of follicular cells and ontogeny of steroidogenesis in ovine
875 fetal ovaries. *Mol Cell Endocrinol* **191**(1), 1-10
876
877 Kaur, J., and Bose, H.S. (2014) Passenger protein determines translocation versus retention in
878 the endoplasmic reticulum for aromatase expression. *Mol Pharmacol* **85**(2), 290-300
879
880 Lee, J.G., Sung, Y.H., and Baek, I.-J. (2018) Generation of genetically-engineered animals
881 using engineered endonucleases. *Arch Pharm Res* **41**(9), 885-897
882
883 Lee, V.H., Britt, J.H., and Dunbar, B.S. (1996) Localization of laminin proteins during early
884 follicular development in pig and rabbit ovaries. *J Reprod Fertil* **108**(1), 115-22
885
886 Mauleon, P., Bezard, J., and Terqui, M. (1977) Very early and transient 17β-estradiol secretion
887 by fetalsheep ovary. In vitro study. . *Ann Biol Anim Bioch Bioph* **17**, 399-401
888
889 Mazen, I., McElreavey, K., Elaidy, A., Kamel, A.K., and Abdel-Hamid, M.S. (2017) Aromatase
890 Deficiency due to a Homozygous CYP19A1 Mutation in a 46,XX Egyptian Patient with
891 Ambiguous Genitalia. *Sex Dev* **11**(5-6), 275-279
892
893 Myers, M., Morgan, F.H., Liew, S.H., Zerafa, N., Gamage, T.U., Sarraj, M., Cook, M., Kapic,
894 I., Sutherland, A., Scott, C.L., Strasser, A., Findlay, J.K., Kerr, J.B., and Hutt, K.J. (2014)
895 PUMA regulates germ cell loss and primordial follicle endowment in mice. *Reproduction*
896 **148**(2), 211-9
897
898 Pannetier, M., Fabre, S., Batista, F., Kocer, A., Renault, L., Jolivet, G., Mandon-Pepin, B.,
899 Cotinot, C., Veitia, R., and Pailhoux, E. (2006) FOXL2 activates P450 aromatase gene
900 transcription: towards a better characterization of the early steps of mammalian ovarian
901 development. *J Mol Endocrinol* **36**(3), 399-413
902
903 Payen, E., Pailhoux, E., Abou Merhi, R., Gianquinto, L., Kirszenbaum, M., Locatelli, A., and
904 Cotinot, C. (1996) Characterization of ovine SRY transcript and developmental expression of
905 genes involved in sexual differentiation. *Int J Dev Biol* **40**(3), 567-75
906
907 Peters, H., Levy, E., and Crone, M. (1965) Oogenesis in Rabbits. *J Exp Zool* **158**, 169-79
908
909 Peyny, M., Jarrier-Gaillard, P., Boulanger, L., Daniel, N., Lavillatte, S., Cadoret, V., Papillier,
910 P., Monniaux, D., Peynot, N., Duranthon, V., Jolivet, G., and Dalbies-Tran, R. (2020)

911 Investigating the role of BCAR4 in ovarian physiology and female fertility by genome editing
912 in rabbit. *Sci Rep* **10**(1), 4992
913

914 Praveen, V.P., Ladjouze, A., Sauter, K.S., Pulickal, A., Katharopoulos, E., Trippel, M., Perren,
915 A., Pandey, A.V., and Fluck, C.E. (2020) Novel CYP19A1 Mutations Extend the Genotype-
916 Phenotype Correlation and Reveal the Impact on Ovarian Function. *J Endocr Soc* **4**(4), bvaa030
917

918 Ross, D.G., Bowles, J., Hope, M., Lehnert, S., and Koopman, P. (2009) Profiles of gonadal
919 gene expression in the developing bovine embryo. *Sex Dev* **3**(5), 273-83
920

921 Sander, J.D., Cade, L., Khayter, C., Reyon, D., Peterson, R.T., Joung, J.K., and Yeh, J.R. (2011)
922 Targeted gene disruption in somatic zebrafish cells using engineered TALENs. *Nat Biotechnol*
923 **29**(8), 697-8
924

925 Shemesh, M. (1980) Estradiol-17 beta biosynthesis by the early bovine fetal ovary during the
926 active and refractory phases. *Biol Reprod* **23**(3), 577-82
927

928 Torley, K.J., da Silveira, J.C., Smith, P., Anthony, R.V., Veeramachaneni, D.N., Winger, Q.A.,
929 and Bouma, G.J. (2011) Expression of miRNAs in ovine fetal gonads: potential role in gonadal
930 differentiation. *Reprod Biol Endocrinol* **9**, 2
931

932 Unal, E., Yildirim, R., Tas, F.F., Demir, V., Onay, H., and Haspolat, Y.K. (2018) Aromatase
933 Deficiency due to a Novel Mutation in CYP19A1 Gene. *J Clin Res Pediatr Endocrinol*
934

935 Vaillant, S., Dorizzi, M., Pieau, C., and Richard-Mercier, N. (2001) Sex reversal and aromatase
936 in chicken. *J Exp Zool* **290**(7), 727-40
937

938 Wilhelm, D., Palmer, S., and Koopman, P. (2007) Sex determination and gonadal development
939 in mammals. *Physiol Rev* **87**(1), 1-28
940

941 Yin, Y., Tang, H., Liu, Y., Chen, Y., Li, G., Liu, X., and Lin, H. (2017) Targeted Disruption of
942 Aromatase Reveals Dual Functions of cyp19a1a During Sex Differentiation in Zebrafish.
943 *Endocrinology* **158**(9), 3030-3041
944

945 Zalewski, A., Cecchini, E.L., and Deroo, B.J. (2012) Expression of extracellular matrix
946 components is disrupted in the immature and adult estrogen receptor beta-null mouse ovary.
947 *PLoS One* **7**(1), e29937
948

949 Zhu, W.J., Cheng, T., Zhu, H., Han, B., Fan, M.X., Gu, T., Zhao, S.X., Liu, Y., Cheng, K.X.,
950 Song, H.D., and Qiao, J. (2016) Aromatase deficiency: a novel compound heterozygous
951 mutation identified in a Chinese girl with severe phenotype and obvious maternal virilization.
952 *Mol Cell Endocrinol* **433**, 66-74
953